

# PRODUCT MONOGRAPH

## **KETODERM**

(Ketoconazole, USP)

**Cream 2%**

**Topical Antifungal Agent**

TaroPharma  
A Division of Taro Pharmaceuticals Inc.  
130 East Drive  
Brampton, Ontario  
L6T 1C1

Preparation Date:  
Sept 02, 2003

Revision Date:  
August 6, 2003

**Control # 086234**

## PRODUCT MONOGRAPH

### KETODERM Cream 2%

Ketoconazole, USP

#### Topical Antifungal Agent

#### ACTION

In vitro studies suggest that the antifungal properties of Ketoconazole may be related to its ability to impair the synthesis of ergosterol, a component of fungal and yeast cell membranes. Without the availability of this essential sterol, there are morphological alterations of the fungal and yeast cell membranes manifested as abnormal membranous inclusions between the cell wall and the plasma membrane. The inhibition of ergosterol synthesis has been attributed to interference with the reactions involved in the removal of the 14-" -methyl group of the precursor of ergosterol, lanosterol (1).

#### INDICATIONS

KETODERM (ketoconazole) cream 2% may be indicated for the topical treatment of tinea pedis, tinea corporis and tinea cruris caused by Trichophyton rubrum, T. mentagrophytes and Epidermophyton floccosum; and in the treatment of tinea versicolor (pityriasis) caused by Malassezia furfur (Pityrosporum orbiculare); and in the treatment of seborrhoeic dermatitis caused by Pityrosporum ovale; and in the treatment of cutaneous candidiasis caused by Candida albicans.

## **CONTRAINDICATIONS**

KETODERM (ketoconazole) cream 2% is contraindicated in persons who have shown hypersensitivity to the active or excipient ingredients of this formulation.

## **WARNINGS**

KETODERM (ketoconazole) cream 2% should never be employed for the treatment of infections of the eye.

To prevent a rebound effect after stopping a prolonged treatment with topical corticosteroids, it is recommended to continue applying a mild topical corticosteroid in the morning and to apply KETODERM cream in the evening, and to subsequently and gradually withdraw the steroid therapy over a period of 2-3 weeks.

## **PRECAUTIONS**

If a reaction suggesting sensitivity or chemical irritation should occur, use of KETODERM (ketoconazole) cream 2% should be promptly discontinued.

Limited short term studies in animals and in human volunteers on whom limited quantities of KETODERM cream 2% were tested have failed to demonstrate absorption of ketoconazole in detectable amounts. Due to the teratogenic nature of the active ingredient, ketoconazole, caution should be exercised when KETODERM cream 2% is administered to pregnant or nursing women.

Cross sensitivity with miconazole and other imidazoles may exist and caution is suggested when KETODERM cream 2% is employed in patients with known sensitivity

to imidazoles.

### **ADVERSE REACTIONS**

Short-term studies indicate that Ketoconazole Cream 2% is well tolerated by the skin. During clinical trials, 43 (5.0%) of 867 patients treated with the cream and 3 (1.8%) of 167 patients treated with placebo reported side effects consisting mainly of severe irritation, pruritus and stinging. One of the patients treated with Ketoconazole Cream 2% developed a painful allergic reaction (swelling of the foot).

In rare circumstances, allergic local skin phenomena such as contact dermatitis have been associated with ketoconazole cream 2% or one of its components, namely sodium sulfite or propylene glycol.

### **SYMPTOMS AND TREATMENT OF OVERDOSAGE**

There has been no experience with overdosage of Ketoconazole Cream 2%. Treatment should include general supportive measures.

### **DOSAGE AND ADMINISTRATION**

When clinically warranted, therapy with KETODERM (ketoconazole) cream 2% may be initiated while results of culture and susceptibility tests are pending. Treatment should be adjusted according to the findings.

KETODERM cream 2% should be applied to the affected and immediate surrounding area in patients with the following conditions:

<u>CONDITIONS</u>	<u>FREQUENCY</u>	<u>DURATION</u>
Tinea pedis	once daily	4-6 weeks
Tinea corporis	once daily	3-4 weeks
Tinea cruris	once daily	2-4 weeks
Tinea versicolor	once daily	2-3 weeks
Cutaneous candidiasis	once daily	2-3 weeks

More resistant cases may be treated twice daily depending on patient response.

Seborrheic dermatitis	twice daily	4 weeks
-----------------------	-------------	---------

The full course of therapy should be followed to reduce the possibility of recurrence. If however, there is no response within the recommended treatment period, the diagnosis should be re-evaluated.

The safety of Ketoconazole cream 2% has not been established with treatment periods exceeding those recommended, therefore, treatment must not exceed the recommended duration of therapy indicated above.


## PHARMACEUTICAL INFORMATION

### Drug Substance

Proper Name: ketoconazole

Chemical Name: cis-1-acetyl-4-[4-[[2-(2,4-dichlorophenyl)-2-(1H-imidazol-1-ylmethyl)-1,3-dioxolan-4-yl] methoxy] phenyl] piperazine

Structural Formula:



Molecular formula: C<sub>26</sub>H<sub>28</sub>Cl<sub>2</sub>N<sub>4</sub>O<sub>4</sub>

Molecular weight: 531.44

Description: Ketoconazole is an almost white to slightly beige coloured powder which is freely soluble in chloroform, methanol and diluted hydrochloric acid; sparingly soluble in 2-propanol and acetone and practically insoluble in water.

### Composition:

KETODERM cream 2% contains the broad-spectrum synthetic antifungal agent, ketoconazole, formulated in an aqueous cream vehicle consisting of propylene glycol,

stearyl and cetyl alcohols, sorbitan monostearate, polysorbate 60, isopropyl myristate, butylated hydroxyanisole (BHA), polysorbate 80 and water.

Stability and Storage Recommendations:

Store at room temperature, between 15° and 25°C. Keep from freezing.

**AVAILABILITY OF DOSAGE FORM**

KETODERM cream 2% is a white odourless cream containing 20 mg ketoconazole per gram and is supplied in 30 g tubes.

**MYCOLOGY**

In Vitro

In yeast and fungal cells, ergosterol is the main sterol regulating membrane permeability. Ketoconazole inhibits the biosynthesis of ergosterol and affects the synthesis of triglycerides and phospholipids.




Diagram 1: Site of action of ketoconazole (K) on the steroid biosynthetic pathway in the fungal cell.

Morphologically, ketoconazole-induced alterations are characterized by the presence of abnormal membranous inclusions between the cell wall and plasma membrane.

Changes in oxidative and peroxidative enzyme activities, leading to an intracellular build up of toxic concentrations of hydrogen peroxide, may contribute to the observed deterioration of subcellular organelles and to cell necrosis.

In vitro studies have shown that ketoconazole is fungistatic at low concentrations (0.01 - 0.1 µg/mL) and fungicidal at very high concentrations (1 - 10 µg/mL) against dermatophytes.

Table 1  
MIC of Ketoconazole against Dermatophytes and Yeasts (2)

Organism	No. of strains tested	Range of minimal inhibitory concentrations ( µg/mL)
<b>Dermatophytes</b>		
Microsporum canis	24	0.1 - 64
Microsporum audouini	4	2 - 64
Microsporum gypseum	9	0.1 - 64
Microsporum cookei	1	1
Trichophyton mentagrophytes	24	0.1 - 20
Trichophyton rubrum	75	10 <sup>-5</sup> - 128
Trichophyton ajelloi	1	1
Trichophyton schoenleini	1	1
Trichophyton tonsurans	35	0.25 - 16
Epidermophyton floccosum	23	0.1 - 8
<b>Yeasts</b>		
Candida albicans	472	0.02 - 80



Candida tropicalis	45	0.1 - 64
Candida pseudotropicalis	2	25 - 50
Candida guilliermondii	4	0.4 - 50
Candida krusei	14	0.2 - 3.1
Candida parapsilosis	18	0.2 - 64
Candida stellatoidea	1	0.8
Cryptococcus neoformans	39	0.1 - 32
Torulopsis glabrata	124	0.8 - 64
Rhodotorula mucilanginosa	1	0.1
Trichosporon cutaneum	1	0.1

---

In vivo studies have shown ketoconazole topical cream to be effective against dermatophyte infections of the skin in guinea pigs. Applied at concentrations > 1.0% in a PEG base once a day for 14 days, ketoconazole cream healed or improved experimental infections due to Trichophyton mentagrophytes and Microsporum canis.

### **PHARMACOLOGY**

The systemic absorption of Ketoconazole cream 2% was not measurable in Beagle dogs. When applied to both intact and abraded skin at a daily dose of 80 mg for 28 days, the cream produced plasma levels of ketoconazole which did not exceed the detection limit of the HPLC- method (0.002 µg/mL).

The systemic absorption of Ketoconazole cream 2% was not detectable in man. A single application of 10 g cream to the back, arms and chest did not produce quantifiable blood levels.

## TOXICOLOGY

### Animal Studies

#### Subacute Dermal Toxicity in New Zealand White Rabbits and Albino Guinea Pigs

Ketoconazole cream 2% at doses of 0, 0.5, 1.0 and 2.0 g/kg was devoid of any irritating effect in 16 male and 16 female New Zealand white rabbits (4 per group). Daily applications to the intact or abraded skin up to 2 g/kg for 30 days produced a barely perceptible irritation that was no different from that with 2 g/kg of the vehicle; a single application of 0.1 mg to the conjunctiva of 6 male New Zealand white rabbits was not irritating.

Ketoconazole cream 2% was also devoid of any allergenic or sensitizing potential in 10 male and 10 female albino guinea pigs. Induction comprised 10 topical applications of 0.5 mg Ketoconazole cream 2% using occlusive patches and 2 intradermal injections of Freund's complete adjuvant over a 4 week period. Subsequently, animals were challenged with a single 48-hour application of the same dose. This dose was applied to a region which had not been previously treated. Neither the patches used for the induction experiment nor those used in the challenge reaction elicited oedema or erythema.

#### Carcinogenicity

When ketoconazole was admixed in the diet, such that daily oral doses up to 80 mg/kg/day were provided to Swiss Albino mice for a period of 18 months and also to Wistar rats for a period of 24 months, there was no evidence of oncogenic activity.

## Mutagenicity

The dominant lethal mutation test in male and female mice revealed that single oral doses of ketoconazole as high as 80 mg/kg produced no mutation at any stage of germ cell development. The Ames Salmonella microsomal activator assay was also negative.

## Reproduction and Teratology

Ketoconazole has been shown to be embryotoxic and teratogenic (syndactylia and oligodactylia) in the rat when given in the diet at 80 mg/kg/day. When ketoconazole was given to rats by gavage, evidence of maternal toxicity and embryotoxicity was seen with doses as low as 10 mg/kg.

## **Human Studies**

### Dermal Irritancy

In 10 female volunteers treated topically with 5  $\mu\text{g}/\text{cm}^2$  Ketoconazole cream 2%, exposure to longwave ultraviolet or visible light did not produce any wheal and flare reaction.

Twenty-five volunteers received a series of 6 applications (2 per week) of 10  $\mu\text{g}/\text{cm}^2$  of Ketoconazole cream 2% on an occlusive dressing applied to the back skin for 24 hours. Following removal of the patch, the skin was exposed to three minimal erythema doses from a Xenon Solar Simulator. Ten days after the last exposure the subjects were challenged with 5  $\mu\text{g}/\text{cm}^2$  Ketoconazole cream 2% applied to an untreated site of the skin for 24 hours and subsequently exposed to long ultraviolet light. Ketoconazole cream 2% was shown to be devoid of any detectable photocontact allergenic potential.

## REFERENCES

1. Borgers M, Van den Bossche H and De Brabander M.: The Mechanism of Action of the New Antimycotic Ketoconazole. American Journal of Medicine. 74: 2-8, January 1983.
2. Levine HB.: Introduction to Antifungal Imidazoles. In: Ketoconazole in the Management of Fungal Disease, Ed. H.B. Levine, Adis Press, Hong Kong, P. 47-53, 1982.
3. Van Cutsem J, Van Gerven F, and Zaman R.: Topical Treatment of Dermatophytosis and Skin Candidosis with Ketoconazole in Guinea Pigs. Janssen Pharmaceutica, Preclinical Research Report R 41 400/45, November 1982.
4. Michiels M, Van Peer A, Van Ravestijn C, et al.: Absorption of Ketoconazole in the dog after Repeated Vaginal Application and after Application on the Intact and Abraded Skin. Janssen Pharmaceutica, Preclinical Research Report R 41 400/41, August 1982.
5. Janssen Pharmaceutica. Subacute Dermal Toxicity in New Zealand White Rabbits. Toxicological Research Report on Ketoconazole 2% Cream, No. 1109, October 1981.
6. Janssen Pharmaceutica. Acute Eye Irritation Study in New Zealand White Rabbits. Toxicological Research Report on Ketoconazole 2% Cream, No. 1103. October 1981.
7. Janssen Pharmaceutica. Sensitizing Potential Study in Guinea Pigs. Toxicological Research Report on Ketoconazole 2% Cream. No. 1106, October 1981.

8. Rawlins M.: Systemic Absorption of Ketoconazole 2% Cream and Ketoconazole 400 mg Vaginal Pessaries. Janssen UK, Clinical Research Report, August 1982.
9. Cunliffe WJ.: A Double-blind Placebo-controlled Cross-over Study to Determine the Incidence and Severity of Dermal Irritancy of Ketoconazole 2% Cream. Janssen UR, Clinical Research Report, August 1982.
10. Leyden JJ.: Determination of the Irritancy Potential of One Material (Ketoconazole Cream 2% R 41 400 - Lot ~ 81L03/F12). IVY Research Laboratories. Inc.. April 1982.
11. Leyden JJ, and Kligman AM.: Determination of the Contact-sensitizing Potential of Ketoconazole Cream 2% by the Maximization Test. IVY Research Laboratories. Inc.. May 1982.
12. Leyden JJ, and Kaidbey KH.: Determination of the Phototoxic Potential of Ketoconazole 2% Cream in Humans. IVY Research Laboratories. Inc., March 1982.
13. Leyden JJ, and Kaidbey KH.: The Photoallergenic Potential of Ketoconazole 2% Cream. IVY Research Laboratories, Inc., May 1982.
14. Taplin D.: Double-blind comparison of 2% Ketoconazole Cream (R 41 400) and Placebo in the Treatment of Tinea Corporis and Tinea Cruris. Janssen Pharmaceutica, Clinical Research Report R 41 400/101, March 1983.
15. Taplin D, and Carmargo G.: Double-blind Comparison of 2% Ketoconazole Cream (R 41 400) and Placebo in the Treatment of Tinea Pedis. Janssen Pharmaceutica, Clinical Research Report R 41 400/98, March 1983.
16. Savin RC.: Double-blind Comparison of 2% Ketoconazole Cream (R 41 400) and Placebo in the Treatment of Tinea Versicolor. Janssen Pharmaceutica,

Clinical Research Report R 41 400/100, March 1983.

17. Jolly HW, and Greer D.: Double-blind Comparison of Two Doses of 2% Ketoconazole Cream (R 41 400) in the Treatment of Chronic and Acute Tinea Pedis. Janssen Pharmaceutica, Clinical Research Report R 41 400/99, March 1983.
18. Satriano RC.: Ketoconazole 2% Cream: an Open Comparison between One and Two Daily Applications in Patients with Dermatophyte Infections of the skin. Janssen Pharmaceutica, Clinical Research Report R 41 400/95, February 1983.
19. Meisel C.: Single-blind, Side-to-Side Comparison of Ketoconazole 2% Cream and Clotrimazole 1% Cream in 8 Patients with Superficial Mycotic Infections. Janssen Pharmaceutica, Clinical Research Report R 41 400/93, March 1983.
20. Qadripur SA, Hellgren L, Gjertsen 8T, and Espedal A.: Double-blind Comparison of Ketoconazole 2% Cream and Clotrimazole 1% Cream in the Treatment of Superficial Fungal Infections. An International Multicenter Study. Janssen Pharmaceutica, Clinical Research Report R 41 400/97, March 1983.
21. Carabelli S, Gimenez MM, and Amdur A.: Topical Treatment of Superficial Dermatophyte or Yeast Infections with Ketoconazole 2% Cream. Janssen Pharmaceutica, Clinical Research Report R 41 400/92, March 1983.
22. Aussems J, Delescluse J, Ghekiere L, et al.: A Belgian Multicentre, Open Evaluation of Ketoconazole 2% Cream in the Treatment of Dermatophyte and Yeast Infections of the skin. Janssen Pharmaceutica. Clinical Research Report R 41 400/94, March 1983.
23. Meisel C, and Konietzko D.: Ketoconazole Cream 2% in the Treatment of Superficial Fungal Infections. Janssen Pharmaceutica, Clinical Research Report R 41 400/96, March 1983.

24. Del Palacia A, and Hernandez B.: Treatment of Superficial Fungal Infections with ketoconazole 2% Cream. Janssen Pharmaceutica, Clinical Research Report R 41 400/91, March 1983.
25. Degreef H et al.: Ketoconazole in the treatment of dermatophyte infections. A clinical, mycological and morphological study. International Journal of Dermatology 20: 662-669, 1981.
26. Smith EB: Ketoconazole 2% Cream. Seminars in Dermatology 6 (1): 62-65, 1987.