

PRODUCT MONOGRAPH

RAMACE[®]

(ramipril)

1.25 mg, 2.5 mg, 5.0 mg and 10.0 mg Tablets

ANGIOTENSIN CONVERTING ENZYME INHIBITOR

sanofi-aventis Canada Inc.
2150 St. Elzear Blvd. West
Laval, Québec H7L 4A8

Date of Revision:
October 24, 2006

Submission Control No. 107870

s-a Version 2.0 dated

**PRODUCT MONOGRAPH
RAMACE®**

(ramipril)

1.25 mg, 2.5 mg, 5.0 mg and 10.0 mg Tablets

PHARMACOLOGIC CLASSIFICATION

Angiotensin Converting Enzyme Inhibitor

ACTION AND CLINICAL PHARMACOLOGY

RAMACE® (ramipril) is an angiotensin converting enzyme (ACE) inhibitor, which is used in the treatment of essential hypertension, following acute myocardial infarction in stabilized patients with clinically confirmed heart failure, and for the management of patients at increased risk of cardiovascular events.

Following oral administration, RAMACE® is rapidly hydrolyzed to ramiprilat, its principal active metabolite.

Angiotensin-converting enzyme catalyzes the conversion of angiotensin I to the vasoconstrictor substance, angiotensin II. Angiotensin II also stimulates aldosterone secretion by the adrenal cortex. Inhibition of ACE activity leads to decreased levels of angiotensin II thereby resulting in decreased vasoconstriction and decreased aldosterone secretion. The latter decrease may result in a small increase in serum potassium (see PRECAUTIONS). Decreased levels of angiotensin II and the accompanying lack of negative feedback on renal renin secretion result in increases in plasma renin activity.

ACE is identical to kininase II. Thus, ramipril may also block the degradation of the vasodepressor peptide bradykinin, which may contribute to its therapeutic effect.

Pharmacokinetics and Metabolism

Following oral administration, ramipril is rapidly absorbed with peak plasma concentrations occurring within 1 hour. The extent of absorption of ramipril is 50-60% and is not significantly altered by the presence of food in the gastrointestinal tract, although the rate of absorption is reduced. Following absorption, ramipril is rapidly hydrolyzed in the liver to its active metabolite, ramiprilat. Peak plasma concentrations of ramiprilat are reached 2-4 hours after drug intake. The serum protein binding of ramipril is about 73% and that of ramiprilat is 56%.

Ramipril is almost completely metabolized to the active metabolite ramiprilat, and to the diketopiperazine ester, the diketopiperazine acid, and the glucuronides of ramipril and

ramiprilat, all of which are inactive. After oral administration of RAMACE[®], about 60% of the parent drug and its metabolites is excreted in the urine, and about 40% is found in the feces. Drug recovered in the feces may represent both biliary excretion of metabolites and/or unabsorbed drug. Less than 2% of the administered dose is recovered in urine as unchanged ramipril.

Following a single administration of up to 5 mg of Ramipril, plasma concentrations of ramipril and ramiprilat increase in a manner that is greater than proportional to dose; after a single administration of 5 mg to 20 mg of ramipril the plasma concentrations for both are dose-proportional. The non-linear pharmacokinetics observed at the lower doses of ramipril can be explained by the saturable binding of ramiprilat to ACE. At steady-state, the 24-hour AUC for ramiprilat is dose-proportional over the recommended dose range. The absolute bioavailabilities of ramipril and ramiprilat were 28% and 44% respectively when 5mg of oral ramipril was compared to 5mg given intravenously.

Plasma concentrations of ramiprilat decline in a triphasic manner. The initial rapid decline, which represents distribution of the drug, has a half life of 2-4 hours. Because of its potent binding to ACE and slow dissociation from the enzyme, ramiprilat shows two elimination phases. The apparent elimination phase has a half-life of 9-18 hours, and the terminal elimination phase has a prolonged half-life of > 50 hours. After multiple daily doses of ramipril 5-10mg, the half-life of ramiprilat concentrations was 13-17 hours, but was considerably prolonged at 2.5mg (27-36 hours).

After once daily dosing, steady state plasma concentrations of ramiprilat are reached by the fourth dose. Steady-state concentrations of ramiprilat are higher than those seen after the first dose of RAMACE[®] especially at low doses (2.5mg).

The urinary excretion of ramipril, ramiprilat, and their metabolites is reduced in patients with impaired renal function. In patients with creatinine clearance < 40 ml/min/1.73m², increases in C_{max} and AUC of ramipril and ramiprilat compared to normal subjects were observed following multiple dosing with 5mg ramipril (see DOSAGE AND ADMINISTRATION).

In patients with impaired liver function, plasma ramipril levels increased about 3-fold, although peak concentrations of ramiprilat in these patients were not different from those seen in patients with normal hepatic function.

A single dose pharmacokinetic study conducted in a limited number of elderly patients indicated that peak ramiprilat levels and the AUC for ramiprilat are higher in older patients (see PRECAUTIONS).

Pharmacodynamics

Administration of RAMACE® to patients with mild to moderate essential hypertension results in a reduction of both supine and standing blood pressure usually with little or no

orthostatic change or change in heart rate. Symptomatic postural hypotension is infrequent, although this may occur in patients who are salt-and/or volume-depleted (see WARNINGS).

In single dose studies, doses of 5-20 mg of RAMACE[®] lowered blood pressure within 1-2 hours, with peak reductions achieved 3-6 hours after dosing. At recommended doses given once daily, antihypertensive effects have persisted over 24 hours.

The effectiveness of RAMACE[®] appears to be similar in the elderly (over 65 years of age) and younger adult patients given the same daily doses.

In studies comparing the same daily dose of RAMACE[®] given as a single morning dose or as a twice daily dose, blood pressure reductions at the time of morning trough blood levels were greater with the divided regimen.

While the mechanism through which RAMACE[®] lowers blood pressure appears to result primarily from suppression of the renin-angiotensin-aldosterone system, RAMACE[®] has an antihypertensive effect even in patients with low-renin hypertension.

The antihypertensive effect of angiotension converting enzyme inhibitors is generally lower in black patients than in non-blacks.

The antihypertensive effect of RAMACE[®] and thiazide diuretics used concurrently is greater than that seen with either agent used alone.

Abrupt withdrawal of RAMACE[®] has not resulted in rapid increase in blood pressure.

The effects of ramipril were assessed in patients who were at high risk for cardiovascular events, but did not have left ventricular dysfunction or heart failure. Heart Outcome Prevention Evaluation (HOPE) study included 9297 patients older than 55 years of age with a history of coronary artery disease, stroke, peripheral vascular disease or diabetes mellitus plus at least one additional cardiovascular risk factor (hypertension, elevated total cholesterol levels, low high-density lipoprotein cholesterol levels, cigarette smoking, or documented microalbuminuria). Patients were excluded if they had heart failure, low ejection fraction (<0.40), were taking an angiotensin converting enzyme inhibitor or vitamin E, had uncontrolled hypertension or overt nephropathy, or had had a myocardial infarction or stroke within four weeks before the study began. The patients were randomly assigned to receive ramipril 10 mg once daily or matching placebo for a mean of five years.

Due to the positive outcome the study was terminated prematurely by an independent monitoring board. The primary end point, the composite of death from cardiovascular causes, myocardial infarction and stroke was reached by a total of 651 ramipril treated patients (14%), as compared to 826 placebo treated patients (17.8%) (relative risk, 0.78; $P < 0.001$). When analyzed separately, the rates of individual component of the composite primary outcome in patients treated with ramipril and placebo were as follows: death from cardiovascular causes 6.1% vs. 8.1% (RR 0.74, $p < 0.001$), myocardial infarction 9.9% vs. 12.3% (RR 0.80, $p < 0.001$) and stroke 3.4% vs. 4.9% of patients (RR 0.68, $p < 0.001$), respectively.

Permanent discontinuation of treatment occurred in 28.9% of the ramipril treated patients versus 27.3% of placebo treated patients. The reasons for stopping the treatment, where the incidence was greater in the ramipril than in the placebo group, were cough (ramipril 7.3%, placebo 1.8%), hypotension/dizziness (ramipril 1.9%, placebo 1.5%) and edema (ramipril 0.4%, placebo 0.2%).

INDICATIONS AND CLINICAL USE

Essential Hypertension

RAMACE® (ramipril) is indicated in the treatment of essential hypertension. It may be used alone or in association with thiazide diuretics.

RAMACE® should normally be used in patients in whom treatment with a diuretic or a beta blocker was found ineffective or has been associated with unacceptable adverse effects.

RAMACE® can also be tried as an initial agent in those patients in whom use of diuretics and/or beta blockers are contraindicated or in patients with medical conditions in which these drugs frequently cause serious adverse effects.

The safety and efficacy of RAMACE® in renovascular hypertension have not been established and therefore, its use in this condition is not recommended.

The safety and efficacy of concurrent use of RAMACE® with antihypertensive agents other than thiazide diuretics have not been established.

Treatment Following Acute Myocardial Infarction

RAMACE® is indicated following acute myocardial infarction in clinically stable patients with signs of left ventricular dysfunction to improve survival and reduce hospitalizations for heart failure.

Sufficient experience in the treatment of patients with severe (NYHA class IV) heart failure immediately after myocardial infarction is not yet available. (See WARNINGS - Hypotension)

Management of Patients at Increased Risk of Cardiovascular Events

RAMACE® may be used to reduce the risk of myocardial infarction, stroke or cardiovascular death in patients over 55 years of age who are at high risk of cardiovascular events because of a history of coronary artery disease, stroke, peripheral artery disease, or diabetes that is accompanied by at least one other cardiovascular risk factor such as hypertension, elevated total cholesterol levels, low high density lipoprotein levels, cigarette smoking, or documented microalbuminuria.

The incidence of the primary outcome (composite of myocardial infarction, stroke and death from cardiovascular causes) was reduced from 17.8% in the placebo-treated group to 14.0% in the ramipril-treated group (see ACTION AND CLINICAL PHARMACOLOGY).

General

In using RAMACE® consideration should be given to the risk of angioedema (see WARNINGS).

CONTRAINDICATIONS

RAMACE® (ramipril) is contraindicated:

- In patients who are hypersensitive to this drug, or to any ingredient in the formulation, or in those patients who have a history of angioedema.
- During pregnancy
- In breast-feeding women

WARNINGS

Serious Warnings and Precautions

When used in pregnancy, angiotensin converting enzyme (ACE) inhibitors can cause injury or even death of the developing fetus. When pregnancy is detected, RAMACE® should be discontinued as soon as possible.

Angioedema:

Angioedema has been reported in patients with ACE inhibitors including RAMACE® (ramipril). Angioedema associated with laryngeal involvement may be fatal. If laryngeal stridor or angioedema of the face, tongue, or glottis occurs, RAMACE® should be discontinued immediately, the patient treated appropriately in accordance with accepted medical care, and carefully observed until the swelling disappears. In instances where swelling is confined to the face and lips, the condition generally resolves without treatment, although antihistamines may be useful in relieving symptoms. Where there is involvement of tongue, glottis, or larynx, likely to cause airway obstruction, appropriate therapy (including, but not limited to 0.3 to 0.5ml of subcutaneous epinephrine solution 1:1000) should be administered promptly (see ADVERSE REACTIONS).

The incidence of angioedema during ACE inhibitor therapy has been reported to be higher in black than in non-black patients.

Patients with a history of angioedema unrelated to ACE inhibitor therapy may be at increased risk of angioedema while receiving an ACE inhibitor (see CONTRAINDICATIONS).

Hypotension:

Symptomatic hypotension has occurred after administration of RAMACE[®], usually after the first or second dose or when the dose was increased. It is more likely to occur in patients who are volume depleted by diuretic therapy, dietary salt restriction, dialysis, diarrhea, or vomiting. In patients with ischemic heart disease or cerebrovascular disease, an excessive fall in blood pressure could result in a myocardial infarction or cerebrovascular accident (see ADVERSE REACTIONS). Because of the potential fall in blood pressure in these patients, therapy with RAMACE[®] should be started under close medical supervision. Such patients should be followed closely for the first weeks of treatment and whenever the dose of RAMACE[®] is increased. In patients with severe congestive heart failure, with or without associated renal insufficiency, ACE inhibitor therapy may cause excessive hypotension and has been associated with oliguria, and/or progressive azotemia, and rarely, with acute renal failure and/or death.

If hypotension occurs, the patient should be placed in a supine position and, if necessary, receive an intravenous infusion of 0.9% sodium chloride. A transient hypotensive response may not be a contraindication to further doses which usually can be given without difficulty once the blood pressure has increased after volume expansion in hypertensive patients. However, lower doses of RAMACE[®] and/or reduced concomitant diuretic therapy should be considered. In patients receiving treatment following acute myocardial infarction, consideration should be given to discontinuation of RAMACE[®] (see ADVERSE REACTIONS - Treatment Following Acute Myocardial Infarction, DOSING & ADMINISTRATION - Treatment Following Acute Myocardial Infarction).

Neutropenia/agranulocytosis:

Agranulocytosis and bone marrow depression have been caused by ACE inhibitors. Several cases of agranulocytosis, neutropenia or leukopenia have been reported in which a causal relationship to RAMACE[®] cannot be excluded. Current experience with the drug shows the incidence to be rare. Periodic monitoring of white blood cell counts should be considered, especially in patients with collagen vascular disease and/or renal disease.

Pregnant Women

ACE inhibitors can cause fetal and neonatal morbidity and mortality when administered to pregnant women. When pregnancy is detected, RAMACE[®] should be discontinued as soon as possible.

The use of ACE inhibitors during the second and third trimesters of pregnancy has been associated with fetal and neonatal injury including hypotension, neonatal skull hypoplasia, anuria, reversible or irreversible renal failure, and death. Oligohydramnios

has also been reported, presumably resulting from decreased fetal renal function, associated with fetal limb contractures, craniofacial deformation, and hypoplastic lung development.

Prematurity, and patent ductus arteriosus and other structural cardiac malformations, as well as neurologic malformations, have also been reported following exposure in the first trimester of pregnancy.

Infants with a history of in utero exposure to ACE inhibitors should be closely observed for hypotension, oliguria, and hyperkalemia. If oliguria occurs, attention should be directed toward support of blood pressure and renal perfusion. Exchange transfusion or dialysis may be required as a means of reversing hypotension and/or substituting for impaired renal function; however, limited experience with those procedures has not been associated with significant clinical benefit.

It is not known if ramipril or ramiprilat can be removed from the body by hemodialysis.

Animal Data: No teratogenic effects of ramipril were seen in studies of pregnant rats, rabbits, and cynomolgus monkeys. The doses used were: 10, 100, or 1000 mg/kg in rats (2500 times maximum human dose), 0.4, 1.0, or 2.5 mg/kg in rabbits (6.25 times maximum human dose), and 5, 50, or 500 mg/kg in cynomolgus monkeys (1250 times maximum human dose). In rats, the highest dose caused reduced food intake in the dams, with consequent reduced birth weights of the pups and weight development during the lactation period. In rabbits, maternal effects were mortalities (high and middle dose) and reduced body weight. In monkeys, maternal effects were mortalities (high and middle dose), vomiting, and reduced weight gain.

Nursing Women

The presence of concentrations of ACE inhibitor have been reported in human milk. The use of RAMACE® is contraindicated during breast-feeding.

PRECAUTIONS

Renal Impairment:

As a consequence of inhibiting the renin-angiotensin-aldosterone system, changes in renal function have been seen in susceptible individuals. In patients whose renal function may depend on the activity of the renin-angiotensin-aldosterone system, such as patients with bilateral renal artery stenosis, unilateral renal artery stenosis to a solitary kidney, or severe congestive heart failure, treatment with agents that inhibit this system has been associated with oliguria, progressive azotemia, and rarely, acute renal failure and/or death. In susceptible patients, concomitant diuretic use may further increase risk.

Use of RAMACE® should include appropriate assessment of renal function.

RAMACE[®] should be used with caution in patients with renal insufficiency as they may require reduced or less frequent doses (see DOSAGE AND ADMINISTRATION). Close monitoring of renal function during therapy should be performed as deemed appropriate in patients with renal insufficiency.

Anaphylactoid reactions during membrane exposure:

Anaphylactoid reactions have been reported in patients dialyzed with high-flux membranes [e.g. polyacrylonitrile (PAN)] and treated concomitantly with an ACE inhibitor. Dialysis should be stopped immediately if symptoms such as nausea, abdominal cramps, burning, angioedema, shortness of breath and severe hypotension occur. Symptoms are not relieved by antihistamines. In these patients consideration should be given to using a different type of dialysis membrane or a different class of antihypertensive agents.

Anaphylactoid reactions during LDL apheresis

Rarely, patients receiving ACE inhibitors during low density lipoprotein apheresis with dextran sulfate have experienced life-threatening anaphylactoid reactions. These reactions were avoided by temporarily withholding the ACE inhibitor therapy prior to each apheresis.

Anaphylactoid reactions during desensitization

There have been isolated reports of patients experiencing sustained life threatening anaphylactoid reactions while receiving ACE inhibitors during desensitization treatment with hymenoptera (bees, wasps) venom. In the same patients, these reactions have been avoided when ACE inhibitors were temporarily withheld for at least 24 hours, but they have reappeared upon inadvertent rechallenge.

Hyperkalemia and Potassium-Sparing Diuretics

Elevated serum potassium (greater than 5.7 mEq/L) was observed in approximately 1% of hypertensive patients in clinical trials treated with RAMACE[®]. In most cases these were isolated values which resolved despite continued therapy. Hyperkalemia was not a cause of discontinuation of therapy in any hypertensive patient. Risk factors for the development of hyperkalemia may include renal insufficiency, diabetes mellitus, and the concomitant use of agents to treat hypokalemia or other drugs associated with increases in serum potassium (see PRECAUTIONS - Drug Interactions).

Surgery/anesthesia:

In patients undergoing surgery or anesthesia with agents producing hypotension, RAMACE[®] may block angiotensin II formation secondary to compensatory renin release. If hypotension occurs and is considered to be due to this mechanism, it may be corrected by volume repletion.

Aortic Stenosis:

There is concern, on theoretical grounds, that patients with aortic stenosis might be at particular risk of decreased coronary perfusion when treated with vasodilators because they do not develop as much afterload reduction.

Patients with Impaired Liver Function:

Hepatitis (hepatocellular and/or cholestatic), elevations of liver enzymes and/or serum bilirubin have occurred during therapy with ACE inhibitors in patients with or without pre-existing liver abnormalities. In most cases the changes were reversed on discontinuation of the drug.

Elevations of liver enzymes and/or serum bilirubin have been reported with RAMACE[®] (see ADVERSE REACTIONS). Should the patient receiving RAMACE[®] experience any unexplained symptoms particularly during the first weeks or months of treatment, it is recommended that a full set of liver function tests and any other necessary investigations be carried out. Discontinuation of RAMACE[®] should be considered when appropriate.

There are no adequate studies in patients with cirrhosis and/or liver dysfunction. RAMACE[®] should be used with particular caution in patients with pre-existing liver abnormalities. In such patients baseline liver function tests should be obtained before administration of the drug and close monitoring of response and metabolic effects should apply.

Pediatric Use:

The safety and effectiveness of RAMACE[®] in children have not been established; therefore use in this age group is not recommended.

Use in Elderly:

Although clinical experience has not identified differences in response between the elderly (> 65 years) and younger patients, greater sensitivity of some older individuals cannot be ruled out (see ACTION AND CLINICAL PHARMACOLOGY, Pharmacokinetics).

Patient Alertness:

RAMACE[®] may lower the state of patient alertness and/or reactivity, particularly at the start of treatment (see ADVERSE REACTIONS).

Cough:

A dry, persistent cough, which usually disappears only after withdrawal or lowering of the dose of RAMACE[®], has been reported. Such possibility should be considered as part of the differential diagnosis of cough (see ADVERSE REACTIONS).

Drug Interactions:

Concomitant Diuretic Therapy: Patients concomitantly taking ACE inhibitors and

diuretics, and especially those in whom diuretic therapy was recently instituted, may occasionally experience an excessive reduction of blood pressure after initiation of therapy. The possibility of hypotensive effects after the first dose of RAMACE® can be minimized by either discontinuing the diuretic or increasing the salt intake prior to initiation of treatment with RAMACE®. If it is not possible to discontinue the diuretic, the starting dose of RAMACE® should be reduced and the patient should be closely observed for several hours following the initial dose and until blood pressure has stabilized (see WARNINGS and DOSAGE AND ADMINISTRATION).

Agents Increasing Serum Potassium: Since RAMACE® decreases aldosterone production, elevation of serum potassium may occur. Potassium sparing diuretics such as spironolactone, triamterene or amiloride, or potassium supplements should be given only for documented hypokalemia and with caution and frequent monitoring of serum potassium, since they may lead to a significant increase in serum potassium. Salt substitutes which contain potassium should also be used with caution. (See also PRECAUTIONS - Drug Interactions - Non-steroidal anti-inflammatory agents)

Agents Causing Renin Release: The antihypertensive effect of RAMACE® is augmented by antihypertensive agents that cause renin release (e.g. diuretics).

Lithium: Increased serum lithium levels and symptoms of lithium toxicity have been reported in patients receiving ACE inhibitors during therapy with lithium. These drugs should be administered with caution, and frequent monitoring of serum lithium levels is recommended. If a diuretic is also used, the risk of lithium toxicity may be further increased.

Antacids: In one open-label, randomized, cross-over single dose study in 24 male subjects, it was determined that the bioavailability of RAMACE® and the pharmacokinetic profile of ramiprilat were not affected by concomitant administration of the antacid, magnesium and aluminum hydroxides.

Digoxin: In one open-label study in 12 subjects, administered multiple doses of both ramipril and digoxin, no changes were found in serum levels of ramipril, ramiprilat, and digoxin.

Warfarin: The co-administration of RAMACE® with warfarin did not alter the anticoagulant effects.

Acenocoumarol: In a multi-dose double-blind, placebo-controlled, pharmacodynamic interaction study with 14 patients with mild hypertension administered both ramipril and therapeutic doses of acenocoumarol, blood pressure, thrombotest time and coagulation factors were not significantly changed.

Non-steroidal anti-inflammatory agents: The antihypertensive effects of ACE inhibitors may be reduced with concomitant administration of non-steroidal anti-inflammatory agents (e.g. indomethacin). Concomitant treatment of ACE inhibitors

and Non-Steroidal Anti-Inflammatory drugs may lead to an increased risk of worsening of renal function and an increase in serum potassium. (See also PRECAUTIONS - Drug Interactions - Agents Increasing Serum Potassium)

Antidiabetic agents (e.g. insulin and sulfonylurea derivatives): ACE inhibitors may reduce insulin resistance. In isolated cases, such reduction may lead to hypoglycaemic reactions in patients concomitantly treated with antidiabetics. Particularly close blood glucose monitoring is, therefore, recommended in the initial phase of co-administration.

INFORMATION FOR THE PATIENT

Angioedema: Angioedema, including laryngeal edema, may occur especially following the first dose of RAMACE[®]. Patients should be so advised and told to report immediately any signs or symptoms suggesting angioedema, such as swelling of face, extremities, eyes, lips, tongue, difficulty in swallowing or breathing. They should immediately stop taking RAMACE[®] and consult with their physician (see WARNINGS).

Hypotension: Patients should be cautioned to report lightheadedness, especially during the first few days of RAMACE[®] therapy. If actual syncope occurs, the patients should be told to discontinue the drug and consult with their physician.

All patients should be cautioned that excessive perspiration and dehydration may lead to an excessive fall in blood pressure because of reduction in fluid volume. Other causes of volume depletion such as vomiting or diarrhea may also lead to a fall in blood pressure, patients should be advised to consult with their physician.

Agranulocytosis/Neutropenia: Patients should be told to report promptly to their physician any indication of infection (e.g. sore throat, fever) as this may be a sign of neutropenia (see WARNINGS and ADVERSE REACTIONS).

Impaired Liver Function: Patients should be advised to return to their physician if they experience any symptoms possibly related to liver dysfunction. This would include "viral-like symptoms" in the first weeks to months of therapy (such as fever, malaise, muscle pain, rash or adenopathy which are possible indicators of hypersensitivity reactions), or if abdominal pain, nausea or vomiting, loss of appetite, jaundice, itching or any other unexplained symptoms occur during therapy.

Hyperkalemia: Patients should be told not to use salt substitutes containing potassium without consulting their physician (see PRECAUTIONS).

ADVERSE REACTIONS

Essential Hypertension

RAMACE[®] (ramipril) has been evaluated for safety in over 4000 hypertensive patients. Almost 500 elderly patients have participated in controlled trials. Long-term safety has been assessed in almost 700 patients treated for 1 year or more. There was no increase in the incidence of adverse events in elderly patients given the same daily dose. The overall frequency of adverse events was not related to duration of therapy or total daily dose.

Serious adverse events occurring in North American placebo-controlled clinical trials with ramipril monotherapy in hypertension (n=972) were: hypotension (0.1%); myocardial infarction (0.3%); cerebrovascular accident (0.1%); edema (0.2%); syncope (0.1%). Among all North American ramipril patients (n=1,244), angioedema occurred in 0.1% patients treated with ramipril and a diuretic.

The most frequent adverse events occurring in these trials with RAMACE[®] monotherapy in hypertensive patients that were treated for at least one year (n=651) were: headache (15.1%); dizziness (3.7%); asthenia (3.7%); chest pain (2.0%); nausea (1.8%); peripheral edema (1.8%); somnolence (1.7%); impotence (1.5%); rash (1.4%); arthritis (1.1%); dyspnea (1.1%). Discontinuation of therapy due to clinical adverse events was required in 5 patients (0.8%).

In placebo-controlled trials, an excess of upper respiratory infection and flu syndrome was seen in the ramipril group. As these studies were carried out before the relationship of cough to ACE inhibitors was recognized, some of these events may represent ramipril-induced cough. In a later 1-year study, increased cough was seen in almost 12% of RAMACE[®] patients, with about 4% of these patients requiring discontinuation of treatment. Approximately 1% of patients treated with RAMACE[®] monotherapy in North American controlled clinical trials (n=972) have required discontinuation because of cough.

Treatment Following Acute Myocardial Infarction

1004 post-AMI patients received RAMACE® in a controlled clinical trial. In both the ramipril and placebo groups, myocardial infarction, heart failure, atrial fibrillation, peripheral vascular disease and urinary tract infection were more common in elderly than in younger patients. Gastrointestinal disturbances were more frequent in elderly patients on ramipril. Cough and hypotension were more frequent in women receiving ramipril.

Adverse events (except laboratory abnormalities) considered possibly/probably related to study drug that occurred in more than one percent of stabilized patients with clinical signs of heart failure treated with RAMACE® following an acute myocardial infarction are shown below. The incidences represent the experiences from the AIRE (Acute Infarction Ramipril Efficacy) study. The follow-up time was between 6 and 48 months for this study (mean follow up= 15 months).

Percentage of Patients with Adverse Events Possibly/Probably Related to Study Drug Placebo-Controlled (AIRE) Mortality Study		
Adverse Event	Ramipril (n=1004)	Placebo (n=982)
Hypotension	10.7	4.7
Cough increased	7.6	3.7
Dizziness/Vertigo	5.6	3.9
Nausea/Vomiting	3.8	1.9
Angina pectoris	2.9	2.0
Postural hypotension	2.2	1.4
Syncope	2.1	1.4
Heart failure	2.0	2.2
Severe/resistant heart failure	2.0	3.0
Myocardial infarct	1.7	1.7
Vomiting	1.6	0.5
Headache	1.2	0.8
Abnormal kidney function	1.2	0.5
Abnormal chest pain	1.1	0.9
Diarrhea	1.1	0.4

Event	Percentage of Patients with Serious Adverse Events Possibly related to Study Drug Placebo-Controlled (AIRE) Mortality Study	
	RAMACE [®] (n = 1004)	Placebo (n = 982)
Hypotension	3.0%	1.1%
Angina pectoris	2.0%	1.2%
Severe/resistant heart failure	1.9%	2.9%
Myocardial infarct	1.7%	1.7%
Heart failure	1.5%	1.5%
Syncope	1.3%	0.8%
Chest pain	0.7%	0.9%
Nausea	0.6%	0.5%
Vomiting	0.5%	0.1%
Dizziness	0.5%	0.5%
Abnormal kidney function	0.5%	0.2%
Chest infection	0.2%	0.0%
Postural hypotension	0.2%	0.2%
Headache	0.1%	0.0%

Isolated cases of death have been reported with the use of ramipril that appear to be related to hypotension (including first dose effects), but many of these are difficult to differentiate from progression of underlying disease (see WARNINGS - Hypotension).

Discontinuation of therapy due to adverse reactions was required in 368/1004 post-AMI patients taking ramipril (36.7%), compared to 401/982 patients receiving placebo (40.8%).

Management of Patients at Increased Risk of Cardiovascular Events

In the Heart Outcome Prevention Evaluation (HOPE) study, based on a total of 4645 patients treated with ramipril, the safety profile of RAMACE[®] was consistent with the post-marketing surveillance experience. The reasons for stopping the treatment, where the incidence was greater in the ramipril than in the placebo group, were cough (ramipril 7.3%, placebo 1.8%), hypotension/dizziness (ramipril 1.9%, placebo 1.5%) and edema (ramipril 0.4%, placebo 0.2%).

Clinical Adverse Events by Body System

Clinical adverse events occurring in less than 1% of patients treated with RAMACE[®] in controlled clinical trials, or seen in post-marketing experience, are listed below by body system:

Body as a whole: anaphylactoid reactions, angioneurotic edema.

Cardiovascular: symptomatic hypotension, syncope, angina pectoris, arrhythmia, chest pain, palpitations, tachycardia, myocardial infarction, cerebrovascular disorders (including ischaemic stroke).

Respiratory: increased cough.

CNS: anxiety, amnesia, confusion, convulsions, depression, hearing loss, insomnia, nervousness, neuralgia, neuropathy, paresthesia, polyneuritis, somnolence, tinnitus, tremor, vertigo, vision disturbances.

Dermatologic: apparent hypersensitivity reactions (with manifestations of urticaria, pruritus, or rash, with or without fever), photosensitivity, purpura, erythema multiforms, pemphigus, Stevens-Johnson syndrome.

In addition, the following cutaneous or mucosal reactions may occur: exacerbation of psoriasis, maculo-papular exanthema, psoriasiform exanthema, pemphigoid exanthema and enanthema, and toxic epidermal necrolysis or onycholysis.

Gastrointestinal: abdominal pain (sometimes with enzyme changes suggesting pancreatitis), pancreatitis, anorexia, constipation, diarrhea, dry mouth, dyspepsia, dysphagia, gastroenteritis, hepatitis, nausea, increased salivation, smell and taste disturbance, vomiting.

Renal: increases in blood urea nitrogen (BUN) and serum creatinine.

Haematologic: agranulocytosis, leucopenia, eosinophilia, thrombocytopenia, pancytopenia and hemolytic anemia.

Other: arthralgia, arthritis, dyspnea, edema, epistaxis, impotence, increased sweating, malaise, myalgia, weight gain.

A symptom complex has been reported which may include fever, vasculitis, myalgia, arthralgia/arthritis, a positive ANA, elevated ESR, eosinophilia and leucocytosis. Rash, photosensitivity or other dermatologic manifestations may also occur.

Clinical Laboratory Test Findings: increased creatinine; increases in blood urea nitrogen (BUN); decreases in hemoglobin or hematocrit; hyponatraemia; elevations of liver enzymes, serum bilirubin, uric acid, blood glucose; proteinuria and significant increases in serum potassium.

SYMPTOMS AND TREATMENT OF OVERDOSAGE

Limited data are available regarding overdosage of RAMACE® (ramipril) in humans. Two cases of overdosage have been reported.

In the case of an overdose with ramipril, the most likely clinical manifestation would be symptoms attributable to severe hypotension, which should normally be treated by intravenous volume expansion with normal saline. It is not known if ramipril or ramiprilat can be removed from the body by hemodialysis.

DOSAGE AND ADMINISTRATION

Essential Hypertension

Dosage of RAMACE® (ramipril) must be individualized. Initiation of therapy requires consideration of recent antihypertensive drug treatment, the extent of blood pressure elevation and salt restriction. The dosage of other antihypertensive agents being used with RAMACE® may need to be adjusted.

Monotherapy:

The recommended initial dosage of RAMACE® in patients not on diuretics is 2.5mg once daily. Dosage should be adjusted according to blood pressure response, generally, at intervals of at least two weeks. The usual dose range is 2.5 to 10mg once daily. A daily dose of 20mg should not be exceeded.

In some patients treated once daily, the antihypertensive effect may diminish towards the end of the dosing interval. This can be evaluated by measuring blood pressure just prior to dosing to determine whether satisfactory control is being maintained for 24 hours. If it is not, either twice daily administration with the same total daily dose, or an increase in dose should be considered. If blood pressure is not controlled with RAMACE® alone, a diuretic may be added. After the addition of a diuretic, it may be possible to reduce the dose of RAMACE®.

Concomitant Diuretic Therapy:

Symptomatic hypotension occasionally may occur following the initial dose of RAMACE® and is more likely in patients who are currently being treated with a diuretic. The diuretic should, if possible, be discontinued for two to three days before beginning therapy with RAMACE® to reduce the likelihood of hypotension (see WARNINGS). If the diuretic cannot be discontinued, an initial dose of 1.25 mg RAMACE® should be used with careful medical supervision for several hours and until blood pressure has stabilized. The dosage of RAMACE® should subsequently be titrated (as described above) to the optimal response.

Use in renal impairment:

For patients with a creatinine clearance below 40ml/min/1.73m² (serum creatinine above 2.5mg/dL), the recommended initial dose is 1.25mg RAMACE[®] once daily. Dosage may be titrated upward until blood pressure is controlled or to a maximum total daily dose of 5mg. In patients with severe renal impairment (creatinine clearance below 10ml/min/1.73m²) the maximum total daily dose of 2.5mg RAMACE[®] should not be exceeded.

Treatment Following Acute Myocardial Infarction

Dosage of RAMACE[®] (ramipril) must be individualized. Initiation of therapy requires consideration of concomitant medication and baseline blood pressure and should be instituted under close medical supervision, usually in a hospital, three to ten days following an acute myocardial infarction in haemodynamically stable patients with clinical signs of heart failure.

The recommended initial dosage of RAMACE[®] is 2.5 mg given twice a day (b.i.d.), one in the morning and one in the evening. If tolerated, and depending on the patient's response, dosage may be increased by doubling at intervals of one to three days. The maximum daily dose of RAMACE[®] should not exceed 5 mg twice daily (b.i.d.).

After the initial dose of RAMACE[®], the patient should be observed under medical supervision for at least two hours and until blood pressure has stabilized for at least an additional hour. If a patient becomes hypotensive at this dosage, it is recommended that the dosage be lowered to 1.25 mg b.i.d. following effective management of the hypotension. (see WARNINGS - Hypotension).

Patients who have been fluid or salt depleted, or treated with diuretics are at an increased risk of hypotension (see WARNINGS-Hypotension). An excessive fall in blood pressure may occur particularly in the following: after the initial dose of RAMACE[®]; after every first increase of dose of RAMACE[®]; after the first dose of a concomitant diuretic and/or when increasing the dose of the concomitant diuretic. If appropriate, the dose of any concomitant diuretic should be reduced which may diminish the likelihood of hypotension (see PRECAUTIONS - Drug Interactions). Consideration should be given to reducing the initial dose to 1.25 mg of RAMACE[®] in these patients.

Use in renal impairment:

In patients with impaired renal function (creatinine clearance of 20-50 mL/min/1.73 m² body surface area), the initial recommended dosage is generally 1.25 mg of RAMACE[®] once daily. This dosage may be increased with caution up to 1.25 mg of RAMACE[®] twice daily, depending upon clinical response and tolerability.

Insufficient data is available concerning the use of ramipril following acute myocardial infarction in patients with heart failure and severe renal failure. (see ACTION & CLINICAL PHARMACOLOGY - Pharmacokinetics & Metabolism, PRECAUTIONS - Renal Impairment).

Use in hepatic impairment:

Insufficient data is available concerning the use of ramipril following acute myocardial infarction in patients with heart failure and hepatic dysfunction. Dose reduction and careful monitoring of these patients is required (see ACTIONS & CLINICAL PHARMACOLOGY- Pharmacokinetics & Metabolism, PRECAUTIONS - Patients with Impaired Liver Function).

Management of Patients at Increased Risk of Cardiovascular Events

Recommended initial dose: 2.5 mg of RAMACE® once daily. Depending on the tolerability, the dose is gradually increased. It is recommended to double the dose after one week of treatment and - after another three weeks - to increase it to 10 mg. Usual maintenance dose: 10 mg of RAMACE® daily (see ACTION AND CLINICAL PHARMACOLOGY, WARNINGS and PRECAUTIONS).

Dosage recommendations for special risk groups such as patients with renal or hepatic impairment, or at an increased risk of hypotension (fluid or salt depletion, treated with diuretics) are to be followed as previously described (SEE WARNINGS and PRECAUTIONS).

PHARMACEUTICAL INFORMATION


I. DRUG SUBSTANCE

Proper name: Ramipril

Chemical name: 2-[N-[(S)-1-ethoxycarbonyl-3-phenylpropyl]-L-alanyl]-(1S,3S,5S)-2-azabicyclo-[3.3.0]octane-3-carboxylic acid

Empirical formula: $C_{23}H_{32}N_2O_5$

Structural formula:



Molecular weight: 416.52

Description: A white to off-white crystalline powder with a melting point of 105°C to 112°C. Slightly soluble in water, and freely soluble in ethanol and methanol.

II. DOSAGE FORM

a) Composition

RAMACE[®] (ramipril) 1.25 mg, 2.5 mg, 5.0 mg, and 10.0 mg tablets contain the medicinal ingredient ramipril in quantities of 1.25 mg, 2.5 mg, 5.0 mg, and 10.0 mg respectively.

The qualitative formulation for all potencies of RAMACE[®] is: ramipril, hydroxypropyl methylcellulose, pregelatinized starch, microcrystalline cellulose, sodium stearyl fumarate.

Yellow ferric oxide is used as colouring agent in the 2.5 mg tablets.

Red ferric oxide is used as colouring agent in the 5 mg tablets.

b) **Stability and storage recommendations**

Store RAMACE® (ramipril) in original container at room temperature, between 15 and 30°C and not beyond the date indicated on the container.

c) **Availability**

RAMACE® (ramipril) is available in the following potencies:

- 1.25 mg white to almost white oblong tablets with score-line. This tablet should not be halved.
Upper stamp: 1.25 and Hoechst
Lower stamp: HMN and 1.25
- 2.5 mg yellowish to yellow oblong tablets with score-line.
Upper stamp: 2.5 and Hoechst logo
Lower stamp: HMR and 2.5
- 5.0 mg Red oblong tablets with score-line.
Upper stamp: 5 and Hoechst
Lower stamp: 5 and HMP
- 10.0 mg white to almost white oblong tablets with score-line.
Upper stamp: HMO/HMO
Lower stamp: Plain

RAMACE® tablets 1.25 mg, 2.5 mg, 5.0 mg and 10.0 mg are packaged in cartons of 28 (2 x 14 blister-packed) tablets.

INFORMATION FOR THE CONSUMER

For Patients with Essential Hypertension

What is hypertension?

Hypertension is the medical term for high blood pressure. When blood flows through the blood vessels it pushes against their walls, almost like water pushing against the sides of a hose. Blood pressure is like that "push". When blood pressure is high (like the water pressure in a hose when the nozzle is partially shut), damage can occur to the heart and blood vessels.

Although you may not feel any symptoms for years, hypertension can lead to stroke, heart attack, kidney disease and other serious conditions.

What causes hypertension?

In most cases, the exact cause of hypertension is not known. But we do know that several factors increase the risk of developing the disease.

Family history: Hypertension, like some other diseases, can run in families. If your parents have high blood pressure, your chances of developing it are greater.

Age: The risk of developing hypertension increases with age.

Race: In North America, there is a higher incidence of hypertension among blacks than among whites.

Diabetes: Diabetics are at higher risk of developing hypertension than non-diabetics.

Weight: The risk of high blood pressure is higher in people who are overweight.

Drinking: Heavy alcohol consumption increases risk of hypertension, as well as stroke and kidney disease.

Sedentary lifestyle: A physically inactive lifestyle can contribute to hypertension.

Smoking: While not a direct cause of hypertension, smoking a cigarette will temporarily increase blood pressure. Smoking also increases the risk of heart disease in people with high blood pressure.

Keeping your blood pressure controlled

Your doctor has prescribed RAMACE[®], a medication that helps to control blood pressure. RAMACE[®] opens blood vessels to reduce blood pressure, like the way opening a hose reduces water pressure. It is not, however, a cure.

But it takes more than just medication to reduce blood pressure. Discuss the risk factors, and how they apply to your lifestyle, with your doctor. You may have to modify some of your daily habits to keep your blood pressure down.

Exercise regularly. It will help to keep your weight down, make you feel more energetic and is a good way to deal with stress. If you are not exercising regularly, be sure to discuss a fitness plan with your doctor.

Remember, hypertension is a long-term disease without symptoms. Just because you feel fine does not mean you can stop taking your medication. If you stop, serious complications of the disease may occur. Therefore, you should continue to take RAMACE[®] regularly, as prescribed by your doctor.

For Patients Following a Recent Heart Attack

Your doctor has prescribed RAMACE[®] to reduce the effort required by your heart to pump blood, to compensate for reduced pumping power that may have resulted from your heart attack.

RAMACE[®] has been shown to improve survival and reduce hospitalizations for heart failure in patients recovering from recent heart attacks.

If you have developed heart failure after a heart attack, you may have to limit your physical activities. Before you begin exercising, be sure to consult with your doctor.

When initiating therapy in patients with heart failure after a heart attack, RAMACE[®] is usually given twice a day, in the morning and in the evening. Take your medication as instructed by your Doctor.

Management of Patients at Increased Risk of Cardiovascular Events

Your doctor has prescribed RAMACE[®] because:

- you have coronary heart disease (such as chest pains or angina, or have had a heart attack in the past)
- you had a stroke
- you have peripheral vascular disease (poor blood circulation)
- you have diabetes and at least one of the following physical conditions: high blood pressure, elevated total cholesterol levels, low high-density lipoprotein (HDL) levels, cigarette smoking or documented microalbuminuria.

RAMACE[®] may lower the risk of heart attack, stroke, or death from a heart disease in some patients who have a heart problem or poor blood circulation. Take your medication as instructed by your Doctor.

GENERAL

When to take your medication

It is important to take it at the same time every day as prescribed by your doctor.

Missed a dose?

If you forgot to take your RAMACE® tablet, take it as soon as possible. However, if it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. Do not double dose.

Managing your lifestyle

The "lifestyle" part of your treatment is as important as your medication. By working as a team with your doctor, you can help reduce the risk of complications to maintain the style of life you are accustomed to.

Alcohol: Avoid alcoholic beverages until you have discussed their use with your doctor. Alcohol consumption may alter your blood pressure and/or increase the possibility of dizziness or fainting.

Diet: Generally, avoid fatty foods and food that is high in salt or cholesterol.

Smoking: Avoid it completely.

Side effects

Along with its intended action, any medication, including RAMACE®, may cause side effects. These include headache, dizziness, fatigue, nausea or coughing. These side effects may disappear once your system becomes used to the medication. If they persist, discuss this with your doctor. Your medication might have to be reduced or changed.

Dizziness or lightheadedness may occur after the first dose of this medicine. Make sure you know how you react to this medicine before you drive, operate machinery, or do anything requiring you to be alert.

If you are suffering from excessive sweating, vomiting or diarrhea, this may cause you to lose too much water and lead to problems with low blood pressure. See your doctor.

A rare, but potentially more serious, side effect is called angioedema - characterized by swollen mouth, lips, tongue, eyes, throat or difficulty in swallowing or breathing. **If you notice swelling or feel pain in these areas, inform your doctor immediately. You should also inform your doctor if you have an unexplained fever, rash or itching.**

Serious Warnings and Precautions

RAMACE[®] should not be used during pregnancy. If you discover that you are pregnant while taking RAMACE[®], stop the medication and please contact your physician as soon as possible.

Keep your doctor informed

Before taking RAMACE[®], it is important that you inform your doctor of the following:

- Are you currently taking any other medications, whether on prescription or otherwise? This is especially important if you are taking diuretics (water pills) or any other medication to reduce blood pressure which may add to the blood pressure lowering effect of RAMACE[®]. You should not be taking salt substitutes, potassium supplements or potassium containing medicine without the advice of your doctor.
- Do you suffer from any other condition? The presence of other medical problems may affect the use of RAMACE[®]. Make sure you tell your doctor if you have any other medical problems, especially if you have diabetes, liver disease, kidney disease, heart or blood vessel disease.

If you are being treated for other conditions by other doctors, keep them all informed of which medications you are taking. Some drugs may have negative effect on RAMACE[®] or RAMACE[®] may have a negative effect on other drugs. If you have to undergo any dental or other surgery, inform the dentist or physician in charge that you are taking this medicine.

- **You are pregnant, breast-feeding or thinking of becoming pregnant?** Taking RAMACE[®] during pregnancy can cause injury and even death to your baby. This medicine should not be used during pregnancy. If you become pregnant while taking RAMACE[®], stop the medication and report to your doctor as soon as possible. It is possible that RAMACE[®] passes into breast milk. You should not breast-feed while taking RAMACE[®].
- Are you possibly allergic to RAMACE[®] (ramipril), including any of its non-medicinal ingredients: hydroxypropyl methylcellulose; pregelatinized starch; microcrystalline cellulose; sodium stearyl fumarate; yellow ferric oxide; and red ferric oxide?

After you have started taking RAMACE[®], it is important that you tell your doctor at once about any unexplained symptom you might experience. Examples of this are unexplained fever, rash, itching, any sign of infection, viral-like symptoms, flu-like symptoms, coughing, sore throat, abdominal pain, loss of appetite, sad mood, or jaundice.

Store in original container at room temperature between 15⁰ and 30⁰ C and not beyond the date indicated on the container. Keep this medication out of the reach of children.

Remember

Use this drug as directed by your doctor. All drugs can have both helpful and harmful effects. Both depend on the person and his or her condition. This leaflet alerts you to some of the times you should call your doctor. Other situations which cannot be predicted can arise. Nothing in this leaflet should stop you from calling your doctor or pharmacist with any questions or concerns you have about RAMACE[®].

PHARMACOLOGY

Mechanism of Action

Study	Species	#/group	Route	Dose	Results
Inhibition of Angiotensin I-induced pressor response after oral ramipril	Rat Dog	n=6 n=3	oral oral	0.1 0.3 1.0 mg/kg	A dose-dependent inhibition was observed, lasting more than 6 hours
Effect of pre-treatment with ramipril on b.p. changes induced by i.v. Angiotensin I, Angiotensin II, and sympathomimetics	Rat	n=5 or n=6	oral	1.0 mg/kg	Effects of Ang. I and indirect-acting sympathomimetics are inhibited, while the effects of Ang. II and direct-acting sympathomimetics are unaffected by ramipril
Effect of ramipril on Na-depleted (furosemide treated) dogs	Dog	n=6	oral	10 mg/kg	Ramipril-induced increase in plasma renin activity is enhanced by furosemide; Ramipril has no influence on heart rate
In vitro inhibition of ACE by ramipril	Rabbit lung		in vitro		IC ₅₀ = 26±8 nmol/L
Effect of ramipril and captopril on renal blood flow, renal vasculature resistance, and blood pressure	Rat	n=5	i.a.	0.1 mg/kg	Ramipril caused a greater increase in renal blood flow and decrease in renal vasculature resistance than a 10-fold higher dose of captopril; this without the decrease in systemic b.p. observed with captopril

Effects on Blood Pressure

Hypertensive Model	Species	#/group	Route	Dose	Duration	Result
Spontaneously hypertensive rats	Rat	n=5	oral	1 mg/kg 0.01,0.1, 1,10 mg/ kg/day	acute 5 weeks	Significant decreases in b.p.(all doses); which persisted for: 2 weeks (chronic) 72 hrs. (acute)
Kidney perinephretic hypertension (no increase in plasma renin activity)	Dog	n=5	oral	10 mg/kg 1 mg/ kg/day	acute 5 days	Significant decrease of systemic blood pressure
2 kidney, 1 clip hypertension	Rat	n=8	oral	1,10 mg/kg	acute	Blood pressure was normalized
Release of an occluded renal pedicle	Rat	n=6	oral	0.1 mg/kg	acute	Hypertension was completely prevented

Pharmacokinetics and Bioavailability

Study Parameter (after oral ramipril)	Results		
	Rat (2 mg/kg)	Dog (2 mg/kg)	Human (10 mg)
GI absorption of ¹⁴ C-ramipril	56%	43%	56%
Maximal blood levels of radioactivity	0.5 hrs	0.5-1 hrs	0.3 hrs
Plasma t _{1/2} of radioactivity	0.6 hrs	1.0 and 3.8 hrs (biphasic)	0.5 and 2.9 hrs (biphasic)
Distribution of radioactivity	High concentration in liver, kidney and particularly lungs. Total fetus: 0.05% Breast milk: 0.25%	-	-
Serum protein binding (concentration range of 0.01-10 µg/ml)	ramipril: - ramiprilat: 41%	ramipril: 72% ramiprilat: 47%	ramipril: 73% ramiprilat: 56%
Metabolism	metabolized to ramiprilat	metabolized to ramiprilat and inactive diketopiperazines	
Excretion of radioactivity	urine: 26% feces: 71% t _{1/2} (both): 1.6-4.8 and 23-42 h	urine: 15% t _{1/2} : 9.3 h feces: 79% t _{1/2} : 8 h	urine: 56% t _{1/2} : 7.2 and 127 h feces: 40% t _{1/2} : 11 and 110 h

TOXICOLOGY

Acute Toxicity:

Below are summarized species-specific LD₅₀ values for both oral and intravenous administrations of ramipril.

Table 1 - Acute Toxicity

Routes	Species	Sex	LD₅₀
Oral	Mouse	Male	10,933 mg/kg
		Female	10,048 mg/kg
	Rat	Male	> 10,000 mg/kg
		Female	> 10,000 mg/kg
	Dog	Male	> 1,000 mg/kg
Intravenous	Mouse	Male	1,194 mg/kg
		Female	1,158 mg/kg
	Rat	Male	688 mg/kg
		Female	609 mg/kg

The symptoms observed in mice were decreased spontaneous activity, crouching, hypothermia, dyspnea, and clonic convulsions; deaths occurred within 30 minutes after intravenous and 24 hours after oral administration. In survivors, the symptoms disappeared by 1 to 5 days after administration; necropsies revealed no abnormality in any of the surviving animals. In rats, reduced spontaneous activity was noted (oral administration), while after intravenous administration similar signs occurred as in mice; the sign of lethal toxicity was clonic convulsions (intravenous administration).

Table 2 - Chronic Toxicity:

Species	Duration	No. of animals per group	Route	Dose (mg/kg/day)	Effects
Mouse	28 days 90 days	2M, 2F 3M, 3F	Oral	1000	Reduced erythrocytes, hemoglobin, hematocrit, increased reticulocytes. Hyperplasia of juxtaglomerular apparatus.
Rat	30 days	10-15M, 10-15F	Oral	2.5, 80, 2500	At all doses: decrease in body weight, reduced liver weight, increased kidney weight. At 80 & 2500 mg/kg/d: Reduced heart weight. At 2500 mg/kg/d: Reduced erythrocytes, hematocrit and bilirubin, increased BUN.
Rat	3 months	10-15M, 10-15F	Oral	2.5, 80, 500	At all doses: Reduced chloride and GOT, increased phosphorus and BUN. At 80 mg/kg/d: Reduced heart, liver, prostate weight, increased kidney weight. Atrophic segments of renal tubules. Increased serum creatinine. At 500 mg/kg/d: Reduced body and heart weight, increased kidney and adrenal weight. Reduced erythrocytes, hemoglobin, hematocrit, increased bilirubin. Increased number of atrophic renal tubular segments. Moderate gastric mucosa necroses.
Rat	3 months	10M, 10F	Oral	500, 1/3 Ringer solution for drinking	Increased number of tubular atrophies.

Table 2 - Chronic Toxicity:

Species	Duration	No. of animals per group	Route	Dose (mg/kg/day)	Effects
Rat	6 months	10-20M, 10-20F	Oral	0.1, 0.25, 3.2, 40, 500	At all doses: Serum bilirubin increased, reduced heart weight. At 40 and 500 mg/kg/d: Increased kidney weight. Reduced erythrocytes, hemoglobin, hematocrit, increased BUN. Distal tubular atrophies, fibromuscular pad formations in gastric mucosa/muscularis not proliferative in nature.
Rat	6 months	20M, 20F	Oral	3.2, 40, 500, 1/3 Ringer solution for drinking	All doses: Fibromuscular or solitary pad formation in gastric fundus mucosa/muscularis.
Rat	18 months	20-25M, 20-25F	Oral	0.25, 3.2, 40, 500	At 3.2 to 500 mg/kg/d: Fibromuscular pads in gastric fundus mucosa, focal atrophies in renal cortex, partly with cysts. At 40 and 500 mg/kg/d: Anemia, increased BUN and serum creatinine, urinary epithelial cells. Reduced heart weight and increased kidney and adrenal weight.
Dog	30 days	2M, 2F	Oral	3.2, 32	No pathological findings.
Dog	3 months	3-4M, 3-4F	Oral	3.2, 32, 320	At 320 mg/kg/d: Anemia, increased BUN and serum creatinine, impaired erythropoiesis. Juxtaglomerular hyperplasia.
Dog	6 months	6M, 6F	Oral	3.2, 32, 320	At 32 mg/kg/d: Anemia, juxtaglomerular hyperplasia. At 320 mg/kg/d: Reduced body weight. Increased BUN and serum creatinine. Distal tubular atrophies with round cell infiltrations. Anemia, juxtaglomerular hyperplasia.

Table 2 - Chronic Toxicity:

Species	Duration	No. of animals per group	Route	Dose (mg/kg/day)	Effects
Dog	12 months	6M, 6F	Oral	2.5, 25, 250	At all doses: Reduced body weight. At 25 and 250 mg/kg/d: Anemia and leukopenia, impaired erythropoiesis, increased hemosiderin deposition in liver and spleen, juxtaglomerular hyperplasia. At 250 mg/kg/d: Increased BUN and serum creatinine.
Monkey	6 months	4-5M, 4-5F	Oral	0.5, 16, 500	At 16 and 500 mg/kg/d: Increased BUN, juxtaglomerular hyperplasia. Reduced body weight. At 500 mg/kg/d: Diarrhea, anemia, increased serum creatinine, some urinary casts, leukocytes and epithelial cells.
Monkey	6 months	5M 5F	Oral	2, 8	No pathological findings.

Table 3 - Reproduction and Teratology:

Species	No. of animals per group	Dose (mg/kg/day)	Duration of dosing	Results
Rat (Wistar)	32M, 32F	5, 50, 500	M 60 days before mating F14 days before mating to end of lactation	At 50 and 500 mg/kg/d: Parents renal pelvis enlargement, off-spring light brown discoloration of kidney tissue and dilatation of renal pelvis. At 500 mg/kg/d: Parents yellow-white coloring and induration of renal marrow. Fertility normal.
Rat (Wistar)	20F	10, 100, 1000	Days 7-17 of gestation	At 1000 mg/kg/d: Reduced food consumption of mothers, reduced body weight gains of young. One young circular non-ossified area in supraoccipital bone, 1 young distortion of right scapula. No teratogenic effects.
Rat (Wistar)	20-30F	0.32, 1.25, 5, 10, 100, 1000	Day 17 of gestation to day 21 of lactation	At 100 and 1000 mg/kg/d: Decreased gestation body weight of young, enlarged to day 21 renal pelvis up to hydronephrosis with light brown coloring of renal cortex and marrow.
Rat (Sprague-Dawley)	20F	100	Day 17 of gestation to day 21 of lactation	Young: Enlarged renal pelvis and light brown coloration of kidney tissue.
Rabbit (Himalayan)	15F	0.4, 1, 2.5	Day 6 to day 18 of gestation	At 0.4 mg/kg/d: One abortion, one fetus with diaphragm hernia. At 1 mg/kg/d: One abortion, one premature delivery, two animals died, no animals gained weight. One dead fetus with possible hydrocephalus. At 2.5 mg/kg/d: Two animals died, no animals gained weight, one fetus with diaphragm hernia, one with first cervical aplasia and aplasia of one thorax vertebra and one rib pair.
Monkey (Cynomolgus)	4-13F	5, 50, 500	Days 20-25 of gestation	At all doses: No sign of terato-genesis. At 5 mg/kg/d: Two abortions, seven diarrhea, two vomiting, ten weight loss. At 50 mg/kg/d: One animal died, three abortions, seven diarrhea, two vomiting, ten weight loss. At 500 mg/kg/d: Three animals died, one abortion, four weight loss, four vomiting, four diarrhea.

Mutagenicity:

Ramipril was not mutagenic in the Ames microbial mutagen test, the HGPRT test in V79 cells, the micronucleus test in mice and the UDS test in human A549 cells.

Carcinogenicity:

There was no evidence of a carcinogenic effect when ramipril was administered for 104 weeks to NMRI mice at doses up to 1000 mg/kg/day and to Wistar rats at doses up to 500 mg/kg/day.

REFERENCES

Benetos A, Vasmant D, Thiéry P, et al. Effects of Ramipril on Arterial Hemodynamics. *J of Cardiovascular Pharmacology* 1991, 18(Suppl 2): S153-S156.

Burris JF. The Effect of Ramipril on Ambulatory Blood Pressure: A Multicenter Trial. *J of Cardiovascular Pharmacology* 1991, 18(Suppl 2): S131-S133.

Carré A, Vasmant D, Elmalem J, et al. Tolerability of Ramipril in a Multicenter Study of Mild-to-Moderate Hypertension in General Practice. *J of Cardiovascular Pharmacology* 1991, 18(Suppl 2): S141-S143.

Hall AS, Winter C, Bogle SM, Mackintosh AF, Murray GD, Ball SG, on behalf of the AIRE Study Investigators: The Acute Ramipril Efficacy (AIRE) study: rationale, design, organization and outcome definitions. *J Cardiovasc Pharmacol* 1991, 18(Suppl.2): S105-S109.

Heidbreder K, Froer K-L, Bauer B et al. Efficacy and Safety of Ramipril in Combination with Hydrochlorothiazide: Results of a Long-Term Study. *J of Cardiovascular Pharmacology* 1991, 18(Suppl 2): S169-S173.

Hosie J and Meredith P. The Pharmacokinetics of Ramipril in a Group of Ten Elderly Patients with Essential Hypertension. *J of Cardiovascular Pharmacology* 1991, 18(Suppl 2): S125-S127.

Lenox-Smith AJ, Street RB and Kendall FD. Comparison of Ramipril Against Atenolol in Controlling Mild-to-Moderate Hypertension. *J of Cardiovascular Pharmacology* 1991, 18(Suppl.2): S150-S152.

Manhem PJO, Ball SG, Morton JJ, Murray GD, Leckie BJ, Fraser R, Robertson JIS. A dose-response study of Hoe 498, a new non-sulphydryl converting enzyme inhibitor, on blood pressure, pulse rate and the renin-angiotensin-aldosterone system in normal man. *Br J Clin. Pharmacol* 1985, 20: 27-35.

McCarron D and The Ramipril Multicenter Study Group. 24-Hour Blood Pressure Profiles in Hypertensive Patients Administered Ramipril or Placebo Once Daily: Magnitude and Duration of Antihypertensive Effects. *Clin Cardiol* 1991, 14: 737-742.

Mills TP. Ramipril: A review of the new ACE inhibitor. *J of the Arkansas Medical Society*, February 1992, 88(9): 437-440.

Reinich W, Hoffmann H, Hoffmann W. Treatment of hypertension with the new ACE-inhibitor Ramipril. (Translation) *Therapiewoche Österreich* 1992, 7: 112-119.

Rosenthal J, Buehler G, Koenig W, et al. Effect of Angiotensin-Converting Enzyme Inhibition on Human Tissue Renin. *J of Cardiovascular Pharmacology* 1991, 18(Suppl 2): S122-S124.

Saalbach R, Wochnik G, Mauersberger H, et al. Antihypertensive Efficacy, Tolerance, and Safety of Ramipril in Young vs. Old Patients: A Retrospective Study. *J of Cardiovascular Pharmacology* 1991, 18(Suppl 2): S134-S136.

Schnaper HW. Dose-Response Relationship of Ramipril in Patients with Mild-to-Moderate Hypertension. *J of Cardiovascular Pharmacology* 1991, 18(Suppl. 2): S128-S130.

Schreiner M, Berendes B, Verho M, et al. Antihypertensive Efficacy, Tolerance, and Safety of Long-Term Treatment with Ramipril in Patients with Mild-to-Moderate Essential Hypertension. *J of Cardiovascular Pharmacology* 1991, 18(Suppl 2): S137-S140.

Vasmant D, Lendresse P, Lemarie J-C, et al. Comparison of Response Rates to the Angiotensin-Converting Enzyme Inhibitor Ramipril in Mild-to-Moderate Hypertension in a Double-Blind, Parallel-Group Study and an Open Single-Blind Study. *J of Cardiovascular Pharmacology* 1991, 18(Suppl 2): S144-S146.

Vierhapper H, Witte U, Waldhausl W. Unchanged pressor effect of norepinephrine in normal man following the oral administration of two angiotensin converting enzyme inhibitors, captopril and Hoe 498. *J Hypertens* 1986, 4: 9-11.

The Acute Ramipril Efficacy (AIRE) Study Investigators: Effect of ramipril on mortality and morbidity of survivors of acute myocardial infarction with clinical evidence of heart failure. *Lancet* 1993, 342: 821-828.

The Heart Outcomes Prevention Evaluation Study Investigators: Effects of an Angiotensin-Converting-Enzyme Inhibitor, Ramipril, on Cardiovascular Events in High-risk Patients. *N Engl J Med* 2000; 342:145-153.