

PRODUCT MONOGRAPH

^{Pr} **NOVO-RIVASTIGMINE**
(Rivastigmine as Rivastigmine Hydrogen Tartrate)

Capsules, 1.5 mg, 3 mg, 4.5 mg, 6 mg

Cholinesterase Inhibitor

Novopharm Limited
30 Novopharm Court
Toronto, Ontario
M1B 2K9

Date of Preparation:
April 1, 2009

Control Numbers: 127374, 108142

Table of Contents

PART I: HEALTH PROFESSIONAL INFORMATION.....	3
SUMMARY PRODUCT INFORMATION	3
INDICATIONS AND CLINICAL USE.....	3
CONTRAINDICATIONS	4
WARNINGS AND PRECAUTIONS.....	4
ADVERSE REACTIONS.....	9
DRUG INTERACTIONS	20
DOSAGE AND ADMINISTRATION	21
OVERDOSAGE	22
ACTION AND CLINICAL PHARMACOLOGY	23
STORAGE AND STABILITY.....	25
DOSAGE FORMS, COMPOSITION AND PACKAGING	25
PART II: SCIENTIFIC INFORMATION	27
PHARMACEUTICAL INFORMATION.....	27
CLINICAL TRIALS.....	28
DETAILED PHARMACOLOGY	38
TOXICOLOGY	40
PART III: CONSUMER INFORMATION.....	46

Pr NOVO-RIVASTIGMINE

(Rivastigmine as Rivastigmine Hydrogen Tartrate)

PART I: HEALTH PROFESSIONAL INFORMATION

SUMMARY PRODUCT INFORMATION

Route of Administration	Dosage Form/Strength	Clinically Relevant Nonmedicinal Ingredients
Oral	Capsules, 1.5 mg, 3 mg, 4.5 mg, 6 mg	Not applicable <i>For a complete listing see Dosage Forms, Composition and Packaging section.</i>

INDICATIONS AND CLINICAL USE

NOVO-RIVASTIGMINE (rivastigmine hydrogen tartrate) is indicated for the symptomatic treatment of patients with mild to moderate dementia of the Alzheimer's type.

NOVO-RIVASTIGMINE is indicated for the symptomatic treatment of patients with idiopathic Parkinson's disease, and mild to moderate dementia, with onset at least 2 years after the initial diagnosis of Parkinson's disease, and in whom other causes of dementia have been ruled out.

Rivastigmine hydrogen tartrate has not been studied in controlled clinical trials for longer than 6 months.

NOVO-RIVASTIGMINE *capsules should only be prescribed by (or following consultation with), clinicians who are experienced in the diagnosis and management of dementia.*

Geriatrics (>85 years of age): see **WARNINGS AND PRECAUTIONS**

Pediatrics: No data is available. See **WARNINGS AND PRECAUTIONS**

CONTRAINDICATIONS

- Patients with known hypersensitivity to rivastigmine, other carbamate derivatives or other components of the formulation. For a complete listing, see the Dosage Forms, Composition and Packaging section of the product monograph.
- Patients with severe liver impairment since it has not been studied in this population.

WARNINGS AND PRECAUTIONS

General

As with other cholinergic substances care must be taken when prescribing Novo-Rivastigmine Hydrogen Tartrate.

- To patients with sick sinus syndrome or conduction defects (sino-atrial block, atrio-ventricular block) (see Cardiovascular).
- To patients with active gastric or duodenal ulcers or patients predisposed to these conditions because gastric acid secretions may be increased (see Gastrointestinal).
- To patients predisposed to urinary obstruction and seizures because cholinomimetics may induce or exacerbate these diseases (see Neurologic).
- To patients with a history of asthma or obstructive pulmonary disease (see Respiratory).

Anesthesia: Rivastigmine hydrogen tartrate as a cholinesterase inhibitor, is likely to exaggerate succinylcholine-type muscle relaxation during anesthesia.

Weight Loss: Cholinesterase inhibitors as well as Alzheimer's disease and dementia associated with Parkinson's disease can be associated with significant weight loss.

In controlled clinical trials of Alzheimer's disease patients, the use of rivastigmine hydrogen tartrate was associated with weight loss. Women exposed to doses of rivastigmine hydrogen tartrate at the higher end of the therapeutic range, i.e. usual maintenance dose range of 6-12 mg/day were at greater risk for weight loss. Approximately 24% of women on 6-12 mg/day doses of rivastigmine hydrogen tartrate had weight loss of equal to or greater than 7% of their baseline weight compared to 6% on placebo. For males, 16% (6-12 mg/day) experienced a similar degree of weight loss compared to 4% on placebo.

In dementia associated with Parkinson's disease, in a single trial of n = 541 patients, at a similar dose range as for Alzheimer's disease patients 16.3% of patients had weight loss equal to or greater than 7% of their baseline weight compared to 14% in the placebo group (21.1 % drug vs. 8.1 % placebo for women, and 13.7 % drug vs. 17.1 % placebo for men). The rates in the drug-treated group are similar to those seen in Alzheimer's disease trial (as above), although the relativities with placebo are not.

Where weight loss may be of clinical concern, body weight should be monitored.

Cardiovascular

Because of their pharmacological action, cholinomimetics may have vagotonic effects on heart rate (e.g. bradycardia). The potential for this action may be particularly important to patients with “sick sinus syndrome” or other supraventricular cardiac conduction conditions. In clinical trials patients with serious cardiovascular disease were excluded. Caution should therefore be exercised in treating patients with active coronary artery disease or congestive heart failure. Syncopal episodes have been reported in association with the use of rivastigmine hydrogen tartrate. It is recommended that rivastigmine hydrogen tartrate not be used in patients with cardiac conduction abnormalities (except for right bundle branch block) including “sick sinus syndrome” and those with unexplained syncopal episodes.

Worsening of Tremor and other Extrapyrarnidal Symptoms

Like other cholinomimetics, rivastigmine hydrogen tartrate may exacerbate or induce extrapyramidal symptoms. Worsening of these symptoms (including bradykinesia, dyskinesia, gait abnormality) and an increased incidence of, or severity of, tremor have been observed in patients with dementia associated with Parkinson’s disease treated with rivastigmine hydrogen tartrate. Particularly in the case of tremor, events were observed shortly after dose increase, and may respond to dose reduction (see also ADVERSE EVENTS, Dementia associated with Parkinson’s Disease, Extrapyrarnidal Symptoms; and DOSAGE AND ADMINISTRATION, Dosing Considerations).

Clinical monitoring is recommended for these adverse events.

Gastrointestinal

Rivastigmine hydrogen tartrate is associated with significant gastrointestinal adverse reactions including nausea, vomiting, anorexia and weight loss.

Treatment with rivastigmine hydrogen tartrate should always be started at a dose of 1.5 mg b.i.d. or 1.5 mg o.d., as clinically indicated, and patients titrated to their maintenance dose. **If treatment with rivastigmine hydrogen tartrate is interrupted for longer than several days, patients should be instructed to reinitiate treatment with the lowest daily dose and be retitrated (see DOSAGE AND ADMINISTRATION) to reduce the possibility of severe vomiting and its potentially serious sequelae (e.g. very rare post-marketing reports of severe vomiting with esophageal rupture).**

Nausea, vomiting and diarrhea appear more frequently at higher doses (see ADVERSE REACTIONS), with nausea and vomiting being more prevalent in women. Females are more sensitive to the cholinergic adverse effects associated with cholinesterase inhibitors and in

general are more likely to experience nausea and vomiting than are males. In most cases these effects were of mild to moderate intensity and transient, and they resolved during continued rivastigmine treatment or upon treatment discontinuation.

Through their primary action, cholinesterase inhibitors may be expected to increase gastric acid secretion due to increased cholinergic activity. Therefore, patients should be monitored for symptoms of active or occult gastrointestinal bleeding, especially those at increased risk for developing ulcers, e.g., those with a history of ulcer disease or those receiving concurrent nonsteroidal anti-inflammatory drugs (NSAIDs). In controlled clinical studies with rivastigmine hydrogen tartrate in Alzheimer's disease patients, patients with a past history (last 2 years) of peptic ulceration and chronic diseases of the gastrointestinal tract were excluded. In the trial population who received rivastigmine hydrogen tartrate there was no significant increase, relative to placebo, in the incidence of peptic ulcer disease. The incidence of GI hemorrhage, in controlled clinical trials was <1% (n = 6/1923) for rivastigmine hydrogen tartrate and 0% (n=0/868) for placebo.

Genetic Polymorphism

The effect of genetic polymorphism of butyrylcholinesterase enzyme on rivastigmine metabolism is unknown.

Genitourinary

Although not reported in clinical trials of rivastigmine hydrogen tartrate, cholinomimetics may cause bladder spasm.

Hepatic/Biliary/Pancreatic

There is limited information on the pharmacokinetics of rivastigmine hydrogen tartrate in hepatically impaired patients (see Pharmacokinetics section). It is therefore recommended that dose escalation with rivastigmine in hepatically impaired patients be undertaken with caution and under conditions of close monitoring for adverse effects (**see DOSAGE AND ADMINISTRATION, Dosing Considerations**). Rivastigmine hydrogen tartrate is contraindicated in patients with severe liver impairment since it has not been studied in this population (**see CONTRAINDICATIONS**).

Laboratory Values

Elevations in Serum Amylase/Lipase levels in Patients with Dementia Associated with Parkinson's Disease:

Of patients with normal levels at baseline, 17% receiving rivastigmine hydrogen tartrate and 10% receiving placebo showed amylase levels beyond the normal range at the second measurement at the end of the Study (Week 24), and 9% and 4% respectively showed lipase levels beyond the normal range. Elevations beyond 2x normal range occurred in 2 of the rivastigmine hydrogen tartrate patients with respect to amylase levels, 7 with respect to lipase

levels, and in 0 placebo patients. Elevations in both amylase and lipase levels occurred in 12 rivastigmine hydrogen tartrate patients, and in 0 placebo patients; pancreatitis was not recorded as an AE for any patient in the study.

Neurologic

Seizures: In placebo controlled clinical trials with rivastigmine hydrogen tartrate cases of seizures were reported. Cholinomimetics are believed to have some potential to cause generalized convulsions. However, seizure activity also may be a manifestation of Alzheimer's Disease. The risk/benefit of rivastigmine hydrogen tartrate treatment for patients with a history of seizure disorder must therefore be carefully evaluated.

Rivastigmine hydrogen tartrate has not been studied in patients with moderately severe or severe Alzheimer's disease, moderately severe or severe dementia associated with Parkinson's disease, or other dementias. The efficacy and safety of rivastigmine hydrogen tartrate in these patient populations is unknown.

Renal

There is limited information on the pharmacokinetics of rivastigmine hydrogen tartrate in renally impaired patients (see Pharmacokinetics section). It is therefore recommended that dose escalation with rivastigmine in renally impaired patients be undertaken with caution and under conditions of close monitoring for adverse effects (see **DOSAGE AND ADMINISTRATION, Dosing Considerations**).

Respiratory

Like other cholinomimetic drugs, rivastigmine hydrogen tartrate should be used with care in patients with a history of asthma or obstructive pulmonary disease. No experience is available in treating patients with these conditions.

Special Populations

Pregnant Women:

The safety of rivastigmine hydrogen tartrate in pregnant women has not been established. Novo-Rivastigmine should not be used in women of childbearing potential unless, in the opinion of the physician, the potential benefit to the patient justifies the potential risk to the fetus.

Nursing Women:

It is not known whether rivastigmine hydrogen tartrate is excreted into human milk, and therefore rivastigmine hydrogen tartrate should not be used in nursing mothers.

Pediatrics:

The safety and effectiveness of rivastigmine hydrogen tartrate in any illness occurring in pediatric patients have not been established.

Geriatrics:

Use in patients > 85 years old: In controlled clinical studies in Alzheimer's disease patients, the number of patients over 85 years old who received rivastigmine hydrogen tartrate in the therapeutic dose range of 6-12 mg/day was 68. Of these patients, 12 received high doses of rivastigmine hydrogen tartrate (>9-12 mg/day). The safety of rivastigmine hydrogen tartrate in this patient population has not been adequately characterized. In Alzheimer's disease patients in controlled clinical trials, nausea, diarrhea, vomiting, dizziness, anorexia, fatigue, dyspepsia and weakness increased with dose. Dose escalation in patients >85 years old should thus proceed with caution (see **DOSAGE AND ADMINISTRATION, Dosing Considerations**).

Use in elderly patients with serious comorbid disease: There is limited information on the safety of rivastigmine hydrogen tartrate treatment in patients with mild to moderate Alzheimer's disease, or mild to moderate dementia associated with Parkinson's disease, and serious comorbidity. The use of rivastigmine hydrogen tartrate in Alzheimer's disease patients, or in patient with dementia associated with Parkinson's disease, with chronic illnesses common among the geriatric population, should be considered only after careful risk/benefit assessment and include close monitoring for adverse events. Dose escalation in this patient population should proceed with caution (see **DOSAGE AND ADMINISTRATION, Dosing Considerations**).

Patients with vascular dementia

Patients diagnosed with probable vascular dementia, according to NINDS-AIREN criteria, were randomized to double-blind treatment with rivastigmine hydrogen tartrate (3-12 mg/day, N=363) or placebo (N=344) for 6 months in a controlled clinical trial. The NINDS-AIREN criteria are designed to identify patients with dementia that appears to be due primarily to vascular causes, and to exclude patients with Alzheimer's disease. Overall, rivastigmine hydrogen tartrate was not shown to be an effective treatment for patients with vascular dementia in this study.

The study also showed that the overall rate of occurrence of treatment emergent adverse events was lower in vascular dementia patients than what was observed previously in Alzheimer's disease patients. However, rates of serious adverse events were generally greater for vascular dementia patients compared to mild to moderate Alzheimer's disease patients for both rivastigmine hydrogen tartrate and placebo groups, and may relate to the greater number of comorbid medical conditions in the vascular dementia population.

In vascular dementia patients, higher rates of all-cause mortality (2.2% on rivastigmine hydrogen tartrate vs. 1.2% on placebo) and certain cardiovascular and cerebrovascular adverse events such as, angina pectoris, myocardial infarction, coronary artery disease, hypertension, dysarthria and cerebrovascular accident were observed in patients who were treated with rivastigmine hydrogen tartrate compared to those who received placebo. The majority of deaths in patients taking either

rivastigmine hydrogen tartrate or placebo resulted from either cardiovascular or cerebrovascular disorders or respiratory failures.

ADVERSE REACTIONS

Adverse Drug Reaction Overview

Clinical Trial Adverse Drug Reactions

Because clinical trials are conducted under very specific conditions the adverse reaction rates observed in the clinical trials may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse drug reaction information from clinical trials is useful for identifying drug-related adverse events and for approximating rates.

Dementia of the Alzheimer’s Type

A total of 1923 patients with mild to moderate Alzheimer’s Disease were treated in controlled clinical studies with rivastigmine hydrogen tartrate. Of these patients, 1417 (74%) completed the studies. The mean duration of treatment for all rivastigmine groups was 154 days (range 1-255 days).

Adverse Events Leading to Discontinuation

Overall, 18% (340/1923) of patients treated with rivastigmine hydrogen tartrate discontinued from Phase III controlled clinical trials due to adverse events compared to 9% (75/868) in the placebo group. During the titration phases of controlled clinical trials the incidence of discontinuations due to adverse events was 5% for placebo, 5% for rivastigmine hydrogen tartrate 1-4 mg/day and 21% for rivastigmine hydrogen tartrate 6-12 mg/day. During the maintenance phases, 3% of patients who received placebo, 3% of patients who received 1-4 mg/day rivastigmine hydrogen tartrate and 6% of patients who received rivastigmine hydrogen tartrate 6-12 mg/day withdrew from studies due to adverse events.

Female patients treated with rivastigmine hydrogen tartrate were approximately twice as likely to discontinue study participation due to adverse events than were male patients [Females 21%; Males: 12%]. Female patients in clinical studies were found to be more susceptible to gastrointestinal adverse drug reactions and weight loss.

The most common adverse events leading to discontinuation, defined as those occurring in at least 2% of patients and at twice the incidence seen in placebo patients, are shown in Table 1.

Table 1. Most frequent adverse events ($\geq 2\%$ and twice the rate in the placebo group) leading to withdrawal from randomized placebo controlled clinical trials B351, B352, and B303 during titration and maintenance phases[†]

	Titration phase (Weeks 1-12)			Maintenance Phase (Weeks 13-26)		
	Placebo n=646	1-4 mg/day n=644	6-12 mg/day n=824	Placebo n=588	1-4 mg/day n=587	6-12 mg/day n=601

	Titration phase (Weeks 1-12)			Maintenance Phase (Weeks 13-26)		
	Placebo n=646	1-4 mg/day n=644	6-12 mg/day n=824	Placebo n=588	1-4 mg/day n=587	6-12 mg/day n=601
All events	5%	5%	21%	3%	3%	6%
Nausea	1%	1%	10%	0%	<1%	1%
Vomiting	0%	<1%	5%	0%	<1%	2%
Anorexia	0%	<1%	3%	<1%	<1%	<1%
Dizziness	<1%	<1%	3%	<1%	0%	1%
Abdominal pain	<1%	<1%	2%	<1%	<1%	<1%
Asthenia	0%	0%	2%	0%	0%	<1%
Fatigue	<1%	<1%	2%	0%	0%	<1%

† All patients who received at least one dose of study medication were included in the results for the titration phase. All patients who entered the maintenance phase were represented in the results for the maintenance phase.

Titration and maintenance dosing should remain flexible and be adjusted according to individual needs.

Most Frequent Adverse Clinical Events Seen in Association with the Use of Rivastigmine Hydrogen Tartrate

The most common adverse events, defined as those occurring at a frequency of at least 5% and twice the placebo rate, are largely predicted by rivastigmine hydrogen tartrate's cholinomimetic effects. These include nausea, vomiting, dizziness, diarrhea, anorexia, weight loss of > 7% of baseline weight, and abdominal pain.

Table 2 presents a comparison of common adverse events ($\geq 5\%$ incidence and twice the placebo rate) by treatment group during titration (Weeks 1-12) and maintenance (Weeks 13-26). The adverse events were generally mild in intensity, more frequent at higher doses, of short duration, and attenuated with continued dosing or discontinuation of drug.

Table 2. Common adverse events ($\geq 5\%$ and twice the rate in the placebo group) in randomized placebo controlled clinical trials B351, B352, and B303 during titration and maintenance phases[†]

Adverse event	Titration phase (Weeks 1-12)			Maintenance phase (Weeks 13-26)		
	Placebo n=646	1-4 mg/day n=644	6-12 mg/day n=824	Placebo n=588	1-4 mg/day n=587	6-12 mg/day n=601
Nausea	9%	15%	40%	4%	8%	15%
Vomiting	3%	5%	23%	3%	5%	14%
Dizziness	10%	10%	19%	4%	6%	10%
Diarrhea	9%	8%	16%	4%	5%	9%
Anorexia	2%	5%	13%	1%	2%	4%
Abdominal pain	4%	5%	10%	3%	3%	4%
Fatigue	4%	4%	8%	1%	2%	3%
Asthenia	2%	1%	6%	1%	2%	3%
Somnolence	2%	4%	5%	1%	1%	1%

† All patients who received at least one dose of study medication were included in the results for the titration phase. All patients who entered the maintenance phase were represented in the results for the maintenance phase.

Titration and maintenance dosing should remain flexible and be adjusted according to individual needs.

In an open label study involving 305 patients with Alzheimer’s disease the tolerability of a 1.5 mg b.i.d. (3 mg/day) starting dose and dose escalation of 1.5 mg b.i.d. (3 mg/day) at a minimum interval of every two weeks were assessed. A total of 40 of these patients (13%) discontinued the study due to adverse events. The type and incidence of common adverse events reported did not appear to differ substantially from those noted in placebo-controlled studies.

Adverse Events Reported in Controlled Trials

The events cited reflect experience gained under closely monitored condition of clinical trials in a highly selected patient population. In actual clinical practice or in other clinical trials, these frequency estimates may not apply, as the conditions of use, reporting behavior, and the kinds of patients treated may differ. Table 3 lists treatment emergent signs and symptoms that were reported in at least 2% of patients in Phase 3 placebo-controlled trials for which the rate of occurrence was greater for rivastigmine hydrogen tartrate assigned than placebo assigned patients. There were too few non Caucasian patients enrolled to assess the effect of race on the incidence of adverse events in the Phase III controlled studies. Similarly, there were too few patients aged more than 85 years to systematically assess the effect of advanced age. Female patients were more susceptible to nausea, vomiting, loss of appetite and weight loss.

Table 3. Adverse Events Reported in Controlled Clinical Trials in at Least 2% of Patients Receiving Rivastigmine Hydrogen Tartrate and at a Higher Frequency than Placebo-treated Patients

Body System/Adverse Event	Placebo (N=868)	RIVASTIGMINE (N=1923)
Percent of Patients with any Adverse Event	79	87
Autonomic Nervous System		
Sweating increased	1	3
Body as a Whole		
Fatigue	5	7
Asthenia	2	5
Malaise	2	4
Weight Decrease	<1	2
Cardiovascular Disorders, General		
Hypertension	2	3
Central and Peripheral Nervous System		
Dizziness	11	19
Headache	12	15
Somnolence	3	5
Tremor	1	3
Gastrointestinal System		
Nausea	12	37
Vomiting	6	23
Diarrhea	11	16
Anorexia	3	13
Abdominal Pain	6	11
Dyspepsia	4	8
Constipation	4	5

Body System/Adverse Event	Placebo (N=868)	RIVASTIGMINE (N=1923)
Percent of Patients with any Adverse Event	79	87
Flatulence	2	4
Eructation	1	2
Psychiatric Disorders		
Insomnia	7	8
Depression	4	5
Anxiety	3	4
Hallucination	3	4
Nervousness	3	4
Aggressive Reaction	2	3
Respiratory System		
Rhinitis	3	4
Dyspnea	1	2
Skin and Appendages		
Pruritus	1	2
Urinary System		
Urinary Incontinence	2	3
Micturition Frequency	1	2
Vision Disorders		
Vision Abnormal	1	2

Other Adverse Events Observed During Clinical Trials

Rivastigmine hydrogen tartrate has been administered to over 5297 individuals during clinical trials worldwide. Of these, 4326 patients have been treated for at least 3 months, 3407 patients have been treated for at least 6 months, 2150 patients have been treated for 1 year, 1250 have been treated for 2 years, and 168 have been treated for over 3 years. With regard to exposure to the highest dose, 1679 patients were exposed to mean daily doses of 10-12 mg, 1659 patients treated for 3 months, 1504 patients treated for 6 months, 885 patients treated for 1 year, 629 patients treated for 2 years, and 86 treated for over 3 years.

Treatment emergent signs and symptoms that occurred during 8 controlled clinical trials and 9 open-label trials in North America, Western Europe, Australia, South Africa and Japan were recorded as adverse events by the clinical investigators using terminology of their own choosing. To provide an overall estimate of the proportion of individuals having similar types of events, the events were grouped into a smaller number of standardized categories using a modified WHO dictionary, and event frequencies were calculated across all studies. These categories are used in the listing below. The frequencies represent the proportion of 5297 patients from these trials who experienced that event while receiving rivastigmine hydrogen tartrate. All adverse events occurring at least 6 times are included, except for those already listed in Table 3, WHO terms too general to be informative, or events less likely to be drug caused. Events are classified by body system and listed using the following definitions: frequent adverse events - those occurring in at least 1/100 patients; infrequent adverse events - those occurring in 1/100 to 1/1000 patients. These adverse events are not necessarily related to rivastigmine hydrogen tartrate treatment and in most cases were observed at a similar frequency in placebo-treated patients in the controlled studies.

Autonomic Nervous System: *Frequent:* Syncope. *Infrequent:* Cold clammy skin, dry mouth, flushing, increased saliva.

Body as a Whole: *Frequent:* Accidental trauma, allergy, chest pain, edema, fever, hot flushes, influenza-like symptoms, overdose, rigors. *Infrequent:* Allergic reaction, chest pain substernal, edema periorbital, facial edema, feeling cold, halitosis, hypothermia, inflammatory reaction unspecified, pain, pallor, tumor unspecified, unspecified eyelid disorder, weight increase.

Cardiovascular System: *Frequent:* Angina pectoris, cardiac failure, hypotension, myocardial infarction, peripheral edema, postural hypotension. *Infrequent:* Chest pain, coronary artery disorder, ECG abnormal, edema, generalized edema, heart sounds abnormal, myocardial ischemia.

Central and Peripheral Nervous System: *Frequent:* Abnormal gait, ataxia, convulsions, extrapyramidal disorder, paresthesia, vertigo. *Infrequent:* Abnormal coordination, aphasia, apraxia, cerebrovascular accident, coma, dysphonia, hyperkinesia, hyperreflexia, hypertonia, hypoesthesia, hypokinesia, hyporeflexia, involuntary muscle contractions, migraine, neuralgia, neuropathy, nystagmus, paresis, peripheral neuropathy, speech disorder.

Collagen Disorders: *Infrequent:* Rheumatoid arthritis.

Endocrine System: *Infrequent:* Goiter, hypothyroidism.

Gastrointestinal System: *Frequent:* Fecal incontinence, gastritis, tooth disorder. *Infrequent:* Colitis, colorectal polyp, diverticulitis, duodenal ulcer, dysphagia, esophagitis, gastric ulcer, gastroenteritis, gastroesophageal reflux, GI hemorrhage, gingivitis, glossitis, hematemesis, hernia, hiccup, increased appetite, intestinal obstruction, melena, pancreatitis, peptic ulcer, rectal disorder, rectal hemorrhage, tenesmus, tooth caries, ulcerative stomatitis.

Hearing and Vestibular Disorders: *Frequent:* Tinnitus. *Infrequent:* Deafness, earache, ear disorder unspecified, vestibular disorder.

Heart Rate and Rhythm Disorders: *Frequent:* Bradycardia, fibrillation atrial, palpitation. *Infrequent:* Arrhythmia, AV block, bundle branch block, cardiac arrest, extrasystoles, sick sinus syndrome, supraventricular tachycardia, tachycardia.

Injury, Poisoning and Procedural Complications: *Infrequent:* Fall.

Liver and Biliary System Disorders: *Infrequent:* Abnormal hepatic function, cholecystitis, cholelithiasis, increased gamma-glutamyl transferase, increased hepatic enzymes.

Metabolic and Nutritional Disorders: *Frequent:* Dehydration, hypokalemia. *Infrequent:* Cachexia, diabetes mellitus, gout, hypercholesterolemia, hyperglycemia, hyperlipemia, hypoglycemia, hyponatremia, thirst.

Musculoskeletal Disorders: *Frequent:* Arthralgia, arthritis, back pain, bone fracture, leg cramps, leg pain, myalgia, pain. *Infrequent:* Arthropathy, arthrosis, bone disorder, bone pain, bursitis, cramps, hernia, joint malformation, muscle weakness, osteoporosis, spine malformation, stiffness, tendinitis, tendon disorder, vertebral disc disorder.

Neoplasms: *Frequent:* Basal cell carcinoma. *Infrequent:* Bladder carcinoma, carcinoma, colon carcinoma, malignant breast neoplasm (female), malignant skin neoplasm, unspecified adenocarcinoma, unspecified neoplasm.

Platelet, Bleeding, and Clotting Disorders: *Frequent:* Epistaxis. *Infrequent:* Hematoma, purpura, thrombocytopenia, unspecified hemorrhage.

Psychiatric Disorders: *Frequent:* Agitation, behavioral disturbance, confusion, delusion, paranoid reaction, paroniria. *Infrequent:* Abnormal dreaming, amnesia, apathy, decreased libido, delirium, dementia, depersonalization, emotional lability, impaired concentration, increased libido, neurosis, psychosis, sleep disorder, stress reaction, suicidal ideation.

Red Blood Cell Disorders: *Frequent:* Anemia. *Infrequent:* Anemia B₁₂ deficiency, hypochromic anemia.

Reproductive Disorders (Female & Male): *Frequent:* Prostatic disorder. *Infrequent:* Atrophic vaginitis, breast pain (female), impotence, intermenstrual bleeding, unspecified uterine disorder, vaginal hemorrhage, vaginitis.

Resistance Mechanism Disorders: *Frequent:* Infection, pneumonia, upper respiratory tract infection, urinary tract infection, viral infection. *Infrequent:* Bacterial infection, cellulitis, cystitis, fungal infection, herpes simplex, herpes zoster, moniliasis, onychomycosis, otitis media, parasitic infection, sepsis.

Respiratory System: *Frequent:* Bronchitis, coughing, pharyngitis, sinusitis. *Infrequent:* Abnormal chest sounds, apnea, bronchospasm, emphysema, hyperventilation, increased sputum, laryngitis, pleural effusion, pulmonary disorder, pulmonary edema, respiratory disorder, respiratory insufficiency.

Skin and Appendages: *Frequent:* Rash, skin disorder, skin ulceration. *Infrequent:* Abscess, acne, alopecia, bullous eruption, contact dermatitis, dermatitis, dry skin, eczema, erythematous rash, furunculosis, genital pruritus, hyperkeratosis, maculo-papular rash, nail disorder, otitis externa, psoriaform rash, seborrhea, skin cyst, skin discoloration, skin exfoliation, skin hypertrophy, sunburn, urticaria, verruca.

Special Senses: *Infrequent:* Loss of taste, perversion of taste.

Urinary System Disorders: *Frequent:* Hematuria. *Infrequent:* Acute renal failure, albuminuria, dysuria, micturition disorder, micturition urgency, nocturia, polyuria, pyuria, renal calculus, renal cyst, renal function abnormal, unspecified bladder disorder, urethral disorder, urinary retention.

Vascular (extracardiac) Disorders: *Frequent:* Cerebrovascular disorder. *Infrequent:* Aneurysm, circulatory disorder, hemorrhoids, intracranial hemorrhage, peripheral ischemia, phlebitis, pulmonary embolism, thrombophlebitis deep, thrombosis, varicose vein, vascular disorder.

Vision Disorders: *Frequent:* Cataract, conjunctivitis. *Infrequent:* Abnormal lacrimation, blepharitis, conjunctival hemorrhage, diplopia, eye abnormality, eye pain, glaucoma.

White Cell and Resistance Disorders: *Infrequent:* Leukocytosis, lymphadenopathy.

Dementia Associated with Parkinson's Disease

In the 24 week, double-blind, placebo-controlled trial, n= 541 patients were randomized to drug or placebo (2:1 ratio). Of these, 73% of patients in the drug arm completed the study (i.e. did not discontinue from drug treatment), and 82% in the placebo arm. The mean duration of treatment for rivastigmine-treated patients was 144 days (range 4-197 days).

The overall AE profile of rivastigmine in this study was consistent with the known profile of patients with Alzheimer's disease, with the exception that frequency of tremor and worsening of Parkinson's disease symptoms in general, is greater compared to placebo. A number of factors beyond that of differing patient populations may affect comparison of AE rates between Alzheimer's disease and Parkinson's disease, including the protocol-specified differences in dosing between the Alzheimer's disease studies, and the sole Parkinson's disease study: a) greater time between dose escalations for the Parkinson's disease patients (4 weeks for Parkinson's disease patients vs 1-2 weeks for Alzheimer's disease patients) and b) lower doses specified for some Alzheimer's disease patients (minimum of 3 mg/day for the Parkinson's disease patients, vs minimum of 1 mg/day for some Alzheimer's disease patients).

Adverse Events leading to discontinuation

The rate of discontinuation due to adverse events in the single controlled trial of rivastigmine was 18.2% for patients receiving 3-12 mg/day compared to 11.2% for patients on placebo during the 24 week study. The most frequent adverse events that led to discontinuation from this study, defined as those occurring in at least 1% of patients receiving rivastigmine and more frequent than those receiving placebo, were nausea (3.6% rivastigmine vs. 0.6% placebo), vomiting (1.9% rivastigmine vs 0.6% placebo), and tremor (1.7% rivastigmine vs. 0.0% placebo).

Most Frequent Adverse Clinical Events

The most common adverse events, defined as those occurring at a frequency of at least 5% and twice the placebo rate, are largely predicted by rivastigmine's cholinergic effects. These include nausea, vomiting, tremor, anorexia, and dizziness.

Table 4 presents a comparison of common adverse events ($\geq 5\%$ incidence and twice the placebo rate) by treatment group during titration (Weeks 1-16) and maintenance (Weeks 17-24).

Table 4: Common adverse events ($\geq 5\%$ and twice the rate in the placebo group) during the single controlled clinical trial, breakdown by titration and maintenance phases

Adverse event	Titration phase (Weeks 1-16)		Maintenance phase (Weeks 14-24)	
	Placebo n=179	Rivastigmine n=362	Placebo n=158	Rivastigmine n=281
Nausea	11%	27%	1%	5%
Vomiting	2%	14%	0%	4%
Tremor	3%	9%	1%	1%
Dizziness	1%	6%	0%	<1%
Anorexia	2%	5%	1%	2%

† All patients who received at least one dose of study medication were included in the results for the titration phase. All patients who entered the maintenance phase were represented in the results for the maintenance phase.

Adverse Events Reported in the Controlled Trial

The events cited reflect experience gained under the closely monitored condition of a clinical trial, in a highly selected patient population. In actual clinical practice or in other clinical trials, these frequency estimates may not apply, as the conditions of use, reporting behavior, and the kinds of patients treated may differ. Table 5 lists treatment emergent signs and symptoms that were reported in at least 2% of patients in the rivastigmine arm (doses of 3-12 mg/day) of placebo-controlled trials and for which the rate of occurrence was greater for patients treated with rivastigmine than for those treated with placebo. There were too few non-Caucasian patients enrolled to assess the effect of race on the incidence of adverse events in the study. Similarly, there were too few patients aged more than 85 years to systematically assess the effect of advanced age. Female patients were more susceptible to nausea, vomiting, loss of appetite and weight loss.

In general, adverse reactions were less frequent later in the maintenance phase of the treatment.

Table 5: Adverse Events Reported in the Single Controlled Clinical Trial in at Least 2% of Patients Receiving Rivastigmine (3-12 mg/day) and at a Higher Frequency than Placebo-treated Patients

Body System/Adverse Event	Placebo (n=179)	Rivastigmine (n=362)
Percent of Patients with any Adverse Event	71	84
Gastrointestinal Disorders		
Nausea	11	29
Vomiting	2	17
Diarrhea	4	7
Upper Abdominal Pain	1	4
General Disorders and Administrative Site Conditions		
Fatigue	3	4
Asthenia	1	2
Metabolism and Nutritional Disorders		
Anorexia	3	6
Dehydration	1	2
Nervous System Disorders		
Tremor	4	10

Dizziness	1	6
Headache	3	4
Somnolence	3	4
Parkinson's Disease (worsening)	1	3
Bradykinesia	2	2
Parkinsonism	1	2
Psychiatric Disorders		
Anxiety	1	4
Insomnia	2	3
Restlessness	2	2
Skin and Subcutaneous Tissue Disorders		
Hyperhidrosis	1	2

Extrapyramidal symptoms

Like other cholinomimetics, rivastigmine may exacerbate or induce extrapyramidal symptoms. Worsening of these symptoms (including bradykinesia, dyskinesia, gait abnormality) and an increased incidence of, or severity of, tremor have been observed in patients with dementia associated with Parkinson's disease treated with rivastigmine. These events led to discontinuation of rivastigmine in some cases (eg discontinuations due to tremor 1.7% on rivastigmine vs 0% on placebo).

Table 6 lists the number and percentage of patients, from the single controlled study, who experienced pre-defined adverse events that may reflect worsening of Parkinson symptoms. Percentages are listed for those events for which rivastigmine was numerically higher than placebo; the remaining pre-defined events are listed at the end.

Table 6: Pre-defined adverse events that may reflect worsening of parkinsonian symptoms in patients

	Rivastigmine n (%)	Placebo n (%)
Total patients studied	362 (100)	179 (100)
Total patients with pre-defined AE(s)	99 (27.3)	28 (15.6)
Tremor	37 (10.2)	7 (3.9)
Worsening of PD/parkinsonism	20 (5.5)	3 (1.7)
Bradykinesia	9 (2.5)	3 (1.7)
Salivary hypersecretion	5 (1.4)	0
Dyskinesia	5 (1.4)	1 (0.6)
Gait abnormality	5 (1.4)	0
Dystonia	3 (0.8)	1 (0.6)
Musculoskeletal stiffness	3 (0.8)	0
Extrapyramidal Disorder	1 (0.3)	0
Hypokinesia	1 (0.3)	0
Movement disorder	1 (0.3)	0
Muscle rigidity	1 (0.3)	0

Rigors	1 (0.3)	0
Motor dysfunction	1 (0.3)	0

† Pre-defined events that were observed in the rivastigmine group, but not at a higher rate than for the placebo group, include fall, balance disorders, drooling, on and off phenomena, freezing phenomenon, hypertonia and dysarthria.

Of the reported tremor cases, approx 90% represent one episode, and 47.5% occurred within a week of a dose increase

Analysis of data on extrapyramidal symptoms utilizing sub-populations of mild (Hoehn and Yahr stage 1.0 to 2.5) vs moderate to severe (Hoehn and Yahr stage 3.0 to 5.0) Parkinson’s disease showed no apparent difference between the two sub-groups, except for

- a) From Table 6 above, a greater percentage of rivastigmine -treated patients in the moderate/severe group experienced the symptoms reflective of worsening of parkinsonian symptoms (32% vs 23%); this was due primarily to the AE “fall” (10% vs 1.6%, respectively) and the AE “worsening of parkinsonian symptoms” (4% vs 0.5%). This pattern was not apparent in the patients who were treated with placebo.
- b) The percentage of patients for whom the AE of tremour was resolved during the study was less for the moderate to severe Parkinson’s disease group compared to the mild group (47% vs 62%)

Clinical monitoring is recommended for these adverse events (see also WARNING AND PRECAUTIONS, Worsening of Extrapyramidal Symptoms and DOSAGE AND ADMINISTRATION, Dosing Considerations).

Other Adverse Events Observed During Clinical Trials of Dementia Associated with Parkinson’s Disease

Rivastigmine hydrogen tartrate has been administered to 485 individuals with dementia associated with Parkinson’s disease during clinical trials worldwide. Of these, 413 patients have been treated for at least 3 months, 253 patients have been treated for at least 6 months, and 113 patients have been treated for 1 year.

Additional treatment emergent adverse events in patients with dementia associated with Parkinson’s disease, occurring in at least 1 patient (approximately 0.3%) are listed below, excluding events that are already listed above for the dementia of the Alzheimer’s type or elsewhere in labeling, WHO terms too general to be informative, relatively minor events, or events unlikely to be drug-caused. Events are classified by body system and listed using the following definitions: frequent adverse events – those occurring in at least 1/100 patients; infrequent adverse events – those occurring in 1/100 to 1/1,000 patients. These adverse events are not necessarily related to rivastigmine treatment and in most cases were observed at a similar frequency in placebo-treated patients in the controlled studies.

Cardiovascular System: *Frequent:* Chest pain. *Infrequent:* Sudden cardiac death.

Central and Peripheral Nervous System: *Frequent:* Dyskinesia, transient ischemic attack.
Infrequent: Dystonia, hemiparesis, epilepsy, restless leg syndrome.

Endocrine System: *Infrequent:* Elevated prolactin level.

Gastrointestinal System: *Frequent:* Dyspepsia. *Infrequent:* Faecaloma, dysphagia, diverticulitis, peritonitis.

Hearing and Vestibular Disorders: *Frequent:* Vertigo. *Infrequent:* Meniere's disease.

Heart Rate and Rhythm Disorders: *Infrequent:* Adam-Stokes syndrome.

Liver and Biliary System Disorders: *Infrequent:* Elevated alkaline phosphatase level, elevated gamma-glutamyltransferase level.

Musculoskeletal Disorders: *Frequent:* Back pain *Infrequent:* Muscle stiffness, myoclonus, freezing phenomenon.

Psychiatric Disorders: *Frequent:* Agitation, depression. *Infrequent:* Delusion, insomnia.

Reproductive Disorders (Female & Male): *Infrequent:* endometrial hypertrophy, mastitis, prostatic adenoma.

Respiratory System: *Frequent:* Dyspnoea. *Infrequent:* Cough.

Urinary System Disorders: *Infrequent:* Urinary incontinence, neurogenic bladder.

Vascular (extracardiac) Disorders: *Infrequent:* Vasovagal syncope, vasculitis.

Vision Disorders: *Infrequent:* Blurred vision, blepharospasm, conjunctivitis, retinopathy.

Post-Market Adverse Drug Reactions

Voluntary reports of adverse events temporally associated with rivastigmine hydrogen tartrate that have been received since market introduction that are not listed above, and that may or may not be causally related to the drug include the following.

Skin and Appendages. Stevens-Johnson syndrome.

Gastrointestinal System: Severe vomiting with esophageal rupture (see WARNINGS AND PRECAUTIONS, Gastrointestinal).

DRUG INTERACTIONS

Overview

Use with Anticholinergics: Because of their mechanism of action, cholinesterase inhibitors have the potential to interfere with the activity of anticholinergic medications.

Use with Cholinomimetics and Other Cholinesterase Inhibitors: A synergistic effect may be expected when cholinesterase inhibitors are given concurrently with succinylcholine, similar neuromuscular blocking agents or cholinergic agonists such as bethanechol.

Use with other Psychoactive Drugs: In controlled clinical trials with rivastigmine hydrogen tartrate few patients received neuroleptics, antidepressants or anticonvulsants, there is thus limited information concerning the interaction of rivastigmine hydrogen tartrate with these drugs.

Effect of Rivastigmine on the Metabolism of Other Drugs: Rivastigmine hydrogen tartrate is mainly metabolized through hydrolysis by esterases. No *in vivo* studies have investigated the effects of rivastigmine on the clearance of drugs metabolised by CYP450. Based on *in vitro* studies, no pharmacokinetic drug interactions with drugs metabolised by the following isoenzyme systems are expected: CYP1A2, CYP2D6, CYP3A4/5, CYP2E1, CYP2C9, CYP2C8, or CYP2C19.

Rivastigmine hydrogen tartrate may inhibit the butyrylcholinesterase mediated metabolism of other drugs (see **ACTIONS AND CLINICAL PHARMACOLOGY, Pharmacokinetics, Metabolism**).

Effect of Other Drugs on the Metabolism of Rivastigmine Hydrogen Tartrate: Drugs which induce or inhibit CYP450 metabolism are not expected to alter the metabolism of rivastigmine hydrogen tartrate. Formal pharmacokinetic studies to assess the potential for drug interaction with other medications commonly taken by the elderly were not done. Population-pharmacokinetic analyses of a subset (n = 359; 6-12 mg/day) of patients with Alzheimer's disease in controlled clinical trials do not suggest that the administration of rivastigmine hydrogen tartrate with some commonly prescribed medications is associated with an alteration in the kinetics of rivastigmine hydrogen tartrate, or an increased risk of clinically relevant untoward effects. However, the number of patients who received concomitant medications chronically was as follows: anilides (e.g. acetaminophen) (10%), antacids (12%), antianginals (6%), antihistamines (2%), antihypertensives (12%), benzodiazepines (<1%), β - blockers (7%), calcium channel blockers (12%), digitalis glycosides (5%), non-steroidal anti-inflammatory drugs (13%), oral hypoglycemics (3%), and salicylic acid and derivatives (28%).

Drug-Drug Interactions

Studies to assess the potential of rivastigmine hydrogen tartrate for interaction with digoxin, warfarin, diazepam or fluoxetine were limited to short term, single-dose studies in young healthy volunteers. No significant effects on the pharmacokinetics of these drugs or on the metabolism of rivastigmine hydrogen tartrate were observed. Similar studies in elderly patients were not done.

DOSAGE AND ADMINISTRATION

Dosing Considerations

- For elderly patients (>85 years old) with low body weight (especially females) or serious comorbid diseases (see **WARNINGS AND PRECAUTIONS**) it is recommended to start treatment with less frequent dosing (1.5 mg once a day) and to escalate dosage at a slower rate than for adults.
- For patients with renal or hepatic impairment (see **WARNINGS AND PRECAUTIONS**) it is recommended that treatment be started with less frequent dosing (1.5 mg once a day) and that dose escalation be slower than that recommended for adults.
- In a population of cognitively-impaired individuals, safe use of this and all other medications may require supervision.
- Adverse effects (eg. Hypertension and hallucinations in patients with Alzheimer's dementia, and worsening of extrapyramidal symptoms, in particular tremor, in patients with dementia associated with Parkinson's disease) have been observed shortly after dose increase. They may respond to a dose reduction. In other cases, rivastigmine has been discontinued.

Recommended Dose and Dosage Adjustment

Novo-Rivastigmine (rivastigmine hydrogen tartrate) *capsules should only be prescribed by (or following consultation with,) clinicians who are experienced in the diagnosis and management of dementia.*

Adults:

The usual maintenance dose range for Novo-Rivastigmine is 6-12 mg/day. The following dosage escalation recommendations, derived from clinical trial data, are provided as a guide only, as individual tolerance to dose increases will vary. The incidence of cholinergic adverse events associated with rivastigmine hydrogen tartrate increase with dose and are more prevalent in females (see **ADVERSE REACTIONS**).

Dementia of the Alzheimer's Type

The starting dose of Novo-Rivastigmine is 1.5 mg b.i.d. (3 mg/day). If this initial dose is well tolerated, after a minimum of 2 weeks the dose may be increased to 3 mg b.i.d. (6 mg/day). Dose increases above 6 mg/day should proceed cautiously. Increases to 4.5 mg b.i.d., (9 mg/day) and then 6 mg b.i.d. (12 mg/day) should also be based on good tolerability of the current dose and should only be considered after a minimum of two weeks treatment at that dose level. The maximum dose should not exceed 6 mg b.i.d (12 mg/day).

Dementia Associated with Parkinson's Disease

In dementia associated with Parkinson's disease, the starting dose of rivastigmine hydrogen tartrate is 1.5 mg b.i.d.; subsequently, the dose may be increased to 3 mg b.i.d.; and further to 4.5 mg b.i.d.; and 6 mg b.i.d.; based on tolerability, with a minimum of 4 weeks at each dose.

Dose Interruption:

Following initiation of therapy or any dosage increase, patients should be closely monitored for adverse effects. If adverse effects (e.g. nausea, vomiting, abdominal pain, loss of appetite) are observed during treatment, the patient should be instructed to stop treatment for several doses and then restart at the same dose level, or lower, as clinically indicated. Anytime treatment is interrupted for longer than several days, patients should be instructed to reinstate treatment with the lowest daily dose (i.e. 1.5 mg b.i.d or 1.5 mg o.d, as clinically indicated) and be re-titrated to their maintenance dose as described above (see **WARNINGS AND PRECAUTIONS**). If side effects persist, the drug should be discontinued.

Novo-Rivastigmine should be taken with food in divided doses in the morning and evening.

OVERDOSAGE

Symptoms: Overdosage with cholinesterase inhibitors can result in cholinergic crisis characterised by severe nausea, vomiting, diarrhoea, hypertension, hallucinations, salivation, sweating, hypotension, respiratory depression, collapse and convulsions. Increasing muscle weakness is a possibility and may result in death if respiratory muscles are involved. Due to the known vagotonic effect of cholinesterase inhibitors on heart rate, bradycardia and/or syncope may also occur.

Treatment: Rivastigmine hydrogen tartrate has a short plasma half-life (about 1- 2 hours) and a moderate duration of cholinesterase inhibition of 8-12 hours. It is recommended that in cases of asymptomatic overdoses, no further dose of rivastigmine hydrogen tartrate should be administered for the next 24 hours and that patients be monitored.

As in any case of overdose, general supportive measures should be utilized. Tertiary anticholinergics such as atropine may be used as an antidote for rivastigmine hydrogen tartrate overdose. Intravenous atropine sulfate titrated to effect is recommended: an initial dose of 1.0 to 2.0 mg IV with subsequent doses based upon clinical response. Atypical responses in blood pressure and heart rate have been reported with other cholinomimetics when co-administered with quaternary anticholinergics such as glycopyrrolate. Due to the short half-life of rivastigmine hydrogen tartrate, dialysis (hemodialysis, peritoneal dialysis, or hemofiltration) would not be clinically indicated in the event of an overdose.

In overdoses accompanied by severe nausea and vomiting, the use of antiemetics should be considered. In a documented case of a 46 mg overdose with rivastigmine hydrogen tartrate, a 69

year old female patient experienced vomiting, incontinence, hypertension, psychomotor retardation and loss of consciousness. The patient was managed conservatively with only supportive measures and fully recovered within 24 hours.

Dose-related signs of toxicity in animals included lacrimation, excessive salivation, vomiting, decreased locomotor activity, ataxia, twitches/flutterers, tremors and clonic convulsions.

For management of a suspected drug overdose, contact your regional Poison Control Centre.

ACTION AND CLINICAL PHARMACOLOGY

Mechanism of Action

Pathological changes in Dementia of the Alzheimer type involve cholinergic neuronal pathways that project from the basal forebrain to the cerebral cortex and hippocampus. A decrease in the function of these cholinergic pathways has been proposed to account for some of the clinical manifestations of dementia. Rivastigmine, a reversible cholinesterase inhibitor of the carbamate type, is thought to enhance cholinergic neurotransmission by slowing the degradation of acetylcholine released by cholinergic neurons through the inhibition of acetylcholinesterase. If this proposed mechanism of action is correct, rivastigmine's effect may lessen as the disease process advances and fewer cholinergic neurons remain functionally intact.

There is no evidence that rivastigmine alters the course of the underlying dementing process.

Pharmacokinetics

Absorption: Rivastigmine is well absorbed and peak plasma concentrations (C_{max}) are reached in approximately 1 hour. A doubling of the dose within the recommended dose range yields an increase in bioavailability by approximately 3 times the expected increase indicating non-linear pharmacokinetics. The estimated absolute bioavailability for a 3 mg dose in healthy young patients is low (<35%). The elimination half-life ($t_{1/2}$) of rivastigmine is about 1 to 2 hours in both the young and elderly. Plasma clearance is dose dependent and is approximately 1 L/h/kg at 3 mg in healthy young subjects. In healthy elderly male patients, plasma rivastigmine levels are approximately 30% higher than that noted in young subjects (see **Pharmacokinetics, Special Populations and Conditions: Age**). When administered with food to healthy young subjects the absorption (T_{max}) of rivastigmine was delayed by 90 min, and C_{max} was lowered while the $AUC_{0-\infty}$ increased by approximately 25%.

Distribution: Rivastigmine is approximately 40% bound to plasma proteins over a concentration range of 1 - 400 ng/mL. Rivastigmine distributes equally between blood and plasma with a blood-to-plasma partition ratio of 0.9 at concentrations which cover the therapeutic range (1 - 400 ng/mL). The apparent volume of distribution is 5 ± 3 L/kg. Rivastigmine can be detected in the CSF, reaching peak concentrations in 1-4 hours. Mean AUC_{0-12hr} ratio of CSF/plasma averaged $40 \pm 0.5\%$ following 1-6 mg bid doses.

Metabolism: Rivastigmine is subject to first pass clearance and is rapidly and extensively metabolised, primarily via esterase-, including acetylcholinesterase-, mediated hydrolysis to a decarbamylated phenolic metabolite. *In vitro* preclinical studies suggest that the decarbamylated phenolic metabolite has approximately 10% the activity of the parent compound. The plasma half-life of the decarbamylated phenolic metabolite ranges from 2.5 to 4 hours. Additional metabolites include a sulphate conjugate, a demethylated sulfate conjugate and several unidentified minor metabolites.

Evidence from *in vitro* studies suggest that the major cytochrome P450 isozymes are minimally involved in rivastigmine metabolism (see **DRUG INTERACTIONS, Overview**).

Rivastigmine inhibits acetylcholinesterase (AChE) and butyrylcholinesterase (BuChE) activity. In patients with Alzheimer's Disease significant dose-dependent inhibition of AChE and BuChE activity were noted in cerebrospinal fluid, with comparable maximum mean inhibition (62%). In plasma, significant inhibition of BuChE activity is generally observed from 1.5 hours post-dose up to 8 hours post-dose, with a maximum observed inhibition of 51% at 5 mg b.i.d. Rivastigmine may therefore inhibit the butyrylcholinesterase mediated metabolism of other drugs (see **DRUG INTERACTIONS, Overview**).

Excretion: Unchanged rivastigmine is not found in the urine; renal excretion is the major route of elimination of the metabolites. Following administration of a single 1 mg or 2.5 mg dose of ¹⁴C-labelled rivastigmine, excretion of radioactivity in the urine (expressed as a percent of the administered dose) is over 90% within 24 hours. Approximately 7% of the decarbamylated phenolic metabolite is found in the urine. The sulfate conjugates account for about 40% of the dose. Less than 1% of the administered dose is excreted in the faeces. The accumulation potential of rivastigmine and its decarbamylated phenolic metabolite in patients with Alzheimer's disease has not been systematically studied however, population pharmacokinetic analyses suggest that no accumulation is expected.

Special Populations and Conditions

Age: In a study in which the effect of age on the pharmacokinetics of rivastigmine was assessed, 24 healthy male elderly (age range: 61 - 71 years) and 24 healthy young patients (age range: 19 - 40 years) received 1.0 mg or 2.5 mg single oral doses of rivastigmine under fasted conditions. Plasma concentrations of rivastigmine exhibited a wider range of values and tended to be higher in the elderly as compared to young subjects after the 1 mg dose. This difference was more pronounced with the higher dose (2.5 mg) at which rivastigmine plasma concentrations were 30% greater in the elderly than in young subjects. Plasma levels of the decarbamylated phenolic metabolite were not substantially affected by age.

Gender and Race: No specific pharmacokinetic study was conducted to investigate the effect of gender and race on the disposition of rivastigmine. However, retrospective pharmacokinetic analyses suggest that gender and race (Blacks, Oriental, and Caucasians) will not affect the clearance of rivastigmine.

Hepatic Insufficiency: In a single dose study of 10 subjects with biopsy proven liver impairment (Child-Pugh score of 5-12), plasma concentrations of rivastigmine were increased, while that of the decarbamylated phenolic metabolite were decreased by about 60% compared to an age, weight and gender matched control group. The safety and efficacy of rivastigmine in Alzheimer's Disease patients with hepatic impairment have not been studied (see **WARNINGS AND PRECAUTIONS, Hepatic/Biliary/Pancreatic**).

Renal Insufficiency: In a single-dose study of 8 subjects with moderate renal impairment (GFR = 10-50 mL/min) mean peak plasma concentrations of rivastigmine were increased by almost 2.5 fold and overall plasma levels (AUC) of the decarbamylated phenolic metabolite were increased by approximately 50% compared to levels in age, weight, and gender matched control subjects. In this same study, patients with severe renal impairment (GFR <10 mL/min, n = 8) showed no difference in rivastigmine blood levels compared to controls. The reason for this discrepancy is unclear. The safety and efficacy of rivastigmine in Alzheimer's Disease patients with renal impairment have not been studied (see **WARNINGS AND PRECAUTIONS, Renal**).

Genetic Polymorphism: The pharmacokinetics of rivastigmine in patients with butyrylcholinesterase enzyme deficiency are unknown (see **WARNINGS AND PRECAUTIONS, Genetic Polymorphism**).

Nicotine Use: Population PK analysis showed that nicotine use increases the clearance of oral rivastigmine by 23% (Smokers: n = 75; Nonsmokers: n = 549).

STORAGE AND STABILITY

Capsules: Store at room temperature (15 - 30°C). Protect from moisture.

DOSAGE FORMS, COMPOSITION AND PACKAGING

NOVO-RIVASTIGMINE (rivastigmine hydrogen tartrate) is supplied as hard-gelatin capsules containing either 1.5 mg, 3.0 mg, 4.5 mg, or 6.0 mg of rivastigmine base.

The 1.5 mg capsules are yellow, printed in red with "N" on one portion and "1.5" on the other. Available in bottles of 60 and 100 and in cartons of 30 unit dose.

The 3.0 mg capsules are orange, printed in red with "N" on one portion and "3" on the other. Available in bottles of 60 and 100 and in cartons of 30 unit dose.

The 4.5 mg capsules are red, printed in white with "N" on one portion and "4.5" on the other. Available in bottles of 60 and 100 and in cartons of 30 unit dose.

The 6.0 mg capsules are orange and red, printed in white with "N" on one portion and "6" on the other. Available in bottles of 60 and 100 and in cartons of 30 unit dose.

Composition of NOVO-RIVASTIGMINE

Capsules: Each hard gelatin capsule contains 1.5, 3.0, 4.5, or 6.0 mg of rivastigmine base. Inactive ingredients are: colloidal silicon dioxide; hydroxypropyl methylcellulose; magnesium stearate, microcrystalline cellulose; hard gelatin capsules contain gelatin, titanium dioxide and red and/or yellow iron oxides.

PART II: SCIENTIFIC INFORMATION


PHARMACEUTICAL INFORMATION

Drug Substance

Common name: Rivastigmine hydrogen tartrate
Chemical name: (+)(S)-N-ethyl-N-3-[1-(dimethylamino)ethyl]-N-methyl-phenyl carbamate hydrogen tartrate

Molecular formula and molecular mass: $C_{14}H_{22}N_2O_2 \cdot C_4H_6O_6$ and 400.42

Structural formula:



Physicochemical properties: Description: White to off-white, fine crystalline powder, very hygroscopic and deliquescent.

Melting Range: 124.0-128.0°C

Solubilities: Soluble in ethanol

pKa in n-octanol/phosphate buffer solution at pH 7: 8.85

CLINICAL TRIALS

A Randomized, Two-Way Crossover, Single-Dose, Blinded Study to Evaluate the Bioequivalence of a Test Capsule Formulation of Rivastigmine Hydrogen Tartrate (6 mg), Compared to an Equivalent Dose of a Commercially Available Reference Drug Product (^{Pr}Exelon*, Novartis Pharmaceutical Canada Inc.) in 26 Healthy, Adult Male Subjects under Fasting Conditions. A summary of the results is presented below:

Rivastigmine (1 x 6 mg capsule) From measured data uncorrected for potency Geometric Mean Arithmetic Mean (CV %)				
Parameter	Test*	Reference†	% Ratio of Geometric Means	Confidence Interval, 90%
AUC _T ‡ (ng.h/mL)	40.68 47.77 (58.96)	38.27 43.82 (53.76)	106.32	98.83 – 114.36
AUC ₁ (ng.h/mL)	41.59 48.61 (58.15)	39.08 44.59 (53.32)	106.44	99.10 – 114.33
C _{max} (ng/mL)	17.46 19.38 (47.63)	17.15 18.74 (43.20)	101.81	94.66 – 109.49
T _{max} § (h)	1.22 (36.02)	1.18 (34.26)		
T _{1/2} § (h)	1.26 (15.55)	1.28 (16.44)		

* Novo-Rivastigmine 6 mg Capsules (Novopharm Limited, Canada)

† Exelon® 6 mg Capsules (Novartis Pharma Canada, Inc., Canada.; and purchased in Canada)

§ Expressed as the arithmetic mean (CV%) only

A Randomized, Two-Way Crossover, Single-Dose, Blinded Study to Evaluate the Bioequivalence of a Test Capsule Formulation of Rivastigmine Hydrogen Tartrate (6 mg), Compared to an Equivalent Dose of a Commercially Available Reference Drug Product (PrExelon[®], Novartis Pharmaceutical Canada Inc.) in 28 Healthy, Adult Male Subjects under Fed Conditions. A summary of the results is presented below:

Rivastigmine (1 x 6 mg capsule) From measured data uncorrected for potency Geometric Mean Arithmetic Mean (CV %)				
Parameter	Test*	Reference†	% Ratio of Geometric Means	Confidence Interval, 90%
AUC _T ‡ (ng·h/mL)	29.85 33.65 (46.08)	29.65 34.42 (54.13)	100.69	94.66 – 107.10
AUC ₁ (ng·h/mL)	30.82 34.57 (45.25)	30.61 35.25 (53.04)	100.69	94.87 – 106.85
C _{max} (ng/mL)	9.41 10.40 (41.49)	9.47 10.64 (47.77)	99.32	92.44 – 106.71
T _{max} § (h)	2.57 (27.95)	2.62 (32.63)		
T _{1/2} § (h)	1.36 (17.13)	1.33 (18.92)		

* Novo-Rivastigmine 6 mg Capsules (Novopharm Limited, Canada)

† Exelon[®] 6 mg Capsules (Novartis Pharma Canada, Inc., Canada., and purchased in Canada)

§ Expressed as the arithmetic mean (CV%) only

Dementia of the Alzheimer's Type

Study demographics and trial design

Efficacy data for rivastigmine in the symptomatic treatment of patients with mild to moderate dementia of the Alzheimer's type (diagnosed by DSM-IV and NINCDS criteria, Mini- Mental State Examination ≥ 10 and ≤ 26) were derived from four clinical trials. These studies were randomized, double-blind, and placebo controlled. The mean age of patients was 73 years (range: 41 to 95). Approximately 59% of the patients were women and 41% were men, while the racial distribution was 87% Caucasian, 4% Black and 9% Other. In these clinical studies, the effectiveness of rivastigmine was evaluated using the following criteria: for primary efficacy two measures were used, (1) the cognitive subscale of the Alzheimer's Disease Assessment Scale (ADAS-Cog), a widely used and well validated multi-item instrument which samples cognitive domains affected by the disease and (2) the CIBIC-Plus (Clinician Interview Based Impression of Change that required caregiver information). The CIBIC-Plus evaluates four major areas of functioning: general, cognition, behaviour and activities of daily living. As a secondary efficacy measure, the Progressive Deterioration Scale (PDS) was used. The PDS is a caregiver-rated evaluation which yields a compound score derived from a visual analogue scale of 29 items concerning participation in activities of daily living.

Study results

Results for two of these studies, in which a flexible maintenance-dose regimen was used, are presented here. The data shown below were obtained from the Intent-to-Treat population (ITT analysis, i.e., all patients who were randomized to treatment, regardless of whether or not they were able to complete the study. For patients unable to complete the study, their last observation while on treatment was carried forward and used at endpoint).

STUDY I (B352, USA, 26 week trial)


This trial was of 26 weeks duration and was conducted in the USA. The study was subdivided into two phases, a forced titration phase, which could last up to 12 weeks, followed by a 14 week maintenance flexible-dose phase. A total of 699 patients were randomized to a 1-4 mg daily dose ($n = 233$) or a 6-12 mg daily dose ($n = 231$) of rivastigmine or placebo ($n = 235$) to be taken with food in two divided doses. Patients in the active treatment groups must have been able to tolerate the minimum dose in their assigned group (i.e. 0.5 mg bid or 3 mg bid) by titration Week 7 or they were discontinued. The dose escalation rate for the 1-4 mg/day group was: Starting dose 0.5 mg bid with 0.5 mg bid increases every one or two weeks according to tolerability. The dose escalation rate for the 6-12 mg/day group was: Starting dose 1 mg bid increased to 1.5 mg bid after 3 days. Subsequent dose increases were at 0.5 mg bid or 0.75 mg bid every one or two weeks according to patient tolerability. The baseline mean Mini Mental State Exam (MMSE) score of patients was 19.7 and the mean score on the Global Deterioration Scale (GDS) was 4.0.

Effects on ADAS-cog: At baseline, mean ADAS-cog scores (mean \pm SE) were for the placebo group: 20.88 ± 0.72 units; for the 1-4 mg/day group: 22.65 ± 0.79 units and for the 6-12 mg/day group: 22.70 ± 0.84 units. At the first measurement of efficacy (Week 12) mean ADAS-cog

change scores from placebo (mean \pm standard error) were: 0.82 ± 0.52 units for the 1-4 mg/day group and 3.24 ± 0.54 units for the 6-12 mg/day dose groups. Differences from placebo were statistically significantly different only for the 6-12 mg/day group. At Week 18, mean change scores from placebo were significant for both rivastigmine dose groups (1-4 mg/day: 1.67 ± 0.54 units; 6-12 mg/day: 3.83 ± 0.57 units). Both rivastigmine treated groups also showed significant differences from placebo in ADAS-cog mean change scores at Week 26: (1-4 mg/day: 1.66 ± 0.57 units; 6-12 mg/day: 4.32 ± 0.60 units). A greater treatment effect size is noted for the 6-12 mg/day treatment. At the end of the 26-week treatment period, either no evidence of deterioration or an improvement was observed in 27% of the placebo group, 35% (1-4 mg/day) and 51% (6-12 mg/day) in the rivastigmine groups. The difference between the 6-12 mg/day group and the placebo group was statistically significant. A 4 point improvement in ADAS-cog score from baseline was observed in 6% of placebo patients, 12% (1-4 mg/day) and 23% (6-12 mg/day) of rivastigmine treated patients at the end of the 26 week period. Statistical significance from placebo for this categorical measure was noted for both the 1-4 mg/day and 6-12 mg/day group.


Figure 1 illustrates the time course for the change from baseline in ADAS-cog scores for all three dose groups over the 26 weeks of the study. At 26 weeks of treatment, the mean differences in the ADAS-cog change scores for the rivastigmine-treated patients compared to the patients treated with placebo were 1.7 and 4.3 units for the 1-4 mg and 6-12 mg treatments, respectively. Both treatments were statistically significantly superior to placebo and the 6-12 mg/day range was significantly superior to the 1-4 mg/day range.

Figure 1: Time-course of the Change from Baseline in ADAS-cog Score (ITT-LOCF Population)



Effects on CIBIC-Plus. At Week 26 the mean drug-placebo differences were 0.22 ± 0.11 units for the 1-4 mg/day group and 0.36 ± 0.12 units for the 6-12 mg/day group. Differences from placebo were statistically significant. However, there was no statistically significant difference between the two active treatments. A histogram of the frequency distribution of CIBIC-Plus scores achieved at Week 26 by patients assigned to each of the three treatment groups is shown in Figure 2.

Figure 2: Frequency Distribution of CIBIC-Plus Scores at Week 26 (ITT-LOCF)



Effects on PDS: The progressive deterioration scale was used as a secondary efficacy measure. At baseline, mean PDS scores (mean \pm SE) were for the placebo group 53.7 ± 1.2 units; for the 1-4 mg/day group: 54.7 ± 1.2 units; for the 6-12 mg/day group: 52.0 ± 1.2 units. At Week 26, the placebo group declined an average of 5.2 ± 0.7 units, the 1-4 mg/day group declined 5.3 ± 0.7 units and the 6-12 mg/day group deteriorated minimally (1.0 ± 0.8 units). The difference between the 6-12 mg/day group and the placebo group was statistically significant.

STUDY II (B303, Multinational, 26 week trial)


This trial of 26 weeks duration was a multinational study (Austria, Canada, France, Germany, Switzerland and USA). A total of 725 patients were randomized into three different treatment arms. Placebo: n = 239, 1-4 mg/day rivastigmine: n = 243, 6-12 mg/day rivastigmine: n = 243. As in Study I, this trial was comprised of two phases, a forced titration phase, which could last up to 12 weeks, followed by a maintenance flexible-dose phase. Patients in the active treatment groups must have been able to tolerate the minimum dose in their assigned group (i.e. 0.5 mg bid or 3 mg bid) by titration Week 7 or they were discontinued. The baseline mean Mini Mental State Exam (MMSE) score was 20 and the mean score on the Global Deterioration Scale (GDS) was 4.0.

Effects on ADAS-cog: At baseline, mean ADAS-cog scores (mean \pm SE) were for the placebo group: 23.22 ± 0.75 units; for the 1-4 mg/day group: 24.05 ± 0.77 units and for the 6-12 mg/day

group: 23.73 ± 0.84 units. At the first measurement of efficacy (Week 12) the difference in mean ADAS-cog change scores (mean \pm standard error) for rivastigmine treated patients compared to placebo treated patients for the ITT-LOCF population were for the 1-4 mg/day group: 0.19 ± 0.55 units and for the 6-12 mg/day group: 1.71 ± 0.57 units. Only the difference between the 6-12 mg/day group and placebo was significant at this time point. At Weeks 18 and 26 mean ADAS-cog change scores from placebo were for the 1-4 mg/day group: 0.57 ± 0.59 (Week 18); 0.22 ± 0.67 units (Week 26) and for the 6-12 mg/day group: 1.77 ± 0.60 units (Week 18); 2.29 ± 0.69 units (Week 26). As for Week 12, only the difference between the 6-12 mg/day group and placebo was statistically significant. At the end of the 26-week treatment period, either no evidence of deterioration or an improvement was observed in 40% of the placebo group, 45% (1-4 mg/day) and 52% (6-12 mg/day) in the rivastigmine groups. A 4 point improvement in ADAS-cog score from baseline was observed in 18% of patients who received placebo, 16% (1-4 mg/day) and 27% (6-12 mg/day) of rivastigmine treated patients at Week 26. Differences between rivastigmine (6-12 mg/day) and placebo treated groups were significant for both categorical measures.


Figure 3 illustrates the time course for the change from baseline in ADAS-cog scores for all three dose groups over the 26 weeks of the study. At 26 weeks of treatment, the mean differences in the ADAS-cog change scores for the rivastigmine-treated patients compared to the patients treated with placebo were 0.2 and 2.3 units for the 1-4 mg and 6-12 mg treatments, respectively. The 6-12 mg/day group was statistically significantly superior to placebo, as well as to the 1-4 mg/day group. The difference between the 1-4 mg/day group and placebo was not statistically significant.

Figure 3: Time-course of the Change from Baseline in ADAS-cog Score (ITT-LOCF Population)



Effects on CIBIC-Plus: At Week 26 the mean drug-placebo differences were 0.15 ± 0.14 units for the 1-4 mg/day group and 0.44 ± 0.15 units for the 6-12 mg/day group. Differences from placebo were statistically significant only for the 6-12 mg/day dose group. A histogram of the frequency distribution of CIBIC Plus scores achieved at Week 26 by patients assigned to each of the three treatment groups is shown in Figure 4.

Figure 4: Frequency Distribution of CIBIC-Plus Scores at Week 26 (ITT-LOCF)



Effects on PDS: The progressive deterioration scale was used as a secondary efficacy measure. At baseline, mean PDS scores (mean \pm SE) were for the placebo group: 54.8 ± 1.3 units; for the 1-4 mg/day group: 53.8 ± 1.3 units; for the 6-12 mg/day group 55.2 ± 1.2 units. At Week 26, while the placebo group declined an average of 2.2 ± 0.9 units and the 1-4 mg/day group deteriorated by 3.3 ± 0.9 units, the 6-12 mg/day group improved by 0.5 ± 1.0 units, which was a statistically significant difference. The 6-12 mg/day group was statistically significantly superior to placebo as well as the lower dose range.

Data from these controlled clinical trials suggest that rivastigmine doses between 6-12 mg/day are more likely to result in beneficial symptomatic effects.

Dementia Associated with Parkinson’s Disease

The dementia which occurs in patients with an established diagnosis of idiopathic Parkinson’s disease is purportedly characterized by impairments in memory retrieval, executive function, and attention. However, based on clinical pathologic data for 110 cases of “Parkinson’s disease dementia” (PDD) from 3 well-designed studies, it is internationally recognized that the differential diagnosis of this type of dementia from Alzheimer’s disease can reliably be made without the necessity to document the specific deficits described above. Instead, the diagnostic criteria are: patients in whom a progressive dementia syndrome occurs at least 2 years after a diagnosis of Parkinson’s disease has been made, and in whom other causes of dementia have been ruled out (see INDICATIONS).

Study demographics and trial design

The efficacy of rivastigmine in the symptomatic treatment of patients with mild to moderate dementia with onset at least 2 years after the initial diagnosis of idiopathic Parkinson's disease was demonstrated by the results of one 24-week randomized, double-blind, placebo-controlled trial, with n = 541 patients randomized in a ratio of 2:1 to an rivastigmine or placebo arm.

The diagnosis of idiopathic Parkinson's disease was based on the United Kingdom Parkinson's Disease Society Brain Bank clinical criteria. The diagnosis of dementia was based on the criteria stipulated under the DSM-IV category "Dementia Due To Other General Medical Condition" (code 294.1), with the additional requirements, as described above, that the dementia must have occurred at least 2 years after a diagnosis of Parkinson's disease has been made, and alternate causes of dementia were excluded by clinical history, physical and neurological examination, brain imaging, and relevant blood tests. Thus, patients were not required to have a distinctive pattern of cognitive deficits as part of the dementia. Patients enrolled in the study had a MMSE score ≥ 10 and ≤ 24 at entry. At baseline 70% of patients had mild dementia (MMSE 17-24) and in 71% of patients severity of Parkinson's disease was moderate (Hoehn and Yahr stage 2.5 to 4). The mean age of patients participating in this trial was 72.7 years (range: 50 to 91). Approximately 65% of patients were men, and 99.6% were Caucasian.

A flexible maintenance-dose regimen was used, with rivastigmine ranging from 3-12 mg per day, divided doses. The 24-week study was divided into a 16-week titration phase, with dose increases every 4 weeks to achieve a maximum well-tolerated dose, followed by an 8-week maintenance phase. The patients in the active treatment arm of the study were maintained at their highest tolerated dose within the specified dose range.

Efficacy measures

As with the Alzheimer's type dementia studies, the outcome data were obtained from the Intent-to-Treat population (ITT analysis, i.e., all patients who were randomized to treatment, and received at least one dose of medication, regardless of whether or not they were able to complete the study. For patients unable to complete the study, their last observation while on treatment was carried forward and used at endpoint ie LOCF).

This study used a dual outcome assessment strategy to measure the efficacy of rivastigmine,

1. The ability of rivastigmine to improve cognitive performance was assessed with the ADAS-cog. This instrument is validated for assessment of cognitive domains affected by dementia of the Alzheimer's type.
2. The ability of rivastigmine to produce an overall clinical effect was assessed using the Alzheimer's Disease Cooperative Study - Clinician's Global Impression of Change (ADCS-CGIC). The ADCS-CGIC is a more standardized form of CIBIC-Plus that focuses on clinicians' observations of change in the patient's cognitive, functional and behavioral performance.


Secondary efficacy measures that focused on cognitive impairments typically observed in patients with PDD included the Cognitive Drug Research (CDR) Computerized Assessment System for assessment of attentional deficit, and the Delis-Kaplan Executive Function System (D-KEFS) for assessment of executive dysfunction.

Study results

Effects on the ADAS-cog: At baseline, mean ADAS-cog scores (mean \pm SD) were 24.5 ± 10.6 points for the placebo-treated group and 24.0 ± 10.3 points for the rivastigmine-treated group. At the first measurement of efficacy (Week 16), mean ADAS-cog change scores from placebo for the rivastigmine-treated patients was 2.74 (95% C.I. 1.42; 4.06; $p < 0.001$). At 24 weeks of treatment, the mean difference in the ADAS-cog change scores for the rivastigmine-treated patients compared to the patients on placebo was 3.54 (95% C.I. 2.05; 5.04; $p < 0.001$). This treatment difference was statistically significant in favor of rivastigmine when compared to placebo. At the end of the 24-week treatment period, a 4-point improvement in ADAS-cog score from baseline was observed in 29% of placebo-treated patients compared to 40% of rivastigmine-treated patients. Statistical significance from the placebo-treated group for this categorical measure was noted.

Figure 5 illustrates the time course for the change from baseline in ADAS-cog scores for both treatment groups over the 24-week study. At 24 weeks of treatment, the mean difference in the ADAS-cog change scores for the rivastigmine-treated patients compared to the patients on placebo was 3.6 points. This treatment difference was statistically significant in favor of when compared to placebo.


Figure 5: Time course of the Change from Baseline in ADAS-cog (ITT-LOCF Population)



Effects on the ADCS-CGIC: At 24 weeks, the mean difference in change scores between the rivastigmine -treated group and the placebo-treated group from baseline was 0.6 points.

The categorical analysis showed statistically significantly more patients who improved and less patients who had worsening in the rivastigmine treated group, compared to those treated with placebo ($p < 0.001$). A histogram of the distribution of patients' scores on the ADCS-CGIC at Week-24 is shown in Figure 6.

Figure 6: Frequency Distribution of ADCS-CGIC Scores at Week 24 (ITT-LOCF)



Secondary cognitive efficacy measures: Results of the analysis of secondary efficacy measures (change from baseline at 24 weeks) for CDR power of attention score and for D-KEFS verbal fluency test supported the co-primary outcomes.

Comparative Bioavailability Studies

A comparative pharmacokinetic study in 53 patients with Alzheimer disease demonstrated that rivastigmine hydrogen tartrate capsules and oral solution produced comparable serum concentrations of rivastigmine at steady state (Table 7). Statistical analyses also demonstrated comparable pharmacokinetic parameters between the two dosage forms for rivastigmine.

Table:7: Preserved Pharmacokinetic Parameters for Rivastigmine Obtained After Treatments with Capsules and Oral Solution (n=53)

Parameter	Rivastigmine capsules Mean	
	3 mg	6 mg
Cmax (µg/mL)	8.0	22.7
Tmax (h)	1.0	1.0
AUCT (µg•Ch/mL)	17.9	69.1
AUCI (µg•Ch/mL)	18.8	70.5
t½ (h)	1.4	1.7

DETAILED PHARMACOLOGY

Animal Pharmacodynamics

In vitro and *in vivo* pharmacology studies with rivastigmine predominantly focused on the main action of the drug: inhibition of acetylcholinesterase (AChE) activity, accumulation of acetylcholine (ACh) levels and cholinergic effects.

IC₅₀ values for rivastigmine-induced inhibition of AChE activity *in vitro* in various rat brain areas were as follows: Cortex: 1.7 x 10⁻⁵M; Hippocampus: 1.5 x 10⁻⁵M, Striatum: 2.0 x 10⁻⁵M and Pons/Medulla: 2.0 x 10⁻⁵M.

AChE activity measured *ex vivo* was inhibited in several rat brain regions following p.o. administration of single rivastigmine doses. The effect of rivastigmine single p.o. doses on enzyme activity was noted to be more pronounced in the hippocampus and cortex than in the striatum and pons/medulla of these rats (IC₅₀ : Cortex: 0.5 mg/kg, p.o.; Hippocampus: 1 mg/kg, p.o.; Striatum: 1.75 mg/kg, p.o. and Pons/Medulla: 2 mg/kg, p.o.). Physostigmine, administered s.c., inhibited AChE activity to an equal degree in all rat brain regions examined (IC₅₀ : Cortex: 0.22 mg/kg; Hippocampus: 0.27 mg/kg, Striatum: 0.28 mg/kg and Pons/Medulla: 0.27mg/kg).

Single p.o. doses of rivastigmine also resulted in an increased accumulation of ACh levels in the rat brain which were more pronounced in the cortex than the hippocampus or striatum.

When administered s.c., a single dose (0.75 mg/kg) of rivastigmine inhibited AChE activity in the periphery (Heart: 55% control values; Blood: 34% control values) to an equivalent degree as in brain (Cortex: 37% control values; Hippocampus 45% control values).

Chronic continuous dosing with rivastigmine also resulted in diminished selectivity of the drug for AChE activity in brain versus the periphery (heart/blood). Similarly, the apparent selectivity

of rivastigmine for AChE within specific rat brain areas was also lost with chronic continuous dosing (14 days).

Induction of slow rhythmic activity in the hippocampal EEG (synchronization of theta-waves) has been proposed to reflect increased central muscarinic activity. Rivastigmine synchronized rhythmical slow wave activity in the hippocampal EEG in rats at a threshold dose of 75 µg/kg both i.p. and p.o. Similar effects were noted with physostigmine at a dose of 75 µg/kg i.p.

Rivastigmine, in a dose range of 0.01-1.5 mg/kg i.v. had minimal effects on circulatory parameters in the anaesthetized cat, while the effects of physostigmine (0.01 – 1.71 mg/kg, i.v.) on circulatory parameters in this animal model were more potent. At a dose of 0.75 mg/kg i.v. rivastigmine induced central effects manifested by strong tremor or slight cramps. Similar effects were noted with physostigmine doses of 0.14 mg/kg, i.v.

The cardiovascular effect of rivastigmine was studied in awake normotensive male adult rats. Oral administration of rivastigmine (1.88 mg/kg) induced weak bradycardia (14%) which was reversed by methylscopolamine. At higher doses (5.6 mg/kg, p.o.) rivastigmine significantly increased (29%) blood pressure. This effect was blocked by scopolamine (1 µmole/rat) but not the peripheral blocker n-methylscopolamine (1 mg/kg, i.v.).

The pulmonary effects of rivastigmine were assessed using the ventilated guinea-pig model. Rivastigmine at doses of 0.01 to 1 mg/kg i.v. did not affect airway resistance. However, pretreatment with 0.1 mg/kg i.v. rivastigmine resulted in a potentiation of ACh-induced bronchospasm at all ACh doses tested (3.2 µg/kg, 5.6 µg/kg and 10 µg/kg, i.v.).

It was concluded that rivastigmine is an acetylcholinesterase inhibitor of the carbamate type. Its main preclinical properties are:

- high central to peripheral cholinergic activity ratio after a single p.o. dose;
- selectivity for cortical and hippocampal brain regions after a single p.o. dose;
- prolonged duration of action (hours); and
- low activity on cardiovascular system at centrally active doses

Animal Pharmacokinetics

The studies performed to evaluate the pharmacokinetics in animals with rivastigmine allow the following conclusions to be drawn concerning rivastigmine:

- peak blood concentrations rapidly achieved following oral administration;
- good oral absorption in all species studied, including man;
- bioavailability increased disproportionately with increasing dose, due to a saturable first pass metabolism;
- total radioactivity rapidly and widely distributed into tissues in rodents;
- extensive metabolism in all species studied prior to excretion primarily via the renal route;
- metabolism qualitatively, but not quantitatively, similar in all species studied *in vivo* and *in vitro* - the main pathways were decarbamylation, conjugation and N-dealkylation;

- metabolic clearance linear except in the dog;
- accumulation and hepatic enzyme induction not present after repeated oral dosing;
- good brain penetration.

TOXICOLOGY

Acute Toxicology

The estimated oral LD₅₀ values in mice were 5.6 mg/kg (males) and 13.8 mg/kg (females). The estimated oral LD₅₀ values in rats were 8.1 mg/kg (males) and 13.8 mg/kg (females). These dose levels are more than 20 times the maximum recommended human dose of 12 mg/day (assuming a 50 kg body weight). The LD₅₀ values determined in these studies are summarized in Table 8.

Table 8

Species	Strain	Sex	Route	Dose Levels (mg/kg)	LD ₅₀ value (mg/kg)
Mouse	CD-1	M	Oral	0.63, 6.25, 31.25	5.6
		F	Oral	0.63, 6.25, 31.25	13.8
	CD-1	M	i.v.	1.25, 3.13, 3.75	2.8
		F	i.v.	3.13, 3.75, 5.0	4.1
Rat	CD	M	Oral	0.63, 6.25, 31.25	8.1
		F	Oral	0.63, 6.25, 31.25	13.8
Mouse	CD-1	M	i.p.	0.63, 6.25, 31.25	1.9
		F	i.p.	0.63, 6.25, 31.25	1.9
Rat	CD	M	i.p.	0.63, 6.25, 31.25	4.4
		F	i.p.	0.63, 6.25, 31.25	1.9
Dog	Beagle	M	Oral	0.31, 1.25, 5.0	>1 and <5

The results of these studies demonstrate the moderate toxicity of ENA 713 following acute oral, i.v., and i.p. administration to mice, rats or dogs.

Long Term Toxicology

Table 9 outlines the long-term toxicology studies done in rats, mice, dogs and monkeys with rivastigmine.

Table 9

Species	Duration of Study Weeks	Route of Administration	No. of animals/group	Dose Levels (mg/kg/day)
Mouse	8	oral(gav)	5M, 5F	0, 0.38, 0.78, 1.56, 2.5, 3.13, 6.25
	13	oral (diet)	10M, 10F	0, 0.13, 0.5-75.0, 1.5
	104	oral(gav)	70M, 70F	0, 0.25, 0.63, 1.56
Rat	2	oral(gav)	10M	0.03, 0.25, 2.50
	2	i.v.	15M, 15F	0, 0.5, 2.5
	4	oral(gav)	10M, 10F	0, 0.38, 1.5, 3.75
	13	oral(gav)	10M	0, 0.13, 0.5-6.0, 1.50
	26	oral(gav)	15M, 15F	0, 0.11, 0.45, 1.50
	52+	oral(gav)	25M, 25F	0, 0.13, 0.38, 1.13, 1.88

Species	Duration of Study Weeks	Route of Administration	No. of animals/group	Dose Levels (mg/kg/day)
	104	oral(gav)	75M, 75F	0, 0.13, 0.38, 1.13
Dog	2	oral(gav)	1M, 1F	0.06, 0.63, 2.50-1.88
	2	i.v.	2M, 2F	0, 0.09, 0.47
	4	oral(gav)	3M, 3F	0, 0.04, 0.38, 2.25-1.88
	4	oral(gav)	3M, 3F	0, 0.11, 0.19, 0.26
	26	oral(gav)	3M, 3F	0, 0.11, 0.45, 1.58
	52	oral(gav)	4M, 4F	0, 0.19, 0.38, 1.56-1.31
Monkey	2	oral(gav)	1M, 1F	1.88 (days 1-7) 2.50 (days 8-10) 3.75 (days 11-13) 6.25 (day 14)

Mice: In multidose studies in mice, the toxic dose for rivastigmine was 2.5 mg/kg/day by oral gavage; oral admixture doses up to 75 mg/kg/day resulted in one mortality during Week 14 at a dose of 75 mg/kg/day.

Clinical signs were typical of cholinergic stimulation and statistically significant decreases in body weights and food consumption were seen at doses of 2.5 mg/kg/day and higher. Plasma (butyryl) and acetylcholinesterase activities were decreased in the 13-week study in the 0.5-75 mg/kg/day group. Selected tissue cholinesterase activity (liver, brain, and psoas muscle) was reduced at doses of 1.5 and 0.5-75 mg/kg/day.

Rats: One mortality in rats at 0.11 mg/kg/day was of unknown causes and was considered to be of questionable biological significance. There were no treatment-related effects on mortality at doses as high as 1.13 mg/kg/day. Treatment related dose-dependent clinical signs were consistent with excessive cholinergic stimulation of the peripheral and central nervous systems and were observed at a dose as low as 0.11 mg/kg/day. Statistically significant decreases in body weight gains and food consumption were observed at 1.13 mg/kg/day. Statistically significant decreases in triglycerides were observed at doses of 1.13, 1.5, 1.88, and 3.75 mg/kg/day in the 4- and 52-week studies, and were considered to be related to rivastigmine. Significant decrease in butylcholinesterase activities was observed at 2.5 and 3.75 mg/kg/day in the 15-day and 4-week studies; and in urinary pH at 3.75 mg/kg/day in males in the 4-week study, considered to be of minimal biological significance. Effects on plasma cholinesterase activity were not observed at doses below 2.5 mg/kg/day in any oral gavage study.

Dogs: Doses were lowered in three studies due to overt clinical signs. Treatment related unscheduled deaths occurred in two dog studies at doses of 1.56/1.31 or 2.25/1.88 mg/kg/day. Treatment related dose-dependent clinical signs were observed at doses as low as 0.19 mg/kg/day and were typical of excessive cholinergic stimulation. Clonic/tonic convulsion was observed in one 0.38 mg/kg/day male on one episode and one female (1.56/1.31 mg/kg/day) on two episodes. Statistically significant dose-related decreases in butylcholinesterase activity were observed at doses as low as 0.04 mg/kg/day. Statistically significant decreases in liver and brain cholinesterase activity at 2.25/1.88 mg/kg/day and liver cholinesterase at 0.45 and 1.58 mg/kg/day were observed in the 4-week and 26-week studies. In life pathology findings revealed that dogs were very sensitive to rivastigmine, particularly on the GI tract.

Monkeys: There was no mortality in the monkey study, however only 2 animals were treated for a period of 2 weeks (See Table 9). There appeared to be slight reduction in body weight and food consumption. Plasma (butyryl) cholinesterase activity was reduced by 15% or 29% and 6% or 14% on Days 6 and 14, respectively. Erythrocyte cholinesterase activity was reduced by 60% or 90% and 40% or 60% at the same time points. It was concluded that rivastigmine was better tolerated in monkeys for up to 2 weeks, than in rats or dogs.

Teratological and Reproductive Studies

Oral studies in pregnant rats at dose levels up to 2.3 mg-base/kg/day and pregnant rabbits at dose levels up to 2.3 mg-base/kg/day gave no indication of a teratogenic potential for rivastigmine. Similarly, there was no evidence of adverse effects of rivastigmine on fertility and reproductive performance in the rat at dose levels up to 1.1 mg-base/kg/day. A minor delay in development up to mating was noted for the F1 generation, however, no teratological changes were reported.

Mutagenicity

Rivastigmine was not mutagenic in the Ames test, a test for induction of DNA repair synthesis, the *in vivo* micronucleus test in mice, and the HGPRT test in V79 Chinese hamster cells. The *in vitro* chromosomal aberration test in V79 Chinese hamster cells showed an increase in aberrations only in the presence of liver metabolic enzymes and at a concentration at least 10,000 times greater than that likely to be found in human plasma.

Carcinogenicity

No evidence of carcinogenicity was found in studies conducted at dose levels up to 1.1 mg-base/kg/day in rats and 1.6 mg-base/kg/day in mice. Normalized to body surface area, these dose levels are approximately equivalent to 12 mg of rivastigmine base administered to a 70 kg human.

REFERENCES

1. Anand, R., Gharabawi, G, Enz, A. Efficacy and safety results of the early phase studies with Exelon (ENA 713) in Alzheimer's Disease: an overview. *J Drug Dev Clin Pract*; 8:109-116 (1996).
2. Anand, R, and Gharabawi, G.: Clinical Development of Exelon (ENA 713). The ADENA programme. *J Drug Dev Clin Pract*; 8:117-122(1996).
3. Bartus, R.T., Dean, R. L., Pontecorvo, M.J., Flicker, C. The cholinergic hypothesis: a historical overview, current perspective, and future directions. *Ann N. Y. Acad Sci* 444: 332-358 (1985).
4. Becker, R.E. Therapy of the cognitive deficit in Alzheimer's disease. The cholinergic system in "Cholinergic Basis for Alzheimer's Therapy". R. Becker, E. Giacobini, (eds: Birkhauser, Boston, pp. 1-22 (1991).
5. Corey-Bloom, J., Anand, R. and Veach, J. for the ENA 713 B352 Study Group. A randomized trial evaluating the efficacy and safety of ENA 713 (rivastigmine tartrate), a new acetylcholinesterase inhibitor, in patients with mild to moderately severe Alzheimer's disease. *Intl J Ger Psychopharm.* 1:55-65 (1998).¹
6. Cutler, N. R. et al. Dose dependent CSF acetylcholinesterase inhibition by SDZ ENA 713 in Alzheimer's Disease. *Acta Neurol Scand*; 97: 244-250 (1998).
7. Becker, R.E, Giacobini, E. Mechanisms of cholinesterase inhibition in senile dementia of the Alzheimer type: clinical, pharmacological, and therapeutic aspects. *Curr Trends Rev*; 12: 163-195 (1988)
8. Deutsch, J.A., The cholinergic synapse and the site of memory. *Science*; 174:788-794 (1971).
9. Enz, A. and Floersheim, P. Cholinesterase inhibitors: An overview of their mechanisms of action in Alzheimer's Disease. In "Alzheimer's Disease: From Molecular Biology to Therapy" R. Becker, E. Giacobini. Birkhauser, Boston, pp. 211-215 (1996).
10. Enz, A., Boddeke, H., Gray, J., et al. Pharmacologic and clinicopharmacologic properties of SDZ ENA 713, a centrally selective acetylcholinesterase inhibitor. *Ann N Y Acad Sci*, 640: 272-275. (1991).
11. Emre, M., Aarsland, D., Albanese, A., et al.,: Rivastigmine for dementia associated with Parkinson's disease. *N Engl J Med*; 351:2509-2518 (2004).

¹ The title of this article does not reflect the prospectively defined disease severity criteria in this study.

12. Foistein, M.F., Folstein McHugh, P.R. "Mini-mental state": a practical method for grading the cognitive state of patients for the clinician, *J Psychiatry Res*, 12: 189-198 (1975).
13. Holsboer-Trachsler, E., Hatzinger, M., Stohler, R., et al: Effects of the novel acetylcholinesterase inhibitor SDZ ENA 713 on sleep in man. *Neuropsychopharmacology*, 8: 87-92 (1993).
14. Katzman, R. "Alzheimer's Disease". *N. Eng. J. Med.* 314: 964-973, (1986).
15. Katzman, R., Saitoh, T. *Advances in Alzheimer's Disease. FASEB J*; 5:278-286 (1991).
16. Mohs, R.C., Rosen, W.G., Davis, K.L. The Alzheimer's Disease Assessment Scale: An Instrument for Assessing Treatment Efficacy. *Psychopharm. Bull.*; 19:448-450 (1983).
17. O'Brien, R.D. Phosphorylation and carbamylation of acetylcholinesterase. *Ann. N.Y. Acad. Sci.* 180: 204-214 (1969).
18. Perry, E.K., Acetylcholine and Alzheimer's Disease *Br. J Psychiatry*; 152: 737-740 (1988)
19. Perry, E.K., Tomlinson, B.E., Blessed, G. Bergman, K., Gibson, P.H., Perry R.H. Correlation of cholinergic abnormalities with senile plaques and mental test scores in senile dementia. *Br. Med J.* ii: 1457-1459, (1978).
20. Reisberg, B., Ferris, S., de Leon, M.J., et al. The Global Deterioration Scale (GDS): An Instrument for the Assessment of Primary Degenerative Dementia (PDD). *Am J Psych*; 139:1136-1139 (1982).
21. Rosen, W.G., Mohs, R. C., Davis, K.L. A new rating scale for Alzheimer's Disease. *Am J Psychiatry*; 141:1356-1364 (1984).
22. Rosler, M. et al. on behalf of the B303 Exelon Study Group. Efficacy and safety of rivastigmine in patients with Alzheimer's disease: international randomised controlled trial *BMJ*, 318: 633-640 (1999).
23. Sramek, J. J., Anand, R., Wardle, T.S, et al: Safety/tolerability trial of SDZ ENA 713 in patients with probable Alzheimer's disease. *Life Sci*; 58:1201-1207 (1996).
24. Taylor, P. Anticholinesterase agents in Goodman and Gilman's *The Pharmacological Basis of Therapeutics*, 8th Edition, Gilman, A. G., Rall, T.W., Nies, A. S. et al, eds. Pergamon Press, New York, 131-149 (1990).

25. Whitehouse, P.J., Price, D. L., Clark, A.W., Coyle, J.T., De Long, M.R., Alzheimer's Disease. Evidence for selective loss of cholinergic neurons in the nucleus basalis, Ann Neurol; 10 122-126(1981).
26. Product Monograph for ^{Pr}EXELON. Novartis Pharmaceuticals Canada Inc., Dorval, Quebec, Canada. Date of Revision: December 23, 2008.
27. A blinded, single-dose, randomized, crossover comparative bioavailability study of two formulations of Rivastigmine Hydrogen Tartrate capsules, NOVO-RIVASTIGMINE 6 mg capsule and EXELON[®] 6 mg capsules under fasting conditions. Data are on file at Novopharm Ltd.

PART III: CONSUMER INFORMATION**Pr NOVO-RIVASTIGMINE
Rivastigmine Capsules**

This leaflet is part III of a three-part “Product Monograph” published when NOVO-RIVASTIGMINE was approved for sale in Canada and is designed specifically for Consumers. This leaflet is a summary and will not tell you everything about NOVO-RIVASTIGMINE. Contact your doctor or pharmacist if you have any questions about the drug.

ABOUT THIS MEDICATION**What the medication is used for:**

Rivastigmine hydrogen tartrate is one of a group of drugs known as “cholinesterase inhibitors” which is used for the treatment of the symptoms of patients with mild to moderate Alzheimer’s disease or with dementia occurring at least 2 years following the diagnosis of Parkinson’s disease. Although there are differences between the two types of dementias in terms of changes to the brain and to mental function, it is known that with both conditions there are decreased levels of acetylcholine, a substance which is found in the brain and which is thought to be necessary for good cognitive function (memory and other mental function).

What it does:

Rivastigmine hydrogen tartrate works by increasing the amount of acetylcholine in the brain. In clinical studies with Rivastigmine, most patients with Alzheimer’s Disease had improved memory and other mental functions, or showed no further decline, as compared to placebo (sugar tablet) for up to 6 months. However, rivastigmine hydrogen tartrate may take as long as 12 weeks to begin working, and patient response to this medicine will vary. This medication should only be taken after proper diagnosis of your condition has been made by your doctor.

When it should not be used:

You should not take NOVO-RIVASTIGMINE:

- if you know that you are allergic to rivastigmine or to any of the other substances listed in this leaflet (see ‘**What the important nonmedicinal ingredients are**’),
- if you have ever had an allergic reaction to a similar type of medicine,
- if you have severe liver problems.

If this applies to you, tell your doctor without taking Novo-Rivastigmine.

NOVO-RIVASTIGMINE should only be used if prescribed to you by your doctor.

What the medicinal ingredient is:

NOVO-RIVASTIGMINE contains the active substance rivastigmine hydrogen tartrate.

What the important nonmedicinal ingredients are:

NOVO-RIVASTIGMINE capsules contain the following inactive substances: hydroxypropyl methylcellulose, magnesium stearate, microcrystalline cellulose and silicon dioxide. The capsule shell is made of gelatin with red and/or yellow iron oxide and titanium dioxide. The printing ink is based on red iron oxide.

What dosage forms it comes in:

Capsules: Each hard gelatin capsule contains 1.5, 3, 4.5 or 6 mg of rivastigmine as rivastigmine hydrogen tartrate.

WARNINGS AND PRECAUTIONS**BEFORE you use NOVO-RIVASTIGMINE talk to your doctor or pharmacist if:**

- if you have, or have ever had other medical problems, such as conditions affecting your heart or lungs;
- if you have, or have ever had seizures (fits or convulsions);
- if you have a history of stomach ulcers or have an increased risk of developing ulcers (for example you are taking non steroidal anti-inflammatory drugs (NSAIDs) or high doses of acetylsalicylic acid [ASA]);
- if you have, or have ever had difficulties in passing urine;
- if you have, or have ever had liver or kidney problems;
- if you are pregnant, planning on becoming pregnant, or breast feeding.

Your doctor will determine whether you can take NOVO-RIVASTIGMINE and how closely you will need to be monitored.

Can I drive vehicles and operate machinery?

Your doctor will tell you whether your illness allows you to drive vehicles and operate machinery safely.

INTERACTIONS WITH THIS MEDICATION

Make sure your doctor knows if you are taking, or begin to take, any other medicines, including drugs, or herbal (natural) products that you can buy without a prescription.

NOVO-RIVASTIGMINE should not be given at the same time as other medicines with similar effects (cholinomimetic agents) or with anticholinergic medicines (e.g. medicines used to relieve stomach cramps or spasms, to treat Parkinson’s disease or to prevent travel sickness).

Your doctor will tell you if you can also take NOVO-RIVASTIGMINE with your current medications. If you have to undergo surgery while taking NOVO-RIVASTIGMINE, you should inform the doctor before you are given any anaesthetics (drugs that produce a loss of sensation).

PROPER USE OF THIS MEDICATION

This medicine has been prescribed only for you. It must not be given to anybody else or used for any other illnesses.

NOVO-RIVASTIGMINE should be taken with food.

Capsules: Swallow the capsules whole with a drink, without opening or crushing them.

Usual dose

You must take NOVO-RIVASTIGMINE twice a day, once with your breakfast and once with your evening meal.

Your doctor will tell you what dosage of NOVO-RIVASTIGMINE to take, starting with a low dose and gradually increasing, depending on how you respond to the treatment. The highest dose that should be taken is 6 mg twice a day (12 mg/day).

To benefit from your medicine you must take it every day.

Overdose:

If more medication has been taken than what has been prescribed, contact either your doctor, hospital emergency department or the nearest, poison control center immediately. Some people who have accidentally taken too much rivastigmine hydrogen tartrate have experienced nausea, vomiting, diarrhea, high blood pressure and hallucinations. Slow heart beat and fainting may also occur. In case of accidental overdose, consult a physician immediately or contact your regional Poison Control Centre.

Missed Dose

If you find you have forgotten to take your dose of NOVO-RIVASTIGMINE do not worry, wait and take the next dose at the usual time. Do NOT take two doses at once.

If you stop taking NOVO-RIVASTIGMINE for more than several days, do NOT begin to take NOVO-RIVASTIGMINE again without contacting your doctor.

SIDE EFFECTS AND WHAT TO DO ABOUT THEM

In clinical studies most side effects of rivastigmine hydrogen tartrate were of mild to moderate intensity.

The most common side effects noted were:

- nausea, vomiting, diarrhea, heartburn, stomach pains and loss of appetite;
- dizziness, headache, sleepiness, trembling;
- agitation, confusion;
- weakness, fatigue, a general feeling of being unwell,
- sweating,
- weight loss

These side effects will most probably disappear gradually as your body becomes used to the medicine or with a reduction in dose. If they persist, however, you should tell your doctor.

Less frequently patients have experienced difficulty in sleeping, change in liver function tests, accidental falls, and very rarely, patients have experienced worsening of Parkinson’s disease symptoms (such as stiffness, difficulty in carrying out movements).

If you feel unwell in this or any other way or have any symptoms that you do not understand, or find distressing, you should contact your doctor immediately. If you experience severe adverse events and cannot contact your doctor, stop taking the drug until you can discuss your symptoms with your doctor

SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM				
Symptom / effect		Talk with your doctor or pharmacist		Stop taking drug and call your doctor or pharmacist
		Only if severe	In all cases	
Common	Heart attack (crushing chest pain)		√	
Uncommon	Fainting		√	
	Depression		√	
	Stroke (loss of coordination, difficulty in speaking or breathing and signs of brain disorder)			√
Rare	Chest pain		√	
	Seizures (fits or convulsion)		√	
	Rashes		√	
	Gastric (stomach) and duodenal (intestinal) ulcers		√	
Very rare	Blood in stools or when vomiting		√	
	Severe vomiting that can lead to rupture of the esophagus		√	
	Urinary tract infection		√	
	Inflammation of the		√	

	pancreas (severe upper stomach pain, often with nausea and vomiting)			
	Heart problems/fast, slow or irregular heart beats.		√	
	High blood pressure		√	
	Hallucinations		√	

This is not a complete list of side effects. For any unexpected effects while taking rivastigmine hydrogen tartrate, contact your doctor or pharmacist

HOW TO STORE IT

- Do not use NOVO-RIVASTIGMINE after the expiry date. Store NOVO-RIVASTIGMINE at room temperature (15 - 30°C) and protect from moisture.
- Keep NOVO-RIVASTIGMINE in a safe place and out of the reach of children.

REPORTING SUSPECTED SIDE EFFECTS

To monitor drug safety, Health Canada through the Canada Vigilance Program collects information on serious and unexpected side effects of drugs. If you suspect you have had a serious or unexpected reaction to this drug you may notify Canada Vigilance:

By toll-free telephone: 866-234-2345
 By toll-free fax: 866-678-6789
 Online: www.healthcanada.gc.ca/medeffect
 By email: CanadaVigilance@hc-sc.gc.ca

By regular mail:
 Canada Vigilance National Office
 Marketed Health Products Safety and Effectiveness Information Bureau
 Marketed Health Products Directorate
 Health Products and Food Branch
 Health Canada
 Tunney's Pasture, AL 0701C
 Ottawa ON K1A 0K9

NOTE: Should you require information related to the management of the side effect, please contact your health care provider before notifying Canada Vigilance. The Canada Vigilance Program does not provide medical advice.

MORE INFORMATION

This document plus the full product monograph, prepared for health professionals can be found by contacting Novopharm Limited at:
 1-800-268-4127 ext. 5005 (English Canada);
 1-877-777-9117 (French Canada)
 or druginfo@novopharm.com

This leaflet was prepared by:
 Novopharm Limited
 30 Novopharm Court
 Toronto, Ontario
 Canada, M1B 2K9

Date of Revision: April 1, 2009