

## PRODUCT MONOGRAPH

Pr**EPREX**\*

epoetin alfa

Sterile Solution

Single-use vials: 1,000 IU/0.5 mL, 2,000 IU/mL, 4,000 IU/mL, 10,000 IU/mL, 40,000 IU/mL

Multi-use vial: 20,000 IU/mL

Single-use pre-filled syringes with PROTECS\* needle guard:

1,000 IU/0.5 mL, 2,000 IU/0.5 mL,  
3,000 IU/0.3 mL, 4,000 IU/0.4 mL, 5,000 IU/0.5 mL, 6,000 IU/0.6 mL,  
7,000 IU/0.7 mL, 8,000 IU/0.8 mL, 9,000 IU/0.9 mL, 10,000 IU/mL,  
20,000 IU/0.5 mL, 30,000 IU/0.75 mL, 40,000 IU/mL

Erythropoiesis Regulating Hormone

This Product Monograph is the exclusive property of Janssen-Ortho Inc.  
It may not be copied in whole or in part without the written permission of Janssen-Ortho Inc.

Janssen-Ortho Inc.  
19 Green Belt Drive  
Toronto, Ontario  
M3C 1L9

Date of Authorization:  
July 23, 2009

[www.janssen-ortho.com](http://www.janssen-ortho.com)

Submission Control No: 129164

\* All trademark rights used under license

© 2009 JANSSEN-ORTHO Inc.

**Table of Contents**

**PART I: HEALTH PROFESSIONAL INFORMATION..... 3**  
SUMMARY PRODUCT INFORMATION ..... 3  
DESCRIPTION..... 4  
INDICATIONS AND CLINICAL USE..... 4  
CONTRAINDICATIONS ..... 5  
WARNINGS AND PRECAUTIONS..... 7  
ADVERSE REACTIONS..... 16  
DRUG INTERACTIONS ..... 23  
DOSAGE AND ADMINISTRATION..... 23  
OVERDOSAGE ..... 30  
ACTION AND CLINICAL PHARMACOLOGY ..... 30  
STORAGE AND STABILITY..... 34  
SPECIAL HANDLING INSTRUCTIONS ..... 34  
DOSAGE FORMS, COMPOSITION AND PACKAGING ..... 34

**PART II: SCIENTIFIC INFORMATION ..... 36**  
PHARMACEUTICAL INFORMATION..... 36  
CLINICAL TRIALS..... 36  
DETAILED PHARMACOLOGY ..... 48  
TOXICOLOGY ..... 53  
REFERENCES ..... 61

**PART III: CONSUMER INFORMATION..... 68**

**PrEPREX\***

epoetin alfa

Sterile Solution

Single-use vials: 1,000 IU/0.5 mL, 2,000 IU/mL, 4,000 IU/mL, 10,000 IU/mL, 40,000 IU/mL

Multi-use vial: 20,000 IU/mL

Single-use pre-filled syringes with PROTECS\* needle guard:

1,000 IU/0.5 mL, 2,000 IU/0.5 mL,  
3,000 IU/0.3 mL, 4,000 IU/0.4 mL, 5,000 IU/0.5 mL, 6,000 IU/0.6 mL,  
7,000 IU/0.7 mL, 8,000 IU/0.8 mL, 9,000 IU/0.9 mL, 10,000 IU/mL,  
20,000 IU/0.5 mL, 30,000 IU/0.75 mL, 40,000 IU/mL

Erythropoiesis Regulating Hormone

## PART I: HEALTH PROFESSIONAL INFORMATION

### SUMMARY PRODUCT INFORMATION

<b>Route of Administration</b>	<b>Dosage Form: Sterile Colourless Solution / Strength</b>	<b>Clinically Relevant Nonmedicinal Ingredients</b>
<b><u>Polysorbate-80 Containing (Human Serum Albumin [HSA]-free formulation)</u></b> Intravenous / Subcutaneous	<b><u>Single-Use Vial (Polysorbate-80 Containing [HSA-free] formulation)</u></b> 1,000 IU/0.5 mL, 2,000 IU/mL, 4,000 IU/mL, 10,000 IU/mL, 40,000 IU/mL <b><u>Single-Use Pre-filled Syringe with PROTECS needle guard (Polysorbate-80 Containing [HSA-free] formulation)</u></b> 1,000 IU/0.5 mL, 2,000 IU/0.5 mL, 3,000 IU/0.3 mL, 4,000 IU/0.4 mL, 5,000 IU/0.5 mL, 6,000 IU/0.6 mL, 7,000 IU/0.7 mL, 8,000 IU/0.8 mL, 9,000 IU/0.9 mL, 10,000 IU/mL, 20,000 IU/0.5 mL, 30,000 IU/0.75 mL, 40,000 IU/mL	<b><u>Single-Use Vials and Single-Use Pre-filled Syringes with PROTECS needle guard</u></b> Glycine, polysorbate 80, sodium chloride, sodium phosphate monobasic dihydrate, sodium phosphate dibasic dihydrate, in water for injection.
<b><u>HSA-Containing formulation:</u></b> Intravenous / Subcutaneous	<b><u>Multi-Use Vial (HSA-Containing formulation)</u></b> 20,000 IU/mL	<b><u>Multi-Use Vial</u></b> albumin (human) (0.25%), sodium chloride, sodium citrate, citric acid, benzyl alcohol (0.9%) as a preservative, in water for injection.

## DESCRIPTION

Erythropoietin is a glycoprotein which stimulates red blood cell production. It is produced in the kidney and stimulates the division and differentiation of committed erythroid progenitors in the bone marrow. EPREX epoetin alfa is a 165 amino acid glycoprotein manufactured by recombinant DNA technology. It has a molecular weight of 30,400 daltons and is produced by mammalian cells into which the human erythropoietin gene has been introduced. The product contains the identical amino acid sequence of isolated natural erythropoietin.

## INDICATIONS AND CLINICAL USE

EPREX epoetin alfa is indicated to elevate or maintain the red blood cell level (as manifested by the hematocrit or hemoglobin determinations) and to decrease the need for transfusions. EPREX therapy is not intended for patients who require immediate correction of severe anemia. EPREX epoetin alfa may obviate the need for maintenance transfusions but is not a substitute for emergency transfusion. Blood pressure should be adequately controlled prior to initiation of EPREX therapy and must be closely monitored and controlled during treatment. EPREX therapy is not indicated for other specific causes of anemia with established treatments such as iron or folate deficiencies, hemolysis or gastrointestinal bleeding which should be managed appropriately.

### **Treatment of Anemia of Chronic Renal Failure**

**Adults:** EPREX therapy is indicated in the treatment of anemia associated with chronic renal failure, including patients on dialysis (end-stage renal disease) and patients not on dialysis (see **DOSAGE AND ADMINISTRATION, CRF PATIENTS**). Non-dialysis patients with symptomatic anemia considered for therapy should have a hemoglobin less than 100 g/L.

**Pediatrics:** Epoetin alfa is indicated in infants and children from 1 month old up to 16 years of age for the treatment of anemia associated with CRF requiring dialysis. Safety and effectiveness in pediatric patients less than 1 month old have not been established (see **WARNINGS AND PRECAUTIONS** and *Product Monograph Part II, CLINICAL TRIALS, CRF PATIENTS, Pediatric Chronic Renal Failure Patients on Dialysis*).

### **Treatment of Anemia in Zidovudine-Treated/HIV-Infected Patients**

EPREX therapy is indicated for the treatment of transfusion-dependent anemia related to therapy with zidovudine in HIV-infected patients. EPREX epoetin alfa is effective in HIV-infected patients treated with zidovudine, when the endogenous serum erythropoietin level is  $\leq 500$  mU/mL and when patients are receiving a dose of zidovudine  $\leq 4200$  mg/week.

### **Treatment of Anemia Due to Chemotherapy in Patients With Non-Myeloid Malignancies**

EPREX is indicated for the treatment of anemia due to the effect of concomitantly administered chemotherapy based on studies that have shown a reduction in the need for RBC transfusions in patients with advanced or metastatic, non-myeloid malignancies receiving chemotherapy for a minimum of 2 months. Studies to determine whether EPREX increases mortality or decreases progression-free /recurrence-free survival are ongoing.

- In patients with a long life expectancy, the decision to administer ESAs should be based on a benefit-risk assessment with the participation of the individual patient. This should take into account the specific clinical context such as (but not limited to) the type of tumor and its stage, the degree of anemia, life expectancy, the environment in which the patient is being treated and known risks of transfusions and ESAs.
- If appropriate, red blood cell transfusion should be the preferred treatment for the management of anemia in patients with a long life expectancy and who are receiving myelosuppressive chemotherapy.
- EPREX is not indicated for use in patients receiving hormonal agents, therapeutic biologic products, or radiotherapy unless receiving concomitant myelosuppressive chemotherapy.

### **Surgery Patients**

EPREX therapy is indicated in the following elective surgery regimens:

#### **Use of EPREX epoetin alfa to Reduce Allogeneic Blood Exposure**

EPREX therapy is indicated to treat patients who are undergoing major elective surgery (including patients who do not wish to or are not eligible to participate in an autologous blood donation program) and have a pretreatment hemoglobin of  $> 100$  to  $\leq 130$  g/L. EPREX therapy is indicated to reduce allogeneic blood transfusions and hasten erythroid recovery in these patients.

#### **Combined Use of EPREX epoetin alfa and ABD**

EPREX epoetin alfa is indicated to facilitate autologous blood collection within a predeposit program and may decrease the risk of receiving allogeneic blood transfusions in patients with hemoglobin of 100-130 g/L who are scheduled for major elective surgery and are expected to require more blood than that which can be obtained through autologous blood collection techniques in the absence of epoetin alfa.

**Geriatrics (> 65 years of age):** No data available.

## **CONTRAINDICATIONS**

EPREX epoetin alfa is contraindicated in patients:

- who develop pure red cell aplasia (PRCA) following treatment with any erythropoiesis regulating hormone (see **WARNINGS AND PRECAUTIONS, ALL PATIENTS, Immune**);
- with uncontrolled hypertension;
- with known hypersensitivity to mammalian cell-derived products, albumin (human) or any component of the product (EPREX Sterile Solution is available in some formats, formulated without human serum albumin; see **DOSAGE FORMS, COMPOSITION AND PACKAGING**);
- who for any reason cannot receive adequate antithrombotic treatment.

In addition, the 20,000 IU/mL formulation, preserved with benzyl alcohol, is contraindicated in premature infants and newborns.

The use of EPREX epoetin alfa in patients scheduled for elective surgery and not participating in an autologous blood donation program is contraindicated in patients with severe coronary, peripheral arterial, carotid, or cerebral vascular disease, including patients with recent myocardial infarction or cerebral vascular accident.

Contraindications defined by the guidelines and methods of practice for ABD programs should be respected in patients receiving epoetin alfa.

## WARNINGS AND PRECAUTIONS

### Serious Warnings and Precautions

#### ALL PATIENTS

- To minimize the risks for death and serious cardiovascular events, follow the recommended dosage for each indication for EPREX and other erythropoiesis stimulating agents (ESAs) (see **WARNINGS AND PRECAUTIONS: Increased Mortality, Serious Cardiovascular and Thromboembolic Events** and **DOSAGE AND ADMINISTRATION**).
- In surgical patients treated with EPREX for reduction of allogeneic red blood cell transfusions, adequate antithrombotic prophylaxis, as per current standard of care, is recommended in order to reduce the incidence of deep venous thrombosis.
- The multi-use preserved formulation contains benzyl alcohol. Benzyl alcohol has been reported to be associated with an increased incidence of neurological and other complications in premature infants which are sometimes fatal.
- Patients with uncontrolled hypertension should not be treated with EPREX; blood pressure should be controlled adequately before initiation of therapy.
- EPREX should be used with caution in patients with a history of seizures.
- Antibody-mediated pure red cell aplasia (PRCA) has been reported after months to years of treatment with ESAs.

#### CHRONIC RENAL FAILURE PATIENTS

- Patients experienced greater risks for death and serious cardiovascular events when administered ESAs to target higher versus lower hemoglobin levels (135 vs. 113 g/L; 140 vs. 100 g/L) in two clinical studies. Individualize dosing to achieve and maintain hemoglobin levels within the range of 100 to 120 g/L.

#### CANCER PATIENTS

- ESAs increased the risks for death and serious cardiovascular and thromboembolic events in some controlled clinical trials.
- ESAs shortened overall survival and/or increased the risk of tumour progression or recurrence in some clinical studies in patients with breast, head and neck, lymphoid, cervical and non-small cell lung cancers when dosed to target a hemoglobin of  $\geq 120$  g/L.
- To minimize the above risks, use the lowest dose needed to avoid red blood cell (RBC) transfusions.
- Use ESAs only for treatment of anemia due to concomitant myelosuppressive chemotherapy.
- If appropriate, red blood cell transfusion should be the preferred treatment for the management of anemia in patients with a long life expectancy and who are receiving myelosuppressive chemotherapy.
- Discontinue EPREX following completion of a chemotherapy course.

## ALL PATIENTS

### General

#### Creutzfeldt-Jakob Disease (CJD)

Some formats of EPREX epoetin alfa contain albumin, a derivative of human blood. Because of careful donor screening and product manufacturing processes, the risk for transmission of viral diseases is extremely remote. The theoretical risk of transmission of Creutzfeldt-Jakob disease also is considered extremely remote. No cases of viral disease transmission or CJD have ever been identified as a result of albumin use.

#### Carcinogenicity, Mutagenicity

Long-term carcinogenicity studies have not been carried out. Epoetin alfa does not induce bacterial gene mutation (Ames Test), chromosomal aberrations in mammalian cells, micronuclei in mice, or gene mutation at the HGPRT locus.

### Cardiovascular

#### Hypertension

Patients with uncontrolled hypertension should not be treated with EPREX epoetin alfa; blood pressure should be controlled adequately before initiation of therapy. Blood pressure may rise during EPREX therapy, often during the early phase of treatment when the hemoglobin is increasing, especially in CRF patients.

For patients who respond to EPREX therapy with a rapid increase in hemoglobin (e.g., more than 10 g/L in any two-week period), the dose of EPREX epoetin alfa should be reduced because of the possible association of excessive rate of rise of hemoglobin with an exacerbation of hypertension.

All patients on EPREX epoetin alfa should have hematocrit/hemoglobin levels measured at least once a week until a stable level is achieved and periodically thereafter.

In all patients receiving EPREX epoetin alfa, blood pressure should be closely monitored and controlled as necessary. Particular attention should be paid to the development of unusual headaches or an increase in headaches as a possible warning signal.

It may be necessary to initiate or increase antihypertensive treatment during EPREX therapy. If blood pressure cannot be controlled, EPREX epoetin alfa should be discontinued until blood pressure control is re-established. Hypertensive encephalopathy and seizures have been observed in patients with CRF treated with EPREX (see **ADVERSE REACTIONS, ALL PATIENTS, Hypertension**).

### Immune

#### Pure Red Cell Aplasia

Cases of pure red cell aplasia (PRCA) and of severe anemia, with or without other cytopenias, associated with neutralizing antibodies to erythropoietin have been reported in patients treated with EPREX. This has been reported predominantly in CRF patients receiving EPREX by subcutaneous administration. Any patient who develops a sudden loss of response to EPREX, accompanied by severe anemia and low reticulocyte count should be evaluated for the etiology



of loss of effect, including the presence of neutralizing antibodies to erythropoietin. If anti-erythropoietin antibody-associated anemia is suspected, withhold EPREX and other erythropoietic proteins. Contact Janssen-Ortho Inc. at 1-800-567-3331 to perform assays for binding and neutralizing antibodies. EPREX should be permanently discontinued in patients with antibody-mediated anemia. Patients should not be switched to other erythropoietic proteins as antibodies may cross-react (see **ADVERSE REACTIONS**).

### **Increased Mortality, Serious Cardiovascular and Thromboembolic Events**

During hemodialysis, patients treated with EPREX epoetin alfa may require increased anticoagulation with heparin to prevent clotting of the artificial kidney. Clotting of the vascular access (A-V fistula) has occurred at an annualized rate of about 0.25 events per patient-year on EPREX therapy. Overall, for patients with CRF (whether on dialysis or not), other thrombotic events (e.g., myocardial infarction, cerebrovascular accident, transient ischemic attack) have occurred at an annualized rate of less than 0.04 events per patient-year of EPREX therapy. Patients with pre-existing vascular disease should be monitored closely.

Patients with CRF experienced greater risks for death and serious cardiovascular events when administered ESAs to target higher versus lower hemoglobin levels (135 vs. 113 g/L; 140 vs. 100 g/L) in two clinical studies. Patients with CRF and an insufficient hemoglobin response to ESA therapy may be at even greater risk for cardiovascular events and mortality than other patients. EPREX and other ESAs increased the risks for death and for serious cardiovascular and thromboembolic events in controlled clinical trials of patients with cancer. These events included myocardial infarction, stroke, congestive heart failure and an increased risk of serious arterial and venous thromboembolic events including hemodialysis vascular access thrombosis. A rate of hemoglobin rise of >10 g/L over 2 weeks may contribute to these risks.

CRF patients with hypo-responsiveness to ESAs may be at an increased risk for mortality and cardiovascular events. These patients should be evaluated for treatable conditions (see **WARNINGS AND PRECAUTIONS: Lack or Loss of Response** and **DOSAGE AND ADMINISTRATION: CRF PATIENTS**). These risks should be carefully weighed against the benefit to be derived from treatment with ESAs, particularly in cancer patients with increased risk factors of thrombotic vascular events (TVEs), such as patients with a prior history of TVEs (e.g. deep venous thrombosis or pulmonary embolism) (see *Product Monograph Part II, CLINICAL TRIALS* and **DOSAGE AND ADMINISTRATION, CRF PATIENTS; CANCER PATIENTS**).

To minimize the risks for death and serious cardiovascular events, EPREX and other ESAs should follow the recommended dose for each indication. For CRF patients, individualize dosing to achieve and maintain hemoglobin levels within the recommended range of 100-120 g/L. For patients with cancer, use the lowest dose sufficient to avoid blood transfusions. The hemoglobin concentrations should not exceed 120 g/L; the rate of hemoglobin increase should not exceed 10 g/L in any 2-week period (see **DOSAGE AND ADMINISTRATION**).

### **Hematologic**

The safety and efficacy of EPREX therapy have not been established in patients with underlying hematologic disease (e.g., sickle cell anemia, myelodysplastic syndromes, or hypercoagulable disorders).

## **Neurologic**

### **Seizures**

EPREX epoetin alfa should be used with caution in patients with a history of seizures. Additional close monitoring of all possible risk factors is advisable if the decision is made to use EPREX epoetin alfa to treat patients with a history of seizures.

Given the potential for an increased risk of seizures in CRF patients during the first 90 days of therapy, blood pressure and the presence of premonitory neurologic symptoms should be monitored closely, and CRF patients should be cautioned to avoid potentially hazardous activities such as driving or operating heavy machinery during this period.

It is recommended that the dose of EPREX epoetin alfa be decreased if the hemoglobin increase exceeds 10 g/L in any two-week period.

The safety and efficacy of EPREX therapy have not been established in patients with a known history of a seizure disorder.

### **Sensitivity/Resistance**

The parenteral administration of any biologic product should be attended by appropriate precautions in case allergic or other untoward reactions occur. If an anaphylactoid reaction occurs, EPREX epoetin alfa should be immediately discontinued and appropriate therapy initiated.

Hypersensitivity reactions, including cases of rash, urticaria, anaphylactic reaction, and angioneurotic edema have been reported.

### **Lack or Loss of Response**

Inadequate response to EPREX epoetin alfa should prompt an investigation for causative factors. If the patient fails to respond or to maintain a response, the following etiologies should be considered and evaluated:

1. Iron deficiency: Virtually all patients will eventually require supplemental iron therapy (see **Monitoring and Laboratory Tests, Iron Evaluation**);
2. Underlying infectious, inflammatory, or malignant processes;
3. Occult blood loss;
4. Underlying hematologic diseases (i.e., thalassemia, refractory anemia, or other myelodysplastic disorders);
5. Vitamin deficiencies: folic acid or vitamin B<sub>12</sub>;
6. Hemolysis;
7. Aluminum intoxication;
8. Osteitis fibrosa cystica;
9. Inflammatory or traumatic episodes;
10. Pure Red Cell Aplasia (PRCA).

### **Special Populations**

#### **Use in Patients with Known Porphyria:**

Exacerbation of porphyria has been observed rarely in CRF patients treated with EPREX epoetin alfa. However, EPREX epoetin alfa has not caused an increased urinary excretion of porphyrin

metabolites in normal volunteers, even in the presence of a rapid erythropoietic response. Nevertheless, EPREX epoetin alfa should be used with caution in patients with known porphyria.

### **Use in Patients with a History of Gout:**

Increased serum uric acid (and phosphorus) levels have been observed in both normal volunteers and dialysis independent CRF patients treated with EPREX epoetin alfa who experienced a rapid rate of rise of hemoglobin. This effect may be related to an increased rate of nucleic acid synthesis in the bone marrow. Consequently, EPREX epoetin alfa should be administered with caution to patients with a history of gout.

### **Hepatic Dysfunction:**

The safety of EPREX epoetin alfa has not been established in patients with hepatic dysfunction.

### **Use in Pregnancy:**

Although epoetin alfa has been shown to have adverse effects in rats when given in doses greater than five times the human dose, it is not known whether it can affect reproduction capacity or cause fetal harm when administered to pregnant women. EPREX epoetin alfa should be given to a pregnant woman only if potential benefit justifies the potential risk to the fetus.

In some female chronic renal failure patients, menses have resumed following EPREX therapy; the possibility of potential pregnancy should be discussed and the need for contraception evaluated.

### **Nursing Mothers:**

It is not known whether EPREX epoetin alfa is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when EPREX epoetin alfa is administered to a nursing woman.

### **Use in Children:**

#### **Pediatric Patients on Dialysis**

Epoetin alfa is indicated in infants and children from 1 month old up to 16 years of age for the treatment of anemia associated with CRF requiring dialysis. Safety and effectiveness in pediatric patients less than 1 month old have not been established (see *Product Monograph Part II, CLINICAL TRIALS, CRF PATIENTS, Pediatric Chronic Renal Failure Patients on Dialysis*). The safety data from these studies show that there is no increased risk to pediatric CRF patients on dialysis when compared to the safety profile of epoetin alfa in adult CRF patients (see **ADVERSE REACTIONS**). Published literature provides supportive evidence of the safety and effectiveness of epoetin alfa in pediatric patients on dialysis.

### **Monitoring and Laboratory Tests**

#### **Hematology**

All patients receiving EPREX epoetin alfa should have hematocrit/hemoglobin levels measured once a week until hematocrit/hemoglobin has been stabilized, and measured periodically thereafter (see **CHRONIC RENAL FAILURE PATIENTS, Monitoring and Laboratory Tests, Hematology** for additional laboratory monitoring in CRF patients).

There may be a moderate dose-dependent rise in the platelet count, within the normal range, during treatment with EPREX epoetin alfa. This regresses during the course of continued

therapy. In addition, thrombocytopenia above the normal range has been reported. The platelet count should be regularly monitored during the first eight weeks of therapy.

### **Iron Evaluation**

In most chronic renal failure, cancer, and HIV-infected patients, the serum ferritin concentrations fall concomitantly with the rise in packed cell volume. Therefore, prior to and during EPREX therapy, the patient's iron stores, including transferrin saturation (serum iron divided by iron binding capacity) and serum ferritin, should be evaluated. Transferrin saturation should be at least 20%, and serum ferritin levels should be at least 100 ng/mL. Supplemental iron, e.g., oral elemental iron or intravenous iron, is recommended to increase and maintain transferrin saturation to levels that will adequately support EPREX epoetin alfa-stimulated erythropoiesis.

All surgery patients being treated with EPREX epoetin alfa should receive adequate iron replacement throughout the course of therapy in order to support erythropoiesis and avoid depletion of iron stores (see **DOSAGE AND ADMINISTRATION**).

### **Vitamin B<sub>12</sub> and Folate Evaluation**

Prior to starting EPREX therapy, the patient's serum vitamin B<sub>12</sub> and serum folate should be assessed. A deficiency in vitamin B<sub>12</sub> and/or folate may blunt the response and should be investigated as per standard clinical practice.

## **CHRONIC RENAL FAILURE (CRF) PATIENTS**

### **General**

Hemoglobin levels during treatment with ESAs should not exceed 120 g/L in both men and women. The dose of EPREX epoetin alfa should be reduced as the hemoglobin approaches 120 g/L or increases by more than 10 g/L in any 2-week period. Patients with CRF experienced greater risks for death and serious cardiovascular events when administered ESAs to target higher versus lower hemoglobin levels (135 vs. 113 g/L; 140 vs. 100 g/L) in two clinical studies. Individualize dosing to achieve and maintain hemoglobin levels within the range of 100 to 120 g/L (see **DOSAGE AND ADMINISTRATION**).

### **Renal**

#### **Dialysis Management**

Therapy with EPREX epoetin alfa results in an increase in hemoglobin and a decrease in plasma volume, which could potentially affect dialysis efficiency. In studies to date, the resulting increase in hemoglobin did not appear to adversely affect dialyzer function or the efficiency of high flux hemodialysis. During hemodialysis, chronic renal failure patients treated with EPREX epoetin alfa may require increased anticoagulation with heparin to prevent clotting of the artificial kidney.

Chronic renal failure patients who are marginally dialyzed may require adjustments in their dialysis prescription. As with all patients on dialysis, the serum chemistry values [including blood urea nitrogen (BUN), creatinine, phosphorus, and potassium] in patients treated with EPREX epoetin alfa should be monitored regularly to assure the adequacy of the dialysis prescription.

## **Diet**

As the hemoglobin increases and patients experience an improved sense of well-being and quality of life, the importance of compliance with dietary and dialysis prescriptions should be reinforced. In particular, hyperkalemia is not uncommon in patients with CRF.

## **Predialysis Management**

Blood pressure and hemoglobin should be monitored in predialysis patients no less frequently than for ESRD patients maintained on dialysis. Renal function and fluid and electrolyte balance should be closely monitored, as an improved sense of well-being may obscure the need to initiate dialysis in some patients.

Based on information to date, the use of EPREX epoetin alfa in predialysis patients does not accelerate the rate of progression of renal insufficiency.

## **Monitoring and Laboratory Tests**

### **Hematology**

Sufficient time should be allowed to determine a patient's responsiveness to a dosage of EPREX epoetin alfa before adjusting the dose. The hemoglobin should be determined weekly until it has stabilized in the recommended range and the maintenance dose has been established. After any dose adjustment, the hemoglobin should also be measured weekly for at least 2-6 weeks until it has been determined that the hemoglobin has stabilized in response to the dose change. The hemoglobin should then be monitored at regular intervals.

A complete blood count with differential and platelet count should be performed regularly. During clinical trials, modest increases were seen in platelets and white blood cell counts. While these changes were statistically significant, they were not clinically significant and the values remained within normal ranges.

In order to avoid reaching or exceeding the recommended hemoglobin range of 100-120 g/L too rapidly, the guidelines for dose and frequency of dose adjustments (see **DOSAGE AND ADMINISTRATION**) should be followed.

The elevated bleeding time characteristic of chronic renal failure (CRF) decreases toward normal after correction of anemia in patients treated with EPREX epoetin alfa. Reduction of bleeding time also occurs after correction of anemia by transfusion.

### **Biochemistry**

In patients with CRF, serum chemistry values [including blood urea nitrogen (BUN), uric acid, creatinine, phosphorus, and potassium] should be monitored regularly. During clinical trials in patients on dialysis, modest increases were seen in BUN, creatinine, phosphorus, and potassium. In some CRF patients not on dialysis who were treated with EPREX epoetin alfa, modest increases in serum uric acid and phosphorus were observed. While changes were statistically significant, the values remained within the ranges normally seen in patients with CRF.

In adult patients with CRF not on dialysis, renal function and fluid and electrolyte balance should be closely monitored, as an improved sense of well-being may obscure the need to initiate dialysis in some patients. In patients with CRF not on dialysis, placebo-controlled studies of the progression of renal dysfunction over periods of greater than one year have not been completed.

In shorter-term trials in adult patients with CRF not on dialysis, changes in creatinine and creatinine clearance were not significantly different in patients treated with EPREX epoetin alfa, compared with placebo-treated patients. Analysis of the slope of 1/serum creatinine vs. time plots in these patients indicates no significant change in the slope after the initiation of EPREX therapy.

## **SURGERY PATIENTS**

### **General**

#### **Combined Use of EPREX epoetin alfa and ABD**

Warnings and precautions defined by the guidelines and methods of practice for ABD programs should be respected in patients receiving epoetin alfa.

#### **Use of EPREX epoetin alfa to Reduce Allogeneic Blood Exposure**

##### **Thrombotic/Vascular Events**

In patients scheduled for major elective orthopedic surgery, thrombotic events can be a risk and this possibility should be carefully weighed against the benefit to be derived from the treatment in this patient group. Patients scheduled for elective surgery should receive adequate antithrombotic prophylaxis as per current standard of care.

An increased incidence of deep vein thrombosis (DVT) was observed in patients receiving epoetin alfa who were undergoing spinal surgery and not receiving prophylactic anticoagulation. Patients should receive adequate antithrombotic prophylaxis in order to reduce the incidence of DVT (see *Product Monograph Part II, CLINICAL TRIALS, SURGERY PATIENTS, Use of EPREX epoetin alfa to Reduce Allogeneic Blood Exposure*).

In a randomized, placebo-controlled study of epoetin alfa in adult patients who were undergoing coronary artery bypass surgery and not participating in an autologous blood donation program, increased mortality was observed (7 deaths in 126 patients randomized to epoetin alfa versus no deaths among 56 patients receiving placebo). Among the seven deaths in the epoetin alfa-treated patients, three were related to intercurrent infectious episodes and four of these deaths occurred during the period of study drug administration. All four deaths were associated with thrombotic events and a causative role for epoetin alfa cannot be excluded.

### **Cardiovascular**

Rarely, blood pressure may rise in the perioperative period in patients being treated with EPREX epoetin alfa. Therefore, blood pressure should be monitored.

##### **Thrombotic/Vascular Events**

Independent of EPREX treatment, thrombotic and vascular events may occur in surgical patients with underlying cardiovascular disease following repeated phlebotomy. Therefore, routine volume replacement should be performed in such patients in autologous blood donation programs.

## CANCER PATIENTS

### **Increased Mortality and/or Increased Risk of Tumour Progression or Recurrence**

ESAs, when administered to target a hemoglobin of  $> 120$  g/L, shortened the time to tumour progression in patients with advanced head and neck cancer receiving radiation therapy. ESAs also shortened survival in patients with metastatic breast cancer and in patients with lymphoid malignancy receiving chemotherapy when administered to target a hemoglobin of  $\geq 120$  g/L.

In addition, ESAs shortened survival in patients with non-small cell lung cancer and in a study enrolling patients with various malignancies who were not receiving chemotherapy or radiotherapy; in these two studies, ESAs were administered to target a hemoglobin level of  $\geq 120$  g/L.

An increased risk of death was observed in a clinical study when ESAs were administered to target hemoglobin of 120 g/L in patients with active malignant disease not being treated with either chemotherapy or radiation therapy. EPREX is not indicated for use in cancer patients who have anemia that is not associated with chemotherapy (see ***Product Monograph Part II, CLINICAL TRIALS, CANCER PATIENTS, Tumour Progression, Increased Mortality and Thromboembolic Events***).

In view of the above, where appropriate red blood cell transfusion should be the preferred treatment for the management of anemia in patients with cancer. The decision to administer ESAs should be based on a benefit-risk assessment with the participation of the individual patient. This should take into account the specific clinical context such as (but not limited to) the type of tumor and its stage, the degree of anemia, life-expectancy, the environment in which the patient is being treated, and known risks of transfusion and ESAs.

### **Neurologic**

#### **Seizures**

In a placebo-controlled, double-blind trial utilizing once weekly dosing with epoetin alfa, 1.2% (n = 2/168) of safety-evaluable patients treated with epoetin alfa and 0% (n = 0/165) of placebo-treated patients had seizures. Seizures in the patients treated with weekly epoetin alfa occurred in the context of a significant increase in hemoglobin from baseline values but significant increases in blood pressure were not seen. Both patients may have had other CNS pathology which may have been related to the seizures.

## **ADVERSE REACTIONS**

### **ALL PATIENTS**

#### **Hypertension**

The most frequent adverse reaction during treatment with EPREX epoetin alfa is a dose-dependent increase in blood pressure or aggravation of existing hypertension. This occurred most commonly in chronic renal failure patients (see **WARNINGS AND PRECAUTIONS**).

Hypertensive crises with encephalopathy and seizures have occurred in isolated patients, including previously normotensive patients.

#### **Thrombotic/Vascular Events**

Serious adverse drug reactions include venous and arterial thromboses and embolism (including some with fatal outcomes), such as deep venous thrombosis, arterial thrombosis, pulmonary emboli, aneurysms, retinal thrombosis, clotting of vascular access (A-V fistula) and shunt thrombosis (including dialysis equipment). Additionally, cerebrovascular accidents (including cerebral infarction and cerebral hemorrhage) and transient ischemic attacks have been reported in patients receiving epoetin alfa.

In clinical studies conducted in surgery patients with a pretreatment hemoglobin of  $> 100$  to  $\leq 130$  g/L (the recommended population) and not participating in an ABD program, the rate of deep venous thrombosis (DVT) was similar among patients treated with EPREX epoetin alfa and placebo. However, in patients with a pretreatment hemoglobin of  $>130$  g/L, the rate of DVTs was higher in the group treated with EPREX epoetin alfa than in the placebo-treated group, but within the range of that reported in the literature for orthopedic surgery patients (47-74% without anticoagulant therapy and 3-37% with use of anticoagulant therapy).

In a study examining the use of EPREX epoetin alfa in 182 patients scheduled for coronary artery bypass graft surgery, 23% of patients treated with EPREX epoetin alfa and 29% treated with placebo experienced thrombotic/vascular events. There were four deaths among the patients treated with EPREX epoetin alfa that were associated with a thrombotic/vascular event and a causative role of EPREX epoetin alfa cannot be excluded.

#### **Influenza-like Illness**

Influenza-like illness including headaches, joint pains, myalgia and pyrexia may occur, especially at the start of treatment.

#### **Hypersensitivity Reactions**

Hypersensitivity reactions, including cases of rash, urticaria, angioneurotic edema and anaphylactic reaction have been reported.

#### **Immune Reactions**

In post-marketing reports, very rare cases of pure red cell aplasia (PRCA) have been reported predominantly in chronic renal failure patients after months to years of treatment with recombinant erythropoietins including EPREX. Most of these reports have been associated with the subcutaneous route of administration. Cases have been rarely reported in patients with hepatitis C treated with interferon and ribavirin, when ESAs are used concomitantly. ESAs are



not authorized in the management of anaemia associated with hepatitis C. Neutralizing antibodies to erythropoietins have been reported in most of these patients (see **WARNINGS AND PRECAUTIONS, ALL PATIENTS, Immune**).

**Seizures**

Seizures have been reported in patients treated with EPREX epoetin alfa.

**CRF Patients**

Studies analyzed to date indicate that EPREX epoetin alfa is generally well tolerated irrespective of the route of administration. The adverse events reported are frequent sequelae of CRF and are not necessarily attributable to EPREX therapy. In double-blind, placebo-controlled studies involving 335 patients with CRF (both predialysis and treated with dialysis), the events reported in greater than 5% of patients treated with EPREX epoetin alfa (n=200) during the blinded phase were:

**Table 1.1**

Adverse Events	EPREX epoetin alfa (n=200)	Placebo (n=135)
Hypertension	24.0%	18.5%
Headache	16.0%	11.9%
Arthralgias	11.0%	5.9%
Nausea	10.5%	8.9%
Edema	9.0%	10.4%
Fatigue	9.0%	14.1%
Diarrhea	8.5%	5.9%
Vomiting	8.0%	5.2%
Chest Pain	7.0%	8.8%
Skin Reaction at Administration Site	7.0%	11.9%
Asthenia	7.0%	11.9%
Dizziness	7.0%	12.6%
Clotted Access	6.8%	2.3%

**Pediatric CRF Patients on Dialysis:**

Epoetin alfa is generally well tolerated in pediatric dialysis patients. The adverse events reported are typical sequelae for patients on dialysis, and relationship to epoetin alfa therapy has not been established. In double-blind clinical studies involving 123 pediatric dialysis patients, only one event, myalgia, occurred with a statistically significant higher frequency in patients treated with epoetin alfa (20%) than in placebo-treated patients (6%), p=0.03. Adverse events reported in greater than 5% of patients treated with epoetin alfa were:

**Table 1.2**

<b>Adverse Events</b>	<b>Epoetin alfa (n=59)</b>	<b>Placebo (n=64)</b>
Hypertension	25.4%	17.2%
Vomiting	22.0%	21.9%
Headache	22.0%	29.7%
Myalgia	20.3%	6.3%
Abdominal Pain	20.3%	25.0%
Access Infection	20.3%	12.5%
Peritonitis	16.9%	14.1%
Fever	13.6%	15.6%
Pruritus	11.9%	3.1%
Upper Respiratory Infection	11.9%	14.1%
Hyperkalemia	11.9%	10.9%
Nausea	11.9%	12.5%
Pharyngitis	10.2%	7.8%
Cough	10.2%	9.4%
Constipation	10.2%	6.3%
Chest Pain	10.2%	3.1%
Access Complication	10.2%	6.3%
Dizziness	8.5%	10.9%
Convulsions	8.5%	6.3%
Hypotension	8.5%	14.1%
Influenza-like Symptoms	8.5%	6.3%
Access Catheter Clotted	8.5%	1.6%
Thrombosis Vascular Access	6.8%	1.6%
Pneumonia	6.8%	3.1%
Increased Nonprotein Nitrogen	5.1%	0.0%
Arthralgia	5.1%	3.1%
Hyperphosphatemia	5.1%	1.6%
Otitis	5.1%	1.6%
Malaise	5.1%	7.8%

Adverse events reported at > 5% in pediatric patients (Table 1.2) that were not previously reported in adult patients include: myalgia, abdominal pain, access infection, peritonitis, fever, pruritus, upper respiratory infection, hyperkalemia, pharyngitis, cough, constipation, access complication, convulsions, hypotension, influenza-like symptoms, thrombosis vascular access, pneumonia, increased nonprotein nitrogen, hyperphosphatemia, otitis, malaise.

The following adverse experiences have been reported at an incidence greater than 1% and less than 5%. No adverse events were reported at an incidence of <1%.

**Application Site:** injection site reaction

**Body as a Whole:** rigors, organ transplant rejection, dialysis complication, syncope, incisional pain, hot flushes, fatigue, edema peripheral, edema periorbital, edema circumoral, access pain, access erythema, access edema, access drainage, access cellulitis

**Cardiovascular:** hypertension aggravated, cardiac failure

**CNS/PNS:** vertigo, somnolence, paresthesia, migraine, encephalopathy hypertensive

**Endocrine Disorders:** glucocorticoids increased

**Gastrointestinal:** hemorrhage GI, increased saliva, gastroenteritis, gastrointestinal disorder, gastritis, flatulence, dyspepsia, diarrhea, anorexia

**Hearing/Vestibular:** otitis media, earache

**Heart Rate/Rhythm:** cardiac arrest

**Hematologic:** thrombocythemia, lymphedema, ecchymosis

**Liver and Biliary:** infectious hepatitis

**Metabolic/Nutrition:** hypokalemia, hypocalcemia, hypercalcemia, SGPT increased, hypertriglyceridemia

**Musculo-Skeletal:** limb pain, back pain, skeletal pain, musculo-skeletal disorder, muscle weakness, arthrosis

**Myo/Endo/Pericardial:** heart murmur, heart disorder

**Psychiatric Disorder:** hallucination

**Reproductive:** vaginitis (female), breast mass (male)

**Resistance Mechanism:** sepsis, infection, viral infection, fungal infection

**Respiratory:** wheezing, upper respiratory tract congestion, sinusitis, epistaxis, dyspnea, bronchospasm, atelectasis

**Skin and Appendages:** rash, dry skin, maculo-papular rash, folliculitis, erythema, cellulitis, acne

**Urinary Disorders:** uremia, urinary tract infection, albuminuria

**Vascular Disorders:** superior vena cava syndrome

**Vision Disorders:** conjunctivitis, abnormal vision

#### **ZIDOVUDINE-TREATED/HIV-INFECTED PATIENTS**

Adverse experiences reported in clinical trials with EPREX epoetin alfa in zidovudine-treated/HIV-infected patients were consistent with the progression of HIV infection. In double-blind, placebo-controlled studies of 3-month duration involving approximately 300 zidovudine-treated/HIV-infected patients, adverse experiences with an incidence of  $\geq 10\%$  in either patients treated with EPREX epoetin alfa (n=144) or placebo-treated patients were:

**Table 1.3**

Adverse Experience	EPREX epoetin alfa (n=144)	Placebo (n=153)
Pyrexia	38%	29%
Fatigue	25%	31%
Headache	19%	14%
Cough	18%	14%
Diarrhea	16%	18%
Rash	16%	8%
Congestion, Respiratory	15%	10%
Nausea	15%	12%
Shortness of Breath	14%	13%
Asthenia	11%	14%
Skin Reaction at Administration Site	10%	7%
Dizziness	9%	10%

There were no statistically significant differences between treatment groups in the incidence of the above events.

EPREX epoetin alfa does not appear to potentiate progression of HIV disease as measured by: incidence of opportunistic infections; mortality; serum p24 antigen levels; or HIV replication in infected cell lines in vitro.

### CANCER PATIENTS

Adverse experiences reported in clinical trials with EPREX epoetin alfa administered thrice weekly in cancer patients were consistent with the underlying disease state. In double-blind, placebo-controlled studies of up to three months duration involving 413 cancer patients, adverse events with an incidence  $\geq 10\%$  in either patients treated with EPREX epoetin alfa or placebo-treated patients were:

**Table 1.4**

Adverse Experience	EPREX epoetin alfa (n=213)	Placebo (n=200)
Nausea	23%	29%
Pyrexia	22%	21%
Asthenia	17%	16%
Fatigue	15%	20%
Vomiting	15%	18%
Diarrhea	15%	9%
Edema	14%	8%
Dizziness	10%	9%
Skin Reaction at Administration Site	10%	10%
Constipation	10%	9%
Shortness of Breath	8%	15%*
Decreased Appetite	8%	12%
Trunk Pain	8%	12%
Chills	7%	10%

\*Significantly higher incidence for placebo patients (p=0.030).

There were no statistically significant differences in the percentage of patients treated with EPREX epoetin alfa reporting these adverse events compared to the corresponding incidence in placebo-treated patients, except for shortness of breath which occurred at a higher incidence in placebo-treated patients than in patients treated with EPREX epoetin alfa.

Thrombotic vascular events can occur in cancer patients as a consequence of their disease, comorbidities, and treatment. An increased incidence of thromboembolic events has been reported in cancer patients receiving ESAs, including epoetin alfa.

In a placebo-controlled, double-blind trial utilizing once-weekly (QW) dosing with epoetin alfa for up to 4 months involving 333 safety-evaluable cancer patients, adverse events were reported using the NCI Common Toxicity Criteria. Adverse events with an incidence > 10% in either patients treated with epoetin alfa or placebo-treated patients were as indicated in the table below. The safety profile of epoetin alfa administered QW was similar to placebo; adverse events in both groups appeared consistent with events expected in advanced cancer or associated with cancer treatment.

**Table 1.5**

<b>Percent of Patients in the Weekly Dosing Study Reporting Event</b>		
<b>CTC Category Adverse Experience (all grades)</b>	<b>epoetin alfa (n=168)</b>	<b>Placebo (n=165)</b>
Fatigue	51%	52%
Nausea	35%	30%
Alopecia	25%	29%
Neurosensory	23%	23%
Vomiting	20%	16%
Diarrhea - no colostomy	20%	22%
Anorexia	18%	19%
Constipation	18%	22%
Dyspnea	16%	22%
Pain	9%	11%
Anemia	8%	13%

Based on comparable survival data and on the percentage of patients treated with epoetin alfa QW and placebo-treated patients who discontinued therapy due to death (7% vs 5%), disease progression (7% vs 8%), or adverse experiences (1% vs 1%), the clinical outcome in patients treated with epoetin alfa and placebo-treated patients appeared to be similar.

## **SURGERY PATIENTS**

### **Use of EPREX epoetin alfa to Reduce Allogeneic Blood Exposure**

Adverse events were combined for all groups treated with EPREX epoetin alfa and the placebo-treated groups from four orthopedic surgery studies where subjects received EPREX epoetin alfa at a dose of 300 or 100 IU/kg daily, 600 IU/kg weekly, or placebo. Adverse events reported by at least 10% of subjects in any treatment group were:

**Table 1.6: Percent of Patients Reporting Event<sup>a</sup>**

Adverse Event	300 <sup>b</sup> or 100 <sup>c</sup> IU/kg	600 <sup>d</sup> IU/kg	Placebo (n=250)
	EPREX epoetin alfa (Daily) (n=546)	EPREX epoetin alfa (Weekly) (n=73)	
Pyrexia	45%	47%	52%
Skin reaction, injection site	42%	26%	40%
Nausea	37%	45%	34%
Constipation	34%	51%	32%
Vomiting	17%	21%	11%
Skin pain	17%	5%	19%
Insomnia	14%	21%	11%
Headache	13%	10%	9%
Pruritus	12%	14%	10%
Dizziness	10%	11%	9%
Diarrhea	9%	10%	9%
Urinary tract infection	8%	11%	10%
Edema	8%	11%	8%
Arthralgia	8%	10%	6%
Urinary retention	7%	11%	7%
Confusion	5%	12%	6%
Flatulence	4%	10%	4%
Anxiety	3%	11%	6%

<sup>a</sup> All patients participating in orthopedic surgery studies regardless of baseline hemoglobin

<sup>b</sup> 300 IU/kg daily for either 5 or 10 days prior to surgery, on the day of surgery and either 3 or 4 days following surgery (either 9, 14 or 15 daily doses)

<sup>c</sup> 100 IU/kg daily for 10 days prior to surgery, on the day of surgery and 4 days following surgery (15 daily doses)

<sup>d</sup> 600 IU/kg once a week beginning 3 weeks prior to surgery and on the day of surgery (4 weekly doses)

Similar proportions of patients treated with EPREX epoetin alfa and placebo-treated patients reported each adverse event.

### **Combined Use of EPREX epoetin alfa and ABD**

The incidence of adverse events was calculated across five double-blind, placebo-controlled studies and one single-blind study, combining all patients treated with EPREX epoetin alfa (n=402), regardless of dose administered, and all placebo patients (n=242). Adverse experiences with an incidence of  $\geq 5\%$  in either patients treated with EPREX epoetin alfa or placebo-treated patients were:

**Table 1.7**

Event	EPREX epoetin alfa (n = 402)	Placebo (n = 242)
Fatigue	18.41%	19.01%
Dizziness	12.19%	13.64%
Nausea	11.44%	9.09%
Headache	9.20%	11.98%
Asthenia	5.47%	3.72%
Diarrhea	3.48%	7.02%

In general, there were no notable differences between patients treated with EPREX epoetin alfa and placebo-treated patients in the incidence of any adverse event.

## DRUG INTERACTIONS

### Overview

No evidence exists that indicates that treatment with EPREX epoetin alfa alters the metabolism of other drugs. However, since cyclosporine is bound by red blood cells, there is potential for a drug interaction. If EPREX epoetin alfa is given concomitantly with cyclosporine, blood levels of cyclosporine should be monitored and the dose of cyclosporine adjusted as necessary.

### Drug-Drug Interactions

Interactions with other drugs have not been established.

### Drug-Food Interactions

Interactions with food have not been established.

### Drug-Herb Interactions

Interactions with herbal products have not been established.

### Drug-Laboratory Interactions

Interactions with laboratory tests have not been established.

## DOSAGE AND ADMINISTRATION

### Dosing Considerations

**IMPORTANT: See WARNINGS AND PRECAUTIONS: SERIOUS WARNINGS AND PRECAUTIONS Box, and Increased Mortality, Serious Cardiovascular and Thromboembolic Events. Follow the recommended dosage for each indication for EPREX.**

EPREX may be given either as an IV or SC injection (see **Recommended Dose and Dosage Adjustment; CRF PATIENTS**).

### Self-Administration and Information for the Patient

In those situations in which the physician determines that a patient can safely and effectively self-administer EPREX epoetin alfa injection, the patient should be instructed as to the proper dosage and administration. The first few doses should be administered under supervision. Following the initial laboratory and clinical assessment, all patients, including those deemed capable of self-administration, should be monitored for their response to EPREX epoetin alfa, their blood pressure and serum levels as indicated in the **WARNINGS AND PRECAUTIONS** section. Patients should be referred to the full **Product Monograph PART III: CONSUMER INFORMATION** section. It is not a disclosure of all possible effects. Patients should be informed of the signs and symptoms of allergic drug reaction and advised of appropriate actions.

If home use is prescribed, the patient should be thoroughly instructed in the importance of proper disposal and cautioned against the reuse of needles, syringes, or drug product. A puncture-resistant container for the disposal of used syringes and needles should be available to the patient. The full container should be disposed of according to the directions provided by the physician.

## **Recommended Dose and Dosage Adjustment**

### **CRF PATIENTS**

The recommended dose to initiate therapy of EPREX epoetin alfa is 50 to 100 IU/kg three times per week (TIW) for adult patients. The recommended starting dose for pediatric CRF patients on dialysis is 50 IU/kg three times weekly (TIW). The dose of EPREX epoetin alfa should be reduced as the hemoglobin approaches 120 g/L or increases by more than 10 g/L in any 2-week period.

The dosage of EPREX epoetin alfa should be individualized to achieve and maintain the hemoglobin within the recommended range of 100-120 g/L. If hemoglobin excursions outside the recommended range occur, EPREX dose should be adjusted using the dosing recommendations below.

It should be recognized that subcutaneous administration of recombinant human proteins may increase the risk of immunogenicity. In patients on hemodialysis, the IV route is recommended.

When used intravenously, EPREX Sterile Solution usually has been administered as a slow IV bolus three times per week. While the administration of EPREX epoetin alfa is independent of the dialysis procedure, EPREX solution may be administered into the venous port at the end of the dialysis procedure to obviate the need for additional venous access. Following the change from intravenous to subcutaneous route of administration, the patient should be monitored carefully to ensure that the hemoglobin response is appropriate. Available data suggests that patients with a baseline hemoglobin < 60 g/L may require higher maintenance doses than those with a baseline hemoglobin > 80 g/L. EPREX therapy is not intended for patients who require immediate correction of severe anemia.

During therapy, hematological parameters should be monitored regularly.

Virtually all patients will eventually require supplemental iron to increase or maintain transferrin saturation to levels that will adequately support EPREX epoetin alfa-stimulated erythropoiesis.

Table 1.8 provides general therapeutic guidelines.



**Table 1.8**

Starting Dose	Reduce Dose When (Starting with 25 IU/kg/dose decrement)	Increase Dose If (Starting with 25 IU/kg/dose increment)	Maintenance Dose	Recommended Hb Range
Adults: 50-100 IU/kg three times per week IV or SC	1) Hb approaches 120 g/L or 2) Hb increases >10 g/L in any 2-week period.	Hb does not increase by 10 g/L after 8-12 weeks of therapy, and Hb is below range.	Individually titrate.	100-120 g/L (and not to be exceeded)
Pediatric patients: 50 IU/kg three times per week IV or SC				

In patients with chronic renal failure, maintenance hemoglobin concentration should not exceed 120 g/L.

### **Dose Adjustment**

Following EPREX therapy, a period of time is required for erythroid progenitors to mature and be released into circulation resulting in an eventual increase in hemoglobin. Additionally, red blood cell survival time affects hemoglobin and may vary due to uremia. As a result, the time required to elicit a clinically significant change in hemoglobin (increase or decrease) following any dose adjustment may be 2-6 weeks.

Dose adjustment should not be made more frequently than once a month, unless clinically indicated. After any dose adjustment, the hemoglobin should be determined weekly for at least 2-6 weeks (see **WARNINGS AND PRECAUTIONS, CHRONIC RENAL FAILURE (CRF) PATIENTS, Monitoring and Laboratory Tests**).

- If the hemoglobin is increasing and approaching 120 g/L, the dose should be decreased by approximately 25 IU/kg three times per week. If the reduced dose does not stop the rise in hemoglobin, doses should be temporarily withheld until the hemoglobin begins to decrease, at which point therapy should be re-initiated at a lower dose.
- At any time, if the hemoglobin increases by more than 10 g/L in a two-week period, the dose should be immediately decreased. After the dose reduction, the hemoglobin should be monitored weekly for 2-6 weeks, and further dose adjustments should be made (see **Maintenance Dose**).
- If a hemoglobin increase of 10 g/L is not achieved after an eight-week period and iron stores are adequate (see **Lack or Loss of Response** in this section), the dose of EPREX epoetin alfa may be increased in increments of 25 IU/kg three times per week. Further increases of 25 IU/kg three times per week may be added at 4-6 week intervals until the desired response is attained.

### **Maintenance Dose**

The maintenance dose must be individualized for each chronic renal failure patient.

If the hemoglobin remains below, or falls below, the recommended range, iron stores should be re-evaluated. If the transferrin saturation is less than 20%, supplemental iron should be administered. If the transferrin saturation is greater than 20%, the dose of EPREX epoetin alfa may be increased by 25 IU/kg three times per week. Such dose increases should not be made more frequently than once a month, unless clinically indicated, as the response time of the hemoglobin to a dose increase can be 2-6 weeks. Hemoglobin should be measured for 2-6 weeks following dose increases.

### **Lack or Loss of Response**

Over 95% of patients with chronic renal failure responded with clinically significant increases in hemoglobin; virtually all patients were transfusion-independent within approximately two months of initiation of EPREX therapy.

If a patient fails to respond or maintain a response, other etiologies should be considered and evaluated as clinically indicated (see **WARNINGS AND PRECAUTIONS**).

For patients whose hemoglobin does not attain a level within the range of 100 to 120 g/L despite the use of appropriate EPREX dose titrations over a 12-week period:

- do not administer higher EPREX doses and use the lowest dose that will maintain a hemoglobin level sufficient to avoid the need for recurrent RBC transfusions,
- evaluate and treat for other causes of anemia (see **WARNINGS AND PRECAUTIONS: Lack or Loss of Response**), and
- thereafter, hemoglobin should continue to be monitored and if responsiveness improves, EPREX dose adjustments should be made as described above; discontinue EPREX if responsiveness does not improve and the patient needs recurrent RBC transfusions.

### **ZIDOVUDINE-TREATED/HIV-INFECTED PATIENTS**

Prior to beginning EPREX therapy, it is recommended that the endogenous serum erythropoietin level be determined (prior to transfusion). Available evidence suggests that patients receiving zidovudine with levels > 500 mU/mL are unlikely to respond to therapy with EPREX epoetin alfa unless the dose of zidovudine is reduced or temporarily stopped.

### **Starting Dose**

For patients with serum erythropoietin levels  $\leq$  500 mU/mL, the recommended starting dose of EPREX epoetin alfa is 100 IU/kg as an intravenous or subcutaneous injection three times weekly for eight weeks.

### **Increase Dose**

During the dose adjustment phase of therapy, the hemoglobin should be monitored weekly. If the response is not satisfactory in terms of reducing transfusion requirements or increasing hemoglobin after eight weeks of therapy, the dose of EPREX epoetin alfa can be increased by 50-100 IU/kg three times per week. Response should be evaluated every 4-8 weeks thereafter and the dose adjusted accordingly by 50-100 IU/kg increments three times per week. If patients

have not responded satisfactorily to an EPREX dose of 300 IU/kg three times per week up to month 12 of therapy, further continuation of treatment is not warranted as it is unlikely that they will respond to higher doses of EPREX epoetin alfa.

**Maintenance Dose**

After attainment of the desired response (i.e., reduced transfusion requirements or increased hemoglobin), the dose of EPREX epoetin alfa should be titrated to maintain the response based on factors such as variations in zidovudine dose and the presence of intercurrent infectious or inflammatory episodes. If the hemoglobin exceeds 120 g/L, the dose should be temporarily withheld until the hemoglobin falls below 120 g/L. Resume dosing at 25% less than the previous dose and titrate the dose to maintain the desired hemoglobin.

**Hemoglobin Range**

Maximum benefit from EPREX therapy appears to occur when the hemoglobin is maintained in the range of 120-130 g/L; however, the hemoglobin for zidovudine-treated/HIV-infected patients should not exceed 120 g/L.

**CANCER PATIENTS RECEIVING CHEMOTHERAPY**

EPREX should not be initiated at hemoglobin levels  $\geq 100$  g/L. EPREX administration should be discontinued following completion of a chemotherapy course.

**Starting Dose**

Two EPREX dosing regimens may be used in adults: 150 IU/kg subcutaneously three times per week or 40,000 IU subcutaneously once per week.

**Dose Adjustment**

*Three times per week (TIW)*

Starting dose	150 IU/kg SC (TIW)
Increase dose to 300 IU/kg for four weeks	If the response is not satisfactory i.e., if after four weeks of therapy, no reduction in transfusion requirements or the hemoglobin has not increased by $\geq 10$ g/L
Reduce Dose by 25%	If the hemoglobin reaches a level needed to avoid transfusions or increases $> 10$ g/L in any 2 week period
Withhold dose	If hemoglobin exceeds a level needed to avoid transfusions or exceeds 120 g/L. Restart dosing at 25% below the previous dose.
Discontinue	Discontinue EPREX if after 8 weeks of therapy there is no response as measured by increase hemoglobin levels or decrease need for RBC transfusions.

Once per week (QW)

Starting Dose	40,000 IU SC (QW)
Increase dose to 60,000 IU/week for four weeks	If the response is not satisfactory i.e., if after four weeks of therapy, no reduction in transfusion requirements or the hemoglobin has not increased by $\geq 10$ g/L
Reduce Dose by 25%	If the hemoglobin reaches a level needed to avoid transfusions or increases $> 10$ g/L in any 2 week period
Withhold dose	If hemoglobin exceeds a level needed to avoid transfusions or exceeds 120 g/L. Restart dosing at 25% below the previous dose.
Discontinue	Discontinue EPREX if after 8 weeks of therapy there is no response as measured by increase hemoglobin levels or decrease need for RBC transfusions.

**Endogenous Serum Erythropoietin Levels**

In patients being treated with cyclic chemotherapy, there does not appear to be a significant relationship between the endogenous serum erythropoietin level and response to EPREX therapy.

Use of EPREX epoetin alfa is not recommended in patients with grossly elevated serum erythropoietin levels (e.g.,  $> 200$  mU/mL).

**SURGERY PATIENTS**

**Use of EPREX epoetin alfa to Reduce Allogeneic Blood Exposure**

The recommended dose regimen is 600 IU/kg subcutaneously given once weekly for three weeks (Days -21, -14, and -7) prior to surgery and on the day of surgery.

If the period prior to surgery is less than three weeks, 300 IU/kg subcutaneously may be given as an alternative dosing regimen for 10 consecutive days prior to surgery, on the day of surgery, and for four consecutive days immediately thereafter.

All patients should receive adequate iron replacement. Iron replacement should be initiated no later than the beginning of treatment with EPREX epoetin alfa and should continue throughout the course of therapy.

**Combined Use of EPREX epoetin alfa and ABD**

EPREX epoetin alfa should be administered twice weekly for three weeks prior to surgery if the presurgical predonation interval permits. At each patient visit, a unit of blood is collected and stored for autologous transfusion if the patient has an acceptable hematocrit or hemoglobin for predonation.

The recommended dosage regimen is 600 IU/kg intravenously twice weekly.

Iron status should be evaluated for all patients prior to treatment with EPREX epoetin alfa. Iron deficiency, if present, should be corrected. To maintain erythropoiesis, adequate iron

supplementation is required beginning as soon as possible and should continue throughout the course of therapy. In anemic patients, the cause of anemia should be explored before starting therapy with EPREX epoetin alfa.

### **Administration**

**NOTE:** All strengths of EPREX Pre-filled Syringes now contain a “peelable” label on the barrel of the syringe. The peelable portion of the label should be removed from the syringe barrel and affixed to the patient’s chart. The peelable label will be used to track Lot numbers and use of EPREX Pre-filled syringes.

1. **Do Not Shake.** Shaking may denature the glycoprotein, rendering it biologically inactive.
2. Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration. Do not use any vials or pre-filled syringes exhibiting particulate matter or discoloration.
3. Using aseptic techniques, attach a sterile needle to a sterile syringe. Remove the flip top from the vial containing EPREX Sterile Solution, and wipe the septum with a disinfectant. Insert the needle into the vial, and withdraw into the syringe an appropriate volume of solution.
4. Intravenous Injection: EPREX Sterile Solution should be administered over at least one to five minutes, depending on the total dose. While the administration of EPREX is independent of the dialysis procedure, EPREX may be administered into the venous line at the end of the dialysis procedure to obviate the need for additional venous access. A slower injection may be preferable in patients who develop flu-like symptoms.
5. Subcutaneous Injection: The maximum volume per injection site should be 1 mL. In case of larger volumes, more than one injection site should be used. The injections should be given in the limbs or the anterior abdominal wall. The patient should always alternate the site for each injection.
6. Multi-use Vial: Contains benzyl alcohol as a preservative and can be re-entered. The date of first entry into the vial should be recorded on the inside flap of the box. Discard 30 days after first entry.
7. Single-use Vial and Pre-filled Syringe: Do not re-enter the vial. Contains no preservative. Discard unused portions.
8. Do not administer by intravenous infusion or mix with other drugs.
9. The multi-use vial contains 0.9% benzyl alcohol which acts as a local anesthetic and may ameliorate subcutaneous injection site discomfort.
10. The phosphate-buffered formulation has been found to mitigate injection site discomfort.

## OVERDOSAGE

The maximum amount of EPREX epoetin alfa that can be safely administered in single or multiple doses has not been determined. Doses of up to 1500 IU/kg three times per week for three to four weeks have been administered to adults without any direct toxic effects of EPREX epoetin alfa itself. Humans have received EPREX epoetin alfa doses as high as 3000 IU/kg in a single day without acute toxic effects.

EPREX therapy can result in polycythemia if the hemoglobin is not carefully monitored and the dose appropriately adjusted. If the recommended range is exceeded, EPREX therapy may be temporarily withheld until the hemoglobin returns to the recommended range; EPREX therapy may then be resumed using a lower dose (see **DOSAGE AND ADMINISTRATION**). If polycythemia is of concern, phlebotomy may be indicated to decrease the hemoglobin to within acceptable ranges. Supportive care should be provided for hypertensive or convulsive events that may be related to overdosing with EPREX epoetin alfa.

## ACTION AND CLINICAL PHARMACOLOGY

### Mechanism of Action

Erythropoietin is a glycoprotein which stimulates red blood cell production. It is produced in the kidney and stimulates the division and differentiation of committed erythroid progenitors in the bone marrow.

### Pharmacodynamics

#### CRF PATIENTS

Erythropoietin is a glycoprotein which stimulates red blood cell production. Endogenous production of erythropoietin is normally regulated by the level of tissue oxygenation. Hypoxia and anemia generally increase the production of erythropoietin, which in turn stimulates erythropoiesis. In normal subjects, plasma erythropoietin levels range from 0.01 to 0.03 U/mL and increase up to 100 to 1000-fold during hypoxia or anemia. In contrast, in patients with chronic renal failure (CRF), production of erythropoietin is impaired, and this erythropoietin deficiency is the primary cause of their anemia.

EPREX epoetin alfa has been shown to stimulate erythropoiesis in anemic patients with CRF, including both patients on dialysis and those who do not require regular dialysis. The first evidence of a response to EPREX epoetin alfa is an increase in the reticulocyte count within 10 days, followed by increases in the red cell count, hemoglobin and hematocrit, usually within 2-6 weeks.

**Because several days are required for erythroid progenitors to mature and be released into the circulation, a clinically significant increase in hematocrit is usually not observed in less than 2 weeks and may require up to 6 weeks in some patients.** Once the hematocrit reaches the recommended range (30-36%), that level can be sustained by EPREX epoetin alfa in the absence of iron deficiency and concurrent illnesses.

The rate of hematocrit increase varies between patients and is dependent upon the dose of EPREX epoetin alfa, within a therapeutic range of approximately 50-300 IU/kg three times per week. Other factors affecting the rate and extent of response include availability of iron stores, the baseline hematocrit, and the presence of concurrent medical problems.

Intravenously administered EPREX epoetin alfa is eliminated at a rate consistent with first order kinetics with a circulating half-life ranging from approximately 4 to 13 hours in adult and 6.2 to 8.7 hours in pediatric patients with CRF. Within the therapeutic dose range, detectable levels of plasma erythropoietin are maintained for at least 24 hours. After subcutaneous administration of EPREX epoetin alfa to patients with CRF, peak serum levels are achieved within 5-24 hours after administration and decline slowly thereafter. In comparison with intravenous administration, subcutaneously administered EPREX epoetin alfa is more slowly absorbed and results in lower serum levels which are maintained for 48 hours. The estimated  $AUC_{0-48}$  for subcutaneous administration is approximately 15% of the  $AUC_{0-48}$  for the same dose given intravenously. Despite these differences, EPREX epoetin alfa exhibits a dose-related effect on hematological parameters which is independent of route. There is no apparent difference in half-life between adult patients not on dialysis whose serum creatinine levels were greater than 264  $\mu\text{mol/L}$  (3 mg/dL), and adult patients maintained on dialysis. In normal volunteers, the half-life of intravenously administered EPREX Sterile Solution is approximately 20% shorter than the half-life in CRF patients.

The pharmacokinetic profile of EPREX epoetin alfa in children and adolescents appears to be similar to that of adults. Limited data are available in neonates.

### **ZIDOVUDINE-TREATED/HIV-INFECTED PATIENTS**

Response to EPREX epoetin alfa in zidovudine-treated/HIV-infected patients is manifested by reduced transfusion requirements and increased hematocrit. Responsiveness to EPREX therapy in HIV-infected patients is dependent upon the endogenous serum erythropoietin levels prior to treatment. Zidovudine-treated/HIV-infected patients with endogenous serum erythropoietin levels  $\leq 500$  mU/mL respond to EPREX therapy. Patients with endogenous serum erythropoietin levels  $> 500$  mU/mL do not appear to respond to EPREX therapy. It appears likely that endogenous serum erythropoietin levels in HIV-infected patients receiving zidovudine are related to the severity of the zidovudine-induced damage to erythroid precursors in the bone marrow.

### **CANCER PATIENTS**

Anemia in cancer patients may be related to the effect of concomitantly administered chemotherapeutic agents. EPREX epoetin alfa administered three times a week (TIW) has been shown to increase hemoglobin and decrease transfusion requirements (after the first month of therapy) in anemic cancer patients. In addition, EPREX epoetin alfa administered once weekly (QW) has been shown to increase hemoglobin and decrease transfusion requirements after the first month of therapy (months 2, 3 and 4) in anemic cancer patients.

In a series of clinical trials enrolling 413 anemic cancer patients who received EPREX epoetin alfa TIW, 289 of whom were receiving cyclic chemotherapy, approximately 75 percent of the patients had endogenous serum erythropoietin levels  $\leq 150$  mU/mL, and approximately 5 percent

of patients had endogenous serum erythropoietin levels > 500 mU/mL. In patients who were being treated with cyclic chemotherapeutic regimens, there was not a statistically significant relationship between response to EPREX therapy and the prestudy endogenous serum erythropoietin level; however, treatment of patients with grossly elevated serum erythropoietin levels (e.g., > 200 mU/mL) is not recommended.

In a Phase 1 PK/PD study comparing 150 IU/kg subcutaneous thrice weekly (TIW) dosing to 40,000 IU subcutaneous once weekly (QW) dosing in healthy subjects, the following parameters were estimated using data corrected for predose endogenous erythropoietin concentration during Week 4: mean  $C_{\max}$  (SD) was 191 (100.1) and 785 (427.3) mIU/mL, respectively; mean (SD)  $C_{\min}$  was 39 (17.9) and 13 (9.5) mIU/mL, respectively, and mean  $t_{1/2}$  was 31.8 and 39.3 hours, respectively, after the 150 IU/kg TIW dosing (n=24) and 40,000 IU QW (n=22) dosing regimens were administered. Bioavailability of epoetin alfa after the 40,000 IU/week dosing regimen relative to the 150 IU/kg thrice weekly dosing regimen, based on AUC comparison, was 176%.

In another Phase 1 PK/PD study, pharmacokinetic parameters were estimated using data corrected for predose endogenous erythropoietin concentration comparing 150 IU/kg TIW and 40,000 IU QW dosing regimens in healthy subjects (n=6 per arm) and anemic cancer subjects (n=9 per arm). The respective mean (SD) parameters for the 150 IU/kg TIW and 40,000 IU QW regimens in healthy subjects were:  $C_{\max}$  163 (53.6) and 1036 (237.9) mIU/mL;  $C_{\min}$  29 (10.4) and 9.2 (5.71) mIU/mL;  $t_{\max}$  9.0 (3.29) and 21 (7.1) hours;  $t_{1/2}$  25.2 (6.76) [n=4] and 28.9 (7.98) hours, and CL/F 31.2 (11.48) and 12.6 (3.05) mL/h/kg. The respective mean (SD) parameters for the 150 IU/kg TIW and 40,000 IU QW regimens in the anemic cancer subjects were:  $C_{\max}$  263 (179.1) and 931 (596.9) mIU/mL;  $C_{\min}$  54 (41.8) and 34 (32.3) mIU/mL;  $t_{\max}$  8.3 (8.58) and 43 (16.0) hours;  $t_{1/2}$  29.9 (2.43) [n=4] and 22.9 (7.55) [n=7] hours and CL/F 45.8 (58.81) and 11.3 (6.45) mL/h/kg. It appeared that the pharmacokinetics of epoetin alfa in anemic cancer subjects were similar to those in healthy subjects.

However, the degree of variability associated with the pharmacokinetic parameters was higher in anemic cancer patients.

### **Pharmacodynamics**

In a PK/PD study comparing the 150 IU/kg thrice weekly and 40,000 IU once weekly dosing regimens in healthy subjects (n = 6 per arm) and in anemic cancer subjects (n = 9 per arm), the time profiles of changes in percent reticulocytes, hemoglobin, and total red blood cells were similar between the two dosing regimens in both healthy and anemic cancer subjects. The AUCs of the respective pharmacodynamic parameters were similar between the 150 IU/kg TIW and 40,000 IU QW dosing regimens in healthy subjects and also in anemic cancer subjects, although the extent of increase in hemoglobin and RBC count was slightly lower in anemic cancer subjects than in healthy subjects (in terms of AUC of hemoglobin and RBC over the study period).



## SURGERY PATIENTS

### Use of EPREX epoetin alfa to Reduce Allogeneic Blood Exposure

Patients undergoing major elective surgery frequently require transfusion of allogeneic blood, both intraoperatively and postoperatively, resulting from blood loss experienced during and after surgery. In patients with a pretreatment hemoglobin of  $> 100$  to  $\leq 130$  g/L, EPREX epoetin alfa has been shown to decrease the risk of receiving allogeneic transfusions and hasten erythroid recovery (i.e., increased hemoglobin levels, hematocrit levels, and reticulocyte counts).

### Combined Use of EPREX epoetin alfa and Autologous Blood Donation (ABD)

EPREX epoetin alfa has been shown to stimulate red blood cell production in order to augment autologous blood collection, and to limit the decline in hematocrit in adult patients scheduled for major elective surgery who are not expected to predeposit their complete perioperative blood needs. The greatest effects are observed in patients with low hematocrit ( $\leq 39\%$ ).

### Pharmacokinetics and Hematological Responses

Measurement of epoetin alfa following multiple dose intravenous administration revealed a half-life of approximately 4 hours in normal volunteers and a somewhat more prolonged half-life in renal failure patients, approximately 5 hours (ranging to 13 hours). Within the therapeutic dose range, detectable levels of plasma erythropoietin are maintained for at least 24 hours. There is no apparent difference in half-life between patients not on dialysis whose serum creatinine levels were greater than 264  $\mu\text{mol/L}$  (3 mg/dL), and patients maintained on dialysis.

After subcutaneous administration of EPREX epoetin alfa to normal subjects, peak serum levels are achieved within 5-24 hours after administration. In comparison with intravenous administration, subcutaneously administered EPREX epoetin alfa is more slowly absorbed and results in lower serum levels which are maintained for 48 hours. In normal subjects, there was no significant epoetin alfa accumulation when administered under the clinical trial conditions. After multiple doses, similar blood levels were obtained after initial and subsequent injections in normal subjects. In these subjects, the half-life was estimated to be about 24 hours. The bioavailability of subcutaneous epoetin alfa is approximately 20% of that of the intravenously injected epoetin alfa. Despite these differences, EPREX epoetin alfa exhibits an injection-route independent, dose-related effect on hematological parameters.

In normal volunteers,  $C_{\text{max}}$  values for 40,000 IU once weekly determined in four-week studies were 6 times and  $\text{AUC}_{(0-168\text{ h})}$  were 3 times that of the 150 IU/kg thrice weekly dosing regimen. Mean hemoglobin increases are similar with  $3.1 \pm 0.86$  and  $3.1 \pm 0.84$  g/dL for the 150 IU/kg thrice weekly and 40,000 IU once weekly, respectively. The time profiles of changes in hemoglobin and total red blood cells over the one-month study period were similar between the two dosing regimens.

However, statistically significant differences between genders for the  $\text{AUC}_{\text{HEMO}}$  and  $\text{AUC}_{\text{RBC}}$  pharmacodynamic response parameters at 40,000 IU once weekly dosing were noted. Clinical consequences of these differences were not studied as the normal volunteers were not scheduled subject to surgical procedures. When demographic data for the 40,000 IU once weekly and 150 IU/kg t.i.w. regimens were examined separately, or combined and stratified by gender, a significant difference in weight between males and females was detected in each treatment

group. Dosing should be individualized on an IU/kg basis (see **DOSAGE AND ADMINISTRATION**).

### **Special Populations and Conditions**

**Pediatrics:** No data available.

**Geriatrics:** No data available.

**Gender:** No data available.

**Race:** No data available.

**Hepatic Insufficiency:** The safety of EPREX epoetin alfa has not been established in patients with hepatic dysfunction.

**Renal Insufficiency:** Based on information to date, the use of EPREX epoetin alfa in predialysis patients does not accelerate the rate of progression of renal insufficiency.

### **STORAGE AND STABILITY**

Store at 2° - 8°C. Protect from exposure to light. **Do Not Freeze. Do Not Shake.** Multi-use vial should be discarded 30 days after first entry.

When the pre-filled syringe or single-use vial is about to be used, it may be removed from the refrigerator and stored at room temperature (not above 25°C) for one single period of maximum 7 days.

### **SPECIAL HANDLING INSTRUCTIONS**

Store at 2° - 8°C. Protect from exposure to light. **Do Not Freeze. Do Not Shake.**

### **DOSAGE FORMS, COMPOSITION AND PACKAGING**

#### **HSA-Containing Formulation Multi-Use Vial**

The HSA-Containing Formulation Multi-Use Vial contains the active ingredient epoetin alfa and the following inactive ingredients:

- Stabilizer: Albumin (Human) (0.25%)
- Other Inactives: Sodium chloride, sodium citrate, citric acid, water for injection and the preservative, benzyl alcohol (0.9%).

**Polysorbate-80 Containing (HSA-free) Formulation Single-Use Vials and Pre-filled Syringes with PROTECS needle guard**

Polysorbate-80 Containing (HSA-free) Formulation Single-Use Vials and Pre-filled Syringes contain the active ingredient epoetin alfa and the following inactive ingredients:

Stabilizer: Glycine and polysorbate 80  
Other Inactives: Sodium chloride, sodium phosphate monobasic dihydrate, sodium phosphate dibasic dihydrate, water for injection.

EPREX epoetin alfa Injection is supplied as a sterile solution in the following formats:

**Single-Use Vial (phosphate-buffered)**

1,000 IU/0.5 mL  
2,000 IU/mL  
4,000 IU/mL  
10,000 IU/mL  
40,000 IU/mL

**Multi-Use Vial (preserved, citrate-buffered)**

20,000 IU/mL

**Single-Use Pre-filled Syringe with PROTECS needle guard (phosphate-buffered)**

1,000 IU/0.5 mL  
2,000 IU/0.5 mL  
3,000 IU/0.3 mL  
4,000 IU/0.4 mL  
5,000 IU/0.5 mL  
6,000 IU/0.6 mL  
7,000 IU/0.7 mL  
8,000 IU/0.8 mL  
9,000 IU/0.9 mL  
10,000 IU/mL  
20,000 IU/0.5 mL  
30,000 IU/0.75 mL  
40,000 IU/mL

To reduce the risk of accidental needle sticks to users, each pre-filled syringe is equipped with the PROTECS needle guard that is automatically activated to cover the needle after complete delivery of the syringe content.

For packaging configuration see published price list.

## PART II: SCIENTIFIC INFORMATION

### PHARMACEUTICAL INFORMATION

#### Drug Substance

Proper name: EPREX

Chemical name: epoetin alfa

Molecular formula and molecular mass: Approximately 30,400 daltons when fully glycosylated. (Approximately 18,000 daltons for the peptide chain.)

Structural formula: Peptide chain of 165 amino acids. Sequence identical to that of human urinary erythropoietin.

Physicochemical properties: Bulk product is a clear colourless liquid with a pH of  $6.9 \pm 0.3$  at  $23^{\circ}\text{C} \pm 5^{\circ}\text{C}$  consisting of epoetin alfa in buffer.

#### Product Characteristics

Epoetin alfa is a purified glycoprotein hormone of recombinant DNA origin that stimulates erythropoiesis. It is produced from mammalian Chinese hamster ovary (CHO) cells that have been transfected with the human gene coding for erythropoietin.

EPREX epoetin alfa is formulated as a sterile, colourless liquid for intravenous (IV) or subcutaneous (SC) administration.

### CLINICAL TRIALS

#### CRF PATIENTS

Response to EPREX epoetin alfa was consistent across all studies. In the presence of adequate iron stores (see **WARNINGS AND PRECAUTIONS, Monitoring and Laboratory Tests, Iron Evaluation**), the time to reach the target hematocrit is a function of the baseline hematocrit and the rate of hematocrit rise.

The rate of increase in hematocrit is dependent upon the EPREX dose administered and individual patient variation. In clinical trials at starting doses of 50-150 IU/kg three times per week, patients responded with an average rate of hematocrit rise of:

**Table 2.1: Hematocrit Increase**

Starting Dose Three times/week, IV	Hematocrit percentage points/day	Hematocrit percentage points/2 weeks
50 IU/kg	0.11	1.5
100 IU/kg	0.18	2.5
150 IU/kg	0.25	3.5

Over this dose range, approximately 95% of all patients responded with a clinically significant increase in hematocrit. By the end of approximately two months of therapy, virtually all patients were transfusion-independent. Once the target hematocrit was achieved, the maintenance dose was individualized for each patient.

### **Adult Patients on Dialysis**

Thirteen clinical studies were conducted, involving intravenous administration to a total of 1010 anemic patients on dialysis for 986 patient-years of EPREX therapy. In the three largest of these clinical trials, the median maintenance dose necessary to maintain the hematocrit between 30-36% was approximately 75 IU/kg given three times per week. In a U.S. multicentre Phase III study (n=412), approximately 65% of the patients required doses of 100 IU/kg, or less, three times per week to maintain their hematocrit at approximately 35%. Almost 10% of patients required a dose of 25 IU/kg, or less, and approximately 10% required a dose of more than 200 IU/kg three times per week to maintain their hematocrit at this level.

Among the 99 patients treated in the double-blind, placebo-controlled, Canadian multicentre, quality of life study, clinically and statistically ( $p < 0.05$ ) significant improvement was shown in the patients treated with EPREX epoetin alfa compared to the placebo group when measuring fatigue, relationships and depression (Kidney Disease Questionnaire), distance walked (Six Minute Walk Test) and time on the Naughton Treadmill test after six months of therapy. Sixty-seven patients from the group treated with EPREX epoetin alfa were then enrolled in an open-label extension study where 62 patients maintained their quality of life improvements for an additional 12 months.

A multicentre study was also conducted in 152 patients receiving peritoneal dialysis who self-administered EPREX epoetin alfa Sterile Solution subcutaneously for approximately 132.9 patient-years of experience. Patients responded to epoetin alfa administered subcutaneously in a manner similar to patients receiving intravenous administration.

A randomized prospective trial of 1265 hemodialysis patients with clinically evident cardiac disease (ischemic heart disease or congestive heart failure) was conducted in which patients were assigned to EPREX treatment targeted to a maintenance hematocrit of either  $42 \pm 3\%$  or  $30 \pm 3\%$ . Increased mortality was observed in 634 patients randomized to a target hematocrit of 42% [221 deaths (35% mortality)] compared to 631 patients targeted to remain at a hematocrit of 30% [185 deaths (29% mortality)]. The reason for increased mortality observed in this study is

unknown; however, the incidence of non-fatal myocardial infarctions (3.1% vs. 2.3%), vascular access thrombosis (39% vs. 29%) and all other thrombotic events (22% vs. 18%) was also higher in the group randomized to achieve a hematocrit of 42%.

A randomized prospective trial (CHOIR) evaluated 1432 anemic chronic renal failure patients who were not undergoing dialysis. Patients were assigned to epoetin alfa treatment targeting a maintenance hemoglobin concentration of 135 g/L or 113 g/L. A major cardiovascular event (death, myocardial infarction, stroke or hospitalization for congestive heart failure) occurred among 125 (18%) of the 715 patients in the higher hemoglobin group compared to 97 (14%) among the 717 patients in the lower hemoglobin group (hazard ratio [HR] 1.3, 95% CI: 1.0, 1.7, p = 0.03).

### **Pediatric Chronic Renal Failure Patients on Dialysis**

One hundred twenty-eight children from 2 months to 19 years of age with CRF requiring dialysis were enrolled in 4 clinical studies of epoetin alfa. The largest study (EPO-9002) was a placebo-controlled, randomized trial in 113 children with anemia (hematocrit  $\leq$  27%) undergoing peritoneal dialysis or hemodialysis. The patient demographics for this study are presented in Table 2.2. The initial dose of epoetin alfa was 50 IU/kg IV or SC, TIW. The dose of study drug was titrated to achieve either a hematocrit of 30% to 36% or an absolute increase in hematocrit of 6 percentage points over baseline.

**Table 2.2: Patient Demographics for Study EPO-9002**

Variable		Hemodialysis		Peritoneal Dialysis		Total (N=113)	
		Epoetin alfa (N=24) %	Placebo (N=24) %	Epoetin alfa (N=31) %	Placebo (N=34) %	Epoetin alfa (N=55) %	Placebo (N=58) %
Age Group	0 - < 5	0.0	4.2	16.1	32.4	9.1	20.7
	5 - < 15	54.2	50.0	51.6	55.9	52.7	53.4
	15 - < 18	45.8	45.8	32.3	11.8	38.2	25.9
Sex	Male	54.2	62.5	64.5	55.9	60.0	58.6
	Female	45.8	37.5	35.5	44.1	40.0	41.4
Race	Caucasian	37.5	25.0	35.5	41.2	36.4	34.5
	Black	37.5	41.7	19.4	20.6	27.3	29.3
	Hispanic	25.0	33.3	35.5	38.2	30.9	36.2
	Asian	0.0	0.0	6.5	0.0	3.6	0.0
	Other	0.0	0.0	3.2	0.0	1.8	0.0

At the end of the initial 12 weeks, a statistically significant rise in mean hematocrit (9.4% vs 0.9%) was observed in the epoetin alfa arm. The proportion of children achieving a hematocrit of 30%, or an increase in hematocrit of 6 percentage points over baseline, at any time during the first 12 weeks was higher in the epoetin alfa arm (96% vs 58%). Within 12 weeks of initiating epoetin alfa therapy, 92.3% of the pediatric patients were transfusion-independent as compared to 71% who received placebo. Among patients who received 36 weeks of epoetin alfa, hemodialysis patients required a higher median maintenance dose (167 IU/kg/week [n=28] vs 76 IU/kg/week [n=36]) and took longer to achieve a hematocrit of 30% to 36% (median time to response 69 days vs 32 days) than patients undergoing peritoneal dialysis. Additional information regarding the number of patients who experienced an increase of 6 or more hematocrit points from baseline, versus attaining a target hematocrit of 30% to 36 % can be found in detail in Table 2.3.

**Table 2.3: Achievement of a Hematocrit Increase of 6 or More Points Above Baseline in the Study EPO-9002 (First 12 Study Weeks)**

Dialysis Type	Age Group (Years)	Epoetin alfa			Placebo		
		No. of Patients	No. Achieved	% Achieved	No. of Patients	No. Achieved	% Achieved
Hemodialysis	0 - < 5	0	0	0.0	1	0	0.0
	5 - < 15	13	13	100.0	11	6	54.5
	15 - < 18	11	8	72.7	11	3	27.3
	<b>Subtotal</b>	24	21	87.5	23	9	39.1
Peritoneal Dialysis	0 - < 5	5	5	100.0	11	7	63.6
	5 - < 15	16	16	100.0	19	13	68.4
	15 - < 18	10	10	100.0	4	2	50.0
	<b>Subtotal</b>	31	31	100.0	34	22	64.7
All Dialysis Types	0 - < 5	5	5	100.0	12	7	58.3
	5 - < 15	29	29	100.0	30	19	63.3
	15 - < 18	21	18	85.7	15	5	33.3
	<b>Total</b>	55	52	94.5	57	31	54.4

### **CRF Patients Not Requiring Dialysis**

Four clinical trials were conducted in patients with CRF not on dialysis involving 181 patients treated with EPREX epoetin alfa for approximately 67 patient-years of experience. The average duration of therapy was nearly five months. These patients responded to EPREX therapy in a manner similar to that observed in patients on dialysis. Patients with CRF not on dialysis demonstrated a dose-dependent and sustained increase in hematocrit when EPREX Sterile Solution was administered by either an intravenous or subcutaneous route. Similar rates of rise of hematocrit were noted when EPREX Solution was administered by either route. Moreover, EPREX doses of 75-150 IU/kg per week have been shown to maintain hematocrits of 36-38% for up to six months. Correcting the anemia of progressive renal failure will allow patients to remain active even though their renal function continues to decrease.

### **ZIDOVUDINE-TREATED/HIV-INFECTED PATIENTS**

EPREX epoetin alfa has been studied in four placebo-controlled trials enrolling 297 anemic (hematocrit < 30%) HIV-infected (AIDS) patients receiving concomitant therapy with zidovudine. In the subgroup of patients (n=89) with prestudy endogenous serum erythropoietin levels  $\leq$  500 mU/mL, EPREX epoetin alfa significantly reduced the mean cumulative number of units of blood transfused per patient by approximately 40%. Among those patients who required transfusions at baseline, 43% of patients treated with EPREX epoetin alfa versus 18% of placebo-treated patients were transfusion-independent during the second and third months of therapy. EPREX epoetin alfa also resulted in statistically significant increases in hematocrit in comparison to placebo (6.9% vs. 0.6% in placebo group,  $p=0.0167$ ). When examining the results according to the weekly dose of zidovudine received during Month 3 of therapy, statistically significant effects of EPREX epoetin alfa on Month 3 transfusion requirements were limited to patients with endogenous serum erythropoietin levels  $\leq$  500 mU/mL who were taking  $\leq$  4200 mg/week of zidovudine. Approximately 17% of the patients with endogenous serum erythropoietin levels  $\leq$  500 mU/mL receiving EPREX epoetin alfa in doses from 100-200 IU/kg three times weekly achieved the target hematocrit of 38% unrelated to transfusion or to a significant reduction in zidovudine dose. In the subgroup of patients whose prestudy endogenous serum erythropoietin levels were  $>$  500 mU/mL, EPREX epoetin alfa did not reduce

transfusion requirements or increase hematocrit compared to the corresponding responses in placebo-treated patients.

Clinical experience with EPREX epoetin alfa administered for more than one year to HIV-infected (AIDS) patients receiving concomitant zidovudine therapy suggests that additional benefit, in terms of raising hematocrit and decreasing transfusion requirements, may be obtained by increasing the EPREX dose in a stepwise fashion above initial dose levels in patients whose prestudy endogenous serum erythropoietin levels are  $\leq 500$  mU/mL.

Responsiveness to EPREX therapy may be blunted by intercurrent infectious/inflammatory episodes and by an increase in zidovudine dosage. Consequently, the dose of EPREX epoetin alfa must be titrated based on these factors to maintain the desired erythropoietic response.

## **CANCER PATIENTS**

### *Three times per week dosing*

EPREX epoetin alfa has been studied in a series of placebo-controlled, double-blind trials in a total of 289 anemic cancer patients. Within the group, 157 patients were treated with concomitant non-cisplatin-containing chemotherapy regimens and 132 patients were treated with concomitant cisplatin-containing chemotherapy regimens. Patients were randomized to EPREX doses of 150 IU/kg or placebo subcutaneously three times per week for 12 weeks.

In each of the study populations tested, EPREX therapy was associated with a significantly ( $p < 0.005$ ) greater hematocrit response than in the corresponding placebo-treated patients. In all patients treated with concomitant chemotherapy, transfusion requirements in patients treated with EPREX epoetin alfa (1.04 units in Months 2, 3) were reduced by approximately 43 percent compared to transfusion requirements in placebo-treated patients (1.81 units in Months 2, 3) after the first month of therapy.

Comparable intensity of chemotherapy in the EPREX epoetin alfa and placebo groups in the chemotherapy trials was demonstrated by a similar area under the neutrophil time curve in patients treated with EPREX epoetin alfa and placebo-treated patients, as well as by a similar proportion of patients in groups treated with EPREX epoetin alfa and placebo-treated groups whose absolute neutrophil counts fell below 1000 and 500 cells/ $\mu$ L.

In the above studies, the double-blind phase was followed by an open-label phase during which all patients received EPREX epoetin alfa. During total EPREX epoetin alfa exposure, in each of the studies, a significant improvement ( $p \leq 0.05$ ) in hematocrit from baseline to final reported value was observed. In addition, the percentage of patients requiring transfusion declined from 30.9% and 26.2% in the chemotherapy and cisplatin groups, respectively, during the month prior to initiating therapy to 11.5% and 6.1%, respectively, during Month 5 of therapy. The following table details the transfusion rates in the open-label phase from one month prior to therapy through to Month 5 of therapy.



**Table 2.4: Proportion of Patients Transfused During Open-Label Phase**

<u>Period</u>	<u>Chemotherapy</u>		<u>Cisplatin</u>	
	N <sup>a</sup>	Percent <sup>b</sup> Transfused	N <sup>a</sup>	Percent <sup>b</sup> Transfused
<u>Baseline</u>				
Month 1	139	30.94	103	26.21
<u>On-Therapy</u>				
Month 1	139	25.18	103	42.72
Month 2	117	17.09	85	15.29
Month 3	107	14.02	71	15.49
Month 4	78	12.82	45	6.67
Month 5	61	11.48	33	6.06

<sup>a</sup> Number of patients participating during the interval.

<sup>b</sup> Percent of patients requiring transfusion during that interval.

Available evidence suggests that patients with hematologic and solid cancers respond equivalently to EPREX therapy, and that patients with or without tumour infiltration of the bone marrow respond equivalently to EPREX therapy.

EPREX epoetin alfa significantly improved energy level (by 21% relative to placebo), increased daily activities (by 24% relative to placebo) and improved overall quality of life (by 25% relative to placebo), as measured on a 100 mm Visual Analog Scale, in cancer patients whose hematocrit increased at least 6 percentage points from baseline.

EPREX epoetin alfa was also studied in a large open label trial of 2030 anemic cancer patients on chemotherapy who were treated with EPREX doses of 150 IU/kg subcutaneously three times a week for up to four months. Functional capacity parameters, i.e., energy level, activity level and overall quality of life, improved significantly ( $p < 0.001$ ) over baseline levels (38%, 32%, 24% improvement, respectively); these improvements correlated ( $p < 0.001$ ) directly with hemoglobin change from baseline. Patients whose hemoglobin increased the most, experienced the greatest improvements in energy level, activity level, and overall quality of life.

#### Once Weekly Dosing

Epoetin alfa was also studied in a placebo-controlled, double-blind trial utilizing QW dosing in a total of 344 anemic cancer patients. In this trial, 285 patients were treated with concomitant non cisplatin-containing chemotherapy regimens and 59 patients were treated with concomitant cisplatin-containing chemotherapy regimens.

Of those patients treated with non cisplatin-containing chemotherapy regimens, 79 received carboplatin and 11 received oxaliplatin. Patients were randomized to epoetin alfa 40,000 IU/week or placebo SC for a planned treatment period of 16 weeks. If after 4 weeks of therapy hemoglobin had not increased by  $> 1$  g/dL independent of RBC transfusion or the patients received RBC transfusion during the first 4 weeks of therapy, study drug was to be increased to 60,000 IU/week. 43% of patients in the epoetin alfa group had their dose of study drug increased to 60,000 IU over the entire study.

Results demonstrated that epoetin alfa therapy was associated with a significantly greater hemoglobin response than in the corresponding placebo-treated patients. The mean change from baseline to final hemoglobin value was 2.8 g/dL in patients treated with epoetin alfa and 0.9 g/dL in patients treated with placebo ( $p < 0.0001$ ) (see Table 2.5).


**Table 2.5: Hemoglobin (g/dL): Mean Change from Baseline to Last Value**

	<u>Epoetin alfa (n)</u>	<u>Placebo (n)</u>
Baseline	9.5 (174)	9.4 (170)
Last Value	12.2 (166)	10.3 (164)
Mean Change	2.8 (166)	0.9 (164)

In addition, the mean number of units of blood transfused per 100 patient-days on the study was significantly ( $p < 0.0001$ ) lower in patients treated with epoetin alfa (0.76 units per 100 patient-days on study) than in corresponding placebo-treated patients (1.54 units per 100 patient-days on study). Moreover, the proportion of patients transfused after study day 29 through week 16 (day 112) was significantly ( $p = 0.0011$ ) lower in the patients treated with epoetin alfa than in the corresponding placebo-treated patients. Estimates of the proportion of patients receiving transfusions over this period, using Kaplan-Meier analysis, indicated transfusion rates of 16% for patients in the epoetin alfa group versus 33% for patients randomized to placebo (see Figure 2.1).

**Figure 2.1**

Time to First Transfusion After Day 28 (From Day 29 to End of Treatment)  
(Intent-to-Treat Population)



Vertical marks represent subjects whose data was "censored" at that point, i.e., no longer available for analysis because the subjects were off study without transfusion. It should be noted that because this analysis starts after day 28; the horizontal lines are at 0 for the first 4 weeks of treatment. The vertical bars after Week 16 indicate the points at which subjects whose treatment extended beyond 16 weeks discontinued treatment.

Comparable intensity of chemotherapy for patients enrolled in the two study arms was suggested by similarities in mean dose and frequency of administration for the 10 most commonly administered chemotherapy agents, and similarity in the incidence of changes in chemotherapy during the trial in the two arms.

#### *Tumour Progression, Increased Mortality and Thromboembolic Events*

A randomized, placebo-controlled trial was conducted in 224 chemotherapy-naïve, non-anemic patients with small cell lung cancer receiving cisplatin-based combination chemotherapy, to investigate whether the concurrent use of epoetin alfa stimulated tumour growth as assessed by impact on overall response rate. Patients were randomized to receive epoetin alfa 150 units/kg or placebo subcutaneously TIW during chemotherapy. The overall response rates, after 3 cycles of treatment, were 72% and 67%, in the epoetin alfa and placebo arms, respectively. Complete response rates (17% vs. 14%) and median overall survival (10.5 mos vs. 10.4 mos) were similar in the epoetin alfa and placebo arms.

A systematic review of 57 randomized controlled trials (including the BEST and ENHANCE studies) evaluating 9353 patients with cancer compared ESAs plus transfusion with transfusion alone for prophylaxis or treatment of anaemia in cancer patients with or without concurrent antineoplastic therapy. An increased relative risk of thromboembolic events (RR 1.67, 95% CI: 1.35, 2.06, 35 trials and 6769 patients) was observed in ESA-treated patients. An overall survival hazard ratio of 1.08 (95% CI: 0.99, 1.18; 42 trials and 8167 patients) was observed in ESA-treated patients. Use of ESAs for treatment of anaemia in cancer patients on chemotherapy was associated with an overall survival odds ratio of 0.99 (95% CI: 0.72, 1.36). Sub-group analysis of anaemic cancer patients receiving chemotherapy from this systematic review demonstrates an odds ratio of 0.92 (95% CI: 0.78, 1.09) for platinum based chemotherapy and 1.10 (95% CI: 0.96, 1.24) for chemotherapy without platinum.

Studies of ESAs in patients with cancer exploring the effect on survival and/or progression of administrations of exogenous erythropoietin with higher hemoglobin targets beyond the treatment of anemia, and in patients not receiving chemotherapy or radiation therapy, and showing safety signals are presented below.

A randomized controlled clinical study EPO-INT-76 (referred to as the 'BEST' study) evaluated 939 mainly non-anemic women with metastatic breast cancer receiving chemotherapy. Patients received either weekly epoetin alfa or placebo for up to a year. This study was designed to show that survival was superior when epoetin alfa was administered to prevent anemia (maintain hemoglobin levels between 120 and 140 g/L). The study was terminated prematurely when interim results demonstrated that higher mortality at 4 months (8.7% epoetin alfa vs 3.4% placebo) and a higher rate of fatal thrombotic events (1.1% epoetin alfa vs 0.2% placebo) were observed among patients treated with epoetin alfa. Based on Kaplan-Meier estimates, at the time of study termination, the 12-month survival was lower in the epoetin alfa group than in the placebo group (70% vs 76%, HR 1.37, 95% CI: 1.07, 1.75, p = 0.012).

In another placebo-controlled study using epoetin beta in 351 patients with head and neck cancer (referred to as the ENHANCE study), study drug was administered to maintain the hemoglobin levels of 140 g/L in women and 150 g/L in men. Locoregional progression-free survival was significantly shorter in patients receiving epoetin beta.

A preliminary report from a clinical study (DAHANCA 10) evaluating 522 patients with primary squamous cell carcinoma of the head and neck receiving radiation therapy demonstrated a 10% increase in locoregional failure rate among darbepoetin alfa-treated patients ( $p=0.01$ ) in an interim analysis in 484 patients. Patients were randomized to darbepoetin alfa or placebo. At the time of study termination, there was a trend toward worse survival in the darbepoetin group ( $p = 0.08$ ).

In a multicentre, randomized, double-blind placebo-controlled trial (EPO-CAN-20), patients with advanced non-small-cell lung cancer unsuitable for curative therapy were treated with epoetin alfa targeting hemoglobin levels between 120 and 140 g/L or placebo. Following an interim analysis of 70 of 300 patients planned, a significant difference in median survival in favour of patients in the placebo group was observed (63 vs 129 days, HR 1.84,  $p = 0.04$ ).

A recently-completed phase III double-blind, randomized, placebo-controlled trial evaluating 989 cancer patients with active malignant disease but not being treated with either chemotherapy or radiation therapy showed no statistically significant reduction in the proportion of patients receiving red blood cell transfusions, and more deaths in the darbepoetin alfa group than in the placebo group (26% vs 20%) at 16 weeks (completion of treatment phase). With a median survival follow-up of 4.3 months, the absolute number of deaths at the end of the study was also higher in the darbepoetin alfa group (49% vs 46%, HR 1.29, 95% CI: 1.08, 1.55).

Randomized, controlled trials with decreased survival and/or decreased locoregional control are summarized in Table 2.6.

**Table 2.6: Randomized, Controlled Trials with Decreased Survival and/or Decreased Locoregional Control**

Study / Tumor / (n)	Hemoglobin Target	Achieved Hemoglobin (Median Q1,Q3)	Primary Endpoint	Adverse Outcome for ESA-containing Arm
<b>Chemotherapy</b>				
BEST Metastatic breast cancer (n=939)	120-140 g/L	129 g/L 122, 133 g/L	12-month overall survival	Decreased 12-month survival
20000161 Lymphoid malignancy (n=344)	130-150 g/L (M) 130-140 g/L (F)	110 g/L 98, 121 g/L	Proportion of patients achieving a hemoglobin response	Decreased overall survival
PREPARE Early breast cancer (n=733)	125-130 g/L	131 g/L 125, 137 g/L	Relapse-free and overall survival	Decreased 3 yr. relapse-free and overall survival
GOG 191 Cervical Cancer (n=114)	120-140 g/L	127 g/L 121, 133 g/L	Progression-free and overall survival and locoregional control	Decreased 3 yr. progression-free and overall survival and locoregional control
<b>Radiotherapy Alone</b>				
ENHANCE Head and neck cancer (n=351)	≥150 g/L (M) ≥140 g/L (F)	Not available	Locoregional progression-free survival	Decreased 5-year locoregional progression-free survival Decreased overall survival
DAHANCA 10 Head and neck cancer (n=522)	140-155 g/L	Not available	Locoregional disease control	Decreased locoregional disease control
<b>No Chemotherapy or Radiotherapy</b>				
EPO-CAN-20 Non-small cell lung cancer (n=70)	120-140 g/L	Not available	Quality of life	Decreased overall survival
20010103 Non-myeloid malignancy (n=989)	120-130 g/L	106 g/L 94, 118 g/L	RBC transfusions	Decreased overall survival

**SURGERY PATIENTS****Use of EPREX epoetin alfa to Reduce Allogeneic Blood Exposure**

EPREX epoetin alfa has been studied in a placebo-controlled, double-blind trial enrolling 316 patients scheduled for major elective orthopedic hip or knee surgery who were expected to require  $\geq 2$  units of blood. Patients were stratified into one of three groups based on their

pretreatment hemoglobin [ $\leq 100$  (n=2),  $>100$  to  $\leq 130$  (n=96), and  $>130$  to  $\leq 150$  g/L (n=218)] and then randomly assigned to receive EPREX doses of 300 IU/kg, 100 IU/kg or placebo by subcutaneous injection for 10 days before surgery, on the day of surgery, and for four days after surgery.

Treatment with EPREX epoetin alfa 300 IU/kg significantly ( $p=0.024$ ) reduced the risk of allogeneic transfusion in patients with a pretreatment hemoglobin of  $>100$  to  $\leq 130$  g/L; 16% of EPREX epoetin alfa 300 IU/kg, 23% of EPREX epoetin alfa 100 IU/kg and 45% of placebo-treated patients were transfused. There was no significant difference in number of patients transfused between EPREX epoetin alfa (9% 300 IU/kg, 6% 100 IU/kg) and placebo (13%) in the  $> 130$  to  $\leq 150$  g/L hemoglobin stratum. In the  $>100$  to  $\leq 130$  g/L pretreatment stratum, the mean number of units transfused per patient treated with EPREX epoetin alfa (0.45 units 300 IU/kg, 0.42 units 100 IU/kg) was less than the mean transfused per placebo-treated patient (1.14 units) (overall  $p=0.028$ ).

During the pretreatment to presurgery period, for both the  $> 100$  to  $\leq 130$  g/L and  $> 130$  to  $\leq 150$  g/L strata, mean increases in hemoglobin (among-group difference,  $p=0.0001$ ) and hematocrit (among-group difference,  $p=0.0001$ ) were observed in both groups treated with EPREX epoetin alfa, while no changes were observed in the placebo-treated group. In addition, greater mean increases in reticulocyte counts were observed in the groups treated with EPREX epoetin alfa compared to the placebo-treated groups ( $p=0.0001$ ). These increases were maintained postoperatively.

EPREX epoetin alfa has also been studied in two additional placebo-controlled trials enrolling patients scheduled to undergo major elective orthopedic surgery. In one study (n=208), patients received either EPREX epoetin alfa 300 IU/kg or placebo for 9 or 14 days. In the second study (n=200) patients received either EPREX epoetin alfa 300 IU/kg, 100 IU/kg, or placebo for 15 days. Results of analyses comparing the risk of transfusion according to baseline hemoglobin for intent-to-treat patients are as follows:

**Table 2.7: Percent of Patients Who Required Transfusions, Stratified by Baseline Hemoglobin**

Baseline Hemoglobin	300 IU/kg 14/15 Days	300 IU/kg 9 Days	100 IU/kg 15 Days	Placebo 14/15 Days
<u>EPREX epoetin alfa 300 IU/kg or placebo for 9 or 14 days</u>				
$>100$ to $130$ g/L	(N=25) 32%	(N=18) 61%	-	(N=23) 74%
$>130$ g/L	(N=52) 19%	(N=35) 14%	-	(N=55) 31%
<u>EPREX epoetin alfa 300 IU/kg, 100 IU/kg or placebo for 15 days</u>				
$>100$ to $130$ g/L	(N=22) 14%	-	(N=25) 44%	(N=27) 78%
$>130$ g/L	(N=29) 14%	-	(N=37) 11%	(N=39) 36%

Results from both orthopedic surgery studies indicate that in patients with a baseline hemoglobin level of  $> 100$  to  $\leq 130$  g/L, the risk that placebo-treated patients will require an allogeneic blood transfusion is two- to three-fold greater than that of patients treated with EPREX epoetin alfa.

EPREX epoetin alfa was also studied in an open-label, parallel-group trial enrolling 145 subjects with a pretreatment hemoglobin level of  $\geq 100$  to  $\leq 130$  g/L who were scheduled for major orthopedic hip or knee surgery. Subjects were randomly assigned to receive one of two dosing regimens of EPREX epoetin alfa (600 IU/kg once weekly for three weeks prior to surgery and on the day of surgery or 300 IU/kg once daily for 10 days prior to surgery, on the day of surgery and for four days after surgery), administered subcutaneously. All subjects received oral iron and appropriate pharmacologic anticoagulation therapy.

From pretreatment to presurgery, the mean increase in hemoglobin in the 600 IU/kg weekly group (14.4 g/L) was greater than observed in the 300 IU/kg daily group (7.3 g/L). The mean increase in absolute reticulocyte count was smaller in the weekly group ( $0.11 \times 10^6/\text{mm}^3$ ) compared to the daily group ( $0.17 \times 10^6/\text{mm}^3$ ). Mean hemoglobin levels were similar for the two treatment groups throughout the postsurgical period.

The erythropoietic response observed in both treatment groups resulted in similar transfusion rates (16% in the 600 IU/kg weekly group and 20% in the 300 IU/kg daily group). The mean number of units transfused per EPREX-treated subject was approximately 0.3 units in both treatment groups. The transfusion rates were substantially lower than that for placebo-treated patients (45%), and similar to the transfusion rate for the EPREX epoetin alfa 300 IU/kg-treated patients (16%) whose baseline hemoglobin was  $\geq 100$  to  $\leq 130$  g/L in the first study referenced in this section.

A randomized controlled study (SPINE) in which 681 adult patients not receiving prophylactic anticoagulation and undergoing spinal surgery received either 4 doses of 600 U/kg epoetin alfa (7, 14 and 21 days before surgery and on the date of surgery) and standard of care treatment, or standard of care treatment alone. Preliminary analysis showed a higher incidence of deep vein thrombosis, determined by either Colour Flow Duplex Imaging or by clinical symptoms in the epoetin alfa group [16 patients (4.7%)] compared to the standard of care group [7 patients (2.1%)]. In addition, 12 patients in the epoetin group and 7 patients in the standard of care group had other thrombotic vascular events.

### **Combined Use of EPREX epoetin alfa and ABD**

A double-blind study was conducted in 204 patients scheduled to undergo elective orthopedic surgery with hematocrits  $\leq 39\%$  and no underlying anemia due to iron deficiency. On average, patients treated with EPREX epoetin alfa 600 IU/kg were able to predeposit significantly more units of blood (4.5 units) than placebo-treated patients (3.0 units) ( $p < 0.001$ ). Also, significantly more patients treated with EPREX epoetin alfa ( $p \leq 0.05$ ) were able to predeposit between 3 to 6 units, inclusively, of autologous blood than the corresponding placebo-treated patients. Virtually all (98%) of the patients treated with EPREX epoetin alfa predeposited 3 or more units compared with 69% of placebo-treated patients. While 37% of the placebo patients were able to predeposit 4 or 5 units, 81% of the EPREX patients predeposited 4 or more units. Among the evaluable patients, fewer patients who received EPREX epoetin alfa required allogeneic transfusions (19.8%) than patients who received placebo (31.0%).

## DETAILED PHARMACOLOGY

### Pharmacodynamics

Table 2.8 summarizes the in vitro biological activity of epoetin alfa.

**Table 2.8**

<u>Test System</u>	<u>Species</u>	<u>Concentration Tested</u>	<u>Observations</u>
Iron incorporation into bone marrow cells in culture	Rat	0.5 - 5.0 IU/mL	Increased <sup>59</sup> Fe incorporation into heme in cell cultures treated with epoetin alfa. Dose response curve identical to that of u-HuEPO.
Colony forming units (CFU-E) assay	Mouse	0.075 - 0.75 IU/mL	The number of CFU-E units formed was directly proportional to amount of epoetin alfa
Burst forming units (BFU-E) assay	Mouse	0.1 - 3.0 IU/mL	Epoetin alfa produced dose-dependent increases in the number of BFU-E.
BFU-E Assay	Human	0.5 - 3.0 IU/mL	Epoetin alfa was able to support the growth and differentiation of human erythroid progenitors.
Assay for colony stimulating factor (CSF) activity	Mouse	10 IU/mL	No increase in non-erythroid colony formation with epoetin alfa.
Bursting-promoting activity assay	Human	0.63 and 1.25 IU/mL	No effect of epoetin alfa on the burst promoting activity in human mononuclear cells.
Megakaryocyte development assay	Mouse	Up to 3.2 IU/mL	Increased number of megakaryocytes in response to epoetin alfa in mouse bone marrow cell culture.
Radioimmunoassay (RIA)	Rabbit Antisera (U/anti-U) (R/anti-U) (R/anti-R)		Parallel curves suggesting immunological identity of epoetin alfa and u-HuEPO.
Non-pregnant uterus	Rat	10 - 1000 IU/mL	No significant effect on spontaneous contractions compared to vehicle.
Pregnant uterus	Rat	10 - 1000 IU/mL	No significant difference between drug and vehicle effects on spontaneous or oxytocin-induced contractions.
Ileum	Guinea Pig	10 - 1000 IU/mL	Epoetin alfa occasionally produced ileal contraction at 1000 IU/mL.
Trachea	Guinea Pig	10 - 1000 IU/mL	No effect on the resting tension or response to histamine.
Vas deferens	Guinea Pig	10 - 1000 IU/mL	No effect on the response to exogenous adrenaline.
Atria	Guinea Pig	10 - 1000 IU/mL	No difference between drug and vehicle treatments on contractile force or rate.
Blood	Rabbit	10 - 1000 IU/mL	No increase in hemolysis in blood of rabbits treated <i>in vitro</i> .



In general pharmacological testing, epoetin alfa is devoid of pharmacological activity up to 1000 IU/mL in *in vivo* studies on the cardiopulmonary system, isolated smooth muscle and the blood clotting system.

In addition, epoetin alfa reacts identically to u-HuEPO in RIAs using a variety of different antisera. Urinary erythropoietin (u-HuEPO) and recombinant epoetin alfa appear indistinguishable from each other in the parameters of biological and immunological reactivity measured by these assay systems. Table 2.9 summarizes the *in vivo* biological activity of epoetin alfa.

**Table 2.9**

<u>Test System</u>	<u>Species</u>	<u>Doses Tested</u>	<u>Observations</u>
Exhypoxic Poly-cythemia	Mouse	1.5 IU/mL	Increased hematopoiesis due to epoetin alfa administration.
Hematocrit	Mouse	50 - 1500 IU/kg i.p.	Increased hematocrit with epoetin alfa administration which was statistically significant at doses equal to or greater than 150 IU/kg.
Sub-total Nephrectomy	Rat	10 IU/day, 5 days/week for 2 weeks, i.m.	Increased hematocrit with epoetin alfa treatment compared to vehicle control.
Hematocrit	Dog	280 or 2800 IU/kg, i.v.	Dose-related increases in hematocrit, reticulocyte and RBC count with epoetin alfa treatment as compared to vehicle control.
r-HuEPO effect on platelet levels	Mouse	300 - 1200 IU/kg i.p. daily for 5 days	Increased hematocrit with no effect on platelet numbers. Increased megakaryocytes in the spleen.
Melphalan - Induced Anemia	Dog	100 or 300 IU/kg	Melphalan-induced anemia is attenuated by epoetin alfa as evidenced by prevention of decreased RBC and reticulocytes
Intestinal Charcoal Meal Propulsion	Mouse	20 - 2000 IU/kg i.v.	No significant effect observed.
Renal Function	Rat	20 - 2000 IU/kg i.v.	No significant effect on urine volume or electrolytes.
Blood Coagulation	Rat	20 - 2000 IU/kg i.v.	No effect on prothrombin time, activated partial thromboplastin time or recalcification time.
Fibrinolysis	Rat	20 - 2000 IU/kg i.v.	No significant effect on euglobulin lysis time.
Cardiopulmonary	Anesthetized Dog	20 - 2000 IU/kg i.v.	No significant alterations observed on blood pressure, heart rate, ECG, blood flow, respiratory rate.
Cardiopulmonary	Conscious Dog	20 - 2000 IU/kg i.v.	No significant difference in blood pressure, heart rate, ECG between vehicle and drug treatments.
CNS	Mouse	20 - 2000 IU/kg i.v.	No effect on general behaviour, motor coordination, acetic acid-induced writhing, hexobarbital sleeping time anticonvulsant effect. No statistically significant effect on locomotor activity. Significant decrease in body temperature produced by 2000 IU/kg at 4 hrs. after treatment.
CNS	Rat	20 - 2000 IU/kg i.v.	No effect on pain threshold, mono- or polysynaptic spinal reflexes, no antipyretic effect.
CNS	Rabbit	20 - 2000 IU/kg i.v.	No effect on E.E.G.

In general pharmacological testing, epoetin alfa is devoid of significant pharmacological activity up to 2000 IU/kg *in vivo*. The only noted pharmacological effect was a decrease in body temperature in mice after the 2000 IU/kg i.v. dose of epoetin alfa. This dose is in excess of the clinically effective dose (20-500 IU/kg i.v.) and therefore this finding should pose no problem to the use of epoetin alfa clinically. Therefore, epoetin alfa, as tested in various pharmacological systems, appears to be devoid of significant pharmacological activity which could affect its clinical safety.

### **Non-Clinical Pharmacokinetics**

#### **Administration:**

In the primary studies of the pharmacokinetics and metabolism of exogenously administered epoetin alfa in animals, the intravenous (i.v.) route was the mode of drug administration. Consequently, the rate and extent of absorption was not a parameter under investigation. However, it has been shown in a uremic rabbit model that <sup>125</sup>I-epoetin alfa is almost completely absorbed from the empty peritoneal cavity.

In one multiple dose study in cynomolgus monkeys, enhanced serum levels of epoetin alfa over baseline were found after subcutaneous (s.c.) administration at 1000 IU/kg, the only s.c. dose given. Epoetin alfa was absorbed slowly from the injection site, which resulted in a much slower decline in serum epoetin alfa levels than observed after i.v. administration. These issues and their clinical relevance are examined in the human clinical pharmacology section, where the kinetics of epoetin alfa after administration by i.v. and s.c. routes are compared.

#### **Distribution:**

In studies in which radiotracer techniques were utilized, tissue concentrations of radioactivity following i.v. administration of <sup>125</sup>I-epoetin alfa to rats were highest in the bone marrow, spleen and kidneys.

Other organ systems in which radioactivity was detected (in addition to the plasma) included the pituitary, lungs, liver and adrenals.

Radioactivity levels in the thyroid increased over time. This was attributed to the uptake of free <sup>125</sup>I and was not unexpected since no measures were taken to block uptake into the thyroid gland.

The initial volume of distribution of i.v. administered <sup>125</sup>I-epoetin alfa approximated that of the plasma volume.

The distribution of exogenously administered epoetin alfa was examined in the rat by whole-body autoradiography and monitoring of individual tissue levels.

In the whole-body autoradiography experiments animals were sacrificed at 0.5, 1, 4 and 24 hours post-dose. The highest radioactivity was detected in the bone marrow 30 minutes after treatment. Relatively high radioactivity was also present in the thyroid, blood, renal cortex, spleen, lungs, liver and adrenals. Almost no activity was present in the brain, gastrointestinal tract, muscle or spinal cord. The radioactivity distribution pattern was similar 30 minutes and 1 hour after treatment. Twenty-four hours after treatment, radioactivity decreased to the background level in all tissues other than the thyroid.

These findings were very consistent with measured radioactivity levels in various tissues, with animals being sacrificed at 0.5, 1, 4, 8 and 24 hours. Thirty minutes after treatment, the highest radioactivity was determined in the bone marrow. Radioactivity was also present at high levels in the spleen and kidneys 30 minutes after treatment, followed in descending order by the pituitary, liver and adrenals. Radioactivity in these organs decreased over time in parallel with plasma radioactivity levels. In contrast, radioactivity levels were low in the brain, thymus, skin and mesenteric lymph nodes. No radioactivity was detected in erythrocytes at any time point. Radioactivity levels in the thyroid increased over the 24-hour study period. This was attributed to the uptake of free  $^{125}\text{I}$  and was not unexpected since no measures were taken to block uptake into the thyroid gland.

Since erythropoietin is the primary regulator of erythropoiesis, it is not surprising that labelled epoetin alfa was promptly transferred to the bone marrow.

The initial volume of distribution of i.v. administered iodinated epoetin alfa approximated that of the plasma volume. Thereafter, the labelled hormone was cleared from the plasma in a biphasic manner. Using labelled albumin as a marker for vascular space, there was insufficient accumulation of  $^{125}\text{I}$ -epoetin alfa in the liver, kidney, spleen or bone marrow to account for the initial component of the plasma disappearance curve. This suggests that this component was due to distribution of the hormone between the plasma and the extravascular space from which elimination subsequently occurred.

### **Metabolism and Excretion:**

The metabolism of both endogenous native and exogenously administered epoetin alfa appears to occur primarily in the liver. Studies in the rat have shown this is due to the fact that desialylated hormone is cleared very efficiently by hepatic galactosyl receptors. Consequently, the bulk of the desialylated hormone is sequestered in the liver where it is rapidly catabolized to smaller, unidentified products which are released back into the plasma and subsequently excreted by the kidney.

Urinary excretion data substantiate the hepatic clearance observations in the rat. Approximately 90% of the administered dose was recovered in the urine but only about 3% of this constituted immunoreactive material.

Endogenous erythropoietin is synthesized as a pro-form that contains a 27 amino acid leader which is processed during secretion. The biologically active hormone has a molecular weight of around 34,000 daltons, is heavily glycosylated and is particularly rich in sialic acid residues. The sialic acid residues of erythropoietin are not required for the expression of biological activity *in vitro* but are required for its expression *in vivo*. The hormone's sialic acid residues serve to prevent its premature removal from the circulation. Any reduction of the sialic acid content results in a drastically reduced biological half-life. This is due to the fact that desialylated erythropoietin/epoetin alfa is cleared very efficiently by galactosyl receptors in the liver. These aspects of the metabolism of erythropoietin have been reviewed and the data obtained in rats for epoetin alfa are described below.

On administration of  $^{125}\text{I}$  labelled desialylated epoetin alfa to male Sprague-Dawley rats, only 24% of the material was recovered in the plasma as TCA-precipitable radioactivity within 2 minutes after injection and 96% of the hormone was cleared from the plasma with a half-life of

2 minutes. There was a reciprocal increase in hepatic acid-precipitable radioactivity as plasma acid-precipitable radioactivity declined. At the peak organ accumulation, 87% of the acid-precipitable radioactivity was sequestered in the liver.

Following the uptake of radiolabelled protein by the liver, there was a rapid increase in the intracellular radioactivity which was acid-soluble and this was followed by a similar increase in plasma acid-soluble radioactivity after a delay of approximately 14 minutes.

Asialo-orosomucoid, but not orosomucoid yeast mannan or Dextran sulfate 500, inhibited the rapid clearance and hepatic accumulation of desialylated erythropoietin. This suggests that uptake by the liver is regulated in part by hepatocyte galactosyl receptors. Oxidation of the desialylated hormone restored its plasma recovery and clearance to that of epoetin alfa but rendered it biologically inactive. Other determinants which may serve as recognition sites for the metabolism of epoetin alfa are as yet unknown.

Metabolism has also been studied by incubating  $^{125}\text{I}$ -epoetin alfa with various rat tissue homogenates at  $37^{\circ}\text{C}$ . Almost 100% of the added radioactivity was recovered from the TCA-precipitable fraction in liver, spleen and muscle at all sampling times (0, 0.5 hr, 1 hr, 2 hr, 4 hr). In the kidneys, however, acid-precipitable radioactivity decreased over time, and the recovery rate was 80.7% four hours after the start of incubation.

The recovery of immunoreactive radioactivity decreased over the course of the experiment in all tissues when incubated at  $37^{\circ}\text{C}$ . Decreases were relatively fast in the kidneys and liver.

The fact that intact hormone and not desialylated epoetin alfa was used in these *in vitro* studies probably influenced the rate at which epoetin alfa was metabolized. Also, although the *in vitro* results implicate the kidney as a metabolic organ in the rat, it necessitates that epoetin alfa is accumulated into kidney cells for this organ to have an inactivation role *in vivo*.

Urinary excretion data for  $^{125}\text{I}$ -epoetin alfa substantiated the hepatic clearance observations presented above. The excretion of  $^{125}\text{I}$  radioactivity in urine and feces of normal and partially nephrectomized rats has been compared. In normal rats, 86.9% and 5.7% of the dose was excreted into urine and feces, respectively, by 96 hours after treatment. Only 5.6% of the dose was recovered in the TCA-precipitable fraction of the urine and only 2.8% of the total dose was found as immunoreactive radioactivity in the urine. In partially nephrectomized rats, 81.5% and 4.8% of the total dose was excreted into urine and feces, respectively, over the same time frame.

Excretion was slightly delayed compared to normal rats. In urine, acid-precipitable radioactivity accounted for 3.6% and 3.3% of the dose, respectively.

The erythropoietin content of 24-hour urine collections has been assayed in a number of clinical conditions and healthy subjects. Low erythropoietin levels (mean values) were found in polycythemia vera ( $<10$  mU/mL), and anemia associated with uremia ( $<16$  mU/mL) and rheumatoid arthritis ( $<20$  mU/mL). Higher levels were found in association with hypoxic polycythemia and anemia ( $>75$  mU/mL) due to iron deficiency (80 mU/mL), blood loss ( $>100$  mU/mL) and marrow hypoplasia (aplastic anemia;  $>14,000$  mU/mL). Hence, urinary excretion of intact endogenous erythropoietin also appears to contribute minimally to the overall clearance of the hormone in man under normal conditions.

## TOXICOLOGY

Epoetin alfa has a very low potential for acute toxicity when administered intravenously to rats, mice and dogs, and orally or intramuscularly to rats and mice. There was no mortality or other signs of toxicity when administered at single doses up to 20,000 IU/kg.

**Table 2.10: Acute Toxicity Studies**

Strain Species	# Animal Group	Route	Vehicle	Dose Level IU/KG	Lethality	Summary Toxic Signs
Mice	8/sex	i.v.	0.25%/HSA	0 (vehicle)	0/16	None
				20000	0/16	None
				20000	0/16	None
				20000	0/16	None
Rats	8/sex	i.v.	0.25%/HSA	0 (vehicle)	0/16	None
				20000	0/16	None
				20000	0/16	None
				20000	0/16	None
Rats	10/sex	i.v.	0.25%/HSA	0 (vehicle)	0/20	None
				80	0/20	None
				240	0/20	None
				800	0/20	None
				2400	0/20	None
				8000	0/20	None
Dogs	2 males	i.v.	0.25%/HSA	0 (vehicle)	0/20	None
				10000	0/20	None
				20000	0/20	None

Most of the changes associated with intravenous (i.v.) and subcutaneous (s.c.) multidose studies represent the expected pharmacologic actions of epoetin alfa, as shown in Table 2.11

**Table 2.11: Multidose Studies Results**

<u>Key to Abbreviations</u>			
RBC	Red blood cells	SGOT	Serum glutamic-oxaloacetic transaminase
Hct	Hematocrit	LDH	Lactic acid dehydrogenase
Hgb	Hemoglobin	CPK	Creatinine phosphokinase
MCH	Mean corpuscular hemoglobin	A/G Ratio	Albumin/Globulin Ratio
MCV	Mean corpuscular volume	M:E Ratio	Myeloid:Erythrocytes Ratio
MCHC	Mean corpuscular hemoglobin concentration	TL	Total lipids
APTT	Activated partial thromboplastin time	TCHO	Total cholesterol
		PL	Phospholipid
		PT	Prothrombin time

Study	Lethality	Toxic Signs	Clinical Path	Gross/Micro
Rats i.v. 4weeks/ 4week recovery	No drug-related deaths	Reddening of pinnae, and extremities in females @ 400 and in both sexes @ 2000 IU/kg. Reduced body weight gains and food consumption in males @ 2000 IU/kg	<u>Increased</u> RBC, Hct, Hgb, reticulocytes, SGOT, LDH, bilirubin. MCH and MCV @ 16, 80 IU/kg in females and 80 IU/kg in males. <u>Decreased</u> MCV, MCHC and MCH @ higher doses except MCHC in females in all treated groups. Platelets in most treated groups, glucose, cholesterol and prolongation of APTT and PT. Urine spec. grav. in males @ 400, 2000 IU/kg. All changes were partially reversible.	Enlarged spleens. Increased erythropoiesis of bone marrow. Extra-medullary erythropoiesis in spleen and liver.
Rats i.v. 13 weeks/ 5week recovery	2 rats @ 500 IU/kg on Days 85 and 87	Decreased body weight gains @ 500 IU/kg. Dilation of vessels of eyes, reddening of retinas @ 100, 500 IU/kg.	<u>Increased</u> RBC, Hct, Hgb, reticulocytes, MCV, MCH, SGOT, CPK, LDH, bilirubin, potassium and phosphorus. Urine occult blood and RBCs in females @ 500 IU/kg. Prolongation of APTT. <u>Decreased</u> urine spec. grav. MCHC, platelets, glucose, albumin, total protein, cholesterol and calcium. Everything reversible.	Enlarged spleens, heart, kidneys, adrenals. Erosion of glandular stomach. Increased erythropoiesis in bone marrow. Extra-medullary hematopoiesis. Reversible.
Rats s.c. 2 weeks	None	None	<u>Increased</u> RBC, Hct, Hgb, reticulocytes, WBC, MCV, nucleated RBC, bilirubin, total protein, calcium, chloride, sodium, albumin and globulin. <u>Decreased</u> MCH, MCHC, BUN, glucose and cholesterol. Low titer antibodies on day 15 in 4 rats @ 200 and 1 rat @ 2000 IU/kg.	Enlarged spleens. Erythroid hyperplasia of bone marrow. Extra-medullary hematopoiesis.

**Table 2.11: Multidose Studies Results (cont'd)**

Study	Lethality	Toxic Signs	Clinical Path	Gross/Micro
Dogs i.v. 4 weeks	None	Conjunctival injection and reddening of oral mucosa @ 200, 2000 IU/kg.	<u>Increased</u> RBC, Hct, Hgb, reticulocytes, LDH, total protein, triglycerides, potassium, bilirubin and giant platelets. <u>Decreased</u> MCH, MCHC, MCV, glucose @ 200, 2000 IU/kg.	Enlarged spleens. Myelofibrosis @ 2000 IU/kg. Extramedullary erythropoiesis and megakaryopoiesis. Prostates smaller @ 2000 IU/kg.
Dogs i.v. 4 weeks	None	Conjunctival injection @ 2000 IU/kg. Decreased weight gain, food consumption @ 2000 IU/kg. Anaphylactic response on Day 14.	<u>Increased</u> RBC, Hct, Hgb, reticulocytes, LDH, total protein, triglycerides, bilirubin, potassium, cholesterol, CPK, alpha-2 globulin and beta globulin. Prolongation of APTT @ 200 IU/kg in males. <u>Decreased</u> MCH in males @ 200, 2000 IU/kg, MCHC @ 2000 IU/kg in females and males. Glucose and A/G ratios @ 200 or 2000 IU/kg.	Enlarged spleens. Extra-medullary erythropoiesis. Decreased lymphocytes in cortex of thymus, partial atrophy of seminiferous tubules @ 2000 IU/kg.
Dogs i.v. 3x/week/3 weeks 3 week Recovery	None	Red, swollen lips hypoactivity, pale mucous membranes and labored respiration in females @ 2800 IU/kg from Day 11 to 17 then decreased.	<u>Increased</u> RBC, Hgb, PCV, MCHC, reticulocytes, total protein, cholesterol, potassium and globulin. <u>Decreased</u> Glucose, A/G ratio, BUN, MCV and M:E ratios.	Enlarged spleens. Extra-medullary hematopoiesis.
Dogs i.v. 13 weeks/5 week recovery	None	Emesis, mucous stool. Decreased spontaneous activity. Prostration and tachypnea. Conjunctival injection. Redness of oral mucosa and auricle. Circumoral and circumpalebral swelling. Reversible.	<u>Increased</u> Urine volume, RBC, Hct, Hgb, reticulocytes, LDH, potassium, total protein and urinary occult blood. Prolongation of APTT and prothrombin. <u>Decreased</u> MCV, MCHC, MCH, glucose, albumin, A/G ratio, urine spec. Grav. And erythrocyte sedimentation rates. Reversible except MCV and MCH.	Enlarged spleens, kidneys, myelofibrosis, extramedullary erythropoietin. Reversible except fibrosis.

**Table 2.11: Multidose Studies Results (Cont'd)**

Study	Lethality	Toxic Signs	Clinical Path	Gross/Micro
Dogs i.v. 52 weeks	5 @ 100 IU/kg 6 @ 5000 IU/kg	Collapse, laboured respiration, salivation, tremors in dogs prior to death. Conjunctival hyperemia. Increased water consumption and decreased body weight gain in 500 IU/kg males.	<u>Increased</u> RBC, Hgb, Hct platelet, reticulocyte, urine volume, urine occult blood, SGOT, LDH, BUN, TL, TCHO, PL, CPK @ 100, 500 IU/kg. <u>Decreased</u> Glucose	Enlarged spleens, kidneys swollen and increased weight @ 100, 500 IU/kg. Extramedullary hematopoiesis and megakaryopoiesis in spleen. Moderate to marked congestion of kidneys associated with glomerular dilation and slight to moderate round-cell infiltration. Dilation and thickening of Bowman's capsule with or without fibrosis. Slight to marked fibrosis of the femoral and sternal marrow cavities @ 100, 500 IU/kg.
Dogs (Neonates) i.v. 4 weeks/ 4 week recovery	No dose-related deaths.	None	<u>Increased</u> Hct in females. Slight RBC through Day 28. During withdrawal phase, RBC decreased @ 500 IU/kg but by Day 55 comparable to controls.	Enlarged spleens. Extra-medullary hematopoiesis. Erythroid hyperplasia of bone marrow.
Cynomolgus monkeys i.v. and s.c. (@ 1000 IU/kg only) 13 week/10 week recovery	1 @ 0 IU/kg 1 @ 500 IU/kg 1 @ 1000 IU/kg	Reddening of facial skin in females @ 100, 500, 1000 IU/kg. Minimal reduction in body weight and food consumption in females @ 1000 IU/kg (i.v.). Dilation of eye vessels @ 500, 1000 IU/kg (i.v. and s.c.).	<u>Increased</u> RBC, Hgb, PCV and reticulocytes.	Necropsy report not included in interim report.
Rhesus Monkeys (Adults Neonates) i.v. 6 week/10 week recovery	None	None	<u>Increased</u> RBC - adults only. Reversible.	Necropsy not performed.



The carcinogenic potential of epoetin alfa has not been evaluated.

There was no indication of teratogenicity in rats or rabbits and no toxicity in F1 and F2 generations at doses up to 500 IU/kg. In male and female rats treated intravenously with epoetin alfa, there was a trend for slightly increased fetal wastage at doses of 100 and 500 IU/kg/day. Fertility, teratology and reproductive performance studies are summarized in Tables 2.12 and 2.13

**Table 2.12: Fertility and Reproductive Performance and Teratology Studies Protocol Summary/Results**

Study	Toxicity	Embryo/Fetal Toxicity	Teratogenicity
Segment I Intravenous Fertility and Reproductive Performance in Sprague-Dawley Rats 0 (vehicle), 20, 100, 500 IU/kg/day. Treated animals mated to treated animals	<u>Yes</u> - Lethality in 1 male @ 100 IU/kg and 8 males @ 500 IU/kg. Mild redness of pinna and limbs, reduced food consumption and body weight gain, gastric erosion @ 100, 500 IU/kg Increased weight of heart, lungs, kidneys, spleen and adrenals due to pharmacologic effect of epoetin alfa @ 100 and 500 IU/kg. Decrease in thymus weights in males @ 500 IU/kg.	Reduced body weights and delays in fetal ossification @ 20, 100, 500 IU/kg were associated with reductions in maternal weight gain. Increase in resorptions @ 500 IU/kg and placental remnants @ 100, 500 IU/kg and dead implantations @ 500 IU/kg.	<u>No</u> - No external, visceral or skeletal abnormalities.
Segment II Intravenous Teratogenicity in Sprague-Dawley rats, 0 (vehicle), 20, 100, 500 IU/kg	<u>No</u> - Increased spleen, liver, left adrenal weights, @ 500 IU/kg due to pharmacologic effect epoetin alfa.	<u>Yes</u> - Fetal body weights and delayed ossification @ 500 IU/kg.	<u>No</u> - No increase in anomalies.
Intravenous Teratogenicity in NZW rabbits, 0 (vehicle), 20, 100, 500 IU/kg	<u>Yes</u> - Slight decrease in body weight and food consumption @ 500 IU/kg.	No	<u>No</u> - No increase in anomalies.

**Table 2.13: Perinatal and Postnatal Studies Protocol Summary/Results**

Study	Maternal/Toxicity	Embryo/Fetal Toxicity	Parturition/Neonatal Growth and Survival
Segment III Intravenous Peri- and Postnatal in Slc:SD rats. 0 (Vehicle), 20, 100, 500 IU/kg	<u>Yes</u> Redness of limbs and pinnae. Slight increase in body weight @ 100 and 500 IU/kg during lactation. Decrease in food consumption @ 500 IU/kg. Increased spleen and lung weights @ 100 and 500 IU/kg and dark red spots in stomach and increased heart weight @ 500 IU/kg.	<u>Yes</u> Reduced body weights and delayed ossification, appearance of abdominal hair, eyelid opening and decrease in the number of caudal vertebrae (day 22 postpartum) @ 500 IU/kg.	All F <sub>1</sub> , F <sub>2</sub> generation normal.

In a series of mutagenicity studies, epoetin alfa failed to induce bacterial gene mutations (Ames Test), chromosomal aberrations in mammalian cells, gene mutation at the HGPRT locus or micronuclei in mice (see Table 2.14). Epoetin alfa was not pyrogenic when tested in rabbits.

**Table 2.14**

Test Type	Study	Dose Levels	Response
Ames	<i>Escherichia coli</i> (WP2, Hcr <sup>-</sup> ) <i>Salmonella typhimurium</i> (TA1535, TA100, TA1537, TA98)	312.5, 625, 1250, 5000 IU/plate in presence and absence of S-9 mix.	Negative
Chromosomal Aberration	Chinese Hamster Lung Cells	1136, 2273, 4545 IU/mL in the absence of, and 947, 1894, 3788 IU/mL in the presence of S-9 mix.	Negative
<i>in vivo</i> Micro-nucleus	Slc:ddY male Mice	1.25 x 10 <sup>5</sup> , 2.5 x 10 <sup>5</sup> , 5 x 10 <sup>5</sup> IU/kg	
Gene Mutation at the HGPRT locus	Chinese Hamster Ovary Cells	625, 1250, 2500, 5000 IU/mL with and without S-9 mix.	Negative

Results of antigenicity studies indicated weak to strong evidence of antibody formation in rabbits and guinea pigs, but not in mice. The reason for this difference in the mouse is not clear, but anaphylaxis is not anticipated when administered to humans since the test material is of human origin. Seizure threshold potential in mice, water content of brain tissue and electrolyte distribution in rats were unaffected by intraperitoneal doses of epoetin alfa. The results of these and other special studies are summarized in Table 2.15.

**Table 2.15: Special Studies**

<u>Key to Abbreviations</u>			
RBC	Red blood cells	SGOT	Serum glutamic-oxaloacetic transaminase
Hct	Hematocrit	LDH	Lactic acid dehydrogenase
Hgb	Hemoglobin	CPK	Creatinine phosphokinase
MCH	Mean corpuscular hemoglobin	A/G Ratio	Albumin/Globulin Ratio
MCV	Mean corpuscular volume	M:E Ratio	Myeloid:Erythrocytes Ratio
MCHC	Mean corpuscular hemoglobin concentration	TL	Total lipids
APTT	Activated partial thromboplastin time	TCHO	Total cholesterol
		PL	Phospholipid
		PT	Prothrombin time

Test Type	Study	Dose Levels	Average Response
Seizure Threshold	CF1 mice 20/group ip 3x/week for 3 weeks. Once a week after the third dose, i.v. infusion with 0.5% metrazole.	0 (saline), 0 (vehicle + saline), 450 or 1500 IU/kg	No changes in seizure threshold.
Electrolyte and Brain Water Distribution	Sprague Dawley Male Rats 15/group i.p. 3x/week for 3 weeks 5/group sacrificed after 3, 6, and 9 doses.	0 (saline), 0 (vehicle + saline), 450 or 1500 IU/kg.	No effect in electrolyte or water content of brain tissue, plasma or spinal fluid. Increased spleen and liver weight and Hct @ 1500 IU/kg. Increased Hct @ 450 IU/kg.
Antigenicity 72-Hr Passive Cutaneous Anaphylaxis (PCA) Test	C <sub>3</sub> H/He and BALB/C mice 10/group i.p. 1x/week for 2 weeks	100, 1000 IU/kg animals + 2 mg alum/animal	Only one mouse (C <sub>3</sub> H/He) in 1000 IU/animal gave positive response.
4-Hr Passive Cutaneous Anaphylaxis (PCA) Test	Hartley Guinea Pigs 5/group, i.v. 2x/week for 2 weeks 5/group s.c. 2x/week for 4 weeks.	100, 1000 IU/kg 14000 IU/kg + Freund's Complete Adjuvant (FCA)	Negative PCA test for 100 and 1000 IU/kg but 50 positive for 14000 IU/kg + FCA.
Active Systemic Anaphylaxis test	Hartley Guinea Pigs 5/group i.v. 2x/week for 2 weeks s.c. 2x/week for 4 weeks then challenged 27 (100, 1000 IU/kg groups) or 20 (14000 IU/kg + Freund's Complete Adjuvant [FCA] group) later with 50,000 IU/animal.	100, 1000 IU/kg 14000 IU/kg + FCA	Remarkable anaphylactic reaction at 1000 IU/kg and 14000 IU/kg.
Cutaneous Reaction	Hartley Guinea Pigs from PCA test were challenged 19 (100, 1000 IU/kg groups) or 26 (14000 IU/kg + FCA group) days later with intradermal doses of 1000 or 10,000 IU/animal.	100, 1000 IU/kg 14000 IU/kg + FCA	Positive reaction at 1000 IU/kg and 14000 IU/kg + FCA
Antigenicity study Passive Hemagglutination (PHA) + 4-Hr Passive Cutaneous Anaphylaxis (PCA) tests	New Zealand White Female Rabbits; 4/group i.m. and s.c. 2x/week for 4 weeks i.m. and s.c. 1x week for 4 weeks	100, 1000 IU/kg 2000 IU/kg + FCA	Hemagglutinating antibodies were detected in sera of 2000 IU/kg + FCA. Antibodies detected 4-Hr PCA in sera of almost all epoetin alfa-treated rabbits.

**Table 2.15: Special Studies (cont'd)**

Test Type	Study	Dose Levels	Average Response
Pyrogenicity Study	New Zealand White Rabbits i.v. 3/group single dose Pyrogenicity measured by temperature and limulus amoebocyte lysate test.	0 (saline), 0 (vehicle), 200, 2000, or 20000 IU/kg Positive control <i>E. coli</i> , LPS.	Negative for both temperature and endotoxin.
Primary Eye Irritation	New Zealand White Rabbits 3/group single ocular dose followed by irrigation 6/group single ocular dose no irrigation.	12,100 U/animal	Classified as non- irritating.
Hematologic	Dogs Beagle 3/sex/group, i.v. administration for 4 weeks then 15 week recovery.	0 (vehicle), 28, 2800 IU/kg 3x week/4 weeks Group 4 pretreated with 40 mg/kg iron dextran for 2 weeks prior to first dose of epoetin alfa @ 2800 IU/kg	No drug-related deaths. <u>Increased</u> RBC, PCV, Hgb, reticulocytes, and platelets. <u>Decreased</u> MCV/MCHC/ MCH, serum glucose and BUN @ 2800 IU/kg with and without iron supplement. Low dose only slight increase RBC, PCV, and Hgb. No other dose- related effects.

Treatment with 1000 IU/kg/day epoetin alfa for 90 days did not cause myelofibrosis in Cynomolgus monkeys. Gastric erosion of the glandular stomach has been observed in the rat and in one high-dose (1000 IU/kg/day) monkey. Kidney changes involving the Bowman's capsule were also observed in dogs after daily injection for a year. Anaphylaxis noted in dogs, rabbits and guinea pigs is thought to be due to a natural reaction to any foreign protein in these animals.

## REFERENCES

1. Egrie JC, Stickland TW, Lane J, et al. Characterization and biological effects of recombinant human erythropoietin. *Immunobiology* 1986;72:213-224.
2. Graber SE, Krantz SB. Erythropoietin and the control of red cell production. *Ann Rev Med* 1978; 29:51-66.
3. Eschbach JW, Adamson JW. Anemia of end-stage renal disease (ESRD). *Kidney Int* 1985;28:1-5.
4. Eschbach JW, Egrie JC, Downing MR, Browne JK, Adamson JW. Correction of the anemia of end-stage renal disease with recombinant human erythropoietin. *N Engl J Med* 1987;316:73-78.
5. Eschbach JW, Adamson JW, Cooperative Multicenter r-HuEPO Trial Group. Correction of the anemia of hemodialysis (HD) patients with recombinant human erythropoietin (r-HuEPO): results of a multicenter study. *Kidney Int* 1988;33:189.
6. Eschbach JW, Egrie JC, Downing MR, Browne JK, Adamson JW. The use of recombinant human erythropoietin (r-HuEPO): effect in end-stage renal disease (ESRD). In: Friedman, Beyer, DeSanto, Giordano, editors. *Prevention of chronic uremia*. Philadelphia, PA: Field and Wood Inc, 1989:148-155.
7. Egrie JC, Eschbach JW, McGuire T, Adamson JW. Pharmacokinetics of recombinant human erythropoietin (r-HuEPO) administered to hemodialysis (HD) patients. *Kidney Int* 1988:262.
8. Lundin AP, Delano BG, Stein R, Quinn RM, Friedman EA. Recombinant human erythropoietin (r-HuEPO) treatment enhances exercise tolerance in hemodialysis patients. *Kidney Int* 1988;33:200.
9. Paganini E, Garcia J, Ellis P, Bodnar D, Magnussen M. Clinical sequelae of correction of anemia with recombinant human erythropoietin (r-HuEPO); urea kinetics, dialyzer function and reuse. *Am J Kidney Dis* 1988;11:16.
10. Delano BG, Lundin AP, Golansky R, Quinn RM, Rao TKS, Friedman EA. Dialyzer urea and creatinine clearances not significantly changed in r-HuEPO treated maintenance hemodialysis (MD) patients. *Kidney Int* 1988;33:219.
11. Stivelman J, Van Wyck D, Ogden D. Use of recombinant human erythropoietin (r-HuEPO) with high flux dialysis (HFD) does not worsen azotemia or shorten access survival. *Kidney Int* 1988;33:239.
12. Lim VS, DeGowin RL, Zavala D, Kirchner PT, Abels R, Perry P, Fangman J. Recombinant human erythropoietin treatment in pre-dialysis patients: a double-blind placebo controlled trial. *Ann Int Med* 1989;110:108-114.

13. Stone WJ, Graber SE, Krantz SB, et al. Treatment of the anemia of pre-dialysis patients with recombinant human erythropoietin: a randomized, placebo-controlled trial. *Am J Med Sci* 1988;296:171-179.
14. Eschbach JW, Abdulhadi MH, Browne JK, et al. Recombinant human erythropoietin in anemic patients with end-stage renal disease. *Ann Intern Med* 1989;111:12.
15. Evans RW, Radar B, Manninen DL, et al. The quality of life of hemodialysis recipients treated with recombinant human erythropoietin. *JAMA* 1990;263:6.
16. Williams SV, Nash DB, Goldfarb N. Differences in mortality from coronary artery bypass graft surgery at five teaching hospitals. *JAMA* 1991;266:810-815.
17. Surgenor DM, Wallace EL, Churchill WH, et al. Red cell transfusions in coronary artery bypass surgery. *Transfusion* 1992;32:458-464.
18. Hannan EL, Kilburn H, O'Donnell JF, et al. Adult open heart surgery in New York State--An analysis of risk factors and hospital mortality rates. *JAMA* 1990;264:2768-2774.
19. Imperiale TF, Speroff T. A meta-analysis of methods to prevent venous thromboembolism following total hip replacement. *JAMA* 1994;271:1780-1785.
20. The RD Heparin Arthroplasty Group. RD Heparin combined with warfarin for prevention of venous thromboembolic disease following total hip or knee arthroplasty. *J Bone Jt Surg* 1994;76:1174-1185.
21. Hull R, Raskob G, Pineo G, et al. A comparison of subcutaneous low-molecular-weight heparin with warfarin sodium for prophylaxis against deep-vein thrombosis after hip or knee implantation. *N Engl J Med* 1993;329:1370-1376.
22. Kraay MJ, Goldberg VM, Herbener TE. Vascular ultrasonography for deep venous thrombosis after total knee arthroplasty. *Clin Orthop Related Res* 1993;286:18-26.
23. Mohr DN, Silverstein MD, Murtaugh PA, Harrison JM. Prophylactic agents for venous thrombosis in elective hip surgery: meta-analysis of studies using venographic assessment. *Arch Intern Med* 1993;153:2221-2228.
24. Muirhead N, Churchill DN, Goldstein M, et al. Comparison of subcutaneous and intravenous recombinant human erythropoietin for anemia in hemodialysis patients with significant comorbid disease. *Am J Nephrol* 1992;12:303-10.
25. Canaud B, Benhold I, Delons S, et al. What is the optimum frequency of administration of r-HuEPO for correcting anemia in hemodialysis patients. *Dialysis & Transplantation* 1995;24(6):306-329.
26. Muirhead N, Laupacis A, Won C. Erythropoietin for anaemia in haemodialysis patients: results of a maintenance study (The Canadian Erythropoietin Study Group). *Nephrol Dial Transplant* 1992;7: 811-816.

27. Amgen Inc. The effects of normal versus anemic hematocrit on outcomes of cardiac disease in dialysis patients. Data on File.
28. Lundin AP, Akerman MJH, Chesler RM, Delano BG, Goldberg N, Stein RA, et al. Exercise in hemodialysis patients after treatment with recombinant human erythropoietin. *Nephron* 1991;58:315-319.
29. Eschbach JW, Kelly MR, Galey NR, Abels RI, Adamson JU. Treatment of the anemia of progressive renal failure with recombinant human erythropoietin. *N Engl J Med* 1989;321:158-163.
30. The US Recombinant Human Erythropoietin Predialysis Study Group. Double-blind placebo-controlled study of the therapeutic use of recombinant human erythropoietin for anemia associated with chronic renal failure in predialysis patients. *Am J Kidney Dis* 1991;18(1):50-59.
31. Abels RI, Larholt KM, Krantz KD, Bryant EC. Recombinant human erythropoietin (r-HuEPO) for the treatment of the anemia of cancer. In: Murphy MJ Jr. ed. *Proceedings of the Beijing Symposium*. Dayton, Ohio: Alpha Med. Press. 1991:121-141.
32. Henry DH, Abels RI. Recombinant human erythropoietin in the treatment of cancer and chemotherapy-induced anemia: Results of double-blind and open-label follow-up studies. *Seminars in Oncology* 1994;21(2) Suppl 3:21-28.
33. Besarab A, Bolton WK, Browne JK, et al. The effects of normal as compared with low hematocrit values in patients with cardiac disease who are receiving hemodialysis and epoetin. *NEJM*. 1998;339:584-90.
34. Singh AK, Szczech L, Tang KL, et al. Correction of anemia with epoetin alfa in chronic kidney disease. *N Engl J Med* 2006;335(20):2085-2098.
35. Leyland-Jones B, Semiglazov V, Pawlicki M, et al. Maintaining normal hemoglobin levels with epoetin alfa in mainly nonanemic patients with metastatic breast cancer receiving first-line chemotherapy: a survival study. *J Clin Oncol* 2005;23(25):5960-5972.
36. Bohlius J, Wilson J, Seidenfeld J, et al. Recombinant Human Erythropoietins and Cancer Patients: Updated Meta-Analysis of 57 Studies Including 9353 Patients. *J Natl Cancer Inst*. 2006; 98:708-14.
37. Henke, M, Laszig, R, Rube, C, et al. Erythropoietin to treat head and neck cancer patients with anaemia undergoing radiotherapy: randomized, double-blind, placebo-controlled trial. *The Lancet*. 2003;362:1255-1260.
38. Wright JR, Ung YC, Julian JA, et al. Randomized, double-blind, placebo-controlled trial of Erythropoietin in non-small cell lung cancer with disease-related anemia. *J Clin Oncol* 2007;25(9):1-6.
39. [http://download.veritasmedicine.com/PDF/CR004621\\_ToplineResults.pdf](http://download.veritasmedicine.com/PDF/CR004621_ToplineResults.pdf)

## OTHER RELEVANT LITERATURE

1. Abels RI, Rudnick SA, and Tsai HC. Erythropoietin for zidovudine-induced anemia [reply]. *New Engl J Med* 1990;323:1070.
2. American College of Physicians. Practice strategies for elective red blood cell transfusion. *Ann Int Med* 1992;116(5):403-406.
3. Blumberg N, and Heal JM. Transfusion and host defenses against cancer recurrence and infection. *Transfusion* 1989;29:236-245.
4. Braun A, Ding R, Seidel C, et al. Pharmacokinetics of recombinant human erythropoietin applied subcutaneously to children with chronic renal failure. *Pediatr Nephrol* 1993;7:61-64.
5. Busch ORC, Hoyneck van Papendrecht MAW, Marquet RL, Jeekel J: Experimental and clinical results of perioperative treatment with recombinant human erythropoietin. *Pathophysiology Pharmacology of Erythropoietin*, Springer Verlag, Berlin, 1992; pp. 315-320.
6. Canadian Erythropoietin Study Group. The effect of recombinant human erythropoietin (r-HuEPO) upon quality of life and exercise capacity in anemic patients on chronic hemodialysis. *Clin Invest Med* 1989;12:B66 (R404).
7. Canadian Erythropoietin Study Group. The clinical effects and side-effects of recombinant human erythropoietin (r-HuEPO) in anemic patients on chronic hemodialysis. *Clin Invest Med* 1989;12:B66 (R403).
8. Canadian Erythropoietin Study Group. The effect of recombinant human erythropoietin (r-HuEPO) on quality of life and exercise capacity of patients on hemodialysis. *Br Med J* 1989;300:573-578.
9. Canadian Orthopedic Perioperative Erythropoietin Study Group. Effectiveness of perioperative recombinant human erythropoietin in elective hip replacement. *Lancet* 1993;341:1227-1232.
10. Carey J, Jackson B, Wallace M, Lederman M, Abels R. The effect of R-Hu erythropoietin on HIV P24 antigen expression in anemic zidovudine treated AIDS patients [abstract]. In: *Sixth International Conference on AIDS*, 1990:195. San Francisco: Univ. of California.
11. Churchill DN, ed. The Canadian Hemodialysis Morbidity Study Group. Morbidity in chronic hemodialysis patients. *Clin Invest Med* 1989;12:B66 (R407).
12. Cotes PM, Pippard MJ, Reid CDL, Winearls CG, Oliver DO, Royston JP. Characterization of the anemia of chronic renal failure and the mode of its correction by a preparation of human erythropoietin (r-HuEPO). An investigation of the pharmacokinetics of intravenous erythropoietin and its effects on erythrokinetics. *Quart J Med* 1989;70(262):113-137.



13. Danna RP, Rudnick SA, Abels RI. Erythropoietin therapy for the anemia associated with AIDS and AIDS therapy and cancer. In: Garnick MB, ed. Erythropoietin in clinical applications an international perspective, New York: Marcel Dekker 1990:301-324.
14. Egrie JC, Browne J, Lai P, Lin PK. Characteristics of recombinant monkey and human erythropoietin: Experimental approaches for the study of hemoglobin switching, Alan R Liss, Inc. 1985:339-350.
15. Egrie JC, Cotes PM, Lane J, Gaines Das RE, Tam RC. Development of radioimmunoassays for human erythropoietin using recombinant erythropoietin as tracer and immunogen. *J Immunol* 1987;99:235-241.
16. Erythropoietin reaches the pharmacy [editorial]. *Lancet* 1989;2:1252-1254.
17. Eschbach JW, Egrie JC, Downing MR, , et al. The Use of Recombinant Human Erythropoietin (r-HuEPO): Effect in End-Stage Renal Disease (ESRD). In: Friedman, Beyer, DeSanto, Giordano, editors. *Prevention Of Chronic Uremia.* , Philadelphia: Field and Wood Inc.; 1989. p. 148-155.
18. Eschbach JW, Kelly MR, Haley R, Abels RI, Adamson JW. Treatment of the anemia of progressive renal failure with recombinant human erythropoietin. *N Engl J Med* 1989;321:158-163.
19. Fischl M, Galpin JE, Levine JD, et al. Recombinant human erythropoietin for patients with AIDS treated with zidovudine. *NEJM* 1990;322:1488-1493.
20. Geva P and Sherwood JB. Pharmacokinetics of recombinant human erythropoietin (rHuEPO) in pediatric patients on chronic cycling peritoneal dialysis (CCPD). *Blood* 1991;78 (Suppl 1):91a.
21. Gough IR, Furnival CM, Schilder L, and Grove W. Assessment of the quality of life of patients with advanced cancer. *Eur J Cancer Clin Oncol* 1983;19:1161-1165.
22. Grimm P, Levana S, Neal MS, et al. Effects of recombinant erythropoietin on HLA sensitization and cell mediated immunity. *Kidney International* 1990;8:12-18.
23. Jabs K, Grant JR, Harmon W, et al. Pharmacokinetics of Epoetin alfa (rHuEPO) in pediatric hemodialysis (HD) patients. *J Am Soc Nephrol* 1991;2:380.
24. Kling PJ, Widness JA, Guillery EN, et al. Pharmacokinetics and pharmacodynamics of erythropoietin during therapy in an infant with renal failure. *J Pediatr* 1992;121:822-825.
25. Lin FK, Suggs S, Lin C-H, Browne JK, Smalling R, Egrie JC, et al. Cloning and expression of the human erythropoietin. *Gene. Proc. Natl. Acad. Sci.* 1985;82:7580-7584.

26. London GM, Zins B, Pannier B, Naret C, Berthelot J-M, Jacquot C, et al. Vascular changes in hemodialysis patients in response to recombinant human erythropoietin. *Kid. Int.* 1989;36:878-882.
27. Muirhead N, Wong C, editors. The Canadian Erythropoietin Study Group. Results of a maintenance study. *Clin Invest Med* 1989;12:B66 (R405).
28. Muller-Wiefel DE and Scigalla P. Specific problems of renal anemia in childhood. *Contrib Nephrol* 1988;66:71-84.
29. National Institutes of Health Consensus Development Conference: Perioperative red cell transfusion. *Transfusion Medicine Reviews* Jan 1989;III(1):63-68.
30. Neff MS, Kim KE, Persoff M, Onesti G, Swartz C. Hemodynamics of uremic anemia. *Circulation* 1971;43:876-883.
31. Porter S. Human Immune Response to Recombinant Human Proteins. *J Pharm Sci* 2001; 90: 1-11.
32. Richman DD, Fischl MA, Grieco MH, et al. The toxicity of azidothymidine (azt) in the treatment of patients with AIDS and AIDS-related complex. *N Engl J Med* 1987;317:192-197.
33. Rothstein P, Roye D, Verdisco L, Stern L. Preoperative use of erythropoietin in an adolescent Jehovah's Witness. *Anesthesiology* 1990;75:568-570.
34. Rutherford CJ, Schneider TJ, Dempsey H, Kirn DH, Brugnara C, Goldberg. Efficacy of different dosing regimens for recombinant human erythropoietin in a simulated perisurgical setting: the importance of iron availability in optimizing response. *Am J Med* 1994;96:139-45.
35. Scharer K, Klare B, Dressel P, et al. Treatment of renal anemia by subcutaneous erythropoietin in children with preterminal chronic renal failure. *Acta Paediatr* 1993;82:953-958.
36. Spivak JL. Recombinant human erythropoietin and its role in transfusion medicine. *Transfusion* 1994; 34(1):1-4.
37. Spivak JL, Barnes DC, Fuchs E, Quinn TC. Serum immunoreactive erythropoietin in HIV-infected patients. *JAMA* 1989;21(261):3104-3106.
38. Tartter I. Does blood transfusion predispose to cancer recurrence? *Am J Clin Oncol* 1989;12:169-172.
39. Van Wyck DB, Stivelman JC, Ruiz J, Kirilin LF, Katz MA, Ogden DA. Iron status in patients receiving erythropoietin for dialysis-associated anemia. *Kid Int* 1989;35:712-726.

40. Ward JW, Bush TJ, Perkins HA, et al. The natural history of transfusion-associated infections with human immunodeficiency virus. *NEJM* 1989;321:947-952.
41. Waymack JP, Alexander JW. Blood transfusions as an immunomodulator - a review. *Comp Immun Microbiol Infect Dis* 1986;9:177-183.
42. Winearls CG. Treatment of the anemia of chronic renal disease with recombinant human erythropoietin. *Drugs* 1989;38:342-345.
43. Winearls CG, Oliver DO, Pippard MJ, Reid C, Downing M, Cotes PM. Effect of human erythropoietin derived from recombinant DNA on the anemia of patients maintained by chronic hemodialysis. *The Lancet* 1986;2:1175-1178.

**PART III: CONSUMER INFORMATION**

**PrEPREX\***  
epoetin alfa

Sterile Solution

This leaflet is Part III of a three-part "Product Monograph" published when EPREX was approved for sale in Canada and is designed specifically for Consumers. Please read this carefully before you start to take your medicine. This leaflet is a summary and will not tell you everything about EPREX. Contact your doctor or pharmacist if you have any questions about the drug.

**INFORMATION ABOUT ANEMIA****What is anemia:**

Anemia is a condition where your blood contains a lower than normal number of red blood cells. Red blood cells contain a substance called hemoglobin that is responsible for transporting oxygen in your blood. As the number of red blood cells in your blood decreases, so does the amount of oxygen delivered to your body. Development of anemia is characterized by several symptoms, such as weakness, fatigue, shortness of breath, dizziness, poor concentration and chills. These symptoms over time can negatively impact your overall quality of life.

**What causes anemia:**

Anemia may be the result of several causes, including blood loss or nutritional deficiencies. Anemia due to blood loss must be investigated and treated by a physician. Nutritional anemia caused by a deficiency of iron, vitamin B<sub>12</sub> or folic acid can be treated with dietary supplements. Anemia can also be caused by chronic disease, such as cancer or kidney (renal) disease.

**Anemia in Chronic Kidney Disease:**

In chronic kidney disease, anemia results when the kidneys are unable to manufacture enough natural erythropoietin to stimulate the bone marrow to produce more red blood cells.

**Anemia in Cancer:**

Anemia in cancer can be caused by the chemotherapy regimens used to treat the cancer. The toxic effects of chemotherapy reduce the body's ability to produce erythropoietin as well as the bone marrow's ability to respond and make more red blood cells. As a result, not enough red blood cells are produced and the patient becomes anemic.

**ABOUT THIS MEDICATION****What the medication is used for:**

EPREX is used to increase the production of red blood cells and to decrease the need for red blood cell transfusion. The dose should be gradually adjusted to achieve this goal. EPREX may be used in adults and children with chronic kidney disease; adults who have HIV-infection and are receiving a drug called zidovudine; adult cancer patients receiving chemotherapy; and adults scheduled to undergo major elective surgery.

**What it does:**

EPREX is a protein made in the laboratory which acts like a substance naturally made in the human body called erythropoietin. Erythropoietin controls the production of red blood cells in the body.

**When it should not be used:**

You should not use EPREX if:

- you are allergic to any of the ingredients in the product;
- you make antibodies (develop pure red cell aplasia [PRCA]) following treatment with any erythropoiesis-stimulating agent;
- you have uncontrolled high blood pressure;
- you cannot for any reason receive adequate antithrombotic treatment.

**What the medicinal ingredient is:**

epoetin alfa

**What the nonmedicinal ingredients are:**

albumin (human) or glycine and polysorbate 80 as stabilizers, sodium chloride, water for injection and either sodium citrate and citric acid, or sodium phosphate

Single-use vials and pre-filled syringes do not contain preservatives. The 20,000 IU/mL multi-use vial contains benzyl alcohol as a preservative.

**What dosage forms it comes in:**

**Single-use vials:** 1,000 IU/0.5 mL, 2,000 IU/mL, 4,000 IU/mL, 10,000 IU/mL, 40,000 IU/mL

**Multi-use vial:** 20,000 IU/mL

**Single-use pre-filled syringes with PROTECS\* needle**

**guard:** 1,000 IU/0.5 mL, 2,000 IU/0.5 mL, 3,000 IU/0.3 mL, 4,000 IU/0.4 mL, 5,000 IU/0.5 mL, 6,000 IU/0.6 mL, 7,000 IU/0.7 mL, 8,000 IU/0.8 mL, 9,000 IU/0.9 mL, 10,000 IU/mL, 20,000 IU/0.5 mL, 30,000 IU/0.75 mL, 40,000 IU/mL

To reduce the risk of accidental needle sticks to users, each pre-filled syringe is equipped with the PROTECS needle guard that is automatically activated to cover the needle after complete delivery of the syringe content.

## WARNINGS AND PRECAUTIONS

### Serious Warnings and Precautions

#### ALL PATIENTS

- To minimize the risks for death and serious cardiovascular (heart and blood vessel-related) side effects, your doctor will follow the recommended dosage for each indication.
- The multi-use preserved formulation contains benzyl alcohol. Benzyl alcohol has been reported to be associated with an increased incidence of neurological and other complications in premature infants which are sometimes fatal.
- Patients with uncontrolled hypertension should not be treated with EPREX epoetin alfa; blood pressure should be controlled adequately before initiation of therapy.
- EPREX epoetin alfa should be used with caution in patients with a history of seizures.
- During hemodialysis, patients treated with EPREX epoetin alfa may require increased anticoagulation with heparin to prevent clotting of the artificial kidney. Patients with pre-existing vascular disease should be monitored closely.
- The multi-use formulation of EPREX epoetin alfa contains albumin, a derivative of human blood. The theoretical risk of transmission of Creutzfeldt-Jakob disease is considered extremely remote. No cases of viral disease transmission or CJD have ever been identified as a result of albumin use.
- Hemoglobin levels during EPREX treatment should not be higher than 120 g/L (may not apply to surgery patients)
- If you undergo surgery while taking EPREX, your physician may prescribe a blood thinner, as appropriate for the surgical procedure, to prevent blood clots.
- Antibody-mediated pure red cell aplasia (PRCA) has been reported after months to years of treatment with recombinant erythropoietins. If you develop PRCA, you may suddenly become severely anemic and this could result in a dependency on blood transfusions.

#### CHRONIC RENAL FAILURE PATIENTS

- If your hemoglobin is kept too high, you have an increased chance of heart attack, stroke, heart failure, blood clots and death. Your doctor should try to keep your hemoglobin between 100 and 120 g/L

#### CANCER PATIENTS

- If you are a cancer patient and your hemoglobin is kept too high (over 120 g/L),
  - your tumor may grow faster,
  - you have an increased chance of heart attack, stroke, blood clots and death.
- Your doctor should use the lowest dose of EPREX needed to avoid red blood cell transfusions.
- In some instances, red blood cell transfusion should be the preferred treatment option.
- Once you have finished your chemotherapy course, EPREX should be discontinued.

#### BEFORE taking EPREX:

- Tell your doctor about any medical problems and about any allergies you have or have had in the past.
- Tell your doctor if you have or have had high blood pressure, seizures, blood clots, liver disease, porphyria, or gout.
- While you are treated with EPREX your doctor will need to check your blood pressure. Your blood pressure will be monitored carefully and any changes outside of the guidelines that your doctor has given you must be reported. If your blood pressure increases, you may need medication to lower it. If you already take blood pressure medication, your doctor may increase the amount.
- Your doctor will also measure your serum iron levels, red blood cell levels and other factors in your blood, prior to starting and during treatment with EPREX, as deemed appropriate.
- If you are on dialysis, your dialysis prescription may need to be changed while you are being treated with EPREX. Your doctor will take blood tests to determine if any change is needed. Your doctor may also need to adjust any medication you receive to prevent blood clotting.
- Tell your doctor if you are pregnant, if you think you might be pregnant, or if you are trying to become pregnant.
- Tell your doctor if you are breast-feeding.
- In many women with severe kidney failure, the monthly period may stop. When these women take EPREX they may restart their monthly cycle. If you are a woman with kidney disease, you should discuss contraception with your doctor.
- In patients with kidney disease, due to the possibility of an increase in blood pressure, there is a small chance of having a seizure when therapy starts. During the initial phase of treatment, your doctor may advise you to avoid driving, using machines or doing anything else that could be dangerous if you are not alert.
- If you are a home dialysis patient, you should continue to check your access, as your doctor or nurse has shown you, to make sure it is working. Be sure to let

your healthcare professional know right away if there is a problem.

- If you are a cancer patient you should be aware that EPREX is a red blood cell growth factor and in some circumstances your cancer may grow faster. You should discuss treatment options for your anemia with your doctor.

## INTERACTIONS WITH THIS MEDICATION

Tell your doctor about all medications you are using, including those obtained without a prescription and any other remedies or dietary supplements. It is especially important that your doctor know if you are taking high blood pressure medication.

## PROPER USE OF THIS MEDICATION

**Follow your doctor's instructions about when and how to take this medication.**

EPREX Sterile Solution is injected under the skin (subcutaneously) into the arms, legs or abdomen.

While you are receiving EPREX, your doctor will measure your red blood cells. Your doctor will use this information to adjust the dose to the amount right for you.

**Do not shake** EPREX Sterile Solution. The solution in the vial or pre-filled syringe should always be clear and colourless. Do not use EPREX Sterile Solution if the contents of the vial or pre-filled syringe appear discoloured or cloudy, or if the vial or pre-filled syringe appears to contain lumps, flakes, or particles. If the vial or pre-filled syringe has been shaken vigorously, the solution may appear to be frothy and should not be used.

Single-use vials and pre-filled syringes of EPREX Sterile Solution do not contain preservatives and therefore are to be used once and discarded. Any unused portion of a vial or pre-filled syringe should not be used. Multi-use vials contain benzyl alcohol as a preservative and may be re-entered. Multi-use vials should be discarded 30 days after first entry.

### **Overdose:**

If you accidentally take too much EPREX, obtain immediate medical advice.

### **Missed dose:**

If you miss a dose, contact your doctor for instructions.

## **Preparing and Injecting the Dose**

**Important: To help avoid contamination and possible infection, follow these instructions exactly.**


## **Single-Use and Multi-Use Vials**

**Use the Correct Syringe:** Your doctor has instructed you on how to give yourself the correct dosage of EPREX. This dosage will usually be measured in units per milliliter or ccs. It is important to use a syringe that is marked in tenths of milliliters (for example, 0.1, 0.2, etc., mL or cc). Failure to use the proper syringe can lead to a mistake in dosage, and you may receive too much or too little EPREX. Only use disposable syringes and needles as they do not require sterilization; they should be used once and disposed of as instructed by your doctor.


1. **Take a vial out of the refrigerator.** The liquid needs to come to room temperature. This usually takes between 15 to 30 minutes.
2. **Check the vial,** to make sure it is the right dose, has not passed its expiry date, is not damaged, and the liquid is clear and not frozen. When using the multi-use vial, record the date of first entry for each vial on the inside flap of the box.
3. **Assemble the other items** you will need for your injection: a syringe, needle and antiseptic swabs.
4. **Wash your hands.**
5. **Remove the cap from the top of the vial but do not remove the rubber stopper.** Clean the rubber stopper with an antiseptic swab.



6. **Remove the needle cover from the syringe.** Pull the syringe plunger out to fill the syringe with an amount of air equal to the amount of liquid prescribed by your doctor.
7. **Put the vial on a flat surface and push the syringe needle through the rubber stopper.** Push the syringe plunger down to add the air into the vial.
8. **Turn the vial and the syringe upside down.** Pull on the syringe plunger to fill the syringe with the correct amount of liquid from the vial. It is important that the needle is always in the liquid in order to prevent air bubbles forming in the syringe.
9. **Remove the syringe and needle from the vial.** Hold the syringe with the needle pointing up to see if it has any air bubbles inside. If there are air bubbles gently press the plunger until all of the air (but none of the liquid) has been removed.



10. **Choose an injection site.** Good sites are the top of the thigh and around the tummy (abdomen) but away from the navel. Vary the site from day to day.



11. **Use an antiseptic swab on the injection site,** to disinfect it.

12. **Pinch a fold of skin** between your thumb and index finger. Don't squeeze it.




13. **Push the syringe needle in fully.** Your doctor or nurse may have shown you how to do this.



14. **Check that you haven't punctured a blood vessel.** Pull back slightly on the plunger. If you see blood, take the syringe out and choose a new injection site.

15. **Push the plunger with your thumb as far as it will go to inject the liquid.** Push it slowly and evenly, keeping the skinfold pinched.

16. **When the plunger is pushed as far as it will go,** take out the needle and let go of the skin.



17. **Press an antiseptic swab** over the injection site for a few seconds after the injection.


18. **Dispose of your used syringe** in a safe container (see **Disposal of Syringes**) and dispose of your vial.

### Single-Use Pre-filled Syringe with PROTECS needle guard

To reduce the risk of accidental needle sticks to users, each pre-filled syringe is equipped with the PROTECS needle guard that is automatically activated to cover the needle after complete delivery of the syringe content.

1. **Take a syringe out of the refrigerator.** The liquid needs to come to room temperature. This usually takes between 15 to 30 minutes.
2. **Check the syringe** to make sure it is the right dose, has not passed its expiry date, is not damaged, and the liquid is clear and not frozen.


3. **Choose an injection site.** Good sites are the top of the thigh and around the tummy (abdomen) but away from the navel. Vary the site from day to day.



4. **Wash your hands.** Use an antiseptic swab on the injection site, to disinfect it.
5. **Take the cover off the syringe** by holding the barrel and pulling the cover off carefully without twisting it.

Don't push the plunger, touch the needle or shake the syringe.

6. **Pinch a fold of skin** between your thumb and index finger. Don't squeeze it.




7. **Push the needle in fully.** Your doctor or nurse may have shown you how to do this.



8. **Check that you haven't punctured a blood vessel.** Pull back slightly on the plunger. If you see blood, take the syringe out and choose a new injection site.

9. **Push the plunger with your thumb as far as it will go to inject all of the liquid.** Push it slowly and evenly, keeping the skinfold pinched. The needle guard will not activate unless the entire dose is given.

10. **When the plunger is pushed as far as it will go,** take out the needle and let go of the skin.



11. **Take your thumb off the plunger.** Allow the syringe to move up until the entire needle is covered by the needle guard.

12. **Press an antiseptic swab** over the injection site for a few seconds after the injection.

13. **Dispose of your used syringe** in a safe container (see **Disposal of Syringes**).

14. **Only take one dose of EPREX from each syringe.** If any liquid remains in the syringe after an injection, the syringe should be properly disposed of, not reused (see **Disposal of Syringes**).

### Disposal of Syringes

1. Place all used needles and syringes in a hard plastic container with a screw-on cap, or a metal container with a plastic lid, such as a coffee can properly labelled as to content. If a metal container is used, cut a small hole in the plastic lid and tape the lid to the metal container. If a hard-plastic container is used, always screw the cap on tightly after each use. When the container is full, tape around the cap or lid, and dispose of according to your doctor's instructions.

2. Do not use glass or clear plastic containers (or any other container) that will be recycled or returned to a store.
3. Always store the container out of the reach of children.
4. Please check with your doctor, nurse, or pharmacist for other suggestions.

## **SIDE EFFECTS AND WHAT TO DO ABOUT THEM**

Any medicine may have unwanted effects. The side effects you might experience may vary depending on the reason you are taking EPREX. Tell your doctor or pharmacist about any unusual sign or symptom whether listed or not. The side effects reported most often in all patients receiving EPREX are flu-like symptoms such as dizziness, drowsiness, fever, headache, muscle and joint pain and weakness, and gastrointestinal disturbances such as nausea, vomiting and diarrhea. Redness, burning and pain at the place where EPREX is given have also been reported.

Be sure to tell your doctor if you have an increase in headaches or develop unusual headaches, as this may be a sign of increased blood pressure.

There is a possible association of a worsening of increased blood pressure if red blood cell production occurs too rapidly. Your doctor may need to reduce your dose of EPREX and initiate or increase blood pressure medication. Signs and symptoms of serious cardiovascular events may include but are not limited to: chest pain, leg pain and swelling, shortness of breath, sudden weakness, numbing or tingling of face, arm or leg, an increase in headaches or severe sudden headaches, loss of vision or loss of speech, lightheadedness.

Patients experiencing these signs or symptoms should contact their doctor immediately.

Patients should NOT discontinue their medication without consulting their doctor first.

## **SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM**

Additional side effects which have been reported more often in chronic renal failure patients than other patients include increases in blood pressure, clotted access, seizures and pure red cell aplasia (PRCA). PRCA is a condition in which severe and sudden anemia (characterized by symptoms such as severe tiredness/fatigue, and shortness of breath on mild exertion) develops due to failure of the bone marrow to produce red blood cells. PRCA could result in a dependency on blood transfusions. Should you be diagnosed with PRCA, your doctor will stop your EPREX therapy and may initiate treatment with drugs and/or blood transfusions to help increase your red blood cell count.

If you develop signs of allergy such as difficulty breathing, hives, itching, rash, or swelling of the throat, face, eyelids, mouth or tongue, discontinue the use of EPREX and contact your doctor or obtain medical help immediately.

*This is not a complete list of side effects. For any unexpected effects while taking EPREX, contact your doctor or pharmacist.*

## **HOW TO STORE IT**

- Store unopened in a refrigerator between 2 and 8 degrees centigrade. Do not freeze. Protect from light.
- Do not use this product after the expiry date written on the package.
- Keep this and all medicines in a safe place away from children.
- If you are using EPREX at home, it is important that the vial or syringe be stored in your refrigerator although not in the freezer compartment. EPREX should not be frozen. Do not use EPREX if it has been frozen. Leave the EPREX vial or syringe to stand for about 15 to 30 minutes until it reaches room temperature prior to using it.
- When the pre-filled syringe or single-use vial is about to be used, it may be removed from the refrigerator and stored at room temperature (not above 25°C) for one single period of maximum 7 days.
- The multi-use vial may be re-entered and therefore should be stored in the refrigerator following each injection.
- Discard the multi-use vial 30 days after the date of first entry into the vial, which you recorded on the inside flap of the box.



## **REPORTING SUSPECTED SIDE EFFECTS**

To monitor drug safety, Health Canada through the Canada Vigilance Program collects information on serious and unexpected side effects of drugs. If you suspect you have had a serious or unexpected reaction to this drug you may notify Canada Vigilance :

By toll-free telephone: 866-234-2345  
By toll-free fax: 866-678-6789

Online: [www.healthcanada.gc.ca/medeffect](http://www.healthcanada.gc.ca/medeffect)  
By email: [CanadaVigilance@hc-sc.gc.ca](mailto:CanadaVigilance@hc-sc.gc.ca)

By regular mail:  
Canada Vigilance National Office  
Marketed Health Products Safety and Effectiveness  
Information Bureau  
Marketed Health Products Directorate  
Health Products and Food Branch  
Health Canada  
Tunney's Pasture, AL 0701C  
Ottawa ON K1A 0K9

***NOTE: Should you require information related to the management of the side effect, please contact your health care provider before notifying Canada Vigilance. The Canada Vigilance Program does not provide medical advice.***

## **MORE INFORMATION**

This document plus the full Product Monograph, prepared for health professionals, can be found at:

<http://www.janssen-ortho.com>

or by contacting the sponsor, Janssen-Ortho Inc., at:

1-800-567-3331

This leaflet was prepared by

Janssen-Ortho Inc.  
Toronto, Ontario M3C 1L9

Last revised: July 2009