

PRODUCT MONOGRAPH

Pr

MELOXICAM

Meloxicam Tablets BP

7.5 mg and 15 mg

Non-Steroidal Anti-Inflammatory Drug (NSAID)

Manufactured by:
Cobalt Pharmaceuticals Inc.
6500 Kitimat Road
Mississauga, Ontario
L5N 2B8

Date of Preparation:
February 3, 2010

Control# : 136262

Table of Contents

PART I: HEALTH PROFESSIONAL INFORMATION.....	3
SUMMARY PRODUCT INFORMATION	3
INDICATIONS AND CLINICAL USE.....	3
CONTRAINDICATIONS	4
WARNINGS AND PRECAUTIONS.....	5
ADVERSE REACTIONS.....	14
DRUG INTERACTIONS	20
DOSAGE AND ADMINISTRATION.....	24
OVERDOSAGE	25
ACTION AND CLINICAL PHARMACOLOGY	25
STORAGE AND STABILITY.....	30
DOSAGE FORMS, COMPOSITION AND PACKAGING	30
PART II: SCIENTIFIC INFORMATION.....	31
PHARMACEUTICAL INFORMATION.....	31
CLINICAL TRIALS.....	32
DETAILED PHARMACOLOGY	35
TOXICOLOGY	35
REFERENCES	37
PART III: CONSUMER INFORMATION.....	40

Pr **MELOXICAM**

Meloxicam Tablets BP

PART I: HEALTH PROFESSIONAL INFORMATION

SUMMARY PRODUCT INFORMATION

Route of Administration	Dosage Form / Strength	Clinically Relevant Nonmedicinal Ingredients
Oral	Tablet, 7.5 mg & 15 mg	Colloidal silicon dioxide, crospovidone, lactose monohydrate, magnesium stearate, microcrystalline cellulose, povidone, sodium citrate.

INDICATIONS AND CLINICAL USE

MELOXICAM (meloxicam) is indicated for the symptomatic treatment of:

- Rheumatoid arthritis in adults and
- Painful osteoarthritis (arthrosis, degenerative joint disease) in adults.

Throughout this document, the term NSAIDs refers to both non-selective NSAIDs and selective COX-2 inhibitor NSAIDs, unless otherwise indicated.

For patients with an increased risk of developing CV and/or GI adverse events, other management strategies that do NOT include the use of NSAIDs should be considered first. (See CONTRAINDICATIONS and WARNINGS AND PRECAUTIONS).

Use of MELOXICAM should be limited to the lowest effective dose for the shortest possible duration of treatment in order to minimize the potential risk for cardiovascular or gastrointestinal adverse events. (See CONTRAINDICATIONS and WARNINGS AND PRECAUTIONS).

MELOXICAM, as a NSAID, does NOT treat clinical disease or prevent its progression.

MELOXICAM, as a NSAID, only relieves symptoms and decreases inflammation for as long as the patient continues to take it.

Geriatrics (> 65 years of age):

Evidence from clinical studies and postmarket experience suggests that use in the geriatric population is associated with differences in safety. (See WARNINGS AND PRECAUTIONS – Special Populations – Geriatrics and DOSAGE AND ADMINISTRATION – Recommended Dose and Dosage Adjustment – Geriatrics (>65 years of age)).

Pediatrics (< 18 years of age):

Safety and efficacy have not been established in the pediatric population. (See CONTRAINDICATIONS).

CONTRAINDICATIONS

MELOXICAM is contraindicated in:

- the peri-operative setting of coronary artery bypass graft surgery (CABG). Although meloxicam has NOT been studied in this patient population, a selective COX-2 inhibitor NSAID studied in such a setting has led to an increased incidence of cardiovascular/thromboembolic events, deep surgical infections and sternal wound complications.
- pregnancy. The risks during the third trimester are premature closure of the ductus arteriosus and prolonged parturition. (See WARNINGS AND PRECAUTIONS – Special Populations – Pregnant Women)
- women who are breastfeeding because of the potential for serious adverse reactions in nursing infants. NSAIDs are known to pass into mother's milk.
- severe uncontrolled heart failure;
- known or suspected hypersensitivity to meloxicam or to any of the components/excipients
- history of acute asthmatic attacks or symptoms of asthma, urticaria, nasal polyps, anaphylaxis, rhinitis, angioedema or other allergic manifestations that are precipitated by ASA or other NSAIDs, since cross-sensitivity may exist. Fatal anaphylactoid reactions may occur in such individuals. Individuals with the above medical problem are at risk of a severe reaction even if they have taken NSAIDs in the past without any adverse reaction. (See WARNINGS AND PRECAUTIONS – Hypersensitivity Reactions – Anaphylactoid Reactions, ASA-Intolerance).
- active or recent gastro-intestinal/gastric/duodenal/peptic ulceration/perforation, active GI bleeding;
- cerebrovascular bleeding or other bleeding disorders;
- inflammatory bowel disease (Crohn's Disease or Ulcerative Colitis);
- severe liver impairment or active liver disease;
- severe renal impairment (creatinine clearance < 30 mL/min or 0.5 mL/sec) or deteriorating renal disease (individuals with lesser degrees of renal impairment are at risk of deterioration of their renal function when prescribed NSAIDs and must be monitored (See WARNINGS AND PRECAUTIONS – Renal);
- known hyperkalemia (see WARNINGS AND PRECAUTIONS – Renal - Fluid and Electrolyte Balance);
- children and adolescents aged less than 18 years.

In case of rare hereditary conditions that may be incompatible with an excipient of the product (please refer to WARNINGS AND PRECAUTIONS) the use of the product is contraindicated.

WARNINGS AND PRECAUTIONS

Risk of Cardiovascular (CV) Adverse Events: Ischemic Heart Disease, Cerebrovascular Disease, Congestive Heart Failure (NYHA II-IV) (see WARNINGS AND PRECAUTIONS – Cardiovascular)

MELOXICAM is a non-steroidal anti-inflammatory drug (NSAID). Use of some NSAIDs is associated with an increased incidence of cardiovascular adverse events (such as myocardial infarction, stroke or thrombotic events) which can be fatal. The risk may increase with duration of use. Patients with cardiovascular disease or risk factors for cardiovascular disease may be at greater risk.

Caution should be exercised in prescribing MELOXICAM to any patient with ischemic heart disease (including but NOT limited to acute myocardial infarction, history of myocardial infarction and/or angina), cerebrovascular disease (including but NOT limited to stroke, cerebrovascular accident, transient ischemic attacks and/or amaurosis fugax) and/or congestive heart failure (NYHA II-IV).

Use of NSAIDs, such as MELOXICAM, can promote sodium retention in a dose-dependant manner, through a renal mechanism, which can result in increased blood pressure and/or exacerbation of congestive heart failure. (See WARNINGS AND PRECAUTIONS – Renal – Fluid and Electrolyte Balance).

Randomized clinical trials with meloxicam have not been designed to detect differences in cardiovascular events in a chronic setting. Therefore, caution should be exercised when prescribing MELOXICAM.

Risk of Gastrointestinal (GI) Adverse Events (see WARNINGS AND PRECAUTIONS – Gastrointestinal)

Use of NSAIDs, such as MELOXICAM, is associated with an increased incidence of gastrointestinal adverse events (such as peptic/duodenal ulceration, perforation, obstruction and gastrointestinal bleeding).

General

For relevant drug interactions that require particular attention, See DRUG INTERACTIONS section.

Frail or debilitated patients may tolerate side effects less well and therefore special care should be taken in treating this population. **To minimize the potential risk for an adverse event, the lowest effective dose should be used for the shortest possible duration.** As with other NSAIDs, caution should be used in the treatment of elderly patients who are more likely to be suffering from impaired renal, hepatic or cardiac function. For high risk patients, alternate therapies that do not involve NSAIDs should be considered.

MELOXICAM is NOT recommended for use with other NSAIDs, with the exception of low-dose ASA for cardiovascular prophylaxis, because of the absence of any evidence demonstrating synergistic benefits and the potential for additive adverse reactions. (See DRUG INTERACTIONS – Drug/Drug Interactions – Acetylsalicylic Acid (ASA) or other NSAIDs)

MELOXICAM tablets contain lactose. Patients with rare hereditary problems of galactose intolerance, the Lapp-lactase deficiency or glucose-galactose malabsorption should not take this medicine.

Carcinogenesis and Mutagenesis

See TOXICOLOGY section.

Cardiovascular

MELOXICAM is a non-steroidal anti-inflammatory drug (NSAID). Use of some NSAIDs is associated with an increased incidence of cardiovascular adverse events (such as myocardial infarction, stroke or thrombotic events) which can be fatal. The risk may increase with duration of use. Patients with cardiovascular disease or risk factors for cardiovascular disease may be at greater risk.

Caution should be exercised in prescribing MELOXICAM to patients with risk factors for cardiovascular disease, cerebrovascular disease or renal disease, such as any of the following (NOT an exhaustive list):

- **Hypertension**
- **Dyslipidemia/Hyperlipidemia**
- **Diabetes Mellitus**
- **Congestive Heart Failure (NYHA I)**
- **Coronary Artery Disease (Atherosclerosis)**
- **Peripheral Arterial Disease**
- **Smoking**
- **Creatinine Clearance (< 60 mL/min or 1 mL/sec)**

Use of NSAIDs, such as MELOXICAM, can lead to new hypertension or can worsen pre-existing hypertension, either of which may increase the risk of cardiovascular events as described above. Thus blood pressure should be monitored regularly. Consideration should be given to discontinuing MELOXICAM should hypertension either develop or worsen with its use.

Use of NSAIDs, such as MELOXICAM, can induce fluid retention and edema, and may exacerbate congestive heart failure, through a renally-mediated mechanism. (See WARNINGS AND PRECAUTIONS – Renal – Fluid and Electrolyte Balance).

For patients with a high risk of developing an adverse CV event, other management strategies that do NOT include the use of NSAIDs should be considered first. **To minimize the potential risk for an adverse CV event, the lowest effective dose should be used for the shortest possible duration.**

Endocrine and Metabolism

Corticosteroids: MELOXICAM is NOT a substitute for corticosteroids. It does NOT treat corticosteroid insufficiency. Abrupt discontinuation of corticosteroids may lead to exacerbation of corticosteroid responsive illness. Patients on prolonged corticosteroid therapy should have their therapy tapered slowly if a decision is made to discontinue corticosteroids. (See DRUG INTERACTIONS – Drug-Drug Interactions – Glucocorticoids).

Gastrointestinal (GI)

Serious GI toxicity (sometimes fatal), such as peptic/duodenal ulceration, inflammation, perforation, obstruction and gastrointestinal bleeding, can occur at any time, with or without warning symptoms in patients treated with NSAIDs, such as MELOXICAM. Minor upper GI problems, such as dyspepsia, commonly occur at any time. Health care providers should remain alert for ulceration and bleeding in patients treated with MELOXICAM, even in the absence of previous GI tract symptoms. Most spontaneous reports of fatal GI events are in elderly or debilitated patients and therefore special care should be taken in treating this population. **To minimize the potential risk for an adverse GI event, the lowest effective dose should be used for the shortest possible duration.** For high risk patients, alternate therapies that do not involve NSAIDs should be considered. (See WARNINGS AND PRECAUTIONS – Special Populations – Geriatrics).

Patients should be informed about the signs and/or symptoms of serious GI toxicity and instructed to discontinue using MELOXICAM and seek emergency medical attention if they experience any such symptoms. The utility of periodic laboratory monitoring has NOT been demonstrated, nor has it been adequately assessed. Most patients who develop a serious upper GI adverse event on NSAID therapy have no symptoms. Upper GI ulcers, gross bleeding or perforation, caused by NSAIDs, appear to occur in approximately 1% of patients treated for 3-6 months, and in about 2-4% of patients treated for one year. These trends continue, thus, increasing the likelihood of developing a serious GI event at some time during the course of therapy. Even short-term therapy has its risks.

Caution should be taken if prescribing MELOXICAM to patients with a prior history of peptic/duodenal ulcer disease or gastrointestinal bleeding as these individuals have a greater than 10-fold higher risk for developing a GI bleed when taking a NSAID than patients with neither of these risk factors. Other risk factors for GI ulceration and bleeding include the following: *Helicobacter pylori* infection, increased age, prolonged use of NSAID therapy, excess alcohol intake, smoking, poor general health status or concomitant therapy with any of the following:

- Anti-coagulants (e.g. warfarin)
- Anti-platelet agent (e.g. ASA, clopidogrel)
- Oral corticosteroids (e.g. prednisone)
- Selective Serotonin Reuptake Inhibitors (SSRIs) (e.g. citalopram, fluoxetine, paroxetine, sertraline)

Prospective, long-term studies required to compare the incidence of serious clinically significant upper gastrointestinal adverse events among patients taking meloxicam versus other NSAID products have not been performed.

There is no definitive evidence that the concomitant administration of histamine H₂ receptor antagonists and/or antacids will either prevent the occurrence of gastrointestinal adverse events or allow continuation of therapy when and if these adverse reactions appear.

Genitourinary

Some NSAIDs are associated with persistent urinary symptoms (bladder pain, dysuria, urinary frequency), hematuria or cystitis. The onset of these symptoms may occur at any time after the initiation of therapy with a NSAID.

Should urinary symptoms occur, in the absence of an alternate explanation, treatment with MELOXICAM should be stopped to ascertain if symptoms disappear. This should be done before any urological investigations or treatments are carried out.

Hematologic

NSAIDs inhibiting prostaglandin biosynthesis interfere with platelet function to varying degrees; patients who may be adversely affected by such an action, such as those on anti-coagulants or suffering from hemophilia or platelet disorders should be carefully observed when MELOXICAM is administered.

Anti-coagulants

Numerous studies have shown that the concomitant use of NSAIDs and anti-coagulants increases the risk of bleeding. Concurrent therapy of MELOXICAM with warfarin requires close monitoring of the international normalized ratio (INR).

Even with therapeutic INR monitoring, increased bleeding may occur. (See DRUG INTERACTIONS – Drug-Drug Interactions – Anticoagulants).

Anti-platelet Effects

NSAIDs inhibit platelet aggregation and have been shown to prolong bleeding time in some patients. Unlike acetylsalicylic acid (ASA) their effect on platelet function is quantitatively less, or of shorter duration, and reversible.

Meloxicam and other NSAIDs have no proven efficacy as anti-platelet agents and should NOT be used as a substitute for ASA or other anti-platelet agents for prophylaxis of cardiovascular thromboembolic diseases. Anti-platelet therapies (e.g. ASA) should NOT be discontinued. There is some evidence that the use of NSAIDs with ASA can markedly attenuate the cardioprotective effects of ASA. (See DRUG INTERACTIONS – Drug-Drug Interactions – Acetylsalicylic Acid or other NSAIDs).

Concomitant administration of meloxicam with low dose ASA increases the risk of GI ulceration and associated complications.

For information on interaction between low dose ASA and meloxicam and any other interaction, see DRUG INTERACTIONS – Acetylsalicylic Acid (ASA) or Other NSAIDs.

Blood dyscrasias

Blood dyscrasias (such as neutropenia, leukopenia, thrombocytopenia, aplastic anemia, and agranulocytosis) associated with the use of NSAIDs are rare, but can occur with severe consequences.

Anemia is sometimes seen in patients receiving NSAIDs, including meloxicam. This may be due to fluid retention, GI blood loss, or an incompletely described effect upon erythropoiesis. Patients on long-term treatment with NSAIDs, including meloxicam, should have their hemoglobin or hematocrit checked if they exhibit any signs or symptoms of anemia or blood loss.

The incidence of treatment-related anemia is more frequent than 1%. The incidence of disturbances of blood count, including differential white cell count, leukopenia and thrombocytopenia, is between 0.1 and 1%.

Concomitant administration of a potentially myelotoxic drug, in particular methotrexate, appears to be a predisposing factor to the onset of cytopenia.

Hepatic/Biliary/Pancreatic

As with other NSAIDs, borderline elevations of one or more liver enzyme tests (AST, ALT, alkaline phosphatase) may occur in up to 15% of patients. These laboratory abnormalities may progress, may remain essentially unchanged, or may be transient with continued therapy. Notable elevations of ALT or AST (approximately three or more times the upper limit of normal) have been reported in approximately 1% of patients in clinical trials with NSAIDs.

A patient with signs and/or symptoms suggesting liver dysfunction, or in whom an abnormal liver function test has occurred, should be evaluated for evidence of the development of a more severe hepatic reaction while on therapy with MELOXICAM. Severe hepatic reactions including jaundice and cases of fatal hepatitis, liver necrosis and hepatic failure, some of them with fatal outcomes, have been reported with NSAIDs.

Although such reactions are rare, if abnormal liver tests persist or worsen, if clinical signs and symptoms consistent with liver disease develop (e.g. jaundice), or if systemic manifestations occur (e.g., eosinophilia, associated with rash, etc.), MELOXICAM should be discontinued.

If there is a need to prescribe MELOXICAM in the presence of impaired liver function, it must be done under strict observation.

Hypersensitivity Reactions

Anaphylactoid Reactions

As with NSAIDs in general, anaphylactoid reactions have occurred in patients without known prior exposure to meloxicam. In post-marketing experience, rare cases of anaphylactic/anaphylactoid reactions and angioedema have been reported in patients receiving meloxicam. MELOXICAM should NOT be given to patients with the ASA-triad. This symptom complex typically occurs in asthmatic patients who experience rhinitis with or without nasal polyps, or who exhibit severe, potentially fatal bronchospasm after taking ASA or other NSAIDs (see CONTRAINDICATIONS). Emergency help should be sought in cases where anaphylactoid reaction occurs.

ASA-Intolerance

MELOXICAM should NOT be given to patients with complete or partial syndrome of ASA-intolerance (rhinosinusitis, urticaria/angioedema, nasal polyps, asthma) in whom asthma, anaphylaxis, urticaria/angioedema, rhinitis or other allergic manifestations are precipitated by ASA or other NSAIDs. Fatal anaphylactoid reactions have occurred in such individuals. As well, individuals with the above medical problems are at risk of severe fatal reaction even if they have taken NSAIDs in the past without any adverse reaction (See CONTRAINDICATIONS).

Cross-sensitivity

Patients sensitive to any one of the NSAIDs may be sensitive to any of the other NSAIDs as well.

Serious Skin Reactions

See WARNINGS AND PRECAUTIONS – Skin.

Immune

See WARNINGS AND PRECAUTIONS – Infection – Aseptic Meningitis.

Infection

MELOXICAM, in common with other NSAIDs, may mask signs and symptoms of an underlying infectious disease.

Aseptic Meningitis

Rarely, with some NSAIDs, the symptoms of aseptic meningitis (stiff neck, severe headaches, nausea and vomiting, fever or clouding of consciousness) have been observed. Patients with autoimmune disorders (systemic lupus erythematosus, mixed connective tissues diseases, etc.) seem to be pre-disposed. Therefore, in such patients, the health care provider must be vigilant to the development of this complication.

Neurologic

Some patients may experience drowsiness, dizziness, blurred vision, vertigo, tinnitus, hearing loss, insomnia or depression with the use of NSAIDs, such as meloxicam. If patients experience these side effects, they should exercise caution in carrying out activities that require alertness.

Ophthalmologic

Blurred and/or diminished vision has been reported with the use of NSAIDs. If such symptoms develop MELOXICAM should be discontinued and an ophthalmologic examination performed. Ophthalmologic examination should be carried out at periodic intervals in any patient receiving MELOXICAM for an extended period of time.

Peri-Operative Considerations

See CONTRAINDICATIONS – Coronary Artery Bypass Graft Surgery.

Psychiatric

See WARNINGS AND PRECAUTIONS – Neurologic.

Renal

Long-term administration of NSAIDs to animals has resulted in renal papillary necrosis and other abnormal renal pathology. In humans, there have been reports of acute interstitial nephritis, hematuria, low grade proteinuria, glomerulonephritis, renal medullary necrosis and occasionally nephrotic syndrome.

Renal insufficiency due to NSAID use is seen in patients with pre-renal conditions leading to reduction in renal blood flow or blood volume. Under these circumstances, renal prostaglandins help maintain renal perfusion and glomerular filtration rate (GFR). In these patients, administration of a NSAID may cause a reduction in prostaglandin synthesis leading to impaired renal function. Patients at greatest risk of this reaction are those with pre-existing renal insufficiency (GFR < 60 mL/min or 1 mL/s), dehydrated patients, patients on salt restricted diets, those with congestive heart failure, cirrhosis, liver dysfunction, taking angiotensin-converting enzyme inhibitors, angiotensin-II receptor blockers, cyclosporine, diuretics and those who are elderly. Serious or life-threatening renal failure has been reported in patients with normal or impaired renal function after short-term therapy with NSAIDs. Even patients at risk who demonstrate the ability to tolerate a NSAID under stable conditions may decompensate during periods of added stress (e.g. dehydration due to gastroenteritis). Discontinuation of NSAIDs is usually followed by recovery to the pre-treatment state.

The extent to which metabolites may accumulate in patients with renal failure has not been studied with meloxicam. As with other NSAIDs, metabolites of which are excreted by the kidney, patients with significantly impaired renal function should be more closely monitored.

Caution should be used when initiating treatment with NSAIDs, such as meloxicam, in patients with considerable dehydration. Such patients should be rehydrated prior to initiation of therapy. Caution is also recommended in patients with pre-existing kidney disease. No dose reduction is required in patients with mild or moderate renal impairment (i.e. in patients with a creatinine clearance of greater than 15 mL/min or 0.25 mL/s).

Advanced Renal Disease

See CONTRAINDICATIONS.

Fluid and Electrolyte Balance

Use of NSAIDs, such as MELOXICAM, can promote sodium retention in a dose-dependent manner, which can lead to fluid retention and edema, and consequences of increased blood

pressure and exacerbation of congestive heart failure. Thus, caution should be exercised in prescribing MELOXICAM in patients with a history of congestive heart failure, compromised cardiac function, hypertension, increased age or other conditions predisposing to fluid retention. For patients at risk, clinical monitoring is recommended. (See WARNINGS AND PRECAUTIONS – Cardiovascular).

Use of NSAIDs, such as MELOXICAM, can increase the risk of hyperkalemia, especially in patients with diabetes mellitus, renal failure, increased age, or those receiving concomitant therapy with adrenergic blockers, angiotensin-converting enzyme inhibitors, angiotensin-II receptor antagonists, cyclosporine, or some diuretics.

Electrolytes should be monitored periodically. (See CONTRAINDICATIONS).

Respiratory

ASA-induced asthma is an uncommon but very important indication of ASA and NSAID sensitivity. It occurs more frequently in patients with asthma who have nasal polyps.

Sexual Function/Reproduction

The use of meloxicam, as with any drug known to inhibit cyclooxygenase / prostaglandin synthesis, may impair fertility and is not recommended in women attempting to conceive. Therefore, in women who have difficulties conceiving, or who are undergoing investigation of infertility, withdrawal of meloxicam should be considered.

Skin

In rare cases, serious skin reactions such as Stevens-Johnson syndrome, toxic epidermal necrolysis, exfoliative dermatitis and erythema multiforme have been associated with the use of some NSAIDs. Because the rate of these reactions is low, they have usually been noted during post-marketing surveillance in patients taking other medications also associated with the potential development of these serious skin reactions. Thus, causality is NOT clear. These reactions are potentially life threatening but may be reversible if the causative agent is discontinued and appropriate treatment instituted. Patients appear to be at highest risk of these reactions early in the course of therapy, the onset of the reaction occurring in the majority of cases within the first month of treatment. MELOXICAM should be discontinued at the first appearance of skin rash, mucosal lesions, or any other sign of hypersensitivity. Patients should be advised that if they experience a skin rash they should discontinue their NSAID and contact their physician for assessment and advice, including which additional therapies to discontinue.

Special Populations

Pregnant Women: MELOXICAM is CONTRAINDICATED for use during pregnancy. The risks during the third trimester are premature closure of the ductus arteriosus and prolonged parturition.

Inhibition of prostaglandin-synthesis may adversely affect pregnancy and/or the embryo-foetal development. Data from epidemiological studies suggest an increased risk of miscarriage and of cardiac malformation and gastroschisis after use of a prostaglandin synthesis inhibitor in early pregnancy. The absolute risk for cardiovascular malformation was increased from less than 1%, up to approximately 1.5%. The risk is believed to increase with dose and duration of therapy.

In animals, administration of a prostaglandin synthesis inhibitor has been shown to result in increased pre- and post implantation loss and embryo-foetal lethality. In addition, increased incidences of various malformations, including cardiovascular, have been reported in animals given a prostaglandin synthesis inhibitor during the organogenetic period.

During the third trimester of pregnancy all prostaglandin-synthesis inhibitors may expose the foetus to:

- cardiopulmonary toxicity (with premature closure of the ductus arteriosus and pulmonary hypertension)
 - renal dysfunction, which may progress to renal failure with oligo-hydroamniosis;
- the mother and the neonate, at the end of pregnancy, to:
- possible prolongation of bleeding time, an anti-aggregating effect which may occur even at very low doses
 - inhibition of uterine contractions resulting in delayed or prolonged labour

See CONTRAINDICATIONS and TOXICOLOGY.

Nursing Women: MELOXICAM is CONTRAINDICATED in nursing women.

MELOXICAM is contraindicated for use in women who are breastfeeding because of the potential for serious adverse reactions in nursing infants. NSAIDs are known to pass into mother's milk.

Pediatrics (<18 years of age): See CONTRAINDICATIONS.

Safety and effectiveness of meloxicam in pediatric patients below the age of 18 years have not been evaluated.

Geriatrics (> 65 years of age): Patients older than 65 years (hereafter referred to as older or elderly) and frail or debilitated patients are more susceptible to a variety of adverse reactions from NSAIDs. The incidence of these adverse reactions increases with dose and duration of treatment. In addition, these patients are less tolerant to ulceration and bleeding. Most reports of fatal GI events are in this population, older patients are also at risk of a lower esophageal injury including ulceration and bleeding. For such patients, consideration should be given to a lower starting dose than the one usually recommended, with individual adjustment when necessary and under close supervision.

Monitoring and Laboratory Tests

The following monitoring criteria and laboratory tests are recommended for patients taking MELOXICAM. This is not an exhaustive list.

Laboratory Testing:

- Potassium (Renal function, Hyperkalemia)
- INR/effects of anticoagulants (Co-prescription of oral anticoagulants)
- Serum transaminases and other liver function tests (liver function)
- Renal function parameters such as serum creatinine and serum urea (in case of Methotrexate, Diuretics, Cyclosporine, ACE-Inhibitor or ARB co-prescription, and in susceptible patients re: the renal effects of meloxicam, e.g. impaired renal function or

- dehydration)
- Lithium plasma concentrations (in case of Lithium co-prescription)
- Blood cell count, including differential white cell count (in case of Methotrexate co-prescription)

Monitoring Activities:

- Patients with GI symptoms
- Patients with oral anticoagulation (see above)
- Blood pressure (in case of Antihypertensives co-prescription, and in susceptible patients with fluid retention)
- Periodic Ophthalmologic evaluation (in patients on extended treatment)

For more information, please refer to the WARNINGS AND PRECAUTIONS and DRUG INTERACTIONS sections.

ADVERSE REACTIONS

Adverse Drug Reaction Overview

The information to compile the following data is based on clinical trials involving 14,325 patients who have been treated with daily oral doses of 7.5 and 15 mg meloxicam tablets or capsules. In these clinical trials, the following indications were studied: osteoarthritis and rheumatoid arthritis (approved indications); ankylosing spondylitis, sciatica and low back pain (unapproved indications).

In the overall clinical trial database of 14325 patients, treatment exposure up to 6 weeks was obtained in 14313* patients, while exposure up to 3 months was in 2185 patients. Exposure up to 6 months was in 1642 patients, exposure up to one year was obtained in 1031 patients and 471 patients were exposed for more than one year to meloxicam.

* For 12 patients treated with Meloxicam, information is missing to categorize the duration of exposure.

Frequent Adverse Events

The following adverse events, which may be casually related to the administration of meloxicam, have a frequency of $\geq 1\%$.

Gastrointestinal:	dyspepsia, nausea, abdominal pain, diarrhoea
Skin and Appendages:	skin rash
Central Nervous System:	light-headedness; headache

Serious Adverse Drug Reactions:

The following serious adverse drug reactions have been reported in association with meloxicam use:

- Gastrointestinal ulceration, perforation or bleeding (see WARNINGS AND PRECAUTIONS, Gastrointestinal (GI) and DRUG INTERACTIONS, Drug-Drug

- Interactions – Selective Serotonin Reuptake Inhibitors (SSRIs);
- Asthma, bronchospasm (see WARNINGS AND PRECAUTIONS, Hypersensitivity Reactions – Anaphylactoid Reactions and WARNINGS AND PRECAUTIONS, Respiratory);
 - Hypersensitivity reactions including angioedema, skin rash, pruritus (see WARNINGS AND PRECAUTIONS, Hypersensitivity Reactions – Anaphylactoid Reactions and Skin);
 - Renal failure, hematuria (see WARNINGS AND PRECAUTIONS, Genitourinary and Renal, DRUG INTERACTIONS, Drug-Drug Interactions – Anti-hypertensives, Cyclosporine or Tacrolimus, Diuretics and Methotrexate);
 - Visual disturbances including blurred vision (see WARNINGS AND PRECAUTIONS, Neurologic and Ophthalmologic).
 - Vomiting or persistent dyspepsia, nausea, abdominal pain or diarrhea (see WARNINGS AND PRECAUTIONS, Gastrointestinal (GI) and Infection – Aseptic Meningitis);
 - Micturition disorders;
 - Oedema (see WARNINGS AND PRECAUTIONS, Cardiovascular and Renal – Fluid and Electrolyte Balance);
 - Jaundice (see WARNINGS AND PRECAUTIONS, Hepatic/Biliary/Pancreatic);
 - Malaise, fatigue;
 - Aseptic Meningitis (see WARNINGS AND PRECAUTIONS, Infections – Aseptic Meningitis);
 - Confusion, depression, lightheadedness (see WARNINGS AND PRECAUTIONS, Neurologic)
 - Tinnitus (see WARNINGS AND PRECAUTIONS, Neurologic).

Clinical Trial Adverse Drug Reactions

Because clinical trials are conducted under very specific conditions the adverse reaction rates observed in the clinical trials may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse drug reaction information from clinical trials is useful for identifying drug-related adverse events and for approximating rates.

Adverse drug reactions (ADRs) occurring with a frequency of $\geq 1\%$ in a 12 week, double-blind, randomized, placebo- and active-controlled clinical trial in osteoarthritis (Study 107.181) are presented in Table 1. Diclofenac was used as a comparator in a dose of 100 mg/day.

TABLE 1 ADVERSE DRUG REACTIONS IN A PLACEBO- AND ACTIVE-CONTROLLED TRIAL IN OSTEOARTHRITIS (107.181) WITH INCIDENCE \geq 1% IN ANY TREATMENT DISPLAYED ON PREFERRED TERM LEVEL

MedDRA system organ class MedDRA preferred term	Treatment at Onset											
	Placebo		Meloxicam 3.75 mg		Meloxicam 7.5 mg		Meloxicam 15 mg		Diclofenac 100 mg		Total	
	N	%	N	%	N	%	N	%	N	%	N	%
Summary Data	157	100.0	154	100.0	154	100.0	156	100.0	153	100.0	774	100.0
Total Treated												
Total with any Adverse Event	28	17.8	34	22.1	31	20.1	26	16.7	34	22.2	153	19.8
Ear and labyrinth disorders	1	0.6	1	0.6	1	0.6	1	0.6	2	1.3	6	0.8
Tinnitus	1	0.6	1	0.6	0	0.0	0	0.0	2	1.3	4	0.5
Gastrointestinal disorders	15	9.6	19	12.3	21	13.6	14	9.0	22	14.4	91	11.8
Abdominal pain	3	1.9	2	1.3	2	1.3	3	1.9	2	1.3	12	1.6
Constipation	3	1.9	1	0.6	2	1.3	0	0.0	4	2.6	10	1.3
Diarrhea	4	2.5	1	0.6	8	5.2	2	1.3	4	2.6	19	2.5
Dry mouth	0	0.0	1	0.6	0	0.0	0	0.0	2	1.3	3	0.4
Dyspepsia	5	3.2	6	3.9	5	3.2	6	3.8	6	3.9	28	3.6
Flatulence	5	3.2	5	3.2	4	2.6	3	1.9	4	2.6	21	2.7
Gastroesophageal reflux disease	0	0.0	0	0.0	1	0.6	3	1.9	2	1.3	6	0.8
Nausea	0	0.0	7	4.5	4	2.6	1	0.6	5	3.3	17	2.2
General disorders and administration site conditions	4	2.5	3	1.9	4	2.6	5	3.2	4	2.6	20	2.6
Gravitational oedema	1	0.6	1	0.6	0	0.0	2	1.3	1	0.7	5	0.6
Oedema peripheral	0	0.0	0	0.0	1	0.6	2	1.3	0	0.0	3	0.4
Metabolism and nutrition disorders	2	1.3	1	0.6	2	1.3	2	1.3	1	0.7	8	1.0
Increased appetite	1	0.6	1	0.6	0	0.0	2	1.3	0	0.0	4	0.5
Musculoskeletal and connective tissue disorders	2	1.3	4	2.6	1	0.6	2	1.3	1	0.7	10	1.3
Arthralgia	0	0.0	2	1.3	0	0.0	1	0.6	1	0.7	4	0.5
Nervous system disorders	9	5.7	9	5.8	5	3.2	5	3.2	5	3.3	33	4.3
Dizziness	3	1.9	3	1.9	1	0.6	2	1.3	1	0.7	10	1.3
Headache	6	3.8	4	2.6	1	0.6	1	0.6	3	2.0	15	1.9
Skin and subcutaneous tissue disorders	2	1.3	3	1.9	4	2.6	5	3.2	2	1.3	16	2.1
Hyperhidrosis	0	0.0	1	0.6	0	0.0	2	1.3	0	0.0	3	0.4
Rash	1	0.6	1	0.6	2	1.3	0	0.0	0	0.0	4	0.5
Vascular disorders	2	1.3	1	0.6	0	0.0	2	1.3	1	0.7	6	0.8
Hypertension	2	1.3	1	0.6	0	0.0	1	0.6	0	0.0	4	0.5

Adverse drug reactions (ADRs) occurring with a frequency of \geq 1% in a 12-week double-blind, randomized, placebo- and active-controlled trial in rheumatoid arthritis (Study 107.183) are presented in Table 2. Diclofenac was used as a comparator in a dose of 150 mg/day (n = 182).

TABLE 2 ADVERSE DRUG REACTIONS IN A PLACEBO- AND ACTIVE-CONTROLLED TRIAL IN RHEUMATOID ARTHRITIS (107.183) WITH INCIDENCE \geq 1% IN ANY TREATMENT DISPLAYED ON PREFERRED TERM LEVEL

MedDRA system organ class MedDRA preferred term	Treatment at Onset											
	Placebo		Meloxicam 7.5 mg		Meloxicam 15 mg		Meloxicam 22.5 mg		Diclofenac 150 mg		Total	
	N	%	N	%	N	%	N	%	N	%	N	%
Summary Data												
Total Treated	178	100.0	176	100.0	185	100.0	177	100.0	182	100.0	898	100.0
Total with any Adverse Event	34	19.1	38	21.6	38	20.5	31	17.5	40	22.0	181	20.2
Blood and lymphatic system disorders	1	0.6	0	0.0	2	1.1	0	0.0	0	0.0	3	0.3
Red blood cell abnormality	1	0.6	0	0.0	2	1.1	0	0.0	0	0.0	3	0.3
Ear and labyrinth disorders	0	0.0	0	0.0	2	1.1	1	0.6	0	0.0	3	0.3
Tinnitus	0	0.0	0	0.0	2	1.1	1	0.6	0	0.0	3	0.3
Eye disorders	2	1.1	0	0.0	1	0.5	1	0.6	1	0.5	5	0.6
Visual disturbance	2	1.1	0	0.0	0	0.0	1	0.6	0	0.0	3	0.3
Gastrointestinal disorders	21	11.8	27	15.3	25	13.5	22	12.4	29	15.9	124	13.8
Abdominal pain	2	1.1	8	4.5	5	2.7	2	1.1	6	3.3	23	2.6
Constipation	1	0.6	2	1.1	3	1.6	2	1.1	4	2.2	12	1.3
Diarrhea	8	4.5	6	3.4	9	4.9	2	1.1	6	3.3	31	3.5
Dyspepsia	6	3.4	9	5.1	5	2.7	5	2.8	6	3.3	31	3.5
Flatulence	2	1.1	3	1.7	6	3.2	7	4.0	6	3.3	24	2.7
Gastritis	0	0.0	0	0.0	0	0.0	1	0.6	3	1.6	4	0.4
Gastrointestinal hemorrhage	0	0.0	2	1.1	1	0.5	0	0.0	1	0.5	4	0.4
Melena	0	0.0	2	1.1	1	0.5	0	0.0	0	0.0	3	0.3
Mouth ulceration	0	0.0	0	0.0	0	0.0	4	2.3	0	0.0	4	0.4
Nausea	4	2.2	7	4.0	7	3.8	1	0.6	3	1.6	22	2.4
Vomiting	2	1.1	0	0.0	1	0.5	0	0.0	0	0.0	3	0.3
General disorders and administration site conditions	3	1.7	4	2.3	3	1.6	4	2.3	7	3.8	21	2.3
Fatigue	0	0.0	1	0.6	0	0.0	0	0.0	4	2.2	5	0.6
Gravitational edema	0	0.0	1	0.6	0	0.0	2	1.1	1	0.5	4	0.4

TABLE 2 (Continued): ADVERSE DRUG REACTIONS IN A PLACEBO- AND ACTIVE CONTROLLED TRIAL IN RHEUMATOID ARTHRITIS (107.183) WITH INCIDENCE \geq 1% IN ANY TREATMENT DISPLAYED ON PREFERRED TERM LEVEL

MedDRA system organ class MedDRA preferred term	Treatment at Onset											
	Placebo		Meloxicam 7.5 mg		Meloxicam 15 mg		Meloxicam 22.5 mg		Diclofenac 150 mg		Total	
	N	%	N	%	N	%	N	%	N	%	N	%
Hepatobiliary disorders	0	0.0	0	0.0	0	0.0	1	0.6	2	1.1	3	0.3
Hepatic function abnormal	0	0.0	0	0.0	0	0.0	1	0.6	2	1.1	3	0.3
Investigations	1	0.6	0	0.0	2	1.1	3	1.7	2	1.1	8	0.9
Blood urea increased	0	0.0	0	0.0	1	0.5	1	0.6	2	1.1	4	0.4
Creatinine renal clearance decreased	0	0.0	0	0.0	1	0.5	2	1.1	0	0.0	3	0.3
Nervous system disorders	8	4.5	8	4.5	4	2.2	2	1.1	7	3.8	29	3.2
Dizziness	4	2.2	4	2.3	1	0.5	0	0.0	1	0.5	10	1.1
Headache	6	3.4	2	1.1	3	1.6	0	0.0	4	2.2	15	1.7
Somnolence	0	0.0	0	0.0	0	0.0	1	0.6	2	1.1	3	0.3
Psychiatric disorders	0	0.0	2	1.1	1	0.5	0	0.0	0	0.0	3	0.3
Insomnia	0	0.0	2	1.1	1	0.5	0	0.0	0	0.0	3	0.3
Skin and subcutaneous tissue disorders	2	1.1	8	4.5	5	2.7	2	1.1	3	1.6	20	2.2
Hyperhidrosis	0	0.0	2	1.1	0	0.0	0	0.0	0	0.0	2	0.2
Rash	1	0.6	2	1.1	5	2.7	0	0.0	1	0.5	9	1.0
Rash erythematous	0	0.0	2	1.1	0	0.0	1	0.6	0	0.0	3	0.3
Vascular disorders	1	0.6	0	0.0	1	0.5	1	0.6	2	1.1	5	0.6
Hypertension	1	0.6	0	0.0	1	0.5	1	0.6	2	1.1	5	0.6

Less Common Clinical Trial Adverse Drug Reactions (<1%)

The following is a list of adverse drug reactions regardless of causality occurring in <1% of patients receiving 7.5 or 15 mg meloxicam in clinical trials involving approximately 14,325 patients. In these clinical trials, the following indications were studied: osteoarthritis and rheumatoid arthritis (approved indications); ankylosing spondylitis, sciatica and low back pain (unapproved indications).

Body as a Whole: allergic reaction, face edema, fatigue, fever, hot flushes, malaise, syncope, weight decrease, weight increase.

Cardiovascular: angina pectoris, cardiac failure, hypertension (increase of blood pressure), hypotension, myocardial infarction, vasculitis, edema, flushes; **Heart Rate and Rhythm:** arrhythmia, palpitation, tachycardia.

Central and Peripheral Nervous System: convulsions, dizziness, paresthesia, tremor, vertigo, tinnitus, drowsiness.

Psychiatric Disorders: abnormal dreaming, anxiety, appetite increased, confusion, depression, nervousness, somnolence.

Gastrointestinal: colitis, dry mouth, duodenal ulcer, eructation, esophagitis, gastric ulcer, gastritis, gastroesophageal reflux, gastrointestinal hemorrhage (occult or macroscopic gastrointestinal bleeding), hematemesis, hemorrhagic duodenal ulcer, hemorrhagic gastric ulcer, gastro-intestinal perforation, melena, pancreatitis, perforated duodenal ulcer, perforated gastric ulcer, stomatitis ulcerative, vomiting, constipation, flatulence, gastroduodenal ulcer.

Gastro-intestinal bleeding, ulceration or perforation may potentially be fatal (See WARNINGS AND PRECAUTIONS – Gastrointestinal).

Hematologic: disturbances of blood count, including differential white cell count, leukopenia, purpura, thrombocytopenia and anemia.

Liver and Biliary System: hepatitis, liver function test abnormal (e.g. raised transaminases or bilirubin).

Metabolic and Nutritional: dehydration.

Respiratory: asthma, bronchospasm, dyspnea.

Skin and Appendages: alopecia, angioedema, bullous eruption, dermatitis bullous, photosensitivity reaction (photosensitisation), pruritus, sweating increased, stomatitis, urticaria.

Special Senses: abnormal vision (including blurred vision), conjunctivitis, taste perversion, tinnitus.

Urinary System: albuminuria, abnormal renal function parameters (increased serum creatinine and/or serum urea), hematuria, acute renal failure.

Abnormal Hematologic and Clinical Chemistry Findings

Few patients in clinical trials in osteoarthritis (study 107.181) and rheumatoid arthritis (study 107.183) experienced abnormal hematologic or clinical chemistry findings with potential clinical significance. There were a few instances of decreased red blood cells in both meloxicam-treated (1.1%) and placebo-treated patients (0.7%). Increased red blood cells were experienced in placebo-treated patients (0.7%). Increased serum potassium was experienced in both meloxicam-treated patients (7.5 mg-0.7%, 15 mg-1.7%) and placebo-treated patients (1.3%). Increased blood urea nitrogen and increased serum creatinine was experienced in meloxicam-treated patients (1.3% and 2.0% respectively).

Post-Market Adverse Drug Reactions

Additional reports of serious adverse events temporally associated with meloxicam during worldwide post-marketing experience are included below. Because these events are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or clearly establish a casual relationship to meloxicam exposure.

Gastro-intestinal: hepatitis, gastritis;

Hematologic: agranulocytosis.

Liver and Biliary System: jaundice, liver failure.

Dermatological: bullous reactions, erythema multiforme, photosensitivity reaction (photosensitization), Stevens Johnson Syndrome, toxic epidermal necrolysis.

Respiratory: Onset of asthma attacks in individual allergic to aspirin or other NSAIDs.

Central nervous system: confusion and disorientation, alteration of mood.

Genitourinary: acute renal failure, interstitial nephritis, micturition disorders, acute urinary retention.

Vision disorders: conjunctivitis, visual disturbances including blurred vision.

Hypersensitivity reactions: angio-oedema and immediate hypersensitivity reactions, including anaphylactoid / anaphylactic reactions including shock.

DRUG INTERACTIONS

Overview

Cytochrome P450 Interactions:

Meloxicam is eliminated almost entirely by hepatic metabolism, of which approximately two thirds are mediated by cytochrome (CYP) P450 enzymes (CYP 2C9 major pathway and CYP 3A4 minor pathway) and one-third by other pathways, such as peroxidase oxidation. The potential for a pharmacokinetic interaction should be taken into account when meloxicam and drugs known to inhibit, or to be metabolised by, CYP 2C9 and/or CYP 3A4 are administered concurrently.

Drug-Drug Interactions

Acetylsalicylic acid (ASA) or other NSAIDs:

Concomitant administration of aspirin (1000 mg TID) to healthy volunteers tended to increase the AUC (10%) and C_{max} (24%) of meloxicam. The clinical significance of this interaction is not known.

The use of meloxicam in addition to any other NSAIDs, including over-the-counter ones (such as ASA and ibuprofen) for analgesic and/or anti-inflammatory effect is NOT recommended because of the absence of any evidence demonstrating synergistic benefits and the potential for additive adverse reactions (e.g. increased risk of gastro-intestinal ulcers and bleeding).

The exception is the use of low dose ASA for cardiovascular protection, when another NSAID is being used for its analgesic/anti-inflammatory effect, keeping in mind that combination NSAID therapy is associated with additive adverse reactions. Concomitant administration of low-dose aspirin with meloxicam may result in an increased rate of GI ulceration or other complications,

compared to use of meloxicam alone. MELOXICAM is not a substitute for aspirin for cardiovascular prophylaxis.

Some NSAIDs (e.g. ibuprofen) may interfere with the anti-platelet effects of low dose ASA, possibly by competing with ASA for access to the active site of cyclooxygenase-1.

Antacids:

No pharmacokinetic interaction was detected with concomitant administration of antacids. MELOXICAM tablets can be administered without regard to timing of antacids. (See ACTION AND CLINICAL PHARMACOLOGY -Pharmacokinetics)

Anti-coagulants:

Anticoagulant activity should be monitored, particularly in the first few days after initiating or changing MELOXICAM therapy in patients receiving warfarin or similar agents, since these patients are at an increased risk of bleeding. The effect of meloxicam on the anticoagulant effect of warfarin was studied in a group of healthy subjects receiving daily doses of warfarin that produced an INR (International Normalized Ratio) between 1.2 and 1.8. In these subjects, meloxicam did not alter warfarin pharmacokinetics and the average anticoagulant effect of warfarin as determined by prothrombin time. However, one subject showed an increase in INR from 1.5 to 2.1. Caution should be used when administering MELOXICAM with warfarin since patients on warfarin may experience changes in INR and an increased risk of bleeding complications when a new medication is introduced. (See WARNINGS AND PRECAUTIONS – Hematologic – Anti-coagulants).

Anti-Hypertensives:

NSAIDs may diminish the anti-hypertensive effect of Angiotensin Converting Enzyme (ACE) inhibitors.

Combinations of ACE inhibitors, angiotensin-II antagonists, diuretics and NSAIDs might have an increased risk for acute renal failure and hyperkalemia. Blood pressure and renal function (including electrolytes) should be monitored more closely in this situation, as occasionally there can be a substantial increase in blood pressure.

NSAIDs and ACE inhibitors or angiotensin-II receptor antagonists exert a synergistic effect on the decrease of glomerular filtration. In patients with pre-existing renal impairment this may lead to acute renal failure.

Anti-Platelet Agents (including ASA):

There is an increased risk of bleeding, via inhibition of platelet function, when anti-platelet agents, oral anticoagulants, systemically administered heparin and thrombolytics are combined with NSAIDs, such as MELOXICAM. If such co-prescribing cannot be avoided, close monitoring of the effects of anti-coagulants is required. (See WARNINGS AND PRECAUTIONS - Hematologic section).

Cholestyramine:

Pretreatment for four days with cholestyramine significantly increased the clearance of meloxicam by 50%. This resulted in a decrease in $t_{1/2}$, from 19.2 hours to 12.5 hours, and a 35%

reduction in AUC. This suggests the existence of a recirculation pathway for meloxicam in the gastrointestinal tract. The clinical relevance of this interaction has not been established.

Cimetidine:

Concomitant administration of 200 mg cimetidine QID did not alter the single-dose pharmacokinetics of 30 mg meloxicam.

Cyclosporine or Tacrolimus:

Nephrotoxicity of cyclosporine or tacrolimus may be enhanced by NSAIDs via renal prostaglandin mediated effects. During combined treatment with either of these drugs, renal function should be monitored.

Digoxin:

Meloxicam 15 mg once daily for 7 days did not alter the plasma concentration profile of digoxin after b-acetyldigoxin administration for 7 days at clinical doses. *In vitro* testing found no protein binding drug interaction between digoxin and meloxicam.

Diuretics:

Clinical studies, as well as post-marketing observations, have shown that NSAIDs can reduce the natriuretic effect of furosemide and thiazide diuretics in some patients. This effect has been attributed to inhibition of renal prostaglandin synthesis. Studies with furosemide agents and meloxicam have not demonstrated a reduction in natriuretic effect. Furosemide single and multiple dose pharmacodynamics and pharmacokinetics are not affected by multiple doses of meloxicam. Nevertheless, during concomitant therapy with furosemide and meloxicam, patients should be observed closely for signs of declining renal function (see WARNINGS AND PRECAUTIONS – Renal Function), as well as to assure diuretic efficacy.

Glucocorticoids:

Some studies have shown that the concomitant use of NSAIDs and oral glucocorticoids increases the risk of GI adverse events such as ulceration and bleeding via a synergistic effect. This is especially the case in older (>65 years of age) individuals.

Lithium:

Monitoring of plasma lithium concentrations is advised when stopping or starting a NSAID, as increased lithium concentrations can occur.

In clinical trials, NSAIDs have produced a reduction in renal lithium clearance and an elevation of plasma lithium levels, which may reach toxic values. The concomitant use of lithium and NSAIDs is not recommended. In a study conducted in healthy subjects, mean pre-dose lithium concentration and AUC were increased by 21% in subjects receiving lithium doses ranging from 804 to 1072 mg BID with meloxicam 15 mg QD as compared to subjects receiving lithium alone. These effects have been attributed to inhibition of renal prostaglandin synthesis by meloxicam. If this combination appears necessary, lithium plasma concentrations should be monitored carefully during the initiation, adjustment and withdrawal of meloxicam treatment.

Methotrexate:

A study in 13 rheumatoid arthritis (RA) patients evaluated the effects of multiple doses of meloxicam on the pharmacokinetics of methotrexate taken once weekly. Meloxicam did not

have a significant effect on the pharmacokinetics of single doses of methotrexate. *In vitro*, methotrexate did not displace meloxicam from its human serum binding sites.

In case combination treatment with methotrexate and NSAIDs is necessary, blood cell count and the renal function should be monitored. Caution should be taken in case both NSAID and methotrexate are given within 3 days, in which case the plasma level of methotrexate may increase and cause increased toxicity. Although the pharmacokinetics of methotrexate (15 mg/week) were not relevantly affected by concomitant meloxicam treatment, it should be considered that the haematological toxicity of methotrexate can be amplified by treatment with NSAID drugs.

Concomitant administration of NSAIDs with a potentially myelotoxic drug, such as methotrexate, appears to be a predisposing factor to the onset of cytopenia.

NSAIDs can reduce the tubular secretion of methotrexate thereby increasing the plasma concentrations of methotrexate. For this reason, for patients on high dosages of methotrexate (more than 15 mg/week) the concomitant use of NSAIDs is not recommended. The risk of an interaction between NSAID preparations and methotrexate, should be considered also in patients on low dosage of methotrexate, especially in patients with impaired renal function.

Oral Contraceptives:

No drug interaction information is available for meloxicam co-administered with oral contraceptives. A decrease of the efficacy of intrauterine devices by NSAIDs has been previously reported but needs further confirmation.

Oral Hypoglycemics:

An interaction with oral hypoglycemic agents has been noted with some NSAIDs, however no interaction data is available for the co-administration of these agents with meloxicam.

Selective Serotonin Reuptake Inhibitors (SSRIs):

Concomitant administration of NSAIDs and SSRIs may increase the risk of gastrointestinal ulceration and bleeding. (See WARNINGS AND PRECAUTIONS – Gastrointestinal).

Other Drug Interactions:

A population kinetics study with meloxicam indicated a lack of relevant interaction of sulfasalazine, gold compounds and glucocorticoids on the pharmacokinetics of meloxicam. No drug interaction data is available for meloxicam and the co-administration of the following products: phenytoin, acetaminophen, alcohol, aminoglycosides, butemide, colchicine, cyclosporin, indapamide, insulin, nephrotoxic agents, NSAIDs (other than ASA), oral contraceptives, potassium supplements, probenecid, valproic acid, zidovudine.

Drug-Food Interactions

Interactions with food have not been established.

Drug-Herb Interactions

Interactions with herbal products have not been established.

Drug-Laboratory Interactions

Interactions with laboratory tests have not been established.

Drug-Lifestyle Interactions

There are no specific studies about effects on the ability to drive vehicles and to use machinery. Patients who experience visual disturbances, drowsiness or other central nervous system disturbances should refrain from these activities.

DOSAGE AND ADMINISTRATION

Dosing Considerations

In patients with increased risks of adverse reactions, treatment should be started at the dose of 7.5 mg once daily.

The maximum recommended daily dose of MELOXICAM (meloxicam) tablets is 15 mg.

Recommended Dose and Dosage Adjustment

Use of MELOXICAM is restricted to adults 18 years of age and older and should be limited to the lowest effective dose for the shortest possible duration of treatment. (See CONTRAINDICATIONS and WARNINGS AND PRECAUTIONS).

Osteoarthritis: 7.5 mg once daily. If necessary, the dose may be increased to 15 mg once daily.

Rheumatoid arthritis: 15 mg once daily. According to the therapeutic response, the dose may be reduced to 7.5 mg once daily.

MELOXICAM may be taken without regard to timing of meals.

Hepatic Impairment: No dose adjustment is necessary in patients with mild to moderate hepatic insufficiency.

No dose reduction is required in patients with clinically stable liver cirrhosis.

MELOXICAM is contraindicated in patients with severe liver impairment or active liver disease.

See CONTRAINDICATIONS, WARNINGS AND PRECAUTIONS – Hepatic/Biliary/Pancreatic and ACTION AND CLINICAL PHARMACOLOGY – Special Populations and Conditions – Hepatic Insufficiency.

Renal Impairment: No dose reduction is required in patients with mild to moderate renal impairment (i.e., in patients with creatinine clearance of greater than 15 mL/min or 0.25 mL/s).

In dialysis patients with severe renal failure, the dose should not exceed 7.5 mg/day.

MELOXICAM is contraindicated in patients with severe renal impairment (creatinine clearance < 30 mL/min or 0.5 mL/s) or deteriorating renal disease.

See CONTRAINDICATIONS, WARNINGS AND PRECAUTIONS – Renal and ACTION AND CLINICAL PHARMACOLOGY – Special Populations and Conditions – Renal Insufficiency.

Geriatrics (> 65 years of age): For elderly, frail or debilitated patients, considerations should be given to a starting dose lower than the one usually recommended, with individual adjustment when necessary and under close supervision.

See CONTRAINDICATIONS, WARNINGS AND PRECAUTIONS – General and Special Populations – Geriatrics and ACTION AND CLINICAL PHARMACOLOGY – Special Populations and Conditions – Geriatrics.

Missed Dose

If a dose is missed, the usual schedule must be resumed the following day. An extra dose must not be taken.

OVERDOSAGE

There is limited experience with meloxicam overdose. Four cases have taken 6 to 11 times the highest recommended dose; all recovered. Cholestyramine is known to accelerate the clearance of meloxicam.

Symptoms following acute NSAID overdose are usually limited to lethargy, drowsiness, nausea, vomiting, and epigastric pain, which are generally reversible with supportive care.

Gastrointestinal bleeding can occur. Severe poisoning may result in hypertension, acute renal failure, hepatic dysfunction, respiratory depression, coma, convulsions, cardiovascular collapse, and cardiac arrest. Anaphylactoid reactions have been reported with therapeutic ingestion of NSAIDs, and may occur following an overdose.

Patients should be managed with symptomatic and supportive care following an NSAID overdose. In cases of acute overdose, gastric lavage followed by activated charcoal is recommended. Gastric lavage performed more than one hour after overdose has little benefit in the treatment of overdose. Administration of activated charcoal is recommended for patients who present 1-2 hours after overdose. For substantial overdose or severely symptomatic patients, activated charcoal may be administered repeatedly. Accelerated removal of meloxicam by 4 gm oral doses of cholestyramine given three times a day was demonstrated in a clinical trial. Administration of cholestyramine may be useful following an overdose. Forced diuresis, alkalinization of urine, hemodialysis, or hemoperfusion may not be useful due to high protein binding.

For management of a suspected drug overdose, contact your regional Poison Control Centre.

ACTION AND CLINICAL PHARMACOLOGY

Mechanism of Action

Meloxicam is a nonsteroidal anti-inflammatory drug (NSAID) that exhibits anti-inflammatory, analgesic, and antipyretic properties in animals. Meloxicam showed potent anti-inflammatory activity in all standard models of inflammation. NSAIDs are believed to exert their

pharmacologic effects primarily through inhibition of the enzyme cyclooxygenase (COX). In turn, inhibition of this enzyme leads to an inhibition of biosynthesis of prostaglandins and other autacoids, substances which are potent biological mediators involved in diverse physiologic functions as well as pathologic conditions.

To date, two isozymes of COX have been identified and characterized, namely, COX-1 and COX-2 which have different intrinsic properties, expression controls and localization. COX-1, the constitutive form, has been described as a constitutive enzyme occurring in many tissues including the gastrointestinal tract, kidney, lungs, brain and platelets. COX-1 is found in blood vessels, platelets, stomach and kidney. In contrast, COX-2, the inducible form, is mostly an inducible enzyme, limited in distribution and expressed in high levels in inflamed tissues. COX-2 is thought to be involved in inflammatory responses. Recent studies have shown that differential inhibition of these two isozymes is associated with a different biological profile. Meloxicam has shown a selective inhibition of COX-2 in several *in vitro* test systems, as demonstrated by a greater dose dependent inhibition of COX-2 over COX-1 at levels similar to those seen in plasma at therapeutic steady state concentrations. The prostaglandins produced by the cyclooxygenases are not the only factors involved in the protection of the gastric mucosa.

A human pharmacology study compared the effects of meloxicam 7.5 mg once daily and indomethacin 25 mg three times daily on platelet aggregation and platelet thromboxane formation, which are exclusively COX-1 dependent, and renal prostaglandin (PGE₂) excretion. Platelet aggregation and thromboxane formation were almost completely inhibited by indomethacin but remained unaffected by meloxicam. Meloxicam showed no significant effects on urinary PGE₂ excretion whereas indomethacin reduced urinary PGE₂ excretion by 43%.

In another study, meloxicam (7.5 and 15 mg) demonstrated a greater inhibition of COX-2 *ex vivo*, as demonstrated by a greater inhibition of lipopolysaccharide-stimulated PGE₂ production (COX-2) as compared with serum thromboxane production (COX-1).

Meloxicam has been shown to inhibit COX-2 in several *in vitro* and *ex vivo* test systems. The inhibition of thromboxane in platelets, and consequently platelet aggregation, occurs via inhibition of COX-1. Meloxicam inhibition of thromboxane in platelets (via COX-1) is dose dependent and incomplete at anti-inflammatory doses. No significant inhibition of platelet aggregation has been observed with meloxicam at the recommended therapeutic doses of 7.5 and 15 mg once daily.

Inhibition of COX-2 also inhibits the production of systemic prostacyclin. Inhibition of prostacyclin may have a pro-thrombotic effect.

Prospective, controlled, long-term (>3 months) studies required to establish the clinical significance of these results have not been performed.

Pharmacodynamics

See ACTION AND CLINICAL PHARMACOLOGY – Mechanism of Action.

Pharmacokinetics

Absorption: The absolute bioavailability of meloxicam capsules was 89% following a single oral dose of 30 mg compared with 30 mg IV bolus injection. Meloxicam capsules have been shown to be bioequivalent to meloxicam tablets. Following single intravenous doses, dose-proportional pharmacokinetics were shown in the range of 5 mg to 60 mg. After multiple oral doses the pharmacokinetics of meloxicam capsules were dose-proportional over the range of 7.5 mg to 15 mg. Mean C_{max} was achieved within four to five hours after a 7.5 mg meloxicam tablet was taken under fasted conditions, indicating a prolonged drug absorption. The rate or extent of absorption was not affected by multiple dose administration, suggesting linear pharmacokinetics. With multiple dosing, steady state conditions were reached by day 5. A second meloxicam concentration peak occurs around 12 to 14 hours post-dose suggesting gastrointestinal recirculation.

Once daily dosing leads to drug plasma concentrations with a relatively small peak-trough fluctuation in the range of 0.4 - 1.0 $\mu\text{g/mL}$ for 7.5 mg doses and 0.8 - 2.0 $\mu\text{g/mL}$ for 15 mg doses, respectively (C_{min} and C_{max} at steady state, respectively).

Continuous treatment for longer periods (e.g. six months,) did not point to any changes in pharmacokinetics compared to steady state pharmacokinetics after two weeks of oral treatment with 15 mg meloxicam/day. Any differences after treatment longer than six months are thus rather unlikely.

Table 3 SINGLE DOSE AND STEADY STATE PHARMACOKINETIC PARAMETERS FOR ORAL 15 MG MELOXICAM (MEAN AND % CV)¹

Pharmacokinetic Parameters (% CV)		Steady State						Single Dose			
		Healthy male adults (Fed) ²		Elderly males (Fed) ²		Elderly females (Fed) ²		Renal Failure (Fasted)		Hepatic insufficiency (Fasted)	
		15 mg ³		15 mg		15 mg		15 mg		15 mg	
N		24		5		8		12		12	
C_{max}	[$\mu\text{g/mL}$]	1.9	-25	2.3	-59	3.2	-24	0.59	-36	0.84	-29
t_{max}	[h]	6.5	-37	5	-12	6	-27	4	-65	10	-87
$t_{1/2}$	[h]	15	-45	21	-34	24	-34	18	-46	16	-29
CL/f	[mL/min]	8.3	-32	9.9	-76	5.1	-22	19	-43	11	-44
Vz/f^4	[L]	10	-36	15	-42	10	-30	26	-44	14	-29

1) The parameter values in the Table are from various studies; 2) not under high fat conditions; 3) meloxicam tablets; 4) $Vz/f = \text{Dose}/(\text{AUC} \cdot \text{Kel})$

Food and Antacid Effects: Drug intake after a high fat breakfast (75 g of fat) did not affect extent of absorption of meloxicam capsules, but led to 22% higher C_{max} values. Mean C_{max} values were achieved between five and six hours. No pharmacokinetic interaction was detected

with concomitant administration of antacids. Meloxicam tablets can be administered without regard to timing of meals and antacids.

Distribution: The mean volume of distribution (V_{ss}) of meloxicam is approximately 10 L. Meloxicam is ~ 99.4% bound to human plasma proteins (primarily albumin) within the therapeutic dose range. The fraction of protein binding is independent of drug concentration, over the clinically relevant concentration range, but decreases to ~99% in patients with renal disease. Meloxicam penetration into human red blood cells, after oral dosing, is less than 10%. Following a radiolabeled dose, over 90% of the radioactivity detected in the plasma was present as unchanged meloxicam.

Meloxicam concentrations in synovial fluid, after a single oral dose, range from 40% to 50% of those in plasma. The free fraction in synovial fluid is 2.5 times higher than in plasma, due to the lower albumin content in synovial fluid as compared to plasma. The significance of this penetration is unknown.

Metabolism: Meloxicam is almost completely metabolized to four pharmacologically inactive metabolites. The major metabolite, 5'-carboxy meloxicam (60% of dose), from P-450 mediated metabolism was formed by oxidation of an intermediate metabolite 5'-hydroxymethyl meloxicam which is also excreted to a lesser extent (9% of dose). *In vitro* studies indicate that cytochrome P-450 2C9 plays an important role in this metabolic pathway with a minor contribution of the CYP 3A4 isozyme. Patients' peroxidase activity is probably responsible for the other two metabolites which account for 16% and 4% of the administered dose, respectively.

Excretion: Meloxicam excretion is predominantly in the form of metabolites, and occurs to equal extents in the urine and feces. Only traces of the unchanged parent compound are excreted in the urine (0.2%) and feces (1.6%). The extent of the urinary excretion was confirmed for unlabeled multiple 7.5 mg doses: 0.5%, 6% and 13% of the dose were found in urine in the form of meloxicam, and the 5'-hydroxymethyl and 5'-carboxy metabolites, respectively. There is significant biliary and/or enteral secretion of the drug. This was demonstrated when oral administration of cholestyramine following a single IV dose of meloxicam decreased the AUC of meloxicam by 50%.

The mean elimination half-life ($t_{1/2}$) ranges from 15 hours to 20 hours. The elimination half-life is constant across dose levels indicating linear metabolism within the therapeutic dose range. Plasma clearance ranges from 7 to 9 mL/min.

Special Populations and Conditions

Pediatrics: In a study of 36 children, kinetic measurements were made in 18 children at doses of 0.25 mg/kg BW. Maximum plasma concentration C_{max} (-34%) as well as $AUC_{0-\infty}$ (-28%) tended to be lower in the younger age group (aged 2 to 6 years, $n = 7$) as compared to the older age group (7 to 14 years, $n = 11$) while weight normalized clearance appeared to be higher in the younger age group. A historical comparison with adults revealed that plasma concentrations were at least similar for older children and adults. Plasma elimination half-lives (13 h) were similar for both groups and tended to be shorter than in adults (15-20 h).

Geriatrics: Elderly males (≥ 65 years of age) exhibited meloxicam plasma concentrations and steady state pharmacokinetics similar to young males. Elderly females (≥ 65 years of age) had a 47% higher AUC_{ss} and 32% higher C_{max} ss as compared to younger females (<55 years of age) after body weight normalization. Despite the increased total concentrations in the elderly females, the adverse event profile was comparable for both elderly patient populations. A smaller free fraction was found in elderly female patients in comparison to elderly male patients.

Mean plasma clearance at steady state in elderly subjects was slightly lower than that reported for younger subjects.

Gender: Young females exhibited slightly lower plasma concentrations relative to young males. After single doses of 7.5 mg meloxicam, the mean elimination half-life was 19.5 hours for the female group as compared to 23.4 hours for the male group. At steady state, the data were similar (17.9 hours vs. 21.4 hours). This pharmacokinetic difference due to gender is likely to be of little clinical importance. There was linearity of pharmacokinetics and no appreciable difference in the C_{max} or T_{max} across genders.

Race: Pharmacokinetic data in Japanese subjects suggest a lower clearance of meloxicam in comparison to Caucasian subjects, but is not considered to require dose-adjustment due to the high intra-individual variability observed.

Hepatic Insufficiency: Following a single 15 mg dose of meloxicam there was no marked difference in plasma concentrations in subjects with mild (Child-Pugh Class I) and moderate (Child-Pugh Class II) hepatic impairment compared to healthy volunteers. Protein binding of meloxicam was not affected by hepatic insufficiency. No dose adjustment is necessary in mild to moderate hepatic insufficiency. Patients with severe hepatic impairment (Child-Pugh Class III) have not been adequately studied.

Renal Insufficiency: Meloxicam pharmacokinetics have been investigated in subjects with different degrees of renal insufficiency. Total drug plasma concentrations decreased with the degree of renal impairment while free AUC values were similar. Total clearance of meloxicam increased in these patients probably due to the increase in free fraction leading to an increased metabolic clearance. There is no need for dose adjustment in patients with mild to moderate renal failure (CrCL >15 mL/min or >0.25 mL/s). Patients with severe renal insufficiency have not been adequately studied. The use of MELOXICAM in subjects with severe renal impairment is not recommended (see WARNINGS AND PRECAUTIONS, Advanced Renal Disease).

In terminal renal failure, the increase in the volume of distribution may result in higher free meloxicam concentrations, and a daily dose of 7.5 mg must not be exceeded.

Hemodialysis: Following a single dose of meloxicam, the free C_{max} plasma concentrations were higher in patients with renal failure on chronic hemodialysis (1% free fraction) in comparison to healthy volunteers (0.3% free fraction). Hemodialysis did not lower the total drug concentration in plasma; therefore, additional doses are not necessary after hemodialysis. Meloxicam is not dialyzable.

STORAGE AND STABILITY

Store at controlled room temperature (15- 30°C), safely out of the reach of children. Store in a dry place.

DOSAGE FORMS, COMPOSITION AND PACKAGING

Tablets, 7.5 mg and 15 mg

7.5 mg: Light yellow, round, bevelled-edged, biconvex tablet with “ > ” on one side and “ML” on the other, contains 7.5 mg meloxicam;
7.5

15 mg: Light yellow, round tablet, convex and bevelled-edged with “ > ” on one side and “ML” on the other, contains 15 mg meloxicam;
15

Each of the above strengths is available in sealed clear PVC/PE/PVDC blisters (3 x 10 tablets) in aluminum foil in a carton of 30 and HDPE bottles of 100's and 500's.

Each tablet contains 7.5 mg or 15 mg meloxicam

Non-medicinal Ingredients: colloidal silicon dioxide (Cab-O-Sil M5), crospovidone (polyplasdone XL), lactose monohydrate (Impalpable #312), magnesium stearate, microcrystalline cellulose (Vivapur type 102), povidone (plasdone), sodium citrate (dihydrate).

PART II: SCIENTIFIC INFORMATION

PHARMACEUTICAL INFORMATION

Drug Substance


Proper name: Meloxicam BP

Chemical name: 4-hydroxy-2-methyl-*N*-(5-methyl-1,3-thiazol-2-yl)-2*H*-1,2-benzothiazine-3-carboxamide-1,1-dioxide

Molecular formula: C₁₄H₁₃N₃O₄S₂

Molecular mass: 351.4

Structural formula:



Physicochemical properties: Meloxicam is a light yellow to yellow coloured crystalline powder. It is soluble in chloroform, acetic acid, practically insoluble in water. Meloxicam has pK_a value of 4.08.

CLINICAL TRIALS

A randomized, two-way single dose crossover comparative bioavailability study of MELOXICAM Tablets 7.5 mg and Mobicox[®] 7.5 mg tablets has been performed in healthy male volunteers in the fasting state. A summary of the bioavailability data is tabulated below.

Table – 4 A Summary Table of Comparative Bioavailability Data, Fasted Study (1 x 7.5 mg)

Meloxicam (1 x 7.5 mg) From measured data				
Geometric Mean Arithmetic Mean (CV %)				
Parameter	Meloxicam 7.5 mg Tablets	MobicoxTM* 7.5 mg	% Ratio of Geometric LS Means	90% Confidence Interval
AUC ₀₋₇₂ (ng•hr/mL)	17235.9 17789.1 (27.8)	16927.0 17334.8 (23.6)	101.82	98.40 – 105.37
AUC _∞ (ng•hr/mL)	19306.7 20495.3 (41.5)	18617.2 19325.1 (30.6)	103.70	99.10 – 108.52
C _{MAX} (ng/mL)	709.3 717.1 (15.3)	669.9 679.3 (16.9)	105.87	99.40 – 112.76
T _{MAX} ** (h)	4.50 (2.00 – 6.00)	4.50 (3.00 – 5.00)		
T _½ *** (h)	20.75 (46.4)	19.07 (29.0)		

* The reference product, MobicoxTM (meloxicam) 15 mg Tablets (Boehringer Ingelheim (Canada) Ltd.) was purchased in Canada.

** The T_{max} parameter is expressed as median (range) only.

*** The T_½ parameter is expressed as arithmetic mean (% CV) only.

A randomized, two-way single dose crossover comparative bioavailability study of MELOXICAM Tablets 15 mg and Mobicox[®] 15 mg tablets has been performed in healthy male volunteers in the fasting state. A summary of the bioavailability data is tabulated below.

Table – 5 Summary Table of the Comparative Bioavailability Data, Fasted Study (1 x 15 mg)

Meloxicam (1 x 15 mg) From measured data				
Geometric Mean Arithmetic Mean (CV %)				
Parameter	Meloxicam 7.5 mg Tablets	Mobicox™* 7.5 mg	% Ratio of Geometric LS Means	90% Confidence Interval
AUC ₀₋₇₂ (ng•hr/mL)	33736.1 34928.0 (25.1)	31163.5 32478.6 (27.3)	108.26	104.04 – 112.64
AUC _∞ (ng•hr/mL)	36528.3 38050.2 (27.6)	34531.7 36583.9 (33.7)	105.78	101.33 – 110.42
C _{MAX} (ng/mL)	1559.0 1597.5 (22.3)	1249.7 1294.1 (22.8)	124.75	116.07 – 134.09
T _{MAX} ** (h)	4.50 (2.00 – 12.00)	5.00 (3.00 – 12.00)		
T _½ *** (h)	17.66 (20.4)	19.61 (31.5)		

* The reference product, Mobicox™ (meloxicam) 15 mg Tablets (Boehringer Ingelheim (Canada) Ltd.) was purchased in Canada.

** The Tmax parameter is expressed as median (range) only.

*** The T½ parameter is expressed as arithmetic mean (% CV) only.

Randomized clinical trials with meloxicam have NOT been designed to detect differences in cardiovascular adverse events in a chronic setting.

Prospective, long-term studies required to compare the incidence of serious, clinically significant upper gastrointestinal adverse events among patients taking meloxicam versus other NSAID products have not been performed.

OSTEOARTHRITIS

Study 107.181

The use of meloxicam for the treatment of the signs and symptoms of osteoarthritis of the knee and hip was evaluated in a double-blind controlled trial involving a total of 774 patients randomized and treated with meloxicam (N=464), placebo (N=157) or diclofenac (N=153) for 12 weeks. Meloxicam (3.75 mg, 7.5 mg and 15 mg daily) was compared to placebo and diclofenac (100 mg) (refer to Table 6). The four primary endpoints were investigator's global assessment, patient global assessment, patient pain assessment, and total WOMAC score (a self-administered questionnaire addressing pain, function and stiffness) (refer to Table 7). Patients on meloxicam 7.5 mg daily and meloxicam 15 mg daily showed significant improvement in each of these endpoints compared with placebo (refer to Table 7).

RHEUMATOID ARTHRITIS

Study 107.183

A 12-week double-blind placebo- and active-controlled comparison to investigate doses of meloxicam 7.5 mg, 15 mg and 22.5 mg in the treatment of rheumatoid arthritis was also performed to investigate the full dose range of meloxicam in one trial (refer to Table 6). In this study, a total of 894 patients were randomized and treated with placebo (N=177), meloxicam (7.5 mg, 15 mg, 22 mg) (N=536) or diclofenac (2 x 75 mg) (N=181) (refer to Table 6). Diclofenac 2 x 75 mg was included as active control to assess trial sensitivity. Meloxicam 7.5 mg and 22.5 mg were statistically superior to placebo in all primary endpoints, whereas 15 mg was statistically superior in three out of five primary endpoints (refer to Table 7). Diclofenac was superior to placebo in four of the five primary endpoints. All active treatments were significantly superior to placebo in secondary endpoints such as withdrawal due to lack of efficacy, patient's and investigator's final global assessment of efficacy, the patient's assessment of status with regard to a change in the arthritic condition and after adjustment for baseline also for the modified health assessment questionnaire. Assessment of efficacy after 4 weeks revealed significant differences between the higher doses of meloxicam and placebo but not between 7.5 mg meloxicam and placebo thus indicating that 7.5 mg may be a valuable dose for the treatment of RA but that acute flares might require a higher starting dose.

Study demographic and trial design

Table 6 Summary of patient demographics for clinical trials in Osteoarthritis and Rheumatoid Arthritis

Study #	Trial design	Dosage, route of administration, duration and comparator	Study subjects randomized and treated (n=number)	Mean age (Range)	Gender
Osteoarthritis Clinical Trial					
107.181	Multicenter, double-blind, double-dummy, randomized, parallel-group	3.75 mg/day, 7.5 mg/day and 15 mg/day, oral, 12 weeks, diclofenac, 100 mg, placebo	774	62 – 64 years	506 females, 268 males
Rheumatoid Arthritis Clinical Trial					
107.183	Multicenter, double-blind, double-dummy, randomized, parallel-group	7.5 mg/day, 15 mg/day and 22.5 mg/day, oral, 12 weeks, diclofenac, 150 mg, placebo	894	54.74 years for diclofenac group and 56.2 years for meloxicam 22.5 mg group.	681 females, 213 males

Study results

Table 7 Results of Osteoarthritis and Rheumatoid Arthritis Studies

Study #	Primary Endpoint(s)	Efficacy results
Osteoarthritis Clinical Trial		
107.181	<ul style="list-style-type: none">Investigator's global assessment of disease activityPatient's global assessment of disease activityPatient's overall assessment of painWOMAC (Western Ontario and McMaster University Osteoarthritis) index	Meloxicam in doses of 15 mg and 7.5 mg was statistically significantly more effective than placebo
Rheumatoid Arthritis Clinical Trial		
107.183	<ul style="list-style-type: none">Number of painful or tender joints out of 28 core jointsNumber of swollen joints out of 28 core jointsPatient's global assessment of disease activityInvestigator's global assessment of disease activityPatient's assessment of pain	At study end, meloxicam 7.5 mg and meloxicam 22.5 mg were significantly better than placebo for all five primary efficacy endpoints. Meloxicam 15 mg was significantly better than placebo for the patient's and investigator's global assessments and patient pain.

DETAILED PHARMACOLOGY

See ACTION AND CLINICAL PHARMACOLOGY section.

TOXICOLOGY

Oral LD₅₀ values ranged from about 98 mg/kg in female rats up to >800 mg/kg in minipigs. Intravenous values ranged from about 52 mg/kg in rats to 100 - 200 mg/kg in minipigs. Main signs of toxicity included reduced motor activity, anemia, and cyanosis. Most deaths occurred as a consequence of gastric ulcers and subsequent perforative peritonitis.

Repeated dose toxicity studies in rats and minipigs showed characteristic changes reported with other NSAIDs e.g. gastrointestinal ulceration and erosions and in the long term studies renal papillary necrosis. Gastrointestinal side effects were observed at oral doses of 1 mg/kg and higher in rats and of 3 mg/kg and above in minipigs. After intravenous administration doses of 0.4 mg/kg in rats and 9 mg/kg in minipigs caused gastrointestinal lesions. Renal papillary necrosis occurred only in rats at doses of 0.6 mg/kg or higher after lifetime exposure to meloxicam.

Oral reproductive studies in the rat have shown a decrease of ovulations and inhibition of implantations and embryotoxic effects (increase of resorptions) at maternotoxic dose levels at 1 mg/kg and higher. The affected dose levels exceeded the clinical dose (7.5-15 mg) by a factor of 6.6 to 3.3-fold on a mg/kg dose basis (50 kg person). Fetotoxic effects at the end of gestation, shared by all prostaglandin synthesis inhibitors, have been described.

Studies of toxicity on reproduction in rats and rabbits did not reveal teratogenicity up to oral doses of 4 mg/kg in rats and 80 mg/kg in rabbits. Doses of 2.5 mg/kg in rats and 20 mg/kg and higher in rabbits were embryotoxic. Prolongation of gestation and labour and an increased incidence of stillbirths, which is a well-known phenomenon of prostaglandin inhibition, occurred in the peri- and postnatal study at doses of 0.125 mg/kg and above.

Meloxicam was not mutagenic in the Ames test, the host-mediated assay and a mammalian gene mutation assay (V79/HPRT), nor clastogenic in a chromosome aberration assay in human lymphocytes and an in vivo micronucleus test in mouse bone marrow.

Carcinogenicity studies in rats and mice did not show any carcinogenic potential up to a dose level of 0.8 mg/kg in rats and 8 mg/kg in mice.

In the above mentioned life-time studies in rats and mice meloxicam did not damage articular cartilage, it was considered to be chondroneutral in these species.

Meloxicam did not induce immunogenic reactions in tests on mice and guinea pigs. In several tests, meloxicam proved to be less phototoxic than some older NSAIDs but similar in this respect to both piroxicam and tenoxicam.

In local tolerance studies meloxicam was well tolerated by all tested routes of administration: intravenous, intramuscular, rectal, dermal, and ocular administration.

REFERENCES

- 1) Bigby M. Nonsteroidal anti-inflammatory drugs. Uses in dermatology and cutaneous toxicity. In: Nonsteroidal anti-inflammatory drugs. Mechanisms and clinical uses. 2nd Edition. Eds. Lewis AJ, Furst DE. New York: Marcel Dekker Inc., 207-215:207-215.
- 2) Bird HA, Bevis PJR, Lapham G: An open study to assess the safety and tolerability of meloxicam 15 mg in subjects with rheumatoid disease and mild renal impairment. Data on file, Boehringer Ingelheim.
- 3) Bush U, Heinzl G, Narjes H: The effect of cholestiramine on the pharmacokinetics of meloxicam, a new non-steroidal anti-inflammatory drug (NSAID) in man. *Eur J Pharmacol* 1995; 48: 269-72.
- 4) Degner FL, Heinzl G, Narjes H et al.: The effect of meloxicam on the pharmacokinetics of beta-acetyl-digoxin. *Br J Clin Pharmacol* 1995; 40: 486-8.
- 5) Distel M, Muller H, Bluhmki E et al.: Global analysis of safety of a new NSAID: meloxicam. Data on file, Boehringer Ingelheim.
- 6) Engelhardt G, Homma D, Schlegel K et al.: Anti-inflammatory, analgesic, antipyretic and related properties of meloxicam, a new non-steroidal anti-inflammatory agent with favourable gastrointestinal tolerance. *Inflamm Res* 1995; 44: 423-33.
- 7) Engelhardt G, Homma D, Schnitzler C: Meloxicam: a potent inhibitor of adjuvant arthritis in the rat. *Inflamm Res* 1995; 44: 548-55.5
- 8) Engelhardt G, Bogel R, Schnitzler C et al.: Meloxicam: Influence on Arachidonic Acid Metabolism - Part I. *IN VITRO FINDINGS*. *Biochem Pharmacol* 1996; 51:21-8.
- 9) Ericson A. Nonsteroidal anti-inflammatory drugs in early pregnancy. *Reprod Toxicol* 2001;15:371.
- 10) European Commission Enterprise Directorate-General. Volume 3B Guidelines: Medicinal Products for Human Use: Safety, Environment and Information – Excipients in the Labelling and Package Leaflet of Medicinal Products for Human Use. July 2003. CPMP/463/00
- 11) Hosie J, Distel M, Bluhmki E: Meloxicam in osteoarthritis: a six-months, double-blind comparison with diclofenac sodium. Data on file, Boehringer Ingelheim
- 12) Huskisson EC, Ghozlan R, Kurthen R et al.: A long term study to evaluate the safety and efficacy of meloxicam therapy in patients with rheumatoid arthritis. Data on file, Boehringer Ingelheim.
- 13) Linden B, Distel M, Bluhmki E: A double-blind study to compare the efficacy and safety of meloxicam 15 mg with piroxicam 20 mg in patients with osteoarthritis of the hip or knee. Data on file, Boehringer Ingelheim.

- 14) Muller FO, Schall R, de Vaal AC et al.: Influence of meloxicam on furosemide pharmacokinetics and pharmacodynamics in healthy volunteers. *Eur J Pharmacol* 1995; 48: 247-51.
- 15) Noble S, Balfour JA: Meloxicam -New Drug Profile. *Drugs* 1996; 51(3):424-430.
- 16) Patoia L, Santucci L, Furno P et al.: A 4-week, double-blind, parallel-group study to compare the gastrointestinal effects of meloxicam 7.5 mg, meloxicam 15 mg, piroxicam 20 mg and placebo by means of faecal blood loss, endoscopy and symptoms evaluation in healthy volunteers. Data on file, Boehringer Ingelheim.
- 17) Schmid J, Busch U, Heinzl G et al.: Meloxicam: pharmacokinetics and metabolic pattern after intravenous infusion and oral administration to healthy subjects. *Drug Metab Dispos* 1995; 23(11): 1206-13.
- 18) Takeuchi K, Smale S, Premchand P, Maiden L, Sherwood R, Thjodleifsson B, Bjornsson E, Bjarnason I. Department of Internal Medicine, Guy's, King's, St. Thomas' Medical School, London, United Kingdom. Prevalence and mechanisms of nonsteroidal anti-inflammatory drug-induced clinical relapse in patients with inflammatory bowel disease. *Clin Gastroenterol Hepatol*. 2006 Feb;4(2):196-202.
- 19) Tata LJ, Fortun PJ, Hubbard RB, Smeeth L, Hawkey CJ, Smith CJP, et al. Does concurrent prescription of selective serotonin reuptake inhibitors and non-steroidal anti-inflammatory drugs substantially increase the risk of upper gastrointestinal bleeding? *Aliment Pharmacol Ther* 2005;22(3):175-181.
- 20) Turck D, Busch U, Heinzl G et al.: Clinical pharmacokinetics of meloxicam. *Eur J Rheumatol Inflamm* 1996; 15: 23-30.
- 21) Turck D, Busch U, Heinzl G et al.: Effect of food on the pharmacokinetics of meloxicam after oral administration. *Clin Drug Invest* 1995; 9: 270-6.
- 22) Vane JR, Botting RM: New insights into the mode of action of anti-inflammatory drugs. *Inflamm Res* 1995; 44: 1-10.
- 23) Verhamme KMC, Dieleman JP, Wijk MAM van, Lei J van der, Bosch JLHR, Stricker BHC, et al. Nonsteroidal anti-inflammatory drugs and increased risk of acute urinary retention. *Arch Intern Med* 2005;165(13):1547-1551.
- 24) Warner TD et al.: Nonsteroid drug selectivities for cyclo-oxygenase-1 rather than cyclo-oxygenase-2 are associated with human gastrointestinal toxicity: A full *in vitro* analysis. *Proc. Natl. Acad. Sci.* 1999; 96: 7563-7568.
- 25) Wojtulewski JA, Schattenkirchner M, Barcelo P et al.: A six-months double-blind trial to compare the efficacy and safety of meloxicam 7.5 mg daily and naproxen 750 mg daily in patients with rheumatoid arthritis. Data on file, Boehringer Ingelheim.

- 26) Product Monograph for ^{Pr}MOBICOX™ (meloxicam), 7.5 & 15.0 mg Tablets,
Manufactured by Boehringer Ingelheim (Canada) Ltd., Control# 120640, Date of
Revision: May 28, 2008.

PART III: CONSUMER INFORMATION

Pr **MELOXICAM**
Meloxicam Tablets BP

This leaflet is part III of a three-part "Product Monograph" published when MELOXICAM was approved for sale in Canada and is designed specifically for Consumers. This leaflet is a summary and will not tell you everything about MELOXICAM. Contact your doctor or pharmacist if you have any questions about the drug.

ABOUT THIS MEDICATION**What the medication is used for:**

MELOXICAM (meloxicam) belongs to a class of drugs known as nonsteroidal anti-inflammatory drugs (NSAIDs).

Your health care provider has prescribed MELOXICAM for you for symptomatic relief of one or more of the following medical conditions:

- Adult rheumatoid arthritis;
- Painful osteoarthritis (arthrosis, degenerative joint disease) in adults.

NSAID medicines may increase the chance of heart attack or stroke that can lead to death. This chance increases:

- With longer use of NSAID medicines
- In people who have heart disease or cardiovascular disease
- In people with risk factors for heart disease or cardiovascular disease

NSAID medicines can cause ulcers and bleeding in the stomach and intestines at any time during treatment. Ulcers and bleedings:

- Can happen without warning symptoms
- May cause death

The chance of a person getting an ulcer or bleeding increases with:

- Taking medicines called "corticosteroids" and "anticoagulants"
- Longer use
- Smoking
- Drinking alcohol
- Older age
- Having poor health

NSAID medicines should only be used:

- Exactly as prescribed
- At the lowest dose possible for your treatment
- For the shortest time needed

What it does:

MELOXICAM (meloxicam), as a nonsteroidal anti-inflammatory drug (NSAID), can reduce the chemicals produced by your body which can cause pain and swelling.

MELOXICAM, as a nonsteroidal anti-inflammatory drug (NSAID), does NOT cure your illness or prevent it from getting worse. MELOXICAM can only relieve pain and reduce swelling as long as you continue to take it.

MELOXICAM helps to relieve joint swelling, redness and pain of arthritis. Your body produces chemicals called prostaglandins. Some of these prostaglandins help line the stomach with a protective layer. In arthritis, other prostaglandins cause pain and swelling. At the dose prescribed by your doctor, MELOXICAM reduces the type that causes pain and swelling.

When it should not be used:

DO NOT TAKE MELOXICAM if you have, or previously had any of the following medical conditions:

- **Heart bypass surgery (planning to have or recently had);**
- **Severe, uncontrolled heart failure;**
- **Congestive heart failure;**
- **Bleeding in the brain or other bleeding disorders;**
- **Current Pregnancy;**
- **Currently breastfeeding (or planning to breastfeed);**
- **Allergy to meloxicam or any other component of MELOXICAM;**
- **Allergy to ASA (Acetylsalicylic Acid) or other NSAIDs (Nonsteroidal Anti-Inflammatory Drugs);**
- **Ulcer (active);**
- **Bleeding from the stomach, or gut (active);**
- **Inflammatory Bowel Disease (Crohn's Disease or Ulcerative Colitis);**
- **Liver Disease (active or severe);**
- **Kidney Disease (severe or worsening);**
- **High potassium in the blood.**
- **Rare hereditary conditions that may be incompatible with the non-medicinal ingredient, lactose.**

Do not take ASA (acetylsalicylic acid), ASA-containing compounds or other drugs used to relieve symptoms of arthritis while taking MELOXICAM unless directed to do so by your physician.

Patients who took a drug in the same class as MELOXICAM after a type of heart surgery (coronary artery bypass grafting (CABG)) were more likely to have heart attacks, strokes, blood clots in the leg(s) or lung(s), and infections or other complications than those who did NOT take that drug.

MELOXICAM should NOT be used in patients under 18 years of age since safety and effectiveness have NOT been established.

What the medicinal ingredient is:

Meloxicam BP

What the important nonmedicinal ingredients are:

Colloidal Silicon Dioxide, Crospovidone, Lactose Monohydrate, Magnesium Stearate, Microcrystalline Cellulose, Povidone, and Sodium Citrate.

What dosage forms it comes in:

7.5 mg and 15 mg tablets.

WARNINGS AND PRECAUTIONS

If you have, or previously had, any of the following medical conditions, see your health care provider to discuss treatment options other than MELOXICAM:

- Heart Attack or angina;
- Stroke or mini-stroke
- Loss of vision
- Current Pregnancy
- Congestive Heart Failure

Before taking this medication, tell your health care provider if you:

- or a family member are allergic to or have had a reaction to MELOXICAM or other anti-inflammatory drugs (such as acetylsalicylic acid (ASA), celecoxib, diclofenac, diflunisal, etodolac, fenoprofen, flurbiprofen, ibuprofen, indomethacin, ketoprofen, ketorolac, mefenamic acid, meloxicam, nabumetone, naproxen, oxaprozin, piroxicam, rofecoxib, sulindac, tenoxicam, tiaprofenic acid, tolmetin, or valdecoxib (NOT a complete list)). Allergic reactions may appear as increased sinus symptoms, hives (rash), new or worsened asthma or anaphylaxis (sudden collapse);
- or a family member has had asthma, nasal polyps, long-term swelling of the sinus (chronic sinusitis) or chronic urticaria (hives);
- have a history of stomach upset, ulcers or bleeding from the stomach or gut;
- have liver disease;
- had bleeding in the brain;
- have kidney disease or urine problems;
- have bleeding problems;
- have high cholesterol;
- have high blood pressure;
- have atherosclerosis;
- have poor circulation to your extremities (hands and feet);
- have diabetes mellitus;
- are a smoker or ex-smoker;
- are on any special diet, such as a low-sodium or low-sugar diet;
- have been told by a physician that you have intolerance to some sugars;
- have any other medical problem(s) such as alcohol

abuse, etc.

Also before taking this medication, tell your health care provider if you are planning to get pregnant.

While taking this medication:

- Tell any other physician, dentist, pharmacist or other health care professional that you see, that you are taking this medication, especially if you are planning to have heart surgery;
- Do NOT drink alcoholic beverages while taking this medication because you would be more likely to develop stomach problems;
- Fertility may be decreased. The use of MELOXICAM is not recommended in women trying to get pregnant. In women who have difficulty conceiving, stopping MELOXICAM should be considered;
- Your regular medical checkups are essential

INTERACTIONS WITH THIS MEDICATION

Talk to your health care provider and pharmacist if you are taking any other medication (prescription or non-prescription) such as any of the following (NOT a complete list):

- Acetylsalicylic Acid (ASA) or Other NSAIDs
 - e.g. ASA, celecoxib, diclofenac, ibuprofen, indomethacin, ketorolac, naproxen
- Anti-depressants
 - Selective Serotonin Reuptake Inhibitors (SSRIs)
 - e.g. citalopram, fluoxetine, paroxetine, sertraline
- Blood pressure medications
 - Angiotensin Converting Enzyme (ACE) Inhibitors
 - e.g. enalapril, lisinopril, perindopril, ramipril
 - Angiotensin II receptor blockers (ARBs)
 - e.g. candesartan, irbesartan, losartan, valsartan
- Blood thinners
 - e.g. warfarin, ASA, clopidogrel
- Cholestyramine
- Cimetidine
- Corticosteroids (including Glucocorticoids)
 - e.g. prednisone
- Cyclosporine
- Digoxin
- Diuretics
 - e.g. furosemide, hydrochlorothiazide
- Intrauterine Devices
- Lithium
- Methotrexate
- Oral contraceptives
- Oral hypoglycemics (diabetes medications)
- Tacrolimus

Your health care provider may prescribe low dose ASA

(acetylsalicylic acid) as blood thinner to reduce your risk of having a heart attack or stroke while you are taking MELOXCIAM. Take only the amount of ASA prescribed by your health care provider. You are more likely to upset or damage your stomach than if you take both MELOXCIAM and ASA than if you took MELOXCIAM alone.

PROPER USE OF THIS MEDICATION

DOSE:

MELOXCIAM is intended for use in patients greater than 18 years of age for the shortest possible duration.

Medical Condition	Starting Dose	Maximum Dose (per day)
Symptomatic treatment of rheumatoid arthritis in adults	15 mg once daily. According to therapeutic response, the dose may be reduced to 7.5 mg once daily	15 mg once daily. In dialysis patients with severe renal failure, the dose should not exceed 7.5 mg/day
Painful osteoarthritis (arthrosis, degenerative joint disease) in adults	7.5 mg once daily. If necessary, the dose may be increased to 15 mg once daily. In patients with increased risks of adverse reactions, treatment should be started at the dose of 7.5 mg once daily.	15 mg once daily. In dialysis patients with severe renal failure, the dose should not exceed 7.5 mg/day.

Take MELOXCIAM only as directed by your health care provider. **Do NOT take more of it, do NOT take it more often and do NOT take it for a longer period of time than your health care provider recommended. If possible, you should take the lowest dose of this medication for the shortest time period.** Taking too much MELOXCIAM increases your chances of unwanted and sometimes dangerous side effects, especially if you are elderly, have other diseases or take other medications.

If you will be using MELOXCIAM for more than 7 days, see your health care provider regularly to discuss whether this medicine is working for you and if it is causing you any unwanted effects. In some types of arthritis, up to two weeks may pass before you feel the full effects of this medicine.

This medication has been prescribed specifically for you. Do NOT give it to anyone else. It may harm them, even if their symptoms seem to be similar to yours.

MELOXCIAM is NOT recommended for use in patients under 18 years of age since safety and effectiveness have NOT been established.

MELOXCIAM tablets may be taken with or without food.

Missed Dose:

You should take MELOXCIAM as your doctor has prescribed. However, if you miss a dose, just resume your usual schedule the following day. Do not take an extra dose.

Overdose:

For management of a suspected drug overdose, contact your regional Poison Control Center.

SIDE EFFECTS AND WHAT TO DO ABOUT THEM

Along with its beneficial effects, MELOXCIAM, like other NSAID drugs, may cause some undesirable reactions especially when used for a long time or in large doses. Elderly, frail or debilitated patients often seem to experience more frequent or more severe side effects. Although not all of these side effects are common, when they do occur, they may require medical attention. Report all symptoms or side effects to your health care provider.

MELOXCIAM may cause you to become drowsy or tired. Be careful about driving or participating in activities that require you to be alert. If you become drowsy, dizzy or light-headed after taking MELOXCIAM, do NOT drive or operate machinery.

MELOXCIAM may cause you to become more sensitive to sunlight. Any exposure to sunlight or sunlamps may cause sunburn, skin blisters, skin rash, redness, itching or discoloration, or vision changes. If you have a reaction from the sun, check with your health care provider.

Check with your health care provider if you are not getting any relief of your arthritis or if any problems develop.

Check with your health care provider IMMEDIATELY if you develop chills, fever, muscle aches or pains, or other flu-like symptoms, especially if they occur before or together with a skin rash. These symptoms may be the first signs of a SERIOUS ALLERGIC REACTION to this medication.

SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM

Symptom / effect	STOP taking MELOXICAM and get emergency medical attention IMMEDIATELY	STOP taking MELOXICAM and talk to your physician or pharmacist
Bloody or black tarry stool;	✓	
Shortness of breath, wheezing, any trouble in breathing or tightness in the chest	✓	
Allergic reactions, such as: skin rash, hives or swelling, itching, chills, fever, muscle aches and pains, or other flu-like symptoms	✓	
Blurred vision, or any visual disturbance	✓	
Any change in the amount of colour of your urine (red or brown)	✓	
Any pain or difficulty experienced while urinating		✓
Swelling of the feet, lower legs, weight gain		✓
Vomiting or persistent indigestion, nausea, stomach pain or diarrhea		✓
Yellow discoloration of the skin or eyes, with or without itchy skin		✓
Malaise, fatigue, loss of appetite		✓
Headaches, stiff neck		✓
Mental confusion, depression		✓
Dizziness, lightheadedness		✓
Hearing problems		✓

This is NOT a complete list of side effects. For any unexpected effects while taking MELOXICAM, contact your health care provider or pharmacist immediately, so that these effects may be properly addressed.

HOW TO STORE IT

Store at room temperature (15-30°C). Store in a dry place.

Do NOT keep outdated medicine or medicine no longer needed. Any outdated or unused medicine should be returned to your pharmacist.

Keep out of reach of children.

This medication has been prescribed for your medical problem. Do not give it to anyone else.

If you require more information on this drug, consult your doctor or pharmacist.

REPORTING SUSPECTED SIDE EFFECTS

You can report any suspected adverse reactions associated with the use of health products to the Canada Vigilance Program by one of the following 3 ways:

- Report online at www.healthcanada.gc.ca/medeffect
- Call toll-free at 1-866-234-2345
- Complete a Canada Vigilance Reporting Form and:
 - Fax toll-free to 1-866-678-6789, or
 - Mail to: Canada Vigilance Program
Health Canada
Postal Locator 0701C
Ottawa, ON K1A 0K9

Postage paid labels, Canada Vigilance Reporting Form and the adverse reaction reporting guidelines are available on the MedEffect™ Canada Web site at www.healthcanada.gc.ca/medeffect.

NOTE: Should you require information related to the management of side effects, contact your health professional. The Canada Vigilance Program does not provide medical advice.

MORE INFORMATION

This document plus the full product monograph, prepared for health professionals can be found at:
<http://www.cobaltpharma.com>
 or by contacting the sponsor, Cobalt Pharmaceuticals Inc.,
 at: 1-866-254-6111.

This leaflet was prepared by:
 Cobalt Pharmaceuticals Inc.
 Mississauga, ON
 Canada, L5N 2B8

Last revised: February 3, 2010