

PRODUCT MONOGRAPH

^{Pr}Zym-AMLODIPINE

(Amlodipine Besylate Tablets)

5 mg and 10 mg Amlodipine (from amlodipine besylate)

Antihypertensive - Antianginal Agent

ZYMCAN PHARMACEUTICALS INC.
6111 Royalmount Ave., Suite 100
Montreal, Quebec H4P 2T4

Date of Preparation:
April 01, 2010

Control No. 135893

PrZym-Amlodipine	3
PART I: HEALTH PROFESSIONAL INFORMATION	3
SUMMARY PRODUCT INFORMATION	3
INDICATIONS AND CLINICAL USE	3
CONTRAINDICATIONS	4
WARNINGS AND PRECAUTIONS	4
ADVERSE REACTIONS	6
DRUG INTERACTIONS	8
DOSAGE AND ADMINISTRATION	9
OVERDOSAGE	10
ACTION AND CLINICAL PHARMACOLOGY	11
STORAGE AND STABILITY	14
DOSAGE FORMS, COMPOSITION AND PACKAGING	14
PART II: SCIENTIFIC INFORMATION	15
PHARMACEUTICAL INFORMATION	15
CLINICAL TRIALS	15
DETAILED PHARMACOLOGY	16
TOXICOLOGY	19
REFERENCES	24
PART III: CONSUMER INFORMATION	26

PrZym-AMLODIPINE

(Amlodipine Besylate Tablets)

5 mg and 10 mg Amlodipine (from amlodipine besylate)

Antihypertensive - Antianginal Agent

PART I: HEALTH PROFESSIONAL INFORMATION

SUMMARY PRODUCT INFORMATION

Route of Administration	Dosage Form/ Strength	Clinically Relevant Nonmedicinal Ingredients
Oral	Tablet 5 mg & 10 mg	None <i>For a complete listing see Dosage Forms, Composition and Packaging section.</i>

INDICATIONS AND CLINICAL USE

Hypertension

Zym-AMLODIPINE (amlodipine besylate) is indicated in the treatment of mild to moderate essential hypertension.

Combination of Zym-AMLODIPINE with a diuretic, a beta-blocking agent, or an angiotensin converting enzyme inhibitor has been found to be compatible and showed additive antihypertensive effect.

Chronic Stable Angina

Zym-AMLODIPINE is indicated for the management of chronic stable angina (effort-associated angina) in patients who remain symptomatic despite adequate doses of beta-blockers and/or organic nitrates or who cannot tolerate those agents.

Zym-AMLODIPINE may be tried in combination with beta-blockers in chronic stable angina in patients with normal ventricular function. When such concomitant therapy is introduced, care must be taken to monitor blood pressure closely since hypotension can occur from the combined effects of the drugs.

CONTRAINDICATIONS

- Patients who are hypersensitive to this drug or other dihydropyridines, or to any ingredient in the formulation or component of the container. For a complete listing, see the **Dosage Forms, Composition and Packaging** section of the product monograph.
- Patients with severe hypotension (less than 90 mmHg systolic).

WARNINGS AND PRECAUTIONS

General

Beta-blocker withdrawal

Amlodipine gives no protection against the dangers of abrupt beta-blocker withdrawal and such withdrawal should be done by the gradual reduction of the dose of beta-blocker.

Peripheral Edema

Mild to moderate peripheral edema was the most common adverse event in the clinical trials (see **ADVERSE REACTIONS**). The incidence of peripheral edema was dose-dependent and ranged in frequency from 3.0 to 10.8% in 5 to 10 mg dose range. Care should be taken to differentiate this peripheral edema from the effects of increasing left ventricular dysfunction.

Cardiovascular

Increased Angina and/or Myocardial Infarction

Rarely, patients, particularly those with severe obstructive coronary artery disease, have developed documented increased frequency, duration and/or severity of angina or acute myocardial infarction on starting calcium channel blocker therapy or at the time of dosage increase. The mechanism of this effect has not been elucidated.

Outflow Obstruction (Aortic Stenosis)

Amlodipine should be used with caution in a presence of fixed left ventricular outflow obstruction (aortic stenosis).

Use in Patients With Congestive Heart Failure

Although generally calcium channel blockers should only be used with caution in patients with heart failure, it has been observed that amlodipine had no overall deleterious effect on survival and cardiovascular morbidity in both short-term and long-term clinical trials in these patients. While a significant proportion of the patients in these studies had a history of ischemic heart disease, angina or hypertension, the studies were not designed to evaluate the treatment of angina or hypertension in patients with concomitant heart failure.

Hypotension

Amlodipine may occasionally precipitate symptomatic hypotension. Careful monitoring of blood pressure is recommended, especially in patients with a history of cerebrovascular insufficiency, and those taking medications known to lower blood pressure.

Hepatic/biliary/Pancreatic

Use in Patients with Impaired Hepatic Function

There are no adequate studies in patients with liver dysfunction and dosage recommendations have not been established. In a small number of patients with mild to moderate hepatic impairment given single dose of 5 mg, amlodipine half-life has been prolonged (see **ACTION AND CLINICAL PHARMACOLOGY, Pharmacokinetics and Metabolism**). Amlodipine should, therefore, be administered with caution in these patients and careful monitoring should be performed. A lower starting dose may be required (see **DOSAGE AND ADMINISTRATION**).

Special Population

Pregnant Women:

Although amlodipine was not teratogenic in the rat and rabbit some dihydropyridine compounds have been found to be teratogenic in animals. In rats, amlodipine has been shown to prolong both the gestation period and the duration of labor. There is no clinical experience with amlodipine in pregnant women. Amlodipine should be used during pregnancy only if the potential benefit outweighs the potential risk to the mother and fetus.

Nursing Women:

It is not known whether amlodipine is excreted in human milk. Since amlodipine safety in newborns has not been established, amlodipine should not be given to nursing mothers.

Pediatrics (0 to 16 years of age):

The use of amlodipine is not recommended in patients less than 6 years of age since safety and efficacy have not been established in that population. Pediatric studies beyond 8 weeks of duration have not been conducted.

The effect of amlodipine on blood pressure in patients less than 6 years of age is not known. The pediatric administration should be based on a careful risk/benefit assessment of the limited available information. The risk/benefit assessment should be conducted by a qualified physician.

Geriatrics (>65 years of age):

In elderly patients clearance of amlodipine is decreased with a resulting increase in AUC (see **ACTION AND CLINICAL PHARMACOLOGY, Pharmacokinetics and Metabolism**). In clinical trials the incidence of adverse reactions in elderly patients was approximately 6% higher than that of younger population (<65 years). Adverse reactions include edema, muscle cramps and dizziness. Amlodipine should be used cautiously in elderly patients. Dosage adjustment is advisable (see **DOSAGE AND ADMINISTRATION**).

ADVERSE REACTIONS

Clinical Trial Adverse Drug Reactions

Because clinical trials are conducted under very specific conditions the adverse reaction rates observed in the clinical trials may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse drug reaction information from clinical trials is useful for identifying drug-related adverse events and for approximating rates.

Amlodipine has been administered to 1,714 patients (805 hypertensive and 909 angina patients) in controlled clinical trials (vs placebo alone and with active comparative agents). Most adverse reactions reported during therapy were of mild to moderate severity.

Hypertension

In the 805 hypertensive patients treated with amlodipine in controlled clinical trials, adverse effects were reported in 29.9% of patients and required discontinuation of therapy due to side effects in 1.9% of patients. The most common adverse reactions in controlled clinical trials were: edema (8.9%), and headache (8.3%).

The following adverse reactions were reported with an incidence of $\geq 0.5\%$ in the controlled clinical trials program (n=805):

Cardiovascular: edema (8.9%), palpitations (2.0%), tachycardia (0.7%), postural dizziness (0.5%).

Skin and Appendages: pruritus (0.7%).

Musculoskeletal: muscle cramps (0.5%).

Central and Peripheral Nervous System: headache (8.3%), dizziness (3.0%), paresthesia (0.5%).

Autonomic Nervous System: flushing (3.1%), increased sweating (0.9%), dry mouth (0.7%).

Psychiatric: somnolence (1.4%).

Gastrointestinal: nausea (2.4%), abdominal pain (1.1%), dyspepsia (0.6%), constipation (0.5%).

General: fatigue (4.1%), pain (0.5%).

Angina

In the controlled clinical trials in 909 angina patients treated with amlodipine, adverse effects were reported in 30.5% of patients and required discontinuation of therapy due to side effects in 0.6% of patients. The most common adverse reactions reported in controlled clinical trials were: edema (9.9%) and headache (7.8%).

The following adverse reactions occurred at an incidence of $\geq 0.5\%$ in the controlled clinical trials program (n=909);

Cardiovascular: edema (9.9%), palpitations (2.0%), postural dizziness (0.6%).

Skin and Appendages: rash (1.0%), pruritus (0.8%).

Musculoskeletal: muscle cramps (1.0%).

Central and Peripheral Nervous System: headache (7.8%), dizziness (4.5%), paresthesia (1.0%), hypoesthesia (0.9%).

Autonomic Nervous System: flushing (1.9%).

Psychiatric: somnolence (1.2%), insomnia (0.9%), nervousness (0.7%).

Gastrointestinal: nausea (4.2%), abdominal pain (2.2%), dyspepsia (1.4%), diarrhea (1.1%), flatulence (1.0%), constipation (0.9%).

Respiratory System: dyspnea (1.1%).

Special Senses: vision abnormal (1.3%), tinnitus (0.6%).

General: fatigue (4.8%), pain (1.0%), asthenia (1.0%).

Less Common Clinical Trial Adverse Drug Reactions (<1%)

Amlodipine has been evaluated for safety in about 11,000 patients with hypertension and angina. The following events occurred in <1% but >0.1% of patients in comparative clinical trials (double-blind comparative vs placebo or active agents; n = 2,615) or under conditions of open trials or marketing experience where a causal relationship is uncertain.

Cardiovascular: arrhythmia (including ventricular tachycardia and atrial fibrillation), bradycardia, hypotension, peripheral ischemia, syncope, tachycardia, postural dizziness, postural hypotension, vasculitis.

Central and Peripheral Nervous System: hypoesthesia, peripheral neuropathy, tremor, vertigo.

Gastrointestinal: anorexia, constipation, dysphagia, vomiting, gingival hyperplasia.

General: allergic reaction, asthenia⁺, back pain, hot flushes, malaise, rigors, weight gain.

Musculoskeletal System: arthralgia, arthrosis, myalgia.

Psychiatric: sexual dysfunction (male⁺ and female), insomnia, nervousness, depression, abnormal dreams, anxiety, depersonalization.

Respiratory System: epistaxis.

Skin and Appendages: pruritus⁺, rash erythematous, rash maculopapular, erythema multiforme.

Special Senses: conjunctivitis, diplopia, eye pain, tinnitus.

Urinary System: micturition frequency, micturition disorder, nocturia.

Autonomic Nervous System: dry mouth, sweating increased.

Metabolic and Nutritional: hyperglycemia, thirst.

Hemopoietic: leucopenia, purpura, thrombocytopenia.

⁺These events occurred in less than 1% in placebo controlled trials, but the incidence of these side effects was between 1% and 2% in all multiple dose studies.

The following events occurred in ≤0.1% of patients: cardiac failure, skin discoloration, urticaria, skin dryness, Stevens-Johnson syndrome, alopecia, twitching, ataxia, hypertonia, migraine, apathy, amnesia, gastritis, pancreatitis, increased appetite, coughing, rhinitis, parosmia, taste perversion, and xerophthalmia.

Isolated cases of angioedema have been reported. Angioedema may be accompanied by breathing difficulty.

Post-Market Adverse Drug Reaction

In postmarketing experience, jaundice and hepatic enzyme elevations (mostly consistent with cholestasis or hepatitis) in some cases severe enough to require hospitalization have been reported in association with use of amlodipine.

DRUG INTERACTIONS

Drug-Drug Interactions

As with all drugs, care should be exercised when treating patients with multiple medications. Dihydropyridine calcium channel blockers undergo biotransformation by the cytochrome P450 system, mainly via CYP 3A4 isoenzyme. Coadministration of amlodipine with other drugs which follow the same route of biotransformation may result in altered bioavailability of amlodipine or these drugs. Dosages of similarly metabolized drugs, particularly those of low therapeutic ratio, and especially in patients with renal and/or hepatic impairment, may require adjustment when starting or stopping concomitantly administered amlodipine to maintain optimum therapeutic blood levels.

Drugs known to be inhibitors of the cytochrome P450 system include: azole antifungals, cimetidine, cyclosporine, erythromycin, quinidine, terfenadine, warfarin.

Drugs known to be inducers of the cytochrome P450 system include: phenobarbital, phenytoin, rifampin.

Drugs known to be biotransformed via P450 include: benzodiazepines, flecainide, imipramine, propafenone, theophylline.

Amlodipine has a low (rate of first-pass) hepatic clearance and consequent high bioavailability, and thus, may be expected to have a low potential for clinically relevant effects associated with elevation of amlodipine plasma levels when used concomitantly with drugs that compete for or inhibit the cytochrome P450 system.

Cimetidine, Warfarin, Cyclosporin, Digoxin

Pharmacokinetic interaction studies with amlodipine in healthy volunteers have indicated:

- **cimetidine** did not alter the pharmacokinetics of amlodipine.
- amlodipine did not change **warfarin**-induced prothrombin response time.
- amlodipine does not significantly alter the pharmacokinetics of **cyclosporin**.
- amlodipine did not change serum **digoxin** levels or **digoxin** renal clearance.

Antacids

Concomitant administration of Maalox[®] (magnesium hydroxide and aluminum hydroxide) had no effect on the disposition of a single 5 mg dose of amlodipine in 24 subjects.

Beta-blockers

When beta-adrenergic receptor blocking drugs are administered concomitantly with amlodipine, patients should be carefully monitored since blood pressure lowering effect of beta-blockers may be augmented by amlodipine's reduction in peripheral vascular resistance.

Sildenafil

A single 100 mg dose of sildenafil (VIAGRA) in subjects with essential hypertension had no effect on AUC_t or C_{max} of amlodipine. When sildenafil (100 mg) was co-administered with amlodipine, 5 or 10 mg in hypertensive patients, the mean additional reduction of supine blood pressure was 8 mm Hg systolic and 7 mm Hg diastolic.

Atorvastatin

In healthy volunteers, co-administration of multiple 10 mg doses of amlodipine with 80 mg of atorvastatin resulted in no significant change in the AUC_t, C_{max} or T_{max} of atorvastatin.

Drug-Food Interactions**Interaction with Grapefruit Juice**

Published data indicate that through inhibition of the cytochrome P450 system, grapefruit juice can increase plasma levels and augment pharmacodynamic effects of some dihydropyridine calcium channel blockers. Following oral administration of 10 mg amlodipine to 20 male volunteers, pharmacokinetics of amlodipine were similar when amlodipine was administered with and without grapefruit juice. (See **ACTION AND CLINICAL PHARMACOLOGY - Pharmacokinetics**)

Drug-Herb Interactions

Interactions with herbal products have not been established.

Drug-Laboratory Test Interactions

There are no known interactions of Amlodipine Besylate with commonly used laboratory tests.

DOSAGE AND ADMINISTRATION**Dosing Considerations**

- Elderly patients or patients with impaired renal function
- Patients with impaired hepatic functions

Dosage should be individualized depending on patient's tolerance and responsiveness.

For both hypertension and angina, the recommended initial dose of Zym-AMLODIPINE is 5 mg once daily. If necessary, dose can be increased after 1-2 weeks to a maximum dose of 10 mg once daily.

Use in the Elderly or in Patients with Impaired Renal Function

The recommended initial dose in patients over 65 years of age or patients with impaired renal function is 5 mg once daily. If required, increasing in the dose should be done gradually and with caution (see **WARNINGS AND PRECAUTIONS**).

Use in Patients with Impaired Hepatic Function

Dosage requirements have not been established in patients with impaired hepatic function. When Zym-AMLODIPINE is used in these patients, the dosage should be carefully and gradually adjusted depending on patients tolerance and response. A lower starting dose of 2.5 mg once daily should be considered (see **WARNINGS AND PRECAUTIONS**).

Use in Children

The effective antihypertensive oral dose in pediatric patients ages 6-17 years is 2.5 mg to 5 mg once daily. Doses in excess of 5 mg daily have not been studied; dose should be determined based upon the medical need of the patients. See **ACTION AND CLINICAL PHARMACOLOGY**.

OVERDOSAGE

Symptoms

Overdosage can cause excessive peripheral vasodilation with marked and probably prolonged hypotension and possibly a reflex tachycardia. In humans, experience with overdosage of amlodipine is limited. When amlodipine was ingested at doses of 105-250 mg some patients remained normotensive with or without gastric lavage while another patient experienced hypotension (90/50 mmHg) which normalized following plasma expansion. A patient who took 70 mg of amlodipine with benzodiazepine developed shock which was refractory to treatment and died. In a 19 month old child who ingested 30 mg of amlodipine (about 2 mg/kg) there was no evidence of hypotension but tachycardia (180 bpm) was observed. Ipecac was administered 3.5 hrs after ingestion and on subsequent observation (overnight) no sequelae were noted.

Treatment

Clinically significant hypotension due to overdosage requires active cardiovascular support including frequent monitoring of cardiac and respiratory function, elevation of extremities, and attention to circulating fluid volume and urine output. A vasoconstrictor (such as norepinephrine) may be helpful in restoring vascular tone and blood pressure, provided that there is no contraindication to its use. As amlodipine is highly protein bound, hemodialysis is not likely to be of benefit. Intravenous calcium gluconate may be beneficial in reversing the effects of calcium channel blockade. Clearance of amlodipine is prolonged in elderly patients and in patients with impaired liver function. Since amlodipine absorption is slow, gastric lavage may be worthwhile in some cases.

ACTION AND CLINICAL PHARMACOLOGY

Amlodipine besylate is a calcium ion influx inhibitor (calcium entry blocker or calcium ion antagonist). Amlodipine is a member of the dihydropyridine class of calcium antagonists.

Mechanism of Action

The therapeutic effect of this group of drugs is believed to be related to their specific cellular action of selectively inhibiting transmembrane influx of calcium ions into vascular smooth muscle and cardiac muscle. The contractile processes of these tissues are dependent upon the movement of extracellular calcium ions into these cells through specific ion channels. Amlodipine inhibits calcium ion influx across cell membranes selectively, with a greater effect on vascular smooth muscle cells than on cardiac muscle cells. Serum calcium concentration is not affected by amlodipine. Within the physiologic pH range, amlodipine is an ionized compound and its kinetic interaction with the calcium channel receptor is characterized by the gradual association and dissociation with the receptor binding site. Experimental data suggest that amlodipine binds to both dihydropyridine and nondihydropyridine binding sites.

- A. Hypertension** The mechanism by which amlodipine reduces arterial blood pressure involves direct peripheral arterial vasodilation and reduction in peripheral vascular resistance.
- B. Angina** The precise mechanism by which amlodipine relieves angina has not been fully delineated. Amlodipine is a dilator of peripheral arteries and arterioles which reduces the total peripheral resistance and, therefore, reduces the workload of the heart (afterload). The unloading of the heart is thought to decrease ischemia and relieve effort angina by reducing myocardial energy oxygen consumption and oxygen requirements.

Pharmacokinetics and Metabolism

After oral administration of therapeutic doses of amlodipine, absorption occurs gradually with peak plasma concentration reached between 6 and 12 hours. Absolute bioavailability has been estimated to be between 64 and 90%. The bioavailability of amlodipine is not altered by the presence of food.

Amlodipine is metabolized through the cytochrome P450 system, mainly via CYP 3A4 isoenzyme. Amlodipine is extensively (about 90%) converted to inactive metabolites (via hepatic metabolism) with 10% of the parent compound and 60% of the metabolites excreted in the urine. *Ex vivo* studies have shown that approximately 93% of the circulating drug is bound to plasma proteins in hypertensive patients. Elimination from the plasma is biphasic with a terminal elimination half-life of about 35-50 hours. Steady state plasma levels of amlodipine are reached after 7 to 8 days of consecutive daily dosing.

The pharmacokinetics of amlodipine are not significantly influenced by renal impairment. Plasma concentrations in the patients with moderate to severe renal failure were higher than in the normal subjects. Accumulation and mean elimination half-life in all patients were within the range of those observed in other pharmacokinetic studies with amlodipine in normal subjects.

In elderly hypertensive patients (mean age 69 years) there was a decrease in clearance of amlodipine from plasma as compared to young volunteers (mean age 36 years) with a resulting increase in the area under the curve (AUC) of about 60%.

Following single oral administration of 5 mg of amlodipine, patients with chronic mild-moderate hepatic insufficiency showed about 40% increase in AUC of amlodipine as compared to normal volunteers. This was presumably due to a reduction in clearance of amlodipine as the terminal elimination half-life was prolonged from 34 hrs in young normal subjects to 56 hrs in the elderly patients with hepatic insufficiency.

Following oral administration of 10 mg amlodipine to 20 male volunteers, pharmacokinetics of amlodipine were similar when amlodipine was administered with and without grapefruit juice. Geometric mean C_{max} of amlodipine was 6.2 ng/mL when the drug was administered with grapefruit juice and 5.8 ng/mL when administered with water. Mean T_{max} of amlodipine was 7.6 hours with grapefruit juice and 7.9 hours with water. Geometric mean AUC_{0-∞} was 315 ng/hr/mL with grapefruit juice and 293 ng/hr/mL with water. Geometric mean bioavailability of amlodipine was 85% when administered with grapefruit juice and 81% when administered with water.

Pediatric Patients

Two studies were conducted to evaluate the use of amlodipine in a pediatric population.

In one study (pharmacokinetic), sixty-two hypertensive patients aged greater than 6 years received doses of amlodipine between 1.25 mg and 20 mg. Weight-adjusted clearance and volume of distribution were similar to values in adults (see **DOSAGE AND ADMINISTRATION**). The mean absorption rate constant (K_a) in children (0.85 hr⁻¹) is approximately 50% higher than that in healthy adults (0.55 hr⁻¹, range of 0.28-1.09hr⁻¹).

Gender effect: In a second trial (clinical), a pattern of greater reductions in both systolic and diastolic blood pressure in females than in males was observed. Mean change in systolic blood pressure from baseline to end of study: amlodipine 2.5 mg: males, -6.9 mmHg (n=51); females, -8.9 mmHg (n=32); amlodipine 5.0 mg: males, -6.6 mmHg (n=63); females, -14.0 mmHg (n=23); placebo males, -2.5 mmHg (n=54), females, -3.8 mmHg (n=33).

Pharmacodynamics

Hemodynamics

Following administration of recommended doses to patients with hypertension, amlodipine produces vasodilation resulting in a reduction of supine and standing blood pressures. These decreases in blood pressure are not accompanied by any significant change in heart rate or plasma catecholamine levels with chronic dosing. With chronic once daily oral administration (5 and 10 mg once daily), antihypertensive effectiveness is maintained throughout the 24 hours dose interval with minimal peak to trough differences in blood pressure reduction. Since the vasodilation induced by amlodipine is gradual in onset, acute hypotension has rarely been reported after oral administration of amlodipine. In normotensive patients with angina

amlodipine has not been associated with any clinically significant reductions in blood pressure or changes in heart rate.

Negative inotropic effects have not been observed when amlodipine was administered at the recommended doses to man, but has been demonstrated in animal models. Hemodynamic measurements of cardiac function at rest and during exercise (or pacing) in angina patients with normal ventricular function have generally demonstrated a small increase in cardiac index without significant influence on dP/dt or on left ventricular end diastolic pressure or volume.

In hypertensive patients with normal renal function, therapeutic doses of amlodipine resulted in a decrease in renal vascular resistance and an increase in glomerular filtration rate and effective renal plasma flow without change in filtration fraction.

Electrophysiologic Effects

Amlodipine does not change sinoatrial nodal function or atrioventricular conduction in intact animals, or man. In patients with chronic stable angina, intravenous administration of 10 mg of amlodipine and a further 10 mg of amlodipine after a 30 min. interval produced peripheral vasodilation and afterload reduction, but did not significantly alter A-H and H-V conduction and sinus node recovery time after pacing. Similar results were obtained in patients receiving amlodipine and concomitant beta-blockers. In clinical studies in which amlodipine was administered in combination with beta-blockers to patients with either hypertension or angina, no adverse effects on electrocardiographic parameters were observed. In clinical trials with angina patients, amlodipine as monotherapy did not alter electrocardiographic intervals.

Effects in Hypertension

Pediatric Patients

Two hundred sixty-eight hypertensive patients aged 6 to 17 years were randomized first to amlodipine 2.5 or 5 mg once daily for 4 weeks and then randomized again to the same dose or to placebo for another 4 weeks. Patients receiving 5 mg at the end of 8 weeks had lower blood pressure than those secondarily randomized to placebo. The magnitude of the treatment effect is difficult to interpret, but it is probably less than 5 mmHg systolic on the 5 mg dose. Adverse events were similar to those seen in adults.

Pediatric safety and efficacy studies beyond 8 weeks of duration have not been conducted. In addition, the long-term effect of amlodipine on growth and development, myocardial growth and vascular smooth muscles has not been studied.

STORAGE AND STABILITY

Zym-AMLODIPINE tablets should be stored between 15° to 30°C, protected from light.

DOSAGE FORMS, COMPOSITION AND PACKAGING

Availability of Dosage Forms

Zym-AMLODIPINE is available as octagonal tablets in the following strengths as described below:

5 mg: White, octagonal tablets with “P” logo and “5” debossed on each side of the score on one side and nothing on the other side.

10 mg: White, octagonal tablet with “P” logo debossed on one side and “10” on the other side.

Supplied: 5 mg and 10 mg: in HDPE bottles of 100 tablets and 500 tablets.

Composition

Zym-AMLODIPINE 5 mg and 10 mg Tablets contain 5 mg and 10 mg Amlodipine, as Amlodipine Besylate, respectively. The tablets also contain the following non-medical ingredients (alphabetically): Dibasic calcium phosphate anhydrous, Magnesium stearate, Microcrystalline Cellulose and Sodium starch glycolate.

Product Monograph available upon request.

Zymcan Pharmaceuticals Inc.
Montréal, Québec
H4P 2T4

PART II: SCIENTIFIC INFORMATION

PHARMACEUTICAL INFORMATION


Drug Substance

Proper Name: Amlodipine Besylate

Chemical Name: 3-Ethyl 5-methyl (\pm)-2-[(2-aminoethoxy)methyl]-4-(o-chlorophenyl)-1,4-dihydro-6-methyl-3,5-pyridinedicarboxylate monobenzenesulfonate

Molecular formula and molecular mass: $C_{20}H_{25}ClN_2O_5 \cdot C_6H_6O_3S$, 567.1 g/mol

Structural formula:



and enantiomer

Physicochemical properties: Amlodipine is a white or almost white powder. Amlodipine is slightly soluble in water, freely soluble in methanol, and sparingly soluble in ethanol. The melting range is 201°C to 205°C with decomposition.

CLINICAL TRIALS

Comparative Bioavailability Studies

A single-dose, crossover, comparative bioavailability study was performed in healthy male volunteers (n=19) under fasting conditions on Amlodipine tablets using Zymcan Pharmaceuticals Inc. Zym-AMLODIPINE 10 mg tablets (Lot # P-1288) versus the reference product, NORVASC[®] 10 mg Tablets (Lot # 402-05281), by Pfizer Canada Inc. The pharmacokinetic data calculated for the Zym-AMLODIPINE 10 mg tablets and NORVASC[®] 10 mg tablets formulation are tabulated below:

SUMMARY TABLE OF THE COMPARATIVE BIOAVAILABILITY DATA

Amlodipine (1 x 10 mg) From measured data uncorrected for potency Geometric Mean Arithmetic Mean (CV %)				
Parameter	Zym-Amlodipine	Norvasc [†]	% Ratio of Geometric Means	90% Confidence Interval
AUC ₀₋₇₂ (ng.h/mL)	237.238 246.992 (27.6)	224.294 234.314 (29.6)	105.77	101.17 - 110.58
AUC _I (ng.h/mL)	349.389 370.719 (35.4)	341.369 361.387 (34.2)	102.35	95.25 - 109.98
C _{max} (ng/mL)	6.123 6.371 (26.2)	5.797 6.033 (27.0)	105.63	99.98 - 111.59
T _{max} [§] (h)	7.00 (5.00 - 11.00)	7.00 (5.00 - 14.00)		
T _{1/2} [€] (h)	41.83 (27.2)	45.15 (21.0)		

[†] Norvasc was manufactured by Pfizer Canada Inc. and was purchased in Canada.

[§] Expressed as the median (range) only.

[€] Expressed as the arithmetic mean (CV%) only

DETAILED PHARMACOLOGY

ANIMAL

a. Mechanism of Action Studies - *In Vitro*

Amlodipine inhibited both calcium-induced and potassium-depolarization-induced contractions of rat aorta. The inhibitory effect was gradual. The potency of amlodipine was more than 10-fold greater against Ca²⁺-responses than against K⁺-responses. Studies in both rat aorta and dog coronary artery indicated that amlodipine was a competitive antagonist. Radioligand binding experiments designed to characterize the interactions of amlodipine with calcium channel binding sites in bovine brain and in cardiac membranes from dog and rat

showed that amlodipine interacts competitively and at high affinity with the dihydropyridine (DHP) recognition site.

Amlodipine has been demonstrated to block constriction of coronary arteries and arterioles in response to calcium, potassium, epinephrine, serotonin, and thromboxane A₂ analog in experimental animal models and in human coronary vessels *in vitro*.

Electrophysiological experiments conducted using isolated papillary muscles from guinea pig hearts confirmed that amlodipine was a highly selective calcium channel blocker which inhibited cardiac slow action potentials in a non-use-dependent manner and with no effect on the fast Na⁺-channel.

In Langendorff-perfused guinea pig hearts, amlodipine showed negative inotropic activity, the concentration producing a 50% inhibition of cardiac contraction being approximately 10 times greater (20.2 nM) than for a 50% inhibition of vascular muscle contraction (1.9 nM). The drug displayed modest negative chronotropic effect (approximately 20%) at a concentration of 50 nM, approximately twice that required for 50% inhibition of cardiac contraction in the same preparation. Using Langendorff-perfused rat hearts the concentration producing a 50% inhibition of cardiac contraction was 300 times greater than for inhibition of coronary artery contraction.

b. Cardiovascular Activity - *In Vivo*

In anesthetized dogs, amlodipine (i.v. 25-1600 µg/kg) was a potent coronary and peripheral vasodilator; ED₅₀ values were 103 and 212 µg/kg for reductions of coronary and systemic vascular resistances respectively. The reductions in vascular resistance were associated with corresponding increases in cardiac output, coronary flow, heart rate and myocardial contractility. Amlodipine possessed slow onset of action, minimal effect on blood pressure, and a long duration of action. Amlodipine caused slight, transient negative inotropic responses only at the highest dose, in excess of that required to cause maximal vasodilatation. The drug did not adversely affect atrial ventricular conduction, as assessed by PR interval.

Oral administration of amlodipine (0.5 to 2.0 mg/kg) to conscious dogs produced dose-related reductions in systemic vascular resistance (max. of 78%) and reflexly-induced increases in heart rate cardiac output and myocardial contractility; maximum effects were achieved much later (3 to 5 h) than after parenteral administration (5 to 30 min) which may explain the dose-related modest blood pressure reductions (max. change of 25%) observed by the oral route.

c. Antihypertensive Efficacy - *In Vivo*

Amlodipine produced dose-related reductions in blood pressure of spontaneously hypertensive rats (SHR) after oral administration. The antihypertensive effect was maintained for at least 6 h after each one of the 3 doses used (1, 3, and 10 mg/kg). In young SHR the development of hypertension was attenuated by 60% over a 12 week period when amlodipine was added to the diet to provide the dose of 8 mg/kg/day. In mature SHR receiving amlodipine for 8 weeks, a marked antihypertensive effect was evident by day 2 and attained a maximum by day 5. This effect was maintained for the remaining treatment period with no

change in heart rate. In addition, treated animals showed a small, but statistically significant, reduction in ventricular weight and marked elevation in plasma renin activity.

In conscious renal-hypertensive dogs, oral administration of single doses of amlodipine (0.25, 0.5 and 1.0 mg/kg) produced dose-related reductions in blood pressures with maximum effects occurring at 5 h after dose. These responses were accompanied by dose-related increases in heart rate.

The slow onset and long-lasting antihypertensive effects of amlodipine were confirmed in conscious renal-hypertensive dogs in which blood pressure was recorded continuously for 24 h.

In conscious renal-hypertensive dogs, orally-administered amlodipine (0.025, 0.05 and 0.25 mg/kg/day) for 10-14 days produced progressive reductions in the daily, resting, pre-dose blood pressure which stabilized after 4 or 5 days. The minimum blood pressures achieved each day were approximately equivalent and tolerance did not develop. Heart rate was inconsistently affected.

d. General Pharmacology

In both normotensive (fluid-loaded) and spontaneously hypertensive rats (SHR) amlodipine produced diuresis and natriuresis. A diuretic effect was also observed in saline loaded conscious or anesthetized dogs treated with low intravenous doses (less than 0.4 mg/kg) of amlodipine; increases in potassium excretion were not significant. Also in the conscious rat amlodipine produced dose-related reduction of basal gastric acid secretion and a small but significant reduction in gastro-intestinal motility. Experiments in anesthetized dogs indicated that phenylephrine was an effective antidote to the hypotensive effect of a supra-maximal dose of amlodipine.

TOXICOLOGY

Acute Toxicity - Amlodipine (as maleate unless otherwise indicated)

			LD ₅₀	Range of Lethal Doses (mg/kg)	
SPECIES	SEX	ROUTE	base/mg/kg	No Deaths	All Dead
Mice	M	p.o.	N.D.	10	40
	F	p.o.	N.D.	10	40
	M	i.v.	N.D.	2.5	10
	F	i.v.	N.D.	2.5	10
Rats	M	p.o.	150	2/10 at 100	400
	F	p.o.	140	2/10 at 100	250
	M	i.v.	N.D.	1	10
	F	i.v.	N.D.	1	10
Rats*	M	p.o.	393**		
	F	p.o.	686**		

* Sprague Dawley Rats from Shizouka Lab Animal Centre, Hamamatsu, Japan

** Besylate Salt

+ Dogs from Interfauna, France

++ Dogs from Japan

N.D. Not Determined: The result did not permit calculations of LD₅₀ values. Thus, range of lethal doses is given.

The main clinical signs in the oral studies were somnolence, decreased spontaneous movement and for rats salivation, dyspnea, ptosis, lacrimation, blanching, cyanosis, rough coat, abdominal distension, and eventually coma. After i.v. injection, the animals died rapidly showing only somnolence, tachypnea or ptosis.

TOXICOLOGY

SPECIES	ROUTE	DOSE base mg/kg/day	ANIMAL PER DOSE LEVEL	DURATIO N	FINDINGS
---------	-------	------------------------	-----------------------------	--------------	----------

MAXIMUM TOLERATED DOSE (SINGLE)

Dog	Oral (gavage)	4 8 16	2 M	Single Dose	<p><u>At all dose levels:</u> Vasodilation and increases in plasma aldosterone levels.</p> <p><u>At 4 mg/kg:</u> Compensatory tachycardia.</p> <p><u>At 8 mg/kg:</u> In 1 of 2 dogs vomiting, sedation, respiratory distress and diarrhea 48 hr post-dose; normal at day 5. Compensatory tachycardia.</p> <p><u>At 16 mg/kg:</u> Moribund with hyperthermia within 24 hours; low blood pressure returned to normal over 2-6 days; transient raise in heart rate.</p> <p><u>Histological examination</u> showed congestion, edema and hemorrhage of the right atrial wall in the 2 dogs at 16 mg/kg. The hemorrhage in the right atrial wall corresponds to the right atrial lesions seen in long-term studies with amlodipine and other vasodilators (see long-term toxicity). One of 2 dogs at each dose showed fibrosis of the left ventricle in the subendocardial region and the posterior papillary muscle.</p> <p>The maximum tolerated dose was not determined.</p>
Dog (Japanese Study)	Oral	3.5 7	1 M 1 F	Single Dose	<p><u>Mortality:</u> 1 male dog at 7 mg/kg. Decreased spontaneous movement and flushing of palpebral conjunctiva and buccal cavity.</p> <p><u>At 7 mg/kg:</u> 1 female vomiting; 1 male hypothermia, lying prone.</p> <p>Hematology/Clinical Chemistry: Increase in WBC and BUN at 10 and 5 mg/kg (males).</p> <p>The maximum tolerated dose was not determined.</p>

SUBACUTE AND CHRONIC TOXICITY

Mouse	Oral (diet)	0 2.5 5 10	10 M 10 F	2 Months	<u>At 10 mg/kg/day</u> : Mice died during week 2 of the study. <u>At 5 mg/kg/day (males and females) and 2.5 mg/kg/day (males)</u> : Increase in water consumption. <u>At 5 mg/kg/day - Pathology</u> : Drug-related increases in heart and liver weights.
Rat (Japanese Study)	Oral (gavage)	0 4 16 32 64	12 M 12 F	1 Month	<u>At 64 mg/kg/day</u> : All rats died within 9 days. <u>At 32 mg/kg/day</u> : 12/24 rats died; decreased food consumption, growth inhibition, ptosis, decreased spontaneous movement. <u>At 16 and 32 mg/kg/day</u> : The pattern of results on heart weights, increased urinary volume, effect on electrolyte balance and the adrenals was similar to that of the 6 month study below; increase in BUN at 16 mg/kg (males) and at 32 mg/kg (males and females).
Rat (Japanese Study)	Oral (gavage)	0 2 7 21	16 M 16 F	3 Months followed by 1 Month drug withdrawal	<u>21 mg/kg/day</u> : Salivation, growth inhibition, increased BUN, increased urinary volume, effect on electrolyte balance and adrenals was similar to that of the 6 month study below. Also <u>post-mortem</u> dilation of small intestine without morphological lesions. <u>At 7 mg/kg/day</u> : Alterations in urinary electrolytes excretion. No drug related effects at the end of 1 month drug withdrawal phase.
Rat	Oral (gavage)	0 2.5 5 10	20 M 20 F	6 Months	<u>At all dose levels</u> : Renal effects: increased urinary volume and/or Na/K/Cl excretion, decreased plasma Na/K and/or Ca/Cl and increased urea; <u>Post-mortem</u> : Increase in heart weights. <u>At 10 mg/kg/day</u> : Renal effects: increased kidney weight. <u>Histopathology</u> : Thickening of zona glomerulosa at 5 and 10 mg/kg/day.
Rat (Japanese Study)	Oral (gavage)	1.4 7 18	30 M 30 F	12 Months (interim sacrifice 5/sex/group after 6 months)	<u>Mortality</u> : 3 rats (2 males and 1 female) at 18 mg/kg/day. <u>At 18 mg/kg/day</u> : Salivation, growth inhibition; Renal effects: increase in urinary volume with increased electrolytes excretion and decreased serum electrolytes; increase in BUN. <u>At 7 mg/kg/day</u> : Growth inhibition (males); Renal effects: increases of urinary volume and electrolyte excretion. <u>Post-mortem</u> : Increases of adrenal weights (at 18/mg/kg), increases of relative heart weight (18 and 7 mg/kg), dilated small intestines without morphological change (18 mg/kg). <u>Histopathology - Main Finding</u> : Enlargement of the zona glomerulosa of the adrenals (18 and 7 mg/kg).

Dog	Oral (gavage)	0.5 to 4	2 M 2 F	10 Days Supplementary Dose Escalation Study (0.5 mg/kg/day)	<u>At 4 mg/kg:</u> Death of all (4/4) dogs preceded in 3 dogs by low systolic blood pressure, bradycardia, disturbances of heart rhythm and conduction. Clinical signs included pale skin, hypothermia and prostration. <u>Histopathology:</u> Showed foci of myocyte necrosis and sarcoplasmic vacuolation in the left ventricle, papillary muscle and left and right atria. Congestion and/or edema in several organs (i.e. gastrointestinal tract/gall bladder wall and surrounding tissues as well as the connective tissue surrounding both kidneys).
Dog	Oral	0 0.25 0.5 1	3 M 3 F	6 Months	<u>At all dose levels:</u> Increase in urinary volume and urinary excretion of electrolytes (not dose-related). Reduction in blood pressure and increases in heart rate. <u>At 1 mg/kg/day - Pathology:</u> Increase in relative heart weights in 4/6 dogs, inflammatory lesion of the right atrial wall was seen which was considered to be consequence of excessive hemodynamic changes.
Dog	Oral	0 0.125 0.25 0.5	4 M 4 F	12 Months	<u>At 0.5 mg/kg/day:</u> Reduction in blood pressure and increases in heart rate; increase in urinary volume and urinary excretion of electrolytes (females). <u>At 0.5 mg/kg/day - Pathology:</u> Showed inflammatory lesions of the right atrial wall in 1/8 dogs, similar to that of the 6 month study above, and diffuse gingival hyperplasia.

MUTAGENICITY

Study	Test Organism	Dose	Route	Major Findings
<u>Ames Test</u> (modified) Quantitative Plate Assay (QAP) and Metabolic Activation (MA) with Hepatic Microsomes	<u>salmonella typhimurium:</u> Strains TA 1535, TA 1537, TA 98 and TA 100	10-0.02 mg/plate (QAP) 0.2-0.0005 mg/plate (MA)	<i>In-vitro</i>	No evidence of mutation frequency.
<u>In-vivo Cytogenetic Tests</u>	mouse bone marrow	20 mg/kg single dose 10 mg/kg/day for 5 days	<i>In-vivo</i> p.o. s.c.	No indication of chromosome breakage or mutagenicity observed.
<u>In vitro Cytogenetic Tests</u> with or without metabolic activation [rat liver microsomal enzymes (S-9)]	human lymphocytes	Without metabolic activation: 0.01 to 1000 µg/ML of culture medium With metabolic activation: 1.0 to 25 µg/ML of culture medium.	<i>In-vitro</i>	<u>Non-activation:</u> No evidence of induced chromosome breakage observed at levels of 1.0 µg/mL and below. At levels higher than 1.0 µg/mL, compound produced mitotic inhibition. <u>Activation:</u> No drug induced clastogenic activity observed at levels up to 10 µg/mL. Higher levels produced mitotic inhibition.
Quantitative Plate Assay (QAP) of Mouse Urine	<u>Salmonella typhimurium</u> Strains: TA 1535, TA 1537, TA 98 and TA 100.	0, 1, 10 and 20 mg/kg	<i>In-vivo</i> p.o.	No incidence of an excreted mutagen.
L 5178Y/TK +/- Gene Mutation Assay with and without liver S-9 fraction	mouse lymphoma cells	1.2 - 38 µg/mL	<i>In-vitro</i>	No evidence of gene mutational activity.

CARCINOGENICITY

There was no evidence of a carcinogenic effect when amlodipine was administered in the diet for up to 24 months to rats up to 2.5 mg/kg/day. Amlodipine was also administered for up to 24 months of dietary administration to mice at doses up to 2.5 mg/kg/day and no evidence of carcinogenicity was observed.

REPRODUCTION AND TERATOLOGY

Species	Route	Dose base/mg/kg/day	Animal per Dose Level	Duration	Findings
Fertility					
Rat (SD) (Japanese Study)	Oral (gavage)	0 1.4 7 18	24 M + 24 F	Males 71 days prior to and during mating. Females 14 days prior to and during mating and up to 7 days of gestation.	<u>At 18 mg/kg</u> : Impairment of body weight gain (females). There were no effects of the drug on copulation or pregnancy rates, nor any evidence of embryotoxicity or teratogenicity.
Teratology					
Rat (Charles River CD/SD)	Oral (gavage)	0 2 5 10	20 F	Days 6-15 post insemination. Hysterectomies on day 20 of gestation.	No effects were observed.
Rat (SD) Japanese Study	Oral (gavage)	0 3 7 18	34 F	Days 7-17 post-insemination. % of dams sacrificed on day 21 of gestation. F ₁ generation followed.	No effects were observed except in the dams. <u>At 18 mg/kg</u> : Reduction in food intake and body weight gain.
Rabbit (Japanese White) Japanese Study	Oral	0 3 7 18	18 or 19 F	Day 6 to day 18 of gestation.	<u>At 18 and 7 mg/kg</u> : Decrease in maternal body weight (18 mg/kg) decrease in food consumption (18 and 7 mg/kg). No evidence of drug induced fetotoxicity or teratogenicity.
Peri- and Post-Natal					
Rat (SD) Japanese Study	Oral (gavage)	0 1.4 2.8 7.0	25 F	Day 17 of gestation to day 21 post-partum.	As in the combined Fertility/Perinatal Study above; at the high dose level (7.0 mg/kg/day) adverse effects were observed on parturition and number of viable pups at birth and day 4 post-partum.

REFERENCES

1. Abernethy DR, Gutkowska J, Lambert MD. Amlodipine in elderly hypertensive patients: pharmacokinetics and pharmacodynamics. *J Cardiovasc Pharmacol* 1988;12(Suppl 7):S67-71.
2. Bernink PJLM, de Weerd P, ten Cate FJ, Remme WJ, Barth J, Enthoven R, Haagen FDM, Holwerda NJ, Klomps HC, and Coinvestigators. An 8-week double-blind study of amlodipine and diltiazem in patients with stable exertional angina pectoris. *J Cardiovasc Pharmacol* 1991;17(Suppl 1):S53-6.
3. Burris JF, Ames RP, Applegate WB, Ram CVS, Davidov ME, Mroczek WJ. Double-blind comparison of amlodipine and hydrochlorothiazide in patients with mild to moderate hypertension. *J Cardiovasc Pharmacol* 1988;12(Suppl 7):S98-102.
4. Cappuccio FP, Markandu ND, Sagnella GA, Singer DRJ, Buckley MG, Miller MA, MacGregor GA. Effects of amlodipine on urinary sodium excretion, renin-angiotensin-aldosterone system, atrial natriuretic peptide and blood pressure in essential hypertension. *J Hum Hypertens* 1991;5:115-19.
5. Flynn JT. A Randomized, Placebo-Controlled Trial of Amlodipine in Children with Hypertension *J Pediat* 2004; Vol 145/3 pp 353-359.
6. Frishman WH, Brobyn R, Brown RD, Johnson BF, Reeves RL, Wombolt DG. A randomized placebo-controlled comparison of amlodipine and atenolol in mild to moderate systemic hypertension. *J Cardiovasc Pharmacol* 1988;12(Suppl 7):S103-6.
7. Glasser SP, Chrysant SG, Graves J, Rofman B, Koehn DK. Safety and efficacy of amlodipine added to hydrochlorothiazide therapy in essential hypertension. *Am J Hypertens* 1989;2:154-7.
8. Glasser SP, West TW. Clinical safety and efficacy of once-a-day amlodipine for chronic stable angina pectoris. *Am J Cardiol* 1988;62:518-22.
9. Hosie J, Bremner AD, Fell PJ, James IG, Saul PA, Taylor SH. Comparison of early side effects with amlodipine and nifedipine retard in hypertension. *Cardiology* 1992;80(Suppl 1):54-9.
10. Klein W, Mitrovic V, Neuss H, Schlepfer M, Cocco G, Prager G, Fitscha P, Meisner W. A 6-week double-blind comparison of amlodipine and placebo in patients with stable exertional angina pectoris receiving concomitant β -blocker therapy. *J Cardiovasc Pharmacol* 1991;17(Suppl 1):S50-2.
11. Lopez LM, Thorman AD, Mehta JL. Effects of amlodipine on blood pressure, heart rate, catecholamines, lipids and responses to adrenergic stimulus. *Am J Cardiol* 1990;66:1269-71.

12. Lorimer AR, Smedsrud T, Walker P, Tyler HM. Comparison of amlodipine and verapamil in the treatment of mild to moderate hypertension. *J Cardiovasc Pharmacol* 1988;12(Suppl 7):S89-93.
13. Lund-Johansen P, Omvik P, White W, Digraanes O, Helland B, Jordal O, Stray T. Long-term haemodynamic effects of amlodipine at rest and during exercise in essential hypertension. *J Hypertens* 1990;8(12):1129-36.
14. Meredith PA, Elliott HL. Clinical pharmacokinetics of amlodipine. *Clin Pharmacokinet* 1992;22(1):22-31.
15. Murdoch D, Heel RC. Amlodipine: a review of its pharmacodynamic and pharmacokinetic properties, and therapeutic use in cardiovascular disease. *Drugs* 1991;41(3):478-505.
16. Taylor SH. Amlodipine in post-infarction angina. *Cardiology* 1992;80(Suppl 1):26-30.
17. Taylor SH, Lee P, Jackson N, Cocco G. A double-blind, placebo-controlled, parallel dose-response study of amlodipine in stable exertional angina pectoris. *J Cardiovasc Pharmacol* 1991;17(Suppl 1):S46-9.
18. Toupance O, Lavaud S, Canivet E, Bernaud C, Hotton JM, Chanard J. Antihypertensive effect of amlodipine and lack of interference with cyclosporine metabolism in renal transplant recipients. *Hypertension* 1994;24(3):297-300.
19. Treatment of Mild Hypertension Research Group. The treatment of mild hypertension study: a randomized, placebo-controlled trial of a nutritional-hygienic regimen along with various drug monotherapies. *Arch Intern Med* 1991;151:1413-23.
20. Varonne J, Investigators of Study AML-NY-86-002. The efficacy and safety of amlodipine in the treatment of mild and moderate essential hypertension in general practice. *J Cardiovasc Pharmacol* 1991;17(Suppl 1):S30-3.
21. Vetrovec G, Dailey S, Kay GN, Epstein A, Plumb V. Hemodynamic and electrophysiologic effects of amlodipine, a new long-acting calcium antagonist: preliminary observations. *Am Heart J* 1989;118(5Pt2):1104-5.
22. NORVASC™ Product Monograph (Pfizer Canada Inc.) Revision date: June 8, 2005, Control Number: 093100.

IMPORTANT: PLEASE READ

PART III: CONSUMER INFORMATION

^{Pr}**Zym-AMLODIPINE**
Amlodipine Besylate Tablets
5 mg and 10 mg

This leaflet is part III of a three-part “Product Monograph” published when Zym-AMLODIPINE was approved for sale in Canada and is designed specifically for Consumers. This leaflet is a summary and will not tell you everything about Zym-AMLODIPINE. Contact your doctor or pharmacist if you have any questions about the drug.

ABOUT THIS MEDICATION

What the medication is used for:

Zym-AMLODIPINE has been prescribed to you for the treatment of mild to moderate essential hypertension (high blood pressure) or for the management of chronic stable angina (chest pain).

What it does:

Zym-AMLODIPINE relaxes the blood vessels that carry blood away from the heart and increases the supply of blood and oxygen to the heart, while reducing its workload.

When it should not be used:

You should not use Zym-AMLODIPINE if you:

- are hypersensitive to this drug or other dihydropyridines, or to any ingredient in the formulation or component of the container (consult list of non-medicinal ingredients).
- have a low blood pressure (less than 90 mmHg systolic).

What the medicinal ingredient is:

Amlodipine Besylate

What the nonmedicinal ingredients are:

Dibasic calcium phosphate anhydrous, Magnesium stearate, Microcrystalline Cellulose and Sodium starch glycolate.

What dosage forms it comes in:

Tablets: 5 mg & 10 mg amlodipine (from amlodipine besylate)

WARNINGS AND PRECAUTIONS

BEFORE you use Zym-AMLODIPINE, talk to your doctor or pharmacist if you / the patient you are caring for:

- have/has a history of stroke, mini-strokes or high cholesterol;
- have/has a history of kidney or liver problems;
- have/has a history of problems with your heart and/or blood vessels;
- are/is pregnant or planning to become pregnant;
- are/is breast-feeding;
- are taking or planning to take any other medicines (prescription, over-the-counter medicines or herbal remedies);
- drink/drinks alcoholic beverages or use/uses drugs.

Beta-blocker withdrawal:

Zym-AMLODIPINE should be discontinued gradually by slowly reducing the dose of the drug.

Use in Children: The use of Zym-AMLODIPINE is not recommended in children less than 6 years of age.

Use in the elderly: Adverse reactions include swelling, muscle cramps and dizziness. Zym-AMLODIPINE should be used cautiously in elderly patients.

INTERACTIONS WITH THIS MEDICATION

If you are taking any medication, prescription, non-prescription and/or herbal, please inform your doctor before starting treatment with this medicine. Concomitant use of Zym-AMLODIPINE with other drugs may require surveillance and dosage adjustments.

Taking amlodipine with other drugs which are eliminated by the body through the same mechanism may result in altered bioavailability of amlodipine or these drugs. Dosages of similarly metabolized drugs, and especially in patients with renal and/or hepatic impairment, may require adjustment when starting or stopping medication.

Medications known or susceptible to interact with this medication include: azole antifungals, cimetidine, cyclosporine, erythromycin, quinidine, terfenadine,

warfarin, Phenobarbital, phenytoin, rifampin, benzodiazepines, flecainide, imipramine, propafenone and theophylline.

When beta-adrenergic receptor blocking drugs are administered concomitantly with amlodipine, patients should be carefully monitored since blood pressure lowering effect of beta-blockers may be augmented by amlodipine.

PROPER USE OF THIS MEDICATION

Take this medication exactly as prescribed by your physician. Do not stop taking your medication without having first informed your doctor.

The dose of Zym-AMLODIPINE will be different for different patients. Do not give it to anybody else, **even if their condition seems similar to yours**, as they may experience undesirable effects, which may be serious.

For both high blood pressure and chest pain, the initial dose of Zym-AMLODIPINE is 5 mg once daily. If necessary, the dose can be increased after 1-2 weeks to a maximum dose of 10 mg once daily.

Use in the Elderly or in Patients with Impaired Renal Function

The recommended initial dose in patients over 65 years of age or patients with impaired renal function is 5 mg once daily. If required, increasing in the dose should be done gradually and with caution (see **WARNING AND PRECAUTIONS**).

Use in Patients with Impaired Hepatic Function

When Zym-AMLODIPINE is used in these patients, the dosage should be carefully and gradually adjusted depending on patients tolerance and response. A lower starting dose of 2.5 mg once daily should be considered (see **WARNINGS AND PRECAUTIONS**).

Use in Children

The effective antihypertensive oral dose in pediatric patients ages 6-17 years is 2.5 mg to 5 mg once daily.

Overdose

If you take / give too much, immediately contact the doctor or go / take the patient to the nearest hospital emergency department, Do this even if there are no signs of discomfort.

One or more of the following signs may occur in an overdose: low blood pressure and rapid heartbeat.

Missed Dose

If you miss a dose, try not to miss any more. **DO NOT TAKE / GIVE TWO DOSES AT ONCE.**

SIDE EFFECTS AND WHAT TO DO ABOUT THEM

The most common side effects of Zym-AMLODIPINE are: edema (swelling), headache, fatigue, dizziness and nausea.

SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM				
Symptom / effect		Talk with your doctor or pharmacist		Stop taking drug and call your doctor or pharmacist
		Only if severe	In all cases	
Common	Edema (swelling of the extremities)	√		
	headache	√		
	fatigue	√		
	dizziness	√		
	nausea	√		
Uncommon	rapid or irregular heartbeat			√
	low blood pressure			√
	liver problems such as jaundice (yellowing of skin or eyes)			√
	symptoms of allergic reactions			√

This is not a complete list of side effects. If you have any unexpected effects while taking this drug, contact your doctor or pharmacist.

REPORTING SIDE EFFECTS

You can report any suspected adverse reactions associated with the use of health products to the Canada Vigilance Program by one of the following 3 ways:

- Report online at www.healthcanada.gc.ca/medeffect
- Call toll-free at 1-866-234-2345
- Complete a Canada Vigilance Reporting Form and:
 - Fax toll-free to 1-866-678-6789, or
 - Mail to: Canada Vigilance Program
Health Canada
Postal Locator 0701C
Ottawa, ON K1A 0K9

Postage paid labels, Canada Vigilance Reporting Form and the adverse reaction reporting guidelines are available on the MedEffect™ Canada Web site at www.healthcanada.gc.ca/medeffect.

NOTE: Should you require information related to the management of side effects, contact your health professional. The Canada Vigilance Program does not provide medical advice.

HOW TO STORE IT

- Do not use this product after the expiry date written on the package.
- Store between 15-30°C, protected from light.
- Keep this and all medicines in a safe place away from children.

MORE INFORMATION

This document plus the full product monograph, prepared for health professionals can be obtained by contacting Zymcan Pharmaceuticals Inc.:

By post: Zymcan Pharmaceuticals Inc.
6111 avenue Royalmount, #100
Montreal, QC H4P 2T4

This leaflet was prepared by
Zymcan Pharmaceuticals Inc.

Date of Preparation: April 1, 2010.