

## PRODUCT MONOGRAPH

Pr **AVA-METFORMIN**

metformin hydrochloride tablets

House standard

500 and 850 mg

Oral antihyperglycemic agent

Avanstra Inc.  
10761 – 25th NE  
Suite 110, Building “B”  
Calgary, Alberta  
T2C 3C2  
Canada

Date of Preparation:  
February 21, 2011

Submission Control No: 145186

**Table of Contents**

**PART I: HEALTH PROFESSIONAL INFORMATION ..... 3**

SUMMARY PRODUCT INFORMATION..... 3

INDICATIONS AND CLINICAL USE ..... 3

CONTRAINDICATIONS..... 3

WARNINGS AND PRECAUTIONS ..... 4

ADVERSE REACTIONS ..... 11

DRUG INTERACTIONS..... 12

DOSAGE AND ADMINISTRATION ..... 15

OVERDOSAGE..... 15

ACTION AND CLINICAL PHARMACOLOGY ..... 16

STORAGE AND STABILITY ..... 16

DOSAGE FORMS, COMPOSITION AND PACKAGING..... 17

**PART II: SCIENTIFIC INFORMATION..... 18**

PHARMACEUTICAL INFORMATION ..... 18

CLINICAL TRIALS ..... 19

DETAILED PHARMACOLOGY ..... 23

TOXICOLOGY ..... 25

REFERENCES..... 27

**PART III: CONSUMER INFORMATION..... 30**

## AVA-METFORMIN

metformin hydrochloride

### PART I: HEALTH PROFESSIONAL INFORMATION

#### SUMMARY PRODUCT INFORMATION

Route of Administration	Dosage Form / Strength	Clinically Relevant Nonmedicinal Ingredients
oral	tablet 500 mg and 850 mg	<i>None (For a complete listing see Dosage Forms, Composition and Packaging section).</i>

#### INDICATIONS AND CLINICAL USE

**AVA-METFORMIN** (metformin hydrochloride) is indicated to control hyperglycemia in **AVA-METFORMIN** (metformin hydrochloride) responsive, stable, mild, non-ketosis prone, maturity onset type of diabetes (Type II) which cannot be controlled by proper dietary management, exercise and weight reduction or when insulin therapy is not appropriate.

**AVA-METFORMIN** (metformin hydrochloride) can be of value for the treatment of obese diabetic patients.

#### CONTRAINDICATIONS

- Unstable and/or insulin-dependent (Type I) diabetes mellitus.
- Acute or chronic metabolic acidosis, including diabetic ketoacidosis, with or without coma, history of ketoacidosis with or without coma. Diabetic ketoacidosis should be treated with insulin.
- In patients with a history of lactic acidosis, irrespective of precipitating factors.
- In the presence of renal impairment or when renal function is not known, and also in patients with serum creatinine levels above the upper limit of normal range. Renal disease or renal dysfunction (e.g., as suggested by serum creatinine levels  $\geq 136 \mu\text{mol/L}$  (males),  $\geq 124 \mu\text{mol/L}$  (females) or abnormal creatinine clearance) which may result from conditions such as cardiovascular collapse (shock), acute myocardial infarction, and septicemia (see also **WARNINGS AND PRECAUTIONS**).

- Congestive heart failure requiring pharmacologic treatment.
- In excessive alcohol intake, acute or chronic.
- In patients suffering from severe hepatic dysfunction, since severe hepatic dysfunction has been associated with some cases of lactic acidosis, **AVA-METFORMIN** (metformin hydrochloride) should generally be avoided in patients with clinical or laboratory evidence of hepatic disease.
- **AVA-METFORMIN** should be temporarily discontinued in patients undergoing radiologic studies involving parenteral administration of iodinated contrast materials, because use of such products may result in acute alteration of renal function (see **WARNINGS AND PRECAUTIONS**).
- In cases of cardiovascular collapse and in disease states associate with hypoxemia such as cardiorespiratory insufficiency, which are often associated with hyperlactacidemia.
- During stress conditions, such as severe infections, trauma or surgery and the recovery phase thereafter.
- In patients suffering from severe dehydration.
- Known hypersensitivity or allergy to metformin HCl or any of the excipients. For a complete listing, see the Dosage Forms, Composition and Packaging section of the product monograph.
- During pregnancy and breastfeeding.

## **WARNINGS AND PRECAUTIONS**

### **Serious Warnings and Precautions**

- Lactic acidosis is a rare, but serious, metabolic complication that occurs due to metformin accumulation during treatment with AVA-METFORMIN (see Endocrine and Metabolism, Lactic Acidosis.)
- Patients should be cautioned against excessive alcohol intake, either acute or chronic, when taking AVA-METFORMIN, since alcohol intake potentiates the effect of metformin on lactate metabolism (see Endocrine and Metabolism, Lactic Acidosis.)

## **General**

### **Patient selection and follow-up:**

Careful selection of patients is important. It is imperative that there be rigid attention to diet and careful adjustment of dosage. Regular thorough follow-up examinations are necessary.

If vomiting occurs, withdraw drug temporarily, exclude lactic acidosis, then resume dosage cautiously (see **ADVERSE REACTIONS**).

Particular attention should be paid to short range and long range complications which are peculiar to diabetes. Periodic cardiovascular, ophthalmic, hematological, hepatic and renal assessments are advisable.

Use of metformin must be considered as treatment in addition to proper dietary regimen and not as a substitute for diet.

Care should be taken to ensure that AVA-METFORMIN is not given when a contraindication exists.

If during AVA-METFORMIN therapy the patient develops acute intercurrent disease such as: clinically significant hepatic dysfunction, cardiovascular collapse, congestive heart failure, acute myocardial infarction, or other conditions complicated by hypoxemia, the drug should be discontinued.

### **Change in Clinical Status of Previously Controlled Diabetes Patients**

A diabetic patient previously well controlled on AVA-METFORMIN who develops laboratory abnormalities or clinical illness (especially vague and poorly defined illness) should be evaluated promptly for evidence of ketoacidosis or lactic acidosis. Evaluation should include serum electrolytes and ketones, blood glucose and, if indicated, blood pH, lactate, pyruvate and metformin levels. If acidosis of either form occurs, AVA-METFORMIN must be stopped immediately and appropriate corrective measures initiated.

### **Hypoxic states**

Cardiovascular collapse (shock) from whatever cause, acute congestive heart failure, acute myocardial infarction and other conditions characterized by hypoxemia have been associated with lactic acidosis and may also cause prerenal azotemia. When such events occur in patients on AVA-METFORMIN therapy, the drug should be promptly discontinued.

## Endocrine and Metabolism

### Lactic acidosis:

Lactic acidosis is a rare, but serious, metabolic complication that occurs due to metformin accumulation during treatment with metformin; when it occurs, it is fatal in approximately 50% of cases. Lactic acidosis may also occur in association with a number of pathophysiologic conditions, including diabetes mellitus, and whenever there is significant tissue hypoperfusion and hypoxemia. Lactic acidosis is characterized by elevated blood lactate levels ( $>5$  mmol/L), decreased blood pH, electrolyte disturbances with an increased anion gap, and an increased lactate/pyruvate ratio. When metformin is implicated as the cause of lactic acidosis, metformin plasma level  $>5$   $\mu\text{g/mL}$  are generally found.

The reported incidence of lactic acidosis in patients receiving metformin is very low (approximately

0.03 cases/1000 patient/year, with approximately 0.015 fatal cases/1000 patient/year) and occurred primarily in diabetic patients with significant renal insufficiency, including both intrinsic renal disease and renal hypoperfusion, often in the setting of multiple concomitant medical/surgical problems and multiple concomitant medications. Patients with congestive heart failure requiring pharmacologic management, in particular those with unstable or acute congestive heart failure who are at risk of hypoperfusion and hypoxemia, are at increased risk of lactic acidosis. In particular, treatment of the elderly should be accompanied by careful monitoring of renal function. AVA-METFORMIN treatment should not be initiated in patients  $\geq 80$  years of age, unless measurement of creatinine clearance demonstrates that renal function is not reduced, as the patients are more susceptible to developing lactic acidosis. The risk of lactic acidosis increases with the degree of renal dysfunction and the patient's age. The risk of lactic acidosis may, therefore, be significantly decreased by regular monitoring of renal function in patients taking metformin and by use of the minimum effective dose of metformin. In addition, metformin should be promptly withheld in the presence of any condition associated with hypoxemia, dehydration or sepsis. Because impaired hepatic function may significantly limit the ability to clear lactate, metformin should generally be avoided in patients with clinical or laboratory evidence of hepatic disease. Patients should be cautioned against excessive alcohol intake, either acute or chronic, when taking metformin since alcohol intake potentiates the effect of metformin hydrochloride on lactate metabolism. In addition, metformin should be temporarily discontinued prior to any intravascular radiocontrast study and for any surgical procedure. The onset of lactic acidosis often is subtle, and accompanied only by nonspecific symptoms such as malaise, myalgias, respiratory distress, increasing somnolence and non-specific abdominal distress. There may be associated hypothermia, hypotension and resistance bradyarrhythmias with more marked acidosis. The patient and the patient's physician must be aware of the possible importance of such symptoms and the patient should be instructed to notify the physician immediately if they occur.

Metformin should be withdrawn until the situation is clarified. Serum electrolytes, ketones, blood glucose and, if indicated, blood pH, lactate levels and even blood metformin levels may be useful. Once a patient is stabilized on any dose level of metformin, gastrointestinal symptoms, which are common during initiation of therapy, are unlikely to be drug related. Later occurrence of gastrointestinal symptoms could be due to lactic acidosis or other serious disease. Levels of fasting venous plasma lactate above the upper limit of normal but less than 5 mmol/L in patients taking metformin do not necessarily indicate impending lactic acidosis and may be explainable by other mechanisms, such as poorly controlled diabetes or obesity, vigorous physical activity or technical problems in sample handling. Lactic acidosis should be suspected in any diabetic patient with metabolic acidosis lacking evidence of ketoacidosis (ketonuria and ketonemia).

Lactic acidosis is a medical emergency that must be treated in a hospital setting. In a patient with lactic acidosis who is taking metformin, the drug should be discontinued immediately and general supportive measures promptly instituted. Because metformin hydrochloride is dialysable (with clearance of up to 170 mL/min under good hemodynamic conditions), prompt hemodialysis is recommended to correct the acidosis and remove the accumulated metformin. Such management often results in prompt reversal of symptoms and recovery (see **CONTRAINDICATIONS**).

**NOTE:** When used as indicated, there has not been a single case of lactic acidosis in Canada. Metformin should be immediately discontinued in the presence of acidosis.

Physicians should instruct their patients to recognize the symptoms which could be signal onset of lactic acidosis. If acidosis of any kind develops, metformin should be discontinued immediately.

### **Loss of control of blood glucose**

When a patient stabilized on any diabetic regimen is exposed to stress such as fever, trauma, infection, or surgery, a temporary loss of glycemic control may occur. At such times, it may be necessary to withhold metformin and temporarily administer insulin. Metformin may be reinstated after the acute episode is resolved.

The effectiveness of oral antidiabetic drugs in lowering blood glucose to a targeted level decreases in many patients over a period of time. This phenomenon, which may be due to progression of the underlying disease or to diminished responsiveness to the drug, is known as secondary failure, to distinguish it from primary failure in which the drug is ineffective during initial therapy.

Should secondary failure occur with metformin, therapeutic alternatives should be considered.

### **Vitamin B12 levels**

Impairment of vitamin B<sub>12</sub> absorption has been reported in some patients. Therefore, measurements of serum vitamin B<sub>12</sub> are advisable at least every one to two years in patients on long-term treatment with metformin.

A decrease to subnormal levels of previously normal serum Vitamin B<sub>12</sub> levels, without clinical manifestations, is observed in approximately 7% of patients receiving metformin in controlled clinical trials of 28 weeks duration. Such decrease, possibly due to interference with B<sub>12</sub> absorption from B<sub>12</sub>-intrinsic factor complex is, however, very rarely associated with anemia and appears to be rapidly reversible with discontinuation of metformin or vitamin B<sub>12</sub> supplementation. Measurement of hematologic parameters on an annual basis is advised in patients on metformin and any apparent abnormalities should be appropriately investigated and managed (see **MONITORING and LABORATORY TESTS**). Certain individuals (those with inadequate vitamin B<sub>12</sub> or calcium intake or absorption) appear to be predisposed to developing subnormal vitamin B<sub>12</sub> levels.

### **Hepatic/Biliary/Pancreatic**

Since impaired hepatic function has been associated with some cases of lactic acidosis, metformin should generally be avoided in patients with clinical or laboratory evidence of hepatic disease.

### **Hypoglycemia**

Hypoglycemia does not occur in patients receiving metformin alone under usual circumstances of use, but could occur when caloric intake is deficient, when strenuous exercise is not compensated by caloric supplementation, or during concomitant use with other glucose lowering agents or ethanol.

Elderly, debilitated or malnourished patients, and those with adrenal or pituitary insufficiency or alcohol intoxication are particularly susceptible to hypoglycemic effects. Hypoglycemia may be difficult to recognize in the elderly, and in people who are taking beta-adrenergic blocking drugs.

The patients should be warned about driving a vehicle or operating machinery under these conditions where risk of hypoglycaemia is present.

### **Peri-Operative Considerations**

Metformin therapy should be temporarily suspended for any surgical procedure (except minor procedures not associated with restricted intake of food and fluids). Metformin should be discontinued 2 days before surgical intervention and should not be restarted until the patient's oral intake has resumed and renal function has been evaluated as normal.

### **Renal**



Metformin is known to be substantially excreted by the kidney, and the risk of metformin accumulation and lactic acidosis increases with the degree of impairment of renal function. Thus, patients with serum creatinine levels above the upper limit of the normal range for their age should not receive metformin. In patients with advanced age, metformin should be carefully titrated to establish the minimum dose for adequate glycemic effect, because aging is associated with reduced renal function. In elderly patients, renal function should be monitored regularly and generally, metformin should not be titrated to the maximum dose (see **DOSAGE AND ADMINISTRATION**).

Before initiation of metformin therapy and every 6 months while on metformin therapy, renal function should be assessed and verified as being within normal range.

In patients in whom development of renal dysfunction is anticipated, renal function should be assessed more frequently and metformin discontinued if evidence of renal impairment is present.

Special caution should be exercised in situations where renal function may become impaired, for example when initiating antihypertensive therapy or diuretic therapy and when starting therapy with an NSAID.

### **Use of Concomitant Medications That May Affect Renal Function or Metformin Disposition**

Concomitant medication(s) that may affect renal function or result in significant hemodynamic change or may interfere with disposition of metformin, such as cationic drugs that are eliminated by renal tubular secretion (see **DRUG INTERACTIONS**), should be used with caution.

Radiologic studies involving the use of intravascular iodinated contrast materials (for example, intravenous urogram, intravenous cholangiography, angiography and computed tomography (CT) scans with intravascular contrast materials.

Intravascular contrast studies with iodinated materials can lead to acute renal failure and have been associated with lactic acidosis in patients receiving metformin (see **CONTRAINDICATIONS**). Therefore, in patients in whom any such study is planned, metformin should be discontinued at the time of or prior to the procedure, and withheld for 48 hours subsequent to the procedure and reinstated only after renal function has been re-evaluated and found to be normal.

### **Special Populations**

#### **Pregnant Women**

Safety in pregnant women has not been established. Metformin was not teratogenic in rats and

rabbits at doses up to 600 mg/kg/day, or about two times the maximum recommended human daily dose on a body surface area basis. Determination of fetal concentrations demonstrated a partial placental barrier to metformin. Because animal reproduction studies are not always predictive of human response, the use of AVA-METFORMIN is not recommended during pregnancy.

Because recent information suggests that abnormal blood glucose levels during pregnancy are associated with a higher incidence of congenital abnormalities, there is a consensus among experts that insulin be used during pregnancy to maintain blood glucose levels as close to normal as possible.

### **Nursing Women:**

Studies in lactating rats show that metformin is excreted into milk and reaches levels comparable to those in plasma. Similar studies have not been conducted in nursing mothers, but caution should be exercised in such patients, and a decision should be made whether to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother.

### **Pediatrics**

Safety and effectiveness in pediatric patients have not been established.

### **Geriatrics**

Controlled clinical studies of metformin did not include sufficient numbers of elderly patients to determine whether they respond differently from younger patients, although other reported clinical experience has not identified differences in responses between the elderly and younger patients. Metformin is known to be substantially excreted by the kidney and because the risk of serious adverse reactions to the drug is greater in patients with impaired renal function, it should only be used in patients with normal renal function (see **CONTRAINDICATIONS and WARNINGS and PRECAUTIONS**). Because aging is associated with reduced renal function, metformin should be used with caution as age increases. Care should be taken in dose selection and should be based on careful and regular monitoring of renal function. Generally, elderly patients should not be titrated to the maximum dose of metformin.

### **Monitoring and Laboratory Tests**

Response to all diabetic therapies should be monitored by periodic measurements of fasting blood glucose and glycosylated hemoglobin levels, with a goal of decreasing these levels toward the normal range. During initial dose titration, fasting glucose can be used to determine therapeutic response. Thereafter, both glucose and glycosylated hemoglobin should be monitored. Measurements of glycosylated hemoglobin may be especially useful for evaluating

long-term control (see **DOSAGE AND ADMINISTRATION**).

Initial and periodic monitoring of hematologic parameters (e.g., hemoglobin/hematocrit and red blood cell indices) and renal function (serum creatinine) should be performed, at least on an annual basis. While megaloblastic anemia has rarely been seen with metformin therapy, if this is suspected, vitamin B<sub>12</sub> deficiency should be excluded.

## **ADVERSE REACTIONS**

### **Adverse Drug Reaction Overview**

The adverse events most commonly associated with AVA-METFORMIN (metformin Hcl) are diarrhea, nausea, and upset stomach. Lactic acidosis is a rare, but serious side effect. Lactic acidosis is fatal in approximately 50% of cases.

#### **Lactic Acidosis**

Very rare (< 1/10 000 and isolated reports). See **WARNINGS, PRECAUTIONS, and OVERDOSAGE** sections.

#### **Gastrointestinal Reactions**

Very common: (>1/10). Gastrointestinal symptoms (diarrhea, nausea, vomiting, abdominal bloating, flatulence, and anorexia) are the most common reactions to metformin and are approximately 30% more frequent in patients on metformin monotherapy than in placebo-treated patients, particularly during initiation of metformin therapy. These symptoms are generally transient and resolve spontaneously during continued treatment. Occasionally, temporary dose reduction may be useful.

Because gastrointestinal symptoms during therapy initiation appear to be dose-related, they may be decreased by gradual dose escalation and by having patients take metformin with meals (see **DOSAGE AND ADMINISTRATION**).

Because significant diarrhea and/or vomiting can cause dehydration and prerenal azotemia, metformin should be temporarily discontinued, under such circumstances.

For patients who have been stabilized on metformin, nonspecific gastrointestinal symptoms should not be attributed to therapy unless intercurrent illness or lactic acidosis have been excluded.

## **Special senses**

Common: ( $\geq 1/100$ ). During initiation of metformin therapy, complaints of taste disturbance are common, i.e. metallic taste.

## **Dermatologic reactions**

Very rare: ( $< 1/10\ 000$  and isolated reports). The incidence of rash/dermatitis in controlled clinical trials was comparable to placebo for metformin monotherapy and to sulfonylurea for metformin / sulfonylurea therapy. Reports of skin reactions such as erythema, pruritus, and urticaria are very rare.

## **Hematologic**

During controlled clinical trials of 29 weeks duration, approximately 9% of patients on metformin monotherapy and 6% of patients on metformin / sulfonylurea therapy developed asymptomatic subnormal serum vitamin B<sub>12</sub> levels; serum folic acid levels did not decrease significantly. However, only five cases of megaloblastic anemia have been reported with metformin administration (none during U.S. clinical studies) and no increased incidence of neuropathy has been observed. (see also **WARNINGS AND PRECAUTIONS**).

Decrease of vitamin B<sub>12</sub> absorption with decrease of serum levels during long-term use of metformin is rare ( $\geq 1/10\ 000$  and  $< 1/1\ 000$ ). Consideration of such aetiology is recommended if a patient presents with megaloblastic anemia.

## **Hepatic**

Very rare: ( $< 1/10\ 000$  and isolated reports): Liver function tests abnormalities or hepatitis resolving upon metformin discontinuation has been documented in isolated reports.

## **DRUG INTERACTIONS**

### **Overview**

Certain drugs may potentiate the effect of metformin, particularly sulfonylurea type of drugs in the treatment of diabetes. The simultaneous administration of these two types of drugs could produce a hypoglycemic reaction, especially if they are given in patients already receiving other drugs which, themselves, can potentiate the effect of sulfonylureas. These drugs can be: long-acting sulfonamides, tuberculostatics, phenylbutazone, clofibrate, monoamine oxidase inhibitors, salicylates, probenecid and propranolol.

In healthy volunteers, the pharmacokinetics of propranolol and ibuprofen were not affected by

metformin when co-administered in single-dose interaction studies.

Metformin is negligibly bound to plasma proteins and is, therefore, less likely to interact with highly protein-bound drugs such as salicylates, sulfonamides, chloramphenicol, and probenecid, as compared to sulfonylureas, which are extensively bound to serum proteins.

## **Drug-Drug Interactions**

### Glyburide:

In a single-dose interaction study in NIDDM subjects, co-administration of metformin and glyburide did not result in any changes in either metformin pharmacokinetics or pharmacodynamics. Decreases in glyburide AUC and  $C_{max}$  were observed, but were highly variable. The single-dose nature of this study and the lack of correlation between glyburide blood levels and pharmacodynamic effects, makes the clinical significance of this interaction uncertain.

### Furosemide:

A single-dose study, metformin-furosemide drug interaction study in healthy subjects demonstrated that pharmacokinetic parameters of both compounds were affected by co-administration. Furosemide increased the metformin plasma and blood  $C_{max}$  by 22% and blood AUC by 15%, without any significant change in metformin renal clearance. When administered with metformin, the  $C_{max}$  and AUC of furosemide were 31% and 12% smaller, respectively, than when administered alone, and the terminal half-life was decreased by 32%, without any significant change in furosemide renal clearance. No information is available about the interaction of metformin and furosemide when co-administered chronically.

### Nifedipine:

A single-dose, metformin-nifedipine drug interaction study in healthy volunteers demonstrated that co-administration of nifedipine increased plasma metformin  $C_{max}$  and AUC by 20% and 9%, respectively, and increased the amount excreted in the urine.  $T_{max}$  and half-life were unaffected. Nifedipine appears to enhance the absorption of metformin. Metformin had minimal effects on nifedipine.

### Cationic drugs:

Cationic drugs (e.g., amiloride, digoxin, morphine, procainamide, quinidine, quinine, ranitidine, triamterene, trimethoprim, and vancomycin) that are eliminated by renal tubular secretion, theoretically have the potential for interaction with metformin by competing for common renal tubular transport systems. Such an interaction has been observed between metformin and oral cimetidine in normal healthy volunteers in both single and multiple-dose, metformin-cimetidine

drug interaction studies, with a 60% increase in peak metformin plasma and whole blood concentrations and a 40% increase in plasma and whole blood metformin AUC was observed. There was no change in elimination half-life in the single-dose study. Metformin had no effect on cimetidine pharmacokinetics. Therefore, careful patient monitoring and dose adjustment of metformin or the interfering drug is recommended in patients who are taking cationic medications that are excreted via renal tubular secretion.

Other:

Other drugs tend to produce hyperglycemia and may lead to a loss of blood sugar control. These include thiazide and other diuretics, corticosteroids, phenothiazines, thyroid products, estrogens, estrogen plus progestogen, oral contraceptives, phenytoin, nicotinic acid, sympathomimetics, calcium channel blocking drugs, isoniazid and beta-2-agonists. *ACE-inhibitors* may decrease the blood glucose levels. When such drugs are administered to patients receiving metformin, the patient should be closely observed to maintain adequate glycemic control.

Elimination rate of the anticoagulant phenprocoumon has been reported to be increased by 20% when used concurrently with metformin. Therefore, patients receiving phenprocoumon or other antivitamin K anticoagulants should be monitored carefully when both types of drugs are used simultaneously. In such cases, an important increase of prothrombin time may occur upon cessation of metformin therapy, with an increased risk of hemorrhage.

**Drug-Food Interactions**

Interactions with food have not been established.

**Drug-Herb Interactions**

Interactions with herbal products have not been established.

**Drug-Laboratory Test Interactions**

Intravascular contrast studies with iodinated materials can lead to acute alteration of renal function and have been associated with lactic acidosis in patients receiving metformin. (See **CONTRAINDICATIONS and WARNINGS AND PRECAUTIONS.**)

**Drug-Lifestyle Interactions**

Patients should be cautioned against excessive alcohol intake, either acute or chronic, when taking metformin, since alcohol intake potentiates the effect of metformin on lactate metabolism (see **CONTRAINDICATIONS**).

## **DOSAGE AND ADMINISTRATION**

### **Dosing Considerations**

In diabetic patients, individual determination of the minimum dose that will lower blood glucose adequately should be made, aiming for glycemic targets as close to normal as possible. A lower recommended starting dose and gradually increased dosage is advised to minimize gastrointestinal symptoms.

Over a period of time, patients may become progressively less responsive to therapy with oral hypoglycemic agents because of deterioration of their diabetic state. Patients should therefore be monitored with regular clinical and laboratory evaluations, including blood glucose and glycosylated hemoglobin (A<sub>1c</sub>) determinations, to determine the minimum effective dosage and to detect primary failure or secondary failure (see WARNINGS and PRECAUTIONS).

In patients in whom the maximum dose fails to lower the blood glucose adequately, therapeutic alternatives should be considered.

The usual dose is 500 mg three to four times a day or 850 mg two or three times a day. Maximal dose should not exceed 2.55 g a day. To minimize gastric intolerance such as nausea and vomiting, AVA-METFORMIN should be taken with food whenever possible.

### **Transfer from Other Antidiabetic Therapy**

When transferring patients from standard oral hypoglycaemic agents, other than chlorpropamide, to metformin, no transition period generally is necessary. When transferring patients from chlorpropamide, care should be exercised during the first two weeks because of the prolonged retention of chlorpropamide in the body, leading to overlapping drug effects and possible hypoglycaemia.

#### Missed Dose

In case the patient forgets to take AVA-METFORMIN tablets, he/she should wait for the next dose at the usual time. He/she should not double the dose to make up the forgotten dose.

## **OVERDOSAGE**

Available information concerning treatment of a massive overdose of metformin is very limited. It would be expected that adverse reactions of a more intense character including epigastric discomfort, nausea and vomiting followed by diarrhea, drowsiness, weakness, dizziness, malaise and headache might be seen. Should those symptoms persist, lactic acidosis

should be excluded. The drug should be discontinued and proper supportive therapy instituted.

Overdose of Metformin hydrochloride has occurred, including ingestion of amounts greater than 50 grams. Hypoglycemia was reported in approximately 10% of cases, but no causal association with metformin hydrochloride has been established. Lactic acidosis has been reported in approximately 32% of metformin overdose cases (see **WARNINGS AND PRECAUTIONS**). Metformin is dialysable with a clearance of up to 170 mL/min under good hemodynamic conditions. Therefore, hemodialysis may be useful for removal of accumulated drug from patients in whom metformin overdosage is suspected.

## **ACTION AND CLINICAL PHARMACOLOGY**

### **Mechanism of Action**

Metformin hydrochloride is a biguanide derivative producing an antihyperglycemic effect which can only be observed in man or in the diabetic animal and only when there is insulin secretion. Metformin, at therapeutic doses, does not cause hypoglycemia when used alone in man or in the non-diabetic animal, except when using a near lethal dose. Metformin has no effects on the pancreatic beta cells.

The mode of action of metformin is not fully understood. It has been postulated that metformin might potentiate the effect of insulin or that it might enhance the effect of insulin on the peripheral receptor site. This increased sensitivity seems to follow an increase in the number of insulin receptors on cell surface membranes.

Metformin absorption is relatively slow and may extend over about 6 hours. The drug is excreted in urine at a high renal clearance rate of about 450 mL/min. The initial elimination of metformin is rapid with a half-life varying between 1.7 and 3 hours. The terminal elimination phase accounting for about 4 to 5 % of the absorbed dose is slow with a half-life between 9 and 17 hours. Metformin is not metabolized. Its main sites of concentration are the intestinal mucosa and the salivary glands. The plasma concentration at steady-state ranges about 1 to 2 mcg/mL. Certain drugs may potentiate the effects of metformin (see **WARNINGS AND PRECAUTIONS and DRUG INTERACTIONS**).

## **STORAGE AND STABILITY**

Store in a dry place between 15-30°C in well closed containers. Protect from light.



## DOSAGE FORMS, COMPOSITION AND PACKAGING

**AVA-METFORMIN 500 mg** tablets are white, round, biconvex tablets, film-coated tablets, partial scored line on one face and rph M12 on the other. Bottles of 100 and 500 tablets.

**AVA-METFORMIN 850 mg** tablets are white, oval, film-coated biconvex tablets, with rph M11 on one side and plain on the other side. Bottles of 100 tablets.

### **Composition:**

**AVA-METFORMIN 500 mg tablet contains:** 500 mg metformin hydrochloride. **Non-medicinal ingredients in alphabetical order:** Crospovidone, hydroxypropyl methylcellulose, magnesium stearate, maltodextrin, pregelatinised maize starch, polyethylene glycol and povidone.

**AVA-METFORMIN 850 mg tablet contains:** 850 mg metformin hydrochloride. **Non-medicinal ingredients in alphabetical order:** Crospovidone, hydroxypropyl methylcellulose, magnesium stearate, maltodextrin, pregelatinised maize starch, polyethylene glycol and povidone.

## PART II: SCIENTIFIC INFORMATION

### PHARMACEUTICAL INFORMATION


#### Drug Substance

Proper name: Metformin hydrochloride

Chemical name: 1,1-dimethyl biguanide hydrochloride

Molecular formula and molecular mass:  $C_4H_{11}N_5.HCl$  and 165.63

Structural formula:



Physicochemical properties: Metformin hydrochloride is a white crystalline powder. Metformin hydrochloride is soluble in water and in 95% ethyl alcohol. It is practically insoluble in ether and in chloroform.

Melting point: 222-226°C

pKa: 2.8 and 11.5 (32°C)

pH: Between 6 and 7 (1 g / 40 mL of H<sub>2</sub>O at 25°C)

### CLINICAL TRIALS

The data which formed the basis of approval for the original metformin submission are not available. Rather, this section presents data from a published study which investigated the safety and efficacy of metformin.

The prospective randomised (UKPDS) study has established the long-term benefit of intensive blood glucose control in adult patients with type 2 diabetes. Analysis of the results for overweight patients treated with metformin after failure of diet alone showed:

- a significant reduction of the absolute risk of any diabetes-related complication in the metformin group (29.8 events/1000 patient-years) versus diet alone (43.3 events/1000 patient-years),  $p=0.0023$ , and versus the combined sulfonylurea and insulin monotherapy groups (40.1 events/1000 patient-years),  $p=0.0034$ ;
- a significant reduction of the absolute risk of diabetes-related mortality: metformin 7.5 events/1000 patient-years, diet alone 12.7 events/1000 patient-years,  $p=0.017$ . There was no significant difference between the metformin group and those assigned intensive therapy with sulfonylurea or insulin.
- a significant reduction of the absolute risk of overall mortality: metformin 13.5 events/1000 patient-years versus diet alone 20.6 events/1000 patient-years, ( $p=0.011$ ), and versus the combined sulfonylurea and insulin monotherapy groups 18.9 events/1000 patient-years ( $p=0.021$ ).
- a significant reduction in the absolute risk of myocardial infarction: metformin 11 events/1000 patient-years, diet alone 18 events/1000 patient-years, ( $p=0.01$ ). There was no significant difference between the metformin group and those assigned intensive therapy with sulfonylurea or insulin.
- There were no significant difference between the metformin group and the diet alone in the other aggregate endpoints (stroke, peripheral vascular disease and microvascular complications).

**Comparative Bioavailability Studies (using the originally marketed formulation of AVA-METFORMIN)**

The originally marketed formulation of **AVA-METFORMIN** was used to conduct two randomized, two-way crossover, single-dose bioavailability studies in fasting (table 1) and fed (table 2), healthy, adult male subjects. The bioavailability of **AVA-METFORMIN** Tablets, 500 mg, relative to Glucophage® Tablets, 500 mg was determined following a single 1 x 500 mg dose. The average values of the pharmacokinetic parameters as well as ratio of means (with 90% confidence intervals) are listed in the following tables:

**Table 1:**

Summary Table of the Comparative Bioavailability Study of <b>AVA-METFORMIN</b> vs Glucophage® (Metformin Hydrochloride) 500 mg tablets conducted under fasting conditions in 18 healthy adult male volunteers (from measured data)			
Parameter	Geometric Mean Arithmetic Mean (CV%)		Ratio of Geometric Means (%) (90% Confidence Limits)
	<b>AVA- METFORMIN</b>	Glucophage®**	
AUC <sub>t</sub> (ng.h/mL)	6168.95 6391.98 (28.0)	6341.16 6469.09 (19.7)	97 (90-105)
AUC <sub>inf</sub> (ng.h/mL)	6327.60 6546.47 (27.5)	6417.17 6534.24 (18.9)	95 (89-102)
C <sub>max</sub> (ng/mL)	995.45 1017.06 (22.7)	1045.43 1068.75 (21.9)	95 (88-103)
T <sub>max</sub> * (h)	2.92 (39.1)	2.61 (38.6)	---
T <sub>1/2</sub> * (h)	2.53 (13.8)	2.50 (14.4)	---

\*The T<sub>max</sub> and T<sub>1/2</sub> parameters are expressed as the arithmetic means.

\*\*Glucophage® is manufactured by Hoechst Marion Roussel and was purchased in Canada.

**Table 2:**

Summary Table of the Comparative Bioavailability Study of AVA-METFORMIN vs Glucophage® (Metformin Hydrochloride) 500 mg tablets conducted under fed conditions in 17 healthy adult male volunteers (from measured data)			
Parameter	Geometric Mean Arithmetic Mean (CV%)		Ratio of Geometric Means (%) (90% Confidence Limits)
	AVA- METFORMIN	Glucophage®**	
AUC <sub>t</sub> (ng.h/mL)	5949.82 6089.05 (23.2)	6116.07 6251.65 (22.0)	97 (92-103)
AUC <sub>inf</sub> (ng.h/mL)	6220.58 6370.00 (23.3)	6341.49 6482.56 (21.9)	98 (93-104)
C <sub>max</sub> (ng/mL)	898.66 927.02 (26.2)	906.14 932.09 (24.5)	99 (94-104)
T <sub>max</sub> * (h)	4.24 (21.3)	4.35 (29.8)	---
T <sub>1/2</sub> * (h)	2.96 (20.3)	2.87 (18.4)	---

\*The  $T_{max}$  and  $T_{1/2}$  parameters are expressed as the arithmetic means.

\*\*Glucophage® is manufactured by Hoechst Marion Roussel and was purchased in Canada.

**Conclusion:** The 90% confidence intervals for the ln-transformed parameters AUC<sub>t</sub>, AUC<sub>inf</sub> and C<sub>max</sub> for metformin were within the 80-125% TPD acceptance range both before and after correction for measured content. Based on these results, **AVA-METFORMIN** and Glucophage® (metformin hydrochloride) 500 mg tablets are considered bioequivalent under both single-dose fasting and fed conditions.

### Comparative Bioavailability Studies (using a new formulation of AVA-METFORMIN)

A new formulation of **AVA-METFORMIN** (metformin hydrochloride) 500 mg tablets was used

to conduct two single dose crossover, comparative bioavailability studies versus Glucophage® (metformin hydrochloride) 500 mg tablets manufactured by Aventis Pharma Inc., Canada (currently Sanofi-Aventis Canada Inc.) in healthy, adult male subjects under fasting (aged 18-55 years; N=22; Table 3) and fed (aged 21-52 years; N=21; Table 4) conditions. The results for both studies are presented below:

**Table 3: Summary Table of the Comparative Bioavailability Data - Fasting**

Metformin (1 x 500 mg) From measured data uncorrected for potency			
Parameter	Geometric Mean Arithmetic Mean (CV%)		
	AVA- METFORMIN 500 mg tablets	Glucophage®† 500 mg tablets	(%) Ratio of Geometric Means (90% Confidence Interval)
AUC <sub>t</sub> (ng.h/mL)	5790.9 5965.8 (26.2)	5591.3 5821.5 (27.9)	103.57 (96.61-111.02)
AUC <sub>inf</sub> (ng.h/mL)	5994.2 6169.5 (25.6)	5781.7 6010.9 (27.4)	103.68 (96.91-110.91)
C <sub>max</sub> (ng/mL)	1057.0 1091.5 (27.7)	1002.5 1052.0 (32.2)	105.44 (96.63-115.04)
T <sub>max</sub> * (h)	2.00 (1.00-3.50)	2.50 (1.00-4.00)	---
T <sub>1/2</sub> ** (h)	3.18 (17.0)	2.98 (12.0)	---

\* T<sub>max</sub> expressed as the median (range) only

\*\* T<sub>1/2</sub> expressed as the arithmetic mean (CV%) only

† Glucophage® 500 mg tablets manufactured by Aventis Pharma Inc. (currently known as Sanofi-Aventis Canada Inc.) were purchased in Canada

**Table 4: Summary Table of the Comparative Bioavailability Data - Fed**

Metformin (1 x 500 mg) From measured data uncorrected for potency			
Parameter	Geometric Mean Arithmetic Mean (CV%)		
	AVA- METFORMIN 500 mg tablets	Glucophage®† 500 mg tablets	(%) Ratio of Geometric Means (90% Confidence Interval)
AUC <sub>t</sub> (ng.h/mL)	5367.1 5530.8 (27.0)	5424.3 5554.3 (23.2)	98.95 (94.81-103.26)
AUC <sub>inf</sub> (ng.h/mL)	5538.2 5703.6 (26.8)	5600.5 5729.6 (22.8)	98.89 (94.86-103.09)
C <sub>max</sub> (ng/mL)	776.8 800.7 (27.2)	786.8 809.0 (25.4)	98.72 (94.21-103.45)
T <sub>max</sub> * (h)	3.50 (1.00-4.50)	3.50 (1.50-5.00)	---
T <sub>1/2</sub> ** (h)	3.64 (18.8)	3.63 (24.6)	---

\* T<sub>max</sub> expressed as the median (range) only

\*\* T<sub>1/2</sub> expressed as the arithmetic mean (CV%) only

† Glucophage® 500 mg tablets manufactured by Aventis Pharma Inc. (currently known as Sanofi-Aventis Canada Inc.) were purchased in Canada

**Conclusion:** The 90% confidence intervals of the relative geometric mean of the Test to the Reference formulation for AUC<sub>t</sub>, was within the acceptance range of 80-125% for both the measured and potency-corrected data. Based on these results, **AVA-METFORMIN** and Glucophage® (metformin hydrochloride) 500 mg tablets are considered bioequivalent under both single-dose fasting and fed conditions with a good tolerance of the subjects.

## DETAILED PHARMACOLOGY

Metformin absorption is relatively slow and may extend over about 6 hours.

Animal studies with metformin, labeled with  $^{14}\text{C}$  have shown that the drug is neither concentrated by liver cells nor is it excreted in the bile; it is concentrated in the intestinal mucosa and salivary glands.

It has been shown that, following a 2 g dose of metformin, the blood level remains under 10 mcg/mL even at the peak, occurring 2 hours after absorption. During the experiments, metformin was shown to be devoid of any notable action in the body, apart from its specific metabolic activity.

In the healthy animal, metformin lowers blood sugar only at a nearly lethal dose. Different animal species are of unequal sensitivity. On the other hand, the animal with experimental diabetes, is sensitive to a much lower dosage, providing some insulin is still secreted.

The antihyperglycemic action of metformin is probably mediated through insulin:

Metformin improves the K co-efficient of glucose assimilation.

Metformin improves the co-efficient of insulin efficiency.

In the obese diabetic with hyperinsulinemia, metformin is reported to normalize insulin output. This normalizing effect is concurrent to that of glycemias.

Metformin has little effect on liver glycogen of the healthy animal. In low and average doses, no change occurs. In high doses nearing lethal levels, liver glycogen decreases. This lowering precedes the fall in blood sugar. This reaction represents a defense mechanism tending to mobilize body reserves in order to combat hypoglycemia.

In the diabetic animal with a low liver glycogen reserve, the opposite occurs and metformin builds up glycogen stores of the liver. *In vitro*, on muscular tissue isolated in Warburg's apparatus, metformin increases glucose uptake by the muscle. This action follows an aerobic pathway. Even in high concentration, contrary to phenethyl-biguanide, metformin apparently does not block respiration or change carbohydrate metabolism via the anaerobic pathway.

Metformin is eliminated in feces and urine. It is rapidly excreted by the kidneys in an unchanged form.

Renal clearance is 450 mL/minute; this appears to explain the absence of accumulation.

Metabolites of metformin have not been identified, neither by radio-active nor by chemical



methods. A single Rf spot is always present following radiochromatographic study of urine and always corresponds to that of pure metformin. Administration during 10 consecutive days has not shown any sign of accumulation.

Inhibition of glycogenesis has been observed in animals following its stimulation by fasting, cortisol, alcohol or other substrates such as alanine lactate or pyruvate. However, such an effect varies according to the type and dosage of the biguanide used, nutritional state of the animal species and design of experimental model.

This inhibition of glycogenesis is observed only in the presence of insulin and it does not appear to play an important role in man.

Inhibition of intestinal absorption of sugars, which is not related to a malabsorption phenomenon has been observed with biguanides under certain experimental conditions in animal and in man. In one study, a 20 % retardation of galactose absorption was observed in man receiving metformin. However, such an effect of metformin could not be confirmed in another study in man.

Recent findings appear to indicate that most of the metabolic effects of the biguanides are exerted through a single mechanism, namely inhibition of fatty acid oxidation and of acetyl-CoA generation.

However, inhibition of insulin-stimulated lipogenesis which has also been observed appears to be due to the inhibition of acetyl-CoA carboxylase by the biguanides. Such an effect may explain, at least partly, the weight-reducing effect exerted by these drugs in obese diabetic patients.

## TOXICOLOGY

### 1. Animal toxicity

<b>Acute toxicity (LD<sub>50</sub>)</b>	<b>Subcutaneously</b>	<b>Orally</b>
<b>Animal</b>	<b>(mg/kg)</b>	<b>(mg/kg)</b>
Mouse	225	3500
Chicken	150	
Rat	300	1000
Rabbit	150	350
Guinea Pig	150	500

#### Chronic toxicity:

- A) The following doses of metformin produced no organ toxicity:
- |         |           |                            |
|---------|-----------|----------------------------|
| Rats    | 125 mg/kg | per os for one year        |
| Rabbits | 100 mg/kg | per os for one year        |
| Dogs    | 50 mg/kg  | subcutaneously for 2 years |

Acute or chronic organ toxicity was not produced in the animal species involved.

- B) A study was carried out during 9 months with 80 rats, male and female, divided in 4 groups, with the following dosage regimen:
- |           |  |
|-----------|--|
| 1st Group | Control  |
| 2nd Group | 150 mg/kg per os   |
| 3rd Group | 300 mg/kg per os   |
| 4th Group | 300 mg/kg per os, dose increased by 100 mg/kg/day every 15 days. |

In summary, the authors report the excellent tolerance of metformin by rats, even when administered in very high doses. No drug related lesion has been observed.

- C) Chronic toxicity studies of 9 months duration were carried through with 16 beagle dogs, although the complete intolerance of this animal species to oral hypoglycemic agents is a well established fact. Trophic and neurologic disorders with cachexia rapidly lead to the dog's death. During the periods of metformin administration, laboratory findings were within normal limits. The levels of enzymes were somewhat elevated, but it is difficult to ascribe a pathological significance to their values, since subjects in the control group were at the same level as treated animals.

Pathological studies show an extreme degree of undernutrition in all metformin treated animals.

Profound wasting especially marked in fat tissues was evident in all organs. Cachexia appears as the common cause of death of these animals.

## **2. Human toxicity**

In man, no adverse effect has been reported on liver or kidney function, the hematopoietic system or on the blood vessels.

The reported incidence of lactic acidosis in patients receiving metformin is very low (approximately 0.03 cases / 1000 patient/years with approximately 0.015 fatal cases / 1000 patients/year).

The consecutive administration of both phenformin and metformin to the same patient has allowed for the demonstration of a fundamental difference between these two biguanides in relation to lactacidemia. In some instances, patients developed hyperlactacidemia with phenformin when the same patients were presenting normal lactic acid levels while being treated with metformin. In other instances, hyperlactacidemia observed during a treatment with phenformin did regress when metformin was substituted for phenformin. Metformin may increase lactacidemia but to a degree that is clinically less significant than the elevation seen after phenformin.

## **3. Teratology**

Teratological studies were carried out in albino rats divided in three groups: No abnormalities were found, even when high doses were administered. The number of animals was the same in each group.

Death rate in the three groups of treated animals and controls was approximately the same. However, the number of living animals in each group treated was slightly lower than in the control group. Also, the frequency of litters exceeding 10 live animals was slightly higher in the control group. A loss of weight at the time of weaning has been observed when compared to the control group.

Nevertheless, on a statistical basis, differences were shown to be non-significant. There is no difference between the groups of treated animals and the control group regarding the number of stillborn. The number of deaths after birth was slightly higher in metformin treated groups than in the control group, but the comparison of average death rates is not significant ( $p < 0.05$ ).

## REFERENCES

1. Alberti, K.G.M. *et al*: Lactic Acidosis, *The Lancet*, 1977;July 2:25-29.
2. Beckmann, R.: Résorption, distribution dans l'organisme et élimination de la Metformine. *Diabétologia*, 1969;5:318-324.
3. Benoit, R. *et al*: Acidose lactique et phenformine. *L'Union Médicale du Canada*, - 1976;105:1810-1814.
4. Berger, W *et al*: Problèmes d'actualité concernant le mécanisme d'action des biguanides. *Jour. Diab. Hôtel-Dieu Paris*, 1975:239-258.
5. Bermond, P.: The coefficient of insulin efficacy. Effect of Metformin on this parameter. XIème Congrès Fédération Int. Diabétologie, Stockholm, 1967. *Ed. Excerpta medica F. Amsterdam*, 1968.
6. Biron, P.: Metformin monitoring. *C.M.A.J.*, 1980;123:11-12.
7. Bouaziz, Pl.: Apport à l'étude de l'épreuve d'hyperglycémie provoquée par voie veineuse sous thérapie diabétique. Thèse de doctorat en Médecine, Paris, 1966.
8. Canadian Diabetes Association. Nephropathy. *Canadian Journal of Diabetes*. 2003 Clinical Practice Guidelines;2003:S66-71.
9. Canadian Diabetes Association. Pharmacologic management of type 2 diabetes. *Canadian Journal of Diabetes*. 2003 Clinical Practice Guidelines;2003:S37-S42.
10. Canadian Diabetes Association. Physical activity and diabetes. *Canadian Journal of Diabetes*. 2003 Clinical Practice Guidelines;2003:S24-S26.
11. Canadian Diabetes Association. Targets for glycemic control. *Canadian Journal of Diabetes*. 2003 Clinical Practice Guidelines;2003:S18-S20.
12. Canadian Diabetes Association. Pre-existing diabetes and pregnancy. *Canadian Journal of Diabetes*. 2003 Clinical Practice Guidelines Expert Committee;2003:S94-S98.
13. Chan JCN, et al. Drug-induced disorders of glucose metabolism. Mechanisms and management. *Drug Safety*, 1996 Aug; 15(2):135-157.
14. Cohen, R.D.: The relative risks of different biguanides in the causation of lactic acidosis. *Research and Clinical Forums*, 1979;1(4):125-134.
15. Cohen, Y. *et al*: Etude autoradiographique chez la souris d'un antidiabétique oral, le N.N. Diméthylbiguanide, marqué au C<sup>14</sup>, *Thérapie*, 1961:109-120.

16. Cohen, Y. *et al*: Etude autoradiographique chez la souris d'un antidiabétique oral marqué au C<sup>14</sup>, le N.N. Diméthylbiguanide, après administrations répétées. *Thérapie* 1968; XXIII:1185-1191.
17. Cox D, et al. The effects of glucose fluctuation on cognitive function and QOL: the functional costs of hypoglycaemia and hyperglycaemia among adults with Type 1 or Type 2 diabetes. *IJCP* 2002 July; (Suppl.129):20-26.
18. Daubresse, J.C. *et al*: Acidose lactique et thérapeutique par biguanides. *Méd. et Hyg.* 1975:1168.
19. Debry, G. *et al*: Etude du mode d'excrétion du N.N. Diméthylbiguanide chez le diabétique adulte. *Thérapie*, 1963;XX:351-358.
20. Derot, M. *et al*: Retrospective study of the cardiovascular fate of 190 patients treated for 5 years or more with biguanides alone. Abstracts, 11<sup>th</sup> Annual Meeting, Munich Sept., 1975.
21. Duval, D.: Contribution à l'étude de l'action hypoglycémisante des biguanides. Thèse de Doctorat en Médecine, Paris, 1960.
22. Duwoos, H. *et al*: Hyperlactacidémie réversible induite par la phenformine avec asthénie musculaire et signes cardio-respiratoires. *Presse Méd.* 1970;78:23-26.
23. Glueck CJ, Wang P, Kobayashi S, Phillips H, Sieve-Smith L. Metformin therapy throughout pregnancy reduces the development of gestational diabetes in women with polycystic ovary syndrome. *Fertility and Sterility*. 2002;77(3):520-525.
24. Hermann, L.S.: Metformin: A review of its pharmacological properties and therapeutic use. *Diabète et Métabolisme*, 1979;3:233-245.
25. Hermann, L.S.: Metabolic effects of Metformin in relation to clinical effects and side effects in Biguanide Therapy Today. International Congress and Symposium, series published by the Royal Society of Medicine, 1981;48:17-43.
26. Holle, A. *et al*: Biguanide treatment increases the number of insulin receptor sites on human erythrocytes. *The New Engl. J. Med.* 1981;305(10): 563-566.
27. Hunt, J.A. *et al*: The use of phenformin and metformin. *Letter to the Editor, C.M.A.J.*-1977;117(5):429-430.
28. Irsigler, K.: Glucoseutilisation and Plasmaliporide bei adiposen Patienten unter dem Einfluss von Dimethylbiguanide (GLUCOPHAGE®). *Wiener med. Wsch.* 1969;119:191-194.

29. Isnard, F. *et al*: Acidose lactique et biguanides. Etat actuel de la question en France. Journées Annuelles de diabétologie de l'Hôtel-Dieu 1977:362-375.
30. Joncas, F: Evaluation clinique de GLUCOPHAGE<sup>®</sup> pour le traitement du diabète de l'adulte. Hôpital *Maisonneuve*, Montréal. L'Union Médicale du Canada, Jan. Issue, 1972.
31. Laurendeau, Ed. *et al*: Traitement du diabète sucré chez des patients âgés, hospitalisés avec le N.N. Diméthylbiguanide (GLUCOPHAGE<sup>®</sup>). Hôpital Notre-Dame de la Merci, Montréal, 1970. Ref Lab. Franca (non publié).
32. Lefebvre, P. *et al*: Le mécanisme d'action des biaguanides. Biguanides et sécrétion insulinique. Congrès International de Diabétologie de Rémini, 1968.
33. Le Jeune C, et al: Les effets hyperglycémiantes des médicaments. *Sem. Hop. Paris*. January 1994: 100-107.
34. Mainguet, P. *et al*: *Le diabète*,-1972;20(1):39
35. McKlish, A. Toxicité du N.N. Diméthylbiguanide chez le chien Beagle. Centre de recherches Laval, Québec (1970). Ref. Laboratoires Franca Inc. (non publié).
36. Meyer, F. *et al*: Données nouvelles sur le mécanisme d'action des biguanides hypoglycémiantes. Journées annuelles de diabétologie de l'Hôtel-Dieu, 1967:341-347
37. Pelletier, G. *et al*: Etude de toxicité chronique de N.N. Diméthylbiguanide chez le rat. Centre de Recherche Laval, Québec. Ref. Laboratoires Franca inc. (non publiée).
38. Pelletier, G. *et al*: Etude tératologique avec le N.N. Diméthylbiguanide chez le rat. Centre de Recherche Laval, Québec. Ref. Laboratoires Franca Inc. (non publiée) 1970.
39. Pignard, P.: Dosage spectrométrique du N.N. Diméthylbiguanide dans le sang et l'urine. *Annales de Biologie Clinique*. 1962:20:25-233
40. Sterne, J.: Oral Hypoglycemic agents. *Medicinal Chemistry*, 1969;9(5):193-294
41. Sterne, J.M. *et al*: Oral hypoglycemic agents: Clinical pharmacology and therapeutic use. *Drugs*, 1977;14:41-56
42. Sterne, J.: Pharmacology and mechanism of action of the antidiabetic biguanides, Paper read in Moscow, April 1977, Unpublished.
43. Stowers, J.M.: Long-term therapy with biguanides in: Biguanide Therapy Today, International Congress and Symposium, Series published by the Royal Society of Medecin, 1981;48:49-57

44. Stowers, J.M. *et al*: Oral hypoglycemic agents: Clinical pharmacology and therapeutic use. *Drugs*, 1977;14:41-56
45. Vague, P. Effet d'une dose unique de metformine sur la tolérance au glucose des sujets normaux obèses. *Le Diabète*, 1970;18:35-39
46. Vermulen, A. *et al*: Influence of dimethylbiguanide (metformin) on carbohydrate metabolism in obese, non diabetic women. *Diabetologia*,1972;8:8-11
47. UK Prospective Diabetes Study (UKPDS) Group. Effect of intensive blood-glucose control with metformin on complications in overweight patients with type 2 diabetes (UKPDS 34). *The Lancet*. 1998;352:854-865.
48. UK Prospective Diabetes Study (UKPDS) Group. Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS 33). *The Lancet*. 1998;352:837-853.
49. Product Monograph for Glucophage (Metformin HCl) by Sanofi-Aventis Canada Inc., June 20, 2006, Control #: 101051.

**PART III: CONSUMER INFORMATION****AVA-METFORMIN**  
metformin hydrochloride

**This leaflet is part III of a three-part "Product Monograph" published when AVA-METFORMIN was approved for sale in Canada and is designed specifically for Consumers. This leaflet is a summary and will not tell you everything about AVA-METFORMIN. Contact your doctor or pharmacist if you have any questions about the drug.**

**ABOUT THIS MEDICATION****What the medication is used for:**

**AVA-METFORMIN** (metformin hydrochloride) is used to treat Type II diabetes which cannot be controlled by diet, exercise and weight reduction.

**What it does:**

AVA-METFORMIN is used to treat type 2 diabetes. People with type 2 diabetes are not able to make enough insulin or respond normally to the insulin their bodies make. When this happens, sugar (glucose) builds up in the blood. This can lead to serious medical problems including kidney damage, amputations, and blindness. Diabetes is also closely linked to heart disease. The main goal of treating diabetes is to lower your blood sugar to a normal level.

High blood sugar can be lowered by diet and exercise, by a number of medicines taken by mouth, and by insulin shots. While you take your diabetes medicine, continue to exercise and follow the diet advised for your diabetes, by your doctor.

No matter what your recommended diabetes management plan is, studies have shown that maintaining good blood sugar control can prevent or delay, complications of diabetes, such as blindness.

AVA-METFORMIN helps control your blood sugar although the mode of action is not fully understood, it is believed to help your body respond better to the insulin it makes naturally by:

- decreasing the amount of sugar your liver makes, and
- decreasing the amount of sugar your intestines absorb.

**When it should not be used:**

Do not take AVA-METFORMIN if you

- have unstable and/or insulin-dependent (Type I) diabetes mellitus,
- have metabolic acidosis (including diabetic ketoacidosis, history of ketoacidosis or lactic acidosis-too much acid in the blood),
- have congestive heart failure,
- drink a lot of alcohol,
- have liver or kidney problems,
- are going to have x-ray procedure with injection of dyes,
- are stressed, have severe infections, are experiencing trauma, prior to surgery or during the recovery phase,
- suffer from severe dehydration (have lost a lot of water from your body)

- are hypersensitive or allergic to metformin hydrochloride or any ingredient in the formulation or component of the container
- are breastfeeding
- are pregnant or planning to become pregnant
- have cardiovascular collapse (abrupt failure of blood circulation) or cardiorespiratory insufficiency.

**What the medicinal ingredient is:**

The medicinal ingredient for AVA-METFORMIN is metformin hydrochloride

**What the non-medicinal ingredients are:**

AVA-METFORMIN tablets contain the following non-medicinal ingredients: Crospovidone, hydroxypropyl methylcellulose, magnesium stearate, maltodextrin, polyethylene glycol pregelatinised maize starch and povidone. This is a complete list of all non-medicinal ingredients.

**What dosage forms it comes in:**

AVA-METFORMIN is formulated into tablets of 500 mg and 850 mg strengths for oral administration.

**WARNINGS AND PRECAUTIONS****Serious Warnings and Precautions**

AVA-METFORMIN may rarely cause a serious, life-threatening condition called lactic acidosis (see section Lactic Acidosis below).

You should not drink a lot of alcohol if you take AVA-METFORMIN (see section Lactic Acidosis below).

**Lactic Acidosis**

AVA-METFORMIN may rarely cause a serious, life-threatening condition called lactic acidosis.

You should not take AVA-METFORMIN due to greater risk for lactic acidosis if you:

- have kidney problems
- are 80 years or older and you have NOT had a kidney function tested
- are taking medication for heart failure
- are seriously dehydrated (have lost a lot of water from your body)
- have heart disease
- have liver disease
- drink a lot of alcohol (regularly drink alcohol or sometimes drink a lot of alcohol, binge drinking)
- have an x-ray procedure with injection of dyes (contrast agents)
- prior to surgery or during recovery phase
- develop a serious medical condition, such as heart attack, severe infection, or a stroke



## IMPORTANT PLEASE READ

Signs and symptoms of lactic acidosis include: discomfort, muscle pain, difficult or fast breathing, extreme tiredness, weakness, upset stomach, stomach pain, feeling cold, low blood pressure or slow heartbeat.

If any of the above side effects occur, consult your doctor immediately.

Tell your doctor if you are pregnant or plan to become pregnant. AVA-METFORMIN should not be used during pregnancy and insulin treatment is recommended during pregnancy. Talk with your doctor about your choices. You must not take AVA-METFORMIN if you are nursing a child.

Tell your doctor of any other medical condition including: vitamin B12 deficiency or anemia, excessive alcohol use, allergies.

Do not start or stop any medicine without doctor or pharmacist approval.

### INTERACTIONS WITH THIS MEDICATION

Some drugs may interact with AVA-METFORMIN. Careful monitoring is advised. Tell your doctor if you are taking:

- other diabetes drugs such as glyburide
- furosemide
- nifedipine
- cationic drugs (e.g., amiloride, digoxin, morphine, procainamide, quinidine, quinine, ranitidine, triamterene, trimethoprim, and vancomycin)
- other drugs tend to produce hyperglycemia (high blood sugar) and may lead to a loss of blood sugar control. Some example of drugs that can increase the blood sugar include:
  - Thiazide and other diuretics (water pills)
  - Corticosteroids
  - Phenothiazines
  - Thyroid products
  - Estrogens or estrogens plus progestogen
  - Oral contraceptives
  - Phenytoin
  - Nicotinic Acid
  - Sympathomimetics
  - Calcium channel blocking drugs
  - Isoniazid
  - Beta-2-agonists
- ACE inhibitors drugs may lower blood glucose and the combination with AVA-METFORMIN should be carefully monitored.

Before using any drugs or herbal products, check with your doctor or your pharmacist.

### PROPER USE OF THIS MEDICATION

#### Usual dose:

Your doctor will tell you how much medicine to take and when to take it. Follow the directions provided by your doctor for using this

medicine. Taking this medicine with food will decrease symptoms such as nausea and vomiting.

#### Overdose:

In general, an overdose may lead to increased symptoms as described under "SIDE EFFECTS AND WHAT TO DO ABOUT THEM" including stomach discomfort, nausea, vomiting, diarrhea, drowsiness, weakness, dizziness, malaise and headache.

A serious, life-threatening condition called lactic acidosis may also occur (see **WARNINGS and PRECAUTIONS**, Lactic Acidosis).

If you have taken too much AVA-METFORMIN, immediately see your doctor, contact the poison control or go to the nearest hospital emergency department. Do this even if there are no signs of discomfort or poisoning.

#### Missed Dose:

If you forget to take AVA-METFORMIN tablets, do not take a double dose to make up for forgotten individual doses. Take the next dose at the usual time.

### SIDE EFFECTS AND WHAT TO DO ABOUT THEM

Common side effects of AVA-METFORMIN include

- diarrhea
- nausea
- upset stomach
- abdominal bloating
- gas
- loss of appetite

These side effects generally go away after you take the medicine for a while. Taking your medicine with meals can help reduce these side effects. Tell your doctor if the side effects bother you a lot, last for more than a few weeks, come back after they've gone away, or start later in therapy. You may need a lower dose or need to stop taking the medicine for a short period or for good.

AVA-METFORMIN rarely cause hypoglycemia (low blood sugar) by themselves. However, hypoglycemia can happen if you do not eat enough, if you drink alcohol, or if you take other medicines to lower blood sugar.

**Lactic Acidosis. In a rare cases, AVA-METFORMIN can cause a serious side effects called lactic acidosis. This is caused by a buildup of lactic acid in your blood. This buildup can cause serious damage.** Lactic acidosis caused by AVA-METFORMIN is rare and has occurred mostly in people whose kidneys were not working normally. Although rare, if lactic acidosis does occur, it can be fatal in up to half the people who develop it.

It is also important for your liver to be working normally when you take AVA-METFORMIN. Your liver helps remove lactic acid from your blood.

Make sure you tell your doctor before you use AVA-METFORMIN if you have kidney or liver problems.

You should also **stop using AVA-METFORMIN and call your**

## IMPORTANT PLEASE READ

**doctor right away if you have signs of lactic acidosis. Lactic acidosis is a medical emergency that must be treated in a hospital.**

If your medical condition suddenly changes, stop taking AVA-METFORMIN and call your doctor right away. This may be a sign of lactic acidosis or another serious side effect.

<b>SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM</b>	
Symptom / effect	Stop taking drug and call your doctor or pharmacist
<b>Uncommon</b>	
Feeling very weak, tired or uncomfortable	√
Unusual muscle pain	√
Trouble breathing	√
Unusual or unexpected stomach discomfort	√
Feeling cold	√
Feeling dizzy or lightheaded	√
Suddenly developing a slow or irregular heartbeat	√
<b>Rare</b>	
<b>Lactic acidosis and symptoms which may be</b> <ul style="list-style-type: none"> <li>• Feeling very weak, tired or uncomfortable</li> <li>• Unusual muscle pain</li> <li>• Trouble breathing</li> <li>• Unusual or unexpected stomach discomfort</li> <li>• Feeling cold</li> <li>• Feeling dizzy or lightheaded</li> <li>• Suddenly developing a slow or irregular heartbeat</li> </ul>	√

*This is not a complete list of side effects. For any unexpected effects while taking AVA-METFORMIN, contact your doctor or*

*pharmacist.*

### HOW TO STORE IT

Store in a dry place between 15°C-30°C in well closed containers. Protect from light. Throw away any medication that is outdated or no longer needed. Talk to your pharmacist about the proper disposal of your medication.

Keep out of reach of children.

### REPORTING SUSPECTED SIDE EFFECTS

**To monitor drug safety, Health Canada collects information on serious and unexpected effects of drugs . If you suspect you have had a serious or unexpected reaction to this drug you may notify Health Canada by:**

**toll-free telephone: 866-234-2345**

**toll-free fax 866-678-6789**

**By email: [cadmp@hc-sc.gc.ca](mailto:cadmp@hc-sc.gc.ca)**

**By regular mail:**

**National AR Centre**

**Marketed Health Products Safety and Effectiveness**

**Information Division**

**Marketed Health Products Directorate**

**Tunney's Pasture, AL 0701C**

**Ottawa ON K1A 0K9**

***NOTE: Before contacting Health Canada, you should contact your physician or pharmacist.***

### MORE INFORMATION

This document plus the full Product Monograph, prepared for health professionals can be found by contacting the sponsor at: 1-855-708-3678 or [medinfo@avanstra.com](mailto:medinfo@avanstra.com)

This leaflet was prepared by:

Avanstra Inc.

10761 – 25th NE

Suite 110, Building “B”

Calgary, Alberta

T2C 3C2, Canada

Last revised: February 21, 2011