

PRODUCT MONOGRAPH

Riva-Ondansetron
(Ondansetron Tablets USP)

4 mg and 8 mg Ondansetron (as Ondansetron Hydrochloride)

Antiemetic
(5-HT₃ receptor antagonist)

**Laboratoire Riva Inc.,
660 Boul. Industriel,
Blainville Québec
J7C 3V4**

**Date of Preparation:
July 22, 2011**

Submission Control No.: 148445

Table of Contents

PART I	HEALTH PROFESSIONAL INFORMATION	3
	SUMMARY PRODUCT INFORMATION	3
	INDICATIONS AND CLINICAL USE	3
	CONTRAINDICATIONS	4
	WARNINGS AND PRECAUTIONS	4
	ADVERSE REACTIONS	5
	DRUG INTERACTIONS	7
	DOSAGE AND ADMINISTRATION	8
	OVERDOSAGE	11
	ACTION AND CLINICAL PHARMACOLOGY	11
	STORAGE AND STABILITY	12
	DOSAGE FORMS, COMPOSITION AND PACKAGING	12
PART II	SCIENTIFIC INFORMATION	14
	PHARMACEUTICAL INFORMATION	14
	CLINICAL TRIALS	15
	DETAILED PHARMACOLOGY	18
	MICROBIOLOGY	20
	TOXICOLOGY	20
	REFERENCES	22
PART III:	CONSUMER INFORMATION	25

PRODUCT MONOGRAPH
RIVA-ONDANSETRON
(Ondansetron tablets USP)

Antiemetic
(5-HT₃ receptor antagonist)

PART I: HEALTH PROFESSIONAL INFORMATION

SUMMARY PRODUCT INFORMATION

Route of Administration	Dosage Form / Strength	Clinically Relevant Non-medicinal Ingredients
Oral	Tablets/ 4 mg and 8 mg Ondansetron (as ondansetron hydrochloride)	Lactose For a complete listing see Dosage Forms, Composition and Packaging section.

INDICATIONS AND CLINICAL USE

Adults

RIVA-ONDANSETRON (Ondansetron Hydrochloride) is indicated for:

- the prevention of nausea and vomiting associated with emetogenic chemotherapy, including high dose cisplatin, and radiotherapy.
- the prevention and treatment of post-operative nausea and vomiting

Pediatrics (< 18 years of age)

Post-Chemotherapy

Clinical experience of Ondansetron in children is currently limited, however, Ondansetron hydrochloride was effective and well tolerated when given to children 4-12 years of age (see DOSAGE AND ADMINISTRATION). Ondansetron hydrochloride is not indicated for the treatment of children 3 years of age or younger.

Post-Radiotherapy

Safety and efficacy of RIVA-ONDANSETRON in any age group in this population following radiotherapy has not been established and is therefore not indicated for use in this population.

Post-Operative Nausea and Vomiting

Safety and efficacy of RIVA-ONDANSETRON in any age group in this population for the prevention and treatment of post-operative nausea and vomiting has not been established and is not indicated for use in this group.

Geriatrics (> 65 years of age)

Post-Chemotherapy and Radiotherapy

Efficacy and tolerance of Ondansetron hydrochloride were similar to that observed in younger adults (see DOSAGE AND ADMINISTRATION).

Post-Operative Nausea and Vomiting

Clinical experience in the use of RIVA-ONDANSETRON in the prevention and treatment of postoperative nausea and vomiting is limited and is not indicated for use in this population.

CONTRAINDICATIONS

- RIVA-ONDANSETRON (Ondansetron hydrochloride) is contraindicated in patients with a history of hypersensitivity to the drug or any components of its formulations. For a complete listing, see the **Dosage Forms, Composition and Packaging** section of the product monograph.

WARNINGS AND PRECAUTIONS

General

Cross-reactive hypersensitivity has been reported between different 5-HT₃ antagonists. Patients who have experienced hypersensitivity reactions to one 5-HT₃ antagonist have experienced more severe reactions upon being challenged with another drug of the same class. The use of a different 5-HT₃ receptor antagonist is not recommended as a replacement in cases in which a patient has experienced even a mild hypersensitivity type reaction to another 5-HT₃ antagonist.

Rarely and predominantly with intravenous ondansetron, transient ECG changes including QT interval prolongation have been reported (see Post-Market Adverse Drug Reactions).

Riva-Ondansetron is not effective in preventing motion-induced nausea and vomiting.

Hepatic/Biliary/Pancreatic

There is no experience in patients who are clinically jaundiced. The clearance of an 8mg intravenous dose of ondansetron was significantly reduced and the serum half-life significantly prolonged in subjects with severe impairment of hepatic function. In patients with moderate or severe impairment of hepatic function, reductions in dosage are therefore recommended and a total daily dose of 8 mg should not be exceeded. This may be given as a single intravenous or oral dose. As ondansetron is known to increase large

bowel transit time, patients with signs of subacute intestinal obstruction should be monitored following administration.

Ondansetron does not itself appear to induce or inhibit the cytochrome P₄₅₀ drug-metabolizing enzyme system of the liver. Because ondansetron is metabolized by hepatic cytochrome P₄₅₀ drug-metabolizing enzymes, inducers or inhibitors of these enzymes may change the clearance and, hence, the half-life of ondansetron. On the basis of available data no dosage adjustment is recommended for patients on these drugs.

Special Populations

Pregnant Women:

The safety of ondansetron for use in human pregnancy has not been established. Ondansetron is not teratogenic in animals. However, as animal studies are not always predictive of human response, the use of ondansetron in pregnancy is not recommended.

Nursing Women: Ondansetron is excreted in the milk of lactating rats. It is not known if it is excreted in human milk, however, nursing is not recommended during treatment with ondansetron.

Pediatrics: (< 3 years of age): Insufficient information is available to provide dosage recommendations for children 3 years of age or younger.

ADVERSE REACTIONS

Clinical Trial Adverse Drug Reactions

Because clinical trials are conducted under very specific conditions the adverse reaction rates observed in the clinical trials may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse drug reaction information from clinical trials is useful for identifying drug-related adverse events and for approximating rates.

Ondansetron has been administered to over 2500 patients worldwide in controlled clinical trials and has been well tolerated.

The most frequent adverse events reported in controlled clinical trials were headache (11%) and constipation (4%). Other adverse events include sensations of flushing or warmth (< 1%).

Cardiovascular:

There have been rare reports of tachycardia, angina (chest pain), bradycardia, hypotension, syncope and electrocardiographic alterations.

Central Nervous System:

There have been rare reports of seizures. Movement disorders and dyskinesia have been reported in two large clinical trials of ondansetron at a rate of 0.1 – 0.3%.

Dermatological:

Rash has occurred in approximately 1% of patients receiving ondansetron.

Hypersensitivity:

Rare cases of immediate hypersensitivity reactions sometimes severe, including anaphylaxis, bronchospasm, urticaria and angioedema have been reported.

Local Reactions:

Pain, redness and burning at the site of injection have been reported.

Metabolic:

There were transient increases of SGOT and SGPT of over twice the upper limit of normal in approximately 5% of patients. These increases did not appear to be related to dose or duration of therapy. There have been reports of liver failure and death in patients with cancer receiving concurrent medications including potentially hepatotoxic cytotoxic chemotherapy and antibiotics. The etiology of the liver failure is unclear. There have been rare reports of hypokalemia.

Other:

There have been reports of abdominal pain, weakness and xerostomia.

Special Senses:

Rare cases of transient visual disturbances (e.g. blurred vision) have been reported during or shortly after intravenous administration of ondansetron, particularly at rates equal to or greater than 30 mg in 15 minutes.

Post-Market Adverse Drug Reactions

Over 250 million patient treatment days of ondansetron hydrochloride have been supplied since the launch of the product worldwide. The following events have been spontaneously reported during post-approval use of ondansetron hydrochloride, although the link to ondansetron cannot always be clearly established.

General Disorders:

Rare cases of hypersensitivity reactions, such as, laryngeal edema, stridor, laryngospasm and cardiopulmonary arrest have also been reported.

Cardiovascular Disorders:

There have been rare reports (< 0.01%) of myocardial infarction, myocardial ischemia, angina, chest pain with or without ST segment depression, arrhythmias (including ventricular or supraventricular tachycardia, premature ventricular contractions, and atrial fibrillation), electrocardiographic alterations (including second degree heart block), palpitations and syncope.

Rarely and predominantly with intravenous ondansetron, transient ECG changes including QT interval prolongation have been reported (see WARNINGS AND PRECAUTIONS).

Eye Disorder:

There have been very rare cases of transient blindness following ondansetron treatment, generally within the recommended dosing range and predominantly during intravenous administration.

The majority of blindness cases reported resolved within 20 minutes. Although most patients had received chemotherapeutic agents, including cisplatin a few cases of transient blindness occurred following ondansetron administration for the treatment of post-operative nausea or vomiting and in the absence of cisplatin treatment. Some cases of transient blindness were reported as cortical in origin.

Hepatobiliary Disorders:

Occasional asymptomatic increases in liver function tests have been reported.

Nervous System Disorders:

Transient episodes of dizziness (< 0.01%) have been reported during or upon completion of IV infusion of ondansetron.

Uncommon reports (<1%) suggestive of extrapyramidal reactions including oculogyric crisis/dystonic reactions (e.g. oro-facial dyskinesia, opisthotonos, tremor, etc.), movement disorders and dyskinesia have been reported without definitive evidence of persistent clinical sequelae.

Respiratory, Thoracic and Mediastinal Disorders:

There have also been rare reports of hiccups.

Very rare reports have been received for bullous skin and mucosal reactions (including Stevens-Johnson syndrome and toxic epidermal necrolysis). These reports have occurred in patients taking other medications that can be associated with bullous skin and mucosal reactions.

DRUG INTERACTIONS**Drug-Drug Interactions**

Specific studies have shown that there are no pharmacokinetic interactions when ondansetron is administered with alcohol, temazepam, frusemide, tramadol or propofol.

Ondansetron is metabolized by multiple hepatic cytochrome P₄₅₀ enzymes: CYP3A4, CYP2D6 and CYP1A2. Despite the multiplicity of metabolic enzymes capable of metabolizing ondansetron which can compensate for an increase or decrease in enzyme activity, it was found that patients treated with inducers of CYP3A4 (i.e. phenytoin, carbamazepine, and rifampicin) demonstrated an increase in oral clearance of ondansetron and a decrease in ondansetron blood concentrations. No effect in ondansetron clearance secondary to enzyme inhibition or reduced activity (e.g. CYP2D6 genetic deficiency) has been identified to date.

Data from small studies indicate that ondansetron may reduce the analgesic effect of tramadol.

DOSAGE AND ADMINISTRATION

Dosing Considerations

Chemotherapy Induced Nausea and Vomiting:

Ondansetron hydrochloride should be given as an initial dose prior to chemotherapy, followed by a dosage regimen tailored to the anticipated severity of emetic response caused by different cancer treatments. The route of administration and dose of ondansetron hydrochloride should be flexible in the range of 8-32 mg a day. The selection of dose regimen should be determined by the severity of the emetogenic challenge (See Recommended Dose and Dosage Adjustment).

Recommended Dose And Dosage Adjustment

As Riva-Ondansetron is not available as a solution for injection, please consult a product monograph for ondansetron hydrochloride injection for dosage instructions when required as noted below.

Chemotherapy Induced Nausea and Vomiting:

Use in Adults:

Highly Emetogenic Chemotherapy (e.g. regimens containing cisplatin)

Ondansetron hydrochloride has been shown to be effective in the following dose schedules for the prevention of emesis during the first 24 hours following chemotherapy:

Post-chemotherapy:

After the first 24 hours, RIVA-ONDANSETRON 8 mg orally every 8 hours for up to 5 days.

No significant differences in terms of emesis control or grade of nausea have been demonstrated between the 32 mg single dose, the 8 mg single dose, or the 8 mg dose followed by the 24 hour 1 mg/h continuous infusion.

However, in some studies conducted in patients receiving medium or high dose of cisplatin chemotherapy, the 32 mg single dose has demonstrated a statistically significant superiority over the 8 mg single dose with regard to control of emesis.

The efficacy of Riva-Ondansetron in highly emetogenic chemotherapy may be enhanced by the addition of a single intravenous dose of dexamethasone sodium phosphate, 20mg administered prior to chemotherapy.

-
- i) The efficacy of twice daily dosage regimens for the treatment of post-chemotherapy emesis has been established only in adult patients receiving less emetogenic chemotherapy. The appropriateness of twice versus three times

daily dosage regimens for other patient groups should be based on an assessment of the needs and responsiveness of the individual patient.

Less Emetogenic Chemotherapy (e.g. regimens containing cyclophosphamide, doxorubicin, epirubicin, fluorouracil and carboplatin)

Initial Dose:

RIVA-ONDANSETRON 8 mg infused intravenously over 15 minutes, given 30 minutes prior to chemotherapy, or ondansetron hydrochloride 8 mg orally 1 to 2 hours prior to chemotherapy

Post-chemotherapy:

RIVA-ONDANSETRON 8 mg orally twice daily for up to 5 days.

Use in Children:

Clinical experience of ondansetron in children is currently limited however, ondansetron was effective and well tolerated when given to children 4-12 years of age. Ondansetron injection should be given intravenously at a dose of 3-5 mg/m² over 15 minutes immediately before therapy. After chemotherapy, RIVA-ONDANSETRON 4mg should be given orally every 8 hoursⁱ for up to 5 days. For children 3 years of age and younger, there is insufficient information available to make dosage recommendations (see INDICATIONS AND CLINICAL USE).

Use in Elderly:

Efficacy and tolerance in patients aged over 65 years were similar to that seen in younger adults indicating no need to alter dosage schedules in this population.

Radiotherapy Induced Nausea and Vomiting:

Use in Adults:

Initial Dose:

RIVA-ONDANSETRON 8 mg orally 1 to 2 hours before radiotherapy

Post-radiotherapy:

RIVA-ONDANSETRON 8 mg orally every 8 hoursⁱ for up to 5 days after a course of treatment

Use in Children:

There is no experience in clinical studies in this population.

Use in Elderly:

Efficacy and tolerance in patients aged over 65 years were similar to that seen in younger adults indicating no need to alter dosage schedules in this population.

Post-Operative Nausea and Vomiting:

Use in Adults:

For prevention of post-operative nausea and vomiting RIVA-ONDANSETRON may be administered as a single dose of 16 mg given orally one hour prior to anaesthesia. Alternatively, a single dose of 4 mg may be given by slow intravenous injection at induction of anaesthesia.

For the treatment of established post-operative nausea and vomiting, a single dose of 4mg given by slow intravenous injection is recommended.

Use in Children:

There is no experience in the use of ondansetron hydrochloride in the prevention and treatment of postoperative nausea and vomiting in children (see INDICATIONS AND CLINICAL USE).

Use in Elderly:

There is limited experience in the use of ondansetron hydrochloride in the prevention and treatment of post-operative nausea and vomiting in the elderly (see INDICATIONS AND CLINICAL USE).

Patients with Renal/Hepatic Impairment:

Use in Patients with Impaired Renal Function:

No alteration of daily dosage, frequency of dosing, or route of administration is required.

Use in Patients with Impaired Hepatic Function:

The clearance of an 8 mg intravenous dose of ondansetron was significantly reduced and the serum half-life significantly prolonged in subjects with severe impairment of hepatic function. In patients with moderate or severe impairment of hepatic function, reductions in dosage are therefore recommended and a total daily dose of 8 mg should not be exceeded. This may be given as a single intravenous or oral dose.

No studies have been conducted to date in patients with jaundice.

Patients with Poor Sparteine/Debrisoquine Metabolism:

The elimination half-life and plasma levels of a single 8 mg intravenous dose of ondansetron did not differ between subjects classified as poor and extensive metabolisers of sparteine and debrisoquine. No alteration of daily dosage or frequency of dosing is recommended for patients known to be poor metabolisers of sparteine and debrisoquine.

OVERDOSAGE

At present there is little information concerning overdosage with ondansetron. Individual doses of 84 mg and 145 mg and total daily doses as large as 252 mg have been administered with only mild side effects. There is no specific antidote for ondansetron, therefore, in cases of suspected overdosage, symptomatic and supportive therapy should be given as appropriate.

The use of Ipecac to treat overdosage with ondansetron is not recommended as patients are unlikely to respond due to the anti emetic action of ondansetron itself.

“Sudden blindness” (amaurosis) of 2 to 3 minutes duration plus severe constipation occurred in one patient that was administered 72 mg of ondansetron intravenously as a single dose. Hypotension (and faintness) occurred in another patient that took 48 mg of oral ondansetron. Following infusion of 32 mg over only a 4-minute period, a vasovagal episode with transient second degree heart block was observed. In all instances, the events resolved completely.

For management of a suspected drug overdose, contact your regional poison control centre.

ACTION AND CLINICAL PHARMACOLOGY

Mechanism of Action

Ondansetron hydrochloride is a selective antagonist of the serotonin receptor subtype, 5-HT₃. Its precise mode of action in the control of chemotherapy induced nausea and vomiting is not known.

Cytotoxic chemotherapy and radiotherapy are associated with the release of serotonin (5-HT) from enterochromaffin cells of the small intestine, presumably initiating a vomiting reflex through stimulation of 5-HT₃ receptors located on vagal afferents. Ondansetron may block the initiation of this reflex. Activation of vagal afferents may also cause a central release of serotonin from the chemoreceptor trigger zone of the area postrema, located on the floor of the fourth ventricle. Thus, the antiemetic effect of ondansetron is

probably due to the selective antagonism of 5-HT₃ receptors on neurons located in either the peripheral or central nervous systems, or both.

The mechanisms of ondansetron's antiemetic action in post-operative nausea and vomiting are not known.

Pharmacodynamics

In vitro metabolism studies have shown that ondansetron is a substrate for human hepatic cytochrome P₄₅₀ enzymes, including CYP1A2, CYP2D6 and CYP3A4. In terms of overall ondansetron turnover, CYP3A4 played the predominant role. Because of the multiplicity of metabolic enzymes capable of metabolizing ondansetron, it is likely that inhibition or loss of one enzyme (e.g. CYP2D6 enzyme deficiency) will be compensated by others and may result in little change in overall rates of ondansetron clearance.

Pharmacokinetics

Pharmacokinetic studies in human volunteers showed peak plasma levels of 20-30 ng/mL at around 1½ hours after an 8 mg oral dose of ondansetron. An 8 mg infusion of Ondansetron resulted in peak plasma levels of 80-100 ng/mL. Repeat dosing of an 8 mg tablet every 8 hours for 6 days increased the peak plasma value to 40 ng/mL. A continuous intravenous infusion of 1 mg/hour after the initial 8 mg loading dose of Ondansetron maintained plasma levels over 30 ng/mL during the following 24 hour period.

The absolute bioavailability of ondansetron in humans was approximately 60% and the plasma protein binding was approximately 73%.

Following oral or IV administration, Ondansetron is extensively metabolized and excreted in the urine and faeces. In humans, less than 10% of the dose is excreted unchanged in the urine. The major urinary metabolites are glucuronide conjugates (45%), sulphate conjugates (20%) and hydroxylation products (10%).

The half-life of Ondansetron after either an 8 mg oral dose or intravenous dose was approximately 3-4 hours and may be extended to 6-8 hours in the elderly.

In a pharmacokinetic study of 16 epileptic patients maintained chronically on carbamazepine or phenytoin, reduction in AUC, C_{max} and T_½ of Ondansetron was observed. This resulted in a significant increase in clearance. However, on the basis of available data, no dosage adjustment is recommended (see WARNINGS AND PRECAUTIONS).

STORAGE AND STABILITY

RIVA-ONDANSETRON tablets store at room temperature (15-30°C) in a dry place.

DOSAGE FORMS, COMPOSITION AND PACKAGING

RIVA-ONDANSETRON Tablets 4 mg:

Oval shaped, yellow, film-coated tablets, engraved '4' on one face. Each tablet contains 4 mg ondansetron (as ondansetron hydrochloride) and the following excipients: lactose, microcrystalline cellulose, pregelatinised starch, magnesium stearate, methyl hydroxypropyl cellulose, triacetin, croscarmellose sodium, iron oxide yellow and titanium dioxide

Available in a tamper-evident polyethylene container of 30 tablets and blister pack of 6 tablets

RIVA-ONDANSETRON Tablets 8 mg:

Oval shaped, yellow, film-coated tablets, engraved '8' on one face. Each tablet contains 8 mg ondansetron (as ondansetron hydrochloride) and the following excipients: lactose, microcrystalline cellulose, pregelatinised starch, magnesium stearate, methyl hydroxypropyl cellulose, triacetin, croscarmellose sodium, iron oxide yellow and titanium dioxide

Available in a tamper-evident polyethylene container of 30 tablets and blister pack of 6 tablets

PART II. SCIENTIFIC INFORMATION

PHARMACEUTICAL INFORMATION

Drug Substance

Proper Name : USP: Ondansetron hydrochloride
Ph.Eur: Ondansetron hydrochloride dihydrate.


Chemical Name : (±)-1,2,3,9-tetrahydro-9-methyl-3-[(2-methyl-1H-imidazol-1-yl)methyl]-4H-carbazol-4-one, hydrochloride, dihydrate.

Molecular formula and molecular mass:

Ondansetron hydrochloride (dihydrate): $C_{18}H_{19}N_3O \cdot HCl \cdot 2H_2O$, 365.9

Ondansetron base: $C_{18}H_{19}N_3O$, 293.4

Structural formula:



. HCl. 2H₂O

Physicochemical properties:

Description and Solubility:

Ondansetron hydrochloride is sparingly soluble in water and in alcohol, soluble in methanol, slightly soluble in isopropyl alcohol and in dichloromethane, very slightly soluble in acetone, chloroform and ethyl acetate.

Solubility profile of Ondansetron Hydrochloride in water is 25mg/ml. Quantitative aqueous pH solubility profile of Ondansetron Hydrochloride is 28.57mg/ml at pH 2.0, 23.53mg/ml at pH 3.0 and 20.00mg/ml at pH 6.0.

CLINICAL TRIALS

Study results

Clinical trial results showing the number and percentage of patients exhibiting a complete response to ondansetron (0 emetic episodes) are shown in the tables below for both postoperative and chemotherapy induced emesis.

PREVENTION OF CHEMOTHERAPY INDUCED EMESIS – RESPONSE OVER 24 HOURS		
DOSE	Ondansetron[@] 3 doses of 0.15 mg/kg	Placebo[@] 3 doses of placebo
# of patients	14	14
Treatment Response		
0 emetic episodes	2 (14%)	0 (0%)
1-2 emetic episodes	8 (57%)	0 (0%)

@Results are from an initial study using a different dosing regimen.

PREVENTION OF POST-OPERATIVE EMESIS – RESPONSE OVER 24 HOURS □			
	ORAL PREVENTION		
DOSE	Ondansetron 16 mg od	Placebo	pValue
# of patients	253	250	
Treatment Response			
0 emetic episodes	126 (50%)	79 (32%)	<0.001

The majority of patients included in the prevention and treatment of post-operative nausea and vomiting studies using Ondansetron Hydrochloride have been adult women receiving balanced anaesthesia for gynaecological surgery.

Pivotal Comparative Bioavailability Studies

Open labeled, randomized two-treatment, two-period, two-sequence single dose, crossover bioequivalence study of ondansetron hydrochloride 8 mg tablets (Laboratoire

Riva Inc.), compared with Zofran™ containing ondansetron hydrochloride 8mg tablets (GlaxoSmithKline Inc., USA) in 24 (+2 stand by) healthy adult, human subjects under fasted and fed conditions.

**SUMMARY TABLE OF THE COMPARATIVE BIOAVAILABILITY DATA
(FASTING CONDITIONS)**

Ondansetron (one x 8 mg) From measured data uncorrected for potency Geometric Mean Arithmetic Mean (CV %)					
Parameter	Riva- Ondansetron*	Zofran†	% Ratio of Geometric Means	90% Confidence Interval**	
				Lower	Upper
AUC _T (ng.hr/mL)	225.712 248.841 (44.88)	210.399 229.313 (44.57)	107.28	99.00	116.25
AUC _∞ (ng.hr/mL)	233.538 256.740 (44.21)	218.660 236.960 (43.41)	106.80	98.73	115.54
C _{max} (ng/mL)	32.154 34.451 (36.96)	31.405 33.788 (41.04)	102.38	94.0	111.51
T _{max} § (hr)	2.19 (30.19)	2.19 (30.56)			
T _{1/2} € (hr)	6.65 (39.07)	6.48 (31.72)			

* Riva-Ondansetron, Manufactured by Laboratoire Riva Inc.

† Zofran manufactured by GlaxoSmithKline Inc, USA

§ Expressed as the arithmetic mean (CV%) only

€ Expressed as the arithmetic mean (CV%) only

** Indicate % Confidence Interval (i.e., 90% or 95%) in the column heading and list for the AUC_T, AUC_I and C_{max}

DETAILED PHARMACOLOGY

Animal Pharmacology

Pharmacodynamics:

The ferret provides an excellent model for demonstrating the antiemetic action of drugs. Emesis can be induced by antineoplastic drugs or whole body irradiation. Behavioural changes associated with these treatments are noted in these animals and may also provide a parallel for the human experience of nausea.

The antiemetic action of ondansetron has been evaluated in both male and female ferrets given cisplatin (9-10 mg/kg), cyclophosphamide (200 mg/kg) or irradiation (2 and 8 Gy, 250 kV). Intravenous doses of ondansetron (0.1-1 mg/kg) abolished cisplatin-induced emesis for up to 2 hours. In cyclophosphamide-induced emesis, subcutaneous doses of 0.5 mg/kg ondansetron completely eliminated vomiting, significantly reduced retching and delayed the onset of these responses.

The radiation-induced emesis, 0.5 mg/kg ondansetron alone completely and rapidly eliminated retching and vomiting.

The antiemetic effects of ondansetron (0.1 mg/kg) in combination with dexamethasone (2-5 mg/kg) were potentiated in ferrets with cyclophosphamide-induced emesis, compared with ondansetron alone. Ondansetron with dexamethasone produced a significant reduction in retching (65%) and vomiting (72%).

Serotonin receptors of the 5HT₃ type are present both peripherally and on vagal nerve terminals. Ondansetron probably acts by preventing activation of these receptors or receptors located in other regions of the central nervous system. Both the peripheral and central nervous systems appear to be involved since both abdominal vagotomy and microinjection of Ondansetron and other 5-HT₃ antagonists directly into the area postrema eliminate cisplatin-induced emesis, while 5-HT₁-like (methiothepin maleate) and 5-HT₂ (ketanserin) antagonists have no effect.

Ondansetron is highly selective for 5-HT₃ receptors and shows negligible binding to other receptors such as 5-HT₁-like, 5-HT₂, α 1 and α 2 adrenoceptors, β 1 and β 2 adrenoceptors, D₁ and D₂ muscarinic, nicotinic, GABA_A, H₁ and H₂ receptors.

The pharmacological specificity of ondansetron may explain the observed lack of extrapyramidal side effects often seen following similar therapy with metoclopramide, which preferentially binds to dopamine receptors of the D₂ subtype.

Among its secondary effects, ondansetron has also been shown to cause a dose-dependent increase in the rate of gastric emptying in the guinea pig which is significant at doses of 0.01-0.1 mg/kg. As gastric stasis is frequently associated with nausea, stimulation of

gastric motility may be a beneficial action of ondansetron. In the cat, dog and monkey, ondansetron has little effect on heart rate, blood pressure or ECG at intravenous doses up to 3 mg/kg.

A study in cloned human cardiac ion channels has shown ondansetron has the potential to affect cardiac repolarisation via blockade of HERG potassium channels. The concentration at which this effect was seen may be attainable with the 32 mg IV dose, however, the clinical relevance of this finding is uncertain.

Pharmacokinetics:

In mice, rats, rabbits and dogs dosed at 1 mg/kg orally and/or intravenously, the plasma half-life of ondansetron was less than 1 hour, but the half-lives of its metabolites were significantly longer. Peak plasma concentrations of ondansetron in rats and dogs ranged from 351 to 419 ng/mL for the IV dose and 8 to 15 ng/mL for the oral dose. Plasma levels were linear over a 30 fold dose range. In repeat dose studies there was no apparent accumulation of ondansetron.

Ondansetron is almost completely absorbed in animals, and is rapidly metabolized by N-demethylation and hydroxylation of the indole ring, followed by conjugation with glucuronic acid and sulphate. There is significant first-pass metabolism after oral doses.

Ondansetron and its metabolites are rapidly and widely distributed in tissues, reaching higher levels than the corresponding plasma levels. In the rat and dog, ondansetron binds reversibly to tissues containing melanin and elastin. In rats and man, plasma protein binding is about 73%, while it is slightly lower in the dog (60%). Ondansetron and its metabolites cross the blood-brain barrier to only a slight extent.

Human Pharmacology

Pharmacodynamics:

In vivo pharmacodynamic studies have investigated the effects of ondansetron on gastric emptying, small bowel transit time and oesophageal motility.

Both oral (16 mg tid) and intravenous (5-10 mg) doses of ondansetron failed to produce a significant effect on gastric emptying in both healthy volunteers and in patients suffering from delayed gastric emptying. However, in one study intravenous doses of 8 mg did increase gastric emptying in over half the volunteers tested.

Intravenous infusion of either 1 mg or 5 mg ondansetron tended to increase small bowel transit times and single intravenous doses of 10 mg ondansetron have been reported to decrease sphincter pressure in the lower oesophagus in some subjects.

In psychomotor testing ondansetron does not impair performance nor cause sedation.

MICROBIOLOGY

Not applicable.

TOXICOLOGY

Acute Toxicity:

Single doses of ondansetron up to the LD₅₀ in mice and in rats were generally well tolerated. Reactions, including tremor and convulsive behaviour, occurred only at near lethal levels.

Species	LD50 (mg/kg)	
	Oral	IV
Mice	10-30	1.0-2.5
Rats	100-150	15-20

All deaths resulted from the acute effects of treatment, the observed clinical signs being consistent with the central nervous system effects associated with behavioural depression. These effects were not associated with any apparent histopathological changes in the brain. No target organ toxicity was identified.

Long term Toxicity:

Subacute Toxicity Studies

Species	Route	Dose (mg/kg/day)	Duration of Study	Results
Rats	Oral	160	7 weeks	Well tolerated
	IV	12	5 weeks	Well tolerated
Dogs	Oral	7.5-25	5 weeks	Transient post-dosing clinical reactions associated with behavioural depression (at highest dose levels)
	IV	2-8	5 weeks	

Maximum daily dose levels in rats were found to be higher when doses were gradually increased. Identical doses were rapidly lethal to rats not previously exposed to ondansetron. Post-dosing reactions, in both rats and dogs, included ataxia, exophthalmia, mydriasis, tremor and respiratory changes. Increases in liver enzymes (SGPT and SGOT) were noted at high dose levels. Dogs dosed at 6.75 mg/kg/day intravenously exhibited vein irritancy in the form of constriction and thickening, creating resistance to needle

penetration. The changes were noted after seven days treatment but were reversed by decreasing the dose concentration.

Chronic Toxicity

Species	Duration	Max. no-effect Dose (mg/kg/day)	Effects
Rat	18 months	1	Usually transient and restricted to highest dose
Dog	12 months	12	

Carcinogenicity Studies

Species	Route	Dose (mg/kg/day)	Duration of Study	Results
Mice	Oral	1-40 (max. oral dose 30)	2 years	No treatment related increases in tumour incidence.
Rats	Oral	1-25 (max. oral dose 150)	2 years	Proportion of benign/malignant tumours also remained Consistent with the pathological background of the Animals studied.

There was no evidence of a tumourigenic effect of ondansetron in any tissue.

Mutagenicity Studies:

No evidence of mutagenicity was observed in microbial mutagen tests using mutant strains of *Salmonella typhimurium*, *Escherichia coli* or *Saccharomyces cerevisiae*, with or without a rat-liver post-mitochondrial metabolizing system.

There was also no evidence of damage to genetic material noted in in-vitro V-79 mammalian cell mutation studies, *in vitro* chromosome aberration tests using human peripheral lymphocytes, or *in vivo* chromosome aberration assays in mouse bone marrow.

Reproduction and Teratology:

Ondansetron was not teratogenic in rats and rabbits at dosages up to the maximum nonconvulsive level, (rat: 15 mg/kg/day, rabbit: 30 mg/kg/day). No adverse effects on pregnancy or foetal and post-natal development were detected in rats and no foetal abnormalities were observed in rabbits after oral administration of ondansetron.

A slight maternal toxicity was observed at the highest dose level in intravenous organogenesis (4.0 mg/kg/day) studies in the rabbit. Effects included maternal body weight loss and increased incidence of early foetal death. In a rat fertility study, there was a dose-related decrease in the proportion of surviving pups of the F2 generation; however, the significance of this is unclear.

Administration of ondansetron to pregnant rats and rabbits, indicated there was foetal exposure to low levels of ondansetron and its metabolites. Ondansetron is retained in the foetal eye presumably bound to melanin. In rats, the transfer of ondansetron and its metabolites into breast milk was extensive. The concentration of unchanged ondansetron in breast milk was higher than in corresponding plasma samples.

Daily administration of ondansetron at dosages up to 15 mg/kg/day to pregnant rats from day 17 of pregnancy to litter day 22 had no effects on pregnancy of the parental generation or on post-natal development and mating of the F1 generation. Foetal development of the F2 generation was comparable to controls; however, the number of implantations and viable foetuses was reduced in the highest dosage group when compared with controls.

REFERENCES

1. Blackwell CP, Harding SM. The clinical pharmacology of Ondansetron. *Eur J Cancer Clin Oncol* 1989; 25(Suppl. 1):S21-S24.
2. Bowman A, Allan SG, Warrington PS, Whelan JM, Smyth JM. Clinical trials and pharmacokinetics of Ondansetron Hydrochloride, a new antiemetic effective against platinum-induced vomiting. *Proceedings of the European Conference of Clinical Oncologists* 1987; 1063.
3. Butler A, Hill JM, Ireland SJ, Jordan CC, Tyers MB. Pharmacological properties of Ondansetron Hydrochloride, a novel antagonist of 5-HT₃ receptors. *Br J Pharmacol* 1988; 94:397-412.
4. Costall B, Naylor RJ, Tyers MB. Recent advances in the neuropharmacology of 5-HT₃ agonists and antagonists. *Reviews in Neurosciences* 1988; 2:41-65.

5. Craig JB, Powell BL: Review. The management of nausea and vomiting in clinical oncology. *Am J Med Sci* 1987; 293:34-44.
6. Cunningham D, Hawthorn J, Pople A, Gazet J-C, Ford HT, Challoner T, Coombes RC. Prevention of emesis in patients receiving cytotoxic drugs by Ondansetron Hydrochloride, a selective 5-HT₃ receptor antagonist. *Lancet* 1987; i:1461-1463.
7. Cunningham D, Turner A, Hawthorn J, Rosin RD. Ondansetron with and without dexamethasone to treat chemotherapy-induced emesis. *Lancet* 1989; i:1323.
8. Green JA, Watkin SW, Hammond P, Griggs J, Challoner T. The efficacy and safety of Ondansetron Hydrochloride in the prophylaxis of ifosfamide- induced nausea and vomiting. *Cancer Chemother Pharmacol* 1989; 24:137-139.
9. Hawthorn J, Cunningham D. Dexamethasone can potentiate the anti-emetic action of a 5HT₃ receptor antagonist on cyclophosphamide induced vomiting in the ferret. *Br J Cancer* 1990; 61(1):56-60.
10. Higgins GA, Kilpatrick GT, Bunce KT, Jones BJ, Tyers MB. 5-HT₃ antagonists injected into the area postrema inhibit cisplatin-induced emesis in the ferret. *Br J Pharmacol* 1989; 97:247-255.
11. Kris MG, Gralla RJ, Clark RA, Tyson LB. Dose-ranging evaluation of serotonin antagonist GR-507/75 (Ondansetron Hydrochloride) when used as an anti-emetic in patients receiving anti-cancer chemotherapy. *J Clin Oncol* 1988; 6:659-662.
12. Kris MG, Gralla RJ, Clark RA, Tyson LB. Phase II trials of the serotonin antagonist GR38032F for the control of vomiting caused by cisplatin. *J Natl Cancer Inst* 1989; 81(1):42-46.

13. Marty M, Droz JP, Pouillart P, Paule B, Brion N, Bons J. Ondansetron Hydrochloride, a 5-HT₃ receptor antagonist, in the prophylaxis of acute cisplatin-induced nausea and vomiting. *Cancer Chemother Pharmacol* 1989; 23:389-391.
14. Priestman T, Challoner T, Butcher M, Priestman S. Control of radiation-induced emesis with Ondansetron Hydrochloride. *Proc Am Soc Clin Oncol* 1988; 7:1089.
15. Priestman TJ. Clinical studies with Ondansetron in the control of radiation-induced emesis. *Eur J Cancer Clin Oncol* 1989; 25(Suppl):S29-S33.
16. Schmoll HJ. The role of Ondansetron in the treatment of emesis induced by noncisplatin-containing chemotherapy regimens. *Eur J Cancer Clin Oncol* 1989; 25(Suppl. 1):S35-S39.
17. Smith DB, Newlands ES, Spruyt OW, Begent RHJ, Rustin GJS, Mellor B, Bagshawe KD. Ondansetron plus dexamethasone: Effective anti-emetic prophylaxis for patients receiving cytotoxic chemotherapy. *Br J Cancer* 1990; 61(2):323-324.
18. Stables R, Andrews PLR, Bailey HE, Costall B, Gunning SJ, Hawthorn J, Naylor RJ, Tyers MB. Antiemetic properties of the 5HT₃-receptor antagonist Ondansetron Hydrochloride. *Cancer Treatment Rev.* 1987; 14:333-336.
19. Tyers MB, Bunce KT, Humphrey PPA. Pharmacological and anti-emetic properties of Ondansetron. *Eur J Cancer Clin Oncol* 1989; 25(Suppl. 1):S15-S19.
20. Van Liessum P, de Mulder P, Kaasa S, Lane-Allman E, Seynaeve C, Verwij J: Ondansetron Hydrochloride in the prophylaxis of nausea and vomiting induced by cisplatin. *Proc European Soc Clin Oncol* 1988; 13:267.
21. Product Monograph, ZOFTRAN®, GlaxoSmithKline Inc., Mississauga, Ontario L5N 6L4, Date of Revision: March 20, 2007.
22. Bioavailability Study, on file.

IMPORTANT : PLEASE READ

PART III: CONSUMER INFORMATION

RIVA-ONDANSETRON (Ondansetron hydrochloride)

This leaflet is part III of a three-part "Product Monograph" published when RIVA-ONDANSETRON was approved for sale in Canada and is designed specifically for Consumers. This leaflet is a summary and will not tell you everything about RIVA-ONDANSETRON. Contact your doctor or pharmacist if you have any questions about the drug.

RIVA-ONDANSETRON can only be obtained with a prescription from your doctor.

ABOUT THIS MEDICATION

What the medication is used for:

The name of your medicine is RIVA-ONDANSETRON Tablets (Ondansetron hydrochloride). This medicine is one of a group called antiemetics.

RIVA-ONDANSETRON is used for:

- the prevention of nausea (feeling of sickness) and vomiting associated with emetogenic chemotherapy, and radiotherapy.
- the prevention and treatment of post-operative nausea and vomiting.

What it does:

Treatments such as general anaesthesia, cancer chemotherapy and radiotherapy are thought to cause the release of a natural substance (serotonin), which can cause you to feel sick and to vomit. RIVA-ONDANSETRON helps to stop this from happening, thus preventing you from vomiting or feeling sick.

When it should not be used:

Do not take RIVA-ONDANSETRON if:

- you have a history of hypersensitivity (an allergic reaction) to any ingredient in RIVA-ONDANSETRON
- you are pregnant, or likely to become pregnant, or if you are breast feeding a baby. However, there may be circumstances when your doctor advises you to use this medicine during pregnancy.

What the medicinal ingredients:

RIVA-ONDANSETRON tablets contain ondansetron hydrochloride as the medicinal ingredient.

What the important non-medicinal ingredients are:

RIVA-ONDANSETRON Tablets contain the following non-medicinal ingredients: lactose, microcrystalline cellulose, pregelatinized starch, magnesium stearate, methyl hydroxypropyl cellulose, triacetin, iron oxide yellow and titanium dioxide.

What dosage forms it comes in:

RIVA-ONDANSETRON tablets are supplied in two strengths, one contains 4 milligrams of ondansetron and the other contains 8 milligrams of ondansetron. Your doctor will decide which strength you need.

WARNINGS AND PRECAUTIONS:

BEFORE you use RIVA-ONDANSETRON talk to your doctor or pharmacist if:

- you have a history of hypersensitivity (an allergic reaction) to any ingredient in RIVA-ONDANSETRON
- you are pregnant or likely to become pregnant.
- you are breast feeding a baby.
- you have liver problems.
- you have signs of intestinal obstruction.
- you have a history of heart problems

If you experience wheezing and tightness of the chest, heart throbbing, swelling of eyelids, face or lips, or develop a skin rash, skin lumps or hives, **contact your doctor immediately.**

Do not take anymore medicine unless your doctor tells you to do so.

INTERACTIONS WITH THIS MEDICATION

It is important that your doctor knows about all your medication so that you get the best possible treatment. Tell your doctor about all the medicines you are taking including those you have bought yourself. If you are taking any medicines containing tramadol (such as TRAMACET), RIVA-ONDANSETRON may decrease its effectiveness.

PROPER USE OF THIS MEDICATION

The label on the container of your medicine should tell you how often to take your medicine and how many doses you should take each time. If not, or if you are not sure, consult your doctor or pharmacist.

Do not take more doses, or take them more often than your doctor prescribes. If, however, you vomit within one hour of taking your medicine, you should take the same amount of medicine again. If vomiting persists, consult your doctor.

Usual dose:

Chemotherapy Induced Nausea and Vomiting

You will receive RIVA-ONDANSETRON prior to chemotherapy. Based on how likely you are to experience nausea and/or vomiting, caused by your cancer treatment, your doctor will tell you the amount you need to take and how frequently.

Adult: The dose of Riva-Ondansetron will be between 8 and 32mg a day depending on the potential of your chemotherapy treatment to cause you to vomit and/or have nausea.

Children (4 to 12 years): After chemotherapy, take 4 mg orally every 8 hours for up to 5 days.

Radiotherapy Induced Nausea and Vomiting

Adult: Take 8 mg orally 1 to 2 hours before radiotherapy. After therapy, take 8 mg orally every 8 hours for up to 5 days after a course of treatment.

Prevention of Post-Operative Nausea and Vomiting

Adult: Take 16 mg orally one hour before anaesthesia.

If you have a liver problem, your dose may be altered. Please follow the instructions of your doctor.

Overdose:

In the event you accidentally take more doses than prescribed, immediately contact your doctor or hospital emergency department or nearest poison control centre.

Missed Dose:

If you miss a dose and do not feel sick, take the next dose when it is due.

If you forget to take your medicine and feel sick or vomit, take a dose as soon as possible.

If your doctor decides to stop the treatment, do not keep any leftover medicine unless your doctor tells you to.

SIDE EFFECTS AND WHAT TO DO ABOUT THEM

You may experience headaches, a feeling of warmth, flushing or constipation, while taking RIVA-ONDANSETRON. There is no need to stop taking your medicine, but you should tell your doctor about these symptoms at your next visit.

If your nausea (feeling of sickness) or vomiting do not improve while taking RIVA-ONDANSETRON, consult your doctor for further advice.

If you feel unwell or have any symptoms that you do not understand, you should contact your doctor immediately.

SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM

Frequency	Side Effect/Symptom	Talk with your Doctor or Pharmacist	Stop taking drug and call your doctor or pharmacist immediately
Uncommon	Heart problems such as fast/slow heart beat, chest pain	X	X
	Seizures Upward rolling of the eyes, abnormal muscular stiffness/body movements/shaking.		X
Rare	Eye problems such as blurred vision	X	X
	Immediate allergic reaction and symptoms such as swelling of the mouth, throat, difficulty in breathing, rash, hives, increased heart rate		
Very Rare	Eye problems such as temporary blindness	X	

This is not a complete list of side effects. For any unexpected effects while taking RIVA-ONDANSETRON, contact your doctor or pharmacist.

HOW TO STORE IT

Keep your medicine in a safe place where children cannot reach it. Your medicine may harm them.

Your RIVA-ONDANSETRON Tablets should be kept at room temperature (15-30°C) in a dry place.

Reporting Suspected Side Effects

To monitor drug safety, Health Canada through the Canada Vigilance Program collects information on serious and unexpected effects of drugs. If you suspect you have had a serious or unexpected reaction to this drug you may notify Health Canada Vigilance:

By toll-free telephone: 866-234-2345

Toll-free fax: 866-678-6789

On-line: www.healthcanada.gc.ca/medeffect

By email: CanadaVigilance@hc-sc.gc.ca

By regular mail:

Canada Vigilance National Office
Marketed Health Products Safety and Effectiveness
Information Division

Marketed Health products Directorate

Health Products and Food Branch

Health Canada

Tunney's Pasture, AL 0701 C

Ottawa ON K1A 0K9

NOTE: Before contacting Canada Vigilance, you should contact your physician or pharmacist.

MORE INFORMATION

Remember: This medicine is for you. Only a doctor can prescribe it for you. Never give it to someone else. It may harm them even if their symptoms are the same as yours.

This leaflet does not contain the complete information about your medicine. If any questions remain unanswered or you are not sure about something, you should ask your doctor or pharmacist.

You may want to read this leaflet again. **Please Do not Throw It Away** until you have finished your medicine.

This document plus the full product monograph, prepared for health professionals can be obtained by contacting the sponsor,

**Laboratoire Riva Inc.,
660 Boul. Industriel,
Blainville, Québec
J7C 3V4
Telephone: 450-434-7482
Fax: 450-434-2500**

Last Revised: July 2011