

## PRODUCT MONOGRAPH

### <sup>Pr</sup>**ratio-TOPIRAMATE**

(topiramate)

25 mg, 100 mg and 200 mg Tablets

Antiepileptic/Migraine Prophylaxis

Teva Canada Limited  
30 Novopharm Court  
Toronto, Ontario  
Canada M1B 2K9  
[www.tevacanada.com](http://www.tevacanada.com)

Date of Preparation:  
July 16, 2012

Control No: 156786

**Table of Contents**

**PART I: HEALTH PROFESSIONAL INFORMATION.....3**  
SUMMARY PRODUCT INFORMATION .....3  
INDICATIONS AND CLINICAL USE.....3  
CONTRAINDICATIONS .....4  
WARNINGS AND PRECAUTIONS.....17  
ADVERSE REACTIONS.....17  
DRUG INTERACTIONS .....33  
DOSAGE AND ADMINISTRATION .....41  
ACTION AND CLINICAL PHARMACOLOGY .....44  
STORAGE AND STABILITY.....47  
DOSAGE FORMS, COMPOSITION AND PACKAGING .....47

**PART II: SCIENTIFIC INFORMATION .....49**  
PHARMACEUTICAL INFORMATION.....49  
CLINICAL TRIALS.....52  
DETAILED PHARMACOLOGY .....57  
TOXICOLOGY .....59  
REFERENCES .....64

**PART III: CONSUMER INFORMATION.....69**

**Pr ratio-TOPIRAMATE**

topiramate  
25 mg, 100 mg and 200 mg Tablets

Antiepileptic/Migraine Prophylaxis

**PART I: HEALTH PROFESSIONAL INFORMATION**

**SUMMARY PRODUCT INFORMATION**

<b>Route of Administration</b>	<b>Dosage Form / Strength</b>	<b>All Nonmedicinal Ingredients</b>
oral	tablet 25 mg, 100 mg and 200 mg	pregelatinized starch, lactose monohydrate, colloidal silicon dioxide, sodium starch glycolate, microcrystalline cellulose and magnesium stearate, hydroxypropyl methylcellulose, titanium dioxide, polydextrose, triacetin and polyethylene glycol. The colourants that are present in the tablets are as follows: 100 mg: yellow iron oxide and 200 mg: red iron oxide.

**INDICATIONS AND CLINICAL USE**

**EPILEPSY**

**ratio-TOPIRAMATE** (topiramate) is indicated as monotherapy for the management of patients (adults and children six years and older) with newly diagnosed epilepsy.

**ratio-TOPIRAMATE** (topiramate) is also indicated as adjunctive therapy for the management of patients (adults and children two years and older) with epilepsy who are not satisfactorily controlled with conventional therapy.

**MIGRAINE PROPHYLAXIS**

**ratio-TOPIRAMATE** (topiramate) is indicated in adults for the prophylaxis of migraine headache. Prophylactic treatment of migraine may be considered in situations such as: adults experiencing four or more migraine attacks per month who fail to respond adequately to acute abortive therapy; recurring attacks that significantly interfere with the patient's daily routine; a pattern of increasing migraine attacks over time, with the risk of developing rebound headache from acute abortive therapies; or failure of, or contraindication to, or troublesome side effects from acute abortive medications. Continuing therapy should be reviewed every six months.

**ratio-TOPIRAMATE** (topiramate) should not be used in the acute treatment of migraine attacks. Safety and efficacy of topiramate in the management or prevention of cluster headache,

hemiplegic, basilar, ophthalmoplegic, or transformed migraine headaches have not been established.

**Geriatrics (> 65 years of age):**

There is limited information in patients over 65 years of age (see **WARNINGS AND PRECAUTIONS, Special Populations, Geriatrics**).

**Pediatrics (< 2 years of age):**

**ratio-TOPIRAMATE** is not indicated in children under two years of age (see **WARNINGS AND PRECAUTIONS, Special Populations, Pediatrics**).

**CONTRAINDICATIONS**

Patients who are hypersensitive to this drug or to any nonmedicinal ingredient in the formulation or component of the container. For a complete listing, see the **DOSAGE FORMS, COMPOSITION AND PACKAGING** section of the Product Monograph.

**ratio-TOPIRAMATE** for the indication prophylaxis of migraine is contraindicated in pregnancy and in women of childbearing potential who are not using an effective method of contraception (see **WARNINGS AND PRECAUTIONS, Special Populations, Pregnancy Registries to Data; MIGRAINE PROPHYLAXIS; and DOSAGE AND ADMINISTRATION, Dosing Considerations**).

**WARNINGS AND PRECAUTIONS**

**General**

Antiepileptic drugs, including **ratio-TOPIRAMATE** (topiramate), should be withdrawn gradually to minimize the potential for seizures or increased seizure frequency. In clinical trials in adult patients with epilepsy, dosages were decreased by 50-100 mg/day at weekly intervals. In clinical trials of children, topiramate was gradually withdrawn over a 2-8 week period. (See **DOSAGE AND ADMINISTRATION, General and Epilepsy**).

In patients without a history of seizures or epilepsy, **ratio-TOPIRAMATE** (topiramate) should be gradually withdrawn to minimize the potential for seizures or increased seizure frequency. In clinical trials in adult patients receiving topiramate for migraine prophylaxis dosages were decreased by 25-50 mg/day at weekly intervals. (See **DOSAGE AND ADMINISTRATION, General and Migraine Propylaxis**).

In situations where rapid withdrawal of **ratio-TOPIRAMATE** (topiramate) is medically required, appropriate monitoring is recommended. (See **DOSAGE AND ADMINISTRATION, General**).

**Hyperammonemia and Encephalopathy**

Topiramate treatment has produced hyperammonemia (in some instances, dose-related) in clinical investigational programs of adolescents (12–16 years) who were treated with topiramate monotherapy for migraine prophylaxis (incidence above the upper limit of normal, placebo: 22%; 50 mg/day: 26%; 100 mg/day: 41%). Pediatric patients under two years of age who were treated with adjunctive topiramate for partial onset epilepsy, also experienced hyperammonemia (placebo: 8%; 5 mg/kg/day: 10%; 15 mg/kg/day: 0%; 25 mg/kg/day: 9%). **ratio-TOPIRAMATE** is not indicated for migraine prophylaxis in patients under 18 years of age. **ratio-TOPIRAMATE** is also not indicated for any use in patients under two years of age (see **INDICATIONS, Pediatrics**).

In some patients, ammonia was markedly increased (> 50% above upper limit of normal). In adolescent patients, the incidence of markedly increased hyperammonemia was 6% for placebo, 6% for 50 mg, and 12% for 100 mg topiramate daily. The hyperammonemia associated with topiramate treatment occurred with and without encephalopathy in placebo-controlled trials and in an open-label, extension trial. Dose-related hyperammonemia was also observed in the extension trial in pediatric patients up to two years old.

Hyperammonemia with and without encephalopathy has also been observed in post-marketing reports in adult patients who were taking topiramate alone or in combination with valproic acid.

#### **Hyperammonemia/Encephalopathy with Concomitant Valproic Acid (VPA)**

Based upon post-marketing reports, concomitant administration of topiramate and valproic acid (VPA) has been associated with hyperammonemia with or without encephalopathy in patients who have tolerated either drug alone. Although hyperammonemia may be asymptomatic, clinical symptoms of hyperammonemic encephalopathy often include acute alterations in level of consciousness and/or cognitive function with lethargy or vomiting. In most cases, symptoms and signs abated with discontinuation of either drug. This adverse reaction is not due to a pharmacokinetic interaction.

Although **ratio-TOPIRAMATE** is not indicated for use in patients under two years of age, in an investigational trial in this population, VPA clearly produced a dose-related increase in the incidence of treatment-emergent hyperammonemia (above the upper limit of normal, placebo: 0%, 5 mg/kg/day: 12%, 15 mg/kg/day: 7%, 25 mg/kg/day: 17%). Markedly increased, dose-related hyperammonemia also occurred in these patients (placebo: 0%; 5 mg/kg/day: 0%, 15 mg/kg/day: 7%, 25 mg/kg/day: 8%). Dose-related hyperammonemia was similarly observed in a long-term extension trial in these very young, pediatric patients.

#### **Monitoring for Hyperammonemia**

Patients with inborn errors of metabolism or reduced hepatic mitochondrial activity may be at an increased risk for hyperammonemia with or without encephalopathy. Although not studied, topiramate treatment or an interaction of concomitant topiramate and valproic acid treatment may exacerbate existing defects or unmask deficiencies in susceptible persons.

Clinical symptoms of hyperammonemic encephalopathy often include acute alterations in level of consciousness and/or cognitive function with lethargy or vomiting. Therefore, in patients who develop unexplained vomiting, lethargy, confusion or other changes in mental status, hyperammonemic encephalopathy should be considered and serum ammonia levels should be measured (see **ADVERSE REACTIONS, Post-Market Adverse Drug Reactions** and **DRUG**

## **INTERACTIONS, Drug-Drug Interactions).**

### **Hypothermia with Concomitant Valproic Acid (VPA) Use**

Hypothermia, defined as an unintentional drop in body core temperature to  $< 35^{\circ}\text{C}$  ( $95^{\circ}\text{F}$ ), has been reported in association with topiramate use with concomitant valproic acid (VPA) both in conjunction with hyperammonemia and in the absence of hyperammonemia. This adverse reaction in patients using concomitant topiramate and valproate can occur after starting topiramate treatment or after increasing the daily dose of topiramate (see **DRUG INTERACTIONS, Valproic Acid**). Consideration should be given to stopping topiramate or valproate in patients who develop hypothermia, which may be manifested by a variety of clinical abnormalities including lethargy, confusion, coma, and significant alterations in other major organ systems such as the cardiovascular and respiratory systems. Clinical management and assessment should include examination of blood ammonia levels.

### **Carcinogenesis and Mutagenesis**

See *Product Monograph Part II: TOXICOLOGY*, **Carcinogenicity** and **Mutagenicity** for discussion on animal data.

### **Endocrine and Metabolism**

#### **Oligohidrosis and Hyperthermia**

Oligohidrosis (decreased sweating) and hyperthermia, infrequently resulting in hospitalization, including fatalities, have been reported in patients treated with topiramate. Oligohidrosis and hyperthermia may have potentially serious sequelae and may be preventable by prompt recognition of symptoms and appropriate treatment. Decreased sweating and elevation of body temperature above normal characterized the cases reported in patients treated with topiramate. Some of the cases were reported after exposure to elevated environmental temperatures.

These reports have primarily involved children. Patients treated with topiramate, especially pediatric patients, should be monitored closely for evidence of decreased sweating and increased body temperature, particularly in hot weather. Proper hydration before and during activities such as exercise or exposure to warm temperatures is recommended.

Caution should be used when topiramate is prescribed with other drugs that predispose patients to heat-related disorders; these drugs include, but are not limited to, other carbonic anhydrase inhibitors and drugs with anticholinergic activity. (See **ADVERSE REACTIONS, Post-Market Adverse Drug Reactions**).

#### **Metabolic Acidosis**

Hyperchloremic, non-anion gap, metabolic acidosis (i.e. decreased serum bicarbonate below the normal reference range in the absence of respiratory alkalosis) is associated with topiramate treatment. This decrease in serum bicarbonate is due to the inhibitory effect of topiramate on renal carbonic anhydrase. Generally, the decrease in bicarbonate occurs early in treatment although it can occur at any time during treatment. These decreases are usually mild to moderate (average decrease of 4 mmol/L at doses of 100 mg/day or above in adults and at approximately 6 mg/kg/day in pediatric patients). Rarely, patients have experienced decreases to values below 10 mmol/L. Conditions or therapies that predispose to acidosis (such as renal disease, severe respiratory disorders, status epilepticus, diarrhea, surgery, ketogenic diet, or certain drugs) may be additive to the bicarbonate lowering effects of topiramate.

In patients > 16 years of age, the incidence of persistent treatment-emergent decreases in serum bicarbonate (levels of <20 mmol/L at two consecutive visits or at the final visit) in controlled clinical trials for adjunctive treatment of epilepsy was 32% for 400 mg/day, and 1% for placebo. Metabolic acidosis has been observed at doses as low as 50 mg/day. The incidence of a markedly abnormally low serum bicarbonate (i.e., absolute value <17 mmol/L and >5 mmol/L decrease from pretreatment) in these trials was 3% for 400 mg/day, and 0% for placebo. In the monotherapy trial, the incidence was 1% for 50 mg/day and 7% for 400 mg/day. Serum bicarbonate levels have not been systematically evaluated at daily doses greater than 400 mg/day.

In pediatric patients two to 16 years of age, the incidence of persistent treatment-emergent decreases in serum bicarbonate in placebo-controlled trials for adjunctive treatment of Lennox-Gastaut Syndrome or refractory partial onset seizures was 67% for topiramate (at approximately 6 mg/kg/day), and 10% for placebo. The incidence of a markedly abnormally low serum bicarbonate (i.e., absolute value <17 mmol/L and >5 mmol/L decrease from pretreatment) in these trials was 11% for topiramate and 0% for placebo.

The incidence of persistent treatment-emergent decreases in serum bicarbonate in placebo-controlled trials for adults for prophylaxis of migraine was 44% for 200 mg/day, 39% for 100 mg/day, 23% for 50 mg/day, and 7% for placebo. The incidence of a markedly abnormally low serum bicarbonate (i.e., absolute value <17 mmol/L and >5 mmol/L decrease from pretreatment) in these trials was 11% for 200 mg/day, 9% for 100 mg/day, 2% for 50 mg/day, and <1% for placebo.

Although not approved for use in patients under two years of age for any indication (see **INDICATIONS**), a controlled trial that examined this population revealed that topiramate produced a metabolic acidosis that was notably greater in magnitude than that observed in controlled trials in older children and adults. The mean treatment difference (25 mg/kg/day topiramate-placebo) was -5.9 mEq/L for bicarbonate. The incidence of metabolic acidosis (defined by a serum bicarbonate < 20 mEq/L) was 0% for placebo, 30% for 5 mg/kg/day, 50% for 15 mg/kg/day, and 45% for 25 mg/kg/day. The incidence of markedly abnormal changes (i.e., < 17 mEq/L and > 5 mEq/L decrease from baseline of > 20 mEq/L) was 0% for placebo, 4% for 5 mg/kg/day, 5% for 15 mg/kg/day, and 5% for 25 mg/kg/day.

Cases of moderately severe metabolic acidosis have been reported in patients as young as five months old, especially at daily doses above 5 mg/kg/day.

In pediatric patients, six to 15 years of age, the incidence of persistent treatment-emergent decreases in serum bicarbonate in the epilepsy controlled clinical trial for monotherapy was 9% for 50 mg/day and 25% for 400 mg/day. The incidence of a markedly abnormally low serum bicarbonate (i.e., absolute value < 17 mEq/L and > 5 mEq/L decrease from pretreatment) in this trial was 1% for 50 mg/day and 6% for 400 mg/day.

In patients ≥16 years of age, the incidence of persistent treatment-emergent decreases in serum bicarbonate in the epilepsy controlled clinical trial for monotherapy was 14% for 50 mg/day and 25% for 400 mg/day. The incidence of a markedly abnormally low serum bicarbonate (i.e.,

absolute value < 17 mEq/L and > 5 mEq/L decrease from pretreatment) in this trial for adults was 1% for 50 mg/day and 6% for 400 mg/day.

Some manifestations of acute or chronic metabolic acidosis may include hyperventilation, nonspecific symptoms such as fatigue and anorexia, or more severe sequelae including cardiac arrhythmias or stupor. Chronic, untreated metabolic acidosis may increase the risk for nephrolithiasis or nephrocalcinosis, and may also result in osteomalacia (referred to as rickets in pediatric patients) and/or osteoporosis with an increased risk for fractures. Chronic metabolic acidosis in pediatric patients may also reduce growth rates. A reduction in growth rate may eventually decrease the maximal height achieved. The effect of topiramate on growth and bone-related sequelae has not been systematically investigated in long-term, placebo-controlled trials. Long-term, open-label treatment of infants/toddlers, with intractable partial epilepsy, for up to one year, showed reductions from baseline in Z SCORES for length, weight, and head circumference compared to age and sex-matched normative data, although these patients with epilepsy are likely to have different growth rates than normal infants. Reductions in Z SCORES for length and weight were correlated to the degree of acidosis. Topiramate treatment that causes metabolic acidosis during pregnancy can possibly produce adverse effects on the fetus and might also cause metabolic acidosis in the neonate from possible transfer of topiramate to the fetus.

Measurement of baseline and periodic serum bicarbonate during topiramate treatment is recommended. If metabolic acidosis develops and persists, consideration should be given to reducing the dose or discontinuing topiramate (using dose tapering). If the decision is made to continue patients on topiramate in the face of persistent acidosis, alkali treatment should be considered.

### **Decreases in Serum Potassium with Concomitant Treatment with Hydrochlorothiazide (HCTZ)**

In a drug interaction study, a greater decrease from baseline in serum potassium values was seen with concomitant treatment than for either drug alone. At the end of each treatment period, 27% (3/11) of subjects on topiramate treatment alone and 25% (3/12) of subjects on HCTZ treatment alone showed a serum potassium value of <3.6 mEq/L, compared to 61% (14/23) of subjects on concomitant drug treatment. One of the subjects who had hypokalemia with concomitant treatment also had an abnormal ECG (nonspecific ST-T wave changes), which may have been related to the decrease in plasma potassium levels. Caution should be used when treating patients who are receiving topiramate and hydrochlorothiazide concomitantly. (See **DRUG INTERACTIONS.**)

### **Nutritional Supplementation**

A dietary supplement or increased food intake may be considered if the patient is losing weight while on this medication.

### **Hepatic/Biliary/Pancreatic Decreased Hepatic Function**

In hepatically impaired patients, topiramate should be administered with caution as the clearance of topiramate was decreased compared with normal subjects.

### **Neurologic**

#### **Central Nervous System Effects**



Adverse events most often associated with the use of topiramate were central nervous system related and were observed in both the epilepsy and migraine populations. In adults, the most significant of these can be classified into three general categories:

- i) psychomotor slowing, difficulty with concentration and speech or language problems, in particular, word-finding difficulties,
- ii) somnolence or fatigue and
- iii) mood disturbances including irritability and depression.

In the controlled epilepsy adjunctive therapy trials, these events were generally mild to moderate, and generally occurred early in therapy. While the incidence of psychomotor slowing does not appear to be dose related, both language problems and difficulty with concentration or attention increased in frequency with increasing dosage in the six double-blind trials, suggesting that these events are dose related (see **ADVERSE REACTIONS, Post-Market Adverse Drug Reactions**).

Central nervous system and psychiatric-related events were also more frequently reported in topiramate-treated subjects in the migraine prophylaxis trials. These included: anorexia, dizziness, difficulty with memory, somnolence, language problems, and difficulty with concentration and attention. Most of the events were mild or moderate in severity, some of which led to withdrawal from treatment (see **ADVERSE REACTIONS, MIGRAINE PROPHYLAXIS**).

Additional non-specific CNS effects occasionally observed with topiramate as add-on epilepsy therapy include dizziness or imbalance, confusion and memory problems. Although the duration of the epilepsy monotherapy studies was considerably longer than the epilepsy adjunctive therapy studies, these adverse events were reported at lower incidences in the monotherapy trials.

### **Paresthesia**

Paresthesia, an effect associated with the use of other carbonic anhydrase inhibitors, appears to be a common effect of topiramate. Paresthesia was more frequently reported in the migraine prophylaxis and epilepsy monotherapy trials versus the adjunctive therapy trials in epilepsy. The higher incidence in the epilepsy monotherapy studies may have been related to the higher topiramate plasma concentrations achieved in the monotherapy studies. In the majority of instances, paresthesia did not lead to treatment discontinuation.

### **Ophthalmologic**

#### **Acute Myopia and Secondary Angle Closure Glaucoma**

A syndrome consisting of acute myopia associated with secondary angle closure glaucoma has been reported in patients receiving topiramate. Symptoms include acute onset of decreased visual acuity and/or ocular pain. Ophthalmologic findings can include myopia, anterior chamber shallowing, ocular hyperemia (redness) and increased intraocular pressure. Mydriasis may or may not be present. This syndrome may be associated with supraciliary effusion resulting in anterior displacement of the lens and iris, with secondary angle closure glaucoma. Symptoms typically occur within a few days to 1 month of initiating topiramate therapy. In contrast to primary narrow angle glaucoma, which is rare under 40 years of age, secondary angle closure glaucoma associated with topiramate has been reported in pediatric patients as well as adults.

The primary treatment to reverse symptoms is discontinuation of topiramate as rapidly as possible, according to the judgment of the treating physician. Other measures, in conjunction with discontinuation of topiramate, may be helpful (see **ADVERSE REACTIONS, Post-Market Adverse Drug Reactions**).

In all cases of acute visual blurring and/or painful/red eye(s), immediate consultation with an ophthalmologist/emergency room is recommended.

Elevated intraocular pressure of any etiology, if left untreated, can lead to serious sequelae including permanent vision loss.

Maculopathy, including visual field defect, has been observed very rarely in post marketing reports (See **ADVERSE REACTIONS, Post-Market Adverse Drug Reactions**).

### **Psychiatric**

#### **Suicidal ideation and behaviour**

Suicidal ideation and behaviour have been reported in patients treated with antiepileptic agents in several indications.

All patients treated with antiepileptic drugs, irrespective of indication, should be monitored for signs of suicidal ideation and behaviour and appropriate treatment should be considered. Patients (and caregivers of patients) should be advised to seek medical advice should signs of suicidal ideation or behaviour emerge.

An FDA meta-analysis of randomized placebo controlled trials, in which antiepileptic drugs were used for various indications, has shown a small increased risk of suicidal ideation and behaviour in patients treated with these drugs. The mechanism of this risk is not known.

There were 43,892 patients treated in the placebo controlled clinical trials that were included in the meta-analysis. Approximately 75% of patients in these clinical trials were treated for indications other than epilepsy and, for the majority of non-epilepsy indications the treatment (antiepileptic drug or placebo) was administered as monotherapy. Patients with epilepsy represented approximately 25% of the total number of patients treated in the placebo controlled clinical trials and, for the majority of epilepsy patients, treatment (antiepileptic drug or placebo) was administered as adjunct to other antiepileptic agents (i.e., patients in both treatment arms were being treated with one or more antiepileptic drug). Therefore, the small increased risk of suicidal ideation and behaviour reported from the meta-analysis (0.43% for patients on antiepileptic drugs compared to 0.24% for patients on placebo) is based largely on patients that received monotherapy treatment (antiepileptic drug or placebo) for non-epilepsy indications. The study design does not allow an estimation of the risk of suicidal ideation and behaviour for patients with epilepsy that are taking antiepileptic drugs, due both to this population being the minority in the study, and the drug-placebo comparison in this population being confounded by the presence of adjunct antiepileptic drug treatment in both arms.

### **Fetal Toxicity**

**ratio-TOPIRAMATE** can cause fetal harm when administered to a pregnant woman. Data from pregnancy registries indicate that infants exposed to topiramate *in utero* have an increased risk for cleft lip and/or cleft palate (oral clefts). When multiple species of pregnant animals received topiramate at clinically relevant doses, structural malformations, including craniofacial defects, and reduced fetal weights occurred in offspring (see **Special Populations, EPILEPSY, Pregnant Women and Pregnancy Registry Data**).

Consider the benefits and the risks of **ratio-TOPIRAMATE** when administering this drug in women of childbearing potential (see **Information for Patients, Fetal Toxicity and Special Populations, EPILEPSY, Women of Childbearing Potential**). **ratio-TOPIRAMATE** should be used during pregnancy only if the potential benefit outweighs the potential risk. If this drug is used during pregnancy, or if the patient becomes pregnant while taking this drug, the patient should be apprised of the potential hazard to a fetus (see **Special Populations, EPILEPSY**).

## **Renal**

### **Kidney Stones**

A total of 32/1,715 (1.9%) of patients exposed to topiramate during its epilepsy adjunctive therapy development reported the occurrence of kidney stones, an incidence about 10 times that expected in a similar, untreated population (M/F ratio: 27/1,092 male; 5/623 female). In double-blind epilepsy monotherapy studies, a total of 8/886 (0.9%) of adults reported the occurrence of kidney stones. In the general population, risk factors for kidney stone formation include gender (male), ages between 20-50 years, prior stone formation, family history of nephrolithiasis, and hypercalciuria. Based on logistic regression analysis of the clinical trial data, no correlation between mean topiramate dosage, duration of topiramate therapy, or age and the occurrence of kidney stones was established; of the risk factors evaluated, only gender (male) showed a correlation with the occurrence of kidney stones. In the pediatric patients studied, there were no kidney stones observed.

Carbonic anhydrase inhibitors, e.g. acetazolamide, promote stone formation by reducing urinary citrate excretion and by increasing urinary pH. Concomitant use of topiramate, a weak carbonic anhydrase inhibitor, with other carbonic anhydrase inhibitors may create a physiological environment that increases the risk of kidney stone formation, and should therefore be avoided (see **DRUG INTERACTIONS, Drug-Drug Interactions**).

Patients, especially those with a predisposition to nephrolithiasis, may have an increased risk of renal stone formation and associated signs and symptoms such as renal colic, renal pain or flank pain. Increased fluid intake increases the urinary output, lowering the concentration of substances involved in stone formation. Therefore, adequate hydration is recommended to reduce this risk. None of the risk factors for nephrolithiasis can reliably predict stone formation during topiramate treatment.

### **Adjustment of Dose in Renal Failure**

The major route of elimination of unchanged topiramate and its metabolites is via the kidney. Renal elimination is dependent on renal function and is independent of age. Patients with impaired renal function ( $CL_{CR} < 70 \text{ mL/min/1.73m}^2$ ) or with end-stage renal disease receiving hemodialysis treatments may take 10 to 15 days to reach steady-state plasma concentrations as compared to 4 to 8 days in patients with normal renal function. As with all patients, the titration

schedule should be guided by clinical outcome (i.e. seizure control, avoidance of side effects) with the knowledge that patients with known renal impairment may require a longer time to reach steady-state at each dose (see **DOSAGE AND ADMINISTRATION, Dosing Considerations**).

### **Information for Patients**

Patients receiving **ratio-TOPIRAMATE** should be given the following instructions by the physician:

#### **Eye Disorders**

Patients taking **ratio-TOPIRAMATE** should be told to seek immediate medical attention if they experience blurred vision, visual disturbances, or periorbital pain (see **WARNINGS AND PRECAUTIONS, Ophthalmologic**).

#### **Oligohydrosis and Hyperthermia**

Patients, especially pediatric patients, treated with **ratio-TOPIRAMATE** should be monitored closely for evidence of decreased sweating and increased body temperature, especially in hot weather (see **WARNINGS AND PRECAUTIONS, Endocrine and Metabolism**).

#### **Metabolic Acidosis**

Patients should be warned about the potential significant risk for metabolic acidosis that may be asymptomatic and may be associated with adverse effects on kidneys (e.g., kidney stones, nephrocalcinosis), bones (e.g., osteoporosis, osteomalacia, and/or rickets in children), and growth (e.g., growth delay/retardation) in pediatric patients, and on the fetus (see **WARNINGS AND PRECAUTIONS, Endocrine and Metabolism, Metabolic Acidosis and Special Populations, EPILEPSY, Pregnant Women; Women of Childbearing Potential**).

#### **Suicidal Behaviour and Ideation**

Patients, their caregivers, and families should be counselled that AEDs, including **ratio-TOPIRAMATE**, may increase the risk of suicidal thoughts and behaviour and should be advised of the need to be alert for the emergence or worsening of the signs and symptoms of depression, any unusual changes in mood or behaviour or the emergence of suicidal thoughts, or behaviour or thoughts about self-harm. Behaviours of concern should be reported immediately to healthcare providers (see **WARNINGS AND PRECAUTIONS, Psychiatric**).

#### **Interference with Cognitive and Motor Performance**

Patients should be warned about the potential for somnolence, dizziness, confusion, difficulty concentrating, or visual effects and should be advised not to drive or operate machinery until they have gained sufficient experience on **ratio-TOPIRAMATE** to gauge whether it adversely affects their mental performance, motor performance, and/or vision (see **WARNINGS AND PRECAUTIONS, Neurologic, Central Nervous System Effects**).

Even when taking **ratio-TOPIRAMATE** or other anticonvulsants, some patients with epilepsy will continue to have unpredictable seizures. Therefore, all patients taking **ratio-TOPIRAMATE** for epilepsy should be told to exercise appropriate caution when engaging in any activities where loss of consciousness could result in serious danger to themselves or those around them (including swimming, driving a car, climbing in high places, etc.). Some patients with refractory epilepsy will need to avoid such activities altogether. Physicians should discuss

the appropriate level of caution with their patients, before patients with epilepsy engage in such activities.

### **Hyperammonemia and Encephalopathy**

Patients should be warned about the possible development of hyperammonemia with or without encephalopathy. Although hyperammonemia may be asymptomatic, clinical symptoms of hyperammonemic encephalopathy often include acute alterations in level of consciousness and/or cognitive function with lethargy or vomiting. This hyperammonemia and encephalopathy can develop with **ratio-TOPIRAMATE** treatment alone or with **ratio-TOPIRAMATE** treatment with concomitant valproic acid (VPA).

Patients should be instructed to contact their physician if they develop unexplained lethargy, vomiting, or changes in mental status (see **WARNINGS AND PRECAUTIONS, Hyperammonemia and Encephalopathy**).

### **Kidney Stones**

Patients, particularly those with predisposing factors, should be instructed to maintain an adequate fluid intake in order to minimize the risk of kidney stone formation (see **WARNINGS AND PRECAUTIONS, Renal, Kidney Stones**).

### **Fetal Toxicity**

Inform pregnant women and women of childbearing potential that use of **ratio-TOPIRAMATE** during pregnancy can cause fetal harm, including an increased risk for cleft lip and/or cleft palate (oral clefts), which occur early in pregnancy before many women know they are pregnant. There may also be risks to the fetus from chronic metabolic acidosis with use of **ratio-TOPIRAMATE** during pregnancy (see **WARNINGS AND PRECAUTIONS, Fetal Toxicity and Special Populations, EPILEPSY, Pregnant Women**). When appropriate, prescribers should counsel pregnant women and women of childbearing potential about alternative therapeutic options.

Prescribers should advise women of childbearing potential who are not planning a pregnancy to use effective contraception while using **ratio-TOPIRAMATE**, keeping in mind that there is a potential for decreased contraceptive efficacy when using estrogen-containing birth control with topiramate (see **DRUG INTERACTIONS, Other Drug Interactions, Oral Contraceptives**).

Patients should be encouraged to enroll in the North American Antiepileptic Drug (NAAED) Pregnancy Registry if they become pregnant. This registry is collecting information about the safety of antiepileptic drugs during pregnancy. To enroll, patients can call the toll free number, 1- 888-233-2334. Information on the registry can also be found at the website <http://www.massgeneral.org/aed/>.

### **MIGRAINE PROPHYLAXIS**

Prophylactic treatment of migraine: Taking **ratio-TOPIRAMATE** to prevent migraine attacks does not outweigh the risk of malformations to the fetus. Consequently, **ratio-TOPIRAMATE** is contraindicated in pregnancy and in women of childbearing potential who are not using an effective method of contraception (see **CONTRAINDICATIONS; WARNINGS AND PRECAUTIONS, Special Populations, EPILEPSY, Pregnant Women and MIGRAINE PROPHYLAXIS; and DOSAGE AND ADMINISTRATION, Dosing Considerations**).

## Special Populations

### EPILEPSY

#### Pregnant Women

**ratio-TOPIRAMATE can cause fetal harm when administered to a pregnant woman. Data from pregnancy registries indicate that infants exposed to topiramate in utero have an increased risk for cleft lip and/or cleft palate (oral clefts). When multiple species of pregnant animals received topiramate at clinically relevant doses, structural malformations, including craniofacial defects, and reduced fetal weights occurred in offspring. ratio-TOPIRAMATE should be used during pregnancy only if the potential benefit outweighs the potential risk. If this drug is used during pregnancy, or if the patient becomes pregnant while taking this drug, the patient should be apprised of the potential hazard to a fetus (see WARNINGS AND PRECAUTIONS, Fetal Toxicity).**

**ratio-TOPIRAMATE** treatment can cause metabolic acidosis. The effect of topiramate-induced metabolic acidosis has not been studied in pregnancy; however, metabolic acidosis in pregnancy (due to other causes) can cause decreased fetal growth, decreased fetal oxygenation, and fetal death, and may affect the fetus' ability to tolerate labour. Pregnant patients should be monitored for metabolic acidosis and treated as in the nonpregnant state. Newborns of mothers treated with **ratio-TOPIRAMATE** should be monitored for metabolic acidosis because of transfer of topiramate to the fetus and possible occurrence of transient metabolic acidosis following birth (see **WARNINGS AND PRECAUTIONS, Endocrine and Metabolism, Metabolic Acidosis**).

#### Women of Childbearing Potential

Data from pregnancy registries indicate that infants exposed to **ratio-TOPIRAMATE** in utero have an increased risk for cleft lip and/or cleft palate (oral clefts) (see **Pregnancy Registry Data** below). Consider the benefits and the risks of **ratio-TOPIRAMATE** when prescribing this drug to women of childbearing potential. Because of the risk of oral clefts to the fetus, which occur in the first trimester of pregnancy before many women know they are pregnant, all women of childbearing potential should be apprised of the potential hazard to the fetus from exposure to **ratio-TOPIRAMATE**. If the decision is made to use **ratio-TOPIRAMATE**, women who are not planning a pregnancy should use effective contraception (see **DRUG INTERACTIONS, Other Drug Interactions, Oral Contraceptives**). Women who are planning a pregnancy should be counselled regarding the relative risks and benefits of **ratio-TOPIRAMATE** use during pregnancy, and alternative therapeutic options should be considered for these patients (see **Information for Patients, Fetal Toxicity**).

To provide information regarding the effects of *in utero* exposure to **ratio-TOPIRAMATE**, physicians are advised to recommend that pregnant patients taking **ratio-TOPIRAMATE** enroll in the North American Antiepileptic Drug (NAAED) Pregnancy Registry. This can be done by calling the toll-free number, 1-888-233-2334, and must be done by patients themselves. Information on the registry can also be found at the website <http://www.massgeneral.org/aed/>.

#### Labour and Delivery

Although the effect of **ratio-TOPIRAMATE** on labour and delivery in humans has not been established, the development of topiramate-induced metabolic acidosis in the mother and/or in the fetus might affect the fetus' ability to tolerate labour (see **Special Populations, EPILEPSY**).

#### Pregnancy Registry Data

Data from the NAAED Pregnancy Registry indicate an increased risk of oral clefts in infants exposed to topiramate monotherapy during the first trimester of pregnancy. The prevalence of oral clefts was 1.2% compared to a prevalence of 0.39% to 0.46% in infants exposed to other AEDs, and a prevalence of 0.12% in infants of mothers without epilepsy or treatment with other AEDs. For comparison, the Centers for Disease Control and Prevention (CDC) reviewed available data on oral clefts in the United States and found a similar background rate of 0.17%. The relative risk of oral clefts in topiramate-exposed pregnancies in the NAAED Pregnancy Registry was 9.6 (95% Confidence Interval = CI 3.6–25.7) as compared to the risk in a background population of untreated women. The UK Epilepsy and Pregnancy Register reported a similarly increased prevalence of oral clefts of 3.2% among infants exposed to topiramate monotherapy. The observed rate of oral clefts was 16 times higher than the background rate in the UK, which is approximately 0.2%.

### **MIGRAINE PROPHYLAXIS**

Prophylactic treatment of migraine: In pregnancy, the occurrence of seizures presents a significant risk for the mother and child. Prescribing ratio-TOPIRAMATE to prevent seizures therefore outweighs the risk of malformations to the fetus. However, taking ratio-TOPIRAMATE to prevent migraine attacks does not outweigh this risk. Consequently, ratio-TOPIRAMATE is contraindicated in pregnancy and in women of child bearing potential who are not using an effective method of contraception (see **CONTRAINDICATIONS** and **DOSAGE AND ADMINISTRATION, Dosing Considerations**).

**Nursing Women:** Limited data on five breastfeeding infants exposed to topiramate showed infant plasma topiramate levels equal to 10–20% of the maternal plasma level. The effects of this exposure on infants are unknown. Caution should be exercised when administered to a nursing woman.

**Pediatrics < 2 years of age:** Safety and effectiveness in patients below the age of two years have not been established for the adjunctive therapy treatment of partial onset seizures, primary generalized tonic-clonic seizures, or seizures associated with Lennox-Gastaut syndrome. In a single randomized, double-blind, placebo-controlled investigational trial, the efficacy, safety, and tolerability of topiramate oral liquid and sprinkle formulations as an adjunct to concurrent antiepileptic drug therapy in infants one to 24 months of age with refractory partial onset seizures were assessed. After 20 days of double-blind treatment, topiramate (at fixed doses of 5, 15, and 25 mg/kg/day) did not demonstrate efficacy compared with placebo in controlling seizures.

Results from the above controlled epilepsy trial and an open-label, long-term extension study in patients under two years of age indicated that some adverse reactions/toxicities occurred in patients under two years of age that had not been previously observed in older pediatric patients and adults for various indications. These events included growth/length retardation, certain clinical laboratory abnormalities, and other adverse reactions/toxicities that occurred with a greater frequency and/or greater severity.

### **Infection**

These very young pediatric patients (< 2 years) appeared to experience an increased risk for infections (any topiramate dose: 12% vs. placebo: 0%) and of respiratory disorders (any

topiramate dose: 40% vs. placebo: 16%). The following adverse reactions were observed in at least 3% of patients on topiramate and were 3% to 7% more frequent than in patients on placebo: viral infection, bronchitis, pharyngitis, rhinitis, otitis media, upper respiratory infection, cough, and bronchospasm. A generally similar profile was observed in older children.

#### Creatinine and BUN

Topiramate resulted in an increased incidence of patients with increased creatinine (any topiramate dose: 5% vs. placebo: 0%), BUN (any topiramate dose: 3% vs. placebo: 0%), protein (any topiramate dose: 34% vs. placebo: 6%), and an increased incidence of decreased potassium (any topiramate dose: 7% vs. placebo: 0%). This increased frequency of abnormal values was not dose-related. The clinical significance of these findings is uncertain.

#### Other Events

Topiramate treatment also produced a dose-related increase in the percentage of patients who had a shift from normal at baseline to high/increased (above the normal reference range) in total eosinophil count at the end of treatment (placebo: 6%; 5 mg/kg/day: 10%; 15 mg/kg/day: 9%; 25 mg/kg/day: 14%; any topiramate dose: 11%).

There was a mean dose-related increase in alkaline phosphatase. The clinical significance of these findings is uncertain.

Topiramate produced a dose-related increased incidence of treatment-emergent hyperammonemia (see **WARNINGS AND PRECAUTIONS, Hyperammonemia and Encephalopathy**).

Treatment with topiramate for up to one year was associated with reductions in Z SCORES for length, weight, and head circumference (see **WARNINGS AND PRECAUTIONS, Endocrine and Metabolism, Metabolic Acidosis and ADVERSE REACTIONS**).

#### Open-label Epilepsy Trial

In an open-label, adjunctive therapy, epilepsy trial, increasing impairment of adaptive behaviour was documented in behavioural testing over time in children under two years of age. There was a suggestion that this effect was dose-related. However, because of the absence of an appropriate control group, it is not known if this decrement in function was treatment-related or reflects the patient's underlying disease. For example, patients who received higher doses may have more severe underlying disease.

In this open-label, uncontrolled study, the mortality was 37 deaths/1,000 patient years. It is not possible to know whether this mortality rate is related to topiramate treatment, because the background mortality rate for a similar, significantly refractory, young pediatric population under two years with partial epilepsy, is not known.

Safety and efficacy of topiramate for the monotherapy treatment of partial onset seizures or any other type of epilepsy in patients under two years of age, have not been established.

#### MIGRAINE PROPHYLAXIS

Although not indicated for migraine prophylaxis in patients under 18 years of age (see



**INDICATIONS**), in a double-blind, placebo-controlled trial of migraine prophylaxis in patients 12 to 16 years, topiramate treatment produced a dose-related increased shift in serum creatinine from normal at baseline to an increased value at the end of 4 months. The incidence of these abnormal shifts was 4% for placebo, 4% for 50 mg, and 18% for 100 mg.

**Weight Loss in Pediatrics (>2 years of age):** Topiramate administration is associated with weight loss in some children that generally occurs early in therapy. Of those pediatric subjects treated in clinical trials for at least a year who experienced weight loss, 96% showed a resumption of weight gain within the period tested. In 2-4 year-olds, the mean change in weight from baseline at 12 months (n=25) was +0.7 kg (range -1.1 to 3.2); at 24 months (n=14), the mean change was +2.2 kg (range -1.1 to 6.1). In 5-10 year olds, the mean change in weight from baseline at 12 months (n=88) was +0.7 kg (range -6.7 to 11.8); at 24 months (n=67), the mean change was +3.3 kg (range -8.6 to 20.0). Weight decreases, usually associated with anorexia or appetite changes, were reported as adverse events for 9% of patients treated with topiramate. The long-term effects of reduced weight gain in pediatric patients are not known.

**Geriatrics (> 65 years of age):** There is limited information in patients over 65 years of age. The possibility of age-associated renal function abnormalities should be considered when using **ratio-TOPIRAMATE** (topiramate). (See **ACTION AND CLINICAL PHARMACOLOGY, Special Populations and Conditions**).

#### **Monitoring and Laboratory Tests**

It has been observed in clinical trials that topiramate treated subjects experienced an average decrease in serum bicarbonate level of 4 mmol/L and an average increase in serum chloride level of 4 mmol/L. (See **WARNINGS AND PRECAUTIONS, Endocrine and Metabolism**)

**Hypokalemia Observed During Concomitant Treatment with Hydrochlorothiazide:** In a drug interaction study with the diuretic hydrochlorothiazide (HCTZ), the percentage of patients with a serum potassium measurement of <3.6 mEq/L was greater at the end of concomitant treatment than at the end of treatment for either drug alone: 27% (3/11) of subjects on topiramate treatment alone and 25% (3/12) of subjects on HCTZ alone versus 61% (14/22) of subjects on concomitant drug treatment (see **WARNINGS AND PRECAUTIONS, Endocrine and Metabolism, Decrease in Serum Potassium with Concomitant Treatment with Hydrochlorothiazide (HCTZ)** and **DRUG INTERACTIONS, *Hydrochlorothiazide (HCTZ)***).

## **ADVERSE REACTIONS**

The majority of the most common adverse events in clinical trials were mild-moderate in severity and dose-related. These dose-related adverse events typically began in the titration phase and often persisted into the maintenance phase, but infrequently began in the maintenance phase. Rapid titration rate and higher initial dose were associated with higher incidences of adverse events leading to discontinuation.

## **EPILEPSY**

### **Adverse Drug Reaction Overview for Monotherapy Adults**

The most commonly observed adverse events associated with the use of topiramate at dosages of 100 to 400 mg/day in controlled trials in adults with newly diagnosed epilepsy were: paresthesia, fatigue, headache, somnolence, dizziness, upper respiratory tract infection, anorexia, weight decrease, depression, and nausea (see Table 1.1).

Approximately 19% of the 886 adult patients who received topiramate as monotherapy in controlled clinical trials for patients with newly diagnosed epilepsy discontinued therapy due to adverse events. Adverse events associated with discontinuing therapy included paresthesia (2.6%), somnolence (2.5%), fatigue (2.3%), nausea (2.0%), and psychomotor slowing (1.6%).

### **Pediatrics**

The most commonly observed adverse events associated with the use of topiramate at dosages of 100 to 400 mg/day in controlled trials in children with newly diagnosed epilepsy were: upper respiratory tract infection, headache, anorexia, difficulty with concentration/attention, weight decrease, somnolence, paresthesia, fever, and fatigue (see Table 1.2).

Approximately 10% of the 245 pediatric patients who received topiramate as monotherapy in controlled clinical trials for patients with newly diagnosed epilepsy discontinued therapy due to adverse events. Adverse events associated with discontinuing therapy included difficulty with concentration/attention (2.0%). No pediatric patients withdrew due to psychomotor slowing.

### **Clinical Trial Adverse Drug Reactions**

*Because clinical trials are conducted under very specific conditions the adverse reaction rates observed in the clinical trials may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse drug reaction information from clinical trials is useful for identifying drug-related adverse events and for approximating rates.*

**Table 1.1: Incidence of Treatment-Emergent Adverse Events in Monotherapy Trials in Adults<sup>a</sup> where rate was  $\geq 2\%$  in any Topiramate Group**

Body System / Adverse Event	Topiramate Dosage (mg/day)		
	50-100 (n=444)	200-400 (n=329)	500 (n=113)
<b>Body as a Whole-General Disorders</b>			
Fatigue	18	18	19
Injury	9	8	4
Asthenia	4	5	4
Back Pain	3	2	5
Pain	3	2	5
Chest Pain	2	2	3
Fever	1	2	3
Syncope	2	1	1
Leg Pain	2	2	1
Peripheral Edema	1	<1	2
<b>Central &amp; Peripheral Nervous System Disorders</b>			
Paresthesia	23	39	38
Headache	23	16	19
Dizziness	16	13	13
Hypoesthesia	5	5	12
Language Problems	4	5	6
Ataxia	3	5	4
Speech Disorder/Related Speech Problems	2	3	3
Vertigo	2	3	4
Tremor	3	2	3
Hypertonia	1	2	2
Involuntary Muscle Contractions	1	2	4
Sensory Disturbances	1	1	4
Migraine	2	1	1
Abnormal Coordination	1	1	3
Convulsions Aggravated	1	0	2
Convulsions Grand Mal	<1	1	2
Gait Abnormal	<1	<1	3
Dyskinesia	0	0	2
<b>Gastrointestinal System Disorders</b>			
Nausea	11	12	12
Diarrhea	6	8	12
Abdominal Pain	6	8	7
Dyspepsia	5	5	4
Vomiting	4	3	2
Constipation	2	3	1
Dry Mouth	1	2	6
Gastroenteritis	2	1	2
Gastritis	1	2	2
Tooth Ache	1	1	2
Gastrointestinal Disorder NOS	<1	<1	2
Hemorrhoids	<1	<1	2
Stomatitis Ulcerative	<1	0	2
<b>Hearing and Vestibular Disorders</b>			
Tinnitus	1	2	2
<b>Heart Rate and Rhythm Disorders</b>			
Palpitation	1	1	4
Tachycardia	1	0	2
<b>Metabolic and Nutritional Disorders</b>			
Weight Decrease	9	14	18
<b>Musculoskeletal System Disorders</b>			
Arthralgia	3	4	4
Myalgia	2	1	2
Muscle Weakness	1	1	2

**Table 1.1: Incidence of Treatment-Emergent Adverse Events in Monotherapy Trials in Adults<sup>a</sup> where rate was  $\geq 2\%$  in any Topiramate Group**

Body System / Adverse Event	Topiramate Dosage (mg/day)		
	50-100 (n=444)	200-400 (n=329)	500 (n=113)
<b>Platelet, Bleeding, and Clotting Disorders</b>			
Epistaxis	1	2	1
Hematoma	0	0	2
<b>Psychiatric Disorders</b>			
Somnolence	11	15	19
Anorexia	8	14	11
Insomnia	9	8	9
Difficulty with Memory NOS	6	10	9
Depression	7	10	4
Difficulty with Concentration/Attention	6	9	8
Nervousness	6	7	8
Mood Problems	5	6	4
Anxiety	4	6	5
Confusion	4	5	7
Psychomotor Slowing	2	5	8
Cognitive Problems NOS	2	3	3
Agitation	2	2	3
Emotional Lability	1	3	2
Aggressive Reaction	2	1	2
Libido Decreased	1	2	1
Depression Aggravated	<1	2	3
Impotence	1	1	2
<b>Reproductive Disorders, Female</b>			
Menstrual Disorder	3	1	8
Dysmenorrhea	2	2	0
Intermenstrual Bleeding	2	1	0
Menorrhagia	1	1	2
Pregnancy Unintended	1	1	2
Mastitis	0	0	2
<b>Reproductive Disorders, Male</b>			
Premature Ejaculation	0	0	2
<b>Resistance Mechanism Disorders</b>			
Infection Viral	5	9	6
Otitis Media	2	1	2
<b>Respiratory System Disorders</b>			
Upper Respiratory Tract Infection	15	13	10
Pharyngitis	5	5	2
Sinusitis	3	4	6
Rhinitis	3	3	5
Bronchitis	2	2	1
Coughing	2	2	2
Dyspnea	1	2	1
Pneumonia	1	<1	3
<b>Skin and Appendages Disorders</b>			
Rash	3	4	3
Alopecia	3	3	1
Acne	1	3	2
Pruritus	1	3	1
Increased Sweating	1	<1	2
Maculopapular Rash	1	0	2
<b>Special Senses Other, Disorders</b>			
Taste Perversion	3	5	6
<b>Urinary System Disorders</b>			
Urinary Tract Infection	2	2	5
Micturition Frequency	1	2	4
Dysuria	<1	2	1
Cystitis	<1	2	1
Renal Calculus	<1	2	2

**Table 1.1: Incidence of Treatment-Emergent Adverse Events in Monotherapy Trials in Adults<sup>a</sup> where rate was  $\geq 2\%$  in any Topiramate Group**

Body System / Adverse Event	Topiramate Dosage (mg/day)		
	50-100 (n=444)	200-400 (n=329)	500 (n=113)
<b>Vision Disorders</b>			
Vision Abnormal	3	4	4
Diplopia	1	1	2

<sup>a</sup> Values represent the percentage of patients reporting a given adverse event. Patients may have reported more than one adverse event during the study and can be included in more than one adverse event category.

**Table 1.2: Incidence of Treatment-Emergent Adverse Events in Monotherapy Trials in Children Ages 6 up to 16 years<sup>a</sup> where rate was  $\geq 2\%$  in any Topiramate Group**

Body System / Adverse Event	Topiramate Dosage (mg/day)		
	50-100 (n=125)	200-400 (n=106)	500 <sup>b</sup> (n=14)
<b>Body as a Whole-General Disorders</b>			
Fatigue	7	10	14
Fever	2	11	7
Injury	4	2	14
Asthenia	0	3	7
Back Pain	2	2	0
Allergic Reaction	1	1	7
Allergy	0	1	7
Influenza-Like Symptoms	0	0	7
<b>Central &amp; Peripheral Nervous System Disorders</b>			
Headache	27	17	29
Dizziness	9	8	0
Paresthesia	4	11	7
Language Problems	0	3	7
Convulsions Grand Mal	2	0	7
Hypertonia	0	0	7
Hyperkinesia	2	0	21
Migraine	2	1	0
Muscle Contractions Involuntary	1	2	0
Tremor	2	0	0
Vertigo	0	3	0
Cramps Legs	2	0	0
Gait Abnormal	2	0	0
<b>Collagen Disorders</b>			
Auto-antibody Response	0	0	7
<b>Gastrointestinal System Disorders</b>			
Diarrhea	9	7	7
Vomiting	8	6	14
Abdominal Pain	6	4	14
Nausea	4	5	14
Gastroenteritis	6	0	7
Constipation	1	0	7
Gastrointestinal Disorder NOS	0	0	7
Dyspepsia	2	1	0
Tooth Ache	1	1	7
<b>Hearing and Vestibular Disorders</b>			
Earache	2	0	0
<b>Metabolic and Nutritional Disorders</b>			
Weight Decrease	5	14	0
Acidosis	0	0	7
<b>Musculoskeletal System Disorders</b>			
Arthralgia	1	2	7
<b>Platelet, Bleeding, &amp; Clotting Disorders</b>			
Epistaxis	2	4	14

**Table 1.2: Incidence of Treatment-Emergent Adverse Events in Monotherapy Trials in Children Ages 6 up to 16 years<sup>a</sup> where rate was  $\geq 2\%$  in any Topiramate Group**

Body System / Adverse Event	Topiramate Dosage (mg/day)		
	50-100 (n=125)	200-400 (n=106)	500 <sup>b</sup> (n=14)
<b>Psychiatric Disorders</b>			
Anorexia	13	13	14
Somnolence	14	9	0
Difficulty with Concentration/Attention	6	13	7
Insomnia	5	4	14
Nervousness	5	6	0
Mood Problems	2	8	0
Difficulty with Memory NOS	4	2	14
Cognitive Problems NOS	1	6	0
Psychomotor Slowing	3	3	0
Aggressive Reaction	2	3	7
Depression	0	5	0
Sleep Disorder	2	2	0
Personality Disorder (Behaviour Problems)	2	2	0
Anxiety	2	1	0
Confusion	0	3	0
Emotional Liability	2	1	0
<b>Red Blood Cell Disorders</b>			
Anemia	1	2	0
<b>Reproductive Disorders, Female</b>			
Vaginitis	0	0	13
Dysmenorrhea	2	2	0
Intermenstrual Bleeding	0	2	0
<b>Reproductive Disorders, Male</b>			
Testis Disorder	2	0	0
<b>Resistance Mechanism Disorder</b>			
Infection Viral	4	7	7
Infection	2	6	0
Otitis Media	2	1	7
<b>Respiratory System Disorders</b>			
Upper Respiratory Tract Infection	26	25	21
Pharyngitis	9	5	21
Rhinitis	5	6	21
Sinusitis	3	6	14
Bronchitis	2	4	0
Asthma	2	1	0
Coughing	2	1	0
<b>Skin and Appendages Disorders</b>			
Rash	3	4	21
Dermatitis	1	0	7
Alopecia	1	3	0
Acne	2	0	0
Nail Disorder	2	0	0
Pruritus	0	2	0
Rash Erythematous	2	0	0
<b>Urinary System Disorders</b>			
Urinary Incontinence	2	2	7
Renal Calculus	0	0	7
Micturition Frequency	0	2	0
Urinary Tract Infection	2	0	0
<b>Vascular Disorders</b>			
Flushing	1	4	7
<b>Vision Disorders</b>			
Conjunctivitis	2	2	0

<sup>a</sup> Values represent the percentage of patients reporting a given adverse event. Patients may have reported more than one adverse event during the study and can be included in more than one adverse event category.

<sup>b</sup> Due to n=14 in the 500 mg topiramate group, an incidence of 7% represents one patient

## **Adverse Drug Reaction Overview for Adjunctive Therapy**

### **Adults**

The most commonly observed adverse events associated with the adjunctive use of topiramate at dosages of 200 to 400 mg/day in controlled trials in adults that were seen at greater frequency in patients treated with topiramate and did not appear to be dose related within this dosage range were: somnolence, dizziness, ataxia, speech disorders and related speech problems, psychomotor slowing, nystagmus, and paresthesia (see Table 1.3).

The most common dose-related adverse events at dosages of 200 to 1,000 mg/day were: nervousness, difficulty with concentration or attention, confusion, depression, anorexia, language problems, and mood problems (see Table 1.4).

### **Pediatrics**

Adverse events associated with the use of topiramate at dosages of 5 to 9 mg/kg/day in worldwide pediatric clinical trials that were seen at greater frequency in patients treated with topiramate were: fatigue, somnolence, anorexia, nervousness, difficulty with concentration/attention, difficulty with memory, aggressive reaction, and weight decrease (see Table 1.5).

### **Clinical Trial Adverse Drug Reactions**

*Because clinical trials are conducted under very specific conditions the adverse reaction rates observed in the clinical trials may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse drug reaction information from clinical trials is useful for identifying drug-related adverse events and for approximating rates.*

**Table 1.3: Incidence (%) of Treatment-Emergent Adverse Events in Placebo-Controlled, Add-On Epilepsy Trials in Adults <sup>a,b</sup> (Events that Occurred in  $\geq 2\%$  of Patients Treated with Topiramate and Occurred More Frequently in Patients Treated with Topiramate Than Placebo-Treated Patients)**

Body System / Adverse Event	Topiramate Dosage (mg/day)		
	Placebo (n=216)	200-400 (n=113)	600-1,000 (n=414)
<b>Body as a Whole</b>			
Asthenia	1.4	8.0	3.1
Back Pain	4.2	6.2	2.9
Chest Pain	2.8	4.4	2.4
Influenza-Like Symptoms	3.2	3.5	3.6
Leg Pain	2.3	3.5	3.6
Hot Flushes	1.9	2.7	0.7
<b>Nervous System</b>			
Dizziness	15.3	28.3	32.1
Ataxia	6.9	21.2	14.5
Speech Disorder/Related Speech Problems	2.3	16.8	11.4
Nystagmus	9.3	15.0	11.1
Paresthesia	4.6	15.0	19.1
Tremor	6.0	10.6	8.9
Language Problems	0.5	6.2	10.4
Coordination Abnormal	1.9	5.3	3.6
Hypoesthesia	0.9	2.7	1.2
Abnormal Gait	1.4	1.8	2.2

**Table 1.3: Incidence (%) of Treatment-Emergent Adverse Events in Placebo-Controlled, Add-On Epilepsy Trials in Adults <sup>a,b</sup> (Events that Occurred in  $\geq 2\%$  of Patients Treated with Topiramate and Occurred More Frequently in Patients Treated with Topiramate Than Placebo-Treated Patients)**

Body System / Adverse Event	Topiramate Dosage (mg/day)		
	Placebo (n=216)	200-400 (n=113)	600-1,000 (n=414)
<b>Gastrointestinal System</b>			
Nausea	7.4	11.5	12.1
Dyspepsia	6.5	8.0	6.3
Abdominal Pain	3.7	5.3	7.0
Constipation	2.3	5.3	3.4
Dry Mouth	0.9	2.7	3.9
<b>Metabolic and Nutritional</b>			
Weights Decrease	2.8	7.1	12.8
<b>Neuropsychiatric</b>			
Somnolence	9.7	30.1	27.8
Psychomotor Slowing	2.3	16.8	20.8
Nervousness	7.4	15.9	19.3
Difficulty with Memory	3.2	12.4	14.5
Confusion	4.2	9.7	13.8
Depression	5.6	8.0	13.0
Difficulty with Concentration/Attention	1.4	8.0	14.5
Anorexia	3.7	5.3	12.3
Agitation	1.4	4.4	3.4
Mood Problems	1.9	3.5	9.2
Aggressive Reaction	0.5	2.7	2.9
Apathy	0	1.8	3.1
Depersonalization	0.9	1.8	2.2
Emotional Lability	0.9	1.8	2.7
<b>Reproductive, Female</b>	(n=59)	(n=24)	(n=128)
Breast Pain, Female	1.7	8.3	0
Dysmenorrhea	6.8	8.3	3.1
Menstrual Disorder	0	4.2	0.8
<b>Reproductive, Male</b>	(n=157)	(n=89)	(n=286)
Prostatic Disorder	0.6	2.2	0
<b>Respiratory System</b>			
Pharyngitis	2.3	7.1	3.1
Rhinitis	6.9	7.1	6.3
Sinusitis	4.2	4.4	5.6
Dyspnea	0.9	1.8	2.4
<b>Skin and Appendages</b>			
Pruritus	1.4	1.8	3.1
<b>Vision</b>			
Diplopia	5.6	14.2	10.4
Vision Abnormal	2.8	14.2	10.1
<b>White Cell and RES</b>			
Leukopenia	0.5	2.7	1.2

<sup>a</sup> Patients in these add-on trials were receiving 1 to 2 concomitant antiepileptic drugs in addition to Topiramate or placebo.

<sup>b</sup> Values represent the percentage of patients reporting a given adverse event. Patients may have reported more than one adverse event during the study and can be included in more than one adverse event category.



**Table 1.4: Incidence (%) of Dose-Related Adverse Events From Placebo-Controlled, Add-On Epilepsy Trials in Adults**

Adverse Event	Topiramate Dosage (mg/day)			
	Placebo (n=216)	200 (n=45)	400 (n=68)	600-1,000 (n=414)
Fatigue	13.4	11.1	11.8	29.7
Nervousness	7.4	13.3	17.6	19.3
Difficulty with Concentration/Attention	1.4	6.7	8.8	14.5
Confusion	4.2	8.9	10.3	13.8
Depression	5.6	8.9	7.4	13
Anorexia	3.7	4.4	5.9	12.3
Language problems	0.5	2.2	8.8	10.1
Anxiety	6	2.2	2.9	10.4
Mood Problems	1.9	0	5.9	9.2

In six double-blind clinical trials, 10.6% of subjects (n=113) assigned to a topiramate dosage of 200 to 400 mg/day in addition to their standard AED therapy discontinued due to adverse events, compared to 5.8% of subjects (n=69) receiving placebo. The percentage of subjects discontinuing due to adverse events appeared to increase at dosages above 400 mg/day. Overall, approximately 17% of all subjects (n=527) who received topiramate in the double-blind trials discontinued due to adverse events, compared to 4% of the subjects (n=216) receiving placebo.

Table 1.5 lists treatment-emergent adverse events that occurred in at least 2% of children treated with 5 to 9 mg/kg/day topiramate in controlled trials that were numerically more common than in patients treated with placebo.

**Table 1.5: Incidence (%) of Treatment-Emergent Adverse Events in Worldwide Pediatric Add-on Epilepsy Clinical Trials Experience (2-16 years of Age)<sup>a,b</sup> (Events that Occurred in ≥2% of Patients Treated with Topiramate and Occurred More Frequently in Patients Treated with Topiramate Than Placebo-Treated Patients)**

Body System / Adverse Event	Placebo (n=101)	Topiramate (n=98)
<b>Body as a Whole - General Disorders</b>		
Fatigue	5	16.3
Injury	12.9	14.3
Allergic Reaction	1	2
<b>Central and Peripheral Nervous System Disorders</b>		
Gait Abnormal	5	8.2
Ataxia	2	6.1
Hyperkinesia	4	5.1
Dizziness	2	4.1
Speech Disorder/Related Speech Problems	2	4.1
Convulsions Aggravated	3	3.1
Hyporeflexia	0	2

**Table 1.5: Incidence (%) of Treatment-Emergent Adverse Events in Worldwide Pediatric Add-on Epilepsy Clinical Trials Experience (2-16 years of Age)<sup>a,b</sup> (Events that Occurred in ≥2% of Patients Treated with Topiramate and Occurred More Frequently in Patients Treated with Topiramate Than Placebo-Treated Patients)**

<b>Body System / Adverse Event</b>	<b>Placebo (n=101)</b>	<b>Topiramate (n=98)</b>
<b>Gastrointestinal System Disorders</b>		
Nausea	5	6.1
Saliva Increased	4	6.1
Constipation	4	5.1
Gastroenteritis	2	3.1
<b>Metabolic and Nutritional Disorders</b>		
Weight Decrease	1	9.2
Thirst	1	2
<b>Platelet, Bleeding and Clotting Disorders</b>		
Purpura	4	8.2
Epistaxis	1	4.1
<b>Nervous Disorders</b>		
Somnolence	15.8	25.5
Anorexia	14.9	24.5
Nervousness	6.9	14.3
Personality Disorder (Behaviour Problems)	8.9	11.2
Difficulty with Concentration/Attention	2	10.2
Aggressive Reaction	4	9.2
Insomnia	6.9	8.2
Mood Problems	6.9	7.1
Difficulty with Memory NOS <sup>c</sup>	0	5.1
Emotional Lability	5	5.1
Confusion	3	4.1
Psychomotor Slowing	2	3.1
<b>Reproductive Disorders, Female</b>		
Leukorrhea	0	2.3
<b>Resistance Mechanism Disorders</b>		
Infection Viral	3	7.1
Infection	3	3.1
<b>Respiratory System Disorders</b>		
Upper Respiratory Tract Infection	36.6	36.7
Pneumonia	1	5.1
<b>Skin and Appendages Disorders</b>		
Skin Disorder	2	3.1
Alopecia	1	2
Dermatitis	0	2
Hypertrichosis	1	2
Rash Erythematous	0	2
<b>Urinary System Disorders</b>		
Urinary Incontinence	2	4.1
<b>Vision Disorders</b>		
Eye Abnormality	1	2
Vision Abnormal	1	2
<b>White Cell and RES Disorders</b>		
Leukopenia	0	2

<sup>a</sup> Patients in these add-on trials were receiving 1 to 2 concomitant antiepileptic drugs in addition to Topiramate or placebo.

<sup>b</sup> Values represent the percentage of patients reporting a given adverse event. Patients may have reported more than one adverse event during the study and can be included in more than one adverse event category.

<sup>c</sup> Not otherwise specified.

None of the pediatric patients who received topiramate adjunctive therapy at 5 to 9 mg/kg/day in controlled clinical trials discontinued due to adverse events. In open extensions of the controlled clinical trials, approximately 9% of the 303 pediatric patients who received topiramate at dosages up to 30 mg/kg/day discontinued due to adverse events. Adverse events associated with discontinuing therapy included aggravated convulsions (2.3%), language problems (1.3%), and

difficulty with concentration/attention (1.3%).

When the safety experience of patients receiving topiramate as adjunctive therapy in both double-blind and open-label trials (1,446 adults and 303 children) was analyzed, a similar pattern of adverse events emerged.

### **Less Common Clinical Trial Adverse Drug Reactions (< 2%)**

Adverse events that occurred less frequently but were considered potentially medically relevant included: taste perversion, cognitive problems (not otherwise specified) and psychosis/psychotic symptoms.

In adult and pediatric patients, nephrolithiasis was reported rarely. Isolated cases of thromboembolic events have also been reported; a causal association with the drug has not been established.

In clinical trials with topiramate, the occurrence rate for all potential cases of oligohidrosis (decreased sweating) was 0.25%.

In clinical trials for topiramate in epilepsy, migraine prophylaxis and other investigational indications (obesity, bipolar disorder and diabetic peripheral neuropathy), suicide-related adverse events\* occurred at a rate of 0.8% (84 reports/10,846 patients) in topiramate versus 0.2% (5 reports/3,150 patients) in placebo groups. Although the average exposure time for patients on topiramate (approximately 10 months) was longer than for those on placebo (approximately 5 months), these adverse events were reported randomly over the exposure period. Suicide attempts occurred in 0.3% (33 reports/10,846 patients) of the topiramate-treated patients compared to 0% in placebo groups. Of these 33 attempts, one completed suicide was reported in a double-blind bipolar disorder trial and 3 in the open-label phase of the bipolar disorder trials (see **WARNINGS AND PRECAUTIONS, Neurologic, Psychiatric**).

\*Suicide-related adverse events include suicidal ideation, suicide attempt, suicide and any evidence of self-harm.

## **MIGRAINE PROPHYLAXIS**

### **Adverse Drug Reaction Overview**

Table 1.6 includes those adverse events reported for patients in four multicentre, randomized, double-blind, placebo-controlled, parallel-group migraine prophylaxis clinical trials where the incidence rate in any topiramate treatment group was at least 2% and was greater than that for placebo patients. Most of the adverse events were mild or moderate in severity and most occurred more frequently during the titration period than during the maintenance period.

### **Clinical Trial Adverse Drug Reactions**

*Because clinical trials are conducted under very specific conditions the adverse reaction rates observed in the clinical trials may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse drug reaction information from clinical trials is useful for identifying drug-related adverse events and for approximating rates.*

**Table 1.6: Incidence (%) of Treatment-Emergent Adverse Events in Placebo-Controlled Migraine Trials Where Rate Was at Least 2% in any Topiramate Group and Greater Than the Rate in Placebo-Treated Patients<sup>a</sup>**

Body System / Adverse Event	Topiramate Dosage (mg/day)			
	Placebo (n=445)	50 (n=235)	100 (n=386)	200 (n=514)
<b>Body as a Whole - General Disorders</b>				
Fatigue	11	14	15	19
Injury	7	9	6	6
Asthenia	1	<1	2	2
Fever	1	1	1	2
Influenza-Like Symptoms	<1	<1	<1	2
Allergy	<1	2	<1	<1
<b>Central and Peripheral Nervous System Disorders</b>				
Paresthesia	6	35	51	49
Dizziness	10	8	9	12
Hypoesthesia	2	6	7	8
Language Problems	2	7	6	7
Involuntary Muscle Contractions	1	2	2	4
Ataxia	<1	1	2	1
Speech Disorder/Related Speech Problems	<1	1	<1	2
<b>Gastrointestinal System Disorders</b>				
Nausea	8	9	13	14
Diarrhea	4	9	11	11
Abdominal Pain	5	6	6	7
Dyspepsia	3	4	5	3
Dry Mouth	2	2	3	5
Vomiting	2	1	2	3
Gastroenteritis	1	3	3	2
<b>Hearing and Vestibular Disorders</b>				
Tinnitus	1	<1	1	2
<b>Metabolic and Nutritional Disorders</b>				
Weight Decrease	1	6	9	11
Thirst	<1	2	2	1
<b>Musculoskeletal System Disorders</b>				
Arthralgia	2	7	3	1
<b>Neoplasms</b>				
Neoplasm NOS	<1	2	<1	<1
<b>Psychiatric Disorders</b>				
Anorexia	6	9	15	14
Somnolence	5	8	7	10
Difficulty with Memory NOS	2	7	7	11
Difficulty with Concentration/Attention	2	3	6	10
Insomnia	5	6	7	6
Anxiety	3	4	5	6
Mood Problems	2	3	6	5
Depression	4	3	4	6
Nervousness	2	4	4	4
Confusion	2	2	3	4
Psychomotor Slowing	1	3	2	4
Libido Decreased	1	1	1	2
Aggravated Depression	1	1	2	2
Agitation	1	2	2	1
Cognitive Problems NOS	1	<1	2	2
<b>Reproductive Disorders, Female</b>				
Menstrual Disorder	2	3	2	2
<b>Reproductive Disorders, Male</b>				
Ejaculation Premature	0	3	0	0
<b>Resistance Mechanism Disorders</b>				
Viral Infection	3	4	4	3
Otitis Media	<1	2	1	1

**Table 1.6: Incidence (%) of Treatment-Emergent Adverse Events in Placebo-Controlled Migraine Trials Where Rate Was at Least 2% in any Topiramate Group and Greater Than the Rate in Placebo-Treated Patients<sup>a</sup>**

Body System / Adverse Event	Topiramate Dosage (mg/day)			
	Placebo (n=445)	50 (n=235)	100 (n=386)	200 (n=514)
<b>Respiratory System Disorders</b>				
Upper Respiratory Tract Infection	12	13	14	12
Sinusitis	6	10	6	8
Pharyngitis	4	5	6	2
Coughing	2	2	4	3
Bronchitis	2	3	3	3
Dyspnea	2	1	3	2
Rhinitis	1	1	2	2
<b>Skin and Appendages Disorders</b>				
Pruritus	2	4	2	2
<b>Special Sense Other, Disorders</b>				
Taste Perversion	1	15	8	12
Taste Loss	<1	1	1	2
<b>Urinary System Disorders</b>				
Urinary Tract Infection	2	4	2	4
Renal Calculus	0	0	1	2
<b>Vision Disorders</b>				
Vision Abnormal	<1	1	2	3
Blurred Vision <sup>b</sup>	2	4	2	4
Conjunctivitis	1	1	2	1

<sup>a</sup> Values represent the percentage of patients reporting a given adverse event. Patients may have reported more than one adverse event during the study and can be included in more than one adverse event category.

<sup>b</sup> Blurred vision was the most common term considered as vision abnormal. Blurred vision was an included term that accounted for >50% of events coded as vision abnormal, a preferred term.

Of the 1,135 patients exposed to topiramate in the placebo-controlled studies, 25% discontinued due to adverse events, compared to 10% of the 445 placebo patients. The most common adverse events associated with discontinuing therapy in the topiramate-treated patients included paresthesia (6.7%), fatigue (4.3%), nausea (4.0%), difficulty with concentration/attention (2.9%), insomnia (2.7%), anorexia (2.1%), and dizziness (2.0%).

In the 6-month migraine prophylaxis controlled trials, the proportion of patients who experienced one or more cognitive-related events was 19% for topiramate 50 mg/day, 22% for 100 mg/day, 28% for 200 mg/day and 10% for placebo. These dose-related adverse reactions typically began in the titration phase and often persisted into the maintenance phase but infrequently began in the maintenance phase.

Table 1.7 shows adverse events that were dose-dependant.

**Table 1.7: Incidence (%) of Dose-Related Adverse Events From Placebo-Controlled Migraine Trials<sup>a</sup>**

Adverse Event	Topiramate Dosage (mg/day)			
	Placebo (n=445)	50 (n=235)	100 (n=386)	200 (n=514)
Paresthesia	6	35	51	49
Fatigue	11	14	15	19
Nausea	8	9	13	14
Anorexia	6	9	15	14
Dizziness	10	8	9	12
Weight decrease	1	6	9	11
Difficulty with Memory NOS	2	7	7	11
Diarrhea	4	9	11	11
Difficulty with Concentration/Attention	2	3	6	10
Somnolence	5	8	7	10
Hypoesthesia	2	6	7	8
Anxiety	3	4	5	6
Depression	4	3	4	6
Mood Problems	2	3	6	5
Dry Mouth	2	2	3	5
Confusion	2	2	3	4
Involuntary Muscle Contractions	1	2	2	4
Abnormal Vision	<1	1	2	3
Renal Calculus	0	0	1	2

<sup>a</sup> The incidence rate of the adverse event in the 200 mg/day group was  $\geq 2\%$  than the rate in both the placebo group and the 50 mg/day group.

### **Other Adverse Events Observed During Migraine Clinical Trials**

For the prophylactic treatment of migraine headache, topiramate has been administered to 1,367 patients in all clinical studies (includes double-blind and open-label extension). During these studies, all adverse events were recorded by the clinical investigators using terminology of their own choosing. To provide a meaningful estimate of the proportion of individuals having adverse events, similar types of events were grouped into a smaller number of standardized categories using modified WHOART dictionary terminology.

The following additional adverse events that were not described earlier were reported by greater than 1% of the 1367 topiramate-treated patients in the controlled clinical trials:

**Body as a Whole:** pain, chest pain, allergic reaction

**Central and Peripheral Nervous System Disorders:** headache, vertigo, tremor, sensory disturbance, migraine aggravated

**Gastrointestinal System Disorders:** constipation, gastroesophageal reflux, tooth disorder

**Musculoskeletal System Disorders:** myalgia

**Platelet, Bleeding and Clotting Disorders:** epistaxis

**Reproductive Disorders, Female:** intermenstrual bleeding

**Resistance Mechanism Disorders:** infection, genital moniliasis

**Respiratory System Disorders:** pneumonia, asthma

**Skin and Appendages Disorders:** rash, alopecia

**Vision Disorders:** abnormal accommodation, eye pain

### **Post-Market Adverse Drug Reactions**

In addition to the adverse events reported during clinical trial testing of topiramate, the following adverse drug reactions have been reported in patients receiving marketed topiramate from worldwide use since approval. Adverse drug reactions from spontaneous reports during the worldwide post-marketing experience with topiramate are included in Table 1.8 below. The adverse drug reactions are ranked by frequency, using the following convention (all calculated per patient-years of estimated exposure):

Very common	$\geq 1/10$
Common	$\geq 1/100$ and $< 1/10$
Uncommon	$\geq 1/1,000$ and $< 1/100$
Rare	$\geq 1/10,000$ and $< 1/1000$
Very rare	$< 1/10,000$

The frequencies provided below reflect reporting rates for adverse drug reactions from spontaneous reports, and do not represent more precise estimates that might be obtained in clinical or experimental studies.

Adverse Event	Reporting Rate			
	Common	Uncommon	Rare	Very Rare
<b>Blood and Lymphatic System Disorders</b>				
• leucopenia and neutropenia				X
• thrombocytopenia				X
<b>Metabolism and Nutrition Disorders</b>				
• anorexia			X	
• metabolic acidosis <sup>1</sup>				X
• hyperammonemia <sup>2</sup>				X
• hypokalemia				X
<b>Musculoskeletal and Connective Tissue Disorders</b>				
• musculoskeletal pain				X
• myalgia				X
• arthralgia				X
<b>Psychiatric Disorders</b>				
• depression <sup>3</sup>			X	
• agitation <sup>3</sup>			X	
• somnolence <sup>3</sup>			X	
• insomnia <sup>3</sup>				X
• mood altered <sup>3</sup>				X
• confusional state <sup>3</sup>				X
• psychotic disorder <sup>3</sup>				X
• aggression <sup>3</sup>				X
• hallucination <sup>3</sup>				X
• suicidal ideation <sup>3</sup>				X
• suicidal attempts <sup>3</sup>				X
• suicide <sup>3</sup>				X
• expressive language disorder				X
• delusion				X
• concentration impaired				X

<b>Nervous System Disorders</b>				
<ul style="list-style-type: none"> <li>• paresthesia<sup>3</sup></li> <li>• convulsion</li> <li>• headache</li> <li>• dizziness</li> <li>• speech disorder</li> <li>• dysgeusia</li> <li>• amnesia</li> <li>• memory impairment</li> <li>• drug withdrawal convulsion</li> <li>• ataxia</li> <li>• hyperkinesias</li> </ul>			<ul style="list-style-type: none"> <li>X</li> <li>X</li> <li>X</li> <li>X</li> </ul>	<ul style="list-style-type: none"> <li>X</li> <li>X</li> <li>X</li> <li>X</li> <li>X</li> <li>X</li> <li>X</li> </ul>
<b>Eye Disorders</b>				
<ul style="list-style-type: none"> <li>• visual disturbance</li> <li>• vision blurred</li> <li>• myopia<sup>4</sup></li> <li>• angle closure glaucoma<sup>4</sup></li> <li>• eye pain</li> <li>• maculopathy (including field vision defect)</li> </ul>			<ul style="list-style-type: none"> <li>X</li> <li>X</li> </ul>	<ul style="list-style-type: none"> <li>X</li> <li>X</li> <li>X</li> <li>X</li> </ul>
<b>Gastrointestinal Disorders</b>				
<ul style="list-style-type: none"> <li>• nausea</li> <li>• diarrhea</li> <li>• abdominal pain</li> <li>• vomiting</li> </ul>			<ul style="list-style-type: none"> <li>X</li> </ul>	<ul style="list-style-type: none"> <li>X</li> <li>X</li> <li>X</li> </ul>
<b>Skin and Subcutaneous Tissue Disorders</b>				
<ul style="list-style-type: none"> <li>• alopecia</li> <li>• rash</li> </ul>			<ul style="list-style-type: none"> <li>X</li> </ul>	<ul style="list-style-type: none"> <li>X</li> </ul>
<b>Renal and Urinary Disorders</b>				
<ul style="list-style-type: none"> <li>• nephrolithiasis<sup>5</sup></li> </ul>			<ul style="list-style-type: none"> <li>X</li> </ul>	
<b>General Disorders and Administration Site Conditions</b>				
<ul style="list-style-type: none"> <li>• fatigue<sup>1</sup></li> <li>• oligohidrosis<sup>1,6</sup></li> <li>• pyrexia</li> <li>• feeling abnormal</li> <li>• asthenia</li> <li>• dehydration</li> <li>• flushing</li> <li>• hot flushes</li> </ul>			<ul style="list-style-type: none"> <li>X</li> <li>X</li> </ul>	<ul style="list-style-type: none"> <li>X</li> <li>X</li> <li>X</li> <li>X</li> <li>X</li> <li>X</li> <li>X</li> </ul>
<b>Investigations</b>				
<ul style="list-style-type: none"> <li>• weight decreased</li> <li>• hepatic enzymes increased</li> </ul>			<ul style="list-style-type: none"> <li>X</li> </ul>	<ul style="list-style-type: none"> <li>X</li> </ul>

<sup>1</sup> see WARNINGS AND PRECAUTIONS, **Endocrine and Metabolism**

<sup>2</sup> see WARNINGS AND PRECAUTIONS, **Hyperammonemia and Encephalopathy** and DRUG INTERACTIONS, **Drug-Drug Interactions**

<sup>3</sup> see WARNINGS AND PRECAUTIONS, **Neurologic**

<sup>4</sup> see WARNINGS AND PRECAUTIONS, **Ophthalmologic**

<sup>5</sup> see WARNINGS AND PRECAUTIONS, **Renal** and DRUG INTERACTIONS, **Drug-Drug Interactions**

<sup>6</sup> The majority of these reports have been in children.

Oligohidrosis (decreased sweating) has been rarely reported with the use of topiramate. The majority of spontaneous post-marketing reports have been in children. Adverse events that may be related to potential cases of oligohidrosis include dehydration, hyperthermia, and heat



intolerance. Adequate hydration prior to activities such as exercise or exposure to warm temperatures is recommended (see **WARNINGS AND PRECAUTIONS, Endocrine and Metabolism**).

To date, there have been rare spontaneous, post-marketing reports of metabolic acidosis. In some cases, acidosis resolved after dosage reduction or upon discontinuation of topiramate (see **WARNINGS AND PRECAUTIONS, Endocrine and Metabolism**).

Rare reports of encephalopathy with or without hyperammonemia have been received for patients treated with topiramate while also taking valproate or other antiepileptic medications (see **WARNINGS AND PRECAUTIONS, General** and **DRUG INTERACTIONS, Drug-Drug Interactions**).

There have been rare spontaneous postmarketing reports of suicide attempts and suicide-related adverse events, including fatalities, in patients treated with topiramate alone or in combination with other medications (see **WARNINGS AND PRECAUTIONS, Neurologic, Psychiatric**).

The following adverse experiences have not been listed above and data are insufficient to support an estimate of their incidence or to establish causation.

Reports of increases in liver function tests in patients taking topiramate with and without other medications have been received. Isolated reports have been received of hepatitis and hepatic failure occurring in patients taking multiple medications while being treated with topiramate.

Isolated reports have also been received for bullous skin and mucosal reactions (including Stevens-Johnson syndrome, toxic epidermal necrolysis, erythema multiforme and pemphigus). The majority of these reports have occurred in patients taking other medications that can be associated with bullous skin and mucosal reactions.

## **DRUG INTERACTIONS**

### **Drug-Drug Interactions**

In all of the studies shown below, except where noted, the maximum topiramate dose administered was 200 mg/day.

#### **Antiepileptic Drugs**

Potential interactions between topiramate and standard AEDs were measured in controlled clinical pharmacokinetic studies in patients with epilepsy. The effects of these interactions on plasma concentrations are summarized in Table 1.9.

**Table 1.9: Drug Interactions with Topiramate Therapy**

<b>AED Co-administered</b>	<b>AED Concentration</b>	<b>Topiramate Concentration</b>
--------------------------------	------------------------------	-------------------------------------

Phenytoin	↔ **	↓ 59%
Carbamazepine (CBZ)	↔	↓ 40%
CBZ epoxide*	↔	NS
Valproic acid	↓ 11%	↓ 14%
Phenobarbital	↔	NS
Primidone	↔	NS
Lamotrigine	↔	13% decrease

at Topiramate doses up to 400 mg/day

*	Is not administered but is an active metabolite of carbamazepine
↔	No effect on plasma concentration (≤ 15% change)
**	Plasma concentrations increased 25% in some patients, generally those on a b.i.d. dosing regimen of phenytoin
↓	Plasma concentrations decrease in individual patients
NS	Not studied
AED	Antiepileptic drug

### Effects of Topiramate on Other Antiepileptic Drugs

The addition of topiramate to other antiepileptic drugs (phenytoin, carbamazepine, valproic acid, phenobarbital, primidone) has no effect on their steady-state plasma concentrations, except in the occasional patient, where the addition of topiramate to phenytoin may result in an increase of plasma concentrations of phenytoin.

The effect of topiramate on steady-state pharmacokinetics of phenytoin may be related to the frequency of phenytoin dosing. A slight increase in steady-state phenytoin plasma concentrations was observed, primarily in patients receiving phenytoin in two divided doses. The slight increase may be due to the saturable nature of phenytoin pharmacokinetics and inhibition of phenytoin metabolism CYP2C19.

The addition of topiramate therapy to phenytoin should be guided by clinical outcome. In general, as evidenced in clinical trials, patients do not require dose adjustments. However, any patient on phenytoin showing clinical signs or symptoms of toxicity should have phenytoin levels monitored. The effects of these interactions on plasma concentrations are summarized in Table 1.9.

### Effects of Other Antiepileptic Drugs on Topiramate

#### Phenytoin and Carbamazepine

Phenytoin and carbamazepine decrease the plasma concentration of topiramate. The addition or withdrawal of phenytoin and/or carbamazepine during adjunctive therapy with topiramate may require adjustment of the dose of topiramate. This should be done by titrating to clinical effect.

#### Valproic Acid

The addition or withdrawal of valproic acid does not produce clinically significant changes in plasma concentrations of topiramate, and therefore, does not warrant dosage adjustment of topiramate. The effects of these interactions on plasma concentrations are summarized in Table 1.9.

Rare post-marketing reports of encephalopathy with or without hyperammonemia have been received for patients treated with topiramate alone or in combination with valproic acid or other antiepileptic medications. The majority of the cases reported concomitant administration of topiramate and valproic acid. Thus, caution is advised when polytherapy is necessary (see

## **ADVERSE REACTIONS, Post-Market Adverse Drug Reactions and WARNINGS AND PRECAUTIONS, General.**

Concomitant administration of **ratio-TOPIRAMATE** with valproic acid has also been associated with hypothermia (with and without hyperammonemia) in patients who have tolerated either drug alone. It may be prudent to examine blood ammonia levels in patients in whom the onset of hypothermia has been reported (see **WARNINGS AND PRECAUTIONS, Hypothermia with Concomitant Valproic Acid (VPA) Use**).

### **Other Drug Interactions**

*Digoxin:* In a single-dose study, serum digoxin AUC decreased 12% due to concomitant topiramate administration (200 mg/day). Multiple-dose studies have not been performed. When topiramate is added or withdrawn in patients on digoxin therapy, careful attention should be given to the routine monitoring of serum digoxin.

*CNS Depressants:* Concomitant administration of topiramate and alcohol or other CNS depressant drugs has not been evaluated in clinical studies. It is recommended that topiramate not be used concomitantly with alcohol or other CNS depressant drugs.

#### *Oral Contraceptives:*

##### *Topiramate (50-200 mg/day) in Healthy Volunteers*

In a pharmacokinetic interaction study in healthy volunteers, subjects were stratified into obese versus non-obese (n=12 versus n=12) with both groups concomitantly administered a combination oral contraceptive product containing 1 mg norethindrone plus 35 µg ethinyl estradiol and topiramate (50 to 200 mg/day) given in the absence of other medications. For the ethinyl estradiol component, both obese and non-obese volunteers showed a decrease in mean AUC and C<sub>max</sub> at 200 mg/day (-10.7% and -9.4% versus -15.2% and -11.3%, respectively) that were not statistically significant. Changes in individual subjects ranged from decreases of approximately 35% to 90% in 5 individuals to increases of approximately 35% to 60% in 3 individuals. At the 50 and 100 mg/day topiramate doses, similar changes in mean C<sub>max</sub> and AUC were observed for non-obese volunteers. The clinical significance of these changes is unknown. For the norethindrone component, only the non-obese group showed a decrease (-11.8%). In view of the dose-dependent decreases seen in the ethinyl estradiol component in epileptic patients receiving topiramate as adjunctive therapy (below), and the fact that the recommended dose is up to 400 mg/day, there may be greater decreases seen at doses above 200 mg/day as monotherapy.

##### *Topiramate as Adjunctive Therapy with Valproic Acid in Epileptic Patients*

In a pharmacokinetic interaction study, epileptic patients received topiramate as adjunctive therapy with valproic acid and a combination oral contraceptive product containing norethindrone (1 mg) plus ethinyl estradiol (35 µg). In this study, topiramate did not significantly affect the oral clearance of norethindrone. The serum levels of the estrogenic component decreased by 18%, 21% and 30% at daily doses of 200, 400 and 800 mg of topiramate, respectively. There are minimal clinical data regarding interaction of valproic acid and oral contraceptives.

In view of both of the above study findings, the efficacy of low-dose (e.g. 20 µg) oral

contraceptives may be reduced in both the monotherapy and adjunctive therapy situation with topiramate. For topiramate doses up to 200 mg/day, which includes the recommended dose for migraine prophylaxis of 100 mg/day, the mean reduction in norethindrone and ethinyl estradiol exposure from topiramate treatment is not significant, although marked changes in individual patients are possible. In the treatment of epilepsy at doses greater than 200 mg/day, significant dose-dependent decreases in ethinyl estradiol exposure are expected. Patients on topiramate doses greater than 200 mg/day who are taking oral contraceptives should receive a preparation containing not less than 30 µg of estrogen.

The possibility of decreased contraceptive efficacy and increased breakthrough bleeding should be considered in patients taking combination oral contraceptive products with **ratio-TOPIRAMATE**. Patients taking oral contraceptives should be asked to report any change in their bleeding patterns. Contraceptive efficacy can be decreased even in the absence of breakthrough bleeding.

*Hydrochlorothiazide (HCTZ):* A parallel-arm drug-drug interaction study conducted in healthy volunteers (12 males, 11 females) evaluated the steady-state pharmacokinetics of the diuretic HCTZ (25 mg q24h) and topiramate (96 mg q12h) when administered alone and concomitantly. The results of this study indicate that mean topiramate  $C_{max}$  increased by 27% and mean AUC increased by 29% when HCTZ was added to topiramate. The clinical significance of this statistically significant change is unknown. Thus, the concomitant use of topiramate and HCTZ may require a downward adjustment of the topiramate dose. The steady-state pharmacokinetics of HCTZ were not significantly influenced by the concomitant administration of topiramate. In addition, greater decreases in serum potassium were seen with concomitant treatment than with either drug alone, both in terms of percentage of patients with a serum potassium measurements of <3.6 mEq/L at the end of each treatment period [61% (14/23) with concomitant treatment versus 27% (3/11) with topiramate alone versus 25% (3/12) with HCTZ alone] and in mean change from baseline (approximately -0.60 mEq/L for concomitant treatment versus -0.25 mEq/L for topiramate alone versus -0.12 mEq/L for HCTZ alone). One of the subjects who had hypokalemia with concomitant treatment also had an abnormal ECG (non-specific ST-T wave changes), which may have been related to the decrease in plasma potassium levels. See also **WARNINGS AND PRECAUTIONS, Endocrine and Metabolism**.

*Metformin:* A drug-drug interaction study conducted in 18 healthy volunteers, ages 18-37, evaluated the steady-state pharmacokinetics of metformin and topiramate in plasma when metformin (500 mg b.i.d.) was given alone and when metformin and topiramate (50, 75 and 100 mg) were given simultaneously for 6 consecutive days. The results of this study indicated that metformin mean  $C_{max}$  and mean  $AUC_{0-12h}$  increased by 18% and 25%, respectively, while mean CL/F decreased 20% when metformin was co-administered with topiramate (up-titrated to 100 mg b.i.d.). Topiramate did not affect metformin  $t_{max}$ . The effects of higher doses of topiramate (>100 mg b.i.d.) on metformin are unknown. The clinical significance of the effect of topiramate on metformin pharmacokinetics is unclear. Oral plasma clearance of topiramate appears to be reduced when administered with metformin. The extent of change in the clearance is unknown. The clinical significance of the effect of metformin on topiramate pharmacokinetics is unclear. When topiramate is added or withdrawn in patients on metformin therapy, careful attention should be given to the routine monitoring for adequate control of their diabetic disease state.

*Glyburide:* A drug-drug interaction study conducted in 28 patients with type 2 diabetes, ages 38-68 years and BMIs 25-40 kg/m<sup>2</sup>, evaluated the steady-state pharmacokinetics of glyburide and topiramate in plasma when glyburide (5 mg/day) was given alone and when glyburide and topiramate (150 mg/day) were given concomitantly for 48 consecutive days. Glyburide systemic exposure was statistically significantly reduced when combined with topiramate such that mean C<sub>max</sub> and mean AUC<sub>24</sub> decreased by 22% and 25%, respectively, while mean CL/F increased by 21%. Systemic exposure of the active metabolites, 4-*trans*-hydroxyglyburide and 3-*cis*-hydroxyglyburide, was also statistically significantly reduced by 13% and 15%, respectively. The steady-state pharmacokinetics of topiramate were unaffected by concomitant administration of glyburide. The clinical significance of the effect of glyburide on topiramate pharmacokinetics is unclear. Mild to moderate declines in serum bicarbonate without metabolic acidosis were associated with the addition of topiramate (see **WARNINGS AND PRECAUTIONS, Endocrine and Metabolism, Metabolic Acidosis**). The effects of higher doses of topiramate (> 150 mg/day) on glyburide are unknown. When topiramate is added to glyburide therapy or glyburide is added to topiramate therapy, careful attention should be given to the routine monitoring of patients for adequate control of their diabetic disease state.

*Pioglitazone:* A drug-drug interaction study conducted in healthy volunteers (26 males, 26 females) evaluated the steady-state pharmacokinetics of topiramate and the antidiabetic agent, pioglitazone, when administered alone and concomitantly. The pharmacokinetic parameters of topiramate were not affected; mean pioglitazone AUC decreased by 15%, and mean C<sub>max</sub> increased non-significantly by 10%, but with individual subjects showing large increases, and 3 of the 4 highest values were recorded by males. In addition, each of the active hydroxy-metabolite and the active keto-metabolite showed mean decreases in C<sub>max</sub> and AUC (approximately 15% for the hydroxy-metabolite and 60% for the keto-metabolite). The clinical significance of these findings is not known. When topiramate is added to pioglitazone therapy or pioglitazone is added to topiramate therapy, careful attention should be given to the routine monitoring of patients for adequate control of their diabetic disease state.

### Lithium:

#### *Healthy Volunteers*

A drug-drug interaction study conducted in twelve healthy volunteers, ages 20-40 years, evaluated the steady-state pharmacokinetics of lithium in plasma when lithium (300 mg q8h) was administered for 14 days and topiramate (up-titrated to 100 mg q12h) was given concomitantly for the last 6 days. Based on the data analysis of twelve subjects, systemic exposure of lithium was statistically significantly reduced in the presence of topiramate such that the C<sub>max</sub> and AUC<sub>0-8h</sub> decreased by 20% and 18%, respectively, while mean CL/F and CL<sub>R</sub> increased by 36% and 12%, respectively. One subject did not have measurable trough lithium concentrations on Day 14, potentially indicating missed dose administration. By excluding this subject from the analyses, systemic exposure of lithium was slightly reduced in the presence of topiramate (12% for C<sub>max</sub>, 10% for AUC<sub>0-8</sub>) while mean CL/F and CL<sub>R</sub> increased by 11% and 16%, respectively. The clinical significance of the effect of topiramate on lithium pharmacokinetics is unclear. The effects of higher doses of topiramate (> 200 mg/day) on the pharmacokinetics of lithium are unknown.

#### *Patients with Bipolar Disorder*

A drug-drug interaction study conducted in 31 patients with various types of bipolar disorder, ages 20-60 years, evaluated the steady-state pharmacokinetics of lithium and topiramate when

administered concomitantly. Subjects were randomized to receive either low doses of topiramate of up to 200 mg/day or high doses of topiramate of up to 600 mg/day. Pharmacokinetic profiles for lithium were obtained following 1 week and 3 weeks of continuous lithium dosing. The pharmacokinetics of lithium were unaffected during treatment with topiramate at doses of up to 200 mg/day, and were unaffected by short-term treatment with topiramate (1 week) at doses up to 600 mg/day. Following treatment with topiramate at doses of up to 600 mg/day for 3 weeks, there was an observed statistically significant increase in systemic exposure of lithium (about 27% for both  $C_{max}$  and AUC). Topiramate exposure for both the low and high dose groups was similar following 1 week and 3 weeks of continuous treatment in the presence of lithium. The effects of higher doses of topiramate (> 600 mg/day) on lithium have not been studied and are unknown. Lithium levels should be monitored when co-administered with topiramate and dose adjustments for lithium should be based on both lithium levels and clinical outcome for the patient.

#### Risperidone:

##### *Healthy Volunteers*

A drug-drug interaction study was conducted in 12 healthy volunteers (6 males, 6 females), ages 28-40 years, with single-dose administration of risperidone (2 mg) and multiple doses of topiramate (titrated up to 200 mg/day). In the presence of topiramate, systemic exposure of the total active moiety (risperidone + 9-hydroxyrisperidone) was reduced such that mean  $AUC_{0-\infty}$  was 11% lower and mean  $C_{max}$  was statistically significantly (18%) lower. In the presence of topiramate, systemic exposure of risperidone was statistically significantly reduced such that mean  $C_{max}$  and  $AUC_{0-\infty}$  were 29% and 23% lower, respectively. The pharmacokinetics of 9-hydroxyrisperidone were unaffected. The effects of a single dose (2 mg/day) of risperidone on the pharmacokinetics of multiple doses of topiramate have not been studied. Therefore, patients receiving risperidone in combination with topiramate should be closely monitored for clinical response to risperidone.

##### *Patients with Bipolar Disorder*

A drug-drug interaction study conducted in 52 patients with various types of bipolar disorder (24 males, 28 females), ages 19-56 years, evaluated the steady-state pharmacokinetics of risperidone and topiramate when administered concomitantly. Eligible subjects were stabilized on a risperidone dose of 1-6 mg/day for 2 to 3 weeks. Topiramate was then titrated up to escalating doses of 100, 250 and 400 mg/day along with risperidone for up to 6 weeks. Risperidone was then tapered and discontinued over 4 weeks while maintaining topiramate (up to 400 mg/day). There was a statistically significant reduction in risperidone systemic exposure (16% and 33% for  $AUC_{12}$  and 13% and 34% for  $C_{max}$  at the 250 and 400 mg/day doses, respectively). Minimal alterations were observed in the pharmacokinetics of the total active moiety (risperidone plus 9-hydroxyrisperidone) and 9-hydroxyrisperidone. Topiramate systemic exposure was slightly reduced (12.5% for mean  $C_{max}$  and 11% for mean  $AUC_{12}$ ) in the presence of risperidone, which achieved statistical significance. There were no clinically significant changes in the systemic exposure of the risperidone total active moiety or of topiramate. The effects of higher doses of topiramate (> 400 mg/day) are unknown. Patients with bipolar disorder receiving risperidone in combination with topiramate should be closely monitored for clinical response to risperidone.

Haloperidol: The pharmacokinetics of a single dose of the antipsychotic haloperidol (5 mg) were not affected following multiple dosing of topiramate (200 mg/day) in 13 healthy adults (6 males,

7 females).

Venlafaxine: A drug-drug interaction study was conducted in 26 healthy volunteers (16 males/10 females, ages 18-40 years, BMI ranging from 25 to 30 kg/m<sup>2</sup>) to evaluate the interaction between venlafaxine and topiramate. Subjects received single 150-mg doses of extended release venlafaxine and multiple doses of topiramate titrated up to 150 mg/day. The single-dose pharmacokinetics of venlafaxine were unaffected by treatment with topiramate. While the  $C_{max}$ ,  $AUC_{\infty}$  and  $CL/F$  of the active metabolite, O-desmethylvenlafaxine were unaffected, the renal clearance of the active metabolite was increased by 53% during the treatment with topiramate. These observed increases in urinary excretion of O-desmethylvenlafaxine during treatment with topiramate did not affect systemic exposure. The steady-state pharmacokinetics of topiramate were unaffected by repeated daily-dose administration of venlafaxine for 5 days. The effects of higher doses of topiramate (> 150 mg/day) on the pharmacokinetics of venlafaxine and higher doses of venlafaxine up to the maximum dose of 375 mg/day on the pharmacokinetics of topiramate are unknown.

Amitriptyline: There was a 12% increase in both AUC and  $C_{max}$  for the tricyclic antidepressant amitriptyline (25 mg/day) in 18 normal subjects (9 males, 9 females) receiving 200 mg/day of topiramate. Individual subjects experienced large changes in amitriptyline concentration, either up or down, in the presence of topiramate; any adjustments in amitriptyline dose should be made according to patients' clinical response and not on the basis of plasma levels.

Pizotifen: Multiple dosing of topiramate (200 mg/day) in 19 healthy volunteers (12 males, 7 females) had little effect on the pharmacokinetics of the anti-histamine pizotifen following daily 1.5 mg doses. There was a mean 12% and 15% decrease respectively in topiramate  $C_{max}$  and AUC in the volunteers (12 males and 7 females) receiving 200 mg/day topiramate and 1.5 mg/day pizotifen. This is not considered to be clinically significant.

Dihydroergotamine: Multiple dosing of topiramate (200 mg/day) in 24 healthy volunteers (12 males, 12 females) had little effect on the pharmacokinetics of a 1 mg subcutaneous dose of dihydroergotamine and a 1 mg subcutaneous dose of dihydroergotamine similarly had little effect on the pharmacokinetics of a 200 mg/day dose of topiramate.

Sumatriptan: Multiple dosing of topiramate (200 mg/day) in 24 healthy volunteers (14 males, 10 females) had little effect on the pharmacokinetics of single doses of the anti-migraine medication sumatriptan, either orally (100 mg) or subcutaneously (6 mg).

Propranolol: Multiple dosing of topiramate (100, then 200, mg/day) in 34 healthy volunteers (17 males, 17 females) had little effect on the pharmacokinetics of propranolol following daily 160 mg doses. There was a 17% increase in  $C_{max}$  of the metabolite 4-OH propranolol at 100 mg/day topiramate. Propranolol doses of 80, then 160, mg/day in 39 volunteers (27 males, 12 females) had a dose-dependent effect on exposure to topiramate (200 mg/day), reaching approximately a 9% and 16% increase for  $C_{max}$  and a 9% and 17% increase for AUC at 80 and 160 mg/day propranolol respectively.

Diltiazem: A drug-drug interaction study was conducted in 28 healthy volunteers (13 males/15 females, ages 18-45 years and BMIs 25-35 kg/m<sup>2</sup>) to evaluate the interaction between topiramate and diltiazem. Eligible subjects received single 240-mg doses of extended-release diltiazem and

multiple doses of topiramate titrated to 150 mg/day. Systemic exposure of diltiazem was statistically significantly reduced during topiramate treatment, where  $C_{\max}$  and  $AUC_{\infty}$  were 10% and 25% lower, respectively, following single-dose administration. There was an increase in diltiazem CL/F by approximately 30%. Systemic exposure of the active metabolite, desacetyl diltiazem, was statistically significantly reduced during treatment with topiramate where  $C_{\max}$  and  $AUC_{36}$  were 27% and 18% lower, respectively. The single-dose pharmacokinetics of the active metabolite, N-demethyl-diltiazem, were unaffected by topiramate. Following repeated daily-dose administration of diltiazem for 5 days, steady-state systemic exposure of topiramate was greater during treatment with diltiazem, where  $C_{\max}$  and  $AUC_{12}$  were approximately 17% and 20% higher, respectively, and CL/F was 16% lower. The effects of higher doses of topiramate (> 150 mg/day) on the pharmacokinetics of diltiazem or its metabolites have not been studied. Overall, the clinical significance of these observations is unclear.

### Flunarizine:

#### *Patients with Migraine – Effects of topiramate on the pharmacokinetics of flunarizine*

The dose of flunarizine used in this study is one-half of the recommended daily dose. A drug-drug interaction study was conducted in forty seven patients with a history of migraine (13 males, 34 females, ages 20-53 years, evaluated the steady-state pharmacokinetics of flunarizine when topiramate was administered concomitantly. Subjects were taking flunarizine for at least 4 weeks before study start. One subgroup was administered only flunarizine (5 mg q24h) for 81 days, and, a second subgroup received flunarizine (5 mg q24h) for 81 days and topiramate (up-titrated to 50 mg/day and then to 100 mg/day) from Day 4 to a.m. dose on Day 82 concomitantly.

Mean  $C_{\max}$  of flunarizine decreased by 22% with concomitant administration of topiramate at 50 mg/day. During concomitant treatment with topiramate at 100 mg/day,  $C_{\max}$  estimates returned to those observed during treatment with flunarizine alone. Mean  $AUC_{0-24}$  for flunarizine was similar with concomitant administration of topiramate at 50 mg/day and 16% higher with topiramate at 100 mg/day compared to treatment with flunarizine alone. Mean CL/F of flunarizine was unaffected by treatment with topiramate. Systemic exposure of topiramate ( $C_{\max}$  and  $AUC_{0-12}$ ) doubled with increasing topiramate dose from 50 mg/day to 100 mg/day. Mean CL/F was similar during both dose periods and was consistent with previously observed estimates in healthy volunteers. These alterations are unlikely to be of clinical significance. However, there are no data on the effects of higher doses of topiramate on flunarizine levels. There is also no information on the interaction of topiramate and flunarizine in patients with history of seizure or epilepsy.

Agents Predisposing to Nephrolithiasis: Topiramate, when used concomitantly with other agents predisposing to nephrolithiasis, such as carbonic anhydrase inhibitors, e.g. acetazolamide, may increase the risk of nephrolithiasis. While using topiramate, agents like these should be avoided since they may create a physiological environment that increases the risk of renal stone formation (see **WARNINGS AND PRECAUTIONS, Renal**).

### **Drug-Food Interactions**

There was no clinically significant effect of food on the bioavailability of topiramate.

### **Drug-Herb Interactions**

Interactions with herbal products have not been established.



### **Drug-Laboratory Interactions**

There are no known interactions of topiramate with commonly used laboratory tests.

## **DOSAGE AND ADMINISTRATION**

### **General**

In patients with or without a history of seizures or epilepsy, **ratio-TOPIRAMATE** (topiramate) should be gradually withdrawn to minimize the potential for seizures or increased seizure frequency. (See **WARNINGS AND PRECAUTIONS, General**).

In clinical trials in adult patients with epilepsy, dosages were decreased by 50-100 mg/day at weekly intervals. In clinical trials of children, topiramate was gradually withdrawn over 2-8 week period.

In clinical trials in adult patients receiving topiramate for migraine prophylaxis dosages were decreased by 25-50 mg/day at weekly intervals. (See **WARNINGS AND PRECAUTIONS, General**).

In situations where rapid withdrawal of topiramate is medically required, appropriate monitoring is recommended. (See **WARNINGS AND PRECAUTIONS, General**).

### **Dosing Considerations**

- Patients with renal impairment
- Patients undergoing hemodialysis
- Patients with hepatic disease
- Prophylactic treatment of migraine: In pregnancy, the occurrence of seizures presents a significant risk for the mother and child. Prescribing **ratio-TOPIRAMATE** to prevent seizures therefore outweighs the risk of malformations to the fetus. However, taking **ratio-TOPIRAMATE** to prevent migraine attacks does not outweigh this risk. Consequently, **ratio-TOPIRAMATE** is contraindicated in pregnancy and in women of child bearing potential who are not using an effective method of contraception (see **CONTRAINDICATIONS**).

### **Recommended Dose and Dosage Adjustment**

**ratio-TOPIRAMATE** (topiramate) Tablets can be taken without regard to meals.

## **EPILEPSY**

### **Monotherapy**

#### **Adults and Children (Age 6 years and older)**

The recommended initial target dose for topiramate monotherapy in adults and children 6 years of age and older is 100 mg/day and the maximum recommended dose is 400 mg/day, administered in two divided doses, as needed and tolerated.

The recommended titration rate for topiramate monotherapy to 100 mg/day is:

	Week 1	Weeks 2-3	Weeks 3-4
Morning Dose	None	25 mg	50 mg
Evening Dose	25 mg	25 mg	50 mg

If doses above 100 mg/day are required, the dose may be increased at weekly intervals in increments of 50 mg/day to a maximum of 400 mg/day. Dose and titration rate should be guided by clinical outcome. Some patients may benefit from a slower titration schedule. Daily doses above 400 mg have not been adequately studied. Only 14 pediatric patients have received 500 mg/day topiramate in controlled clinical trials (see **ADVERSE REACTIONS, EPILEPSY, Clinical Trial Adverse Drug Reactions, Table 1.2**).

### **Adjunctive Therapy**

#### Adults (Age 17 years and older)

It is recommended that topiramate as adjunctive therapy be initiated at 50 mg/day, followed by titration as needed and tolerated to an effective dose. At weekly intervals, the dose may be increased by 50 mg/day and taken in two divided doses. Some patients may benefit from lower initial doses, e.g. 25 mg and/or a slower titration schedule. Some patients may achieve efficacy with once-a-day dosing.

The recommended total daily maintenance dose is 200-400 mg/day in two divided doses. Doses above 400 mg/day have not been shown to improve responses and have been associated with a greater incidence of adverse events. The maximum recommended dose is 800 mg/day. Daily doses above 1,600 mg have not been studied.

#### Children (Ages 2-16 years)

It is recommended that topiramate as adjunctive therapy be initiated at 25 mg (or less, based on a range of 1 to 3 mg/kg/day) nightly for the first week followed by titration as needed and tolerated to an effective dose. The dosage should then be increased at 1- or 2-week intervals by increments of 1 to 3 mg/kg/day (administered in two divided doses). Some patients may benefit from lower initial doses and/or a slower titration schedule.

The recommended total daily maintenance dose is approximately 5 to 9 mg/kg/day in two divided doses.

### **Drug Discontinuation**

In patients with a history of seizures or epilepsy, topiramate should be gradually withdrawn to minimize the potential for seizures or increased seizure frequency. In clinical trials, daily dosages were decreased in weekly intervals by 50-100 mg in adults with epilepsy.

In clinical trials of children, topiramate was gradually withdrawn over a 2-8 week period.

In situations where rapid withdrawal of topiramate is medically required, appropriate monitoring is recommended.

### **MIGRAINE PROPHYLAXIS**

## Adults

The usual total daily dose of topiramate as treatment for prophylaxis of migraine headache is 100 mg/day administered in two divided doses. Dose and titration rate should be guided by clinical outcome. If required, longer intervals between dose adjustments can be used. No extra benefit has been demonstrated from the administration of doses higher than 100 mg/day and the incidence of some adverse events increases with increasing dose (see **ADVERSE REACTIONS, MIGRAINE PROPHYLAXIS, Clinical Trial Adverse Drug Reactions, Table 1.7**).

The recommended titration rate for topiramate for migraine prophylaxis to 100 mg/day is:

	Morning Dose	Evening Dose
Week 1	None	25 mg
Week 2	25 mg	25 mg
Week 3	25 mg	50 mg
Week 4	50 mg	50 mg

## Drug Discontinuation

In patients without a history of seizures or epilepsy, topiramate should be gradually withdrawn to minimize the potential for seizures or increased seizure frequency. In clinical trials, daily dosages were decreased in weekly intervals by 25-50 mg in adults receiving topiramate at doses up to 100 mg/day for migraine prophylaxis.

In situations where rapid withdrawal of topiramate is medically required, appropriate monitoring is recommended.

## Pediatrics

The safety and efficacy of topiramate in the management or prevention of migraine in pediatrics have not been established.

## Patients with Renal Impairment

In renally impaired subjects (creatinine clearance less than 70 mL/min/1.73m<sup>2</sup>), one-half of the usual adult dose is recommended. Such patients will require a longer time to reach steady-state at each dose (see **WARNINGS AND PRECAUTIONS, Renal** and **ACTION AND CLINICAL PHARMACOLOGY, Special Populations and Conditions, Renal Insufficiency**).

## Patients Undergoing Hemodialysis

Topiramate is cleared by hemodialysis at a rate that is 4 to 6 times greater than a normal individual. Accordingly, a prolonged period of dialysis may cause topiramate concentration to fall below that required to maintain an antiseizure effect. To avoid rapid drops in topiramate plasma concentration during hemodialysis, a supplemental dose of topiramate may be required. The supplemental dose should take into account 1) the duration of dialysis, 2) the clearance rate of the dialysis system being used, and 3) the effective renal clearance of topiramate in the patient being dialyzed (see **WARNINGS AND PRECAUTIONS, Renal**).

## Patients with Hepatic Disease

In hepatically impaired patients, topiramate plasma concentrations are increased approximately 30%. This moderate increase is not considered to warrant adjustment of the topiramate dosing regimen. Initiate topiramate therapy with the same dose and regimen as for patients with normal hepatic function. The dose titration in these patients should be guided by clinical outcome, i.e. seizure control, and avoidance of adverse effects. Such patients will require a longer time to reach steady-state at each dose (see **ACTION AND CLINICAL PHARMACOLOGY, Special Populations and Conditions, Hepatic Insufficiency**).

### **Geriatrics**

See **WARNINGS AND PRECAUTIONS** section.

### **Missed Dose**

The missed dose should be taken as soon as possible. If it is almost time for the next dose, the missed dose should not be taken. Instead, the next scheduled dose should be taken. Doses should not be doubled.

### **Administration**

Tablets should not be broken.

## **OVERDOSAGE**

**For management of a suspected drug overdose, contact your regional Poison Control Centre immediately.**

Overdoses of topiramate have been reported. Signs and symptoms included convulsions, drowsiness, speech disturbances, blurred vision, diplopia, mentation impaired, lethargy, abnormal coordination, stupor, hypotension, abdominal pain, agitation, dizziness and depression. The clinical consequences were not severe in most cases but deaths have been reported after polydrug overdoses involving topiramate.

Topiramate overdose can result in severe metabolic acidosis (see **WARNINGS AND PRECAUTIONS, Endocrine and Metabolism, Metabolic Acidosis**).

The highest topiramate overdose reported was calculated to be between 96 and 110 g and resulted in coma lasting 20-24 hours followed by full recovery after 3 to 4 days.

In acute topiramate overdose, if the ingestion is recent, the stomach should be emptied immediately by lavage or by induction of emesis. Activated charcoal has been shown to adsorb topiramate *in vitro*. Treatment should be appropriately supportive. Hemodialysis has been shown to be an effective means of removing topiramate from the body. The patient should be well hydrated.

## **ACTION AND CLINICAL PHARMACOLOGY**

### **Pharmacodynamics**

Topiramate is a novel agent classified as a sulfamate substituted monosaccharide. Three pharmacological properties of topiramate are believed to contribute to its anticonvulsant activity.

First, topiramate reduces the frequency at which action potentials are generated when neurons are subjected to a sustained depolarization indicative of a state-dependent blockade of voltage-sensitive sodium channels. Second, topiramate markedly enhances the activity of GABA at some types of GABA receptors. Because the antiepileptic profile of topiramate differs markedly from that of the benzodiazepines, it may modulate a benzodiazepine-insensitive subtype of GABA<sub>A</sub> receptor. Third, topiramate antagonizes the ability of kainate to activate the kainate/AMPA subtype of excitatory amino acid (glutamate) receptors but has no apparent effect on the activity of N-methyl-D-aspartate (NMDA) at the NMDA receptor subtype.

In addition, topiramate inhibits some isoenzymes of carbonic anhydrase. This pharmacologic effect is much weaker than that of acetazolamide, a known carbonic anhydrase inhibitor, and is not thought to be a major component of topiramate's antiepileptic activity.

### **Pharmacokinetics**

Topiramate exhibits low intersubject variability in plasma concentrations and therefore has predictable pharmacokinetics. The pharmacokinetics of topiramate are linear with plasma clearance remaining constant and area under the plasma concentration curve increasing in a dose-proportional manner over a 100 to 400 mg single oral dose range in healthy subjects. Patients with normal renal function may take 4 to 8 days to reach steady-state plasma concentrations. The mean C<sub>max</sub> following multiple twice-a-day oral doses of 100 mg to healthy subjects was 6.76 µg/mL. The mean plasma elimination half-lives from multiple 50 mg and 100 mg q12h doses of topiramate were approximately 21 hours. The elimination half-life did not significantly change when switching from single dose to multiple dose.

In well-controlled add-on trials, no correlation has been demonstrated between trough plasma concentrations and its clinical efficacy. It is not necessary to monitor topiramate plasma concentrations to optimize therapy with topiramate.

No evidence of tolerance requiring increased dosage has been demonstrated in patients during 5 years of use.

Concomitant multiple-dose administration of topiramate, 100 to 400 mg q12h, with phenytoin or carbamazepine shows dose-proportional increases in plasma concentrations of topiramate.

**Absorption:** Topiramate is rapidly and well absorbed. Following oral administration of 100 mg topiramate to healthy subjects, a mean peak plasma concentration (C<sub>max</sub>) of 1.5 µg/mL was achieved within 2 to 3 hours (T<sub>max</sub>). The mean extent of absorption from a 100 mg oral dose of <sup>14</sup>C-topiramate was at least 81% based on the recovery of radioactivity from the urine.

There was no clinically significant effect of food on the bioavailability of topiramate.

**Distribution:** Approximately 13% to 17% of topiramate is bound to plasma proteins. A low capacity binding site for topiramate in/on erythrocytes that is saturable above plasma concentrations of 4 µg/mL has been observed.

The volume of distribution varied inversely with the dose. The mean apparent volume of distribution was 0.80 to 0.55 L/kg for a single-dose range of 100 to 1200 mg.

**Metabolism:** Topiramate is not extensively metabolized ( $\approx 20\%$ ) in healthy volunteers. It is metabolized up to 50% in patients receiving concomitant antiepileptic therapy with known inducers of drug metabolizing enzymes. Six metabolites formed through hydroxylation, hydrolysis and glucuronidation, have been isolated, characterized and identified from plasma, urine and feces of humans. Each metabolite represents less than 3% of the total radioactivity excreted following administration of  $^{14}\text{C}$ -topiramate.

Two metabolites, which retained most of the structure of topiramate, were tested and found to have little or no pharmacological activity.

**Excretion:** In humans, the major route of elimination of unchanged topiramate and its metabolites is via the kidney (at least 81% of the dose). Approximately 66% of a dose of  $^{14}\text{C}$ -topiramate was excreted unchanged in the urine within 4 days. The mean renal clearance for 50 mg and 100 mg of topiramate, following q12h dosing, was approximately 18 mL/min and 17 mL/min, respectively. Evidence exists for renal tubular reabsorption of topiramate. This is supported by studies in rats where topiramate was co-administered with probenecid, and a significant increase in renal clearance of topiramate was observed. This interaction has not been evaluated in humans. Overall, plasma clearance (CL/F) is approximately 20 to 30 mL/min in humans following oral administration.

### **Special Populations and Conditions**

**Pediatrics:** Pharmacokinetics of topiramate were evaluated in patients ages 4 to 17 years receiving one or two other antiepileptic drugs. Pharmacokinetic profiles were obtained after one week at doses of 1, 3, and 9 mg/kg/day. As in adults, topiramate pharmacokinetics were linear with clearance independent of dose and steady-state plasma concentrations increasing in proportion to dose. Compared with adult epileptic patients, mean topiramate clearance is approximately 50% higher in pediatric patients. Steady-state plasma topiramate concentrations for the same mg/kg dose are expected to be approximately 33% lower in children compared to adults. As with adults, hepatic enzyme-inducing antiepileptic drugs (AEDs) decrease the plasma concentration of topiramate.

**Geriatrics:** Plasma clearance of topiramate is unchanged in elderly subjects in the absence of underlying renal disease.

**Race, Gender and Age:** Although direct comparison studies of pharmacokinetics have not been conducted, analysis of plasma concentration data from clinical efficacy trials have shown that race, gender and age appear to have no effect on the plasma clearance of topiramate. In addition, based on pooled analyses, race and gender appear to have no effect on the efficacy of topiramate.

**Hepatic Insufficiency:** The pharmacokinetics of a single 100 mg oral dose of topiramate were evaluated in subjects with moderate to severe hepatic impairment ( $n=5$ ) and in six healthy subjects in which five of the healthy subjects were demographically matched to the five hepatically impaired subjects. Plasma topiramate concentrations in the hepatically impaired group increased ( $C_{\max}$  28.9% and  $\text{AUC}_{(0-\infty)}$  29.2%) with respect to the healthy subjects, due to an approximate 26% decrease in topiramate oral plasma clearance. The decrease in topiramate oral plasma clearance (CL/F) was primarily due to a 49% decrease in renal clearance. The reason for this decrease in renal clearance in hepatically impaired subjects is not known. Therefore,

**ratio-TOPIRAMATE** should be administered with caution in patients with hepatic impairment (see **DOSAGE AND ADMINISTRATION, Patients with Hepatic Disease**).

**Renal Insufficiency:** The pharmacokinetics of a single 100 mg oral dose of topiramate were evaluated in patients with moderate or severe renal impairment (seven patients per group) and were compared to seven demographically matched subjects with normal renal function. Compared to normal subjects, the overall oral plasma clearance (CL/F) of topiramate was reduced by 42% and 54% in patients with moderate and severe renal impairment, respectively. The respective renal clearance values decreased by 54% and 77%. As a result, mean plasma exposure (AUC) values in moderate and severe renal impairment increased by 1.9- and 2.2-fold, respectively. Overall, higher steady-state topiramate plasma AUC is expected for a given dose in renally impaired patients as compared to those with normal renal function. In addition, patients with renal impairment will require a longer time to reach steady-state at each dose. In patients with moderate and severe renal impairment, half of the usual starting and maintenance dose is recommended (see **WARNINGS AND PRECAUTIONS, Renal** and **DOSAGE AND ADMINISTRATION, Patients with Renal Impairment**).

**Hemodialysis:** Topiramate is effectively removed from plasma by hemodialysis (see **DOSAGE AND ADMINISTRATION, Patients Undergoing Hemodialysis**).

## **STORAGE AND STABILITY**

**ratio-TOPIRAMATE** Tablets should be stored between 15 - 30°C in tightly closed light resistance containers. Protect from moisture.

## **DOSAGE FORMS, COMPOSITION AND PACKAGING**

### **Availability of Dosage Forms**

**ratio-TOPIRAMATE** (topiramate) is available as the following:

#### **25 mg tablets:**

White, round biconvex, beveled-edge coated tablet engraved *T63* on one side and *rph* on the other. Available in bottles of 100 tablets.

#### **100 mg tablets:**

Yellow, round biconvex, beveled-edge coated tablet engraved *T62* on one side and *rph* on the other. Available in bottles of 100 tablets.

#### **200 mg tablets:**

Salmon colored round biconvex coated tablet engraved *T61* on one side and *rph* on the other. Available in bottles of 100 tablets.

### **Composition**

**ratio-TOPIRAMATE** (topiramate) contains topiramate as the active ingredient. The tablets also contain the following inactive ingredients: pregelatinized starch, lactose monohydrate, colloidal silicon dioxide, sodium starch glycolate, microcrystalline cellulose and magnesium

stereate, hydroxypropyl methylcellulose, titanium dioxide, polydextrose, triacetin and polyethylene glycol. The colourants that are present in the tablets are as follows: 100 mg: yellow iron oxide and 200 mg: red iron oxide.



## PART II: SCIENTIFIC INFORMATION

### PHARMACEUTICAL INFORMATION


#### Drug Substance

Proper name: Topiramate

Chemical name: 2,3:4,5-bis-*O*-(1-methylethylidene)- $\beta$ -D-fructopyranose sulfamate

Molecular formula and molecular mass:  $C_{12}H_{21}NO_8S$ , 339.36 g/mol

Structural formula:



Physicochemical properties: Topiramate is a white crystalline powder having a bitter taste. Topiramate is most soluble in alkaline solutions containing sodium hydroxide or sodium phosphate with a pH of 9 to 10. It is freely soluble in acetone, chloroform, dimethylsulfoxide and ethanol. The solubility in water is 9.8 mg/mL. Its saturated solution has a pH of 6.3.

## CLINICAL TRIALS

### Comparative Bioavailability Studies

A single-dose, cross over comparative bioavailability study was performed using 20 healthy adult male volunteers under fasted conditions. The rate and extent of absorption of **ratio-TOPIRAMATE** (topiramate) measured and compared following oral administration of a 100 mg dose of Topiramate (1 x 100 mg) tablet or Topamax<sup>®</sup> (1 x 100 mg) tablet. The results from measured data are summarized as follows:

**Table 2.2: Summary Table of the Comparative Bioavailability Data**

TOPIRAMATE (1 x 100 mg Tablets) From measured data				
Parameter	Geometric Mean Arithmetic Mean (CV%)		% Ratio of Geometric Means	90 % Confidence Interval
	Test ratio-TOPIRAMATE	Reference Topamax <sup>®</sup> †		
AUC <sub>0-72</sub> (mg.h/mL)	51225.872 51950.28 (16.57)	54879.766 <sup>‡</sup> 55510.19 (15.55) <sup>‡</sup>	92.82%	85-53-100.73%
AUC <sub>∞</sub> (mg.h/mL)	62592.45 63574.66 (17.17)	67819.16 68752.37 (17.38)	92.30%	84.42-100.91%
C <sub>MAX</sub> (mg/mL)	1960.1965 1985.83 (17.59)	2020.0954 2054.39 (18.98)	97.03%	89.95-104.68%
T <sub>MAX</sub> <sup>§</sup> (h)	1.27 (57.93)	1.25 (76.55)		
T <sub>1/2</sub> <sup>§</sup> (h)	29.58 (9.50)	30.51 (12.69)		

<sup>‡</sup> n= 18

<sup>§</sup> Expressed as the arithmetic mean (CV%) only.

<sup>†</sup> Topamax<sup>®</sup> is manufactured by Janssen-Ortho Inc. and was purchased in Canada.

Based on these results, the ratiopharm and Janssen-Ortho Inc. (Topamax<sup>®</sup>) 100 mg topiramate tablets are bioequivalent under fasting conditions.

A single-dose, cross over comparative bioavailability study was performed using 20 healthy adult male volunteers under fasted conditions. The rate and extent of absorption of topiramate measured and compared following oral administration of a 200 mg dose of **ratio-TOPIRAMATE** (1 x 200 mg) tablet or Topamax<sup>®</sup> (1 x 200 mg) tablet. The results from measured data are summarized in as follows:

**Table 2.3: Summary Table of the Comparative Bioavailability Data**

TOPIRAMATE (1 x 200 mg Tablets) From measured data				
Parameter	Geometric Mean Arithmetic Mean (CV%)		% Ratio of Geometric Means	90 % Confidence Interval
	Test ratio-TOPIRAMATE	Reference Topamax <sup>®</sup> †		
AUC <sub>0-72</sub> (mg.h/mL)	121188.20 122570.26 (15.19)	121455.11 123591.93 (18.59)	99.78%	96.58-103.08%
AUC <sub>∞</sub> (mg.h/mL)	145146.67 147221.25 (16.74)	144842.18 147638.13 (18.92)	100.21%	96.38-104.19%
C <sub>MAX</sub> (mg/mL)	4713.5659 4825.81 (22.33)	4332.9332 4472.50 (27.50)	108.79%	98.07-120.68%
T <sub>MAX</sub> <sup>§</sup> (h)	1.82 (69.30)	1.81 (81.27)		
T <sub>½</sub> <sup>§</sup> (h)	27.79 (18.05)	27.38 (14.09)		

§ Expressed as the arithmetic mean (CV%) only.

† Topamax<sup>®</sup> is manufactured by Janssen-Ortho Inc. and was purchased in Canada.

Based on these results, the ratiopharm and Janssen-Ortho Inc. (Topamax<sup>®</sup>) 200 mg topiramate tablets are bioequivalent under fasting conditions.

## EPILEPSY

### Monotherapy Controlled Trials

The effectiveness of topiramate as monotherapy in adults and children 6 years of age and older with newly diagnosed epilepsy was established in a multicentre, randomized, double-blind, parallel-group trial that compared the safety and efficacy of 2 doses of topiramate as monotherapy for the treatment of newly diagnosed or recurrent epilepsy.

The trial was conducted in 487 patients (6 to 83 years of age) who had a new diagnosis of epilepsy (partial onset or generalized) or a diagnosis of recurrent epilepsy while not taking antiepileptic drugs (AEDs). Patients who had either 1 or 2 well-documented seizures during the 3-month retrospective baseline phase entered the study and received topiramate 25 mg/day for 7 days in an open-label fashion. Any AED therapy used for temporary or emergency purposes was discontinued prior to randomization. Following that phase, patients were randomized to receive topiramate 50 mg/day or topiramate 400 mg/day. Patients remained in the double-blind phase until they experienced a first partial onset or generalized tonic-clonic seizure, until termination of the double-blind phase 6 months after randomization of the last subject, or until withdrawal for protocol-specified reasons. The primary efficacy assessment was based on the comparison between topiramate dose groups with respect to time to first partial onset or generalized tonic-clonic seizure during the double-blind phase. Comparison of the Kaplan-Meier survival curves of time to first seizure favoured topiramate 400 mg/day over topiramate 50 mg/day (p=0.0002, log rank test). The separation between the groups in favour of the higher dose group occurred early in the titration phase and was statistically significant as early as 2 weeks post randomization

( $p=0.046$ ), when, by following the weekly titration schedule, the subjects in the higher dose group had achieved a maximum topiramate dose of 100 mg/day. The higher dose group was also superior to the lower dose group with respect to the proportion of subjects who remained seizure-free, based on the Kaplan-Meier estimates, for a minimum of 6 months of therapy (82.9% vs. 71.4%;  $p=0.005$ ), and for a minimum of 1 year of therapy (75.7% vs. 58.8%;  $p=0.001$ ). The ratio of hazard rates for time to first seizure was 0.516 (95% confidence interval, 0.364 to 0.733). The treatment effects with respect to time to first seizure were consistent across various subject subgroups defined by age, sex, geographic region, baseline body weight, baseline seizure type, time since diagnosis, and baseline AED use.

### **Adjunctive Therapy Controlled Trials in Adults with Partial Onset Seizures**

The effectiveness of topiramate as adjunctive therapy in adults with refractory partial onset seizures with or without secondarily generalized seizures was established in six multicentre, outpatient, randomized, double-blind, placebo-controlled trials. Patients in all six studies were permitted a maximum of two antiepileptic drugs (AEDs) in addition to topiramate therapy (target doses of 200, 400, 600, 800, or 1,000 mg/day) or placebo.

In all six add-on trials, the primary efficacy measurement was reduction in seizure rate from baseline during the entire double-blind phase; responder rate (fraction of patients with at least a 50% reduction) was also measured. The median percent reductions in seizure rates and the responder rates by treatment group for each study are shown in Table 2.1.

**Table 2.1: Median Percent Seizure Rate Reduction and Percent Responders in Six Double-Blind, Placebo-Controlled, Add-On Trials in Adults with Partial Onset Seizures**

Protocol	Efficacy results	Placebo	Target Topiramate Dosage (mg/day)				
			200	400	600	800	1,000
YD	n	45	45	45	46	–	–
	Median % Reduction	13.1	29.6 <sup>a</sup>	47.8 <sup>c</sup>	44.7 <sup>d</sup>	–	–
	% Responders	18	27	47 <sup>b</sup>	46 <sup>b</sup>	–	–
YE	n	47	–	–	48	48	47
	Median % Reduction	1.2	–	–	40.7 <sup>d</sup>	41.0 <sup>d</sup>	37.5 <sup>d</sup>
	% Responders	9	–	–	44 <sup>d</sup>	40 <sup>c</sup>	38 <sup>c</sup>
Y1	n	24	–	23	–	–	–
	Median % Reduction	1.1	–	40.7 <sup>a</sup>	–	–	–
	% Responders	8	–	35 <sup>b</sup>	–	–	–
Y2	n	30	–	–	30	–	–
	Median % Reduction	-12.2	–	–	46.4 <sup>c</sup>	–	–
	% Responders	10	–	–	47 <sup>c</sup>	–	–
Y3	n	28	–	–	–	28	–
	Median % Reduction	-17.8	–	–	–	35.8 <sup>c</sup>	–
	% Responders	0	–	–	–	43 <sup>c</sup>	–
YF/YG	n	42	–	–	–	–	167
	Median % Reduction	1.2	–	–	–	–	50.8 <sup>d</sup>
	% Responders	19	–	–	–	–	52 <sup>d</sup>

Comparisons with placebo: <sup>a</sup>p > 0.05; <sup>b</sup>p < 0.05; <sup>c</sup>p ≤ 0.01; <sup>d</sup>p ≤ 0.001

Across the six efficacy trials in adults, 232 of the 527 topiramate patients (44%) responded to treatment with at least a 50% seizure reduction during the double-blind phase; by comparison, only 25 of the 216 placebo-treated patients (12%) showed the same level of treatment response. When the treatment response was defined more rigorously as a 75% or greater decrease from baseline in seizure rate during double-blind treatment, 111 of the 527 topiramate patients (21%) in the 200 to 1,000 mg/day groups, but only 8 of the 216 placebo patients (4%), demonstrated this level of efficacy. In addition, 24 (5%) of the patients treated with topiramate became seizure-free, compared with 0% in the placebo group ( $p \leq 0.01$ ). At target dosages of 400 mg/day and higher, the percent of treatment responders was statistically greater for patients treated with topiramate than placebo-treated patients.

Pooled analyses of secondarily generalized seizure rates for all patients who had this seizure type during the studies show statistically significant percent reductions in the topiramate groups when compared with placebo. The median percent reduction in the rate of generalized seizures was 57% for patients treated with topiramate compared with -4% for placebo-treated patients. Among patients treated with topiramate, 109 (55%) of 198 had at least a 50% reduction in generalized seizure rate compared with 24 (27%) of 88 placebo-treated patients.

The dose titration in the original clinical trials was 100 mg/day the first week, 100 mg b.i.d., the second week, and 200 mg b.i.d. the third week. In a 12-week, double-blind trial, this titration rate was compared to a less rapid rate beginning at 50 mg/day. There were significantly fewer adverse experiences leading to discontinuation and/or dosage adjustment in the group titrated at

the less rapid rate. Seizure rate reductions were comparable between the groups at all time points measured.

### **Adjunctive Therapy Controlled Trials in Children with Partial Onset Seizures**

The effectiveness of topiramate as an adjunctive treatment for children with partial onset seizures was established in a multicentre, randomized, double-blind, placebo-controlled trial comparing topiramate and placebo in patients with a history of partial onset seizures, with or without secondarily generalized seizures.

Patients in this study were permitted a maximum of two antiepileptic drugs (AEDs) in addition to topiramate or placebo. Patients were stabilized on optimal dosages of their concomitant AEDs during an 8-week baseline phase. Included were patients who experienced at least six partial onset seizures, with or without secondarily generalized seizures, during the baseline phase.

Following randomization, patients began the double-blind phase of treatment. Patients received active drug beginning at 25 or 50 mg/day; the dose was then increased by 25 to 150 mg/day increments every other week until the assigned dosage of 125, 175, 225, or 400 mg/day, based on patient's weight to approximate a dosage of 6 mg/kg per day, was reached. After titration, patients entered an 8-week stabilization period.

The reduction in seizure rate from baseline during the entire double-blind phase was measured. The median percent reduction in seizure rate and the responder rate (fraction of patients with at least a 50% reduction) were also measured and the key results are shown in Table 2.2.

**Table 2.2: Median Percent Seizure Rate Reduction and Percent Responders in a Double-Blind, Placebo-Controlled, Add-On Trial in Pediatric Patients with Partial Onset Seizures**

Protocol	Efficacy results	Target Topiramate Dosage		
		Placebo	6 mg/kg/day*	p value
YP	n	45	41	
	Median % Reduction	10.5	33.1	0.034
	% Responders	20	39	0.08

\* For Protocol YP, protocol-specified target dosages (<9.3 mg/kg/day) were assigned based on subject's weight to approximate a dosage of 6 mg/kg/day; these dosages corresponded to mg/day dosages of 125, 175, 225, and 400 mg/day.

Forty patients received topiramate during the double-blind study and continued topiramate treatment in the open-label study. During the open-label study, dose escalation was permitted if required. The percent responders increased to 53% at a median average dose of 7.5 mg/kg/day.

### **Additional Adjunctive Therapy Clinical Data**

Some data demonstrating efficacy of topiramate as adjunctive therapy in adults and a small number of pediatric patients for primary generalized tonic-clonic seizures and seizures associated with Lennox-Gastaut syndrome are available from randomized, double-blind, placebo-controlled trials.

In the epilepsy clinical trials in approximately 1300 patients, daily dosages were decreased when required in weekly intervals by 50 to 100 mg in adults and over a 2 to 8 week period in children;

transition was permitted to a new antiepileptic regimen when clinically indicated.

## MIGRAINE PROPHYLAXIS

### **Controlled Trials in the Prophylactic Treatment of Migraine**

The results of 2 multicentre, randomized, double-blind, placebo-controlled, parallel-group clinical trials established the effectiveness of topiramate in the prophylactic treatment of migraine headache. The design of both trials was identical, enrolling patients with a history of migraine, with or without aura, for at least 6 months, according to the International Headache Society diagnostic criteria. Patients with a history of cluster headaches or basilar, ophthalmoplegic, hemiplegic, or transformed migraine headaches were excluded from the trials. Patients were required to have completed a washout of any prior migraine preventive medications before starting the baseline phase.

Patients who experienced 3 to 12 migraine periods (each migraine period was defined as any occurrence of migraine headache that started and ended, or recurred within a 24-hour interval) over the 4 weeks in the baseline phase were equally randomized to either topiramate 50 mg/day, 100 mg/day, 200 mg/day, or placebo and treated for a total of 26 weeks (8-week titration period and 18-week maintenance period). Treatment was initiated at 25 mg/day for one week, and then the daily dosage was increased by 25-mg increments each week until reaching the assigned target dose or maximum tolerated dose (administered twice daily). Up to 2 dose adjustments were allowed after the second week of treatment during the double-blind phase if unacceptable tolerability problems occurred. When needed, rescue medications were allowed for the acute treatment of headache or migraine-associated symptoms.

Effectiveness of treatment was assessed through the reduction in migraine headache frequency, as measured by the change in 4-week migraine period rate from the baseline phase to double-blind treatment in each topiramate treatment group compared to placebo.

In the first study, a total of 469 patients (416 females, 53 males), ranging in age from 13 to 70 years were randomized and provided efficacy data. Two hundred and sixty-five patients completed the entire 26-week double-blind phase. The median average daily dosages were 47.8 mg/day, 88.3 mg/day, and 132.1 mg/day in the target dose groups of topiramate 50, 100, and 200 mg/day, respectively.

The mean migraine headache frequency rate at baseline was approximately 5.5 migraine headaches/28 days and was similar across treatment groups. The change in the mean 4-week migraine headache frequency from baseline to the double-blind phase was -1.3, -2.1, and -2.2 in the topiramate 50, 100, and 200 mg/day groups, respectively, versus -0.8 in the placebo group (see Figure 2.1). The differences between the topiramate 100 and 200 mg/day groups versus placebo were statistically significant ( $p < 0.001$  for both comparisons: confidence intervals vs. placebo: topiramate 100 mg/day [-1.93, -0.55], and topiramate 200 mg/day [-2.04, -0.62]). The changes in migraine frequency represent a median percent reduction of 31%, 53%, and 55% in the topiramate 50, 100, and 200 mg/day groups, respectively, versus 21% in the placebo group.

In the second study, a total of 468 patients (406 females, 62 males), ranging in age from 12 to 65 years, were randomized and provided efficacy data. Two hundred and fifty-five patients completed the entire 26-week double-blind phase. The median average daily dosages were


46.5 mg/day, 85.6 mg/day, and 150.2 mg/day in the target dose groups of topiramate 50, 100, and 200 mg/day, respectively.

The mean migraine headache frequency rate at baseline was approximately 5.5 migraine headaches/28 days and was similar across treatment groups. The change in the mean 4-week migraine headache period frequency from baseline to the double-blind phase was -1.4, -2.1, and -2.4 in the topiramate 50, 100, and 200 mg/day groups, respectively, versus -1.1 in the placebo group (see Figure 2.1). The differences between the topiramate 100 and 200 mg/day groups versus placebo were statistically significant ( $p=0.008$  and  $p < 0.001$ , respectively; confidence intervals vs. placebo: topiramate 100 mg/day [-1.76, -0.27], and topiramate 200 mg/day [-2.06, -0.57]). The changes in migraine frequency represent a median percent reduction of 35%, 49%, and 48% in the topiramate 50, 100, and 200 mg/day groups, respectively, versus 19% in the placebo group.

In both studies, there were no apparent differences in treatment effect within age, gender, or racial subgroups.

In the migraine prophylaxis clinical trials in approximately 900 patients, daily dosages were decreased when required in weekly intervals by 25 to 50 mg in adults receiving topiramate at doses up to 100 mg/day.

**Figure 2.1: Reduction in 4-Week Migraine Headache Frequency (Studies TOPMAT-MIGR-001 and TOPMAT-MIGR-002)**



Additional efficacy measures that were assessed, in both studies, included responder rate, cumulative response rate, change in average monthly migraine attack rate, change in the average monthly rate of rescue medication use, change in the average number of monthly migraine days



and onset of action defined as the earliest month that there was a statistically significant difference between each topiramate treatment group and placebo with respect to the primary efficacy endpoint that was maintained for the remainder of the double-blind phase.

## DETAILED PHARMACOLOGY

### **Preclinical**

#### ***In Vitro* Studies**

Electrophysiological and biochemical studies on cultured neurons have revealed three properties that may contribute to the antiepileptic efficacy of topiramate. Action potentials elicited repetitively by a sustained depolarization of the neurons were blocked by topiramate in a time-dependent manner, suggestive of a state-dependent sodium channel blocking action. Topiramate increased the frequency at which  $\gamma$ -aminobutyrate (GABA) activated GABA<sub>A</sub> receptors, and enhanced the ability of GABA to induce a flux of chloride ions into neurons, suggesting that topiramate potentiates the activity of this inhibitory neurotransmitter.

Because the antiepileptic profile of topiramate differs markedly from that of the benzodiazepines, it may modulate a benzodiazepine-insensitive subtype of GABA<sub>A</sub> receptor. Topiramate antagonized the ability of kainate to activate the kainate/AMPA ( $\alpha$ -amino-3-hydroxy-5-methylisoxazole-4-propionic acid) subtype of excitatory amino acid (glutamate) receptor, but had no apparent effect on the activity of N-methyl-D-aspartate (NMDA) at the NMDA receptor subtype. These effects of topiramate were concentration-dependent over a range of 1  $\mu$ M to 200  $\mu$ M, with minimum activity observed at 1  $\mu$ M to 10  $\mu$ M.

In addition, topiramate inhibits some isoenzymes of carbonic anhydrase. This pharmacologic effect is much weaker than that of acetazolamide, a known carbonic anhydrase inhibitor, and is not thought to be a major component of topiramate's antiepileptic activity.

#### ***In Vivo* Studies**

##### **Pharmacodynamics**

Topiramate was initially found to possess anticonvulsant activity in the maximal electroshock seizure (MES) test in mice. Subsequent studies revealed that topiramate was also highly effective in the MES test in rats. In both species, anticonvulsant activity was evident within 30 minutes after oral administration, reached a peak 1 to 6 hours after dosing, and gradually declined thereafter.

Topiramate's anticonvulsant activity in rodents was further evaluated using chemical convulsants (pentylenetetrazole, bicuculline, picrotoxin, strychnine) to induce clonic or tonic seizures. Topiramate was either weak or inactive in blocking chemically induced seizures.

Topiramate was found to effectively block seizures in mouse and rat models of hereditary epilepsy, in some animal models of kindled epilepsy, and in a rat model of stroke-induced epilepsy. In the spontaneous epileptic rat (SER) model of hereditary epilepsy, topiramate blocked the clonic motor seizures and the absence-like seizures monitored by EEG recordings.

The potency of topiramate in blocking MES seizures is similar to that of phenytoin and carbamazepine, and much greater than that of valproate. The oral ED<sub>50</sub> of topiramate at the time of peak activity was 20 to 50 mg/kg in mice and 5 to 15 mg/kg in rats.

Studies in mice receiving concomitant administration of topiramate and carbamazepine or phenobarbital showed synergistic anticonvulsant activity, while combination with phenytoin showed additive anticonvulsant activity.

An investigation of the possible development of tolerance to the anticonvulsant activity revealed no tolerance in rats dosed orally with topiramate for 14 days at twice the ED<sub>50</sub> value. When mice were dosed orally for 5 days at four times the ED<sub>50</sub> value, a small but significant degree of tolerance did occur.

Topiramate was examined for effects on central nervous system (CNS) function, particularly reflex activity and motor co-ordination. A quantitative measure of CNS impairment was obtained by calculating the dose required to cause a loss of righting reflex (LRR) in either 3% (TD<sub>3</sub>) or 50% (TD<sub>50</sub>) of mice tested, or the dose that caused 50% (TD<sub>50</sub>) of mice or rats to be unable to remain ambulatory on a rotating rod or reel. A protective index (PI) was obtained by calculating the ratio of the TD<sub>50</sub> dose to the ED<sub>50</sub> dose in the MES test (or the TD<sub>3</sub> dose to the ED<sub>97</sub> dose). The calculated PI values for topiramate compared favourably to those of the reference anticonvulsants phenytoin, carbamazepine, valproate (divalproex), and phenobarbital, particularly in rats. An evaluation of acute effects in dogs indicated that impairment of CNS function occurred only at doses several times the ED<sub>50</sub> dose in the MES test in rats and mice.

Topiramate was evaluated for effects on general behaviour in mice, rats, and dogs at doses ranging from 10 to 1,000 mg/kg. Dose-related effects in mice and rats included a decrease in spontaneous motor activity, and a decrease in body tone and respiratory activity. In dogs, emesis occurred in one of three dogs at 100 mg/kg (p.o.), and at 500 mg/kg (p.o.) one of three dogs exhibited preconvulsant activity and one of three convulsed. Recovery was complete at six hours after dosing. When administered i.v. to rats at doses ranging from 1 mg/kg to 10 mg/kg, topiramate had no effect on EEG activity, cerebral pH, spinal reflexes, or neuromuscular conduction. In mice, topiramate at doses of 30 mg/kg (p.o.) or greater prolonged pentobarbital-induced sleep time threefold to eightfold in a dose-dependent manner. In rats pretreated with topiramate at 60 mg/kg or 200 mg/kg (p.o.) one hour prior to inducing sleep with ethanol, sleep time was prolonged 38% and 54%, respectively. When rats were pretreated with these doses of topiramate four hours prior to inducing sleep with ethanol, there was no prolongation of sleep time.

In cardiovascular studies, topiramate, when given i.v. to anesthetized dogs at doses up to 10 mg/kg, caused a small, dose-related increase in blood pressure, which was associated with a slight decrease in heart rate. There was no effect on electrocardiographic measures at these doses. Topiramate, when administered to spontaneously hypertensive rats at doses of 30 mg/kg i.p. and 100 mg/kg p.o. caused a biphasic response in mean arterial pressure; an initial transient increase was followed by a modest decrease in blood pressure that persisted for about 12 hours. Topiramate, at concentrations up to 10 μM, elicited no biologically significant effects on coronary flow, contractile force, or flow rate in the isolated guinea pig heart.

In GI studies, topiramate at concentrations up to 100  $\mu$ M had no effect on basal or pentagastrin-stimulated gastric acid secretion in the isolated mouse stomach assay. Topiramate weakly inhibited gastric acid secretion in rats and dogs.

Topiramate and acetazolamide were examined for effects on renal function using rats anesthetized with pentobarbital. Both compounds were infused i.v. at 9 or 90  $\mu$ M/kg/h. At each dose, both compounds produced changes in renal function, including an increase in urinary flow rate, solute clearance and urinary pH. Also, a decrease in urinary osmolality and decreases in arterial blood pH and plasma bicarbonate concentration were observed. The effects of both dosage levels of topiramate were similar to, but less than, those of acetazolamide. Renal vascular resistance, heart rate, and glomerular filtration rate did not differ from pretreatment control values.

### Pharmacokinetics

Studies performed in rats and dogs employing  $^{14}$ C-topiramate show that topiramate is rapidly and well absorbed after oral administration and that unchanged topiramate is the major component in plasma for several hours after dosing. The absolute bioavailability of topiramate is approximately 100% in male and female rats.

Topiramate is poorly bound to plasma proteins (9% to 17%) in the mouse, rat, rabbit, dog, and monkey, but there appears to be a low capacity erythrocyte binding site for the drug in all species studied. Studies in rats show that following oral administration of  $^{14}$ C-topiramate total radioactivity does not accumulate in any tissue. Topiramate did distribute across the blood-brain barrier, with brain tissue concentrations of total radioactivity being about 40% of plasma concentrations 6 hours after a single oral dose.

The metabolism of topiramate has been investigated in mice, rats, rabbits and dogs. The metabolic pathways, primarily hydroxylation or hydrolysis of the isopropylidene groups and subsequent conjugation, were qualitatively similar in all species studied.

The major route of elimination of unchanged topiramate and its metabolites in all species studied is via the kidney. All species excreted a significant proportion of the dose in urine as intact topiramate; however the proportion of metabolites excreted tended to be higher in species with shorter plasma half-lives.

## **TOXICOLOGY**

In acute and long-term studies conducted in mice, rats, dogs, and rabbits, exposure to topiramate was well tolerated.

### Acute Toxicity

<b>Table 2.4: Acute Toxicity Studies Performed with Topiramate</b>				
<b>Species/Strain</b>	<b>Route of Administration</b>	<b>No. Animals/ Group M/F [Age]</b>	<b>Dose Range (mg/kg)</b>	<b>Estimated LD<sub>50</sub> (mg/kg)</b>

Mouse CrI: COBS CD <sup>®</sup> -1 (ICR)BR	Oral gavage	2/2 or 5/5 [6-8 weeks]	1000-3375	M 2338 F 2915
Mouse CrI: COBS CD <sup>®</sup> -1 (ICR)BR	i.p.	5/5 or 2/2 [6-8 weeks]	500-1700	M 605 F 710
Rat CrI: COBS <sup>®</sup> (WI)BR	Oral gavage	5/5 or 2/2 [7-8 weeks]	1500-4220	M 3745 F 2436
Rat CrI: COBS <sup>®</sup> (WI)BR	i.p.	5/5 or 2/2 [7-8 weeks]	750-2550	M 1633 F 1227
Dog Beagle	p.o.	1/1 or 2/2 [approx. 1 yr.]	270-400	No deaths

## Chronic Toxicity

<b>Table 2.5: Multiple-Dose Toxicity Studies</b>					
<b>Species/Strain Sex</b>	<b>Route of Administration</b>	<b>No. Animals/ Group M/F</b>	<b>Dosage (mg/kg/day)</b>	<b>Duration</b>	<b>Results</b>
Rat CrI: CD <sup>®</sup> (SD) Male & Female	Oral gavage	15/15	10, 90, 750	3 months	Lower body weight and weight gain; CNS signs; diuresis with some hemoconcentration; higher kidney and liver weights with hepatocytic hypertrophy; and urothelial hyperplasia with some microcalculi (few females). Findings principally at 90 and 750 mg/kg per day.
Rat CrI: CD <sup>®</sup> (SD) Male & Female	Oral gavage	16/16 and 6/6 during recovery	10, 90, 750	3 months plus 4-week recovery period	Some slight effects (lower body weight gain and lower urine sodium) occurred at 10 mg/kg, the lowest dosage tested in this study; however, they are considered to be of no toxicological concern. Effects at ≥90 mg/kg were similar to those occurring in other 3-month and/or 12-month rat studies. Recovery occurred for all changes except for the increased water consumption and hyperplasia of the transitional epithelium of the bladder.
Rat CrI: COBS <sup>®</sup> (WI) Male & Female	Oral (diet)	25/25	10, 55, 300	12 months	Lower body weight and weight gain with lower food efficiency (300 mg/kg per day only); lower erythrocyte parameters and triglycerides; higher serum chloride and cholesterol; higher kidney and liver weights with hepatocytic hypertrophy; urothelial hyperplasia with urinary calculi; and gastric epithelial hyperplasia. Only body weight, chloride and gastric changes seen at 10 mg/kg per day.
Rat CrI: COBS <sup>®</sup> (WI) Male & Female	Oral (diet)	26/23 treated 20/21 untreated	Male: > 300 Female: ≥ 450	11 months plus 4-, 9-, and 20-week recovery periods	Lower weight gain; higher gastrin levels and gastric epithelial hyperplasia. No effect on gastric enterochromaffin-like cells (often associated with tumor formation in the presence of high gastrin levels). During the recovery phase, all changes were reversible.

<b>Table 2.5 (cont'd): Multiple-Dose Toxicity Studies Multiple-Dose Toxicity Studies</b>					
<b>Species/Strain Sex</b>	<b>Route of Administration</b>	<b>No. Animals/ Group M/F</b>	<b>Dosage (mg/kg/day)</b>	<b>Duration</b>	<b>Results</b>
Dog/Beagle Male & Female	p.o.	4/4	10, 40, 150	3 months	Lower weight gain, food consumption, and food efficiency; hemodilution; lower transaminases and urine specific gravity; higher urine pH, serum alkaline phosphatase and chloride; and higher liver weights. No significant morphological changes. Observations essentially at 40 and 150 mg/kg per day only.
Dog/Beagle Male & Female	p.o.	4/4	10, 30, 100	12 months	Sporadic emesis at all dosage levels. Lower body weight gain; hemodilution; higher urine pH, serum alkaline phosphatase and chloride; and higher liver weights. No significant morphologic changes. Only findings seen below 100 mg/kg per day were emesis, and higher alkaline phosphatase and chloride.

### **Reproductive Toxicity**

Topiramate has demonstrated selective developmental toxicity, including teratogenicity, in multiple animal species at clinically relevant doses. When oral doses of 20, 100, or 500 mg/kg were administered to pregnant mice during the period of organogenesis, the incidence of fetal malformations (primarily craniofacial defects) was increased at all doses. The low dose is approximately 0.2 times the recommended human dose (RHD) 400 mg/day on a mg/m<sup>2</sup> basis. Fetal body weights and skeletal ossification were reduced at 500 mg/kg in conjunction with decreased maternal body weight gain.

In rat studies (oral doses of 20, 100, and 500 mg/kg or 0.2, 2.5, 30, and 400 mg/kg), the frequency of limb malformations (ectrodactyly, micromelia, and amelia) was increased among the offspring of dams treated with 400 mg/kg (10 times the RHD on a mg/m<sup>2</sup> basis) or greater during the organogenesis period of pregnancy. Embryotoxicity (reduced fetal body weights, increased incidence of structural variations) was observed at doses as low as 20 mg/kg (0.5 times the RHD on a mg/m<sup>2</sup> basis). Clinical signs of maternal toxicity were seen at 400 mg/kg and above, and maternal body weight gain was reduced during treatment with 100 mg/kg or greater.

In rabbit studies (20, 60, and 180 mg/kg or 10, 35, and 120 mg/kg orally during organogenesis), embryo/fetal mortality was increased at 35 mg/kg (2 times the RHD on a mg/m<sup>2</sup> basis) or greater, and teratogenic effects (primarily rib and vertebral malformations) were observed at 120 mg/kg (6 times the RHD on a mg/m<sup>2</sup> basis). Evidence of maternal toxicity (decreased body weight gain, clinical signs, and/or mortality) was seen at 35 mg/kg and above.

When female rats were treated during the latter part of gestation and throughout lactation (0.2, 4, 20, and 100 mg/kg or 2, 20, and 200 mg/kg), offspring exhibited decreased viability and delayed physical development at 200 mg/kg (5 times the RHD on a mg/m<sup>2</sup> basis) and reductions in

preand/ or postweaning body weight gain at 2 mg/kg (0.05 times the RHD on a mg/m<sup>2</sup> basis) and above. Maternal toxicity (decreased body weight gain, clinical signs) was evident at 100 mg/kg or greater.

In a rat embryo/fetal development study with a postnatal component (0.2, 2.5, 30, or 400 mg/kg during organogenesis; noted above), pups exhibited delayed physical development at 400 mg/kg (10 times the RHD on a mg/m<sup>2</sup> basis) and persistent reductions in body weight gain at 30 mg/kg (1 times the RHD on a mg/m<sup>2</sup> basis) and higher.

### **Carcinogenicity**

Tumours of smooth muscle origin in the urinary bladder were seen only in mice (oral dosages up to 300 mg/kg for 21 months) and appear to be unique to the species. Since no human counterpart exists, they were not considered clinically relevant. No such findings occurred in the rat carcinogenicity study (oral dosages up to 120 mg/kg/day for 24 months).

### **Mutagenicity**

In a battery of *in vitro* and *in vivo* mutagenicity assays, topiramate did not show genotoxic potential.

## REFERENCES

1. Rangel RJ, Penry JK, Wilder BJ, et al. Topiramate: A new antiepileptic drug for complex partial seizures - first use in epileptic patients. *Neurology* 1988; 38:234.
2. Ostergaard L, Dam M, Mikkelsen M. Topiramate as add-on therapy in refractory partial epilepsy. *Epilepsia* 1992; 33(Suppl 3):105.
3. Anon. Topiramate, MCN-4853, RWJ-17021-000, EN=105605. *Drugs of the Future*. 1990; 15:432.
4. Aranguiz C, McJilton J, Vega M, Ramsay RE. Safety and effectiveness of three oral doses of topiramate in the treatment of patients with refractory partial epilepsy. *Epilepsia* 1991;32:11.
5. Fincham R, Schottelius D, Faught E, Kuzniecky R, Thompson G. Efficacy and safety of topiramate (TPM) as adjunctive therapy in adults with refractory partial epilepsy (RPE). *Neurology* 1992; 42:311.
6. Rosenfeld WE, Holmes GB, Hunt TL, Schaefer P. Topiramate-effective dosaging enhances potential for success. *Epilepsia* 1992; 33:118.
7. Ben-Menachem E. Double-blind placebo-controlled trial of topiramate as add-on therapy for the treatment of complex partial seizures. *Epilepsia* 1992; 33:105.
8. Ben-Menachem E, Mikkelsen M, Dam M, Engelskjøn T, Henriksen O, Johanessen SI, Schmidt D, Ried S, Proest G. Topiramate add-on treatment in patients with intractable partial epilepsy: a multicenter study. *Epilepsia* 1993; 34:109.
9. Faught E, Wilder BJ, Ramsay RE, et al. Topiramate placebo-controlled dose-ranging trial in refractory partial epilepsy using 200-, 400-, and 600-mg daily dosages. *Neurology* 1996; 46: 1684-1696.
10. Ben-Menachem E. Clinical Efficacy of topiramate as add-on Therapy in Refractory Partial Epilepsy. *Epilepsia* 1997; 38(suppl 1):228-230.
11. Sharief M, Viteri C, Ben-Menachem E. et al. Double-blind placebo-controlled study of topiramate in patients with refractory partial epilepsy. *Epilepsy Res*. 1996; 25(3):217-24.
12. Dose DR, Scott VV, Marchgull BL, et al. Multiple-dose pharmacokinetics of topiramate in healthy male subjects. *Epilepsia* 1988; 29(5):662.
13. Easterling DE, Zakszewski T, Moyer MD, et al. Plasma pharmacokinetics of topiramate, a new anticonvulsant in humans. *Epilepsia* 1988; 29(5):662.



14. Wilder BJ, Penry JK, Rangel RJ, et al. Topiramate: Efficacy, toxicity, and dose/plasma concentrations. *Epilepsia* 1988; 29(5):698.
15. Wilensky AJ, Ojemann LM, Chmelir T, et al. Topiramate pharmacokinetics in epileptic patients receiving carbamazepine. *Epilepsia* 1989; 30(5):645-646.
16. Floren KL, Graves NM, Leppik IE, et al. Pharmacokinetics of topiramate in patients with partial epilepsy receiving phenytoin or valproate. *Epilepsia* 1989; 30(5):646.
17. Marchyanoff BE, Marchgull BL. Topiramate. *Drugs of the Future* 1989; 14(4):342-344.
18. Nakamura J, Tamura S, Ishii A. Inhibition of tonic and absence-like seizures in spontaneously epileptic rats by a novel antiepileptic drug topiramate (KW6485). *Jap J Pharmacol* 1992; 58:321.
19. Tanabe K, Wang Y, Kimishima K. Anticonvulsant activities and properties of topiramate. *J Yonago Medical Assoc* 1991; 42(6):330-346.
20. Edmonds H, Jiang D, Zhang P, Vaught JL. Topiramate in a rat model of post-traumatic epilepsy. *Epilepsia* 1991; 32:15.
21. Kanda T, Nakamura J, Kurokawa M, et al. Inhibition of excessive releases of excitatory amino acids in hippocampus of spontaneously epileptic rat (SER) by topiramate (KW-6485). *Jap J Pharmacol* 1992; 58:92.
22. Coulter DA, Sombati S, DeLorenzo RJ. Selective effects of topiramate on sustained repetitive firing and spontaneous bursting in cultured hippocampal neurons. *Epilepsia* 1993; 34(Suppl 2):123.
23. Brown SD, Wolf HH, Swinyard EA, Twyman RE, White HS. The novel anticonvulsant topiramate enhances GABA-mediated chloride flux. *Epilepsia* 1993; 34(Suppl 2):122-3.
24. Vaught JL, Marchyanoff BE, Shank R. The pharmacological profile of topiramate: A structurally novel, clinically effective anticonvulsant. *Epilepsia* 1991; 32:19.
25. Shank RP, Vaught JL, Raffa RB, Marchyanoff BE. Investigation of the mechanism of topiramate's anticonvulsant activity. *Epilepsia* 1991; 32:7-8.
26. Edmonds HL, Jiang D, Zhang YP, Shank RP. Topiramate as a neuroprotectant and anticonvulsant in post-ischemic injury. *Epilepsia*, December 6-9, 1992; 33:118-119.
27. No Authors listed in citation. Topiramate. *Drugs of the Future* 1993; 18:4:398.

28. Privitera M, Fincham R, Penny J, et al. Topiramate placebo-controlled dose-ranging trial in refractory partial epilepsy using 600-, 800- and 1,000- mg daily dosages. *Neurology* 1996;46: 1678-1683.
29. Ben-Menachem E, Henriksen O, Dam M, et al. Double-blind, placebo-controlled trial of topiramate as add-on therapy in patients with refractory partial seizures. *Epilepsia* 1996;37:539-543.
30. Tassinari CA, Michelucci R, Chauvel P, et al. Double-blind, placebo-controlled trial of topiramate (600 mg daily) for the treatment of refractory partial epilepsy. *Epilepsia* 1996;37:763-768.
31. Rosenfeld B, Abou-Khalil B, Reife R, et al. Placebo-controlled trial of topiramate as adjunctive therapy to carbamazepine or phenytoin for partial-onset epilepsy. *Epilepsia* 1996; 37 (Suppl 5): 153.
32. Special Advisory Committee on Reproductive Physiology to the Drugs Directorate, Health Protection Branch, Health Canada. *Oral Contraceptives* 1994.
33. Dulac O, N'Guyen T. The Lennox-Gastaut syndrome. *Epilepsia* 1993; 34 (Suppl 7):S7-S17.
34. Reife RA. Topiramate: a novel antiepileptic agent. In: Shorvon S, Dreifuss F, Fish D, and Thomas D (eds), *The Treatment of Epilepsy*. Cambridge, MA: Blackwell Science Ltd.; 1996:1-11.
35. Dodson WE, Bourgeois BFD. Pharmacology and therapeutic aspects of antiepileptic drugs in paediatrics. *J Child Neurol* 1994; 9 (Suppl):2S1-2S7.
36. Biton V, Reife R, Kamin M, Bobrosky K. Topiramate (TOPAMAX<sup>®</sup>) as add-on therapy for patients with primary generalized epilepsy. *Neurology* 1996; 46:A176.
37. Glauser TA. Topiramate. *Seminars in Pediatric Neurology* 1997; 4(1):34-42.
38. Elterman RD, Glauser RA, Wyllie E. et al. A double-blind, randomized trial of topiramate as adjunctive therapy for partial-onset seizures in children (Study YP). *Neurology* April 1999; 52(7): 1338-1344.
39. Sachdeo RC, Glauser TA, Ritter F et al. A Double-Blind, randomized trial of topiramate in Lennox-Gastaut Syndrome (Study YL). *Neurology* June 1999; 52(9): 1882-1887.
40. Biton V, Montouris GD, Ritter R et al. A randomized, placebo-controlled study of topiramate in primary generalized tonic-clonic (Study YTC). *Neurology* April 1999; 52(7):1330-1337.

41. Montouris GD et al. Nonfocal generalized tonic-clonic seizures: response during long-term topiramate treatment. Topiramate YTC/YTCE Study Group. *Epilepsia*. 2000;41 Suppl 1:S77-81.
42. JWAS Sander. Practical aspects of the use of topiramate in patients with epilepsy. *Epilepsia* 1997; 38 (Suppl 1): S56-S58.
43. Banta J, Hoffman K, Budenz D, Ceballos E, Greenfield D. Presumed topiramate-induced bilateral acute angle-closure glaucoma. *American Journal of Ophthalmology* 2001;132(1):112-114.
44. Sen H, O'Halloran H, Lee W. Topiramate-induced acute myopia and retinal striae. *Arch. Ophthalmol.* 2001; 119(5): 775-777.
45. Aldenkamp AP, Baker G, Mulder OG, et al. A multicenter, randomized clinical study to evaluate the effect on cognitive function of topiramate compared with valproate as add-on therapy to carbamazepine in patients with partial-onset seizures. *Epilepsia* 2000; 41(9):1167-1178.
46. Baker GA, Currie NG, Light MJ, Schneiderman JH. The effects of adjunctive topiramate therapy on seizure severity and health-related quality of life in patients with refractory epilepsy - a Canadian study. *Seizure* 2002; 11(1): 6-15.
47. Stephen LJ, Sills GJ, Brodie MJ. Topiramate in refractory epilepsy: a prospective observational study. *Epilepsia* 2000; 41(8): 977-980.
48. Rosenfeld, WE, Slater, J. Characterization of topiramate-associated weight changes in adults with epilepsy. *AES Proceedings. Epilepsia* 2001; 42(7):252.
49. Naritoku DK, Hulihan JF, Kamin M et al. Reduction of AED Co-therapy improves tolerability of add-on therapy with topiramate: A novel randomized study. *AES Proceedings. Epilepsia* 2001; 42(7):258.
50. Ben-Menachem E, Smith U, Hellstrom K et al. Predictors of weight loss in patients with epilepsy treated with topiramate. *AES Proceedings. Epilepsia* 2002; 43(7):220-221.
51. International Headache Society. The international classification of headache disorders. *Cephalgia* 2004; 24(Supplement 1):1-160.
52. Silberstein SD, Neto W, Schmitt J, Jacobs DJ. Topiramate in migraine prevention: Results of a large controlled trial. *Arch Neurol* 2004; 61:490-495.
53. Brandes JL, Saper JR, Diamond M et al. Topiramate for migraine prevention: A randomized controlled trial. *JAMA* 2004; 291:965-973.

54. Diener H-C, Tfelt-Hansen P, Dahlöf C et al. Topiramate in migraine prophylaxis: Results from a placebo-controlled trial with propranolol as an active control. *J Neurol* 2004; 251 :943-950.
55. Product Monograph, Topamax<sup>®</sup> (Topiramate), Janssen-Ortho Inc., Revised February 21, 2012.
56. Comparative bioavailability study, Teva Canada Limited, on file.

## PART III: CONSUMER INFORMATION

**Pratio-TOPIRAMATE**  
topiramate

This leaflet is part III of a three-part “Product Monograph” published when ratio-TOPIRAMATE was approved for sale in Canada and is designed specifically for Consumers. This leaflet is a summary and will not tell you everything about ratio-TOPIRAMATE. Contact your doctor or pharmacist if you have any questions about the drug.

**ABOUT THIS MEDICATION****What the medication is used for:**

- **ratio-TOPIRAMATE** has been prescribed to you/your child to control epilepsy.
- **ratio-TOPIRAMATE** may also be prescribed to you to prevent your migraine headaches if you are an adult patient (over 18 years of age) with 4 or more attacks per month and are not responding to acute treatment.

**What it does:**

**ratio-TOPIRAMATE** affects chemicals in the brain that are involved in sending signals to the nerves. **ratio-TOPIRAMATE** belongs to a group of medicines used to treat epilepsy.

**When it should not be used:**

- You/your child should not use **ratio-TOPIRAMATE** if you are allergic to any of the ingredients in the product. Contact your doctor immediately if you experience an allergic reaction (e.g. skin rash, hives) or any severe or unusual side effects.
- You should not use **ratio-TOPIRAMATE** to prevent your migraine headaches if you are pregnant or a woman of childbearing potential and are not using an effective method of birth control.
- For the prevention of other types of headaches that are different from migraine attacks.
- For the acute treatment of migraine headache.

**What the medicinal ingredient is:**

topiramate

**What the nonmedicinal ingredients are:**

The tablets contain the following inactive ingredients: pregelatinized starch, lactose monohydrate, colloidal silicon dioxide, sodium starch glycolate, microcrystalline cellulose and magnesium stearate, hydroxypropyl methylcellulose, titanium dioxide, polydextrose, triacetin and polyethylene glycol. The Colourants that are present in the tablets are as follows: 100 mg: yellow iron oxide and 200 mg: red iron oxide.

**What dosage forms it comes in:**

Tablets: 25 mg, 100 mg, 200 mg

**WARNINGS AND PRECAUTIONS**

BEFORE you use **ratio-TOPIRAMATE** talk to your doctor or pharmacist if:

- you drive a vehicle, use machines, perform hazardous tasks during your work or do anything else that could be dangerous if you are not alert.
- you/your child have or have had kidney stones or kidney disease. Your doctor may want you to increase the amount of fluids you/your child drink(s) while taking this medicine.
- you/your child have or have had liver disease.
- you/your child have or have had depression, mood problems, or suicidal thoughts or behaviour.
- you/your child have a history of metabolic acidosis (too much acid in the blood).
- you/your child have weak, brittle, or soft bones (osteomalacia, osteoporosis, osteopenia, or decreased bone density).
- you/your child have eye problems, especially glaucoma.
- you/your child have diarrhea.
- you/your child are having surgery.
- you/your child have or have had any medical problems and allergies.
- **you are breast-feeding (nursing).**
- you/your child are/is taking medicines that slow down the nervous system (CNS depressants).
- **you/your child are taking oral contraceptives and ratio-TOPIRAMATE Tablets, and tell your doctor about any changes in your bleeding patterns (breakthrough bleeding/spotting).**
- you are taking a ketogenic diet (a diet high in fat and low in protein and sugar).
- you consume alcohol regularly.
- You/your child have a growth problem.

**EPILEPSY ONLY**

- **If you take ratio-TOPIRAMATE during pregnancy, your baby has a higher risk for birth defects called cleft lip and cleft palate. These defects can begin early in pregnancy, even before you know you are pregnant.**
- **Cleft lip and cleft palate may happen even in children born to women who are not taking any medicines and do not have other risk factors.**
- **There may be other medicines to treat your condition that have a lower chance of birth defects.**
- **All women of childbearing age who are being treated for epilepsy should talk to their healthcare providers about using other possible treatments instead of ratio-TOPIRAMATE. If the decision is made to use ratio-TOPIRAMATE, you should use effective birth control (contraception) unless you are planning to become pregnant. You should talk to your doctor about the best kind of birth control to use while you are taking ratio-TOPIRAMATE.**
- **Metabolic acidosis may have harmful effects on your baby. Talk to your healthcare provider if ratio-TOPIRAMATE has caused metabolic acidosis during your pregnancy.**

- **Tell your doctor right away if you become pregnant while taking ratio-TOPIRAMATE. You and your doctor should decide if you will continue to take ratio-TOPIRAMATE while you are pregnant.**

**Pregnancy Registry: If you become pregnant while taking ratio-TOPIRAMATE talk to your doctor about registering with the North American Antiepileptic Drug Pregnancy Registry. You can enroll in this registry by calling 1-888-233-2334. The purpose of this registry is to collect information about the safety of antiepileptic medicine during pregnancy. Information on the registry can also be found at the website <http://www.massgeneral.org/aed/>.**

#### **MIGRAINE PROPHYLAXIS**

**ratio-TOPIRAMATE is not to be used to prevent migraine headaches in pregnant women or women of childbearing potential who are not using an effective method of birth control.**

#### Other Precautions:

**ratio-TOPIRAMATE** may cause some people to be less alert than normal. Make sure you know how you/your child are/is affected by this medication before you drive, use machines, or do anything else that could be dangerous if you are not alert.

**ratio-TOPIRAMATE** may reduce the efficacy of oral contraceptives even in the absence of breakthrough bleeding. Therefore, oral contraceptives containing not less than 30 µg of estrogen should be used.

A very small number of people may have thoughts of suicide.

Rarely, blood tests have shown a slight increase in acidity. In many cases, there are no symptoms from this increased acidity but some people may experience symptoms such as rapid breathing, persistent lack of energy and loss of appetite. Some people may experience more serious symptoms such as heart problems, confused thinking or reduced consciousness.

**Do not stop ratio-TOPIRAMATE without first talking to a healthcare provider.** Stopping ratio-TOPIRAMATE suddenly can cause serious problems.

#### **INTERACTIONS WITH THIS MEDICATION**

Tell your doctor about all medications (prescription and non-prescription) and dietary supplements you/your child are/is using. It is especially important that your doctor know if you/your child are/is taking digoxin, oral contraceptives, glyburide, lithium, risperidone, diltiazem, or any other antiepileptic drugs, such as phenytoin, valproic acid or carbamazepine.

#### **PROPER USE OF THIS MEDICATION**

**ratio-TOPIRAMATE** is usually taken twice a day; however, your

doctor may tell you to take it once a day or at a higher or lower dose.

Never stop taking, increase or decrease the amount of **ratio-TOPIRAMATE** you are taking unless your doctor tells you to.

Swallow the Tablets and take with plenty of water. You/your child can take the Tablets with or without food. Do not break or crush your tablets.

Always check that you have enough Tablets and do not run out. Do not suddenly stop taking this medicine without first checking with your doctor.

#### **EPILEPSY**

It is important that you take **ratio-TOPIRAMATE** exactly as your doctor has instructed. Your doctor will start with a low dose and slowly increase the dose to the lowest amount needed to control your/your child's epilepsy.

#### Usual dose:

**ratio-TOPIRAMATE taken alone:** The usual maintenance dose in adults and children (6 years of age and older) is between 100 mg/day and 400 mg/day. **ratio-TOPIRAMATE** is usually taken twice a day.

**ratio-TOPIRAMATE taken in combination with other antiepileptic drugs:** The usual adult maintenance dose is 200 mg to 400 mg per day.

In children, dosing is based on weight and the dose is approximately 5 to 9 mg/kg/day.

**ratio-TOPIRAMATE** is not indicated for use in patients under 2 years of age.

#### **MIGRAINE PROPHYLAXIS**

It is important that you follow your doctor's instructions carefully to help reduce the chances of getting a migraine headache. Your doctor will start treatment with a dose of 25 mg to be taken at night. Your doctor will then increase your dose to the lowest amount needed to prevent migraine headaches.

#### Usual dose:

The usual adult dose is 100 mg per day. **ratio-TOPIRAMATE** is taken twice a day (50 mg in the morning and 50 mg at night). Your doctor may tell you to use a lower or higher dose.

**ratio-TOPIRAMATE** is not indicated for the prevention of migraine attacks in patients under 18 years of age.

**Remember: This medicine has been prescribed for you/your child. Do not give it to anybody else.**

#### Overdose:

**In case of a drug overdose, immediately go to the nearest emergency room even if you do not feel sick. Make sure you take your medicine bottle with you to show the doctor.**

**Missed Dose:**

If you/your child miss/misses a dose, take it as soon as you remember. But, if it is almost time for the next dose, do not take the missed dose. Instead, take the next scheduled dose. Do not try to make up for the missed dose by taking a double dose next time.

**SIDE EFFECTS AND WHAT TO DO ABOUT THEM**

- Any medicine may have unwanted effects. Tell your doctor or pharmacist about any unusual sign or symptom whether listed or not.
- Contact your doctor immediately or go to the Emergency Room if you/your child experience/experiences sudden worsening of vision, blurred vision or painful/red eye(s).
- Decreased sweating may occur while taking **ratio-TOPIRAMATE**. Make sure you/your child increase/increases and maintain/maintains fluid intake prior to and during activities such as exercise and exposure to warm temperatures. Contact your doctor if you/your child are/is not sweating as usual or show/shows signs of increased body temperature.
- Side effects reported most often in adults are: *co-ordination problems, difficulty concentrating, slow thinking, confusion and forgetfulness, dizziness, tiredness, tingling, headache, upper respiratory tract infection (e.g. colds, bronchitis) and drowsiness*. Less frequently reported side effects are: *agitation, decrease in appetite, speech disorders (e.g. hesitancy or word-finding difficulty), depression, emotional lability, vision disorders (e.g. double vision), mood swings, nausea, taste changes, weight loss and kidney stones (may include symptoms such as blood in the urine, or low back pain or pain in the genital area)*.
- In children, the following side effects were associated with the use of topiramate: *difficulty concentrating, forgetfulness, tiredness, drowsiness, nervousness, decrease in appetite, weight loss, upper respiratory tract infection (e.g. colds, bronchitis), headache, fever, tingling and aggressive behaviour*.

**SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM**

Symptom / effect		Talk with your doctor or pharmacist right away		Seek Urgent medical attention
		Only if severe	In all cases	
<b>Uncommon</b>	Sudden worsening of vision, blurred vision with painful/red eye(s)			✓

**SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM**

Symptom / effect	Talk with your doctor or pharmacist right away	Seek Urgent medical attention
<i>Allergic Reaction (red skin, hives, itching, swelling of the lips, face, tongue, throat, trouble breathing, wheezing, shortness of breath, skin rashes, blisters of the skin, sore mouth or eyes)</i>		✓
Decreased sweating	✓	
Thoughts of suicide or hurting yourself	✓	
<b>Rare</b>		✓
<i>Kidney stones (blood in the urine or pain in the lower back or genital area)</i>		✓
<b>Very rare</b>		✓
<i>Metabolic Acidosis (unexplained tiredness, loss of appetite, irregular heartbeat, and impaired consciousness)</i>		✓
Confusion, problems with concentration, attention, memory, and/or speech	✓	
<i>High blood ammonia (decreased alertness, tiredness, or fatigue, vomiting)</i>		✓

Uncommon side effects – between 1 and 10 reports in every 1000 patients exposed  
 Rare side effects – from 1 to less than 10 reports in every 10,000 patients exposed  
 Very rare side effects – less than 1 report in every 10,000 patients exposed

*This is not a complete list of side effects. If you experience any unexpected effects while taking this drug, contact your doctor or pharmacist.*

**HOW TO STORE IT**

- Do not use this product after the expiry date written on the package.
- Store between 15 - 30°C in tightly closed light resistance containers. Protect from humidity.
- Keep this and all medicines in a safe place away from children.

#### REPORTING SUSPECTED SIDE EFFECTS

You can report any suspected adverse reactions associated with the use of health products to the Canada Vigilance Program by one of the following 3 ways:

- Report online at [www.healthcanada.gc.ca/medeffect](http://www.healthcanada.gc.ca/medeffect)
- Call toll-free at 1-866-234-2345
- Complete a Canada Vigilance Reporting Form and:
  - Fax toll-free to 1-866-678-6789, or
  - Mail to: Canada Vigilance Program  
Health Canada  
Postal Locator 0701E  
Ottawa, ON K1A 0K9

Postage paid labels, Canada Vigilance Reporting Form and the adverse reaction reporting guidelines are available on the MedEffect™ Canada Web site at [www.healthcanada.gc.ca/medeffect](http://www.healthcanada.gc.ca/medeffect).

***NOTE: Should you require information related to the management of side effects, contact your health professional. The Canada Vigilance Program does not provide medical advice.***

#### MORE INFORMATION

This document plus the full product monograph, prepared for health professionals can be found by contacting the sponsor, Teva Canada Limited at:  
1-800-268-4127 ext. 5005  
or [druginfo@tevacanda.com](mailto:druginfo@tevacanda.com)

This leaflet was prepared by:  
Teva Canada Limited  
30 Novopharm Court  
Toronto, Ontario  
Canada, M1B 2K9

Last revised: July 16, 2012