

## **PRODUCT MONOGRAPH**

### **Pr Vinblastine Sulfate Injection**

1 mg/mL  
(10 mg/ 10 mL)  
Sterile

Antineoplastic Agent

Sandoz Canada  
145, Jules-Léger  
Boucherville, QC  
J4B 7K8

Date of Preparation:  
May 15, 2013

Submission Control No.: 140855

## PRODUCT MONOGRAPH

### Pr **Vinblastine Sulfate Injection**

1 mg/mL  
(10 mg/ 10 mL)  
Sterile

### THERAPEUTIC CLASSIFICATION

Antineoplastic Agent

### CAUTION

**VINBLASTINE SULFATE IS A POTENT DRUG AND SHOULD BE USED ONLY BY PHYSICIANS EXPERIENCED WITH CANCER CHEMOTHERAPEUTIC DRUGS. BLOOD COUNTS SHOULD BE TAKEN ONCE OR TWICE WEEKLY. DISCONTINUE OR REDUCE THE DOSAGE UPON EVIDENCE OF ABNORMAL DEPRESSION OF THE BONE MARROW. VINBLASTINE SULFATE SHOULD NOT BE GIVEN INTRATHECALLY.**

### ACTIONS AND CLINICAL PHARMACOLOGY

Tissue culture studies indicate that vinblastine is a selective mitotic inhibitor of certain malignant cells and as such appears to be different from other recognized antineoplastic drugs.

There is evidence that vinblastine interferes with cell metabolism and the entrance of glutamic acid into the citric acid cycle and to urea. There is also evidence that the anti-tumor effect of vinblastine may possibly be due to its effect on cell energy mechanisms and a decreased adenosine diphosphate production resulting from retarded nucleotide production. It has been suggested that the therapeutic ratio may depend on the greater energy needs of cancer cells along with their decreased efficiency in generating energy because of their reliance on the glycolytic pathway. These energy related actions for vinblastine have been brought forward because although it has been demonstrated that vinblastine has a stathmokinetic effect and produces various atypical mitotic figures in treated cells it has been observed clinically and experimentally that these cytological changes may occur in the absence of oncolytic effects.

It has also been demonstrated *in vitro* that vinblastine can prevent the invasion of normal tissue by malignant cells, thus preventing the spread of malignancy.

The relationship of vinblastine to amino acid metabolism has been shown by a reversal of the anti-tumor effect of vinblastine by glutamic acid or tryptophan. As well, aspartic and glutamic acids had protected mice from lethal doses of vinblastine, although aspartic acid was relatively ineffective in reversing the anti-tumor effect of vinblastine.

Vinblastine has been shown in clinical practice to provide palliative treatment for a variety of malignant neoplastic diseases and in susceptible tumors produced a temporary reduction in the size of some tumors. Its use has relieved pain and other symptoms associated with neoplasm and permitted some patients to regain appetite and weight. Remission periods have varied from patient to patient.

Leukopenia is an expected effect of vinblastine and the leukocyte count is an important guide to vinblastine therapy. In general, the larger the dose, the longer lasting and more profound the leukopenia. However, the white cell producing mechanism is not permanently depressed by vinblastine, as it returns to normal on discontinuing treatment. Normally, when the white cells have essentially disappeared from the peripheral blood, the white count would have returned to normal. The nadir in white blood cell count usually occurs 5 - 10 days after the last treatment with vinblastine. Recovery is usually quite rapid and complete within another 7 - 14 days. If small doses are employed for maintenance therapy, leukopenia may not be clinically significant.

The thrombocyte count is not usually significantly reduced by vinblastine therapy, however, in patients whose bone marrow has been recently impaired by radiation therapy or other anti-neoplastic agents, thrombocytopenia may develop (less than 200,000 platelets per mm<sup>3</sup>). Thrombocyte values below 200,000 per mm<sup>3</sup> are rarely encountered when other chemotherapeutic agents or radiation have not previously been used, even though significant leukopenia may be present. If thrombocytopenia does occur, it usually reverses within a few days after termination of treatment.

Vinblastine generally has an insignificant effect on red cell count and hemoglobin. However, patients with a malignant disease may have anemia even in the absence of antineoplastic therapy.

## **INDICATIONS AND CLINICAL USES**

Vinblastine Sulfate is indicated in the palliative treatment of the following neoplastic diseases:

1. Frequently Responsive Malignancies  
Generalized Hodgkin's disease (Stages III and IV, Ann Arbor modification of Rye)  
Lymphocytic lymphoma (modular and diffuse, poorly and well differentiated)  
Histocytic lymphoma  
Mycosis fungoides (advanced stages)  
Advanced carcinoma of the testis  
Kaposi's Sarcoma  
Letterer-Siwe disease (histiocytosis-X)
2. Less Frequently Responsive Malignancies  
Choriocarcinoma resistant to other neoplastic drugs  
Cancer of the breast (unresponsive to endocrine surgery and hormonal therapy)

The simultaneous use of several cancer chemotherapy drugs is common practice. Generally, drugs with different dose limiting clinical toxicities and different mechanisms of action are selected in order to obtain an increase in therapeutic response without added toxicity. Rarely is it

possible to obtain equally as good a response with single antineoplastic treatment. Therefore, vinblastine is often part of polychemotherapy because at the recommended doses it does not cause significant suppression of the bone marrow or neuropathy. This approach to multipletreatment has been used in the chemotherapy of Hodgkin's disease.

### Hodgkin's disease

Vinblastine has been found to be one of the most effective single antineoplastic agents for the treatment of Hodgkin's disease. Successful treatment of advanced Hodgkin's disease has been accomplished by the use of various multiple-drug regimens that have included vinblastine. Patients who have relapsed following treatment with the MOPP-regimen (mechlorethamine hydrochloride [nitrogen mustard], vincristine sulfate, prednisone and procarbazine) have often responded to combination drug therapy that included vinblastine. An alternative therapy that has been used in previously untreated patients with advanced Hodgkin's disease employs cyclophosphamide in place of nitrogen mustard and vinblastine instead of vincristine.

Advanced testicular germinal-cell cancers such as embryonal carcinoma, teratocarcinoma, and choriocarcinoma have been shown to be sensitive to vinblastine alone but a more satisfactory clinical response may be obtained by the concomitant administration of vinblastine with other anti-tumor drugs. The efficacy of bleomycin has been found to be enhanced if vinblastine is given 6 - 8 hours prior to bleomycin. This procedure appears to result in more cells being arrested during metaphase, the stage of cell division in which bleomycin is active.

## **CONTRAINDICATIONS**

The use of vinblastine is contraindicated in patients with leukopenia. Vinblastine should not be administered to patients with bacterial infections. Such infections must be brought under control by the use of antibiotic or antiseptic therapy prior to the initiation of vinblastine treatment.

### Pregnancy

Although no abnormalities of the human fetus have been associated with the use of vinblastine, information on its use during pregnancy is limited. Animal studies suggest that vinblastine may be teratogenic. Therefore, the use of vinblastine during pregnancy is contraindicated unless the expected benefits clearly outweigh the risk of side effects.

## **WARNINGS**

Aspermia has been reported in man. Animal studies have demonstrated degenerative changes in germ cells and arrest of cell division in metaphase.

## **PRECAUTIONS**

If leukopenia with less than 2,000 white blood cells per mm<sup>3</sup> develops following administration of vinblastine, the patient should be monitored carefully for evidence of infection until the white blood cell count returns to normal.

If cachexia or skin ulcers are present, a more profound leukopenia response to the drug may occur. Therefore the use of vinblastine should be avoided in elderly persons with either of these conditions.

In patients with malignant-cell infiltration of the bone marrow, the leukocyte and platelet counts have occasionally fallen precipitously after moderate doses of vinblastine and the administration of additional doses of vinblastine in such patients is not recommended.

The use of daily low doses of vinblastine for prolonged periods is not recommended, even though the total weekly dosage may be similar to the recommended treatment regimen. Little or no added therapeutic benefit has been demonstrated with the use of such low dose regimens. Strict adherence to the recommended dosage schedule is very important. When vinblastine was given in 7 daily injections at a total dose equal to several times the recommended weekly dosage for prolonged periods, convulsions, severe and permanent central nervous system damage and even death occurred.

Avoid contamination of the eye with vinblastine solutions. If accidental contamination does occur, severe irritation may result and if the drug was given under pressure, corneal ulceration may result. The eye should be washed immediately with copious quantities of water.

## **ADVERSE REACTIONS**

### Leukopenia

Bone-marrow depression, especially leukopenia, is the most common adverse effect with vinblastine and tends to be dose-limiting. Before administering the drug, patients should be advised of the possibility of adverse reactions. Maximum depression occurs 4 - 10 days after administration, with recovery in one to three weeks.

Except for epilation and leukopenia the adverse reactions seen with vinblastine usually do not persist for more than 24 hours.

### Gastrointestinal

Nausea, vomiting, constipation, vesiculation of the mouth, ileus, diarrhea, anorexia, abdominal pain, rectal bleeding, pharyngitis, hemorrhagic enterocolitis and bleeding from a dormant peptic ulcer may occur.

### Neurologic

Neurologic effects can involve the autonomic nervous system and include malaise, headache, depression, psychoses, paresthesia, neuromyopathy, loss of deep tendon reflexes, peripheral neuritis, constipation, numbness and convulsions.

### Miscellaneous

Epilation, malaise, weakness, dizziness, pain at the site of the tumor, and vesiculation of the skin may occur. Epilation is frequently not complete and in some instances hair re-growth will occur even though therapy continues.

Cellulitis and phlebitis may result if extravasation occurs during intravenous injection. If the extravasation is excessive, sloughing may occur.

You can report any suspected adverse reactions associated with the use of health products to the Canada Vigilance Program by one of the following 3 ways:

---

Report online at [www.healthcanada.gc.ca/medeffect](http://www.healthcanada.gc.ca/medeffect)

Call toll-free at 1-866-234-2345

Complete a Canada Vigilance Reporting Form and:

- Fax toll-free to 1-866-678-6789, or
- Mail to: Canada Vigilance Program  
Health Canada  
Postal Locator 0701E  
Ottawa, ON K1A 0K9

Postage paid labels, Canada Vigilance Reporting Form and the adverse reaction reporting guidelines are available on the MedEffect™ Canada Web site at [www.healthcanada.gc.ca/medeffect](http://www.healthcanada.gc.ca/medeffect).

*NOTE: Should you require information related to the management of side effects, contact your health professional. The Canada Vigilance Program does not provide medical advice.*

---

## **SYMPTOMS AND TREATMENT OF OVERDOSAGE**

|   |
|---|
| For management of a suspected drug overdose, contact your regional Poison Control Centre immediately. |
|---|

Side effects are dose related. Thus, patients can expect to experience these effects in an exaggerated manner if more than the recommended dose of vinblastine is given. As well, neurotoxicity may occur similar to that seen with vincristine sulfate. Management of suspected over-dosage of vinblastine should include the following:

1. Administer an antiemetic drug which usually controls nausea and vomiting.
2. Administer phenobarbital in anti-convulsant doses.
3. Be alert for the development of ileus which may necessitate non-surgical decompression of the gastrointestinal tract.
4. Monitor the patient's cardiovascular system.
5. Carry out daily blood counts as a guide for transfusion requirements. The most serious effect of an excessive dose of vinblastine, which may be life-threatening, is granulopoiesis.

## DOSAGE AND ADMINISTRATION

There are variations in the depth of the leukopenic response which follows therapy with vinblastine. For this reason, it is recommended that Vinblastine Sulfate Injection be given no more frequently than once every 7 days. It is wise to initiate therapy with a single intravenous dose of  $3.7 \text{ mg/m}^2$  of body surface area (bsa). Thereafter, white blood cell counts should be made to determine the patient's sensitivity to vinblastine.

A simplified and conservative incremental approach to dosage at weekly intervals may be outlined as follows:

### First Dose:

Adults:  $3.7 \text{ mg/m}^2$  bsa  
Children:  $2.5 \text{ mg/m}^2$  bsa

### Second Dose:

Adults:  $5.5 \text{ mg/m}^2$  bsa  
Children:  $3.75 \text{ mg/m}^2$  bsa

### Third Dose:

Adults:  $7.4 \text{ mg/m}^2$  bsa  
Children:  $5.0 \text{ mg/m}^2$  bsa

### Fourth Dose:

Adults:  $9.25 \text{ mg/m}^2$  bsa  
Children:  $6.25 \text{ mg/m}^2$  bsa

### Fifth Dose:

Adults:  $11.1 \text{ mg/m}^2$  bsa  
Children:  $7.5 \text{ mg/m}^2$  bsa

The above mentioned increases may be used until a maximum dose (not exceeding  $18.5 \text{ mg/m}^2$  bsa for adults and  $12.5 \text{ mg/m}^2$  bsa for children) is reached. The dose should not be increased after that dose which has reduced the white cell count to approximately  $3,000 \text{ cells/mm}^3$ . In some adults,  $3.7 \text{ mg/m}^2$  bsa may produce this leukopenia. Other adults may require more than  $11.1 \text{ mg/m}^2$  bsa and, very rarely, as much as  $18.5 \text{ mg/m}^2$  bsa may be necessary. For most adult patients, however, the weekly dosage will prove to be  $5.5$  to  $7.4 \text{ mg/m}^2$  bsa.

When the dose of vinblastine which will produce the above degree of leukopenia has been established, a dose one increment smaller than this should be administered at weekly intervals for maintenance. Thus, the patient is receiving the maximum dose that does not cause leukopenia. It should be emphasized that, even though 7 days have elapsed, the next dose of vinblastine should not be given until the white cell count has returned to at least  $4,000/\text{mm}^3$ . In some cases, oncolytic activity may be encountered before leukopenic effect. When this occurs, there is no need to increase the size or subsequent doses.

The duration of maintenance therapy varies according to the disease being treated and the

combination of antineoplastic agents being used. There are differences of opinion regarding the duration of maintenance therapy with the same protocol for a particular disease; for example, various durations have been used with the MOPP program in treating Hodgkin's disease. Prolonged chemotherapy for maintaining remissions involves several risks, among which are life-threatening infectious diseases, sterility and possibly the appearance of other cancers through suppression of immune surveillance. In some disorders, survival following complete remission may not be as prolonged as that achieved with shorter periods of maintenance therapy. On the other hand, failure to provide maintenance therapy in some patients may lead to unnecessary relapse. Failure to provide maintenance therapy for at least two years after complete remission in patients with testicular cancer often results in early relapse.

The dose of vinblastine (calculated to provide the desired amount) may be injected either into the tubing of a running intravenous infusion or directly into a vein. The latter procedure is readily adaptable to out-patient therapy. In either case, the injection may be completed in about one minute. If care is taken to ensure that the needle is securely within the vein and that no solution containing vinblastine is injected extravascularly, cellulitis and/or phlebitis will not occur.

To minimize further the possibility of extravascular spillage, it is suggested that the syringe and needle be rinsed with venous blood before withdrawal of the needle. The dose should not be diluted in large volumes of diluent (i.e. 100 to 250 mL) or given intravenously for prolonged periods (ranging from 30 to 60 minutes or more), since this frequently results in irritation of the vein and increases the chances of extravasation.

Because of the enhanced possibility of thrombosis, it is considered inadvisable to inject a solution of vinblastine into an extremity in which the circulation is impaired or potentially impaired by such conditions as compressing or invading neoplasm, phlebitis or varicosity.

## **CAUTION**

If leakage into the surrounding tissue should occur during intravenous administration of vinblastine, it may cause considerable irritation. The injection should be discontinued immediately and any remaining portion of the dose should then be introduced into another vein. Local injection of hyaluronidase and the application of moderate heat to the area of leakage help disperse the drug and are thought to minimize discomfort and the possibility of cellulitis.




## PHARMACEUTICAL INFORMATION

### Drug substance

**Proper Name:** Vinblastine Sulfate

**Chemical Name:** Vincalokoblastine (VBL) Sulfate

**Structural Formula:**



**Molecular Formula:**  $C_{46}H_{58}O_9N_4 \cdot H_2SO_4$

**Molecular Weight:** 909.07 g/mol

**Description:** Vinblastine Sulfate is the sulfate salt of an alkaloid obtained from *Vinca rosea Linn* (a flowering herb commonly known as the periwinkle) or *Catharanthus roseus G. Don*. Vinblastine sulfate is a white to slightly yellow, odorless, hygroscopic amorphous or crystalline powder. It is soluble in water and methyl alcohol, slightly soluble in chloroform and alcohol and practically insoluble in ether.

**Composition:** Vinblastine Sulfate Injection contains 1 mg/mL of Vinblastine Sulfate with 0.9% Sodium Chloride in Water for Injection as a sterile unpreserved solution. Sodium Hydroxide and/or Sulphuric Acid as pH adjusters.

**Stability and Storage Recommendations:**

Vinblastine Sulfate Injection should be stored at 2°C-8°C, protected from light. Protect solution from freezing.

Single use vials. Discard unused portion.

**As with all parenteral drug products, intravenous admixtures should be inspected visually for clarity, particulate matter, precipitate, discoloration and leakage prior to administration, whenever solution and container permit. Solutions showing haziness, particulate matter, precipitate, discoloration or leakage should not be used.**

**SPECIAL INSTRUCTIONS FOR HANDLING OF CYTOTOXIC DRUGS**

The following are precautionary measures recommended in the handling and preparation of cytotoxic agents such as vinblastine:

1. The procedure should be carried out in a vertical laminar flow hood (Biological Safety Cabinet - Class II).
2. PVC gloves, safety glasses, disposable gowns and masks should be worn by personnel.
3. All vials, syringes, needles and other materials which have come in contact with vinblastine should be segregated and destroyed by incineration. Sealed containers may explode if a tight seal exists. If incineration is unavailable, neutralization using 5% sodium hypochlorite or 5% sodium thiosulfate should be carried out instead.
4. Bi-annual hematologic examinations should be performed on personnel regularly involved in the handling and preparation of vinblastine.

**DOSAGE FORMS, COMPOSITION AND PACKAGING**

Each mL of Vinblastine Sulfate Injection contains vinblastine sulfate 1 mg, sodium chloride 9 mg, sodium hydroxide and/or sulfuric acid to adjust pH and water for injection.

Vinblastine Sulfate Injection, 1 mg/mL, is available in single use vials of 10 mg/10 mL (single packs).

**REFERENCES**

1. Hospira Healthcare Corporation, Prescribing Information, Vinblastine Sulfate Injection (1 mg/mL), Control No.: 114804, Date of Preparation: June 18, 2007.

**Sandoz Canada Inc.  
Boucherville, Québec**