

## **PRODUCT MONOGRAPH**

**Pr pms-HYDROCHLOROTHIAZIDE**

Hydrochlorothiazide Tablets USP

12.5 mg, 25 mg, 50 mg & 100 mg

**DIURETIC - ANTIHYPERTENSIVE**

**PHARMASCIENCE INC.**

6111 Royalmount Ave., Suite 100

Montréal, Canada

H4P 2T4

[www.pharmascience.com](http://www.pharmascience.com)

Date of Preparation

October 31, 2013

**Control No.** 168496

**Table of Contents**

**PART I: HEALTH PROFESSIONAL INFORMATION** ..... 3

    SUMMARY PRODUCT INFORMATION..... 3

    INDICATIONS AND CLINICAL USE ..... 3

    CONTRAINDICATIONS..... 4

    WARNINGS AND PRECAUTIONS ..... 4

    ADVERSE REACTIONS ..... 7

    DRUG INTERACTIONS ..... 8

    DOSAGE AND ADMINISTRATION ..... 11

    OVERDOSAGE..... 13

    ACTION AND CLINICAL PHARMACOLOGY ..... 13

    STORAGE AND STABILITY ..... 14

    DOSAGE FORMS, COMPOSITION AND PACKAGING..... 14

**PART II: SCIENTIFIC INFORMATION** ..... 16

    PHARMACEUTICAL INFORMATION ..... 16

    DETAILED PHARMACOLOGY ..... 17

    TOXICOLOGY..... 17

    REFERENCES..... 19

**PART III: CONSUMER INFORMATION** ..... 21

# Pr pms-HYDROCHLOROTHIAZIDE

Hydrochlorothiazide Tablets USP

## PART I: HEALTH PROFESSIONAL INFORMATION

### SUMMARY PRODUCT INFORMATION

Route of Administration	Dosage Form/ Strength	All Non-medicinal Ingredients
Oral	tablet 12.5 mg, 25 mg, 50 mg & 100 mg	Alginic acid, colloidal silicon dioxide, FD&C Yellow No. 6 Aluminum Lake, magnesium stearate, microcrystalline cellulose and sodium carboxymethylcellulose.

### INDICATIONS AND CLINICAL USE

pms-HYDROCHLOROTHIAZIDE (hydrochlorothiazide) is indicated for the treatment of:

- Edema
- Hypertension
- Toxemia of Pregnancy

#### Edema

pms-HYDROCHLOROTHIAZIDE (hydrochlorothiazide) is indicated in edema associated with congestive heart failure, hepatic cirrhosis, corticosteroid and estrogen therapy, premenstrual tension with edema and in edema of renal origin (i.e. nephrotic syndrome, acute glomerulonephritis and chronic renal disease). In obese patients in whom fluid retention is a complicating factor, it may help to initiate a loss of fluid and, thus of weight.

#### Hypertension

pms-HYDROCHLOROTHIAZIDE may be used alone or as an adjunct to other antihypertensive drugs. Since it enhances the action of these agents, their dosage must be reduced to avoid an excessive drop in pressure and other unwanted side effects.

#### Toxemia of Pregnancy

pms-HYDROCHLOROTHIAZIDE may be effective in the treatment of toxemia of pregnancy (including eclampsia).

**Geriatrics (> 65 years of age):** No data is available.

**Pediatrics (0 to 16 years of age):** No data is available.

## CONTRAINDICATIONS

- pms-HYDROCHLOROTHIAZIDE (hydrochlorothiazide), as all diuretics, is contraindicated in anuria.
- pms-HYDROCHLOROTHIAZIDE should be discontinued if increasing azotemia and oliguria occur during treatment of severe progressive renal disease.
- pms-HYDROCHLOROTHIAZIDE is contraindicated in persons known to be sensitive to hydrochlorothiazide or to other sulfonamide-derived drugs.
- Patients who are hypersensitive to any ingredient in the formulation of pms-HYDROCHLOROTHIAZIDE or component of the container. For a complete listing, see the **Dosage Forms, Composition and Packaging** section of the product monograph.

## WARNINGS AND PRECAUTIONS

### General

Patients on long therapy with hydrochlorothiazide are required to be on potassium rich diet. Periodic determinations of serum electrolytes to detect possible electrolyte imbalance should be performed.

The antihypertensive effects of the drug may be enhanced in the postsympathectomy patient.

### Carcinogenesis and Mutagenesis

No data available.

### Cardiovascular

No data available.

### Ear/Nose/Throat

No data available.

### Endocrine and Metabolism

**Calcium:** Calcium excretion is decreased by thiazides.

**Chloride:** Chloride deficiency is generally mild and does not require specific treatment except under special conditions such as renal or/and hepatic disease.

**Dilutional Hyponatremia:** Dilutional hyponatremia may occur in edematous patients in hot weather; appropriate therapy is water restriction rather than administration of salt except when hyponatremia is life threatening. In actual salt depletion, appropriate replacement is the therapy

of choice.

**Fluid and electrolyte imbalance:** All patients receiving thiazide should be observed for clinical signs of fluid or electrolyte imbalance: namely hyponatremia, hypochloremic alkalosis, and hypokalemia. Serum and urine electrolyte determinations are particularly important when the patient is vomiting excessively, or receiving parenteral fluids. Warning signs of serum electrolyte imbalance, irrespective of cause are: dryness of mouth, thirst, weakness, lethargy, drowsiness, restlessness, muscle pains or cramps, muscular fatigue, hypotension, oliguria, tachycardia, and gastrointestinal disturbances such as nausea and vomiting. Serum electrolytes may also be influenced by medication such as digitalis.

**Hyperuricemia:** Hyperuricemia may occur or frank gout may be precipitated in certain patients receiving thiazide therapy.

**Hypokalemia:** Hypokalemia may develop, especially with rapid diuresis, when severe cirrhosis is present or during concomitant use of corticosteroids or ACTH. Deficient oral electrolyte intake will also contribute to hypokalemia. Hypokalemia may sensitize or exaggerate the response of the heart to the toxic effects of digitalis (e.g. increased ventricular irritability). Hypokalemia may be avoided or treated by the use of potassium supplements.

**Insulin:** Insulin requirements in diabetic patients may be increased, decreased, or remain unchanged. Latent diabetes mellitus may become manifest during thiazide therapy. Concomitant therapy with lithium is not recommended with diuretics because of the reduction of renal clearance of lithium and therefore an added risk of lithium toxicity.

**Parathyroid gland:** Pathological changes in the parathyroid gland with hypercalcemia and hypophosphatemia have been observed in a few patients on prolonged thiazide therapy. The common complications of hyperparathyroidism such as renal lithiasis, bone resorption, and peptic ulceration have not been reported. Use of thiazides should be discontinued before carrying out tests for parathyroid function.

**Protein bound iodine (PBI):** Thiazides may decrease serum PBI levels without signs of thyroid disturbance.

### **Gastrointestinal**

Non-specific small bowel lesions consisting of stenosis with or without ulceration, may occur in association with the administration of enteric coated potassium salts, alone or with oral diuretics. These small bowel lesions have caused obstruction, hemorrhage and perforation. Surgery was frequently required and deaths have occurred. Available information tends to implicate enteric coated potassium salts, although lesions of this type also occur spontaneously. Such preparations should be used only when adequate dietary supplementation is not practical, and should be discontinued immediately if abdominal pain, distention, nausea, vomiting or gastrointestinal bleeding occur.

### **Genitourinary**

No data available.

### **Hematologic**

No data available.

### **Hepatic/biliary/Pancreatic**

pms-HYDROCHLOROTHIAZIDE should be used with caution in patients with impaired hepatic function or progressive liver disease, since minor alterations of fluid and electrolyte balance or of serum ammonia may precipitate hepatic coma.

### **Immune**

The possibility of exacerbation or activation of systemic lupus erythematosus has been reported.

Sensitivity reactions may occur in patients with or without a history of allergy or bronchial asthma.

### **Neurologic**

No data available.

### **Ophthalmologic**

#### **Acute Myopia and Secondary Angle-Closure Glaucoma**

Hydrochlorothiazide, a sulfonamide, can cause an idiosyncratic reaction, resulting in acute transient myopia and acute angle-closure glaucoma. Symptoms include acute onset of decreased visual acuity or ocular pain and typically occur within hours to weeks of drug initiation. Untreated acute angle-closure glaucoma can lead to permanent vision loss.

The primary treatment is to discontinue hydrochlorothiazide as rapidly as possible. Prompt medical or surgical treatments may need to be considered if the intraocular pressure remains uncontrolled. Risk factors for developing acute angle-closure glaucoma may include a history of sulphonamide or penicillin allergy.

### **Peri-Operative Considerations**

No data available.

### **Psychiatric**

No data available.

### **Renal**

In progressive renal impairment, therapy with pms-HYDROCHLOROTHIAZIDE should be withheld or discontinued.

Hydrochlorothiazide may commence or precipitate azotemia. It should be used with caution in patients with severely impaired renal function to avoid toxic or cumulative effect. If azotemia becomes more severe and oliguria occurs during treatment of patients with severe renal disease, administration of the diuretic must be stopped.

**Respiratory**

No data available.

**Sensitivity/Resistance**

No data available.

**Sexual Function/Reproduction**

No data available.

**Skin**

No data available.

**Special Population****Pregnant Women**

Thiazides cross the placental barrier and appear in cord blood. When hydrochlorothiazide is used in pregnancy or in women of child-bearing age, the potential benefits of the drug should be weighed against the possible hazards to the fetus. These hazards include fetal or neonatal jaundice, thrombocytopenia, and possibly other adverse reactions which have occurred in the adult.

The routine use of diuretics in otherwise healthy pregnant women with or without mild edema is not indicated.

**Nursing Women**

Since thiazides appear in breast milk, hydrochlorothiazide is contraindicated in nursing mothers. If use of the drug is deemed essential, the patient should stop nursing.

**Pediatrics (0 to 18 years of age)**

Safety and effectiveness in children under 18 years of age have not been established.

**Geriatrics**

Safety and effectiveness in adults over 65 years of age have not been established.

**ADVERSE REACTIONS**

Whenever adverse reactions are moderate or severe, thiazide dosage should be reduced or therapy withdrawn.

Cardiovascular: Orthostatic hypotension (may be aggravated by alcohol, barbiturates, or narcotics).

Central nervous system: Dizziness, vertigo, paresthesias, headache, xanthopsia.

Gastrointestinal system: Anorexia, gastric irritation, nausea, vomiting, cramps, diarrhea, constipation, jaundice (intrahepatic cholestatic jaundice), pancreatitis, sialadenitis.

Hematologic: Leukopenia, agranulocytosis, thrombocytopenia, aplastic anemia.

Hypersensitivity: Purpura, photosensitivity, rash, urticaria, necrotizing angitis (vasculitis), fever, respiratory distress including pneumonitis, anaphylactic reactions.

Other: Hyperglycemia, glycosuria, hyperuricemia, muscle spasm, weakness, restlessness, transient blurred vision.

## DRUG INTERACTIONS

Hydrochlorothiazide adds to or potentiates the action of other antihypertensive drugs. Potentiation occurs especially with ganglionic or peripheral adrenergic blocking drugs.

### Drug-Drug Interactions

Proper Name	Ref.	Effect	Clinical comment
Norepinephrine	C	Thiazides may decrease arterial responsiveness to norepinephrine. This diminution is not sufficient to preclude effectiveness of the pressor agent for therapeutic use.	
Alcohol, barbiturates, or narcotics	C	Potentiation of orthostatic hypotension may occur.	Avoid alcohol, barbiturates or narcotics, especially with initiation of therapy.
Amphotericin B	T	Amphotericin B increases the risk of hypokalemia induced by thiazide diuretics	Monitor serum potassium level.
Antidiabetic agents (e.g. insulin and oral hypoglycemic agents)	CT	Thiazide-induced hyperglycemia may compromise blood sugar control. Depletion of serum potassium augments glucose intolerance.	Monitor glycemic control, supplement potassium if necessary, to maintain appropriate serum potassium levels, and adjust diabetes medications as required.



<b>Proper Name</b>	<b>Ref.</b>	<b>Effect</b>	<b>Clinical comment</b>
Antihypertensive drugs	CT	Hydrochlorothiazide may potentiate the action of other antihypertensive drugs (e.g. guanethidine, methyldopa, beta-blockers, vasodilators, calcium channel blockers, ACEI, ARB, and direct renin inhibitors).	
Antineoplastic drugs, including cyclophosphamide and methotrexate	C	Concomitant use of thiazide diuretics may reduce renal excretion of cytotoxic agents and enhance their myelosuppressive effects.	Hematological status should be closely monitored in patients receiving this combination. Dose adjustment of cytotoxic agents may be required.
Bile acid sequestrants, eg. cholestyramine	CT	Bile acid sequestrants bind thiazide diuretics in the gut and impair gastrointestinal absorption by 43-85%. Administration of thiazide 4 hours after a bile acid sequestrant reduced absorption of hydrochlorothiazide by 30-35%.	Give thiazide 2-4 hours before or 6 hours after the bile acid sequestrant. Maintain a consistent sequence of administration. Monitor blood pressure, and increase dose of thiazide, if necessary.
Calcium and vitamin D supplements	C	Thiazides decrease renal excretion of calcium and increase calcium release from bone.	Monitor serum calcium, especially with concomitant use of high doses of calcium supplements. Dose reduction or withdrawal of calcium and/or vitamin D supplements may be necessary.
Carbamazepine	C	Carbamazepine may cause clinically significant hyponatremia. Concomitant use with thiazide diuretics may potentiate hyponatremia.	Monitor serum sodium levels. Use with caution.

<b>Proper Name</b>	<b>Ref.</b>	<b>Effect</b>	<b>Clinical comment</b>
Corticosteroids, and adrenocorticotrophic hormone (ACTH)	T	Intensified electrolyte depletion, particularly hypokalemia, may occur.	Monitor serum potassium, and adjust medications, as required.
Digoxin	CT	Thiazide-induced electrolyte disturbances, i.e. hypokalemia, hypomagnesemia, increase the risk of digoxin toxicity, which may lead to fatal arrhythmic events.	Concomitant administration of hydrochlorothiazide and digoxin requires caution. Monitor electrolytes and digoxin levels closely. Supplement potassium or adjust doses of digoxin or thiazide, as required.
Drugs that alter GI motility, i.e., anticholinergic agents, such as atropine and prokinetic agents, such as metoclopramide, domperidone	CT, T	Bioavailability of thiazide diuretics may be increased by anticholinergic agents due to a decrease in gastrointestinal motility and gastric emptying. Conversely, prokinetic drugs may decrease the bioavailability of thiazide diuretics.	Dosage adjustment of thiazides may be required
Gout medications (allopurinol, uricosurics, xanthine oxidase inhibitors)	T, RC	Thiazide-induced hyperuricemia may compromise control of gout by allopurinol and probenecid. The co-administration of hydrochlorothiazide and allopurinol may increase the incidence of hypersensitivity reactions to allopurinol.	Dosage adjustment of gout medications may be required.
Lithium	CT	Thiazide diuretics reduce the renal clearance of lithium and add a high risk of lithium toxicity.	Concomitant use of thiazide diuretics with lithium is generally not recommended. If such use is deemed necessary, reduce lithium dose by 50% and monitor lithium levels closely.

<b>Proper Name</b>	<b>Ref.</b>	<b>Effect</b>	<b>Clinical comment</b>
Nonsteroidal anti-inflammatory drugs (NSAID)	CT	NSAID-related retention of sodium and water antagonises the diuretic and antihypertensive effects of thiazides.  NSAID-induced inhibition of renal prostaglandins leading to decreases of renal blood flow, along with thiazide-induced decreases in GFR may lead to acute renal failure. Patients with heart failure may be at particular risk.	If combination use is necessary, monitor renal function, serum potassium, and blood pressure closely. Dose adjustments may be required.
Selective serotonin reuptake inhibitors (SSRIs, e.g. citalopram, escitalopram, sertraline)	T,C	Concomitant use with thiazide diuretics may potentiate hyponatremia.	Monitor serum sodium levels. Use with caution.
Skeletal muscle relaxants of the curare family, eg., tubocurare	C	Thiazide drugs may increase the responsiveness of some skeletal muscle relaxants, such as curare derivatives	
Topiramate	CT	Additive hypokalemia. Possible thiazide-induced increase in topiramate serum concentrations.	Monitor serum potassium and topiramate levels. Use potassium supplements, or adjust topiramate dose as necessary.

Legend: C = Case Study; RCS =Retrospective Cohort Study; CT = Clinical Trial; T =Theoretical

### **Drug-Food Interactions**

Interactions with food have not been established.

### **Drug-Herb Interactions**

Interactions with herbal products have not been established.

### **Drug-Laboratory Test Interactions**

There are no known interactions of hydrochlorothiazide with commonly used laboratory tests.

## **DOSAGE AND ADMINISTRATION**

Therapy should be individualized according to the patients requirement. Use the smallest dosage necessary to achieve the required response.

## **Dosing Considerations**

### **Adult patients**

#### **Diuresis**

The recommended adult dosage is 50 mg to 100 mg once or twice a day. Many patients respond to intermittent therapy, i.e. administration on alternate days or on three to five days each week.

With an intermittent schedule, excessive response and the resulting undesirable electrolyte imbalance are less likely to occur.

#### **Toxemia of pregnancy**

The recommended dosage is 100 mg daily or, in severe cases and for brief periods, 200 mg daily (in divided doses). Frequency of administration may range from once every four days to daily.

#### **Premenstrual tension with edema**

The recommended dosage is 25 mg to 50 mg once or twice a day from the first appearance of symptoms until onset of the menses.

#### **Control of Hypertension**

The usual recommended starting dosage is 50 mg or 100 mg a day as a single or divided dose. Dosage is increased or decreased according to the blood pressure response of the patient. Some patients may require doses of 200 mg a day in divided doses.

Careful observation for changes in blood pressure must be made when pms-HYDROCHLOROTHIAZIDE (hydrochlorothiazide) is used with other antihypertensive drugs, especially during initial therapy. The dosage of other agents must be reduced by at least 50%, as soon as it is added to the regimen, to prevent excessive drop in blood pressure. As the blood pressure falls under the potentiating effect of this agent, a further reduction in dosage, or discontinuation of other antihypertensive drugs may be necessary. A single daily dose as low as 12.5 mg of hydrochlorothiazide could be used in combination with another antihypertensive.

In the case of hypertension monotherapy, doses as low as a single daily dose 12.5 mg may be effective (especially in the elderly or as a starting dose), as well as a daily dose of 25 mg given in two divided doses.

### **Infants and Children**

The usual recommended pediatric dosage is based on 1.0 mg of pms-HYDROCHLOROTHIAZIDE per pound of body weight per day in two doses. Infants under 6 months of age may require up to 1.5 mg per pound per day in two doses.

On this basis, infants up to 2 years of age may be given 12.5 mg to 37.5 mg daily in two doses.

Children from 2 to 12 years of age may be given 37.5 mg to 100 mg daily in two doses. Dosage in both age groups should be based on body weight.

## **OVERDOSAGE**

### **Symptoms**

Overdosage of hydrochlorothiazide may produce diuresis accompanied with electrolyte imbalance (hypokalemia, hyponatremia and hypochloremic alkalosis) and dehydration.

The symptoms are as follows: dryness of mouth, thirst, weakness, lethargy, drowsiness, restlessness, muscle pains or cramps, muscular fatigue, hypotension, oliguria, tachycardia, gastrointestinal disturbances, mental confusion, delirium, convulsions, shock, coma.

Hypokalemia can sensitize or exaggerate the response of the heart to the toxic effects of digitalis (e.g. increased ventricular irritability).

Hydrochlorothiazide may precipitate hepatic coma in patients with cirrhosis; increase the effect of other antihypertensive agents and decrease arterial responsiveness to norepinephrine.

### **Treatment**

No specific antidote is available.

Treatment is symptomatic and supportive. Induce emesis or perform gastric lavage. Correct dehydration, electrolyte imbalance, hepatic coma, and hypotension by established procedures. Administer oxygen or artificial respiration for respiratory impairment.

For management of a suspected drug overdose, contact your regional Poison Control Center immediately.

## **ACTION AND CLINICAL PHARMACOLOGY**

### **Mechanism of Action**

Hydrochlorothiazide is a diuretic and an antihypertensive agent. The exact mechanism of the antihypertensive effect is unknown. Hydrochlorothiazide has no effect on normal blood pressure.

Hydrochlorothiazide affects the renal tubular mechanism of electrolyte reabsorption. It increases excretion of sodium and chloride in approximately equivalent amounts and reduces the rate of formation of solute-free water. Natriuresis causes a secondary loss of potassium and bicarbonate.

### **Pharmacokinetics**

**Absorption:** Hydrochlorothiazide is rapidly absorbed from the gastrointestinal tract. Onset of action after oral administration occurs in 2 hours and the peak effect at approximately 4 hours. Duration of action persists for approximately 6 to 12 hours.

**Distribution:** The drug is distributed throughout the extracellular space and does not accumulate in tissues other than the kidney. It passes readily through the placental barrier to the fetus.

**Metabolism:** Hydrochlorothiazide is not metabolized.

**Excretion:** Hydrochlorothiazide is eliminated rapidly by the kidney.

## **STORAGE AND STABILITY**

pms-HYDROCHLOROTHIAZIDE (hydrochlorothiazide) Tablets should be stored between 15°C and 30°C.

## **DOSAGE FORMS, COMPOSITION AND PACKAGING**

pms-HYDROCHLOROTHIAZIDE (hydrochlorothiazide) tablets are available for oral use in four dosage strengths of 12.5 mg, 25 mg, 50 mg and 100 mg hydrochlorothiazide.

### **Availability of Dosage Forms**

pms-HYDROCHLOROTHIAZIDE (hydrochlorothiazide) Tablets are supplied as:

- 12.5 mg tablets: Round, peach colored, flat faced, bevel edged tablets debossed with a “P” logo on one side and plain on the other side. Supplied in bottles of 500 and 1000 tablets.
- 25 mg tablets: Round, peach colored, flat faced, bevel edged tablets debossed with “25” score “P” on one side and plain on the other side. Supplied in bottles of 100, 500 and 1000 tablets.
- 50 mg tablets: Round, peach colored, flat faced, bevel edged tablets debossed with “50” score “P” on one side and plain on the other side. Supplied in bottles of 100, 1000 and 5000 tablets.
- 100 mg tablets: Round, peach-colored, flat faced, bevel edged tablets, debossed with “100” score “P” on one side and plain on the other side. Supplied in bottles of 100 tablets.

## **Composition**

pms-HYDROCHLOROTHIAZIDE (hydrochlorothiazide) 12.5 mg, 25 mg, 50 mg and 100 mg tablets contains 12.5 mg, 25 mg, 50 mg and 100 mg of hydrochlorothiazide, respectively. Non medicinal ingredients (alphabetical): alginic acid, colloidal silicon dioxide, FD&C Yellow No. 6 Aluminum Lake, magnesium stearate, microcrystalline cellulose and sodium carboxymethylcellulose.

## PART II: SCIENTIFIC INFORMATION

### PHARMACEUTICAL INFORMATION


#### Drug Substance

Proper Name: Hydrochlorothiazide

Chemical Name: 6-chloro-3,4-dihydro-2H-1,2,4-benzothiadiazine-7-sulfonamide  
1,1-dioxide

Molecular formula and molecular mass:  $C_7H_8ClN_3O_4S_2$ ; 297.72 g/mol

Structural formula:



Physicochemical properties: Hydrochlorothiazide is a white or almost white odourless crystalline powder with a slightly bitter taste. It is almost insoluble in water, benzene, chloroform, ether and dilute mineral acids, and soluble 1 in 500 of alcohol and 1 in 50 of acetone. It is freely soluble in dimethyl formamide, n-butylamine and solutions of alkali hydroxides..



## DETAILED PHARMACOLOGY

Orally, hydrochlorothiazide is an effective diuretic and antihypertensive agent. Diuresis is effected by inhibition of tubular resorption of electrolytes and an accompanying volume of water. Hydrochlorothiazide increases the excretion of sodium and chloride in approximately equivalent amounts and causes a simultaneous, usually minimal loss of bicarbonate. The excretion of ammonia is reduced slightly as a consequence of which concentrations of ammonia in the blood may be increased. Hydrochlorothiazide slightly increases the excretion of potassium. Calcium excretion is decreased and magnesium excretion is increased.

## TOXICOLOGY

### Acute Toxicity

SPECIES	ROUTE	LD <sub>50</sub> (mg/kg)
MOUSE	ORAL	10,000*
MOUSE	I.V.	884
RAT	ORAL	10,000*
RAT	I.P.	3,130*
RABBIT	I.V.	461
DOG	I.V.	1,000

Dogs tolerated at least 2,000 mg/kg orally without signs of toxicity.

\*Hydrochlorothiazide was administered as a suspension.

### Subacute Toxicity:

#### Rat

Hydrochlorothiazide administered to rats, orally as a suspension at doses of 500, 1,000 and 2,000 mg/kg/day, 5 days/week, for 3 weeks did not produce any toxic symptoms. Three of the ten rats which received 2,000 mg/kg/day of sodium hydrochlorothiazide salt died after the 5<sup>th</sup> day of treatment.

The mortality was attributed to pneumonia.

## Dog

Hydrochlorothiazide administered to dogs, orally at doses of 250, 500 and 1,000 mg/kg, 7 days/week for 8 weeks did not produce any observable adverse effects or gross signs of drug toxicity except for electrolytic imbalance.

## Chronic Toxicity

### Rat and Dog

The results of 6-month chronic oral toxicity on hydrochlorothiazide in rats and dogs indicated no toxicity attributable to the drug administered to rats at doses of up to 2 grams/kg/day and to dogs at doses of up to 250 mg/kg/day. On gross examination the following changes were observed in the dog: slight depression of plasma potassium; small amounts of yellow crystalline precipitate in the bladder in two of twelve dogs tested. Histomorphologic studies did not show any drug related changes.

## REFERENCES

- Applied Pharmacology. The Kidneys, Ch. 310. Classification of Diuretics, pp. 587-589. W.B. Saunders Company, Philadelphia; Toronto, 1976.
- Physicians' Desk Reference (PDR), 1977, pp. 1090. HydroDiuril (hydrochlorothiazide, MSD) U.S.P.
- FROHLICH, E.D. Hypertension pp. 208-213. Current Therapy, 1976. Conn, F. Editor. W.B. Saunders Company, Philadelphia, London, Toronto
- MUDGE, G.H. Diuretics and other agents employed in the mobilization of edema fluid. The Pharmacological Basis of Therapeutics, Ch. 39, pp. 854-858. Goodman & Gilman, 4<sup>th</sup> Edition, 1970.
- SACKNER, M.A.; WALLACK, A.A. BELLETS, S. The Diuretic Effects of hydrochlorothiazide in Congestive Heart Failure, Cirrhosis, Chronic Renal Disease & Hypertension: Preliminary Report based on a study of 28 cases. Am. J. M. Sc., 237; 575-584, May 1959.
- ZATUCHNI, J.; KING, W.; RESINSKI, M. Hydrochlorothiazide, A new Saluretic. Am. J. M. Sc., 237; 479, April 1959.
- FORD, R.V. Comparative Studies of Th e Newer Diuretics. Annals New York Academy of Sciences, 88, 809-814, 1960.
- FUCKS, M; MOYER, J.H.; NEWMAN, B.E. Human Clinical Pharmacology of the Newer Diuretics: Benzothiadiazine & Phthalimidine. Annals New York Academy of Sciences, 88, 795-808, October 1960.
- RENNICK, B. Animal Pharmacology of the New Diuretics: Benzothiadiazines, Spirolactones & Phthalimidines. Annals New York Academy of Sciences, 88, 785-794, October 1960.
- GROLLMAN, A.; FURNESS, F. Present concept of the Mechanism of Urine Formation and of Diuretic Action. Annals of the New York Academy of Sciences, 88 (4): 771-1020, Oct. 1960.
- BORHANI, N.O. Chlorothiazide & Hydrochlorothiazide: A Comparative Study of their hypotensive, Saluretic & Hyperuricemic action. Ann Int. Med., 53, 342, 1960.
- WINER, B.N. The Antihypertensive Actions of Benzothiadiazines. Circulation 23: 211, Feb. 1961.

- ROSENBLOOM, S.E.; SHAPER, R.P.; GOLDSBLOOM, S.; PINCUS, J.; SHAPIRO, A.P. II. Comparison of Chlorothiazide, Hydrochlorothiazide and a Placebo in the Hypertensive Patient. *New England J. Med.*; 264, 164, Jan. 1961.
- KJELLBO, H.; STAKEBERG, H.; MELLGREN, J. Possibly Thiazide - Induced Renal Necrotising Vasculitis. *Lancet*; 1, 1034, May 1965.
- CAFRUNY, E.J. How Diuretics work. *Geriatrics*; 22, 107, Jan. 1967.
- BRYANT, J.M. SCHVARTZ, N.; ROQUE, M.; FLETCHER, L.; FERTIG, H.; LAULER, D.P. The Hypotensive Effects of Chlorothiazide and Hydrochlorothiazide. *Am. J. Cardiol.*; 7, 392, March 1961.
- WOLF, R.L.; MENDLOWITZ, M.; ROBOZ, J.; GITLOW, S.E. Treatment of Hypertension with Antihypertensive Diuretic Drugs. *Am. Heart J.*; 72, 692, Nov. 1966.
- TANNENBAUM, P.J.; CROSLEY, A.P. A Comparison of the Effects of Hydrochlorothiazide and Hydrochlorothiazide on Electrolyte Balance. *Clin. Pharmacol. & Therap.*; 7, 777, Nov.-Dec. 1966.
- KOLODNY, A.L. Technic of Drug Evaluation in Hypertension.. *New York J. Med.*, 62, 1585, May 1962.
- SALERNO, L.J.; STONE, M.L. The Use of Diuretics in the Pregnant Patient. *Annals of the New York, Academy of Sciences*, 88, 881-889, Oct. 1960.

**PART III: CONSUMER INFORMATION**

**Pr** **pms-HYDROCHLOROTHIAZIDE**  
Hydrochlorothiazide Tablets USP

This leaflet is part III of a three-part “Product Monograph” published when pms-HYDROCHLOROTHIAZIDE was approved for sale in Canada and is designed specifically for Consumers.

Read this carefully before you start taking pms-HYDROCHLOROTHIAZIDE and each time you get a refill. This leaflet is a summary and will not tell you everything about pms-HYDROCHLOROTHIAZIDE. Talk to your doctor, nurse, or pharmacist about your medical condition and treatment and ask if there is any new information about pms-HYDROCHLOROTHIAZIDE.

**ABOUT THIS MEDICATION**

**What the medication is used for:**

**Adults:**

- Decreases swelling caused by fluid retention (edema) due to heart failure, liver disease, kidney disease, premenstrual tension, or corticosteroid and estrogen therapy.
- Lowers high blood pressure.
- Lowers pregnancy-induced high blood pressure.

**What it does:**

pms-HYDROCHLOROTHIAZIDE is a diuretic often called “water pill”. It increases urination. This lowers blood pressure and decreases swelling.

This medicine does not cure high blood pressure or edema. It helps to control them. Therefore, it is important to continue taking pms-HYDROCHLOROTHIAZIDE regularly even if you feel fine.

**When it should not be used:**

Do not take pms-HYDROCHLOROTHIAZIDE if you:

- Are allergic to hydrochlorothiazide or to any non-medicinal ingredient in the formulation.
- Are allergic to any sulfonamide-derived drugs (sulfa drugs); most of them have a medicinal ingredient that ends in “-MIDE”.
- Have difficulty urinating or produce no urine.
- Are breastfeeding. Hydrochlorothiazide passes into breast milk.

**What the medicinal ingredient is:**  
Hydrochlorothiazide.

**What the non-medicinal ingredients are:**  
Alginic acid, colloidal silicon dioxide, FD&C Yellow No. 6 Aluminum Lake, magnesium stearate, microcrystalline cellulose and sodium carboxymethylcellulose.

**What dosage forms it comes in:**  
**Tablets:** 12.5 mg, 25 mg, 50 mg & 100 mg

**WARNINGS AND PRECAUTIONS**

Before you use pms-HYDROCHLOROTHIAZIDE talk to your doctor, nurse or pharmacists if you:

- Are allergic to penicillin.
- Have diabetes, liver or kidney disease.
- Have lupus or gout.
- Are dehydrated or suffer from excessive vomiting, diarrhea, or sweating.
- Are less than 18 years old

**Hydrochlorothiazide in pms-HYDROCHLOROTHIAZIDE can cause Sudden Eye Disorders:**

- **Myopia:** sudden nearsightedness or blurred vision.
- **Glaucoma:** an increased pressure in your eyes, eye pain. Untreated, it may lead to permanent vision loss.

These eye disorders are related and can develop within hours to weeks of starting pms-HYDROCHLOROTHIAZIDE.

You may become sensitive to the sun while taking pms-HYDROCHLOROTHIAZIDE. Exposure to sunlight should be minimized until you know how you respond.

**Driving and using machines:** Before you perform tasks which may require special attention, wait until you know how you respond to pms-HYDROCHLOROTHIAZIDE. Dizziness, lightheadedness, or fainting can especially occur after the first dose and when the dose is increased.

**INTERACTIONS WITH THIS MEDICATION**

As with most medicines, interactions with other drugs are possible. Tell your doctor, nurse, or pharmacist about all the medicines you take, including drugs prescribed by other doctors, vitamins, minerals, natural supplements, or alternative medicines.

The following may interact with

pms-HYDROCHLOROTHIAZIDE:

- Alcohol, barbiturates (sleeping pills), or narcotics (strong pain medications). They may cause low blood pressure and dizziness when you go from lying or sitting to standing up.
- Amphotericin B, an antifungal drug.
- Anticancer drugs, including cyclophosphamide and methotrexate.
- Antidepressants, in particular selective serotonin reuptake inhibitors (SSRIs), including citalopram, escitalopram, and sertraline.
- Antidiabetic drugs, including insulin and oral medicines.
- Bile acid resins used to lower cholesterol.
- Calcium or vitamin D supplements.
- Corticosteroids used to treat joint pain and swelling.
- Digoxin, a heart medication.
- Drugs that slow down or speed up your bowels, including atropine, metoclopramide, and domperidone.
- Drugs used to treat epilepsy, including carbamazepine and topiramate.
- Gout medications, including allopurinol and probenecid.
- Lithium used to treat bipolar disease.
- Nonsteroidal anti-inflammatory drugs (NSAIDs), used to reduce pain and swelling. Examples include ibuprofen, naproxen, and celecoxib.
- Other blood pressure lowering drugs. When taken in combination with hydrochlorothiazide, they may cause excessively low blood pressure.
- Skeletal muscle relaxants used to relieve muscle spasms, including tubocurare.

pressure: The usual dose is a 100 mg. The doctor may briefly increase dosage to 200 mg.

Doses may be prescribed:

- once a day or
- every 4 days.
- For the treatment of swelling caused by fluid retention (edema): 25 mg to 50 mg once or twice a day.

**Usual Infant and Child dose:**

- Infants up to 24 months: 12.5 mg to 37.5 mg twice a day.
- Children 2 to 12 years old: 37.5 mg to 100 mg twice a day.

Overdose:

If you think you have taken too much pms-HYDROCHLOROTHIAZIDE contact your doctor, nurse, pharmacist, hospital emergency department or regional Poison control Center immediately, even if there are no symptoms.

Missed Dose:

If you have forgotten to take your dose during the day, carry on with the next one at the usual time. Do not double dose.

**SIDE EFFECTS AND WHAT TO DO ABOUT THEM**

Side effects may include:

- muscle cramps, spasms, and pain, weakness, restlessness
- dizziness, pins and needles in your fingers, headache
- constipation, diarrhea, nausea, vomiting, decreased appetite, upset stomach, enlargement of the glands in your mouth
- reduced libido
- bleeding under the skin, rash, red patches on the skin

**If any of these affects you severely, tell your doctor, nurse or pharmacist.**

pms-HYDROCHLOROTHIAZIDE can cause abnormal blood test results. Your doctor will decide when to perform blood tests and will interpret the results.

**PROPER USE OF THIS MEDICATION**

Take pms-HYDROCHLOROTHIAZIDE exactly as prescribed. It is recommended to take your dose at about the same time everyday.

pms-HYDROCHLOROTHIAZIDE can be taken with or without food. If pms-HYDROCHLOROTHIAZIDE causes upset stomach, take it with food or milk.

In your diet, be sure to include foods that contain potassium such as tomatoes, bananas, and beans.

**Usual Adult dose:**

- For the treatment of high blood pressure: 50 mg or 100 mg, once a day or as a divided dose as directed by your doctor. Your doctor may increase or decrease your dose.
- For the treatment of pregnancy-induced high blood

**SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM**

Symptom / effect	Talk with your doctor, nurse, or pharmacist		Stop taking drug and seek immediate medical help
	Only if severe	In all cases	
<b>Common</b>	<b>Low Blood Pressure:</b> dizziness, fainting, lightheadedness. May occur when you go from lying or sitting to standing up.	√	
	<b>Decreased levels of potassium in the blood:</b> irregular heartbeats, muscle weakness and generally feeling unwell		
<b>Uncommon</b>	<b>Allergic Reaction:</b> rash, hives, swelling of the face, lips, tongue or throat, difficulty swallowing or breathing		√
	<b>Kidney Disorder:</b> change in frequency of urination, nausea, vomiting, swelling of extremities, fatigue		√
	<b>Liver Disorder:</b> yellowing of the skin or eyes, dark urine, abdominal pain, nausea, vomiting, loss of appetite		√
	<b>Increased blood sugar:</b> frequent urination, thirst, and hunger	√	
	<b>Electrolyte Imbalance:</b> weakness, drowsiness, muscle pain or cramps, irregular heartbeat		√
<b>Rare</b>	<b>Decreased Platelets:</b> bruising, bleeding, fatigue and weakness		√
	<b>Decreased White Blood Cells:</b> infections, fatigue, fever, aches, pains, and flu-like symptoms		√

**SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM**

Symptom / effect	Talk with your doctor, nurse, or pharmacist		Stop taking drug and seek immediate medical help
	Only if severe	In all cases	
<b>Very rare</b> <b>Toxic Epidermal Necrolysis:</b> severe skin peeling, especially in mouth and eyes			√
<b>Unknown</b>	<b>Eye Disorders:</b> - <b>Myopia:</b> sudden near sightedness or blurred vision - <b>Glaucoma:</b> increased pressure in your eyes, eye pain		√
	<b>Anemia:</b> fatigue, loss of energy, weakness, shortness of breath.		√
	<b>Inflammation of the Pancreas:</b> abdominal pain that lasts and gets worse when you lie down, nausea, vomiting		√

*This is not a complete list of side effects. For any unexpected effects while taking pms-HYDROCHLOROTHIAZIDE, contact your doctor, nurse, or pharmacist.*

**HOW TO STORE IT**

Stored between 15°C and 30°C.

**Keep out of reach and sight of children.**

**REPORTING SUSPECTED SIDE EFFECTS**

You can report any suspected adverse reactions associated with the use of health products to the Canada Vigilance Program by one of the following 3 ways:

- Report online at [www.healthcanada.gc.ca/medeffect](http://www.healthcanada.gc.ca/medeffect)
- Call toll-free at 1-866-234-2345
- Complete a Canada Vigilance Reporting Form and:
  - Fax toll-free to 1-866-678-6789, or
  - Mail to: Canada Vigilance Program

Health Canada  
Postal Locator

0701E

Ottawa, Ontario

K1A 0K9

Postage paid labels, Canada Vigilance Reporting Form and the adverse reaction reporting guidelines are available on the MedEffect™ Canada Web site at [www.healthcanada.gc.ca/medeffect](http://www.healthcanada.gc.ca/medeffect).

*NOTE: Should you require information related to the management of side effects, contact your health professional. The Canada Vigilance Program does not provide medical advice.*

**MORE INFORMATION**

This document plus the full product monograph, prepared for health professionals, can be obtained by contacting Pharmascience Inc. at 1-888-550-6060.

This leaflet was prepared by  
**Pharmascience Inc.**  
Montréal, Canada  
H4P 2T4

[www.pharmascience.com](http://www.pharmascience.com)

Last revised: October 31, 2013