

**PRODUCT MONOGRAPH**

**Pr DIVALPROEX**

**Divalproex Sodium Enteric-Coated Tablets**

**125 mg, 250 mg and 500 mg valproic acid (as divalproex sodium)**

**House Standard**

**Antiepileptic**

**Sanis Health Inc.  
333 Champlain St. Suite 102  
Dieppe, New Brunswick  
E1A 1P2**

**DATE OF REVISION:**

**June 16, 2014**

**Control Number: 174536**

## Table of Contents

<b>PART I: HEALTH PROFESSIONAL INFORMATION .....</b>	<b>3</b>
SUMMARY PRODUCT INFORMATION .....	3
INDICATIONS AND CLINICAL USE .....	3
CONTRAINDICATIONS .....	4
WARNINGS AND PRECAUTIONS .....	5
ADVERSE REACTIONS .....	16
DRUG INTERACTIONS .....	23
DOSAGE AND ADMINISTRATION .....	29
OVERDOSAGE .....	31
ACTION AND CLINICAL PHARMACOLOGY .....	32
STORAGE AND STABILITY .....	35
DOSAGE FORMS, COMPOSITION AND PACKAGING .....	35
<b>PART II: SCIENTIFIC INFORMATION .....</b>	<b>36</b>
PHARMACEUTICAL INFORMATION .....	36
CLINICAL TRIALS .....	36
DETAILED PHARMACOLOGY .....	37
TOXICOLOGY .....	38
REFERENCES .....	42
<b>PART III: CONSUMER INFORMATION .....</b>	<b>47</b>

<sup>Pr</sup>**DIVALPROEX**  
Divalproex Sodium Enteric-Coated Tablets  
House Standard  
125 mg, 250 mg, and 500 mg

**PART I: HEALTH PROFESSIONAL INFORMATION**

**SUMMARY PRODUCT INFORMATION**

<b>Route of Administration</b>	<b>Dosage Form / Strength</b>	<b>Clinically Relevant Nonmedicinal Ingredients</b>
Oral	Enteric-coated tablets 125 mg, 250 mg and 500 mg	none <i>For a complete listing see <b>DOSAGE FORMS, COMPOSITION AND PACKAGING</b> section.</i>

**INDICATIONS AND CLINICAL USE**

DIVALPROEX (divalproex sodium) enteric-coated tablets are indicated for:

**Epilepsy**

- use as sole or adjunctive therapy in the treatment of simple or complex absence seizures, including petit mal, and is useful in primary generalized seizures with tonic-clonic manifestations.
- use adjunctively in patients with multiple seizure types which include either absence or tonic clonic seizures.

In accordance with the International Classification of Seizures, simple absence is defined as a very brief clouding of the sensorium or loss of consciousness (lasting usually 2 to 15 seconds), accompanied by certain generalized epileptic discharges without other detectable clinical signs.

Complex absence is the term used when other signs are also present.

**Acute Mania**

- the treatment of the manic episodes associated with bipolar disorder (DSM-III-R).

The safety and effectiveness of divalproex sodium for long-term use in mania, that is for more than 3 weeks, has not been evaluated in controlled trials.

DIVALPROEX is not indicated for use as a mood stabilizer in patients under 18 years of age. See (CONTRAINDICATIONS), and (WARNINGS AND PRECAUTIONS, Hepatic/Biliary/Pancreatic, Serious or Fatal Hepatotoxicity) for statement regarding serious or fatal hepatic dysfunction.

**Geriatrics (≥ 65 years of age):**

The safety and efficacy of divalproex sodium in elderly patients with epilepsy and mania has not been evaluated in clinical trials. Caution should thus be exercised in dose selection for an elderly patient, recognizing the more frequent hepatic and renal dysfunctions, and limited experience with divalproex sodium in this population. For a brief discussion, see [WARNINGS AND PRECAUTIONS, Special Populations, Geriatrics (≥ 65 years of age)], (DOSAGE AND ADMINISTRATION, Dosing in Elderly Patients) and (ACTION AND CLINICAL PHARMACOLOGY, Special Populations and Conditions, Geriatrics).

**Pediatrics (< 18 years of age):**

When divalproex sodium is used in children under the age of 2 years; it should be used with extreme caution and as a sole agent. Above the age of 2 years, experience in epilepsy has indicated that the incidence of fatal hepatotoxicity decreases considerably in progressively older patient groups. For a brief discussion, see [WARNINGS AND PRECAUTIONS, Special Populations, Pediatrics (< 18 years of age)]. The safety and effectiveness of divalproex sodium for the treatment of acute mania has not been studied in individuals below the age of 18 years.

**CONTRAINDICATIONS**

- DIVALPROEX (divalproex sodium) enteric-coated tablets should not be administered to patients with hepatic disease or significant hepatic dysfunction.
- DIVALPROEX is contraindicated in patients with known hypersensitivity to the drug. For a complete listing, see the (DOSAGE FORMS, COMPOSITION AND PACKAGING) section of the Product Monograph.
- DIVALPROEX is contraindicated in patients with known urea cycle disorders. See (WARNINGS AND PRECAUTIONS, Endocrine and Metabolism, Urea Cycle Disorders).

## WARNINGS AND PRECAUTIONS

### Serious Warnings and Precautions

- **Hepatotoxicity:** Hepatic failure resulting in fatalities has occurred in patients receiving divalproex sodium. These incidences usually occurred during the first 6 months of treatment with divalproex sodium. Caution should be observed when administering divalproex sodium to patients with a prior history of hepatic disease. Patients on multiple anticonvulsants, children, those with congenital metabolic disorders, those with severe seizure disorders accompanied by mental retardation, and those with organic brain disease may be at particular risk. Experience has indicated that children under the age of 2 years are at a considerably increased risk of developing fatal hepatotoxicity, especially those on multiple anticonvulsants. See (**WARNINGS AND PRECAUTIONS, Hepatic/Biliary/Pancreatic, Serious or Fatal Hepatotoxicity**).
- **Teratogenicity:** Divalproex sodium can produce teratogenic effects such as neural tube defects (e.g., spina bifida). Accordingly, the use of the medication in women of childbearing potential requires that the benefits of its use be weighed against the risk of injury to the fetus. See (**WARNINGS AND PRECAUTIONS, Special Populations, Pregnant Women**).
- **Pancreatitis:** Cases of life-threatening pancreatitis have been reported in both children and adults receiving divalproex sodium. Some of the cases have been described as hemorrhagic with a rapid progression from initial symptoms to death. Some cases have occurred shortly after initial use as well as after several years of use. See (**WARNINGS AND PRECAUTIONS, Hepatic/Biliary/Pancreatic, Pancreatitis**).

### General

#### **Interaction with Carbapenem Antibiotics**

Carbapenem antibiotics (ertapenem, imipenem, meropenem, doripenem) can reduce serum valproic acid concentrations to sub-therapeutic levels. This can result in loss of seizure control in epileptic patients or loss of efficacy in non-epileptics. In some cases of co-administration in epileptic patients, breakthrough seizures have occurred. Increasing valproic acid dose may not be sufficient to overcome this interaction. If co-administration is essential, serum valproic acid concentrations should be monitored daily after initiating carbapenem therapy. Alternative antibacterial or anticonvulsant therapy should be considered if serum valproic acid concentrations drop significantly or seizure control deteriorates. See (**DRUG INTERACTIONS, Drug-Drug Interactions, Table 2**).

#### **Patients with Special Diseases and Conditions**

There are in vitro studies that suggest valproate stimulates the replication of the Human Immunodeficiency Virus and Cytomegalovirus (HIV and CMV) viruses under certain experimental conditions. The clinical relevance of these in vitro data is unknown.

## **Carcinogenesis and Mutagenesis**

Long-term animal toxicity studies indicate that valproic acid is a weak carcinogen or promoter in rats and mice. The significance of these findings for humans is unknown at present. See (TOXICOLOGY, **Mutagenicity and Carcinogenicity**).

## **Endocrine and Metabolism**

### **Urea Cycle Disorders**

DIVALPROEX (divalproex sodium) enteric-coated tablets are contraindicated in patients with known urea cycle disorders. Hyperammonemic encephalopathy, sometimes fatal, has been reported following initiation of divalproex sodium in patients with urea cycle disorders, a group of uncommon genetic abnormalities, particularly ornithine transcarbamylase deficiency. Prior to initiation of divalproex sodium, evaluation for urea cycle disorders (UCD) should be considered in the following patients:

- those with a history of unexplained encephalopathy or coma, encephalopathy associated with protein load, pregnancy-related or postpartum encephalopathy, unexplained mental retardation, or history of elevated plasma ammonia or glutamine;
- those with signs and symptoms of UCD, for example, cyclical vomiting and lethargy, episodic extreme irritability, ataxia, low blood urea nitrogen (BUN), protein avoidance;
- those with a family history of UCD or a family history of unexplained infant deaths (particularly males);
- those with other signs or symptoms of UCD. Patients receiving DIVALPROEX who develop symptoms of unexplained hyperammonemic encephalopathy should receive prompt treatment (including discontinuation of DIVALPROEX) and be evaluated for underlying urea cycle disorders. See (CONTRAINDICATIONS) and (WARNINGS AND PRECAUTIONS, **Endocrine and Metabolism**, **Hyperammonemia and Hyperammonemia and Encephalopathy Associated with Concomitant Topiramate Use**).

### **Hyperammonemia**

Hyperammonemia has been reported in association with divalproex sodium and may be present despite normal liver function tests. In patients who develop unexplained lethargy and vomiting or changes in mental status, hyperammonemic encephalopathy should be considered as a possible cause and serum ammonia level should be measured. Hyperammonemia should also be considered in patients who present with hypothermia. See (WARNINGS AND PRECAUTIONS, **Endocrine and Metabolism**, **Hypothermia**). If serum ammonia is increased, DIVALPROEX should be discontinued. Appropriate interventions for treatment of hyperammonemia should be initiated, and such patients should undergo investigation for underlying urea cycle disorders. See (CONTRAINDICATIONS) and (WARNINGS AND

**PRECAUTIONS, Endocrine and Metabolism, Urea Cycle Disorders and Hyperammonemia and Encephalopathy Associated with Concomitant Topiramate Use).**

Asymptomatic elevations of serum ammonia are more common and, when present, require close monitoring of serum ammonia levels. If the elevation persists, discontinuation of DIVALPROEX should be considered.

**Hyperammonemia and Encephalopathy Associated with Concomitant Topiramate Use**

Concomitant administration of topiramate and divalproex sodium has been associated with hyperammonemia with or without encephalopathy in patients who have tolerated either drug alone. Clinical symptoms of hyperammonemic encephalopathy often include acute alterations in level of consciousness and/or cognitive function with lethargy or vomiting. Hypothermia can also be a manifestation of hyperammonemia. See (**WARNINGS AND PRECAUTIONS, Endocrine and Metabolism, Hypothermia**). In most cases, symptoms and signs abated with discontinuation of either drug. This adverse event is not due to a pharmacokinetic interaction.

It is not known if topiramate monotherapy is associated with hyperammonemia.

**Patients with inborn errors of metabolism or reduced hepatic mitochondrial activity may be at an increased risk for hyperammonemia with or without encephalopathy. Although not studied, an interaction of topiramate and divalproex sodium may exacerbate existing defects or unmask deficiencies in susceptible persons. See (CONTRAINDICATIONS) and (WARNINGS AND PRECAUTIONS, Endocrine and Metabolism, Urea Cycle Disorders and Hyperammonemia).**

**Hypothermia**

Hypothermia, defined as an unintentional drop in core body temperature to < 35°C (95°F), has been reported in association with divalproex sodium therapy both in conjunction with and in the absence of hyperammonemia. This adverse reaction can also occur in patients using concomitant topiramate with divalproex sodium after starting topiramate treatment or after increasing the daily dose of topiramate. See (**WARNINGS AND PRECAUTIONS, Endocrine and Metabolism, Hyperammonemia, Hyperammonemia and Encephalopathy Associated with Concomitant Topiramate Use and DRUG INTERACTIONS, Drug-Drug Interactions, Table 2**). Hypothermia may be manifested by a variety of clinical abnormalities including lethargy, confusion, coma and significant alterations in other major organ systems such as the cardiovascular and respiratory systems. Clinical management and assessment should include examination of blood ammonia levels. Consideration should be given to stopping DIVALPROEX in patients who develop hypothermia. See (**WARNINGS AND PRECAUTIONS, Endocrine and Metabolism, Hyperammonemia and Hyperammonemia and Encephalopathy Associated with Concomitant Topiramate Use**).

## **Hematologic**

### **Thrombocytopenia**

Because of reports of thrombocytopenia, inhibition of the second phase of platelet aggregation, and abnormal coagulation parameters (e.g., low fibrinogen), platelet counts and coagulation tests are recommended before initiating therapy and at periodic intervals. It is recommended that patients receiving DIVALPROEX be monitored for platelet count and coagulation parameters prior to planned surgery. Clinical evidence of hemorrhage, bruising or a disorder of hemostasis/coagulation is an indication for reduction of the dosage or withdrawal of therapy. See also (**WARNINGS AND PRECAUTIONS, Hematologic, Dose-related Adverse Reactions: Thrombocytopenia**).

### **Dose-related Adverse Reactions: Thrombocytopenia**

The frequency of adverse effects (particularly elevated liver enzymes and thrombocytopenia) may be dose-related. In a clinical trial of divalproex sodium as monotherapy in patients with epilepsy, 34/126 patients (27%) receiving approximately 50 mg/kg/day on average, had at least one value of platelets  $\leq 75 \times 10^9/L$ . Approximately half of these patients had treatment discontinued with return of platelet counts to normal. In the remaining patients, platelet counts normalized with continued treatment. In this study, the probability of thrombocytopenia appeared to increase significantly at total valproate concentrations of  $\geq 110$  mcg/mL (females) or  $\geq 135$  mcg/mL (males). The therapeutic benefit which may accompany the higher doses should therefore be weighed against the possibility of a greater incidence of adverse events.

In addition, the findings from a crossover clinical trial conducted with divalproex sodium extended-release tablets, in 44 epilepsy patients, indicate that the frequency of treatment-emergent mild thrombocytopenia (platelet count between  $100$  to  $150 \times 10^9/L$ ) was significantly higher after 12 weeks of treatment with divalproex sodium extended-release tablets than after 12 weeks of treatment with divalproex sodium enteric-coated tablets (7 versus 3 low counts, respectively).

## **Hepatic/Biliary/Pancreatic**

### **Serious or Fatal Hepatotoxicity**

Hepatic failure resulting in fatalities has occurred in patients receiving divalproex sodium and its derivatives. These incidences usually occurred during the first 6 months of treatment with divalproex sodium. Caution should be observed when administering DIVALPROEX to patients with a prior history of hepatic disease. Patients on multiple anticonvulsants, children, those with congenital metabolic disorders, those with severe seizure disorders accompanied by mental retardation, and those with organic brain disease may be at particular risk.



Experience has indicated that children under the age of 2 years are at a considerably increased risk of developing fatal hepatotoxicity, especially those on multiple anticonvulsants, those with congenital metabolic disorders, those with severe seizure disorders accompanied by mental retardation, and those with organic brain disease. The risk in this age group decreased considerably in patients receiving divalproex sodium as monotherapy. Similarly, patients aged 3 to 10 years were at somewhat greater risk if they received multiple anticonvulsants than those who received only divalproex sodium. Above the age of 2 years, experience in epilepsy has indicated that the incidence of fatal hepatotoxicity decreases considerably in progressively older patients. No deaths have been reported in patients over 10 years of age who received divalproex sodium alone.

If DIVALPROEX is to be used for the control of seizures in children 2 years old or younger, it should be used with extreme caution and as a sole agent. The benefits of therapy should be weighed against the risks. See [**WARNINGS AND PRECAUTIONS, Special Populations, Pediatrics (< 18 years of age)**].

Serious or fatal hepatotoxicity may be preceded by nonspecific symptoms such as, malaise, weakness, lethargy, facial edema, anorexia, and vomiting. In patients with epilepsy, a loss of seizure control may also occur. Patients should be monitored closely for appearance of these symptoms. Patients and parents should be instructed to report such symptoms. Because of the nonspecific nature of some of the early signs, hepatotoxicity should be suspected in patients who become unwell, other than through obvious cause, while taking DIVALPROEX.

Liver function tests should be performed prior to therapy and at frequent intervals thereafter especially during the first 6 months. However, physicians should not rely totally on serum biochemistry since these tests may not be abnormal in all instances, but should also consider the results of careful interim medical history and physical examination.

In high-risk patients, it might also be useful to monitor serum fibrinogen and albumin for decreases in concentration and serum ammonia for increases in concentration. If changes occur, DIVALPROEX should be discontinued. Dosage should be titrated to and maintained at the lowest dose consistent with optimal seizure control.

The drug should be discontinued immediately in the presence of significant hepatic dysfunction, suspected or apparent. In some cases, hepatic dysfunction has progressed in spite of discontinuation of drug. The frequency of adverse hepatic effects (particularly elevated liver enzymes) may increase with increasing dose. The therapeutic benefit which may accompany the higher doses should therefore be weighed against the possibility of a greater incidence of adverse effects. (See (**CONTRAINDICATIONS**)).

## **Pancreatitis**

**Cases of life-threatening pancreatitis have been reported in both children and adults receiving divalproex sodium. Some of the cases have been described as hemorrhagic with a rapid progression from initial symptoms to death. Some cases have occurred shortly after initial use as well as after several years of use. The rate based upon the reported cases exceeds that expected in the general population and there have been cases in which pancreatitis recurred after rechallenge with divalproex sodium. In clinical trials, there were 2 cases of pancreatitis without alternative etiology in 2,416 patients, representing 1,044 patient-years experience. Patients and guardians should be warned that abdominal pain, nausea, vomiting, and/or anorexia can be symptoms of pancreatitis that require prompt medical evaluation. If pancreatitis is diagnosed, DIVALPROEX should ordinarily be discontinued. Alternative treatment for the underlying medical condition should be initiated as clinically indicated.**

## **Neurologic**

### **Driving and Hazardous Occupations**

DIVALPROEX may produce central nervous system (CNS) depression, especially when combined with another CNS depressant, such as alcohol. Therefore, patients should be advised not to engage in hazardous occupations, such as driving a car or operating dangerous machinery, until it is known that they do not become drowsy from the drug.

## **Psychiatric**

### **Suicidal Behaviour and Ideation**

Suicidal ideation and behaviour have been reported in patients treated with antiepileptic agents in several indications.

All patients treated with antiepileptic drugs (AEDs), irrespective of indication, should be monitored for signs of suicidal ideation and behaviour and appropriate treatment should be considered. Patients (and caregivers of patients) should be advised to seek medical advice should signs of suicidal ideation or behaviour emerge.

An FDA meta-analysis of randomized placebo controlled trials, in which AEDs were used for various indications, has shown a small increased risk of suicidal ideation and behaviour in patients treated with these drugs. The mechanism of this risk is not known.

There were 43,892 patients treated in the placebo controlled clinical trials that were included in the meta-analysis. Approximately 75% of patients in these clinical trials were treated for indications other than epilepsy and, for the majority of non-epilepsy indication the treatment (AED or placebo) was administered as monotherapy. Patients with epilepsy represented approximately 25% of the total number of patients treated in the placebo controlled clinical trials and, for the majority of epilepsy patients, treatment (AED or placebo) was administered as

adjunct to other antiepileptic agents (i.e., patients in both treatment arms were being treated with one or more AED). Therefore, the small increased risk of suicidal ideation and behaviour reported from the meta-analysis (0.43% for patients on AEDs compared to 0.24% for patients on placebo) is based largely on patients that received monotherapy treatment (AED or placebo) for non-epilepsy indications. The study design does not allow an estimation of the risk of suicidal ideation and behaviour for patients with epilepsy that are taking AEDs, due both to this population being the minority in the study, and the drug-placebo comparison in this population being confounded by the presence of adjunct AED treatment in both arms.

## **Renal**

### **Renal Impairment**

Renal impairment is associated with an increase in the unbound fraction of valproate. In several studies, the unbound fraction of valproate in plasma from renally impaired patients was approximately double that for subjects with normal renal function. Accordingly, monitoring of total concentrations in patients with renal impairment may be misleading since free concentrations may be substantially elevated whereas total concentrations may appear to be normal. Hemodialysis in renally impaired patients may remove up to 20% of the circulating valproate.

## **Sensitivity/Resistance**

### **Multi-organ Hypersensitivity Reaction**

Multi-organ hypersensitivity reactions have been rarely reported in close temporal association to the initiation of divalproex sodium therapy in adult and pediatric patients (median time to detection 21 days; range 1 to 40). Although there have been a limited number of reports, many of these cases resulted in hospitalization and at least one death has been reported. Signs and symptoms of this disorder were diverse; however, patients typically, although not exclusively, presented with fever and rash associated with other organ system involvement. Other associated manifestations may include lymphadenopathy, hepatitis, liver function test abnormalities, hematological abnormalities (e.g., eosinophilia, thrombocytopenia, neutropenia), pruritus, nephritis, oliguria, hepato-renal syndrome, arthralgia, and asthenia. Because the disorder is variable in its expression, other organ system symptoms and signs, not noted here may occur. If this reaction is suspected, DIVALPROEX should be discontinued and an alternative treatment started. Although the existence of cross sensitivity with other drugs that produce this syndrome is unclear, the experience amongst drugs associated with multi-organ hypersensitivity would indicate this to be a possibility.

## **Sexual Function/Reproduction**

### **Fertility**

The effect of divalproex sodium on testicular development and on sperm production and fertility in humans is unknown. See (**TOXICOLOGY, Reproduction and Teratology, Fertility**) for results in animal studies.

### **Skin**

#### **Serious Skin Reactions**

The dose of lamotrigine should be reduced when co-administered with DIVALPROEX. Serious skin reactions (such as Stevens-Johnson syndrome and Toxic Epidermal Necrolysis) have been reported with concomitant lamotrigine and divalproex sodium administration (see Lamotrigine Product Monograph for details on lamotrigine dosing with concomitant divalproex sodium administration).

### **Special Populations**

#### **Pregnant Women**

According to published and unpublished reports in the medical literature, divalproex sodium may produce teratogenic effects, such as neural tube defects (e.g., spina bifida) in the offspring of human females receiving the drug during pregnancy. There are data that suggest an increased incidence of congenital malformations associated with the use of divalproex sodium during pregnancy when compared with some other AEDs. Therefore, DIVALPROEX should be considered for women of childbearing potential only after the risks have been thoroughly discussed with the patient and weighed against the potential benefits of treatment.

Multiple reports in the clinical literature indicate an association between the use of AEDs and an elevated incidence of birth defects in children born to epileptic women taking such medication during pregnancy. The incidence of congenital malformations in the general population is regarded to be approximately 2%; in children of treated epileptic women, this incidence may be increased 2- to 3-fold. The increase is largely due to specific defects such as congenital malformations of the heart, cleft lip and/or palate, craniofacial abnormalities and neural tube defects. Nevertheless, the great majority of mothers receiving antiepileptic medications deliver normal infants.

#### **Neural Tube Defects**

**The data described below were gained almost exclusively from women who received divalproex sodium to treat epilepsy. The incidence of neural tube defects in the fetus is increased in mothers receiving divalproex sodium during the first trimester of pregnancy. Based upon a single report, it was estimated that the risk of divalproex sodium-exposed women having children with spina bifida is approximately 1 to 2%.**

## Congenital Anomalies

Other congenital anomalies (e.g., craniofacial defects, cardiovascular malformations and anomalies involving various body systems), compatible and incompatible with life, have been reported. Sufficient data to determine the incidence of these congenital anomalies are not available.

The higher incidence of congenital anomalies in AED-treated women with seizure disorders cannot be regarded as a cause and effect relationship. There are intrinsic methodological problems in obtaining adequate data on drug teratogenicity in humans; genetic factors or the epileptic condition itself, may be more important than the drug therapy in contributing to congenital anomalies.

## Developmental Delay, Autism and/or Autism Spectrum Disorder

There have been post-marketing reports of developmental delay, autism and/or autism spectrum disorder in the offspring of women exposed to divalproex sodium during pregnancy. Cognitive testing at year 2 or 3 in a 6-year, prospective study of long-term cognitive development in 309 children (born to 303 mothers) exposed in utero to antiepileptic drug monotherapy showed that children exposed to valproic acid had significantly lower IQs (92; 95% confidence interval, 88 to 97) than did children exposed to other antiepileptic drugs (carbamazepine, 98; lamotrigine, 101; phenytoin, 99), whereas IQ scores did not differ significantly among children exposed to the other 3 antiepileptic drugs. There was a significant correlation between the dose of valproic acid during pregnancy and the child's IQ.

Patients taking divalproex sodium may develop clotting abnormalities. A patient who had low fibrinogen when taking multiple anticonvulsants, including divalproex sodium, gave birth to an infant with afibrinogenemia who subsequently died of hemorrhage. If DIVALPROEX is used in pregnancy, the clotting parameters should be monitored carefully.

Hepatic failure, resulting in the death of a newborn and of an infant has been reported following the use of valproate during pregnancy.

AEDs should not be abruptly discontinued in patients to whom the drug is administered to prevent major seizures, because of the strong possibility of precipitating status epilepticus with attendant hypoxia and risks to both the mother and the unborn child. With regard to drugs given for minor seizures, the risks of discontinuing medication prior to or during pregnancy should be weighed against the risk of congenital defects in the particular case and with the particular family history. In individual cases where the severity and frequency of the seizure disorder are such that the removal of medication does not pose a serious threat to the patient, discontinuation of the drug may be considered prior to and during pregnancy, although it cannot be said with any confidence that even minor seizures do not pose some hazard to the developing embryo or fetus.

In summary, current best practice guidelines should be considered in order to provide the optimal counsel to patients regarding the teratogenic risks associated with DIVALPROEX.

Epileptic women of childbearing age should be encouraged to seek the counsel of their physician and should report the onset of pregnancy promptly to him. Where the necessity for continued use of antiepileptic medication is in doubt, appropriate consultation is indicated.

Risk-benefit must be carefully considered when treating or counselling women of childbearing age for bipolar disorder.

If DIVALPROEX is used during pregnancy, or if the patient becomes pregnant while taking this drug, the patient should be made aware of the potential hazard to the fetus.

Tests to detect neural tube and other defects using current accepted procedures should be considered a part of routine prenatal care in childbearing women receiving DIVALPROEX.

### Teratogenicity in Animals

Animal studies have demonstrated valproic acid induced teratogenicity, see (**TOXICOLOGY, Reproduction and Teratology**), and studies in human females have demonstrated placental transfer of the drug. Increased frequencies of malformations, as well as intrauterine growth retardation and death, have been observed in mice, rats, rabbits, and monkeys following prenatal exposure to valproate. Malformations of the skeletal system are the most common structural abnormalities produced in experimental animals, but neural tube closure defects have been seen in mice exposed to maternal plasma valproate concentrations exceeding 230 mcg/mL (2.3 times the upper limit of the human therapeutic range for epilepsy) during susceptible periods of embryonic development.

Administration of an oral dose of 200 mg/kg/day or greater (50% of the maximum human daily dose or greater on a mg/m<sup>2</sup> basis) to pregnant rats during organogenesis produced malformations (skeletal, cardiac and urogenital) and growth retardation in the offspring. These doses resulted in peak maternal plasma valproate levels of approximately 340 mcg/mL or greater (3.4 times the upper limit of the human therapeutic range for epilepsy or greater). Behavioural deficits have been reported in the offspring of rats given a dose of 200 mg/kg/day throughout most of pregnancy.

An oral dose of 350 mg/kg/day (approximately 2 times the maximum human daily dose on a mg/m<sup>2</sup> basis) produced skeletal and visceral malformations in rabbits exposed during organogenesis. Skeletal malformations, growth retardation, and death were observed in rhesus monkeys following administration of an oral dose of 200 mg/kg/day (equal to the maximum human daily dose on a mg/m<sup>2</sup> basis) during organogenesis. This dose resulted in peak maternal plasma valproate levels of approximately 280 mcg/mL (2.8 times the upper limit of the human therapeutic range for epilepsy).

## **Nursing Women**

Divalproex sodium is excreted in breast milk. Concentrations in breast milk have been reported to be 1 to 10% of serum concentrations. As a general rule, nursing should not be undertaken while a patient is receiving DIVALPROEX. It is not known what effect this may have on a nursing infant.

## **Pediatrics (< 18 years of age)**

Experience has indicated that children under the age of 2 years are at a considerably increased risk of developing fatal hepatotoxicity, especially those with the aforementioned conditions. See (**WARNINGS AND PRECAUTIONS, Hepatic/Biliary/Pancreatic, Serious or Fatal Hepatotoxicity**). When DIVAPROEX is used in this patient group, it should be used with extreme caution and as a sole agent. The benefits of therapy should be weighed against the risks.

Above the age of 2 years, experience in epilepsy has indicated that the incidence of fatal hepatotoxicity decreases considerably in progressively older patient groups.

Younger children, especially those receiving enzyme-inducing drugs, will require larger maintenance doses to attain targeted total and unbound valproate concentrations. The variability in free fraction limits the clinical usefulness of monitoring total serum valproate concentrations. Interpretation of valproate concentrations in children should include consideration of factors that affect hepatic metabolism and protein binding.

The safety and effectiveness of divalproex sodium for the treatment of acute mania has not been studied in individuals below the age of 18 years.

## **Geriatrics (≥ 65 years of age)**

Alterations in the kinetics of unbound valproate in the elderly indicate that the initial dosage should be reduced in this population. See (**DOSAGE AND ADMINISTRATION**) and (**ACTION AND CLINICAL PHARMACOLOGY, Special Populations and Conditions, Geriatrics**).

The safety and efficacy of divalproex sodium in elderly patients with epilepsy and mania has not been evaluated in clinical trials. Caution should thus be exercised in dose selection for an elderly patient, recognizing the more frequent hepatic and renal dysfunctions, and limited experience with divalproex sodium in this population.

A study of elderly patients revealed valproate-related somnolence and discontinuation of divalproex sodium for this adverse event. See (**WARNINGS AND PRECAUTIONS, Special Populations, Geriatrics, Somnolence in the Elderly**). The starting dose should be reduced in elderly patients, and dosage reductions or discontinuation should be considered in patients with excessive somnolence. See (**DOSAGE AND ADMINISTRATION**).

### Somnolence in the elderly

In a group of elderly patients (mean age 83 years old, n = 172), divalproex sodium doses were increased by 125 mg/day to a target dose of 20 mg/kg/day. Compared to placebo a significantly higher number of valproate-treated patients had somnolence, and although not statistically significant, a higher number of valproate-treated patients experienced dehydration. Discontinuations for somnolence were also significantly higher in valproate-treated patients compared to placebo. In approximately one-half of the patients with somnolence, there was also associated reduced nutritional intake and weight loss. In elderly patients, dosage should be increased more slowly and with regular monitoring for fluid intake, dehydration, somnolence, urinary tract infection and other adverse events. Dose reductions or discontinuation of DIVALPROEX should be considered in patients with decreased food or fluid intake and in patients with excessive somnolence. See **(DOSAGE AND ADMINISTRATION)**.

### **Monitoring and Laboratory Tests**

Since DIVALPROEX may interact with concurrently administered drugs which are capable of enzyme induction, periodic plasma concentration determinations of valproate and concomitant drugs are recommended during the early course of therapy and whenever enzyme-inducing drugs are introduced or withdrawn. See **(DRUG INTERACTIONS)**.

### **Monitoring Valproate Concentrations**

Protein binding of valproate is reduced in the elderly, in patients with renal impairment, and in the presence of other drugs (e.g., acetylsalicylic acid). Accordingly, measurements of plasma levels of valproate may be misleading in these patients, as actual drug exposure may be higher than measured values. See **(WARNINGS AND PRECAUTIONS, Hepatic/Biliary/Pancreatic), (WARNINGS AND PRECAUTIONS, Endocrine and Metabolism, Hyperammonemia), (WARNINGS AND PRECAUTIONS, Hematologic, Thrombocytopenia) and (DRUG INTERACTIONS, Drug-Drug Interactions, Table 2).**

## **ADVERSE REACTIONS**

### **Adverse Drug Reaction Overview**

#### **Epilepsy**

The most commonly reported adverse reactions are nausea, vomiting and indigestion. Since divalproex sodium enteric-coated tablets has usually been used with other antiepileptics, it is not possible in most cases to determine whether the adverse reactions mentioned in this section are due to divalproex sodium alone or to the combination of drugs.

Adverse events that have been reported with divalproex sodium from epilepsy trials, spontaneous reports, and other sources are listed below by system organ class.



Blood and Lymphatic System Disorders:	Thrombocytopenia and inhibition of the secondary phase of platelet aggregation may be reflected in altered bleeding time, petechiae, bruising, hematoma formation, epistaxis, and frank hemorrhage. See ( <b>WARNINGS AND PRECAUTIONS, <u>Hematologic, Thrombocytopenia</u></b> ). Relative lymphocytosis, macrocytosis and hypofibrinogenemia have been noted. Leukopenia and eosinophilia have also been reported. Anemia, including macrocytic with or without folate deficiency, aplastic anemia, pancytopenia, bone marrow suppression, agranulocytosis and acute intermittent porphyria have been reported.
Cardiac Disorders:	Bradycardia has been reported.
Ear and Labyrinth Disorders:	Hearing loss, either reversible or irreversible, has been reported; however, a cause and effect relationship has not been established. Ear pain has also been reported.
Gastrointestinal Disorders:	Nausea, vomiting and indigestion are the most commonly reported side effects at the initiation of therapy. These effects are usually transient and rarely require discontinuation of therapy. Diarrhea, abdominal cramps, constipation and parotid gland swelling have also been reported.  There have been reports of acute pancreatitis, including rare fatal cases, occurring in patients receiving divalproex sodium. See ( <b>WARNINGS AND PRECAUTIONS, <u>Hepatic/Biliary/Pancreatic, Pancreatitis</u></b> ).
General Disorders and Administration Site Conditions:	Edema of extremities, fever and hypothermia has been reported.
Hepatobiliary Disorders:	Minor elevations of transaminases [e.g., serum glutamic-oxaloacetic transaminase (SGOT) and serum glutamic-pyruvic transaminase (SGPT)] and lactate dehydrogenase (LDH) are frequent and appear to be dose-related. Occasionally, laboratory tests also show increases in serum bilirubin and abnormal changes in other liver function tests. These results may reflect potentially serious hepatotoxicity. See ( <b>WARNINGS AND PRECAUTIONS, <u>Hepatic/Biliary/Pancreatic, Serious or Fatal Hepatotoxicity</u></b> ).
Immune System Disorders:	Allergic reaction, anaphylaxis has been reported.

Infections and Infestations:	Pneumonia and otitis media have been reported.
Investigations:	Abnormal thyroid function tests have been reported. See <b>(DRUG INTERACTIONS, <u>Drug-Laboratory Interactions</u>)</b> .
Metabolism and Nutrition Disorders:	<p>Hyperammonemia, see <b>(WARNINGS AND PRECAUTIONS, <u>Endocrine and Metabolism, Hyperammonemia</u>)</b>, hyponatremia and inappropriate antidiuretic hormone (ADH) secretion. There have been rare reports of Fanconi syndrome occurring primarily in children. Decreased carnitine concentrations have been reported although the clinical relevance is undetermined. Hyperglycinemia (elevated plasma glycine concentration) has been reported and associated with a fatal outcome in a patient with pre-existing non-ketotic hyperglycinemia.</p> <p>Anorexia with some weight loss and increased appetite with some weight gain have also been reported.</p>
Musculoskeletal and Connective Tissue Disorders:	<p>Weakness and bone pain have been reported.</p> <p>Reports have been received of decreased bone mass, potentially leading to osteoporosis and osteopenia, during long-term therapy with some anticonvulsant medications, including divalproex sodium. Some studies have indicated that supplemental calcium and vitamin D may be of benefit to patients who are on chronic divalproex sodium therapy.</p> <p>A lupus erythematosus-like syndrome has been reported rarely.</p>
Nervous System Disorders:	<p>Sedative effects have been noted in patients receiving divalproex sodium alone but occur most often in patients on combination therapy. Sedation usually disappears upon reduction of other antiepileptic medication.</p> <p>Hallucination, ataxia, headache, nystagmus, diplopia, asterixis, "spots before the eyes", tremor (may be dose-related), confusion, dysarthria, dizziness, hypesthesia, vertigo, incoordination and parkinsonism have been noted. Rare cases of coma have been reported in patients receiving divalproex sodium alone or in conjunction with phenobarbital.</p> <p>Encephalopathy, with or without fever or hyperammonemia, has been reported without evidence of hepatic dysfunction or inappropriate valproate plasma levels. Most patients recovered,</p>

with noted improvement of symptoms, upon discontinuation of the drug.

Reversible cerebral atrophy and dementia have been reported in association with divalproex sodium.

Psychiatric Disorders: Emotional upset, depression, psychosis, aggression, hyperactivity, hostility and behavioural deterioration have been reported.

Renal and Urinary Disorders: Enuresis and urinary tract infection.

Reproductive System and Breast Disorders: There have been reports of irregular menses, secondary amenorrhea, breast enlargement and galactorrhea in patients receiving divalproex sodium.

There have been rare spontaneous reports of polycystic ovary disease. A cause and effect relationship has not been established.

Respiratory, Thoracic and Mediastinal Disorders: Increased cough has been reported.

Skin and Subcutaneous Tissue Disorders: Transient increases in hair loss have been observed. Skin rash, photosensitivity, generalized pruritus, erythema multiforme, Stevens-Johnson syndrome (SJS), and petechiae have rarely been noted.

Rare cases of Toxic Epidermal Necrolysis (TEN) have been reported including a fatal case of a 6 month old infant taking divalproex sodium and several other concomitant medications. An additional case of Toxic Epidermal Necrolysis resulting in death was reported in a 35 year old patient with AIDS taking several concomitant medications and with a history of multiple cutaneous drug reactions.

Serious skin reactions have been reported with concomitant administration of lamotrigine and divalproex sodium. See **(DRUG INTERACTIONS, Drug-Drug Interactions, Table 2)**.

Cutaneous vasculitis has also been reported.

## **Divalproex Sodium Extended-Release Tablets versus Divalproex Sodium Enteric-Coated Tablets**

A 24 week cross-over study compared the safety and efficacy of divalproex sodium extended-release tablets administered once daily to that of equal doses of divalproex sodium enteric-coated tablets (administered twice daily or three times daily) in the treatment of adolescent and adult epileptic patients with generalized seizures (n=44). Two adverse events occurred in significantly more patients on divalproex sodium extended-release tablets than on divalproex sodium enteric-coated tablets: asthenia (15.9% versus 6.8% respectively) and treatment-emergent mild thrombocytopenia (16.2% versus 6.8%, respectively).

### **Bipolar Disorder**

The incidence of adverse events has been ascertained based on data from two short-term (21 day) placebo-controlled clinical trials of divalproex sodium in the treatment of acute mania, and from 2 long-term (up to 3 years) retrospective open trials.

#### Most Commonly Observed

During the short-term placebo-controlled trials, the six most commonly reported adverse events in patients (n = 89) exposed to divalproex sodium were nausea (22%), headache (21%), somnolence (19%), pain (15%), vomiting (12%), and dizziness (12%).

In the long-term retrospective trials (634 patients exposed to divalproex sodium), the six most commonly reported adverse events were somnolence (31%), tremor (29%), headache (24%), asthenia (23%), diarrhea (22%), and nausea (20%).

#### Associated With Discontinuation of Treatment

In the placebo-controlled trials, adverse events which resulted in divalproex sodium discontinuation in at least one percent of patients were nausea (4%), abdominal pain (3%), somnolence (2%), and rash (2%).

In the long-term retrospective trials, adverse events which resulted in divalproex sodium discontinuation in at least one percent of patients were alopecia (2.4%), somnolence (1.9%), nausea (1.7%), and tremor (1.4%). The time to onset of these events was generally within the first two months of initial exposure to divalproex sodium. A notable exception was alopecia, which was first experienced after 3 to 6 months of exposure by 8 of the 15 patients who discontinued divalproex sodium in response to the event.

#### Controlled Trials

Table 1 summarizes those treatment-emergent adverse events reported for patients in the placebo-controlled trials when the incidence rate in the divalproex sodium group was at least 5%.

(Maximum treatment duration was 21 days; maximum dose in 83% of patients was between 1,000 to 2,500 mg per day).

System Organ Class/Event	Divalproex Sodium N = 89 (%)	placebo N = 97 (%)
<b>Gastrointestinal Disorders</b>		
Nausea	22.5	15.5
Vomiting	12.4*	3.1
Diarrhea	10.1	13.4
Abdominal Pain	9	8.2
Dyspepsia	9	8.2
Constipation	7.9	8.2
<b>General Disorders and Administration Site Conditions</b>		
Pain	14.6	15.5
Asthenia	10.1	7.2
<b>Injury, Poisoning and Procedural Complications</b>		
Accidental Injury	11.2	5.2
<b>Musculoskeletal and Connective Tissue Disorders</b>		
Back Pain	5.6	6.2
<b>Nervous System Disorders</b>		
Headache	21.3	30.9
Somnolence	19.1	12.4
Dizziness	12.4	4.1
Tremor	5.6	6.2
<b>Respiratory, Thoracic and Mediastinal Disorders</b>		
Pharyngitis	6.7	9.3
<b>Skin and Subcutaneous Tissue Disorders</b>		
Rash	5.6	3.1

\* Statistically significant at P < 0.05 level.

The following adverse events not listed above were reported by at least 1%, but less than 5%, of the 89 patients from the two placebo-controlled clinical trials of divalproex sodium tablets.

Cardiac Disorders: Palpitations, tachycardia.

Congenital, Familial and Genetic Disorders: Vascular anomaly.

Ear and Labyrinth Disorders: Deafness, ear disorder, ear pain, tinnitus, vertigo.

Eye Disorders: Abnormal vision, amblyopia, conjunctivitis, diplopia, dry eyes, eye pain.

Gastrointestinal Disorders: Fecal incontinence, flatulence, gastroenteritis, glossitis.

General Disorders and Administration Site: Abnormal gait, chest pain, chills, chills and fever, cyst, edema, fever, furunculosis, periodontal abscess, peripheral

Conditions:	edema.
Infections and Infestations:	Infection, rhinitis.
Metabolism and Nutrition Disorders:	Anorexia.
Musculoskeletal and Connective Tissue Disorders:	Arthralgia, arthrosis, leg cramps, neck pain, neck rigidity twitching.
Nervous System Disorders:	Ataxia, dysarthria, hypertonia, hypokinesia, paresthesia, reflexes increased, tardive dyskinesia.
Psychiatric Disorders:	Abnormal dreams, agitation, catatonic reaction, confusion, depression, hallucinations, insomnia, thinking abnormalities.
Renal and Urinary Disorders:	Dysurea, urinary incontinence.
Reproductive System and Breast Disorders:	Dysmenorrhea.
Respiratory, Thoracic and Mediastinal Disorders:	Dyspnea.
Skin and Subcutaneous Tissue Disorders:	Alopecia, discoid lupus erythematosus, dry skin, maculopapular rash, seborrhea.
Vascular Disorders:	Ecchymosis, hypertension, hypotension, postural hypotension, vasodilation.

### **Adverse Events in Elderly Patients**

In elderly patients (above 65 years of age), there were more frequent reports of accidental injury, infection, pain, and to a lesser degree, somnolence and tremor, when compared to patients 18 to 65 years of age. Somnolence and tremor tended to be associated with the discontinuation of divalproex sodium.

## DRUG INTERACTIONS

### Serious Drug Interactions

- Rare cases of coma have been reported in patients receiving divalproex sodium alone or in conjunction with phenobarbital. See (**Drug-Drug Interactions, Table 2**).
- Serious skin reactions (such as Stevens-Johnson syndrome and Toxic Epidermal Necrolysis) have been reported with concomitant lamotrigine and divalproex sodium administration. See (**Drug-Drug Interactions, Table 2**).

### Overview

Divalproex sodium has been found to be a weak inhibitor of some P450 isozymes, epoxide hydrolase, and glucuronyl transferases.

Drugs that affect the level of expression of hepatic enzymes, particularly those that elevate levels of glucuronyl transferases, may increase the clearance of valproate. For example, phenytoin, carbamazepine, and phenobarbital (or primidone) can double the clearance of valproate. Thus, patients on divalproex sodium monotherapy will generally have longer half-lives and higher concentrations than patients receiving polytherapy with antiepilepsy drugs.

In contrast, drugs that are inhibitors of cytochrome P450 isozymes, such as antidepressants, may be expected to have little effect on divalproex sodium clearance because cytochrome P450 microsomal mediated oxidation is a relatively minor secondary metabolic pathway compared to glucuronidation and beta-oxidation.

The concomitant administration of DIVALPROEX with drugs that exhibit extensive protein binding (e.g., acetylsalicylic acid, carbamazepine, dicumarol, warfarin, tolbutamide, and phenytoin) may result in alteration of serum drug levels.

Since DIVALPROEX may interact with concurrently administered drugs which are capable of enzyme induction, periodic plasma concentration determinations of valproate and concomitant drugs are recommended during the early course of therapy and whenever enzyme-inducing drugs are introduced or withdrawn.

### Drug-Drug Interactions

**Table 2** provides information about the potential influence of several commonly prescribed medications on valproate pharmacokinetics as well as the potential influence of valproate on the pharmacokinetics and pharmacodynamics of several commonly prescribed medications. The list is not exhaustive nor could it be, since new interactions are continuously being reported. Please note that drugs may be listed under specific name, family or pharmacologic class. Reading the entire section is recommended.

Concomitant Drug	Ref	Effect	Clinical comment
Acetaminophen	CT	↔ acetaminophen	Divalproex sodium had no effect on any of the pharmacokinetic parameters of acetaminophen when it was concurrently administered to three epileptic patients.
Acetylsalicylic Acid	CT	↑ valproate	A study involving the co-administration of acetylsalicylic acid at antipyretic doses (11 to 16 mg/kg) with divalproex sodium to pediatric patients (n = 6) revealed a decrease in protein binding and an inhibition of metabolism of valproate. Valproate free fraction was increased 4-fold in the presence of acetylsalicylic acid compared to divalproex sodium alone. The beta-oxidation pathway consisting of 2-E-valproic acid, 3-OH-valproic acid, and 3-keto valproic acid was decreased from 25% of total metabolites excreted on divalproex sodium alone to 8.3% in the presence of acetylsalicylic acid. Caution should be observed when DIVALPROEX is administered with drugs affecting coagulation, (e.g., acetylsalicylic acid and warfarin). See ( <b>ADVERSE REACTIONS</b> ).
Alcohol	T	No pharmacokinetic (PK) interaction	Divalproex sodium may potentiate the CNS depressant action of alcohol.
Amitriptyline / Nortriptyline	CT	In general: ↓ amitriptyline ↓ nortriptyline  Rarely: ↑ amitriptyline ↑ nortriptyline	Administration of a single oral 50 mg dose of amitriptyline to 15 normal volunteers (10 males and 5 females) who received divalproex sodium (500 mg twice daily) resulted in a 21% decrease in plasma clearance of amitriptyline and a 34% decrease in the net clearance of nortriptyline.  Rare post-marketing reports of concurrent use of divalproex sodium and amitriptyline resulting in an increased amitriptyline and nortriptyline levels have been received. Concurrent use of divalproex sodium and amitriptyline has rarely been associated with toxicity. Monitoring of amitriptyline levels should be considered for patients taking DIVALPROEX concomitantly with amitriptyline. Consideration should be given to lowering the dose of amitriptyline/nortriptyline in the presence of DIVALPROEX.
Antacids	CT	↔ valproate	A study involving the co-administration of divalproex sodium 500 mg with commonly administered antacids (Maalox®, Trisogel and Titalac™ - 160 milliequivalent doses) did not reveal any effect on the extent of absorption of valproate.
Other -Antipsychotics, Monoamine Oxidase Inhibitors (MAOIs) and Tricyclic Antidepressants			In addition to enhancing CNS depression when used concurrently with divalproex sodium, antipsychotics, tricyclic antidepressants and MAOIs may lower the seizure threshold. Dosage adjustments may be necessary to control seizures.
Benzodiazepines			Divalproex sodium may decrease oxidative liver metabolism of some benzodiazepines, resulting in increased serum concentrations. See ( <b>Table 1. Diazepam and Lorazepam</b> ).
Carbamazepine / carbamazepine-10,11-epoxide	CT	↓ CBZ ↑ CBZ-E  ↓ valproate	Concomitant use of carbamazepine (CBZ) with divalproex sodium may result in decreased serum concentrations and half-life of valproate due to increased metabolism induced by hepatic microsomal enzyme activity. Monitoring of serum concentrations is recommended when either medication is added to or withdrawn from an existing regimen. Changes in the serum concentration of the 10,11-epoxide (CBZ-E) metabolite of carbamazepine, however, will not be detected by routine serum carbamazepine assay.  Serum levels of carbamazepine decreased 17% while that of carbamazepine-10,11-epoxide increased by 45% upon co-administration of divalproex sodium and CBZ to epileptic patients.



Concomitant Drug	Ref	Effect	Clinical comment
Carbapenem Antibiotics		↓ valproate	Carbapenem antibiotics (ertapenem, imipenem, meropenem, doripenem) can reduce serum valproic acid concentrations to sub-therapeutic levels. This can result in loss of seizure control in epileptic patients or loss of efficacy in non-epileptics. In some cases of co-administration in epileptic patients breakthrough seizures have occurred. Increasing valproic acid dose may not be sufficient to overcome this interaction. If co-administration is essential, serum valproic acid concentrations should be monitored daily. Alternative antibacterial or anticonvulsant therapy should be considered if serum valproic acid concentrations drop significantly or seizure control deteriorates. See ( <b>WARNINGS AND PRECAUTIONS, General, Interaction with Carbapenem Antibiotics</b> ).
Chlorpromazine	CT	↑ valproate	A study involving the administration of 100 to 300 mg/day of chlorpromazine to schizophrenic patients already receiving divalproex sodium (200 mg twice daily) revealed a 15% increase in trough plasma levels of valproate. This increase is not considered clinically important.
Cimetidine	T	↑ valproate	Cimetidine may decrease the clearance and increase the half-life of divalproex sodium by altering its metabolism. In patients receiving DIVALPROEX, serum valproic acid levels should be monitored when treatment with cimetidine is instituted, increased, decreased, or discontinued. The DIVALPROEX dose should be adjusted accordingly.
Clonazepam	T	No PK interaction	The concomitant use of DIVALPROEX and clonazepam may induce absence status in patients with a history of absence type seizures.
Clozapine	CT	No interaction	In psychotic patients (n = 11), no interaction was observed when divalproex sodium was co-administered with clozapine.
Diazepam	CT	↑ diazepam	Valproate displaces diazepam from its plasma albumin binding sites and inhibits its metabolism. Co-administration of divalproex sodium (1,500 mg daily) increased the free fraction of diazepam (10 mg) by 90% in healthy volunteers (n = 6). Plasma clearance and volume of distribution for free diazepam were reduced by 25% and 20%, respectively, in the presence of valproate. The elimination half-life of diazepam remained unchanged upon addition of valproate.
Ethosuximide	CT	↑ ethosuximide	Valproate inhibits the metabolism of ethosuximide. Administration of a single ethosuximide dose of 500 mg with divalproex sodium (800 to 1,600 mg/day) to healthy volunteers (n = 6) was accompanied by a 25% increase in elimination half-life of ethosuximide and a 15% decrease in its total clearance as compared to ethosuximide alone. Patients receiving DIVALPROEX and ethosuximide, especially along with other anticonvulsants, should be monitored for alterations in serum concentrations of both drugs.
Felbamate	CT	↑ valproate	A study involving the co-administration of 1,200 mg/day of felbamate with divalproex sodium to patients with epilepsy (n = 10) revealed an increase in mean valproate peak concentration by 35% (from 86 to 115 mcg/mL) compared to divalproex sodium alone. Increasing the felbamate dose to 2,400 mg/day increased the mean valproate peak concentration to 133 mcg/mL (another 16% increase). A decrease in DIVALPROEX dosage may be necessary when felbamate therapy is initiated. Lower doses of divalproex sodium may be necessary when used concomitantly with felbamate.
Haloperidol	CT	↔ valproate	A study involving the administration of 6 to 10 mg/day of haloperidol to schizophrenic patients already receiving divalproex sodium (200 mg twice daily) revealed no significant changes in valproate trough plasma levels.

Concomitant Drug	Ref	Effect	Clinical comment
Lamotrigine	CT	↑ lamotrigine ↓ valproate	<p>The effects of divalproex sodium on lamotrigine were investigated in 6 healthy male subjects. Each subject received a single oral dose of lamotrigine alone and with divalproex sodium 200 mg every 8 hours for 6 doses starting 1 hour before the lamotrigine dose was given. Divalproex sodium administration reduced the total clearance of lamotrigine by 21% and increased the plasma elimination half-life from 37.4 hours to 48.3 hours (<math>p &lt; 0.005</math>). Renal clearance of lamotrigine was unchanged. In a steady-state study involving 10 healthy volunteers, the elimination half-life of lamotrigine increased from 26 to 70 hours with divalproex sodium co-administration (a 165% increase).</p> <p>In a study involving 16 epileptic patients, divalproex sodium doubled the elimination half-life of lamotrigine. In an open-labelled study, patients receiving enzyme inducing AEDs (e.g., carbamazepine, phenytoin, phenobarbital, or primidone) demonstrated a mean lamotrigine plasma elimination half-life of 14 hours while the elimination half-life was 30 hours in patients taking divalproex sodium plus an enzyme inducing antiepileptic agent. The latter value is similar to the lamotrigine half-life during monotherapy indicating that valproic acid may counteract the effect of the enzyme inducer. If divalproex sodium is discontinued in a patient receiving lamotrigine and an enzyme inducing antiepileptic serum lamotrigine concentrations may decrease. Patients receiving combined antiepileptic therapy require careful monitoring when another agent is started, stopped or when the dose is altered.</p> <p>Serious skin reactions (such as Stevens-Johnson syndrome and Toxic Epidermal Necrolysis) have been reported with concomitant lamotrigine and divalproex sodium administration.</p>
Lithium	CT	↔ lithium	<p>In a double-blind placebo-controlled multiple dose crossover study in 16 healthy male volunteers, pharmacokinetic parameters of lithium were not altered by the presence or absence of valproate. The presence of lithium, however, resulted in an 11 to 12% increase in the AUC and <math>C_{max}</math> of valproate. <math>T_{max}</math> was also reduced. Although these changes were statistically significant, they are not likely to have clinical importance.</p> <p>Co-administration of divalproex sodium (500 mg twice daily) and lithium carbonate (300 mg three times daily) to normal male volunteers (<math>n = 16</math>) had no effect on the steady-state kinetics of lithium.</p>
Lorazepam	CT	↑ lorazepam	<p>Concomitant administration of divalproex sodium (500 mg twice daily) and lorazepam (1 mg twice daily) in normal male volunteers (<math>n = 9</math>) was accompanied by a 17% decrease in the plasma clearance of lorazepam. This decrease is not considered clinically important.</p>
Oral contraceptive Steroids	CT	No PK interaction	<p>Evidence suggests that there is an association between the use of certain AEDs capable of enzyme induction and failure of oral contraceptives. One explanation for this interaction is that enzyme-inducing drugs effectively lower plasma concentrations of the relevant steroid hormones, resulting in unimpaired ovulation. However, other mechanisms, not related to enzyme induction, may contribute to the failure of oral contraceptives. Divalproex sodium is not a significant enzyme inducer and would not be expected to decrease concentrations of steroid hormones. However, clinical data about the interaction of divalproex sodium with oral contraceptives are minimal.</p> <p>Administration of a single-dose of ethinylestradiol (50 mcg)/levonorgestrel (250 mcg) to 6 women on divalproex sodium</p>

Concomitant Drug	Ref	Effect	Clinical comment
			(200 mg twice daily) therapy for 2 months did not reveal any pharmacokinetic interaction.
Phenobarbital	CT	↑ phenobarbital	<p>Valproate was found to inhibit the metabolism of phenobarbital. Co-administration of divalproex sodium (250 mg twice daily for 14 days) with phenobarbital to normal subjects (n = 6) resulted in a 50% increase in half-life and a 30% decrease in plasma clearance of phenobarbital (60 mg single-dose). The fraction of phenobarbital dose excreted unchanged increased by 50% in the presence of valproate.</p> <p>There is evidence for severe CNS depression, with or without significant elevations of barbiturate or valproate serum concentrations. All patients receiving concomitant barbiturate therapy should be closely monitored for neurological toxicity. Serum barbiturate concentrations should be obtained, if possible, and the barbiturate dosage decreased, if appropriate.</p>
Phenytoin	CT	↑ phenytoin	<p>Valproate displaces phenytoin from its plasma albumin binding sites and inhibits its hepatic metabolism. Co-administration of divalproex sodium (400 mg three times daily) with phenytoin (250 mg) in normal volunteers (n = 7) was associated with a 60% increase in the free fraction of phenytoin. Total plasma clearance and apparent volume of distribution of phenytoin increased 30% in the presence of valproate. Both the clearance and apparent volume of distribution of free phenytoin were reduced by 25%.</p> <p>In patients with epilepsy, there have been reports of breakthrough seizures occurring with the combination of DIVALPROEX and phenytoin. The dosage of phenytoin should be adjusted as required by the clinical situation.</p>
Primidone	T	↑ phenobarbital	Primidone is metabolized into a barbiturate (phenobarbital), and therefore, may also be involved in a similar or identical interaction with divalproex sodium as phenobarbital.
Rifampin	CT	↓ valproate	A study involving the administration of a single dose of divalproex sodium (7 mg/kg) 36 hours after 5 nights of daily dosing with rifampin (600 mg) revealed a 40% increase in the oral clearance of valproate. DIVALPROEX dosage adjustment may be necessary when it is co-administered with rifampin.
Selective Serotonin Re-Uptake Inhibitors (SSRIs)	C	↑ valproate	Some evidence suggests that SSRIs inhibit the metabolism of divalproex sodium, resulting in higher than expected levels of valproate.
Tolbutamide	T	↑ tolbutamide	From in vitro experiments, the unbound fraction of tolbutamide was increased from 20 to 50% when added to plasma samples taken from patients treated with divalproex sodium. The clinical relevance of this displacement is unknown.
Topiramate	CT	Effect unknown	<p><u>Hyperammonemia</u></p> <p>Concomitant administration of divalproex sodium and topiramate has been associated with hyperammonemia with and without encephalopathy. See (CONTRAINDICATIONS) and (WARNINGS AND PRECAUTIONS, <b>Endocrine and Metabolism</b>, Urea Cycle Disorders, Hyperammonemia and Hyperammonemia and Encephalopathy Associated with Concomitant Topiramate Use).</p> <p><u>Hypothermia</u></p>

Concomitant Drug	Ref	Effect	Clinical comment
			Concomitant administration of topiramate with divalproex sodium has also been associated with hypothermia in patients who have tolerated either drug alone. Blood ammonia levels should be measured in patients with reported onset of hypothermia. See ( <b>WARNINGS AND PRECAUTIONS, <u>Endocrine and Metabolism, Hypothermia</u></b> ).
Warfarin	T	Effect unknown	In an in vitro study, valproate increased the unbound fraction of warfarin by up to 32.6%. The therapeutic relevance of this is unknown, however, coagulation tests should be monitored if DIVALPROEX is instituted in patients taking anticoagulants.  Caution is recommended when DIVALPROEX is administered with drugs affecting coagulation. See ( <b>ADVERSE REACTIONS</b> ).
Zidovudine	CT	↑ zidovudine	In 6 patients who were seropositive for HIV, the clearance of zidovudine (100 mg every 8 hours) was decreased by 38% after administration of divalproex sodium (250 or 500 mg every 8 hours); the half-life of zidovudine was unaffected.

Legend: C = Case Study; CT = Clinical Trial; T = Theoretical

### **Drug-Food Interactions**

Co-administration of DIVALPROEX with food should cause no clinical problems in the management of patients with epilepsy.

### **Drug-Herb Interactions**

Interactions with herbal products have not been established.

### **Drug-Laboratory Interactions**

Divalproex sodium is partially eliminated in the urine as a ketone-containing metabolite which may lead to a false interpretation of the urine ketone test.

There have been reports of altered thyroid function tests associated with divalproex sodium; the clinical significance of these is unknown.

### **Drug-Lifestyle Interactions**

Refer to (**WARNINGS AND PRECAUTIONS, Neurologic, Driving and Hazardous Occupations**) for details.

## DOSAGE AND ADMINISTRATION

### Dosing Considerations

#### Epilepsy

Patients receiving combined antiepileptic therapy require careful monitoring when another agent is started, stopped or when the dose is altered. See (**DRUG INTERACTIONS**).

As the dosage of DIVALPROEX (divalproex sodium) is titrated upward, blood concentrations of phenobarbital, carbamazepine and/or phenytoin may be affected. See (**DRUG INTERACTIONS**).

AEDs should not be abruptly discontinued in patients in whom the drug is administered to prevent major seizures because of the strong possibility of precipitating status epilepticus with attendant hypoxia and threat to life.

Any changes in dosage and administration, or the addition or discontinuance of concomitant drugs, should ordinarily be accompanied by close monitoring of clinical status and valproate plasma concentrations.

When changing therapy involving drugs known to induce hepatic microsomal enzymes (e.g., carbamazepine) or other drugs with valproate interactions, see (**DRUG INTERACTIONS**), it is advisable to monitor serum valproate concentrations.

#### Dosing in Elderly Patients

Due to a decrease in unbound clearance of valproate and possibly a greater sensitivity to somnolence in the elderly, the starting dose should be reduced. Dosage should be increased more slowly and with regular monitoring for fluid and nutritional intake, dehydration, somnolence, urinary tract infection, and other adverse events. Dose reductions or discontinuation of DIVALPROEX should be considered in patients with decreased food or fluid intake and in patients with excessive somnolence. The ultimate therapeutic dose should be achieved on the basis of clinical response. See [**WARNINGS AND PRECAUTIONS, Special Populations, Geriatrics (≥ 65 years of age)**].

#### Dose-Related Adverse Events

The frequency of adverse events (particularly elevated liver enzymes and thrombocytopenia) may be dose related. The probability of thrombocytopenia appears to increase significantly at total valproate concentration of  $\geq 110$  mcg/mL (females) or  $\geq 135$  mcg/mL (males). See (**WARNINGS AND PRECAUTIONS, Hematologic, Dose-related Adverse Reactions: Thrombocytopenia**). Therefore, the benefit of improved therapeutic effect with higher doses should be weighed against the possibility of a greater incidence of adverse effects.

## **Recommended Dose and Dosage Adjustment**

### **Epilepsy**

DIVALPROEX (divalproex sodium) enteric-coated tablets are administered orally. The recommended initial dosage is 15 mg/kg/day, increasing at one-week intervals by 5 to 10 mg/kg/day until seizures are controlled or side effects preclude further increases.

The maximal recommended dosage is 60 mg/kg/day. When the total daily dose is 250 mg and over, it should be given in a divided regimen (**Table 3**).

**Table 3. Initial Doses by Weight (based on 15 mg/kg/day)**

Weight		Total Daily Dose (mg)	Dosage (mg) equivalent to valproic acid		
kg	lb		Dose 1	Dose 2	Dose 3
10 to 24.9	22 to 54.9	250	125	0	125
25 to 39.9	55 to 87.9	500	250	0	250
40 to 59.9	88 to 131.9	750	250	250	250
60 to 74.9	132 to 164.9	1,000	250	250	500
75 to 89.9	165 to 197.9	1,250	500	250	500

### **Therapeutic Blood Levels**

A good correlation has not been established between daily dose, total serum valproate concentration and therapeutic effect. However, therapeutic valproate serum concentrations for most patients with epilepsy will range from 50 to 100 mcg/mL (350 to 700 micromole/L). Some patients may be controlled with lower or higher serum concentrations. See (**WARNINGS AND PRECAUTIONS**).

### **Conversion from Valproic Acid to Divalproex Sodium**

Divalproex sodium enteric-coated tablets dissociate to the valproate ion in the gastrointestinal tract. Divalproex sodium tablets are uniformly and reliably absorbed, however, because of the enteric coating, absorption is delayed by an hour when compared to valproic acid capsules.

The bioavailability of divalproex sodium tablets is equivalent to that of valproic acid capsules.

In patients previously receiving valproic acid therapy, divalproex sodium should be initiated at the same daily dosing schedule. After the patient is stabilized on divalproex sodium, a dosing schedule of two or three times a day may be elected in selected patients. Changes in dosage administration of divalproex sodium or concomitant medications should be accompanied by increased monitoring of plasma concentrations of valproate and other medications, as well as the patient's clinical status.

## **Acute Mania**

DIVALPROEX is not indicated for the treatment of the symptoms of mania in patients under 18 years of age.

The recommended initial dose is 250 mg three times a day. The dose should be increased as rapidly as possible to achieve the lowest therapeutic dose which produces the desired clinical effect or the desired range of plasma concentrations.

In placebo-controlled trials, 84% of patients received and tolerated maximum daily doses of between 1,000 to 2,500 mg/day. The maximum recommended dosage is 60 mg/kg/day.

The relationship of plasma concentration to clinical response has not been established for divalproex sodium. In controlled clinical studies, 79% of patients achieved and tolerated serum valproate concentrations between 50 µg/mL and 125 µg/mL.

## **Missed Dose**

The patient should not abruptly stop taking their medication because of the risk of increasing their seizures.

If the patient misses a dose, they should not try to make up for it by doubling up on their next dose. They should take their next regularly scheduled dose and try not to miss any more doses.

## **Administration**

DIVALPROEX may be taken with or without food.

Patients who experience gastrointestinal irritation may benefit from administration of the drug with food or by a progressive increase of the dose from the initial low level. The tablets should be swallowed without chewing. Co-administration of DIVALPROEX with food should cause no clinical problems in the management of patients with epilepsy.

## **OVERDOSAGE**

For management of a suspected drug overdose, contact your regional Poison Control Centre.
---

Overdosage with DIVALPROEX (divalproex sodium) may result in somnolence, heart block, and deep coma. Fatalities have been reported; however, patients have recovered from valproate levels as high as 2,120 mcg/mL.

In a reported case of overdosage with divalproex sodium after ingesting 36 g in combination with phenobarbital and phenytoin, the patient presented in deep coma. An electroencephalogram (EEG) recorded diffuse slowing, compatible with the state of consciousness. The patient made an uneventful recovery.

In overdose situations, the fraction of drug not bound to protein is high and hemodialysis or tandem hemodialysis plus hemoperfusion may result in significant removal of drug. The benefit of gastric lavage or emesis will vary with the time since ingestion. General supportive measures should be applied with particular attention to the prevention of hypovolemia and the maintenance of adequate urinary output.

Naloxone has been reported to reverse the CNS depressant effects of divalproex sodium overdosage. Because naloxone could theoretically also reverse the antiepileptic effects of divalproex sodium it should be used with caution in patients with epilepsy.

## **ACTION AND CLINICAL PHARMACOLOGY**

### **Mechanism of Action**

Divalproex sodium has anticonvulsant properties, and is chemically related to valproic acid. Divalproex sodium dissociates to the valproate ion in the gastrointestinal tract. Although its mechanism of action has not yet been established, it has been suggested that its activity in epilepsy is related to increased brain concentrations of gamma-aminobutyric acid (GABA). The effect on the neuronal membrane is unknown.

### **Pharmacodynamics**

A good correlation has not been established between daily dose, serum level and therapeutic effect. In epilepsy, the therapeutic plasma concentration range is believed to be from 50 to 100 mcg/mL (350 to 700 micromole/L) of total valproate. Occasional patients may be controlled with serum levels lower or higher than this range. See (**DOSAGE AND ADMINISTRATION**). In placebo-controlled clinical studies in acute mania, 79% of patients were dosed to a plasma concentration between 50 and 125 mcg/mL. Protein binding of valproate is saturable ranging from 90% at 50 mcg/mL to 82% at 125 mcg/mL.

### **Pharmacokinetics**

**Table 4. Summary of the Pharmacokinetic Parameters of Divalproex Sodium in Healthy, Fasting Subjects**

Single Dose	Dosage	N	Mean (SD) Pharmacokinetic Parameters					
			C <sub>max</sub> (mg/L)	T <sub>max</sub> (h)	t <sub>½</sub> (h)	AUC <sub>∞</sub> (mg•h/L)	CL (l/h)	V <sub>d</sub> (L)
Divalproex Sodium	2 x 500 mg once daily	28	93.9 (11.7)	4.0 (1.2)	15.2 (15.3)	1818 (345)	--	--



## **Absorption**

Peak serum levels of valproic acid occur in 3 to 4 hours. A slight delay in absorption occurs when the drug is administered with meals but this does not affect the total absorption.

## **Distribution**

Valproic acid is rapidly distributed throughout the body and the drug is strongly bound (90%) to human plasma proteins. Increases in doses may result in decreases in the extent of protein binding and variable changes in valproic acid clearance and elimination.

## Protein Binding

The plasma protein binding of valproate is concentration dependent and the free fraction increases from approximately 10% at 40 mcg/mL to 18.5% at 130 mcg/mL. Protein binding of valproate is reduced in the elderly, in patients with chronic hepatic diseases, in patients with renal impairment, in hyperlipidemic patients, and in the presence of other drugs (e.g., acetylsalicylic acid). Conversely, valproate may displace certain protein-bound drugs (e.g., phenytoin, carbamazepine, warfarin, and tolbutamide). See (**DRUG INTERACTIONS**) for more detailed information on the pharmacokinetic interactions of valproate with other drugs.

## CNS Distribution

Valproate concentrations in cerebrospinal fluid (CSF) approximate unbound concentrations in plasma (ranging from 7 to 25% of total concentration).

## **Metabolism:**

Valproate is metabolized almost entirely by the liver. In adult patients on monotherapy, 30 to 50% of an administered dose appears in urine as a glucuronide conjugate. Mitochondrial (beta)-oxidation is the other major metabolic pathway, typically accounting for over 40% of the dose. Usually, less than 15 to 20% of the dose is eliminated by other oxidative mechanisms. Less than 3% of an administered dose is excreted unchanged in urine.

Due to the saturable plasma protein binding, the relationship between dose and total valproate concentration is nonlinear; concentration does not increase proportionally with the dose, but rather increases to a lesser extent. The kinetics of unbound drug are linear.

## **Excretion:**

Mean plasma clearance and volume of distribution for total valproate are 0.56 L/hr/1.73 m<sup>2</sup> and 11 L/1.73 m<sup>2</sup>, respectively. Mean plasma clearance and volume of distribution for free valproate are 4.6 L/hr/1.73 m<sup>2</sup> and 92 L/1.73 m<sup>2</sup>, respectively. These estimates cited apply primarily to patients who are not taking drugs that affect hepatic metabolizing enzyme systems. For example, patients taking enzyme-inducing AEDs (carbamazepine, phenytoin, and phenobarbital) will clear valproate more rapidly. Because of these changes in valproic acid clearance, monitoring of valproate and concomitant drug concentrations should be intensified whenever enzyme-inducing

drugs are introduced or withdrawn.

Elimination of valproic acid and its metabolites occurs principally in the urine, with minor amounts in the feces and expired air. Very little unmetabolized parent drug is excreted in the urine.

The serum half-life ( $t_{1/2}$ ) of valproic acid is typically in the range of 6 to 16 hours. Half-lives in the lower part of the above range are usually found in patients taking other drugs capable of hepatic enzyme induction.

### **Special Populations and Conditions**

#### **Neonates/Infants**

Within the first 2 months of life, infants have a markedly decreased ability to eliminate valproate compared to children and adults. This is a result of reduced clearance (perhaps due to delay in development of glucuronosyltransferase and other enzyme systems involved in valproate elimination) as well as increased volume of distribution (in part due to decreased plasma protein binding). For example, in one study, the half-life in neonates under 10 days ranged from 10 to 67 hours, compared to a range of 7 to 13 hours in children greater than 2 months.

#### **Pediatrics**

Patients between 3 months and 10 years have 50% higher clearances expressed on weight (i.e., mL/min/kg) than do adults. Over the age of 10 years, children have pharmacokinetic parameters that approximate those of adults.

#### **Geriatrics**

The capacity of elderly patients (age range: 68 to 89 years) to eliminate valproate has been shown to be reduced compared to younger adults (age range: 22 to 26 years). Intrinsic clearance is reduced by 39%; the free fraction is increased by 44%. See (**DOSAGE AND ADMINISTRATION**).

#### **Gender**

There are no differences in unbound clearance (adjusted for body surface area) between males and females ( $4.8 \pm 0.17$  and  $4.7 \pm 0.07$  L/hr per 1.73 m<sup>2</sup>, respectively).

#### **Race**

The effects of race on the kinetics of valproate have not been studied.

#### **Hepatic Insufficiency**

See (**CONTRAINDICATIONS**) and (**WARNINGS AND PRECAUTIONS**),

**Hepatic/Biliary/Pancreatic, Serious or Fatal Hepatotoxicity**) for statements regarding hepatic dysfunction and associated fatalities.

### **Renal Insufficiency**

See (WARNINGS AND PRECAUTIONS, **Renal**, **Renal Impairment**).

### **Genetic Polymorphism**

No data available on genetic polymorphism.

## **STORAGE AND STABILITY**

Store at controlled room temperature 15-30°C (59-86°F) in tight, light-resistant containers.

## **DOSAGE FORMS, COMPOSITION AND PACKAGING**

DIVALPROEX 125 mg enteric coated tablets: Each red-coloured, oblong, enteric-coated tablet, engraved "APO" on one side and "125" on the other contains divalproex sodium equivalent to 125 mg of valproic acid. Available in bottles of 100 tablets.

In addition to the active ingredient, divalproex sodium, each tablet also contains the non-medicinal ingredients: D & C Red No. 30, eudragit, FD & C Yellow No. 6, guar gum, hydroxypropyl cellulose, hydroxypropylmethylcellulose, microcrystalline cellulose, polyethylene glycol, talc, titanium dioxide, triethyl citrate .

DIVALPROEX 250 mg enteric coated tablets: Each peach-coloured, oblong, enteric-coated tablet, engraved "APO" on one side and "250" on the other contains divalproex sodium equivalent to 250 mg of valproic acid. Available in bottles of 100 tablets.

In addition to the active ingredient, divalproex sodium, each tablet also contains the non-medicinal ingredients: eudragit, FD&C Yellow No. 6, guar gum, hydroxypropyl cellulose, hydroxypropylmethylcellulose, microcrystalline cellulose, polyethylene glycol, talc, titanium dioxide, triethyl citrate.

DIVALPROEX 500 mg enteric coated tablets: Each pink-coloured, oblong, enteric-coated tablet, engraved "APO" on one side and "500" on the other contains divalproex sodium equivalent to 500 mg of valproic acid. Available in bottles of 100 tablets.

In addition to the active ingredient, divalproex sodium, each tablet also contains the non-medicinal ingredients: D & C Red No. 33, eudragit, guar gum, hydroxypropyl cellulose, hydroxypropylmethylcellulose, microcrystalline cellulose, polyethylene glycol, red ferric oxide, talc, titanium dioxide, triethyl citrate.

## PART II: SCIENTIFIC INFORMATION

### PHARMACEUTICAL INFORMATION


#### Drug Substance

Proper Name: Divalproex sodium

Chemical Name: 1) Pentanoic acid, 2-propyl-, sodium salt (2:1);  
2) Sodium hydrogen bis(2-propylvalerate).

Molecular formula and molecular weight: Molecular Formula:  $C_{16}H_{31}NaO_4$   
Molecular Weight: 310.41

Structural Formula:



Physicochemical properties: Divalproex sodium is a white powder with a characteristic odor, freely soluble in many organic solvents and in aqueous alkali solutions.

### CLINICAL TRIALS

#### Comparative Bioavailability Studies

Two comparative bioavailability studies were performed: one with food and one without food. The double-blind, randomized, two-way crossover studies were conducted in healthy, adult, male volunteers to evaluate the relative bioavailability of single oral doses of DIVALPROEX 500 mg tablets manufactured by Sanis Health Inc. and Epival® 500 mg tablets, manufactured by Abbott Laboratories. The mean pharmacokinetic parameters of these subjects are summarized in

the following tables:

**Table 5: Summary Table of the Comparative Bioavailability Data (Fasting)**

STUDY 1 (Fasting)				
Divalproex Sodium Enteric-Coated Tablets (Dose: 500 mg)				
From Measured Valproic Acid Data				
Parameter	Geometric Mean Arithmetic Mean (CV%)		Ratio of Geometric Means **	95% Confidence Interval
	DIVALPROEX	Epival <sup>®</sup> †		
AUC <sub>T</sub> (µg•hr/mL)	1118 1138 (20.0)	1158 1177 (19.8)	0.97	93-100
AUC <sub>1</sub> (µg•hr/mL)	1231 1264 (25.0)	1265 1294 (23.2)	0.98	94-101
C <sub>max</sub> (µg/mL)	66.04 66.81 (16.4)	65.36 66.48 (19.0)	1.01	93-108
T <sub>max</sub> (hr)*	3.10 (24.5)	3.67 (27.1)	--	--
t <sub>1/2</sub> (hr)*	17.48 (26.5)	16.86 (20.2)	--	--
* Arithmetic means only (CV%).				
** Based on the least square estimate of the geometric means.				
† Epival <sup>®</sup> is manufactured by Abbott Laboratories Limited, and was purchased in Canada.				

**Table 6: Summary Table of the Comparative Bioavailability Data (With Food)**

STUDY 2 (With food)				
Divalproex Sodium Enteric-Coated Tablets (Dose: 500 mg)				
From Measured Valproic Acid Data				
Parameter	Geometric Mean Arithmetic Mean (CV%)		Ratio of Geometric Means	95% Confidence Interval
	DIVALPROEX	Epival <sup>®</sup> †		
AUC <sub>T</sub> (µg•hr/mL)	986.3 1007 (21.1)	971.0 991.7 (22.7)	1.02	99-105
AUC <sub>1</sub> (µg•hr/mL)	1098 1131 (25.6)	1094 1136 (33.2)	1.00	97-104
C <sub>max</sub> (µg/mL)	50.83 51.38 (15.2)	51.15 51.58 (13.3)	0.99	95-105
T <sub>max</sub> (h)*	5.69 (78.1)	7.27 (40.7)	--	--
t <sub>1/2</sub> (h)*	17.60 (21.9)	17.42 (28.4)	--	--
* Arithmetic means only (CV%).				
† Epival <sup>®</sup> is manufactured by Abbott Laboratories Limited, and was purchased in Canada.				

## DETAILED PHARMACOLOGY

### Animal

Valproic acid has been shown to be effective against several types of chemically and electrically

induced convulsions in a variety of animal species. These included maximal electroshock, low frequency electroshock, CO<sub>2</sub> withdrawal, pentylene tetrazole, cobalt, bemegride, bicuculline and l-glutamate. Many forms of photic and auditory induced seizures are also effectively blocked by valproic acid.

In animal studies, valproic acid at doses of 175 mg/kg or less had no effect on locomotor activity and conditioned responses to positive reinforcement.

Doses greater than 175 mg/kg inhibited spontaneous and conditioned behaviour in mice and rats and interfered with coordination of hind limbs in rats. Suppression of spontaneous and evoked brain potentials was also demonstrated at these higher dose levels.

Valproic acid at doses of 175 mg/kg or less had little or no effect on the autonomic nervous system, cardiovascular system, respiration, body temperature, inflammatory responses, smooth muscle contraction or renal activity. Intravenous doses of 22, 43 and 86 mg/kg in animals caused very transient decreases followed by compensatory increases in blood pressure.

Sodium valproate injectable caused decreased activity, ataxia, dyspnea, prostration and death in rats and mice acutely exposed to dosages exceeding 200 mg/kg.

Divalproex sodium produced plasma valproic acid concentrations comparable to those of valproic acid when the two compounds were administered orally at equimolar doses to mice, rats and a beagle dog.

## **TOXICOLOGY**

The initial animal testing was done with sodium valproate, whereas most of the recent research has been with valproic acid. The conversion factor is such that 100 mg of the sodium salt is equivalent to 87 mg of the acid. References to dosage are in terms of valproic acid activity.

### **Acute Toxicity**

Acute toxicity has been determined in several animal species using oral, intravenous, intraperitoneal and subcutaneous routes. The oral median lethal dose in adult rats and dogs was about 1 to 2 g/kg. Toxicity was similar for both sexes; however, it tended to be greater in newborn and 14-day old rats and in young adult rats. The signs of toxicity were those of central nervous system depression. Specific organ damage was limited to cellular debris in reticuloendothelial tissue and slight fatty degeneration of the liver.

Large oral doses (more than 500 mg/kg) produced irritation of the gastrointestinal tract of rats.

In adult male mice, the oral medial lethal dose of divalproex sodium was 1.66 g/kg (equal to approximately 1.54 g/kg valproic acid).

Pulverized divalproex sodium enteric-coated tablets (equivalent to 250 mg valproic acid), suspended in 0.2% methylcellulose, were administered orally to mice and rats of both sexes

(10/sex/species/group) in dosages ranging from 1.74 to 4.07 g/kg. The oral median lethal dose (LD50) ranged from 2.06 to 2.71 g/kg. No consistent sex-related or species-related differences were observed.

Signs of central nervous system depression, such as decreased activity, ataxia, and sleep, were observed. At necropsy, discolouration and/or thickening of the glandular mucosa were observed in only 2 female rats treated with 2.71 g/kg that died acutely.

When mature rats and dogs were administered up to 240 mg/kg/day or 120 mg/kg/day, respectively, for at least four consecutive weeks, no significant toxicologic effects were reported. However, significant reductions in testicular weights and total white cell counts in rats given 240 mg/kg/day were considered as evidence of subtle toxicity from sodium valproate injectable. Therefore, 90 mg/kg/day in rats and 120 mg/kg/day in dogs were considered the highest nontoxic doses.

The acute intravenous toxicity of sodium valproate injectable formulation containing the equivalent of 100 mg valproic acid/mL was evaluated in both sexes of mice and rats. Groups of mice and rats (five/sex/species/group) were treated at dosages ranging from 0.5 to 9.0 mL/kg (50 to 900 mg valproate/kg). No overt signs of toxicity were present in rats and mice given 0.5 mL/kg (50 mg valproate/kg). LD50 values for the test solution in mice and rats (data combined for both sexes) were 7.3 and 7.0 mL/mg (730 and 700 mg valproate/kg), respectively.

### **Subacute and Chronic Toxicity**

Subacute and chronic toxicity studies consisted of 1, 3, 6 and 18 months studies in rats and 3, 6 and 12 months studies in dogs. Pathologic changes included suppression of the hematopoietic system, depletion of lymphocytes from lymphoid tissues and the loss of germinal epithelial cells from seminiferous tubules. Reduced spermatogenesis and testicular atrophy occurred in dogs at doses greater than 90 mg/kg/day and in rats at doses greater than 350 mg/kg/day. In rats, the first indication of toxicity at 350 mg/kg/day was decreased food consumption and growth.

### **Mutagenicity and Carcinogenicity**

#### **Mutagenicity**

Valproate was not mutagenic in an in vitro bacterial assay (Ames test), did not produce dominant lethal effects in mice, and did not increase chromosome aberration frequency in an in vivo cytogenetic study in rats. Increased frequencies of sister chromatid exchange (SCE) have been reported in a study of epileptic children taking valproate, but this association was not observed in another study conducted in adults. There is some evidence that increased SCE frequencies may be associated with epilepsy. The biological significance of increase in SCE frequency is not known.

#### **Carcinogenicity**

Two hundred rats were given valproic acid in the diet for 107 weeks. Mean doses consumed in

the treatment period were: 81 mg/kg/day (males) and 85 mg/kg/day (females), in the low dose group; 161 mg/kg/day (males) and 172 mg/kg/day (females) in the high dose group (approximately 10 to 50% of the maximum human daily dose on a mg/m<sup>2</sup> basis). Control animals received corn oil in the diet. The chief finding in the study was an increased incidence of skin fibrosarcomas in treated males of the high-dose group. There were 2 such neoplasms in the low dose group, 5 in the high dose group and none in control males. Fibrosarcomas in rats are relatively infrequent, usually occurring in less than 3% of animals.

Valproic acid was also administered in the diet to female mice for nearly 19 months at doses of 81 and 163 mg/kg/day and to male mice for nearly 23 months at doses of 80 and 159 mg/kg/day. A significant dose related trend occurred in male mice in the incidence of bronchoalveolar adenomas, and when the data were adjusted for the times of death, the incidence in the high dose group was significantly increased.

Depending on the method of statistical analysis, the incidence of hepatocellular carcinomas and/or adenomas also showed significant or almost significant increases for the corresponding observations. The results of these two studies indicate that valproic acid in a weak carcinogen or promoter in rats and mice. The significance of these findings for humans is unknown at present.

### **Reproduction and Teratology**

Studies in rats have shown placental transfer of the drug. Doses greater than 65 mg/kg/day given to rats, mice and rabbits produced an increased incidence of skeletal abnormalities of the ribs, vertebrae and palate.

Doses greater than 150 mg/kg/day given to pregnant rabbits produced fetal resorptions and (primarily) soft-tissue abnormalities in the offspring.

In rats, there was a dose related delay in onset of parturition. Post-natal growth and survival of the progeny were adversely affected, particularly when drug administration spanned the entire gestation and early lactation period. Embryoletality or major developmental abnormalities occurred in rats and rabbits at doses of 350 mg/kg/day.

Survival among pups born to the high dose females was very poor but was improved when pups were transferred to control dams shortly after birth.

### **Fertility**

Chronic toxicity studies in juvenile and adult rats and dogs demonstrated reduced spermatogenesis and testicular atrophy at oral doses of valproic acid of 400 mg/kg/day or greater in rats (approximately equivalent to or greater than the maximum human daily dose on a mg/m<sup>2</sup> basis) and 150 mg/kg/day or greater in dogs (approximately 1.4 times the maximum human daily dose or greater on a mg/m<sup>2</sup> basis). Segment I fertility studies in rats have shown that oral doses up to 350 mg/kg/day (approximately equal to the maximum human daily dose on a mg/m<sup>2</sup> basis) for 60 days have no effect on fertility.



The effect of valproate on testicular development and on sperm production and fertility in humans is unknown.

## REFERENCES

- 1) Bartels H, Oldiss HD, Wallis S. Faecal Excretion of non-disintegrated retard-coated dragees of sodium valproate (ersenyl retard). *Neuropediatric* 1979;10:301-302.
- 2) Benoit JM, Besson JM, Bivot JB, Aleonard P. Effects of Na dipropylacetate (DPA) on dorsal root potentials induced by cortical seizures. *J Pharmacol* 1971;2:23.
- 3) Binnie CD, van Emde Boas W, Kasteleijn-Nolste-Trenite DGA et al. Acute effects of lamotrigine (BW430C) in persons with epilepsy. *Epilepsia* 1986;27(3):248-254.
- 4) Bowdle TA, Patel IH, Levy RH, Wilensky AJ. Valproic acid dosage and plasma protein binding and clearance. *Clin Pharmacol Ther* 1980;28(4):486-492.
- 5) Brennan MJW, Sandyk R, Borsook D. Efficacy of valproate in manic-depressive psychosis: mechanism of action and implications for pathophysiology. *Psychiatry: the state of the art* 1985;3:459-465.
- 6) Brodie MJ. Lamotrigine. *Lancet* 1992;339:1397-1400.
- 7) Bruni J, Wilder BJ, Perchalski RJ, Hammond EJ, Villarreal HJ. Valproic acid and plasma levels of phenobarbital. *Neurology* 1980; 30:94-97.
- 8) Brusilow SW, Maestri NE, Urea Cycle Disorders: Diagnosis, Pathophysiology, and Therapy: *Advances in Pediatrics* . 1996; 43: 127-170.
- 9) Calabrese JR, Rappert DJ, Kimmel SE, Reece B, Woysville MJ. Rapid cycling bipolar disorder and its treatment with valproate. *Can J Psychiatry* 1993;38 (suppl 2):S57-S61.
- 10) Cohen AF, Land GS, Breimer DD et al. Lamotrigine, a new anticonvulsant: Pharmacokinetics in normal humans. *Clin Pharmacol Ther* 1987;42:535-541.
- 11) Cramer JA, Mattson RH. Valproic acid: In vitro plasma protein binding and interaction with phenytoin. *Ther Drug Monit* 1979;1(1): 105-116.
- 12) Centers for Disease Control, valproate: a new cause of birth defects - report from Italy and follow-up from France. *Morbidity and Mortality Weekly Report*. 1983;32(33):438-439.
- 13) Dreifuss FE, Langer DH, Moline KA, Maxwell BA. Valproic acid hepatic fatalities. II. US experience since 1984. *Neurology* 1989;39:201-207.
- 14) Eadie MJ. Plasma level monitoring of anticonvulsants. *Clin Pharmacokinet* 1976;1:52-66.

- 15) Freeman TW. A double-blind comparison of valproate and lithium in the treatment of acute mania. *Am J Psychiatry* 1992;149:108-111.
- 16) Frey HH, Loscher W. Di-n-propylacetic acid - Profile of anticonvulsant activity in Mice. *Arzneimittelforschung* 1976;26:299-301.
- 17) Garibaldi RA. Residential care and the elderly: the burden of infection. *J Hosp Infect.* 1999 Dec;43 Suppl:S9-18. Review.
- 18) Gidal B, et. al. Valproate-mediated disturbances of hemostatis. *Neurology* 1994;44:1418- 1422.
- 19) Godin Y, Heiner L, Mark J, Mandel P. Effects of di-n-propylacetate, an anticonvulsive compound, on GABA metabolism, *J Neurochem* 1969;16:869-873.
- 20) Gram L, Wulff K, Rasmussen KE, Flachs H, Wurtz-Jorgensen A, Sommerbeck KW, Lohren V. Valproate sodium: A controlled clinical trial including monitoring of drug levels. *Epilepsia* 1977;18:141-148.
- 21) Gram L, Flachs H, Wurtz-Jorgensen A, Parnas J, Andersen B. Sodium valproate, serum level and clinical effect in epilepsy: A controlled study. *Epilepsia* 1979;20:303-311.
- 22) Gugler R, Schell A, Froscher W, Schulz HU. Oral disposition of valproic acid following single and multiple doses. *Arch Pharmacol* 1977;297(Suppl. 2):R62.
- 23) Gugler R, Von Unruh GE. Clinical pharmacokinetics of valproic acid. *Clin Pharmacokinet* 1980;5:67-83.
- 24) Hurst DL. Expanded therapeutic range of valproate. *Pediatr Neurol.* 1987;3:342-344.
- 25) Jaffe RL, Rives W, Dubin WR, Roemer RA, Siegal L. Problems in maintenance ECT in bipolar disorder: replacement of lithium and anticonvulsants. *Convulsive therapy* 1991;7:288-294.
- 26) Jawad S, Yuen WC, Peck AW et al. Lamotrigine: single-dose pharmacokinetics and initial 1-week experience in refractory epilepsy. *Epilepsy Res* 1987;1(3):194-201.
- 27) Johannessen SI, Henriksen O. Comparative steady state serum levels of valproic acid administered as two different formulations - Deprakine and Orfiril. *Acta Neurol Scand* 1979;60:371-374.
- 28) Jordan BJ, Orwin JM, Nichols JD, Henson R. The bioavailability of sodium valproate (2-PP) from three epilim formulations. *Pharm Weekly* 1977;112:347-349.
- 29) Kastner T, Friedman DL. Verapamil and valproic acid treatment of prolonged mania. *J Am Acad Child Psychiatry* 1992;31:271-275.

- 30) Kupferberg HJ, Lust WD, Penry JK. Anticonvulsant activity of dipropylacetic acid (DPA) in relation to GABA and CGMP brain levels in mice. *Fed Proc* 1975;34:283.
- 31) Kapetamovic IM, Kupferberg HJ. Inhibition of microsomal Phenobarbital metabolism by valproic acid. *Biochem Pharmacol* 1981;30(11): 1361-1363.
- 32) 32. Levy RH, Cenraub B, Loiseau P, Akbaraly R, Brachet-Liermain A, Guyot M, Gomeni R, Morselli PL. Meal-dependent absorption of enteric-coated sodium valproate. *Epilepsia* 1980;21:273-280.
- 33) Lum M, Fontaine R, Elie R, Ontiveros A. Probable interaction of sodium divalproex with benzodiazepines. *Prog Neuropsychopharmacol Biol Psychiatry* 1991;15:269-273.
- 34) McElroy SL, Keck PE. Treatment guidelines for valproate in bipolar and schizoaffective disorders. *Can J Psychiatry* 1993;38(suppl 2):S62-S66.
- 35) McFarland BH, Miller MR, Straumfjord AA. Valproate use in the older manic patient. *J Clin Psychiatry* 1990;51:479-481.
- 36) Mattson RH, et al. Use of oral contraceptives by women with epilepsy. *JAMA* 1986;256(2):238-240.
- 37) Millet Y, Sainty JM, Galland MC, Sidoine R, Jouglard J. Problèmes posés par l'association thérapeutique phénobarbital-dipropylacétate de sodium. A propos d'un cas. *Eur J Toxicol Environ Hyg* 1976;9(6): 381-383.
- 38) Misslin R, Ropartz P, Mandel P. The effects of n-dipropylacetate on the acquisition of conditioned behaviour with negative reinforcement in mice. *Psychopharmacologia* 1975;44:263-265.
- 39) Nicolle LE. Urinary tract infections in long-term-care facilities. *Infect Control Hosp Epidemiol.* 2001 Mar;22(3):167-75.
- 40) Patel IH, Levy RH, Cutler RE. Phenobarbital - valproic acid interaction. *Clin Pharmacol Ther* 1980;27(4):515-521.
- 41) Pinder PM, Brogden RN, Speight TM, Avery GS. Sodium valproate: A review of its pharmacological properties and therapeutic efficacy in epilepsy. *Drugs* 1977;13:81-123.
- 42) Pisani F, Primerano G, D'Agostino AA, et al. Valproic Acid-Amitriptyline Interaction in Man. *Therapeutic Drug Monitoring.* 1986; 8(3): 382-3.
- 43) Pope HG Jr, McElroy SL, Keck PE, Hudson JI. Valproate in the treatment of acute mania: a placebo-controlled study. *Arch Gen Psychiatry* 1991;48:62-68.

- 44) Post RM. Introduction: emerging perspectives on valproate in affective disorders. *J Clin Psychiatry* 1989;50(suppl 3):3-9.
- 45) Potter WZ, Ketter TA. Pharmacological issues in the treatment of bipolar disorder: Focus on mood-stabilizing compounds. *Can J Psychiatry* 1993;38(suppl 2):S51-S56.
- 46) Rai PV. Acute intoxication during a combined treatment of sodium valproate and phenobarbitone. In: Meinardi H and Rowan AJ, editors. *Advances in epileptology*, Swets Zertlinger, Amsterdam, pp. 366-369, 1978.
- 47) Robert E, Guibaud P. Maternal valproic acid and congenital neural tube defects. *Lancet* 1982;II:937.
- 48) Scheyer RD, Cramer JA, Totftness BR, Hochholzer JM, Mattson RH. In vivo determination of valproate binding constants during sole and multi-drug therapy. *The Drug Monitor* 1990;12:117-123.
- 49) Shakir RA, Johnson RH, Lambie DG, Melville ID, Nanda RN. Comparison of sodium valproate and phenytoin as single drug treatment in epilepsy. *Epilepsia* 1981;22:27-33.
- 50) Shakir RA, Nanda RN, Lambie DG, Johnson RH. Comparative trial of valproate sodium and clonazepam in chronic epilepsy. *Arch Neurol* 1979; 36:301-304.
- 51) Shearer DE, Fleming DE, Bigler ED, Wilson CE. Suppression of photically evoked afterdischarge bursting following administration of anticonvulsants in waking rats. *Pharmacol Biochem Behav* 1974; 2:839-842.
- 52) Shuto K, Nishigaki T. The pharmacological studies on sodium dipropylacetate anticonvulsant activities and general pharmacological actions. *Oyo Yakuri (Pharmacometrics)* 1970;4:937.
- 53) Simler S, Ciesielski L, Maitre M, Randrianarisoa, Mandel P. Effects of sodium ndipropylacetate on audiogenic seizures and brain aminobutyric acid level. *Biochem Pharmacol* 1973;22:1701-1708.
- 54) Simon D, Penry JK. Sodium di-n-propylacetate (DPA) in the treatment of epilepsy. A review, *Epilepsia* 1975;16(4):549-573.
- 55) Sovner R. A clinically significant interaction between carbamazepine and valproic acid. *J Clin Psychopharmacol* 1988;8:448-449.
- 56) Tatro DS, Editor. *Drug interaction facts*. The Mediphor Editorial Group, Division of Clinical Pharmacology, Stanford University School of Medicine. St. Louis, MO: Facts and Comparisons Division, J.B. Lippincott Co.; 1992.

- 57) Turnbull DM, Rawlins MD, Weightman D, Chadwick DW. A comparison of phenytoin and valproate in previously untreated adult epileptic patients. *J Neurol Neurosurg Psychiatry* 1982;45:55-59.
- 58) Van de Mortel I. Étude pharmacokinétique du di-npropylacétate de sodium sous ses formes ordinaire et retard. *Acta Neurol Belg* 1979; 76:165-172.
- 59) Van Valkenburg C, Kluznik J, Merrill R, Erickson W. Therapeutic levels of valproate for psychosis. *Psychopharmacol Bulletin* 1990;26:254-255.
- 60) Watanabe J. Plasma protein binding and interaction of valproic acid with acetazolamide and diazepam, both in vivo and in vitro studies. *Hirosaki Igaku (Hirosaki Med J)* 1985;37:611- 625. (English translation of a Japanese article)
- 61) Wilder BJ, Karas BJ, Hammond EJ, Perchalski RJ. Twice-daily dosing of valproate with divalproex. *Clin Pharmacol Ther* 1983;34(4):501-504.
- 62) Wilder BJ, Karas BJ, Penry JK, Asconape J. Gastrointestinal tolerance of divalproex sodium. *Neurology* 1983;33:808-811.
- 63) Wong SL, Cavanaugh J, Shi H, et al. Effects of Divalproex Sodium on Amitriptyline and Nortriptyline Pharmacokinetics. *Clin Pharmacol Ther.* July 1996; 60(1): 48-53.
- 64) Yuen AWC, Land G, Weatherley BC, Peck AW. Sodium valproate acutely inhibits lamotrigine metabolism. *Br J Clin Pharmacol* 1992;33:511
- 65) Product Monograph - Epival® (divalproex sodium) Enteric-Coated tablets (125 mg, 250 mg and 500 mg). Abbott Laboratories, Limited. Date of Revision: September 10, 2010, Control number: 137362

**PART III: CONSUMER INFORMATION**

**Pr DIVALPROEX**  
**Divalproex sodium enteric-coated tablets**  
**House Standard**

This leaflet is part III of a three-part “Product Monograph” published when DIVALPROEX was approved for sale in Canada and is designed specifically for Consumers. This leaflet is a summary and will not tell you everything about DIVALPROEX. Contact your doctor or pharmacist if you have any questions about the drug.

**ABOUT THIS MEDICATION**

**What the medication is used for:**

DIVALPROEX has been prescribed to you to either:

- control your epilepsy
- treat symptoms of mania associated with bipolar disorder, such as aggressiveness, agitation, impulsive behaviour or excessively elevated mood.

DIVALPROEX is not indicated for the treatment of the symptoms of mania in patients under 18 years of age.

Please follow your doctor’s recommendations carefully.

**What it does:**

Divalproex sodium has anticonvulsant properties. The mechanism of action has not yet been established. It has been suggested that its activity in epilepsy is related to increased brain concentrations of gammaaminobutyric acid (GABA).

**When it should not be used:**

DIVALPROEX should not be taken by:

- patients with liver disease or significant liver dysfunction
- patients who are allergic to the drug
- patients with known urea cycle disorders (a genetic disorder)

**What the medicinal ingredient is:**

Divalproex sodium

**What the important nonmedicinal ingredients are:**

Eudragit, guar gum, hydroxypropyl cellulose, hydroxypropylmethyl cellulose, microcrystalline cellulose, polyethylene glycol, talc, titanium dioxide, triethyl citrate,

D & C red No. 30 (125 mg tablets only), D & C red No. 33 (500 mg tablets only), red ferric oxide-orange shade (500 mg tablets only) and FD & C yellow No. 6 (125 mg and 250 mg tablets only)

**What dosage forms it comes in:**

DIVALPROEX is available in 125 mg, 250 mg and 500 mg enteric-coated tablets.

**WARNINGS AND PRECAUTIONS**

**Serious Warnings and Precautions**

- **Hepatotoxicity:** liver failure resulting in death has occurred in patients receiving divalproex sodium. These incidents usually occurred during the first 6 months of treatment with divalproex sodium. Patients taking several anticonvulsant drugs, children, those with a history of liver disease, metabolic disorders, severe seizure disorders accompanied by mental retardation, and those with brain disease may be at particular risk. Experience has indicated that children under the age of 2 years are at a considerably increased risk of developing fatal hepatotoxicity, especially those on multiple anticonvulsants.
- **Teratogenicity:** Divalproex sodium can produce birth defects to an unborn baby. Accordingly, the use of DIVALPROEX in women of childbearing potential requires that the benefits of its use be weighed against the risk of injury to the fetus.
- **Pancreatitis:** cases of life threatening pancreas disorder have been reported in both children and adults receiving divalproex sodium. Some cases have occurred shortly after first use as well as after several years of use. Abdominal pain, nausea, vomiting and/or anorexia can be symptoms of pancreatitis that require immediate medical evaluation.

BEFORE you use DIVALPROEX talk to your doctor or pharmacist if:

- you have a history of, or suffer from a liver disease, such as jaundice (yellowing of the skin and eyes);
- you ever had an unusual or allergic reaction to DIVALPROEX (including fever or rash);
- you are allergic to any component of DIVALPROEX tablets;
- you are pregnant or are planning to become pregnant;
- you are breast-feeding (nursing); DIVALPROEX passes into breast milk;
- you are taking any other prescription or over the counter medicine.;
- you have kidney disease;
- you have other medical conditions including a history of unexplained coma, mental retardation or any type of brain dysfunction
- you have a psychiatric disorder or have thoughts of

## IMPORTANT: PLEASE READ

- suicide;
- you consume alcohol on a regular basis.

### Precautions while taking DIVALPROEX:

- Your doctor will monitor your response to DIVALPROEX on a regular basis. However, if your seizures get worse, you should tell your doctor immediately.
- Since DIVALPROEX may cause poor coordination and/or drowsiness, you should not engage in hazardous activities, such as driving and operating machinery, until you know that you don't become drowsy from the drug.
- You should not stop taking your medication unless directed by your doctor. You should always check that you have an adequate supply of DIVALPROEX. You should remember that this medicine was prescribed only for you; it should never be given to anyone else.

## INTERACTIONS WITH THIS MEDICATION

### Serious Drug Interactions

- Rare cases of coma have been reported in patients receiving divalproex sodium alone or when taken with phenobarbital.
- Serious skin reactions (such as conditions called Stevens-Johnson syndrome and Toxic Epidermal Necrolysis) have been reported when divalproex sodium and lamotrigine were taken together.

Drugs that may interact with divalproex sodium include:

- anticonvulsants such as carbamazepine, lamotrigine, primidone, topiramate, felbamate, phenytoin, ethosuximide, phenobarbital;
- anticoagulants such as acetylsalicylic acid, warfarin, dicumarol;
- benzodiazepines such as diazepam, lorazepam, clonazepam;
- some medicines used to treat infections such as rifampin;
- some medicines used to treat diabetes such as tolbutamide;
- some HIV-antiviral medication such as zidovudine;
- any of the group of antibiotics in the carbapenem class such as doripenem, ertapenem, imipenem, meropenem;
- some medicines used to treat heartburn and peptic ulcers such as cimetidine;
- medicines used to treat depression such as Selective Serotonin Re-Uptake Inhibitors (SSRIs), Monoamine Oxidase Inhibitors (MAOIs), Tricyclic antidepressants such as amitriptyline, nortriptyline;
- antipsychotics.

## PROPER USE OF THIS MEDICATION

Please consult your doctor before taking any other medication, including over-the-counter medicines. Some drugs can produce various side effects when they are used in combination with DIVALPROEX.

It is important to keep your appointments for medical checkups.

The doctor may need to take blood tests to measure the amount of divalproex sodium in your blood when adjusting your medications.

### Usual dose:

It is very important to take DIVALPROEX exactly as instructed by your doctor.

The recommended starting dose of DIVALPROEX will be decided by your doctor based on your weight, your seizures or manic episodes and your concomitant medications. Be sure to tell your doctor all the prescription and over the counter medications that you are currently taking. Your doctor will gradually increase the dosage until your condition is well controlled without experiencing side effects. You should carefully follow the instructions that were given to you and not change your dose without consulting with your doctor.

DIVALPROEX may be taken with or without food.

DIVALPROEX is not indicated for the treatment of the symptoms of mania in patients under 18 years of age.

### Overdose:

**In case of drug overdose, contact a health care practitioner, hospital emergency department or regional Poison Control Centre immediately, even if there are no symptoms.**

If you accidentally take an overdose of DIVALPROEX, you should contact your doctor or nearest hospital emergency, or your Regional Poison Control Centre, even though you may not feel sick.

### Missed Dose:

Do not abruptly stop taking your medicine because of the risk of increasing your epileptic seizures. If you miss a dose, you should not try to make up for it by doubling up on your next dose. You should take your next regularly scheduled dose and try not to miss any more doses.

## SIDE EFFECTS AND WHAT TO DO ABOUT THEM

You should check with your doctor or pharmacist right away if you notice any bothersome or unusual effects while taking DIVALPROEX.

Different side effects have been reported by patients taking divalproex sodium. The most commonly reported adverse reactions are nausea, vomiting and indigestion. You should know that this does not mean that you will experience such effects, because people can react in different ways to the same medicine.



**IMPORTANT: PLEASE READ**

**SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM**

Symptom / effect		Talk with your doctor or pharmacist right away		Seek emergency medical attention
		Only if severe	In all cases	
Common	Nausea	*		
	Vomiting	*		
	Indigestion	*		
	Sedation	*		
	Headache	*		
	Diarrhea	*		
Uncommon	Brain dysfunction with high blood ammonia levels (increased lethargy/drowsiness, vomiting, ataxia [abnormal gait, abnormal walking], episodes of extreme irritability†, combative/bizarre behaviour†† and refusal to eat meat or high protein products††)		*	
	Decreased number of platelets in the blood (may result in easy bruising and bleeding from the skin or other areas)		*	
	Liver disorder (not feeling well, develop weakness, lethargy, facial swelling, loss of appetite, yellowing of the skin or eyes, dark urine, and vomiting)		*	
	Pancreas disorder (abdominal pain, nausea, vomiting, and/or loss of appetite)		*	
	Thoughts of suicide or hurting yourself		*	

† In young children  
 †† In older children or adults

*This is not a complete list of side effects. For any unexpected effects while taking DIVALPROEX, contact your doctor or pharmacist.*

**HOW TO STORE IT**

Store at controlled room temperature 15-30° C (59-86°F) in tight, light-resistant containers.

**REPORTING SUSPECTED SIDE EFFECTS**

**You can report any suspected adverse reactions associated with the use of health products to the Canada Vigilance Program by one of the following 3 ways:**

- Report online at [www.healthcanada.gc.ca/medeffect](http://www.healthcanada.gc.ca/medeffect)
- Call toll-free at 1-866-234-2345
- Complete a Canada Vigilance Reporting Form and:
  - Fax toll-free to 1-866-678-6789, or
  - Mail to : Canada Vigilance Program  
Health Canada  
Address Locator: 0701E  
Ottawa, ON K1A 0K9

Postage paid labels, Canada Vigilance Reporting Form and the adverse reaction reporting guidelines are available on the MedEffect™ Canada Web site at [www.healthcanada.gc.ca/medeffect](http://www.healthcanada.gc.ca/medeffect).

***NOTE: Should you require information related to the management of side effects, contact your health professional. The Canada Vigilance Program does not provide medical advice.***

**MORE INFORMATION**

For more information, please contact your doctor, pharmacist or other healthcare professional.

This document plus the full product monograph, prepared for health professionals can be found by contacting the sponsor, Sanis Health Inc. at: 1-866-236-4076, or by email: [quality@sanis.com](mailto:quality@sanis.com).

This leaflet was prepared by

Sanis Health Inc.  
 333 Champlain Street, Suite 102  
 Dieppe, New Brunswick, E1A 1P2  
 Canada

Last revised: June 16, 2014