

PRODUCT MONOGRAPH

^{Pr}VAN-Irbesartan-HCTZ

Irbesartan and Hydrochlorothiazide Tablets

(House Standard)

150/12.5 mg, 300/12.5 mg and 300/25 mg

Angiotensin II AT₁ Receptor Blocker / Diuretic

Manufacturer and Distributor:

Vanc Pharmaceuticals Inc.
Building 152, 11782 River Road
Richmond, BC V6X 1Z7
www.vancpharm.com

Date of Preparation:
September 10, 2014

Submission Control Number: 177492

Table of Contents

PART I: HEALTH PROFESSIONAL INFORMATION	
SUMMARY PRODUCT INFORMATION	3
INDICATIONS AND CLINICAL USE	3
CONTRAINDICATIONS	4
WARNINGS AND PRECAUTIONS	4
ADVERSE REACTIONS	8
DRUG INTERACTION	12
DOSAGE AND ADMINISTRATION	15
OVERDOSAGE	17
ACTION AND CLINICAL PHARMACOLOGY	17
STORAGE AND STABILITY	21
DOSAGE FORMS, COMPOSITION AND PACKAGING	21
PART II: SCIENTIFIC INFORMATION	
PHARMACEUTICAL INFORMATION	22
CLINICAL TRIALS	22
DETAILED PHARMACOLOGY	26
TOXICOLOGY	27
REFERENCES	33
PART III: CONSUMER INFORMATION	
ABOUT THIS MEDICATION	34
WARNINGS AND PRECAUTIONS	34
INTERACTIONS WITH THIS MEDICATION	35
PROPER USE OF THIS MEDICATION	36
SIDE EFFECTS AND WHAT TO DO ABOUT THEM	36
HOW TO STORE IT	37
MORE INFORMATION	37

PrVAN-Irbesartan-HCTZ

Irbesartan and Hydrochlorothiazide Tablets

PART I: HEALTH PROFESSIONAL INFORMATION

SUMMARY PRODUCT INFORMATION

Route of Administration	Dosage Form / Strength	Non medicinal Ingredients
Oral	Tablet / 150/12.5 mg 300/12.5 mg 300/25 mg	<p>VAN-Irbesartan-HCTZ 150/12.5 mg and 300/12.5 mg tablets: lactose monohydrate, microcrystalline cellulose, starch pregelatinized, croscarmellose sodium, povidone, magnesium stearate, hypromellose, titanium dioxide, polyethylene glycol, iron oxide red and iron oxide yellow.</p> <p>VAN-Irbesartan-HCTZ 300/25 mg tablets also contain: lactose monohydrate, microcrystalline cellulose, starch pregelatinized, croscarmellose sodium, povidone, magnesium stearate, hypromellose, titanium dioxide, polyethylene glycol, iron oxide red and iron oxide black.</p>

INDICATIONS AND CLINICAL USE

VAN-Irbesartan-HCTZ is indicated for the treatment of essential hypertension in patients for whom combination therapy is appropriate (see DOSAGE AND ADMINISTRATION).

VAN-Irbesartan-HCTZ is also indicated as initial therapy in patients with severe essential hypertension (Sitting DBP \geq 110 mmHg) for whom the benefit of a prompt blood pressure reduction exceeds the risk of initiating combination therapy in these patients (see CLINICAL TRIALS and DOSAGE AND ADMINISTRATION).

VAN-Irbesartan-HCTZ is not indicated as initial therapy in patients with mild to moderate essential hypertension.

Geriatrics (> 65 years of age): In clinical studies, no overall differences in safety or efficacy were observed between patients over 65 years of age and younger patients (See WARNINGS AND PRECAUTIONS- Special populations).

Pediatrics (< 18 years of age): The safety and efficacy of VAN-Irbesartan-HCTZ in patients <18 years of age have not been established (See WARNINGS AND PRECAUTIONS- Special populations).

CONTRAINDICATIONS

Patients who are hypersensitive to this drug or to any ingredient in the formulation. For a complete listing, see the DOSAGE FORMS AND PACKAGING section of the Product Monograph.

Because of the hydrochlorothiazide component, VAN-Irbesartan-HCTZ is contraindicated in patients with anuria, and in patients who are hypersensitive to other sulfonamide-derived drugs.

WARNINGS AND PRECAUTIONS

Serious Warnings and Precautions

When used in pregnancy, angiotensin receptor (AT₁) blockers (ARB) can cause injury and even death of the developing fetus. When pregnancy is detected, VAN-Irbesartan-HCTZ should be discontinued as soon as possible. (see WARNINGS AND PRECAUTIONS-Special Populations).

General

The effect of irbesartan on ability to drive and use machines has not been studied, but based on its pharmacodynamic properties, irbesartan is unlikely to affect this ability. When driving vehicles or operating machines, it should be taken into account that occasionally dizziness or weariness may occur during treatment of hypertension.

Carcinogenesis and Mutagenesis

See **TOXICOLOGY** for discussion of animal data.

Cardiovascular

Hypotension

Occasionally, symptomatic hypotension has occurred after administration of irbesartan, in some cases after the first dose. It is more likely to occur in patients who are volume depleted by diuretic therapy, dietary salt restriction, dialysis, diarrhea, or vomiting. In these patients, because of the potential fall in blood pressure, therapy should be started under close medical supervision (see DOSAGE AND ADMINISTRATION). Similar considerations apply to patients with ischemic heart or cerebrovascular disease, in whom an excessive fall in blood pressure could result in myocardial infarction or cerebrovascular accident.

Valvular Stenosis

There is concern on theoretical grounds that patients with aortic stenosis might be at particular risk of decreased coronary perfusion when treated with vasodilators because they do not develop as much afterload reduction.

Endocrine and Metabolism

Thiazides, including hydrochlorothiazide, can cause fluid or electrolyte imbalance (hypokalemia, hyponatremia and hypochloremic alkalosis). Periodic determination of

serum electrolytes to detect possible electrolyte imbalance should be performed at appropriate intervals.

Calcium excretion is decreased by thiazides which may cause intermittent and slight elevation of serum calcium. If calcium or a calcium sparing drug (e.g., vitamin D therapy) is prescribed, serum calcium levels should be monitored and calcium dosage adjusted accordingly. Marked hypercalcemia suggests the possibility of hyperparathyroidism. Thiazides should be discontinued before carrying out tests for parathyroid function.

Thiazides have been shown to increase the urinary excretion of magnesium, which may result in hypomagnesemia.

Hyperuricemia may occur, and an acute attack of gout may be precipitated in certain patients receiving thiazide therapy.

Insulin requirements in diabetic patients may be altered and latent diabetes mellitus may become manifest during thiazide diuretic therapy.

Increases in cholesterol and triglyceride levels may be associated with thiazide diuretic therapy.

Thiazides may decrease serum PBI levels without signs of thyroid disturbance.

Hepatic/Biliary/Pancreatic

Thiazides should be used with caution in patients with impaired hepatic function or progressive liver disease, since minor alterations in fluid and electrolyte balance may precipitate hepatic coma.

Immune

Hypersensitivity Reaction

Sensitivity reactions to hydrochlorothiazide may occur in patients with or without a history of allergy or bronchial asthma.

Systemic Lupus Erythematosus

Thiazide diuretics have been reported to cause exacerbation or activation of systemic lupus erythematosus.

Ophthalmic

Acute Myopia and Secondary Acute Angle-Closure Glaucoma

Hydrochlorothiazide, a sulfonamide, can cause an idiosyncratic reaction, resulting in acute transient myopia and acute angle-closure glaucoma. Symptoms include acute onset of

decreased visual acuity or ocular pain and typically occur within hours to weeks of drug initiation. Untreated acute angle-closure glaucoma can lead to permanent vision loss. The primary treatment is to discontinue drug intake as rapidly as possible. Prompt medical or surgical treatments may need to be considered if the intraocular pressure remains uncontrolled. Risk factors for developing acute angle-closure glaucoma may include a history of sulfonamide or penicillin allergy.

Renal

Azotemia

Azotemia may be precipitated or increased by hydrochlorothiazide. Cumulative effects of the drug may develop in patients with impaired renal function.

If increasing azotemia and oliguria occur during treatment of severe progressive renal disease the diuretic should be discontinued.

Renal Impairment

As a consequence of inhibiting the renin-angiotensin-aldosterone system, changes in renal function have been seen in susceptible individuals. In patients whose renal function may depend on the activity of the renin-angiotensin-aldosterone system, such as patients with bilateral renal artery stenosis, unilateral renal artery stenosis to a solitary kidney, or severe congestive heart failure, treatment with agents that inhibit this system has been associated with oliguria, progressive azotemia, and rarely, acute renal failure and/or death. In susceptible patients, concomitant diuretic use may further increase risk.

Use of irbesartan should include appropriate assessment of renal function.

Thiazides should be used with caution.

Because of the hydrochlorothiazide component, irbesartan/ hydrochlorothiazide is not recommended in patients with severe renal impairment (creatinine clearance \leq 30 mL/min).

Special Populations

Pregnant Women

Drugs that act directly on the renin-angiotensin system (RAAS) can cause fetal and neonatal morbidity and death when administered to pregnant women. When pregnancy is detected, irbesartan and hydrochlorothiazide tablets should be discontinued as soon as possible.

The use of ARB is not recommended during pregnancy. Epidemiological evidence regarding the risk of teratogenicity following exposure to angiotensin converting enzyme inhibitors (another class of therapeutic products interfering with the RAAS) during the first trimester of pregnancy has not been conclusive; however a small increase in risk cannot be excluded. Given the current evidence available on the risk with ARB, similar risks may

exist for this class of drugs. Patients planning pregnancy should be changed to alternative anti-hypertensive treatments which have an established safety profile for use in pregnancy. When pregnancy is diagnosed, treatment with angiotensin II antagonists should be stopped immediately, and, if appropriate, alternative therapy should be started.

The use of ARBs during the second and third trimesters is known to induce human fetotoxicity (decreased renal function, oligohydramnios, skull ossification, retardation) and neonatal toxicity (renal failure, hypotension, hyperkalemia).

Infants with histories of *in utero* exposure to ARBs should be closely observed for hypotension, oliguria, and hyperkalemia. If oliguria occurs, attention should be directed toward support of blood pressure and renal perfusion. Exchange transfusion may be required as means of reversing hypotension and/or substituting for impaired renal function; however, limited experience with those procedures has not been associated with significant clinical benefit. Irbesartan is not removed by hemodialysis.

Thiazides cross the placental barrier and appear in cord blood. The routine use of diuretics in otherwise healthy pregnant women is not recommended and exposes mother and fetus to unnecessary hazard, including fetal or neonatal jaundice, thrombocytopenia, and possibly other adverse reactions that have occurred in adults. Diuretics do not prevent development of toxemia of pregnancy and there is no satisfactory evidence that they are useful in the treatment of toxemia.

Nursing Women

It is not known whether irbesartan is excreted in human milk, but significant levels of radioactivity have been found in the milk of lactating rats. Thiazides appear in human milk. Thiazides in high doses causing intense diuresis can inhibit the milk production. Because many drugs are excreted in human milk and because of their potential for affecting the nursing infant adversely, a decision should be made whether to discontinue nursing or discontinue the drug, taking into account the importance of the drug to the mother.

Pediatrics

Safety and effectiveness have not been established.

Geriatrics

Of the 2650 hypertensive patients receiving VAN-Irbesartan-HCTZ in clinical studies, 618 patients were 65 years of age and over. No overall age-related differences were seen in the adverse effect profile but greater sensitivity in some older individuals cannot be ruled out.

ADVERSE REACTIONS

Adverse Drug Reaction Overview

VAN-Irbesartan-HCTZ has been evaluated for safety in 2746 patients with essential hypertension including 968 patients for 1 year or more.

The most commonly reported adverse events (occurring in $\geq 10\%$ of patients treated with VAN-Irbesartan-HCTZ) was headache (11.0%), which occurred at a significantly higher incidence in the placebo group (16.1%).

The adverse events most frequently resulting in clinical intervention (discontinuation of VAN-Irbesartan-HCTZ) were due to dizziness (0.7%) and headache (0.7%). The adverse event of hypotension is more likely to occur in volume depleted patients (See Warnings and Precautions related to Cardiovascular under Hypotension).

The following potentially serious adverse reactions have been reported rarely with irbesartan in controlled clinical trials: syncope, hypotension.

Clinical Trial Adverse Drug Reactions

Because clinical trials are conducted under very specific conditions the adverse reaction rates observed in the clinical trials may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse drug reaction information from clinical trials is useful for identifying drug-related adverse events and for approximating rates.

Hypertension

In placebo-controlled clinical trials, therapy was discontinued due to a clinical or laboratory adverse event in 3.6 percent of patients treated with irbesartan/hydrochlorothiazide, versus 6.8 percent of patients given placebo.

Adverse events regardless of drug relationship, occurring in $\geq 1\%$ of the irbesartan/hydrochlorothiazide patients in placebo-controlled clinical trials include the following:

Table 1 : Adverse events regardless of drug relationship, occurring in $\geq 1\%$ of the irbesartan/hydrochlorothiazide patients in placebo-controlled clinical trials

	VAN-Irbesartan-HCTZ n = 898 (%)	Irbesartan n = 400 (%)	HCTZ n = 380 (%)	Placebo n = 236 (%)
<i>Cardiovascular</i>				
Edema	3.1	1.5	1.6	2.5
Tachycardia	1.2	0.5	0.5	0.4
<i>Dermatologic</i>				
Rash	1.2	1.8	3.2	1.7
<i>Gastrointestinal</i>				

	VAN-Irbesartan- HCTZ n = 898 (%)	Irbesartan n = 400 (%)	HCTZ n = 380 (%)	Placebo n = 236 (%)
Nausea/vomiting	3.2	1.5	2.4	0.4
Dyspepsia	2.1	0.3	1.6	0.8
Diarrhea	2.1	2.8	1.1	3.4
Abdominal Pain	1.7	1.5	1.6	0.8
<i>General</i>				
Fatigue	6.5	4.0	3.2	3.0
Influenza	2.8	2.0	1.8	1.3
Chest Pain	1.8	1.5	1.6	1.3
<i>Immunology</i>				
Allergy	1.1	0.5	0.5	0
<i>Musculoskeletal</i>				
Musculoskeletal Pain	6.5	6.0	9.7	4.7
Muscle Cramp	1.0	0.8	2.1	1.3
<i>Nervous System</i>				
Headache	11.0	9.3	11.6	16.1
Dizziness	7.6	5.5	4.7	4.2
Orthostatic Dizziness	1.1	1.0	0.8	0.4
Anxiety /Nervousness	1.0	1.0	0.5	1.7
<i>Renal/Genitourinary</i>				
Urination abnormal	1.9	0.5	2.1	0.8
Urinary Tract Infection	1.6	1.5	2.4	2.5
<i>Respiratory</i>				
URTI	5.6	8.3	7.1	5.5
Sinus disorder	2.9	4.5	3.2	4.7
Cough	2.2	2.3	2.6	3.0
Pharyngitis	2.1	2.3	2.9	1.7
Rhinitis	1.9	2.0	1.6	2.5

Severe Hypertension

In a clinical study in patients with severe hypertension (SeDBP \geq 110 mmHg), the overall pattern of adverse events reported through seven weeks of follow-up was similar in patients treated with VAN-Irbesartan-HCTZ as initial therapy and in patients treated with irbesartan as initial therapy.

Table 2: Most common adverse events regardless of drug relationship, occurring in \geq 1% of the irbesartan/ hydrochlorothiazide patients with Severe Hypertension in controlled clinical trial

	Number (%) of Subjects VAN- Irbesartan-HCTZ N = 468	Number (%) of Subjects Irbesartan N= 227
Headache	19 (4.1)	15 (6.6)
Dizziness	16 (3.4)	9 (4.0)
Nasopharyngitis	8 (1.7)	10 (4.4)
Bronchitis	6 (1.3)	6 (2.6)
Fatigue	6 (1.3)	1 (0.4)
Upper Respiratory Tract Infection	6 (1.3)	4 (1.8)
Erectile Dysfunction	5 (1.1)	0
Nausea	5 (1.1)	5 (2.2)
Diarrhaea	4 (0.9)	3 (1.3)

	Number (%) of Subjects VAN-Irbesartan-HCTZ N = 468	Number (%) of Subjects Irbesartan N= 227
Sinusitis	4 (0.9)	3 (1.3)
Cough	3 (0.6)	4 (1.8)
Muscle Spasms	2 (0.4)	3 (1.3)

The incidence of the pre-specified adverse events was: 0% reported cases of syncope in either treatment group; 0.6% and 0% cases of hypotension, 3.6% and 4.0% cases of dizziness, 4.3% and 6.6% cases of headache, 0.2% and 0% cases of hyperkalemia, and 0.6% and 0.4% cases of hypokalemia reported in the group treated with VAN-Irbesartan-HCTZ and the group treated with irbesartan, respectively.

The rate of discontinuation due to adverse events was 1.9% and 2.2% in the group treated with ibesartan/hydrochlorothiazide and the group treated with irbesartan, respectively.

Irbesartan Alone

In addition, the following potentially important events occurred in less than 1% of patients receiving irbesartan, regardless of drug relationship:

Body as a Whole: fever;

Cardiovascular: flushing, hypertension, myocardial infarction, angina pectoris, arrhythmic/conduction disorder, cardio-respiratory arrest, heart failure, hypertensive crisis;

Dermatologic: pruritus, dermatitis, ecchymosis, erythema, urticaria, photosensitivity;

Endocrine: sexual dysfunction, libido disorder, gout;

Gastrointestinal: constipation, gastroenteritis, flatulence, abdominal distention, hepatitis;

Musculoskeletal: muscle cramp, arthritis, myalgia, muscular weakness;

Nervous System: sleep disturbance, numbness, somnolence, vertigo, depression, paresthesia, tremor, transient ischemic attack, cerebrovascular accident.

Renal/Genitourinary: abnormal urination;

Respiratory: epistaxis, tracheobronchitis, pulmonary congestion, dyspnea, wheezing;

Special Senses: visual disturbance, hearing impaired, conjunctivitis, taste disturbance.

Abnormal Hematologic and Clinical Chemistry Findings

VAN-Irbesartan-HCTZ

Liver Function Tests: Occasional elevations of liver enzymes and/or serum bilirubin have occurred. In patients with essential hypertension treated with irbesartan/hydrochlorothiazide alone, one patient was discontinued due to elevated liver enzymes.

Creatinine, Blood Urea Nitrogen: Minor increases in blood urea nitrogen (BUN) or serum creatinine were observed in 2.3% of patient. No patient was discontinued due to increased BUN. One patient was discontinued due to a minor increase in serum creatinine.

IRBESARTAN

Liver Function Tests: In placebo-controlled trials, elevations of AST and ALT $\geq 3X$ upper limit of normal occurred in 0.1% and 0.2%, respectively, of irbesartan treated patients compared to 0.3% and 0.3%, respectively, of patients receiving placebo. The cumulative incidence of AST and/or ALT elevations $\geq 3X$ upper limit of normal was 0.4% in patients treated with irbesartan for a mean duration of over 1 year.

Hyperkalemia: In placebo-controlled trials, greater than a 10% increase in serum potassium was observed in 0.4% of irbesartan treated patients compared to 0.5% of patients receiving placebo.

Creatinine, Blood Urea Nitrogen: Minor increases in blood urea nitrogen (BUN) or serum creatinine were observed in less than 0.7% of patients with essential hypertension treated with irbesartan alone versus 0.9% on placebo.

Hemoglobin: Mean decreases in hemoglobin of 0.16g/dL were observed in patients receiving irbesartan. No patients were discontinued due to anemia.

Neutropenia: Neutropenia (<1000 cells/mm³) was observed in 0.3% of irbesartan treated patients compared to 0.5% of patients receiving placebo.

In clinical trials, the following were noted to occur with an incidence of $< 1\%$, regardless of drug relationship: anemia, thrombocytopenia, lymphocytopenia, and increased CPK.

Post-Market Adverse Drug Reactions

Angioedema (involving swelling of the face, lips, and/or tongue) has been reported rarely in post marketing use. The following adverse reactions, regardless of drug relationship, were reported in post-marketing use: syncope, asthenia, jaundice, myalgia, elevated liver function tests, and impaired renal function including cases of renal failure in patients at risk (see WARNINGS AND PRECAUTIONS – Renal - Renal Impairment).

Cases of muscle pain, muscle weakness, myositis and rhabdomyolysis have been reported in patients receiving angiotensin II receptor blockers.

Other clinical adverse events (regardless of relationship to drug) reported with the use of hydrochlorothiazide alone include: pancreatitis and toxic epidermal necrolysis.

DRUG INTERACTIONS

Drug-Drug Interactions

Table 3- Established or Potential Drug-Drug Interactions

Proper Name	Ref.	Effect	Clinical comment
Alcohol, barbiturates, or narcotics	C	Potiation of orthostatic hypotension may occur.	Avoid alcohol, barbiturates or narcotics, especially with initiation of therapy.
Agents increasing Serum Potassium	RCS	Since irbesartan decreases the production of aldosterone, potassium-sparing diuretics or potassium supplements should be given only for documented hypokalemia and with frequent monitoring of serum potassium. Potassium-containing salt substitutes should also be used with caution. Concomitant thiazide diuretic use may attenuate any effect that irbesartan may have on serum potassium.	
Amphotericin B	T	Amphotericin B increases the risk of hypokalemia induced by thiazide diuretics	Monitor serum potassium level.
Antidiabetic agents (e.g. insulin and oral hypoglycemic agents)	CT	Thiazide-induced hyperglycemia may compromise blood sugar control. Depletion of serum potassium augments glucose intolerance.	Monitor glycemic control, supplement potassium if necessary, to maintain appropriate serum potassium levels, and adjust diabetes medications as required.
Antihypertensive drugs	CT	Hydrochlorothiazide may potentiate the action of other antihypertensive drugs (e.g. guanethidine, methyldopa, beta-blockers, vasodilators, calcium channel blockers, diuretics, ACEI, ARB, and direct renin inhibitors).	Thiazides, including hydrochlorothiazide, can cause fluid or electrolyte imbalance (hypokalemia, hyponatremia and hypochloremic alkalosis). Periodic determination of serum electrolytes to detect possible electrolyte imbalance should be performed at appropriate intervals.
Antineoplastic drugs, including cyclophosphamide and methotrexate	C	Concomitant use of thiazide diuretics may reduce renal excretion of cytotoxic agents and enhance their myelosuppressive effects.	Hematological status should be closely monitored in patients receiving this combination. Dose adjustment of cytotoxic agents may be required.
Bile acid sequestrants, e.g., cholestyramine	CT	Bile acid sequestrants bind thiazide diuretics in the gut and impair gastrointestinal absorption by 43-85%. Administration of thiazide 4 hours after a bile acid sequestrant reduced absorption of hydrochlorothiazide by 30-35%.	Give thiazide 2-4 hours before or 6 hours after the bile acid sequestrant. Maintain a consistent sequence of administration. Monitor blood pressure, and increase dose of thiazide, if necessary.
Calcium and vitamin D supplements	C	Thiazides decrease renal excretion of calcium and increase calcium release from bone.	Monitor serum calcium, especially with concomitant use of high doses of calcium

Proper Name	Ref.	Effect	Clinical comment
			supplements. Dose reduction or withdrawal of calcium and/or vitamin D supplements may be necessary.
Carbamazepine	C	Carbamazepine may cause clinically significant hyponatremia. Concomitant use with thiazide diuretics may potentiate hyponatremia.	Monitor serum sodium levels. Use with caution. If possible, another class of diuretics should be used.
Corticosteroids, and adrenocorticotropic hormone (ACTH)	T	Intensified electrolyte depletion, particularly hypokalemia, may occur	Monitor serum potassium, and adjust medications, as required.
Digoxin	CT	When irbesartan was administered as 150 mg once daily under steady-state conditions, no effect was seen on the pharmacokinetics of digoxin at steady-state. Thiazide-induced electrolyte disturbances, i.e. hypokalemia, hypomagnesemia, increase the risk of digoxin toxicity, which may lead to fatal arrhythmic events.	Concomitant administration of hydrochlorothiazide and digoxin requires caution. Monitor electrolytes and digoxin levels closely. Supplement potassium or adjust doses of digoxin or thiazide, as required.
Drugs that alter GI motility, i.e., anticholinergic agents, such as atropine and prokinetic agents, such as metoclopramide, domperidone	CT,T	Bioavailability of thiazide diuretics may be increased by anticholinergic agents due to a decrease in gastrointestinal motility and gastric emptying. Conversely, prokinetic drugs may decrease the bioavailability of thiazide diuretics.	Dose adjustment of thiazide may be required.
Gout medications (allopurinol, uricosurics, xanthine oxidase inhibitors)	T, RCS	Thiazide-induced hyperuricemia may compromise control of gout by allopurinol and probenecid. The co-administration of hydrochlorothiazide and allopurinol may increase the incidence of hypersensitivity reactions to allopurinol.	Dosage adjustment of gout medications may be required.
Lithium	CT	Thiazide diuretics reduce the renal clearance of lithium and add a high risk of lithium toxicity. As with other drugs which eliminate sodium, lithium clearance may be reduced in the presence of irbesartan.	Concomitant use of thiazide diuretics with lithium is generally not recommended. If such use is deemed necessary, reduce lithium dose by 50% and monitor lithium levels closely. Serum lithium levels should be monitored carefully if lithium salts are to be administered with irbesartan.
Nifedipine	CT	The pharmacokinetics of irbesartan were not affected by coadministration of nifedipine.	
Nonsteroidal anti-inflammatory drugs (NSAID) (including selective COX-2 inhibitors)	CT	In patients who are elderly, volume-depleted (including those on diuretic therapy), or with compromised renal function, coadministration of NSAIDs, including selective COX-2 inhibitors, with angiotensin II	Monitor renal function periodically in patients receiving irbesartan and NSAID therapy.

Proper Name	Ref.	Effect	Clinical comment
		<p>receptor antagonists, including irbesartan, may result in deterioration of renal function, including possible acute renal failure. These effects are usually reversible. The antihypertensive effect of Angiotensin II receptor antagonists, including irbesartan, may be attenuated by NSAIDs including selective COX-2 inhibitors.</p> <p>In some patients, the administration of a non-steroidal anti-inflammatory agent can reduce the diuretic, natriuretic, and antihypertensive effects of loop, potassium-sparing and thiazide diuretics.</p>	When irbesartan/hydrochlorothiazide and non-steroidal anti-inflammatory agents are used concomitantly, the patient should be observed closely to determine if the desired effect of the diuretic is obtained.
Pressor Amines (e.g., Norepinephrine)	CT	In the presence of diuretics, possible decreased response to pressor amines may be seen but not sufficient to preclude their use.	
Selective serotonin reuptake inhibitors (SSRIs, e.g., citalopram, escitalopram, sertraline)	T, C	Concomitant use with thiazide diuretics may potentiate hyponatremia.	Monitor serum sodium levels. Use with caution.
Simvastatin	CT	When irbesartan was administered in a small single-dose study with 12 young, healthy males aged 19 to 39, the single-dose pharmacokinetics of simvastatin were not affected by the concomitant administration of 300 mg irbesartan. Simvastatin values were highly variable whether simvastatin was administered alone or in combination with irbesartan.	
Skeletal muscle relaxants of the curare family, e.g., tubocurare	C	Thiazide drugs may increase the responsiveness of some skeletal muscle relaxants, such as curare derivatives	
Topiramate	CT	Additive hypokalemia. Possible thiazide-induced increase in topiramate serum concentrations.	Monitor serum potassium and topiramate levels. Use potassium supplements, or adjust topiramate dose as necessary.
Warfarin	CT	When irbesartan was administered as 300 mg once daily under steady-state conditions, no pharmacodynamic effect on PT was demonstrated in subjects stabilized on warfarin.	

Legend: C = Case Study; RCS = Retrospective Cohort Study; CT = Clinical Trial; T = Theoretical

Drug-Food Interactions

No statistically significant effects of food were observed on the C_{max} , $AUC_{(inf)}$ or $T_{1/2}$ of irbesartan or on the $AUC_{(inf)}$ or $T_{1/2}$ of hydrochlorothiazide. In the fasted and fed states, T_{max} increased from 1 to 2 hours for irbesartan and from 1.5 to 3.5 hours for hydrochlorothiazide. The C_{max} for hydrochlorothiazide decreased 21% in the fed state relative to the fasted state. None of these changes were considered to be clinically significant.

Drug-Herb Interactions

There have been no clinical studies to assess the possible interaction of any herbal products and VAN-Irbesartan-HCTZ.

DOSAGE AND ADMINISTRATION

Dosing Considerations

- Dosage must be individualized.
- The fixed combination is not for initial therapy except for severe hypertension.
- The dose of VAN-Irbesartan-HCTZ should be determined by the titration of the individual components.
- Use of VAN-Irbesartan-HCTZ in with patients with liver impairment is not advisable.
- Dosage adjustment may be required in hemodialysis patients (see Recommended Dose and Dosage Adjustment - Adjustment in Renal Insufficiency).

Recommended Dose and Dosage Adjustment

Once the patient has been stabilized on the individual components as described below, either one tablet of VAN-Irbesartan-HCTZ 150/12.5 mg, 300/12.5 mg or 300/25 mg once daily may be substituted if the doses on which the patient was stabilized are the same as those in the fixed combination.

VAN-Irbesartan-HCTZ may be administered with or without food, however it should be taken consistently with respect to food intake.

Irbesartan Monotherapy

The recommended dose of irbesartan is 150 mg once daily. In patients whose blood pressure is not adequately controlled, the daily dose may be increased to 300 mg.

SEVERE HYPERTENSION (Sitting DBP \geq 110 mmHg)

The starting dose of VAN-Irbesartan-HCTZ for initial treatment of severe hypertension is one tablet of VAN-Irbesartan-HCTZ 150/12.5 mg once daily (see INDICATIONS AND CLINICAL USE and CLINICAL TRIALS: Severe Hypertension). The dosage may be increased after two to four weeks of therapy to a maximum of one 300/25 mg tablet once daily. VAN-Irbesartan-HCTZ is not recommended as initial therapy in patients with intravascular volume depletion (**see WARNINGS AND PRECAUTIONS: Hypotension**).

DOSE ADJUSTMENT IN SPECIAL POPULATION

Diuretic Treated Patients

In patients receiving diuretics, irbesartan therapy should be initiated with caution, since these patients may be volume-depleted and thus more likely to experience hypotension following initiation of additional antihypertensive therapy. Whenever possible, all diuretics should be discontinued two to three days prior to the administration of irbesartan to reduce the likelihood of hypotension (see WARNINGS AND PRECAUTIONS – Cardiovascular: Hypotension, and DRUG INTERACTIONS). If this is not possible because of the patient's condition, irbesartan should be administered with caution and the blood pressure monitored closely. The recommended starting dose of irbesartan is 75 mg once daily in hypovolemic patients (see WARNINGS AND PRECAUTIONS – Cardiovascular: Hypotension). Thereafter, the dosage should be adjusted according to the individual response of the patient.

Geriatric

No initial dosage adjustment in irbesartan is necessary for most elderly patients. Appropriate caution should nevertheless be used when prescribing to the elderly, as increased vulnerability to drug effect is possible in this patient population. [See WARNINGS AND PRECAUTIONS -Geriatrics].

Hepatic Insufficiency

No initial dosage adjustment in irbesartan is generally necessary in patients with mild to moderate hepatic impairment. Since thiazide diuretics may precipitate hepatic coma, the use of a fixed combination product such as VAN-Irbesartan-HCTZ is not advisable.

Renal Insufficiency

No initial dosage adjustment in irbesartan is generally necessary in patients with renal impairment, although due to the apparent greater sensitivity of hemodialysis patients, an initial dose of 75 mg is recommended in this group of patients.

The usual regimens of therapy with VAN-Irbesartan-HCTZ may be followed as long as the patient's creatinine clearance is > 30mL/min. In patients with more severe renal impairment, loop diuretics are preferred to thiazides so VAN-Irbesartan-HCTZ is not recommended.

Missed Dose

Patients should be instructed to take VAN-Irbesartan-HCTZ tablet at the next scheduled dose and not take two doses at the same time if they miss a dose.

OVERDOSAGE

For management of a suspected drug overdose, contact your regional Poison Control Centre.
--

No specific information is available on the treatment of overdosage with VAN-Irbesartan-HCTZ. The patient should be closely monitored, and the treatment should be symptomatic and supportive, including fluid and electrolyte replacement.

Irbesartan

No data or very little data available in regard to overdosage in humans.

The most likely manifestations of overdosage would be hypotension and/or tachycardia; bradycardia might also occur in this setting. Irbesartan is not removed by hemodialysis.

Hydrochlorothiazide

The most common signs and symptoms observed are those caused by electrolyte depletion (hypokalemia, hypochloremia, hyponatremia) and dehydration resulting from excessive diuresis. If digitalis has also been administered, hypokalemia may accentuate cardiac arrhythmias.

The degree to which hydrochlorothiazide is removed by hemodialysis has not been established.

ACTION AND CLINICAL PHARMACOLOGY

Mechanism of Action

VAN-Irbesartan-HCTZ combines the actions of irbesartan, an angiotensin II AT₁ receptor blocker, and that of a thiazide diuretic, hydrochlorothiazide.

Irbesartan

Irbesartan antagonizes angiotensin II by blocking AT₁ receptors.

Angiotensin II is the primary vasoactive hormone in the renin-angiotensin system. Its effects include vasoconstriction and the stimulation of aldosterone secretion by the adrenal cortex.

Irbesartan blocks the vasoconstrictor and aldosterone-secreting effects of angiotensin II by selectively blocking in a non competitive manner the binding of angiotensin II to the AT₁ receptor found in many tissues. Irbesartan has no agonist activity at the AT₁ receptor. AT₂ receptors have been found in many tissues, but to date they have not been associated with cardiovascular homeostasis. Irbesartan has essentially no affinity for the AT₂ receptors.

Irbesartan does not inhibit angiotensin converting enzyme, also known as kininase II, the enzyme that converts angiotensin I to angiotensin II and degrades bradykinin, nor does it affect renin or other hormone receptors or ion channels involved in cardiovascular regulation of blood pressure and sodium homeostasis.

Hydrochlorothiazide

Hydrochlorothiazide is a thiazide diuretic. Thiazides affect the renal tubular mechanisms of electrolyte reabsorption, directly increasing excretion of sodium and chloride in approximately equivalent amounts. Indirectly, the diuretic action of hydrochlorothiazide reduces plasma volume, with consequent increases in plasma renin activity, increases in aldosterone secretion, increases in urinary potassium loss, and decreases in serum potassium. The renin-aldosterone link is mediated by angiotensin II, so coadministration of an angiotensin II AT₁ receptor blocker tends to reverse the potassium loss associated with these diuretics.

The mechanism of the antihypertensive effect of thiazides is not fully understood.

Pharmacodynamics

Irbesartan

In healthy subjects, single oral doses of irbesartan up to 300 mg produced dose-dependent inhibition of the pressor effect of angiotensin II infusions. The inhibition was complete (100%) 4 hours following oral doses of 150 mg or 300 mg. Partial inhibition of 40% and 60% was still present 24 hours post-dose with 150 mg and 300 mg irbesartan respectively.

In hypertensive patients, angiotensin II receptor inhibition following chronic administration of irbesartan causes a 1.5-2 fold rise in angiotensin II plasma concentration and a 2-3 fold increase in plasma renin levels. Aldosterone plasma concentrations generally decline following irbesartan administration, however serum potassium levels are not significantly affected at recommended doses.

During clinical trials, minimal incremental blood pressure response was observed at doses greater than 300 mg.

The blood pressure lowering effect of irbesartan is apparent after the first dose and substantially present within 1-2 weeks, with the maximal effect occurring by 4-6 weeks. In long-term studies, the effect of irbesartan appeared to be maintained for more than one year. There was essentially no change in average heart rate in patients treated with irbesartan in controlled trials.

There is no rebound effect after withdrawal of irbesartan.

Black hypertensive patients had a smaller blood pressure response to irbesartan monotherapy than caucasians.

Hydrochlorothiazide

Onset of the diuretic action following oral administration occurs in 2 hours and the peak action in about 4 hours. Diuretic activity lasts about 6 to 12 hours.

VAN-Irbesartan-HCTZ

The components of VAN-Irbesartan-HCTZ have been shown to have an additive effect on blood pressure reduction, reducing blood pressure to a greater degree than either component alone.

Pharmacokinetics

Table 4: Pharmacokinetic Parameters for Irbesartan

Irbesartan	T _{max} (h)	t _{1/2} (h)	Clearance (mL/minute)	Volume of distribution (L)
Single dose mean	1.5-2	11-15	Plasma 157 -176 Renal 3.0 – 3.5	53 -93

Table 5: Pharmacokinetic Parameters for hydrochlorothiazide

Hydrochlorothiazide	T _{max} (h)	t _{1/2} (h)	Clearance (mL/minute)	Volume of distribution (L/kg)
Single dose mean	1.5-2	5-15	Plasma 192-343 Predominantly Renal (unchanged)	1.5-4.2

Irbesartan

Absorption

Irbesartan is an orally active agent. The oral absorption of irbesartan is rapid and complete with an average absolute bioavailability of 60% - 80%. Following oral administration, peak plasma concentrations are attained at 1.5-2 hours after dosing. Steady-state concentrations are achieved within 3 days.

Distribution

The average volume of distribution of irbesartan is 53-93 litres.

Irbesartan is approximately 96% protein-bound in the plasma, primarily to albumin and α_1 -acid glycoprotein.

Metabolism

Irbesartan is metabolized via glucuronide conjugation, and oxidation by the cytochrome P-450 system.

Excretion

Irbesartan and its metabolites are excreted by both biliary and renal routes. Following either oral or intravenous administration of ¹⁴C-labeled irbesartan, about 20% of radioactivity is recovered in the urine and the remainder in the feces. Less than 2% of the dose is excreted in urine as unchanged irbesartan. Irbesartan exhibits linear pharmacokinetics over the therapeutic dose range with an average terminal elimination half-life of 11-15 hours.

Total plasma and renal clearances are in the range of 157 - 176 and 3.0 - 3.5 mL/minute, respectively.

Hydrochlorothiazide

Absorption

Hydrochlorothiazide is rapidly absorbed from the gastrointestinal tract. The bioavailability is approximately 65 to 70%.

Distribution

Hydrochlorothiazide crosses the placental but not the blood-brain barrier and is excreted in breast milk.

Metabolism

Hydrochlorothiazide is not metabolized.

Excretion

Hydrochlorothiazide is eliminated rapidly by the kidney. At least 61% of the oral dose is eliminated unchanged within 24 hours. The plasma half-life has been observed to vary from 5.6 and 14.8 hours.

Special Populations and Conditions

Geriatrics

In subjects over the age of 65 years, irbesartan elimination half-life was not significantly altered, but AUC and C_{MAX} values were about 20 - 50% greater than those of young subjects.

Renal Insufficiency

The mean AUC and C_{max} of irbesartan were not altered in patients with any degree of renal impairment, including patients on hemodialysis. However, a wide variance was seen in patients with severe renal impairment.

Hepatic Insufficiency

The pharmacokinetics of irbesartan following repeated oral administration were not significantly affected in patients with mild to moderate cirrhosis of the liver. No data is available in patients with severe liver disease.

STORAGE AND STABILITY

Irbesartan and hydrochlorothiazide tablets can be stored between 15°C and 30°C.

DOSAGE FORMS, COMPOSITION AND PACKAGING

VAN-Irbesartan-HCTZ 150/12.5 mg tablets are peach coloured, biconvex, oval shaped, film coated tablets.

VAN-Irbesartan-HCTZ 300/12.5 mg tablets are peach coloured, biconvex, oval shaped, film coated tablets.

In addition to the active ingredients, irbesartan and hydrochlorothiazide, each tablet contains: lactose monohydrate, microcrystalline cellulose, starch pregelatinized, croscarmellose sodium, povidone, magnesium stearate, hypromellose, titanium dioxide, polyethylene glycol, iron oxide red and iron oxide yellow for the two strengths.

VAN-Irbesartan-HCTZ 300/25 mg tablets are dark pink coloured, biconvex, oval shaped, film coated tablets.

In addition to the active ingredients, irbesartan and hydrochlorothiazide, each tablet contains: lactose monohydrate, microcrystalline cellulose, starch pregelatinized, croscarmellose sodium, povidone, magnesium stearate, hypromellose, titanium dioxide and polyethylene glycol, iron oxide red and iron oxide black.



VAN-Irbesartan-HCTZ 150/12.5 mg, 300/12.5 mg and 300/25 mg tablets are available in bottles of 100 tablets.

PART II: SCIENTIFIC INFORMATION

PHARMACEUTICAL INFORMATION

Drug Substance

Proper Name: irbesartan/hydrochlorothiazide

	Irbesartan	Hydrochlorothiazide
Chemical Name	2-Butyl-3-[[[2'-(1 <i>H</i> -tetrazol-5-yl) biphenyl-4-yl] methyl]-1,3-diazaspiro[4.4] non-1-en-4-one.	6-chloro-3,4-dihydro-2 <i>H</i> -1,2,4-benzothiadiazine-7- sulphonamide-1,1-dioxide.
Molecular Formula	C ₂₅ H ₂₈ N ₆ O	C ₇ H ₈ ClN ₃ O ₄ S ₂
Structural Formula		
Molecular Weight	428.53 g/mol	297.7 g/mol
Physicochemical Properties	Irbesartan is a white or almost white, crystalline powder. Irbesartan is sparingly soluble in methanol and slightly soluble in methylene chloride, practically insoluble in water.	Hydrochlorothiazide is a white or almost white, crystalline powder. It is very slightly soluble in water, soluble in acetone, sparingly soluble in ethanol (96%). It dissolves in dilute solutions of alkali hydroxides.

CLINICAL TRIALS

A randomized, double blind, two-treatment, two-sequence, two-period, single-dose, crossover, bioequivalence study comparing Irbesartan and Hydrochlorothiazide 300/25 mg Tablets (Vanc Pharmaceuticals Inc.) with AVALIDE[®] (irbesartan and hydrochlorothiazide) 300/25mg Tablets (sanofi-aventis Canada Inc.) in 29 healthy adult, human subjects, under fasting conditions. A summary of the bioavailability data for Irbesartan and Hydrochlorothiazide is presented below.

SUMMARY TABLE OF THE COMPARATIVE BIOAVAILABILITY DATA

VAN-Irbesartan-HCTZ 1 x (300 mg/25 mg) VAN-Irbesartan-HCTZ From measured data Geometric Mean Arithmetic Mean (CV %)				
Parameter	Test*	Reference [#]	% Ratio of Geometric Means	90% Confidence Interval
AUC _T (ng*hr/mL)	23776.11 25197.67 (34.78)	22697.08 23773.99 (34.09)	104.8	96.7 – 113.5
AUC _I (ng*hr/mL)	24453.62 25922.32 (35.42)	23218.79 24308.79 (33.96)	105.3	97.1 – 114.2
C _{max} (ng/mL)	3521.25 3694.32 (30.88)	3588.21 3693.86 (23.63)	98.1	90.7 – 106.1
T _{max} [§] (h)	2.10 (57.05)	2.35 (59.52)		

VAN-Irbesartan-HCTZ 1 x (300 mg/25 mg) VAN-Irbesartan-HCTZ From measured data Geometric Mean Arithmetic Mean (CV %)				
Parameter	Test*	Reference [#]	% Ratio of Geometric Means	90% Confidence Interval
T _{1/2} [§] (h)	9.31 (33.58)	8.49 (35.06)		

* Irbesartan (300 mg) [Vanc Pharmaceuticals Inc.]

[#] AVALIDE[®] Irbesartan (300 mg) [sanofi-aventis Canada Inc.] were purchased in Canada.

[§] Expressed as the arithmetic mean (CV%) only

SUMMARY TABLE OF THE COMPARATIVE BIOAVAILABILITY DATA

HYDROCHLOROTHIAZIDE 1 x (300 mg/25 mg) VAN-Irbesartan-HCTZ From measured data Geometric Mean Arithmetic Mean (CV %)				
Parameter	Test *	Reference [#]	% Ratio of Geometric Means	90% Confidence Interval
AUC _T (ng*hr/mL)	1029.95 1074.27 (29.39)	1010.64 1052.85 (27.34)	101.9	93.4 – 111.2
AUC _I (ng*hr/mL)	1081.59 1123.06 (28.00)	1066.01 1102.97 (25.55)	101.5	93.8 -109.7
C _{max} (ng/mL)	164.33 172.12 (30.73)	155.41 162.95 (30.74)	105.7	96.3 – 116.1
T _{max} [§] (h)	2.14 (37.32)	2.30 (41.19)		
T _{1/2} [§] (h)	9.46 (10.28)	9.73 (17.96)		

* Hydrochlorothiazide (25 mg) [Vanc Pharmaceuticals Inc.]

[#] AVALIDE[®] (Hydrochlorothiazide 25 mg) [sanofi-aventis Canada Inc.] were purchased in Canada.

[§] Expressed as the arithmetic mean (CV%) only

Irbesartan-Hydrochlorothiazide

The antihypertensive effects of irbesartan and hydrochlorothiazide tablets were examined in 4 placebo-controlled studies of 8–12 weeks in patients with mild-moderate hypertension. These trials included 1914 patients randomized to fixed doses of irbesartan (37.5 to 300 mg) and concomitant hydrochlorothiazide (6.25 to 25 mg). One factorial study compared all combinations of irbesartan (37.5, 100 and 300 mg or placebo) and hydrochlorothiazide (6.25, 12.5, and 25 mg or placebo). The irbesartan-hydrochlorothiazide combinations of 75/12.5 mg and 150/12.5 mg were compared to their individual components and placebo in a separate study. A third study investigated the ambulatory blood pressure responses to irbesartan-hydrochlorothiazide (75/12.5 mg and 150/12.5 mg) and placebo after 8 weeks of dosing. Another trial investigated the effects of the addition of irbesartan (75 mg) in patients not controlled on hydrochlorothiazide (25 mg) alone.

In controlled trials, the addition of irbesartan 150–300 mg to hydrochlorothiazide doses of 6.25, 12.5 or 25 mg produced further dose-related reductions in blood pressure of 8–10/3–6 mmHg, comparable to those achieved with the same monotherapy dose of irbesartan. The

addition of hydrochlorothiazide to irbesartan produced further dose related reductions in blood pressure at trough (24 hours post-dose) of 5–6/2–3 mmHg (12.5 mg) and 7–11/4–5 mmHg (25 mg), also comparable to effects achieved with hydrochlorothiazide alone. Once-daily dosing with 150 mg irbesartan and 12.5 mg hydrochlorothiazide, 300 mg irbesartan and 12.5 mg hydrochlorothiazide, or 300 mg irbesartan and 25 mg hydrochlorothiazide produced mean placebo-adjusted blood pressure reductions at trough (24 hours post-dosing) of about 13–15/7–9, 14/9–12, and 19–21/11–12 mmHg, respectively. Peak effects occurred at 3–6 hours, with the trough-to peak ratios >65%.

In another study, irbesartan (75–150 mg) or placebo was added on a background of 25 mg hydrochlorothiazide in patients not adequately controlled (SeDBP 93–120 mmHg) on hydrochlorothiazide (25 mg) alone. The addition of irbesartan (75–150 mg) gave an additive effect (systolic/diastolic) at trough (24 hours post-dosing) of 11/7 mmHg.

There was no difference in response for men and women or in patients over or under 65 years of age. Black patients had a larger response to hydrochlorothiazide than non-black patients and a smaller response to irbesartan. The overall response to the combination was similar for black and non-black patients.

Severe Hypertension

The efficacy of irbesartan and hydrochlorothiazide as initial therapy for severe hypertension (defined as a mean seated diastolic blood pressure (SeDBP) \geq 110 mmHg confirmed on two separate occasions off all antihypertensive therapy) was studied in a seven-week, double-blind, randomized, multicenter study. Patients were randomized to either irbesartan and hydrochlorothiazide (150/12.5 mg) or to irbesartan (150 mg) once daily and followed for blood pressure response. These initial study regimens were increased at one week to irbesartan 300 mg/HCTZ 25 mg or to irbesartan 300 mg, respectively. The primary endpoint was a comparison at five weeks of the proportion of patients who achieved trough SeDBP <90 mmHg. An additional supportive endpoint compared the proportion of subjects in each treatment group whose blood pressure was controlled, defined as simultaneous SeDBP <90 mmHg and seated systolic blood pressure (SeSBP) <140 mmHg.

Study demographics and trial design

The study randomized 697 patients, in a 2:1 ratio to receive either combination therapy (irbesartan plus HCTZ, N=468) or irbesartan monotherapy (N=229), and included 296 (42%) females, 101 (14%) blacks, and 92 (13%) \geq 65 years of age. The mean age was 52 years. The mean blood pressure at baseline for the total population was 172/113 mmHg.

Table 6- Summary of patient demographics for clinical trial with irbesartan and hydrochlorothiazide in subjects with severe hypertension

Study #	Trial design	Dosage, route of administration and duration	Study subjects (n=number)	Mean age in years (Range)	Gender
CV131176	Multicenter, randomized, double-blind, active controlled, 7-week, parallel	Oral administration of irbesartan /HCTZ 150/12.5 mg or irbesartan 150 mg increased at one week to irbesartan /HCTZ 300/25	697 Irbesartan: 229 Irbesartan/HCTZ: 468	52.5 (23.0-83.0)	Male 57.5% Female 42.5%

Study #	Trial design	Dosage, route of administration and duration	Study subjects (n=number)	Mean age in years (Range)	Gender
	group study	mg or to irbesartan 300 mg.			

Study results

The study results are summarized in table 7.

After five weeks of therapy, the mean SeDBP was 4.7 mmHg lower ($p \leq 0.0001$) and the mean SeSBP was 9.7 mmHg lower ($p < 0.0001$) in the group treated with VAN-Irbesartan-HCTZ than in the group treated with irbesartan. Mean reductions from baseline for SeDBP and SeSBP at trough were 24.0 mmHg and 30.8 mmHg for VAN-Irbesartan-HCTZ-treated patients and 19.3 mmHg and 21.1 mmHg for irbesartan-treated patients, respectively. A greater proportion of the patients on VAN-Irbesartan-HCTZ achieved a diastolic blood pressure < 90 mmHg (47.2% for VAN-Irbesartan-HCTZ, 33.2% for irbesartan; $p = 0.0005$) and a greater proportion of the patients on VAN-Irbesartan-HCTZ achieved simultaneous control of SeSBP < 140 mmHg and SeDBP < 90 mmHg (34.6% versus 19.2%; $p < 0.0001$). Similar results were seen when the patients were grouped according to gender, race or age (< 65 years, ≥ 65 years). The proportions of subjects with controlled SeDBP, as well as with simultaneous SeDBP/SeSBP control, at each week of the double-blind period were consistently larger and statistically significantly greater for VAN-Irbesartan-HCTZ-treated patients than for irbesartan-treated patients.

Table 7- Results at week 5 of study with VAN-Irbesartan-HCTZ in subjects with severe hypertension

Endpoints	irbesartan /HCTZ 150/12.5 mg force titrated to 300/25 mg	Irbesartan 150 mg force titrated to 300 mg	p value								
Primary Endpoint: Proportion of subjects in each treatment group whose SeDBP was controlled (SeDBP < 90 mmHg).	47.2%	33.2%	0.0005								
Other Endpoints:											
<ul style="list-style-type: none"> Proportion of subjects whose BP was controlled (simultaneous SeDBP < 90 mmHg and SeSBP < 140 mmHg) Mean changes from baseline in trough <table border="0" style="margin-left: 20px;"> <tr> <td>SeDBP</td> <td>-24.0</td> <td>-19.3</td> <td>< 0.0001</td> </tr> <tr> <td>SeSBP</td> <td>-30.8</td> <td>-21.1</td> <td>< 0.0001</td> </tr> </table> 	SeDBP	-24.0	-19.3	< 0.0001	SeSBP	-30.8	-21.1	< 0.0001	34.6%	19.2%	< 0.0001
SeDBP	-24.0	-19.3	< 0.0001								
SeSBP	-30.8	-21.1	< 0.0001								

DETAILED PHARMACOLOGY

Pharmacodynamics

Irbesartan

In healthy subjects, single oral doses of irbesartan up to 300 mg produced dose-dependent inhibition of the pressor effect of angiotensin II infusions. The inhibition was complete (100%) 4 hours following oral doses of 150 mg or 300 mg. Partial inhibition of 40% and 60% was still present 24 hours post-dose with 150 mg and 300 mg irbesartan respectively.

In hypertensive patients, angiotensin II receptor inhibition following chronic administration of irbesartan causes a 1.5-2 fold rise in angiotensin II plasma concentration and a 2-3 fold increase in plasma renin levels. Aldosterone plasma concentrations generally decline following irbesartan administration, however serum potassium levels are not significantly affected at recommended doses.

During clinical trials, minimal incremental blood pressure response was observed at doses greater than 300 mg.

The blood pressure lowering effect of irbesartan is apparent after the first dose and substantially present within 1-2 weeks, with the maximal effect occurring by 4-6 weeks. In long-term studies, the effect of irbesartan appeared to be maintained for more than one year. There was essentially no change in average heart rate in patients treated with irbesartan in controlled trials.

There is no rebound effect after withdrawal of irbesartan.

Black hypertensive patients had a smaller blood pressure response to irbesartan monotherapy than caucasians.

The blood pressure lowering effect of irbesartan in combination with hydrochlorothiazide was apparent after the first dose and substantially present within 1-2 weeks, with the maximal effect occurring by 6-8 weeks. In long-term follow-up studies, the effect of VAN-Irbesartan-HCTZ was maintained for over one year.

There was no significant difference in blood pressure response based on age or gender.

Pharmacokinetics

Irbesartan

Following either oral or intravenous administration of ¹⁴C-labeled irbesartan, more than 80% of the circulating plasma radioactivity is attributable to unchanged irbesartan. The primary circulating metabolite is the inactive irbesartan glucuronide (approximately 6%). The remaining oxidative metabolites do not add appreciably to the pharmacologic activity.

In vitro studies of irbesartan indicate that the oxidation of irbesartan is primarily by cytochrome P-450 isoenzyme CYP 2C9. Metabolism of irbesartan by CYP 3A4 is negligible. Irbesartan is neither metabolized, nor does it substantially induce or inhibit the following isoenzymes: CYP 1A1, 1A2, 2A6, 2B6, 2D6, 2E1. There was no induction or inhibition of CYP 3A4.

The mean AUC and C_{MAX} were not altered in patients with any degree of renal impairment, including patients on hemodialysis. However, a wide variance was seen in patients with severe renal impairment.

The pharmacokinetics of irbesartan following repeated oral administration were not significantly affected in patients with mild to moderate cirrhosis of the liver. No data is available in patients with severe liver disease.

Hydrochlorothiazide

Hydrochlorothiazide is not metabolized but is eliminated rapidly by the kidney. The plasma half-life has been observed to vary between 5.6 and 14.8 hours when the plasma levels can be followed for at least 24 hours. At least 61 percent of the oral dose is eliminated unchanged within 24 hours. Hydrochlorothiazide crosses the placental but not the blood-brain barrier and is excreted in breast milk.

TOXICOLOGY

Acute Toxicity

Irbesartan

Table 8: Acute Toxicity for Irbesartan

Species	Sex (N)	Route	LD ₅₀ (mg/kg)
Mouse	M (5) F (5)	PO	> 2000
Rat	M (5) F (5)	PO	> 2000
Mouse	M (5) F (5)	IV	> 50
Rat	M (5) F (5)	IV	> 50
Mouse	M (5) F (5)	IP	200 - 2000
Rat	M (5) F (5)	IP	200 - 2000

After single administration, toxicity was slight and no target organ was identified. Very few toxic effects, characterized by pilo-erection and/or somnolence were noted at 2000 mg/kg by the oral route, 200 mg/kg by the intraperitoneal route and 50 mg/kg by the intravenous route. Acute oral toxicity studies with irbesartan in mice and rats indicated acute lethal doses were in excess of 2000 mg/kg, about 25 - 50 fold the maximum human dose (300 mg) on a mg/m² basis, respectively.

Irbesartan - hydrochlorothiazide

Table 9: Acute Toxicity for Irbesartan – hydrochlorothiazide

Species	Sex (N)	Route	LD ₅₀ (mg/kg)		
			Irbesartan	HCTZ	Irbesartan/ HCTZ
Mouse	M (5) F (5)	PO	> 2000	> 4000	> 2000/4000
Rat	M (5)	PO	> 3000	> 500	> 3000/500

No mortality occurred following administration of the VAN-Irbesartan-HCTZ combination up to and including the highest dose of irbesartan:hydrochlorothiazide (2000/4000 mg/kg

in mice or 3000/500 mg/kg in rats). No treatment-related clinical signs and body weight changes were observed. At necropsy, performed at the end of the 14-day observation period, pathologic examinations did not reveal any treatment-induced changes.

Subacute and Chronic Toxicity

Irbesartan

Table 10: Subacute and Chronic Toxicity Irbesartan

Species/ Strain	Sex (N/Dose)	Dose (mg/kg/day)	Route	Time	Effects
SUBACUTE TOXICITY					
Rat	M (10) F (10)	0, 30, 70, 150	PO	4 weeks	<ul style="list-style-type: none"> Irbesartan only induced slight decrease in hemoglobin levels (at 150 mg/kg) and slight increase in glucose (≥ 30 mg/kg), urea (≥ 70 mg/kg), creatinine and K⁺ levels (at 150 mg/kg), and slight decrease in Na⁺ and Cl⁻ urinary concentrations and excretions (≥ 30 mg/kg).
Rat	M (10) F (10)	0, 0.8, 2, 5	IV	16 days	<ul style="list-style-type: none"> Very slight increase in Na⁺ and Cl⁻ plasma levels (≥ 0.8 mg/kg/day in males) Very slight increase in K⁺ plasma levels, in ASAT and slight decrease in kidney relative weight at 5 mg/kg/day in males.
Monkey	M (3) F (3)	0, 10, 30, 90	PO	4 weeks	<ul style="list-style-type: none"> Dose-related hyperplasia of the juxtaglomerular apparatus (from 30 mg/kg/day upwards).
Monkey	M (3) F (3)	0, 250, 500, 1000	PO	4 weeks	<ul style="list-style-type: none"> ≥ 250 mg/kg/day: changes in the kidney (hyperplasia of the juxtaglomerular apparatus), heart (myocardial fibrosis) and erythrocytes parameters (slight anemia). At 500 mg/kg/day: increased platelet count, fibrogen and neutrophil levels and at 1000 mg/kg/day, health deterioration were also noted. One animal receiving 250 mg/kg/day presented the most severe heart lesions and marked electrocardiographic modifications on D1 and D29. However, pre-existing lesions could not be excluded.
Monkey	M (3) F (3)	0, 0.8, 2, 5	IV	2 weeks	<ul style="list-style-type: none"> Irbesartan induced only a slight hyperplasia of the juxtaglomerular apparatus in 2/3 females receiving 5 mg/kg/day. One high-dose animal presented a marked heart hypertrophy with marked ECG changes on D1 and D10 suggesting that it was a preexisting lesion.
Rat	M (20) - F (20) [main study]	0, 10, 30, 90	PO	26 weeks	<ul style="list-style-type: none"> Slight reduction of the body weight gain in males at 90 mg/kg/day (- 6 to - 8%).

Species/ Strain	Sex (N/Dose)	Dose (mg/kg/day)	Route	Time	Effects
	M (10) - F (10) [reversibility study for control and high dose groups] M (5) - F (5) [toxicokinetics study]				<ul style="list-style-type: none"> Other changes can be considered to be of pharmacological origin for some of them and have no clear toxicological significance for all of them. The no-observed adverse effect dose was considered to be 30 mg/kg/day.
CHRONIC TOXICITY					
Rat	M (20) - F (20) [main study] M (10) - F (10) [reversibility study for control and high dose groups] M (5) - F (5) [toxicokinetics study]	0, 0, 250, 500, 1000	PO	26 weeks	<ul style="list-style-type: none"> Slight reduction of body weight gain without any dose-relationship reversible. Changes in hematology and blood biochemistry parameters demonstrating an effect on red blood cells and on the renal function, likely associated with the pharmacological activity of irbesartan and reversible. Hyperplasia/hypertrophy of the juxtaglomerular apparatus in males (≥ 250 mg/kg/day) and in females (≥ 500 mg/kg/day) partially reversible.
Monkey	M (5) - F (5) [main study] M (3) - F (3) [reversibility study for control and high dose groups]	0, 10, 30, 90	PO	6 months	<ul style="list-style-type: none"> Dose-related hyperplasia of juxtaglomerular apparatus in all treated animals partially reversible at the end of treatment. Slight dose-related decrease in weight gain from the 30 mg/kg/day dose level upwards and slight anemia from 10 mg/kg/day upwards, both reversible on cessation of treatment.
Monkey	M (5) F (5)	0, 20, 100, 500	PO	52 weeks	<ul style="list-style-type: none"> Irbesartan was well tolerated and most of the changes observed were considered to be due to the pharmacological activity of the drug: Dose-related decrease in blood pressure at doses ≥ 20 mg/kg/day associated with necrosis of the tip of the tail likely due to a decrease in blood flow at 500 mg/kg/day. Dose-related hyperplasia / hypertrophy of the juxtaglomerular apparatus in all treated animals with degenerative kidney changes at 500 mg/kg/day. Slight decrease in body weight gain and erythrocyte parameters at doses ≥ 100 mg/kg/day.

After repeated oral administrations at dose levels up to 1000 mg/kg per day, most of the treatment-related effects noted in all species are linked to the pharmacological activity of irbesartan. The kidney can be considered as the primary target organ:

hyperplasia/hypertrophy of the juxtaglomerular apparatus which was observed in all species, is a direct consequence of the interaction with the renin-angiotensin system. Irbesartan also induced some hematology (slight decrease in erythrocyte parameters) and blood biochemistry variations (slight increase in urea, creatinine, phosphorus, potassium and calcium levels) likely due to a disturbance in the renal blood flow, and a slight decrease in heart weight which could result from a decrease in cardiac work load due to decreased peripheral vascular resistance. At high doses (> 500 mg/kg per day),

degenerative changes of the kidney were noted which could be secondary to prolonged hypotensive effects.

Subacute and Chronic Toxicity (Cont'd)

VAN-Irbesartan-HCTZ

Table 11: Subacute and Chronic Toxicity VAN-Irbesartan-HCTZ

Species/ Strain	Sex (N/Dose)	Dose (mg/kg/day)	Route	Time	Effects
Rat	M (20) F (20)	0*/0**, 10/10, 90/90, 90/0, 0/90	PO	6 months	<ul style="list-style-type: none"> Exposure to HCTZ was greater when administered in combination with irbesartan, than when given alone. Body weight gains in the high dose group (90/90 mg/kg) were slightly decreased in females and moderately decreased in males. Hemoglobin, hematocrit, and erythrocyte counts were slightly decreased in females given the high dose combination (90/90). In the high dose combination, serum urea nitrogen and alkaline phosphatase (males) were slightly elevated; serum potassium and calcium (Week 12) were slightly decreased in males; serum cholesterol and triglycerides were slightly to moderately decreased In the low dose combination, serum cholesterol, triglycerides and potassium were slightly decreased. Slight increases in urine pH; urine protein concentrations markedly lower in high dose combination group. Decreased heart weights in males and females at 10/10, 99/90 and 90/0. Decreased liver weights in males. Juxtaglomerular-cell hypertrophy/hyperplasia. Increased urine output. Increased kidney weights in females. At necropsy, discoloration of the glandular stomach correlated with focal coagulative necrosis or ulceration of the mucosa were noted in all treated groups with an incidence slightly greater in rats given the high-dose combination.
Monkey	M (20) F (20)	0*/0**, 10/10, 90/90, 0/90, 90/0	PO	6 months	<ul style="list-style-type: none"> Exposure to HCTZ was approximately 60% greater when administered in combination with irbesartan than when administered alone. Body weights of males in the high dose combination group (90/90) were mildly decreased. Mean hemoglobin, hematocrit and erythrocyte values were mildly to moderately decreased at the high dose combination (90/90). Moderate increases in BUN; mild to moderate increases in creatinine values; mean sodium, potassium, and chloride values were

Species/ Strain	Sex (N/Dose)	Dose (mg/kg/day)	Route	Time	Effects
					mildly to moderately decreased. • Mild to moderate juxtaglomerular apparatus, hypertrophy/hyperplasia [all treated with irbesartan either alone or in combination]

* Irbesartan

** Hydrochlorothiazide

Reproduction and Teratology

Irbesartan

Fertility and reproductive performance were not affected in studies of male and female rats even at oral doses of irbesartan causing pronounced toxicity (up to 650 mg/kg/day). No significant effects on the number of corpora lutea, implants, or live fetuses were observed. Irbesartan did not affect survival, development, or reproduction of offspring except for a slight decrease of body weight gain during lactation which was reversible after weaning.

In a study of rats receiving maternally toxic doses of irbesartan (650 mg/kg/day), transient effects were observed in fetuses. These effects included increased incidences of renal pelvic cavitation at doses ≥ 50 mg/kg/day and subcutaneous edema at doses ≥ 180 mg/kg/day. Slight decreases in body weight gain were noted (prior to weaning) in offspring of females receiving irbesartan at doses ≥ 50 mg/kg/day. In rabbits, maternally toxic doses of irbesartan (30 mg/kg/day) were associated with maternal mortality and abortion. Surviving females receiving this dose had a slight increase in early resorption. However, no teratogenic effect was observed. Radioactivity was present in the rat and rabbit fetus during late gestation and in rat milk following oral doses of radiolabeled irbesartan. These findings are attributed to drug exposure in late gestation and during lactation.

VAN-Irbesartan-HCTZ

In a Segment II teratology study carried out in rats, a dose of the combination VAN-Irbesartan-HCTZ up to 150/150mg/day/kg did not show any teratogenic potential. There was decreased foetal body weight in the litters of dams given 150/150 mg/kg/day.

Carcinogenicity and Mutagenicity

Irbesartan

No evidence of carcinogenicity was observed when irbesartan was administered at doses of up to 500/1000 mg/kg/day (males/females, respectively) in rats and 1000 mg/kg/day in mice for 2 years. These doses provided systemic exposures of 3.6 - 24.9 times (rats) and 3.8 - 6.2 times (mice) the exposures in humans receiving 300 mg daily.

Irbesartan was not mutagenic in a battery of *in vitro* tests (Ames microbial test, rat hepatocyte DNA repair test, V79 mammalian cell forward gene mutation assay). Irbesartan was negative in several tests for induction of chromosomal aberrations (*in vitro* - human lymphocyte assay; *in vivo* - mouse micronucleus study).

VAN-Irbesartan-HCTZ

No carcinogenicity studies have been conducted with the VAN-Irbesartan-HCTZ combination.

VAN-Irbesartan-HCTZ was not mutagenic in standard *in vitro* tests (Ames microbial test and Chinese hamster mammalian-cell forward gene-mutation assay). VAN-Irbesartan-HCTZ was negative in tests for induction of chromosomal aberrations (*in vitro* - human lymphocyte assay; *in vivo* - mouse micronucleus study).

REFERENCES

1. Cazaubon C, Gougat J, Bousquet F, Guiraudou P, Gayraud R, Lacour C, et al. Pharmacological characterization of SR 47436, a new nonpeptide AT₁ subtype angiotensin II receptor antagonist. *J Pharmacol Exp Ther* 1993; 265:826-834
2. Haggmann M, Burnier M, Nussberger J, Leenhardt AF, Brouard R, Waeber B, et al. Natriuretic and hormonal effects of SR 47436 (BMS 186295), a new angiotensin II receptor antagonist in normotensive volunteers. *Am J Hyperten* 1994; 7 (4,Pt.2): 13A
3. Marino MR, Langenbacher KM, Raymond RH, Whigan D, and Ford NF Pharmacokinetics (pK) and antihypertensive effects of irbesartan (an AII receptor antagonist) in subjects with hypertension. *J Hyperten* 1996; 14(1): S348
4. Martinez F, Schmitt F, Savoie C, Leenhardt AF, Brouard R, Peronnet P, et al. Effect of SR 47436 (BMS 186295) on renal hemodynamics and on glomerular permselectivity in healthy humans. *J Am Soc Nephrol* 1994; 5: 607
5. McIntyre M, McFadyen RJ, Meredith PA, Bruard R, and Reid JL Dose-ranging study of the angiotensin II receptor antagonist irbesartan (SR 47436 / BMS 186295) on blood pressure and neurohormonal effects in salt-deplete men. *J Cardiovas Pharmacol* 1996; 28: 101-106
6. Ribstein J, Sissmann J, Picard A, Bouroudian M, and Mimran A. Effects of the angiotensin II antagonist SR 47436 (BMS 186295) on the pressor response to exogenous angiotensin II and the renin-angiotensin system in sodium replete normal subjects. *J Hypertens* 1994; 12:131
7. Sissmann J, Bouroudian M, Armagnac C, Donazollo Y, Latreille M, and Panis R. Angiotensin II blockade in healthy volunteers: Tolerability and impact on renin angiotensin system components of single and repeated doses of a new angiotensin II receptor antagonist SR 47436 (BMS 186295). *J Hypertens* 1994; 12: S92
8. Pool JL *et al.* Dose-Related Antihypertensive Effects of Irbesartan in Patients with Mild-to-Moderate Hypertension. *American Journal of Hypertension*, 1998, 11: 462-470.
9. Ribstein J, Picard A, Armagnac C, Sissman J and Mimran A. Inhibition of the Acute Effects of Angiotensin II by the Receptor Antagonist Irbesartan in Normotensive Men. *Journal of Cardiovascular Pharmacology*, 2001, 37:449-460.
10. Pr AVALIDE® Product Monograph, sanofi-aventis Canada, Inc. Submission Control No. 159804. Last Revised: February 4, 2013.

PART III: CONSUMER INFORMATION

Pr VAN-Irbesartan-HCTZ
 Irbesartan and Hydrochlorothiazide Tablets

Read this carefully before you start taking VAN-Irbesartan-HCTZ tablets and each time you get a refill. This leaflet is a summary and will not tell you everything about VAN-Irbesartan-HCTZ tablets. Talk to your doctor, nurse, or pharmacist about your medical condition and treatment and ask if there is any new information about VAN-Irbesartan-HCTZ tablets.

ABOUT THIS MEDICATION

What the medication is used for:

VAN-Irbesartan-HCTZ lowers high blood pressure.

What it does:

Irbesartan and hydrochlorothiazide tablet is a combination of 2 drugs, irbesartan and hydrochlorothiazide:

- Irbesartan is an angiotensin receptor blocker (ARB). You can recognize an ARB because its medicinal ingredient ends in “SARTAN”. It lowers blood pressure.
- Hydrochlorothiazide is a diuretic or “water pill” that increases urination. This lowers blood pressure.

This medicine does not cure high blood pressure. It helps to control it. Therefore, it is important to continue taking irbesartan and hydrochlorothiazide tablets regularly even if you feel fine.

When it should not be used:

Do not take VAN-Irbesartan-HCTZ if you:

- Are allergic to irbesartan or hydrochlorothiazide or to any non-medicinal ingredients in the formulation.
- Are allergic to any sulfonamide-derived drugs (sulfa drugs); most of them have a medicinal ingredient that ends in “MIDE”.
- Have experienced an allergic reaction (angioedema) with swelling of the hands, feet, or ankles, face, lips, tongue, throat, or sudden difficulty breathing or swallowing to any ARB (any drug in the same class as irbesartan). Be sure to tell your doctor, nurse, or pharmacist that this has happened to you.
- Have difficulty urinating or produce no urine.
- Are pregnant or plan to become pregnant. Taking VAN-Irbesartan-HCTZ during pregnancy can cause injury and even death to your baby.

- Are breast-feeding. VAN-Irbesartan-HCTZ passes into breast milk.
- Have one of the following rare hereditary diseases:
 - Galactose intolerance
 - Lapp lactase deficiency
 - Glucose-galactose malabsorption

Because lactose is a non-medicinal ingredient in irbesartan and hydrochlorothiazide tablets.

What the medicinal ingredients are:

Irbesartan and hydrochlorothiazide.

What the nonmedicinal ingredients are:

Irbesartan and hydrochlorothiazide 150/12.5 mg and 300/12.5 mg tablets contain: lactose monohydrate, microcrystalline cellulose, starch pregelatinized, croscarmellose sodium, povidone, magnesium stearate, hypromellose, titanium dioxide, polyethylene glycol, iron oxide red and iron oxide yellow.

Irbesartan and hydrochlorothiazide 300/25 mg tablet contains: lactose monohydrate, microcrystalline cellulose, starch pregelatinized, croscarmellose sodium, povidone, magnesium stearate, hypromellose, titanium dioxide, polyethylene glycol, iron oxide red and iron oxide black.

What dosage forms it comes in:

Tablets in three strengths:

VAN-Irbesartan-HCTZ: 150 mg/ 12.5 mg

VAN-Irbesartan-HCTZ: 300 mg/ 12.5 mg

VAN-Irbesartan-HCTZ: 300 mg/ 25 mg

WARNINGS AND PRECAUTIONS

Serious Warnings and Precautions - Pregnancy

VAN-Irbesartan-HCTZ should not be used during pregnancy. If you discover that you are pregnant while taking irbesartan/hydrochlorothiazide stop the medication and contact your doctor, nurse, or pharmacist as soon as possible.

BEFORE you use VAN-Irbesartan-HCTZ tablets tell your doctor, nurse, or pharmacist if you:

- Are allergic to any drug used to lower blood pressure, including angiotensin converting enzyme (ACE) inhibitors, or penicillin.
- Have narrowing of an artery or a heart valve.
- Have heart failure.
- Have diabetes, liver or kidney disease.
- Have gout, or lupus erythematosus.
- Are on dialysis.

- Are dehydrated or suffer from excessive vomiting, diarrhea or sweating.
- Are taking a salt substitute that contains potassium, potassium supplements, or a potassium-sparing diuretic (a specific kind of “water pill”).
- Are on a low-salt diet.
- Are less than 18 years old.
- Have had a heart attack or stroke.

Hydrochlorothiazide in VAN-Irbesartan-HCTZ tablets can cause Sudden Eye Disorders:

- **Myopia:** sudden nearsightedness or blurred vision.
- **Glaucoma:** an increased pressure in your eyes, eye pain. Untreated, it may lead to permanent vision loss.

These eye disorders are related and can develop within hours to weeks of starting irbesartan/hydrochlorothiazide.

You may become sensitive to the sun while taking VAN-Irbesartan-HCTZ. Exposure to sunlight should be minimized until you know how you respond.

Before surgery and general anesthesia (even at the dentist’s office), tell the doctor or dentist that you are taking VAN-Irbesartan-HCTZ, as there may be a sudden fall in blood pressure associated with general anesthesia.

Driving and using machines: Before you perform tasks which may require special attention, wait until you know how you respond to VAN-Irbesartan-HCTZ. Dizziness, lightheadedness, or fainting can especially occur after the first dose and when the dose is increased.

INTERACTIONS WITH THIS MEDICATION

As with most medicines, interactions with other drugs are possible. Tell your doctor, nurse, or pharmacist about all the medicines you take, including drugs prescribed by other doctors, vitamins, minerals, natural supplements or alternative medicines.

The following may interact with VAN-Irbesartan-HCTZ:

- Adrenocorticotrophic hormone (ACTH) used to treat West Syndrome.
- Alcohol, barbiturates (sleeping pills), or narcotics (strong pain medications). They may cause low blood pressure and dizziness

when you go from lying or sitting to standing up.

- Amphotericin B, an antifungal drug.
- Anticancer drugs, including cyclophosphamide and methotrexate.
- Antidepressants, in particular selective serotonin reuptake inhibitors (SSRIs), including citalopram, escitalopram, and sertraline.
- Antidiabetic drugs, including insulin and oral medicines.
- Bile acid resins used to lower cholesterol.
- Other blood pressure lowering drugs. When taken in combination with VAN-Irbesartan-HCTZ, they may cause excessively low blood pressure.
- Calcium or vitamin D supplements.
- Corticosteroids used to treat joint pain and swelling and other conditions.
- Digoxin, a heart medication.
- Drugs that slow down or speed up bowel function, including atropine, metoclopramide, and domperidone.
- Drugs used to treat epilepsy, including carbamazepine and topiramate.
- Gout medications, including allopurinol and probenecid.
- Lithium used to treat bipolar disease.
- Nonsteroidal anti-inflammatory drugs (NSAIDs), used to reduce pain and swelling. Examples include ibuprofen, naproxen, and celecoxib.
- Potassium-sparing diuretics, potassium supplements or potassium-containing salt substitutes.
- Pressor amines such as norepinephrine.
- Skeletal muscle relaxants used to relieve muscle spasms, including tubocurare.
- Warfarin.

PROPER USE OF THIS MEDICATION

Take VAN-Irbesartan-HCTZ tablets every day exactly as prescribed. It is recommended to take your dose at about the same time every day.

VAN-Irbesartan-HCTZ can be taken with or without food but it should be taken the same way each day. If VAN-Irbesartan-HCTZ causes upset stomach, take it with food or milk.

Usual Adult dose:

Patients should be individually titrated for each component separately.

Usual maintenance dose is: 1 tablet daily.

Overdose:

If you think you have taken too much VAN-Irbesartan-HCTZ, contact your doctor, nurse, pharmacist, hospital emergency department or regional Poison Control Centre immediately, even if there are no symptoms.

Missed Dose:

If you have forgotten to take your dose during the day, carry on with the next one at the usual time. Do not double dose.

SIDE EFFECTS AND WHAT TO DO ABOUT THEM

Side effects may include:

- back or leg pain, muscle cramps, spasms and pain, weakness, restlessness
- dizziness, pins and needles in your fingers, headache and fatigue
- constipation, diarrhea, nausea, vomiting, decreased appetite, upset stomach, enlargement of the glands in your mouth, bleeding under the skin, rash, red patches on the skin
- drowsiness, insomnia
- reduced libido

If any of these affects you severely, tell your doctor, nurse or pharmacist.

VAN-Irbesartan-HCTZ can cause abnormal blood test results. Your doctor will decide when to perform blood tests and will interpret the results.

SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM

Symptom / effect		Talk with your doctor, nurse or pharmacist		Stop taking drug and seek immediate medical help
		Only if Severe	In all cases	
Common	Low Blood Pressure: dizziness, fainting, lightheadedness	√		
	May occur when you go from lying or sitting to standing up.			
	Edema: swelling of hands, ankles or feet		√	

SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM

Symptom / effect		Talk with your doctor, nurse or pharmacist		Stop taking drug and seek immediate medical help
		Only if Severe	In all cases	
	Tachycardia: increased heart beats		√	
Uncommon	Decreased or increased levels of potassium in the blood: irregular heartbeats, muscle weakness and generally feeling unwell		√	
	Allergic Reaction: rash, hives, swelling of the face, lips, tongue or throat, difficulty swallowing or breathing			√
	Kidney Disorder: change in frequency of urination, nausea, vomiting, swelling of extremities, fatigue		√	
	Liver Disorder: yellowing of the skin or eyes, dark urine, abdominal pain, nausea, vomiting, loss of appetite		√	
	Increased blood sugar: frequent urination, thirst, and hunger	√		
	Electrolyte Imbalance: weakness, drowsiness, muscle pain or cramps, irregular heartbeat		√	
Rare	Rhabdomyolysis: muscle pain that you cannot explain, muscle tenderness or weakness, dark brown urine		√	
	Decreased White		√	

SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM

Symptom / effect	Talk with your doctor, nurse or pharmacist	Stop taking drug and seek immediate medical help	
		Only if Severe	In all cases
Blood Cells: infections, fatigue, fever, aches, pains, and flu-like symptoms			
Decreased Platelets: bruising, bleeding, fatigue and weakness		√	
Very rare Toxic Epidermal Necrolysis: severe skin peeling, especially in mouth and eyes			√
Unknown Eye disorders: - Myopia: sudden near sightedness or blurred vision - Glaucoma: increased pressure in your eyes, eye pain			√
Anemia: fatigue, loss of energy, weakness, shortness of breath.		√	
Inflammation of the Pancreas: abdominal pain that lasts and gets worse when you lie down, nausea, vomiting		√	

This is not a complete list of side effects. For any unexpected effects while taking VAN-Irbesartan-HCTZ, contact your doctor, nurse or pharmacist.

HOW TO STORE IT

Store at room temperature (15°C to 30°C).

Keep out of the reach and sight of children.

REPORTING SUSPECTED SIDE EFFECTS

You can report any suspected adverse reactions associated with the use of health products to the Canada Vigilance Program by one of the following 3 ways:

Report online at

www.healthcanada.gc.ca/medeffect

Call toll-free at: 1-866-234-2345

Complete a Canada Vigilance Reporting Form and:

- Fax toll-free to 1-866-678-6789, or
- Mail to: Canada Vigilance Program
Health Canada
Postal Locator 0701E
Ottawa, ON K1A 0K9

Postage paid labels, Canada Vigilance Reporting Form and the adverse reaction reporting guidelines are available on the MedEffect™ Canada Web site at www.healthcanada.gc.ca/medeffect.

NOTE: Should you require information related to the management of the side effect, contact your health professional. The Canada Vigilance Program does not provide medical advice.

MORE INFORMATION

This document plus the full product monograph prepared for health professional can be obtained by contacting the distributor, Vanc Pharmaceuticals Inc. at 1-877-929-0699 or by email at: safety@vancpharm.com

This leaflet was prepared by Vanc Pharmaceuticals Inc.

Date of preparation: September 10, 2014