

PRODUCT MONOGRAPH

Pr GRANISETRON HYDROCHLORIDE INJECTION

1 mg/mL granisetron (as granisetron hydrochloride)
1 mL and 4 mL vials

Manufacturer's Standard

STERILE

Antiemetic
(5-HT₃ receptor antagonist)

Sandoz Canada Inc.
145 Jules-Léger
Boucherville, QC, Canada
J4B 7K8

Date of Revision: October 23, 2014

Submission Control No: 177133

Table of Contents

PART I: HEALTH PROFESSIONAL INFORMATION.....	3
SUMMARY PRODUCT INFORMATION.....	3
INDICATIONS AND CLINICAL USE	3
CONTRAINDICATIONS	3
WARNINGS AND PRECAUTIONS	4
ADVERSE REACTIONS.....	6
DRUG INTERACTIONS	8
DOSAGE AND ADMINISTRATION	9
OVERDOSAGE	10
ACTION AND CLINICAL PHARMACOLOGY	10
STORAGE AND STABILITY	12
DOSAGE FORMS, COMPOSITION AND PACKAGING.....	13
PART II: SCIENTIFIC INFORMATION.....	14
PHARMACEUTICAL INFORMATION.....	14
CLINICAL TRIALS	15
DETAILED PHARMACOLOGY	16
TOXICOLOGY	19
REFERENCES	22
PART III: CONSUMER INFORMATION.....	24

Pr GRANISETRON HYDROCHLORIDE INJECTION

1 mg/mL granisetron (as granisetron hydrochloride)
1 mL and 4 mL vials

STERILE

PART I: HEALTH PROFESSIONAL INFORMATION

SUMMARY PRODUCT INFORMATION

Route of Administration	Dosage Form/Strength	Nonmedicinal Ingredients
Intravenous	Injection, 1 mg/mL granisetron (as granisetron hydrochloride)	Citric acid (monohydrate), methylparaben, propylparaben, sodium chloride, hydrochloric acid and/or sodium hydroxide and water for injection

INDICATIONS AND CLINICAL USE

Granisetron Hydrochloride Injection is indicated for:

Adults

- The prevention of nausea and vomiting associated with emetogenic cancer chemotherapy, including high-dose cisplatin.

Geriatrics (>65 years of age):

Chemotherapy-Induced Nausea and Vomiting

Safety and efficacy of granisetron hydrochloride appear to be similar to that observed in younger adults (see WARNINGS AND PRECAUTIONS and DOSAGE AND ADMINISTRATION).

Pediatrics:

Safety and efficacy of granisetron hydrochloride has not been adequately studied in children or adolescents under 18 years of age and it is not indicated for use in this population (see WARNINGS AND PRECAUTIONS and DOSAGE AND ADMINISTRATION).

CONTRAINDICATIONS

- Granisetron Hydrochloride Injection (granisetron hydrochloride) is contraindicated in patients with a known hypersensitivity to the drug or to any component of its formulation. For a complete listing, see the Dosage Forms, Composition and Packaging section of the product monograph.

- The concomitant use of Granisetron Hydrochloride Injection with apomorphine is contraindicated based on reports of profound hypotension and loss of consciousness when apomorphine was administered with another 5-HT₃ antagonist.

WARNINGS AND PRECAUTIONS

Carcinogenesis and Mutagenesis

Granisetron hydrochloride has been associated with an increased occurrence of hepatocellular tumours in carcinogenicity studies performed in rodents at doses in excess of the recommended human dose. Although the clinical significance of these findings has not been determined, the use of this drug should be restricted to the treatment of nausea and vomiting in patients undergoing emetogenic cancer chemotherapy. The recommended dosage of Granisetron Hydrochloride Injection should not be exceeded.

Granisetron was administered to rats in the diet in a 24-month carcinogenicity study. The incidence of hepatocellular carcinomas and adenomas was significantly increased in male rats treated at doses of 5 mg/kg/day and in rats of both sexes treated with 25 mg/kg/day. No increase in the rate of occurrence of liver tumours was observed in the 1 mg/kg/day treatment group (100 times the recommended human dose given intravenously).

In another 24-month carcinogenicity study, mice were administered granisetron in the diet at doses of 1, 5, and 50 mg/kg/day. There was a statistically significant increase in the incidence of hepatocellular carcinomas in males and hepatocellular adenomas in females dosed with 50 mg/kg/day. No statistically significant increase in liver tumours was observed in mice at a dose of 5 mg/kg/day (500 times the recommended human dose given intravenously).

Cardiovascular

As for other 5-HT₃ antagonists, cases of ECG modifications including QT prolongation have been reported with granisetron hydrochloride. These ECG changes with granisetron hydrochloride were minor and generally not of clinical significance, specifically with no evidence of proarrhythmia. However, in patients with pre-existing arrhythmias or cardiac conduction disorders, this might lead to clinical consequences. Therefore, caution should be exercised in patients with cardiac co-morbidities, on cardio-toxic chemotherapy and/or with concomitant electrolyte abnormalities. See also DRUG INTERACTIONS and ADVERSE REACTIONS, Post-marketing Reports of Adverse Events.

Gastrointestinal

Granisetron Hydrochloride Injection is not a drug that stimulates gastric or intestinal peristalsis. It should not be used instead of nasogastric suction. The use of Granisetron Hydrochloride Injection in patients with chemotherapy-induced nausea and vomiting may mask a progressive ileus and/or gastric distention. Patients with signs of subacute intestinal obstruction should be monitored following administration of Granisetron Hydrochloride Injection.

Sensitivity/Resistance

Hypersensitivity reactions may occur in patients who have exhibited hypersensitivity to other selective 5-HT₃ receptor antagonists.

Serotonin Syndrome/Neuroleptic Malignant Syndrome-like Events

Cases of life-threatening serotonin syndrome have been reported with 5-HT₃ receptor antagonist antiemetics, including granisetron hydrochloride, particularly when given in combination with other serotonergic and/or neuroleptic drugs. Serotonin syndrome symptoms may include mental status changes (e.g., agitation, hallucinations, coma), autonomic instability (e.g., tachycardia, labile blood pressure, hyperthermia), neuromuscular aberrations (e.g., hyperreflexia, incoordination) and/or gastrointestinal symptoms (e.g., nausea, vomiting, diarrhea). As this syndrome may result in potentially life-threatening conditions, treatment should be discontinued if such events occur and supportive symptomatic treatment should be initiated. If concomitant treatment of Granisetron Hydrochloride Injection with a drug affecting the serotonergic neurotransmitter system is clinically warranted, careful observations of the patient is advised, particularly during treatment initiation and dose increases (see DRUG INTERACTIONS).

Special Populations

Pregnant Women: The use of granisetron hydrochloride in pregnant women has not been studied and is not recommended. Reproduction studies performed in pregnant rats given granisetron at intravenous dosages up to 9 mg/kg/day and pregnant rabbits at intravenous dosages up to 3 mg/kg/day revealed no evidence of impaired fertility or harm to the fetus due to granisetron (see TOXICOLOGY, Reproduction).

Nursing Women: It is not known whether granisetron is excreted in human milk. Nursing is not recommended during treatment with Granisetron Hydrochloride Injection.

Pediatrics: The safety and efficacy of granisetron hydrochloride has not been adequately studied in children or adolescents under 18 years of age (see INDICATIONS AND CLINICAL USE and DOSAGE AND ADMINISTRATION).

Geriatrics (>65 years of age): During clinical trials, 713 patients 65 years of age or older received intravenous granisetron hydrochloride and of 325 patients 65 years of age or older who received oral granisetron hydrochloride, 298 were 65 to 74 years of age and 27 were 75 years of age or older. The efficacy and safety of granisetron hydrochloride did not appear to be age dependent (see INDICATIONS AND CLINICAL USE and DOSAGE AND ADMINISTRATION).

Information for Patients

Effect on Ability to Drive and Use Machinery

In healthy subjects, no clinically relevant effects on resting EEG or on the performance of psychometric tests were observed after IV granisetron hydrochloride at any dose tested (up to 200 mcg/kg). There are no data on the effect of granisetron hydrochloride on the ability to drive. As there have been occasional reports of somnolence in clinical studies, patients should be advised to avoid driving a car or operating hazardous machinery until they are reasonably certain that the drug treatment does not affect them adversely.

ADVERSE REACTIONS

Adverse Drug Reaction Overview

The most common adverse events reported by patients receiving intravenous granisetron hydrochloride in single-day chemotherapy trials are: headache, asthenia, somnolence, diarrhea, constipation, and abdominal pain (see Table 1 for the percentages of patients with these events). The only two common adverse experiences recognized to be causally related to granisetron hydrochloride are constipation and headache.

As for other 5-HT₃ antagonists, cases of ECG modifications including QT prolongation have been reported with granisetron hydrochloride. These ECG changes with granisetron hydrochloride were minor and generally not of clinical significance, specifically with no evidence of proarrhythmia (see WARNINGS AND PRECAUTIONS, Cardiovascular, DRUG INTERACTIONS, and ADVERSE REACTIONS, Post-marketing Reports of Adverse Events).

Clinical Trial Adverse Drug Reactions

Because clinical trials are conducted under very specific conditions the adverse reaction rates observed in the clinical trials may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse drug reaction information from clinical trials is useful for identifying drug-related adverse events and for approximating rates.

Chemotherapy-Induced Nausea and Vomiting

Intravenous granisetron hydrochloride was given as a single dose. Patients received cancer chemotherapy which consisted primarily of cisplatin or cyclophosphamide regimens. During the 24-hour period following intravenous administration of granisetron hydrochloride, IV fluids were also given. Adverse events were recorded over seven days when granisetron hydrochloride was given on a single day and up to 28 days when granisetron hydrochloride was administered for 7 or 14 days. In the absence of a placebo group, the relationship of observed adverse events to treatment with granisetron hydrochloride is difficult to judge.

Table 1 gives the frequencies of the six adverse events most commonly reported by patients receiving intravenous granisetron hydrochloride in single-day chemotherapy trials. This table does not include those events that are commonly associated with chemotherapy or the underlying malignant disease.

Table 1. Principal Adverse Events in Clinical Trials of Single-Day Chemotherapy

	Percentage of Patients with Event Intravenous granisetron hydrochloride (10-40 mcg/kg) (N=1519)
Headache	14%
Asthenia	5%
Somnolence	4%
Diarrhea	5%
Constipation	4%

	Percentage of Patients with Event Intravenous granisetron hydrochloride (10-40 mcg/kg) (N=1519)
Abdominal Pain	3%

The only two common adverse experiences recognized to be causally related to granisetron hydrochloride are constipation and headache. As with other drugs of this class, rare cases of hypersensitivity reactions, sometimes severe (e.g. anaphylaxis, shortness of breath, hypotension, urticaria) have been reported.

Less Common Clinical Trial Adverse Drug Reactions (≤1%)

Chemotherapy-Induced Nausea and Vomiting

The safety profile of granisetron hydrochloride has been evaluated in 3269 patients receiving intravenous granisetron hydrochloride (2 to 160 mcg/kg) in single-day and multiple-day clinical trials with emetogenic cancer therapies. In the listings which follow, a COSTART-based dictionary terminology has been used to classify reported adverse experiences. The frequencies presented, therefore, represent the proportion of the patients who experienced an event of the type cited on at least one occasion while receiving granisetron hydrochloride.

Experiences are further classified within body system categories and enumerated in order of decreasing frequency using the following definitions:

Frequent experiences are defined as: those occurring on one or more occasion in at least 1/100 patients

Infrequent adverse experiences are defined as: those occurring in less than 1/100 but at least 1/1000 patients

Rare experiences are defined as: those occurring in less than 1/1000 patients

Many adverse experiences are observed in cancer chemotherapy patients. All adverse experiences are included except those for which the drug cause was remote, those reported in general terms that are uninformative and those already listed in Table 1.

Body as a Whole

Frequent: Abdominal pain

Infrequent: Abdomen enlarged, chills, fever, malaise

Rare: Allergic reaction, chest pain

Cardiovascular System

Infrequent: Hypertension, hypotension, migraine, syncope, vasodilatation

Rare: Arrhythmia, bradycardia, palpitation, postural hypotension, tachycardia, ventricular arrhythmia, angina pectoris, and atrial fibrillation

Gastrointestinal System	Frequent: Decreased appetite Infrequent: Dry mouth, dyspepsia, flatulence, jaundice, liver function tests abnormal [Elevation of AST and ALT (>2 times the upper limit of normal)], nausea Rare: Gastrointestinal hemorrhage, hepatic coma, ileus, liver damage, melena, vomiting
Hemic and Lymphatic System	Rare: Coagulation time increased, eosinophilia, leukopenia, anemia, thrombocytopenia
Metabolic and Nutritional	Infrequent: Hypokalemia Rare: Bilirubinemia, edema, hyperphosphatemia, hyponatremia
Nervous System	Infrequent: Agitation, anxiety, dizziness, drugged feeling, insomnia, nervousness, paresthesia, tremor Rare: Coma, depersonalisation, grand mal convulsion, vertigo
Respiratory System	Infrequent: Dyspnea, hiccup Rare: Epistaxis, rhinitis, sinusitis
Skin and Appendages	Infrequent: Pruritus, rash, sweating Rare: Photosensitivity
Special Searches	Rare: Puncture site pain
Special Senses	Infrequent: Taste perversion Rare: Abnormal vision
Urogenital System	Infrequent: Dysuria Rare: Urinary incontinence

Post-marketing Reports of Adverse Events

The post-marketing safety experience in over 4 million patients is consistent with the clinical trial safety information.

Cases of cardiac arrest, ventricular fibrillation, ventricular tachycardia, ECG QT prolonged, sudden death and syncope were reported rarely in post-marketing experience. Some reports showed temporal association with granisetron, most of them also presenting confounding factors such as concomitant medications, medical history, and/or known risk factors. The relationship to granisetron cannot be established or excluded, either as an effect of the drug alone or in combination with other factors (see WARNINGS AND PRECAUTIONS, Cardiovascular, and DRUG INTERACTIONS).

As with other 5-HT₃ antagonists, cases of serotonin syndrome (including altered mental status, autonomic dysfunction and neuromuscular abnormalities) have been reported following the

concomitant use of granisetron hydrochloride and other serotonergic drugs (refer to WARNINGS AND PRECAUTIONS and DRUG INTERACTIONS).

DRUG INTERACTIONS

Overview

No pharmacodynamic interaction was found between single 160 mcg/kg IV doses of granisetron and single oral doses of 2.5 mg lorazepam or 3 mg haloperidol. Pharmacokinetic interactions with these drugs were not investigated.

The pharmacokinetic characteristics of a single 40 mcg/kg IV dose of granisetron were not significantly different whether it was administered alone or following 8 days of treatment with the hepatic enzyme inhibitor, cimetidine (200 mg four times daily).

Granisetron does not induce or inhibit the cytochrome P₄₅₀ drug metabolizing enzyme system.

As for other 5-HT₃ antagonists, cases of ECG modifications including QT prolongation have been reported with granisetron hydrochloride. These ECG changes with granisetron hydrochloride were minor and generally not of clinical significance, specifically with no evidence of proarrhythmia. However, in patients concurrently treated with drugs known to prolong QT interval and/or are arrhythmogenic, this may lead to clinical consequences.

Serotonin Syndrome

As with other serotonergic agents, serotonin syndrome, a potentially life-threatening condition, may occur with 5-HT₃ receptor antagonist antiemetic treatment, particularly with concomitant use of other agents that may affect the serotonergic neurotransmitter system (including triptans, SSRIs, SNRIs, lithium, sibutramine, fentanyl and its analogues, dextromethorphan, tramadol, tapentadol, meperidine, methadone and pentazocine or St. John's Wort [*Hypericum perforatum*]), and with drugs which impair metabolism of serotonin (such as MAOIs, including linezolid [an antibiotic which is a reversible non-selective MAOI], and methylene blue; See WARNINGS AND PRECAUTIONS).

DOSAGE AND ADMINISTRATION

Recommended Dose, Dosage Adjustment and Administration

Emetogenic Chemotherapy

Adults: The recommended dosage of Granisetron Hydrochloride Injection is 10 mcg/kg infused intravenously over 5 minutes, beginning within 30 minutes before initiation of chemotherapy only on the day(s) when chemotherapy is given (see Reconstitution for Diluted Solution).

Geriatrics: Available clinical data suggest that dosage reductions may not be necessary in this patient population (see INDICATIONS AND CLINICAL USE and WARNINGS AND PRECAUTIONS).

Pediatrics: See INDICATIONS AND CLINICAL USE and WARNINGS AND PRECAUTIONS.

Renally Impaired Patients: Available clinical data suggest that dosage reductions may not be necessary in this patient population.

Hepatically Impaired Patients: The clearance of granisetron hydrochloride is reduced by half in patients with hepatic impairment. The dose response of granisetron hydrochloride in patients with hepatic impairment has not been determined.

Reconstitution for Diluted Solutions

Infusion Preparation:

To prepare granisetron hydrochloride injection for IV infusion, aseptically transfer the appropriate amount of Granisetron Hydrochloride Injection to the desired volume of any of the following solutions: sodium chloride 0.9%, dextrose 5%, Hartmann's solution, mannitol (see STORAGE AND STABILITY).

OVERDOSAGE

There is no specific antidote for Granisetron Hydrochloride Injection (granisetron hydrochloride) overdose. In the case of overdose, symptomatic treatment should be given. Overdose has been reported with both the intravenous and oral formulation. Overdosage of up to 38.5 mg of granisetron hydrochloride injection has been reported without symptoms or with the occurrence of a slight headache.

For management of a suspected drug overdose, contact your regional Poison Control Centre.

ACTION AND CLINICAL PHARMACOLOGY

Mechanism of Action

Granisetron hydrochloride is a selective antagonist of 5-hydroxytryptamine (5-HT₃) receptors. Following exposure to emetogenic cancer chemotherapy, mucosal enterochromaffin cells release serotonin which stimulates 5-HT₃ receptors located peripherally on vagal nerve terminals and centrally in the nucleus tractus solitarius. The antiemetic effect of granisetron appears to involve antagonism of the serotonin-induced stimulation of vagal afferent activity.

Radioligand binding studies have demonstrated that granisetron hydrochloride has negligible affinity for other 5-HT receptors or for dopamine D₂ receptor binding sites.

Pharmacodynamics

In healthy subjects, granisetron hydrochloride produced no consistent or clinically significant changes in pulse rate, blood pressure or ECG. There was no evidence of an effect on psychomotor performance at intravenous doses of up to 200 mcg/kg IV. Granisetron

hydrochloride did not affect the plasma levels of prolactin or aldosterone at single intravenous doses of up to 300 mcg/kg or after repeat intravenous doses of 40 mcg/kg for 5.5 days.

Pharmacokinetics

Chemotherapy-Induced Nausea and Vomiting

In adult cancer patients undergoing chemotherapy and in healthy volunteers, infusion of a single 40 mcg/kg dose of granisetron hydrochloride produced the following mean pharmacokinetic data:

Table 2. Pharmacokinetic Parameters in Adult Cancer Patients Undergoing Chemotherapy and in Volunteers, Following a Single Intravenous 40 mcg/kg Dose of granisetron hydrochloride injection

	Peak Plasma Concentration (ng/mL)	Terminal Phase Plasma Half-Life (h)	Area Under Curve (AUC) (ng•h/mL)	Total Clearance (L/h)
Cancer Patients (N=14)				
Mean	63.8*	8.95*	167*	25.8*
Range	18.0 to 176	0.90 to 31.1	26.0 to 294	8.92 to 95.2
Young Adult Volunteers 21 to 42 years (N=20)				
Mean	64.3 ⁺	4.91 ⁺	89.7 ⁺	51.8 ⁺
Range	11.2 to 182	0.88 to 15.2	15.6 to 201	11.3 to 176
Elderly Volunteers 65 to 81 years (N=20)				
Mean	57.0 ⁺	7.69 ⁺	115 ⁺	27.1 ⁺
Range	14.6 to 153	2.65 to 17.7	37.7 to 240	10.9 to 58.4

* 5 minute infusion

+ 3 minute infusion

Distribution: Granisetron hydrochloride is extensively distributed between plasma and red blood cells with a mean volume of distribution of approximately 3 L/kg. Plasma protein binding is approximately 65%.

Metabolism: The clearance of granisetron occurs predominantly through hepatic metabolism. Biotransformation pathways involve N-demethylation and aromatic ring oxidation followed by conjugation.

Excretion: In normal volunteers, the urinary excretion of unchanged granisetron hydrochloride averages 12% of the administered dose over a period of 48 hours, while the remainder of the dose is excreted as metabolites, 47% in the urine and 34% in the feces. The metabolism of granisetron involves N-demethylation and aromatic ring oxidation followed by conjugation.

Special Populations and Conditions

Pediatrics: The safety and efficacy of granisetron hydrochloride has not been adequately studied in children or adolescents under 18 years of age.

Geriatrics: In geriatric (mean age 71 yrs) subjects after single intravenous doses of 40 mcg/kg, pharmacokinetic parameters were within the range found for young subjects (mean age 29 yrs). Although the elimination half-life was prolonged and the total plasma clearance reduced in the geriatric relative to the young subject group, no significant differences were determined between the two groups with regard to maximum plasma concentration or area under the plasma concentration time curve values (see Table 2).

Gender/Race: There were too few male and black patients to adequately assess differences in effect in either population (see CLINICAL TRIALS section).

Hepatic Insufficiency: A pharmacokinetic study in patients with hepatic impairment due to neoplastic liver involvement showed that total clearance was approximately halved and mean area under the plasma concentration time curve (AUC) values were approximately doubled compared to patients without hepatic impairment.

Renal Insufficiency: Although renal clearance was decreased in subjects with severe renal impairment (N=11) relative to normal volunteers (N=12), total plasma clearance was numerically higher in this renally impaired group (43 L/h) than in the normal volunteers (32 L/h). Mean area under the plasma concentration time curve values were similar for the two subject groups.

Cancer Patients: Following intravenous administration, mean terminal elimination half-life values are approximately twice as long in cancer patients as they are in healthy adult volunteers, while clearance values are decreased by approximately 50% (see Table 2).

STORAGE AND STABILITY

Store between 15 and 30°C. Discard 28 days after initial use.

Granisetron Hydrochloride Injection has been shown to be stable for at least 24 hours, at room temperature, and 72 hours, if refrigerated, in the following solutions: 20 mcg/mL with either, Dextrose Injection 5% or Sodium Chloride Injection 0.9% in polyvinyl chloride (PVC) bags.

As with all parenteral drug products, intravenous admixtures should be inspected visually for clarity, particulate matter, precipitate, discoloration and leakage prior to administration, whenever solution and container permit. Appropriate precautions should be taken to maintain the sterility of the infusion solution once prepared.

Pharmaceutical Precautions

As a general precaution, Granisetron Hydrochloride Injection should not be mixed in solution with other drugs.

DOSAGE FORMS, COMPOSITION AND PACKAGING

Each mL of Granisetron Hydrochloride Injection contains: 1 mg granisetron (as granisetron hydrochloride), methylparaben 0.12% and propylparaben 0.015% (as preservatives), citric acid (monohydrate) 0.2%, sodium chloride 0.9%, hydrochloric acid and/or sodium hydroxide to adjust pH and water for injection.

Granisetron Hydrochloride Injection is supplied in 1 mL amber multidose vials, boxes of 10 and in 4 mL amber multidose vials, boxes of 1.

PART II: SCIENTIFIC INFORMATION

PHARMACEUTICAL INFORMATION

Drug Substance


Proper Name: Granisetron Hydrochloride

Chemical Name: 1-Methyl-N-[(1R,3r,5S)-9-methyl-9-azabicyclo[3.3.1]non-3-yl]-1H-indazole-3-carboxamide hydrochloride.

Molecular Formula: $C_{18}H_{25}ClN_4O$

Molecular Mass: 348.88 g/mol

Structural Formula:



Physicochemical Properties: Granisetron hydrochloride is a white to off-white powder.

Solubility: Granisetron hydrochloride is freely soluble in water, sparingly soluble in methylene chloride, slightly soluble in methanol.

Melting Point Range: 290-292°C.

pH Value: the pH in 1% water solution is about 5.4.

CLINICAL TRIALS

Chemotherapy-Induced Nausea and Vomiting

Granisetron hydrochloride has been shown to prevent nausea and vomiting associated with single-day and repeat-cycle cancer chemotherapy.

Single-Day Chemotherapy

In a double-blind, placebo-controlled study in 28 cancer patients, granisetron hydrochloride administered as a single intravenous infusion of 40 mcg/kg, was significantly more effective than placebo in preventing nausea and vomiting induced by cisplatin chemotherapy (see Table 3).

Table 3. Prevention of Chemotherapy-Induced Nausea and Vomiting – Single-Day Cisplatin Therapy¹

	Granisetron hydrochloride	Placebo	P value
Number of Patients	14	14	----
Response Over 24 Hours			
Complete Response ²	93%	7%	<0.001
No Vomiting	93%	14%	<0.001
No More Than Mild Nausea	93%	7%	<0.001

¹ Cisplatin administration began within 10 minutes of granisetron hydrochloride by infusion and continued for 1.5 to 3.0 hours. Mean cisplatin doses were 86 mg/m² in the group of patients treated with granisetron hydrochloride by infusion and 80 mg/m² in the placebo group.

² No vomiting and no moderate or severe nausea.

Granisetron hydrochloride was evaluated in a double-blind, randomized dose response study of 353 patients stratified for high (> 80 to 120 mg/m²) or low (50 to 79 mg/m²) cisplatin dose. Response rates of patients for both cisplatin strata are given in Table 4.

Table 4. Prevention of Chemotherapy-Induced Nausea and Vomiting – Single-Day High-Dose and Low-Dose Cisplatin Therapy¹

Dose of granisetron hydrochloride (mcg/kg)	5	10	20	40
Number of Patients	82	90	88	93
Complete Response ² (%)	23	48*	48*	44*
No Vomiting (%)	32	54*	53*	48*
No Nausea (%)	22	46*	38*	38*

¹ Cisplatin administration began within 10 minutes of granisetron hydrochloride by infusion and continued 2 hours (mean). Mean Cisplatin dose was 82 mg/m².

² No vomiting and no rescue medication.

* p<0.05 vs 5 mcg/kg.

The 10, 20 and 40 mcg/kg doses were more effective than the 5 mcg/kg dose in preventing nausea and vomiting within 24 hours of chemotherapy administration. The 10 mcg/kg dose was at least as effective as the higher doses.

Repeat-Cycle Chemotherapy

Two single-blind, active-controlled studies have been performed in which granisetron hydrochloride was administered to a total of 246 chemotherapy-naive patients with malignant disease receiving cytostatic therapy (≥ 15 mg/m²/day cisplatin, ≥ 1.2 g/m²/day ifosfamide, and ≥ 120 mg/m²/day etoposide) for 5 days. Granisetron hydrochloride was administered as a daily 40 mcg/kg IV dose 5 minutes before the infusion of the cytostatic with up to two additional 40 mcg/kg IV doses permitted over each 24-hour period. In both studies, response rates (percentage of patients with no vomiting and no more than mild nausea in the 24-hour period following administration of granisetron hydrochloride) were observed to decline with repeated treatment, decreasing from 87-90% at day 1 to 70-71% at day 3, and 67-73% at day 5.

Co-Administration with Dexamethasone

A randomized, double-blind, placebo-controlled trial compared the safety and efficacy of intravenous granisetron hydrochloride (3mg) plus 8 mg dexamethasone phosphate with that of IV granisetron hydrochloride alone (3mg) in the prevention of emesis induced by cytotoxic chemotherapy.

A total of 278 patients received one of the following agents as their main cytotoxic therapy, either as a single agent or in combination with other cytotoxic agents: carboplatin >300 mg/m², cisplatin >20 mg/m² to <50 mg/m², dacarbazine >350 mg/m² to <500 mg/m², cyclophosphamide >500 mg/m² in combination, doxorubicin >40 mg/m² as single agent, doxorubicin >25 mg/m² in combination, epirubicin >75 mg/m² as a single agent, epirubicin >50 mg/m² in combination.

Patients received chemotherapy on Day 0 and were followed up to a further 6 days.

For the efficacy parameter of total control, there was a significantly better response in the group of patients treated with granisetron hydrochloride/dexamethasone than in the group of patients treated with granisetron hydrochloride/placebo on Day 0 ($p=0.020$) [95% CI (2.2%, 24.2%)] (see Table 5).

Table 5. Summary of Total Control¹ on Day 0 (Number (%) of Patients)

Treatment Group			
	Granisetron Hydrochloride/Dexamethasone (N=141)	Granisetron Hydrochloride/Placebo (N=137)	P-value
Day 0	103 (73.0%)	82 (59.9%)	0.020

¹Patients with no vomiting, no nausea, no rescue therapy and not withdrawn.

In addition, a statistically significant difference between the treatment groups was observed over the seven day period for the parameters of time to first rescue medication, time to first vomiting episode and time to first moderate/severe nausea.

DETAILED PHARMACOLOGY

Radioligand binding studies have been performed on rat and guinea pig brain membrane preparations. Granisetron appears to possess a high specificity for the 5-HT₃ receptor, while exhibiting negligible affinity for other 5-HT receptor subtypes (5-HT₁, 5-HT₂, 5-HT_{1A}, 5-HT_{1B/C}, and 5-HT_{1C}) or α 1, α 2, or β -adrenoreceptors; dopamine-D₂, histamine-H₁, benzodiazepine, picrotoxin, or opioid-binding sites.

The antagonistic effects of granisetron have been demonstrated in three models of 5-HT₃ receptor dependent activities, as listed below:

- 1) transient bradycardia (the Von Bezold-Jarisch reflex) following the intravenous injection of 5-HT into anesthetized rats ($IC_{50} = 0.7$ mcg/kg).
- 2) 5-HT- induced contractions of the guinea pig isolated ileum ($pA_2 = 8.1$).
- 3) tachycardia following 5-HT injection into the carotid arteries of the rabbit isolated heart ($pA_2 = 10.7$).

Two metabolites of granisetron (7-hydroxy metabolite and desmethyl metabolite) antagonized the Von Bezold-Jarisch reflex in anesthetized rats with potencies similar to that for the parent compound. However, the low plasma concentrations of these metabolites relative to the parent compound suggest that they are not likely to play a significant role following administration of granisetron.

Granisetron has proved efficacious both for the prophylaxis and treatment of emesis induced in the ferret by cisplatin, doxorubicin + cyclophosphamide, or X-irradiation. Maximal antiemetic efficacy in ferrets appeared to be achieved at a dose of 0.5 mg/kg IV administered 15 minutes before chemotherapy or radiation therapy. Furthermore, when a 0.5 mg/kg IV dose of granisetron was administered during emetic episodes occurring 90 minutes after cisplatin treatment, cessation of emesis was observed within 5 to 30 sec. of injection.

Granisetron was however, ineffective as an antiemetic in a canine model of apomorphine-evoked emesis and a ferret model of morphine-induced emesis suggesting that dopamine D₂ and opioid receptor antagonism are not components of its mechanism of action.

Other than for some inhibition of locomotor activity in mice at 10 mcg/kg SC and in rats at 1 to 5 mg/kg SC, granisetron did not exert central nervous system effects in the models studied. At cumulative doses up to 4.3 mg/kg IV administered over 2 hrs, granisetron had no effect on basal blood pressure or heart rate in conscious male rats. In the anesthetized dog, however, granisetron was demonstrated to decrease arterial blood pressure, heart rate, and myocardial contractility in a dose-dependent manner over a 1 to 3 mg/kg dose range.

Gastrointestinal: Granisetron (0.1-1.0 mg/kg SC) was associated with reduced fecal pellet output in conscious mice, suggesting a constipating effect.

Reproductive: Granisetron inhibited 5-HT induced contractions of the non-pregnant rat uterus *in vitro* with an IC_{50} of 5.9 mcM.

Preclinical Pharmacokinetics

Pharmacokinetics and ADME of granisetron have been extensively studied in rat and dog, the main species used in the non-clinical toxicology studies. Information has also been obtained on the mouse (used for carcinogenicity assessment), rabbit (teratology assessment) and ferret (efficacy pharmacology).

In rat and dog after intravenous dosing, granisetron freely diffused between plasma and red cells. Plasma protein binding in rats and dogs was moderate, 57% and 45%, respectively. A volume of distribution equivalent to approximately 3 L/kg in both species reflected the extensive tissue uptake expected of a lipophilic amine. In the rat, low excretion of granisetron in urine (approximately 2% dose) and a total plasma clearance (3.7 L/h/kg) similar to hepatic blood flow classified granisetron as a highly extracted drug whose clearance was flow-rate limited. In the dog, the plasma clearance value (2.6 L/h/kg) and the low urinary excretion (2-4% dose) classified granisetron extraction as medium to high in this species. In both species, relatively short granisetron plasma half-lives were observed (approximately 0.7 hours) and linear kinetics were indicated by the proportionate increases of granisetron plasma AUC with dose. As expected, no accumulation was observed on repeated daily dosing.

Complete absorption of ¹⁴C-granisetron from the gastrointestinal tract in rats, dogs, mice and rabbits was observed. However, oral bioavailability was severely reduced by the large first-pass effect resulting from the high liver extraction. Thus in rats, granisetron bioavailability was estimated at 0.2% of the dose after an oral dose of 5 mg/kg, whilst in dogs oral bioavailability was higher (about 17% at 0.25 and 1.5 mg/kg PO) reflecting lower liver extraction. In both species, bioavailability increased at the high-dose levels used in the toxicology studies (rat: about 10% bioavailability at 100 mg/kg PO; dogs: about 80% at 10 mg/kg PO), as the increased drug input partially saturated the first-pass effect.

Granisetron-related material (radioactivity) was rapidly and widely distributed to tissues after intravenous or oral doses of ¹⁴C-granisetron to rats. Whole body autoradiography and direct measurement revealed relatively high concentrations in excretory organs, liver and kidney, and low concentrations in blood and brain tissue. The time-course of elimination from tissues was similar to that from blood. The radioactivity was readily eliminated, such that only 1% remained in the tissues at 24 hours, though trace amounts were eliminated more slowly. Like many amine drugs, small amounts were taken up by melanin-containing tissues in pigmented animals and slowly released. On daily repeated intravenous dosing, minimal accumulation of radioactivity was observed in blood and tissues.

Granisetron was extensively metabolized, resulting in low excretion of unchanged drug in urine and feces. Metabolites found in mice, rats, rabbits, dogs and ferrets revealed that similar metabolic processes (oxidation at the N-methyl groups, oxidation in the benzenoid ring followed by sulphate and glucuronide conjugation, and a combination of these) were used in all species. However, quantitative differences between species were observed. Notably, 5-hydroxylation was higher than 7-hydroxylation in rats, mice and rabbits, whilst the converse was true for dogs and ferrets. Excretion of granisetron itself was low (no more than 13%) in all species studied. Granisetron excretion in feces accounted for less than 3% of the dose. Granisetron metabolites were readily excreted in both urine and feces of mice, rats, rabbits, dogs and ferrets. On

administering ¹⁴C-granisetron, urinary excretion of radioactivity in rats and dogs amounted to approximately 40% of the dose, irrespective of dose route, and the remainder was excreted in feces. In mice and rabbits, the urinary route accounted for about 60% dose, and in ferrets about 20% dose. The majority of urinary excretion occurred in the first 24 hours after dosing.

TOXICOLOGY

Acute Toxicity

The acute toxicity of granisetron is due primarily to CNS stimulation. LD₅₀ values by the intravenous route are within the range of 14 to 25 mg/kg in rats and mice.

The intravenous LD₅₀ for granisetron hydrochloride in mice is 17 mg/kg in males and 25 mg/kg in females. In rats it is 14 mg/kg in males and 16 mg/kg in females. The oral LD₅₀ is 350 mg/kg in both male and female mice. It is 350 mg/kg in male rats and 1100 mg/kg in female rats.

Long-term Toxicity

Subacute Toxicity Studies:

The subacute toxicity of granisetron was studied in the rat and dog; both species are suitable for the safety evaluation of granisetron hydrochloride on pharmacokinetic and metabolic grounds. In intravenous studies of up to 3 months duration in the rat, signs of acute CNS stimulation were dose-limiting at 9 mg/kg/day. Histopathological examination revealed an increase in the fat content of the liver in the majority of females at 6 mg/kg/day after 13 weeks of treatment but not following a further 4 weeks off-dose. In the dog, intravenous studies of up to 3 months duration resulted in convulsions at near lethal doses (3 mg/kg/day). Increases in aspartate aminotransferase (AST) and alanine aminotransferase (ALT) were seen in some animals at 3 mg/kg/day but there were no histopathological findings to indicate target organ toxicity at this level. Following treatment for 3 months, the intravenous no-toxic-effect level in both species was 0.5 mg/kg/day, representing a factor of approximately 3 over the maximum anticipated daily clinical dose of granisetron hydrochloride.

Chronic Toxicity Studies

Granisetron was administered in oral repeat-dose studies of up to 12 months duration in the rat and dog.

In the 6 and 12 month rat studies, there were changes in plasma enzymes associated with liver function, however, none of these changes was evident in subgroups of high-dose animals maintained for an off-dose period after the treatment. Dose-related increases in liver weights were also seen in rats given granisetron for up to 52 weeks, in the diet; such increases occurred in males dosed at 25 mg/kg/day and above.

Morphometric analysis confirmed there were an increased number of hepatocytes per unit area at the high dose, indicating that the increased liver weights were associated with hepatocyte hyperplasia. There was no evidence of hyperplasia at a dose of 5 mg/kg/day. Although there were increased incidences of rats with foci or areas of acidophilic and/or basophilic hepatocyte alteration in the intermediate and high-dose groups, precise morphometric quantification of the

amount of liver occupied by foci demonstrated that increases compared with the controls were confined to the high dose. These results clearly define 5 mg/kg/day to be a no-effect dose, at which the drug does not cause the production of liver foci or induce hyperplasia. The morphometric analysis also showed that, at the high dose, the amount of liver occupied by foci regressed after cessation of treatment.

In the 6-month dog study, there was a trend toward increased alanine aminotransferase (ALT) and lactate dehydrogenase (LDH) at the high dose of 10 mg/kg/day, although histopathological changes were not observed. Physical signs at the high dose during the initial stages of treatment consisted of prominence of the nictating membrane, black or dark discoloured feces and, in males, an increased incidence of loose feces. Emesis and isolated clonic convulsions were also noted. One high-dose male died on Day 181 having shown no previous signs of ill health; the cause of death could not be established. There were no toxic effects at the mid dose of 1.5 mg/kg/day.

In the 12-month dog study, the high dose (5 mg/kg/day) produced no CNS effects and there were no changes in plasma enzymes indicative of altered hepatic function or treatment related histopathological findings at this dose.

Carcinogenicity Studies

In a 24-month carcinogenicity study, mice were treated orally with granisetron 1, 5 or 50 mg/kg/day. There was a statistically significant increase in the incidence of hepatocellular carcinomas in males and hepatocellular adenomas in females dosed with 50 mg/kg/day (5000 times the recommended human dose given intravenously). No increase in liver tumours was observed in mice at a dose of 5 mg/kg/day (500 times the recommended human dose given intravenously).

In a 24-month carcinogenicity study, rats were treated orally with granisetron 1, 5 or 50 mg/kg/day. Owing to manifestations of toxicity, the 50 mg/kg dose was reduced to 25 mg/kg/day (2500 times the recommended human dose given intravenously) from week 59 of treatment onwards. There was a statistically significant increase in the incidence of hepatocellular carcinomas and adenomas in males dosed with 5 mg/kg/day (500 times the recommended human dose given intravenously) and above, and in females dosed with 50 mg/kg/day (5000 times the recommended human dose given intravenously). No increase in liver tumours was observed in rats at a dose of 1 mg/kg/day (100 times the recommended human dose, given intravenously) in males and 5 mg/kg/day (500 times the recommended human dose given intravenously) in females.

Experimental evidence in rats shows that granisetron exhibits the characteristics of a promoter of liver tumours with a clear no-effect dose of 1 mg/kg (100 times the recommended human dose given intravenously). The probable mechanism for this effect is sustained liver cell hyperplasia. In a study in which rats were treated for 12 months with 100 mg/kg/day (10,000 times the recommended human dose given intravenously), the observed promoting effects were reversible upon cessation of treatment. Additionally, there was no adverse effect on the liver of dogs treated for 12 months with granisetron, 5 mg/kg/day (500 times the recommended human dose given intravenously).

Mutagenicity Studies

The effects of granisetron were investigated in a battery of seven tests for mutagenicity, including an investigation of DNA damage in rat hepatocytes. Granisetron did not cause gene mutation in Ames bacterial assays in *Salmonella* and *E. coli* or in a mouse lymphoma cell assay. No evidence of chromosomal damage was observed in human lymphocytes *in vitro* or in a mouse micronucleus test at doses of up to 1800 times the recommended human dose given intravenously. However, granisetron was associated with a significant increase in the number of cells with polyploidy in an *in vitro* human lymphocyte chromosomal aberration test. There was no evidence of DNA damage and repair in assays of unscheduled DNA synthesis (UDS) in rat hepatocytes *in vitro* (or *in vivo* at doses of up to 35,000 times the recommended human dose given intravenously). There was an apparent increase in UDS in HeLa cells exposed to granisetron *in vitro* when DNA synthesis was measured by scintillation counting of incorporated radioactive thymidine. However, when this test was repeated using a more definitive autoradiographic methodology and microscopic examination of HeLa cells, the test was negative for UDS. It is likely that the apparent UDS in the initial study was, in fact, a reflection of DNA synthesis in cells undergoing normal division.

Reproduction

In a reproduction and fertility study in rats, granisetron at subcutaneous doses up to 6 mg/kg/day (600 times the recommended human dose given intravenously) had no effect on male or female fertility.

At dose levels which produced maternal toxicity, intravenous granisetron [up to 9 mg/kg/day (900 times the recommended human dose given intravenously) in rats, and up to 3 mg/kg/day (300 times the recommended human dose given intravenously) in rabbits] had no adverse effect on the course and outcome of pregnancy. A similar lack of effect was apparent in peri- and postnatal studies and general reproductive studies, in the rat.

REFERENCES

1. Andrews PLR, Rapeport WG, Sanger GJ. Neuropharmacology of emesis induced by anticancer chemotherapy. *TiPS* 1988;9:334-341.
2. Andrews PLR, Hawthorn JA. The neurophysiology of vomiting. *Balliere's Clin Gastroenterol* 1988;2(1): 141-168.
3. Bermudez J, Boyle EA, Miner WD, et al. The antiemetic potential of the 5-hydroxytryptamine receptor antagonist BRL 43694. *Br J Cancer* 1988;58:644-650.
4. Hawthorn J, Ostler KJ, et al. The role of the abdominal visceral innervation and 5-hydroxytryptamine M receptors in vomiting induced by the cytotoxic drugs cyclophosphamide and cisplatin in the ferret. *Q J Exp Physiol* 1988;73:7-21.
5. Goddard PM, Jones M, Pollard LA. The 5-HT₃ antagonist, BRL 43694, does not compromise the efficacy of cisplatin in tumour-bearing mice. *Cancer Chemother Pharmacol* 1990;25(5):377-9.
6. Pratt GD, Bowery NG, Kilpatrick GT et al. Consensus meeting agrees distribution of 5-HT₃ receptors in mammalian hindbrain. *TiPS* 1990;11:135-7.
7. Cupissol D. The efficacy of granisetron as a prophylactic antiemetic and intervention agent in high-dose cisplatin-induced emesis. *Eur J Cancer* 1990;26 (Suppl. 1):S23-S27.
8. Granisetron study group. The antiemetic efficacy and safety of granisetron compared with metoclopramide plus dexamethasone in patients receiving fractionated chemotherapy over 5 days. *J Cancer Res Clin Oncol* 1993;119:555-9.
9. Bremer K. A single-blind study of the efficacy and safety of intravenous granisetron compared with alizapride plus dexamethasone in the prophylaxis and control of emesis in patients receiving 5-day cytostatic therapy. *Eur J Cancer* 1992;28A (Suppl. 6/7):1018-1022.
10. Carmichael J et al. I.V. granisetron versus I.V.. granisetron plus I.V. dexamethasone in the prophylaxis of emesis induced by cytotoxic chemotherapy. Abstract presented at ECCO 7, Jerusalem 1993.
11. Hacking A. Oral granisetron – simple and effective: A preliminary report. *Eur J Cancer* 1992;28A(1):S28-S32.
12. Bleiberg HH-1 et al. Antiemetic treatment with oral granisetron in patients receiving moderately emetogenic chemotherapy: A dose-ranging study. *Clin Ther* 1995;17(1):38-50.
13. Heron JF et al. Oral granisetron alone and in combination with dexamethasone: A double-blind randomized comparison against high-dose metoclopramide plus dexamethasone in prevention of cisplatin-induced emesis. *Ann Oncol* 1994;5(7):579-584.

14. Gralla, RJ. et al. Single-dose oral granisetron has equivalent antiemetic efficacy intravenous ondansetron for highly emetogenic cisplatin-based chemotherapy. *J Clin Oncol* 1998;Vol 16(4): 1568-1573.
15. Perez, EA et al. Comparison of single-dose oral granisetron *versus* intravenous ondansetron in the prevention of nausea and vomiting induced by moderately emetogenic chemotherapy: A multicenter, double-blind, randomized parallel study. *J Clin Oncol* 1998;Vol.16(2):754-760.
16. Mayron, D et al. Stability and compatability of granisetron hydrochloride in IV solutions and oral liquids and during simulated Y-site injection with selected drugs. *Am J Health-Syst Pharm* 1996;Vol 53 :294-304.
17. Hoffmann-La Roche Limited, Mississauga, (ON), Canada, KYTRIL Product Monograph (Tablets and Injection), Control No.: 129213; Date of Revision: October 29, 2009.
18. Hoffmann-La Roche Limited, Mississauga, (ON), Canada, KYTRIL Product Monograph (Tablets), Control No.: 172914; Date of Revision: June 6, 2014.
19. Omega Laboratories Ltd, Montreal, (QC), Canada, Granisetron Hydrochloride Injection Product Monograph, Control No.: 163080; Date of revision: October 29, 2013.

PART III: CONSUMER INFORMATION

Pr Granisetron Hydrochloride Injection
 1 mg/mL granisetron (as granisetron hydrochloride)
 1 mL and 4 mL vials
 STERILE

This leaflet is part III of a three-part "Product Monograph" published when Granisetron Hydrochloride Injection was approved for sale in Canada and is designed specifically for Consumers. This leaflet is a summary and will not tell you everything about Granisetron Hydrochloride Injection. Contact your doctor or pharmacist if you have any questions about the drug.

ABOUT THIS MEDICATION**What the medication is used for:**

Granisetron Hydrochloride Injection is one of a group called antiemetics and it can only be obtained with a prescription from your doctor.

Granisetron Hydrochloride Injection is intended to prevent nausea (feeling sick) and vomiting which may occur after you receive cancer chemotherapy.

What it does:

Cancer chemotherapies are thought to cause the release of serotonin, a natural substance in the body. Serotonin can cause you to feel sick and to vomit. Granisetron, the active ingredient in Granisetron Hydrochloride Injection, will stop the action of serotonin and help prevent you from feeling sick and vomiting.

When it should not be used:

- Do not take this medicine if you are allergic to granisetron or any of the ingredients Granisetron Hydrochloride Injection contains.
- If you are taking apomorphine.

What the medicinal ingredient is:

Granisetron hydrochloride

What the nonmedicinal ingredients are:

Citric acid (monohydrate) 0.2%, methylparaben 0.12% and propylparaben 0.015% (as preservatives), sodium chloride 0.9%, hydrochloric acid and/or sodium hydroxide to adjust pH and water for injection.

What dosage forms it comes in:

Granisetron Hydrochloride Injection is supplied in 1 mL amber multidose vials, boxes of 10 and in 4 mL amber multidose vials, boxes of 1.

WARNINGS AND PRECAUTIONS

BEFORE you use Granisetron Hydrochloride Injection talk to your doctor or pharmacist if:

- you have any allergies to similar antiemetics such as dolasetron mesylate or ondansetron
- you are pregnant, plan to become pregnant or are breastfeeding
- you have liver problems
- you have a history of heart problems
- you have been told by a doctor that you have a blockage of your gut or if you have severe constipation, pain or swelling in your stomach
- you are taking other medications, including drugs you can buy without a prescription and herbal products.

Serotonin Syndrome is a rare but potentially life-threatening reaction that may occur if you take Granisetron Hydrochloride Injection with certain other medications. It may cause serious changes in how your brain, muscles, and digestive system work. Be sure to tell your healthcare professional all the medicines you are taking.

As Granisetron Hydrochloride Injection may cause drowsiness, you should avoid driving a car or operating hazardous machinery until you know it does not affect you.

INTERACTIONS WITH THIS MEDICATION

As with most medicines, interactions with other drugs are possible. To avoid potentially life-threatening reactions, tell your healthcare professional about **ALL** the medicines you take, including drugs prescribed by other doctors, vitamins, minerals, natural supplements, or alternative medicines.

PROPER USE OF THIS MEDICATION

This medicine is only for you, the person for whom the prescription was written. Do not give this medication to others.

Usual adult dose:

Granisetron Hydrochloride Injection will be given to you by hospital staff before and/or after your therapy.

Overdose:

In case of drug overdose, contact a health care practitioner, hospital emergency department or regional Poison Control Centre immediately, even if there are no symptoms.

SIDE EFFECTS AND WHAT TO DO ABOUT THEM

If you experience an allergic reaction (e.g. shortness of breath, drop in blood pressure, skin lumps or hives), contact your doctor immediately. Do not take any more medicine unless instructed to do so by your doctor.

You may experience headaches, constipation, weakness, sleepiness, diarrhea or abdominal pain while taking Granisetron Hydrochloride Injection. You may also experience pain, anemia or fever while on Granisetron Hydrochloride Injection. There is no need to stop the medicine but you should tell your doctor about these symptoms.

SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM

Symptom / effect		Talk with your doctor or pharmacist		Stop taking drug and seek immediate medical help
		Only if severe	In all cases	
Rare	Allergic reaction - symptoms include swelling of the mouth, throat, difficulty in breathing, rash, hives, increased heart rate			✓
	Disturbance in heart rhythm - dizziness, palpitations (fast, pounding or irregular heart beat), chest pain, fainting			✓
	Serotonin Syndrome - symptoms that may be observed when granisetron hydrochloride is taken with certain other medications include: •Fever, sweating, shivering, diarrhea, nausea, vomiting •Muscle shakes, jerks, twitches or stiffness, overactive reflexes, loss of coordination •Fast heartbeat, changes in blood pressure •Confusion, agitation, restlessness, hallucinations, mood changes, unconsciousness, and coma			✓

This is not a complete list of side effects. For any unexpected effects while taking Granisetron Hydrochloride Injection, contact your doctor or pharmacist.

HOW TO STORE IT

Granisetron Hydrochloride Injection should be stored between 15 and 30°C. Discard 28 days after initial use.

The expiry date of this medicine is printed on the label. Do not use the medicine after this date. Keep your medicine in a safe place out of reach of children.

Reporting Side Effects

You can help improve the safe use of health products for Canadians by reporting serious and unexpected side effects to Health Canada. Your report may help to identify new side effects and change the product safety information.

3 ways to report:

- Online at [MedEffect](#);
- By calling 1-866-234-2345 (toll-free);
- By completing a Consumer Side Effect Reporting Form and sending it by:

- Fax to 1-866-678-6789 (toll-free), or
- Mail to: Canada Vigilance Program
Health Canada, Postal Locator 0701E
Ottawa, ON
K1A 0K9

Postage paid labels and the Consumer Side Effect Reporting Form are available at [MedEffect](#).

NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

MORE INFORMATION

This document, plus the full Product Monograph prepared for health professionals, can be obtained by contacting the sponsor, Sandoz Canada Inc., at:

1-800-361-3062

Or by written request at:

145, Jules-Léger
Boucherville, (QC), Canada
J4B 7K8

Patient leaflet also available at www.sandoz.ca

This leaflet was prepared by Sandoz Canada Inc.

Last revised: October 23, 2014