

PRODUCT MONOGRAPH

Pr Zinda-ANASTROZOLE

anastrozole

1 mg tablet

Non-Steroidal Aromatase Inhibitor

Stason Pharmaceuticals, Inc.
11 Morgan
Irvine, CA 92618 USA

Date of Revision:
October 10, 2014

Zinda Pharma Limited
8250 Decarie Suite 110
Montreal, QC H4P 2P5

Submission Control No: 178552

Table of Contents

PART I: HEALTH PROFESSIONAL INFORMATION.....	3
SUMMARY PRODUCT INFORMATION	3
INDICATIONS AND CLINICAL USE.....	3
CONTRAINDICATIONS	3
WARNINGS AND PRECAUTIONS.....	4
ADVERSE REACTIONS.....	7
DRUG INTERACTIONS	19
DOSAGE AND ADMINISTRATION	20
OVERDOSAGE	21
ACTION AND CLINICAL PHARMACOLOGY	21
STORAGE AND STABILITY	25
DOSAGE FORMS, COMPOSITION AND PACKAGING	25
PART II: SCIENTIFIC INFORMATION	24
PHARMACEUTICAL INFORMATION.....	24
CLINICAL TRIALS	27
DETAILED PHARMACOLOGY	44
TOXICOLOGY	45
REFERENCES	50
PART III: CONSUMER INFORMATION.....	54

^{Pr}Zinda-ANASTROZOLE

Anastrozole Tablets
1 mg

PART I: HEALTH PROFESSIONAL INFORMATION

SUMMARY PRODUCT INFORMATION

Route of Administration	Dosage Form / Strength	All Nonmedicinal Ingredients
Oral	Tablet/1mg	croscarmellose sodium, hypromellose, lactose monohydrate, macrogol, magnesium stearate, maltodextrin, and povidone.

INDICATIONS AND CLINICAL USE

Zinda-ANASTROZOLE (anastrozole) is indicated for the adjuvant treatment of postmenopausal women with hormone receptor positive early breast cancer.

Approval is based on superior disease-free survival for anastrozole in comparison to tamoxifen. However, overall survival was not significantly different between the two treatments (see **PART II, CLINICAL TRIALS**).

Zinda-ANASTROZOLE is indicated for hormonal treatment of advanced breast cancer in postmenopausal women.

Geriatrics:

No changes in dose are necessary for elderly patients.

Pediatrics:

Zinda-ANASTROZOLE is not recommended for use in pediatric patients as safety and efficacy have not been established in this group of patients.

CONTRAINDICATIONS

- Patients who are hypersensitive to Zinda-ANASTROZOLE or to any ingredient in the formulation. For a complete listing, see the **Dosage Forms, Composition and Packaging** section of the Product Monograph.
- Pregnant or lactating women.

WARNINGS AND PRECAUTIONS

Serious Warnings and Precautions

- Not recommended for use in pre-menopausal women as safety and efficacy have not been established in these patients (see **ACTION AND CLINICAL PHARMACOLOGY** section).
- Not recommended for use in pediatric patients as safety and efficacy have not been established in these patients (see **Special Populations, Pediatrics** section below).
- Potential risk/benefit should be carefully assessed in patients with severe hepatic and severe renal impairment (see **Hepatic/Biliary** and **Renal** sections below).
- Potential risk/benefit should be carefully assessed in patients with osteoporosis or risk factors for osteoporosis (see **Musculoskeletal** section below).
- Should be administered under the supervision of a qualified physician experienced in the use of anti-cancer agents (see **DOSAGE AND ADMINISTRATION** section).

Body as a Whole

Zinda-ANASTROZOLE is unlikely to impair the ability of patients to drive and operate machinery. However, asthenia and somnolence have been reported with the use of anastrozole and caution should be observed when driving or operating machinery while such symptoms persist.

Cardiovascular

In the ATAC trial, ischemic cardiovascular events were reported more frequently in patients treated with anastrozole compared to those treated with tamoxifen, although the difference was not statistically significant. A retrospective evaluation has shown that this numerical difference was associated with a sub-group of patients with pre-existing ischemic heart disease. A statistical analysis could not be performed on this subgroup evaluation. Serious adverse events continued to be collected during the off-treatment follow-up and the incidence of cardiovascular events reported was similar in the anastrozole and tamoxifen arms (3.9% vs. 3.7%, respectively).

Hepatic/Biliary

Anastrozole pharmacokinetics have been investigated in subjects with stable hepatic cirrhosis related to alcohol abuse. The apparent oral clearance of anastrozole was approximately 30% lower in subjects with hepatic cirrhosis than in control subjects with normal liver function. However, plasma anastrozole concentrations in the subjects with hepatic cirrhosis are within the range of concentrations seen in normal subjects across all clinical trials. Dosage adjustment in patients with mild-to-moderate hepatic dysfunction is not necessary. Zinda-ANASTROZOLE has not been investigated in patients with severe hepatic impairment. The potential risk/benefit to such patients should be carefully considered before administration of Zinda-ANASTROZOLE.

Musculoskeletal

Arthralgia/Arthritis: The use of Aromatase Inhibitors, including anastrozole, may cause arthralgia/arthritis, which may impact on treatment compliance and quality of life.

In the ATAC study, 35.6% of patients on the anastrozole arm reported joint pain/stiffness (includes arthralgia, arthrosis, arthritis and joint disorder) versus 29.4% of patients on the tamoxifen arm. Arthritis alone was reported in 16.6% of patients on the ARIMIDEX arm versus 14.4% of patients on the tamoxifen arm.

Bone Mineral Density: The use of estrogen lowering agents, including Zinda-ANASTROZOLE, may cause a reduction in bone mineral density with a possible consequent increased risk of fracture. Data available from a phase III/IV study [SABRE (Study of Anastrozole with the Bisphosphonate Risedronate)] showed that in postmenopausal women with hormone receptor positive early breast cancer with existing moderate (T-score < -1.0 in either lumbar spine or total hip, provided neither of these was less than -2.0, and with no personal history of a fragility fracture) or high risk (T-score < -2.0 in either the lumbar spine, or hip, or a personal history of fragility fracture) of fragility fracture, bone mineral density (BMD) loss could be inhibited by using anastrozole together with a bisphosphonate (risedronate). All patients in the study received vitamin D and calcium supplementation. Patients at existing low risk (T-score in both the lumbar spine, and total hip, of -1.0 or higher, and no personal history of fragility fracture) of fragility fracture in the study were treated with anastrozole only and did not have a loss of lumbar spine BMD following 12 months of treatment although statistically significant changes were seen following 24 months of treatment (estimated percentage change -2.07%; 95% CI: -3.60, -0.53; p=0.0109). No change in total hip BMD was seen at 12 and 24 months in the low-risk group (see Clinical Trials, Adjuvant treatment of breast cancer in postmenopausal women – assessment of bone). Women should have their osteoporosis risk assessed and managed according to local clinical practice and guidelines.

Myalgia: Myalgia has been associated with both anti-estrogens and estrogen-lowering agents. In the adjuvant setting, muscle pain was reported in the ATAC study at a higher incidence for anastrozole (5.8%) compared to tamoxifen (5.2%).

Other

Zinda-ANASTROZOLE has not been investigated in patients with any degree of brain or leptomeningeal involvement or with pulmonary lymphangitic disseminated disease.

Renal

Anastrozole pharmacokinetics have been investigated in subjects with renal insufficiency. Anastrozole renal clearance decreased proportionately with creatinine clearance and was approximately 50% lower in subjects with severe renal impairment (creatinine clearance less than 30 mL/min/1.73m² or 0.5 mL/sec/1.73m²) compared to controls. Because renal clearance is not a significant pathway of elimination, the apparent oral clearance of anastrozole is unchanged even in severe renal impairment. Dosage adjustment in patients with renal dysfunction is not necessary.

Zinda-ANASTROZOLE has not been investigated in patients with breast cancer and severe renal impairment. The potential risk/benefit to patients with severe renal impairment should be carefully considered prior to the administration of Zinda-ANASTROZOLE.

Special Populations

Pregnant Women: Zinda-ANASTROZOLE is contraindicated in pregnant women.

The extent of exposure in pregnancy to anastrozole during clinical trials and post-marketing is very limited to individual cases only. If a patient becomes pregnant while receiving this drug, the patient should be apprised of the potential hazard to the fetus or potential risk for loss of the pregnancy.

Anastrozole has been found to cross the placenta following oral administration of 0.1 mg/kg in rats and rabbits. Studies in both rats and rabbits at doses equal to or greater than 0.1 and 0.02 mg/kg/day, respectively (about 1 and 1/3, respectively, the recommended human dose on a mg/m² basis), administered during the period of organogenesis showed that anastrozole increased pregnancy loss (increased pre-and/or post-implantation loss, increased resorption and decreased numbers of live fetuses). Effects were dose related in rats. Placental weights were significantly increased in rats at doses of 0.1 mg/kg/day or more.

Evidence of fetotoxicity, including delayed fetal development (i.e. incomplete ossification and depressed fetal body weights), was observed in rats administered doses of 1 mg/kg/day (about 8 times the recommended human dose on a mg/m² basis). There was no evidence of teratogenicity in rats administered doses up to 1 mg/kg/day. In rabbits, anastrozole caused pregnancy failure at doses equal to or greater than 1 mg/kg/day (about 16 times the recommended human dose on a mg/m² basis). There was no evidence of teratogenicity in rabbits administered 0.2 mg/kg/day (about 3 times the recommended human dose on a mg/m² basis).

Nursing Women: Zinda-ANASTROZOLE is contraindicated in lactating women.

Pediatrics: Zinda-ANASTROZOLE is not recommended for use in pediatric patients as safety and efficacy have not been established.

Geriatrics: Anastrozole pharmacokinetics have been investigated in postmenopausal female volunteers and patients with breast cancer. The pharmacokinetics were similar in volunteers and in patients, and no age related effects were seen.

Monitoring and Laboratory Tests

Zinda-ANASTROZOLE has not been observed to interfere with routine clinical laboratory test results.

During the ATAC trial, more patients receiving anastrozole were reported to have elevated serum cholesterol compared to patients receiving tamoxifen (9.0% versus 3.5%, respectively). Lipid profile was assessed as part of the SABRE trial. In this study, treatment for 12 months with anastrozole alone had a neutral effect on lipid profile.

ADVERSE REACTIONS

Adverse Drug Reaction Overview

Anastrozole has generally been well tolerated. Adverse events have usually been mild to moderate with few withdrawals from treatment due to undesirable events.

The pharmacological action of anastrozole may give rise to certain expected effects.

Arthritis/arthralgia, joint pain/stiffness and hot flushes were reported very commonly ($\geq 10\%$).

Common adverse reactions ($\geq 1\%$ - $< 10\%$) are: asthenia, joint pain/stiffness, bone pain, carpal tunnel syndrome, sensory disturbances (including paraesthesia, taste loss and taste perversion), vaginal dryness, hair thinning (alopecia), rash, nausea, diarrhea, headache and increases in alkaline phosphatase, alanine aminotransferase and aspartate aminotransferase.

Uncommonly reported adverse reactions ($\geq 0.1\%$ - 1%) are: vaginal bleeding, trigger finger, anorexia, hypercholesterolaemia, vomiting and somnolence, hepatitis and increases in gamma-GT and bilirubin. Rare cases ($\geq 0.1\%$ - 0.1%) of cutaneous vasculitis have been observed. Very rare cases ($< 0.01\%$) of erythema multiforme, Stevens-Johnson syndrome and allergic reactions including angioedema, urticaria and anaphylaxis have also been reported. These reported frequencies are generated from a number of anastrozole studies as well as post-marketing reports.

In the ATAC trial, ischemic cardiovascular events were reported more frequently in patients treated with anastrozole compared to those treated with tamoxifen, although the difference was not statistically significant. A retrospective evaluation has shown that this numerical difference was associated with a sub-group of patients with pre-existing ischemic heart disease. A statistical analysis could not be performed on this subgroup evaluation. Serious adverse events continued to be collected during the off-treatment follow-up and the incidence of cardiovascular events reported was similar in the anastrozole and tamoxifen arms (3.9% vs. 3.7%, respectively).

Events of carpal tunnel syndrome have been reported in patients receiving anastrozole treatment in clinical trials in greater numbers than those receiving treatment with tamoxifen. The majority of these events occurred in patients with identifiable risk factors for the development of the condition. In the ATAC adjuvant trial, 83 events of carpal tunnel syndrome occurred in 78 patients in the anastrozole monotherapy arm, and 22 events occurred in 22 patients in the tamoxifen arm.

Vaginal bleeding has been reported infrequently, mainly in patients during the first few weeks after changing from existing hormonal therapy to treatment with anastrozole. If bleeding persists, further evaluation should be considered.

Clinical Trial Adverse Drug Reactions

Adjuvant Treatment of Early Breast Cancer in Postmenopausal Women:

Anastrozole was generally well tolerated in the ATAC trial. At the time of the 5-year treatment completion analysis, the median duration of adjuvant treatment was 59.8 months and 59.6 months for patients receiving anastrozole 1 mg and tamoxifen 20 mg, respectively. The combination of anastrozole and tamoxifen did not demonstrate any safety benefits in

comparison to tamoxifen alone after the results from the first analysis (median duration of treatment was approximately 33 months).

Anastrozole was associated with statistically significant fewer discontinuations from treatment as a result of an adverse event compared to tamoxifen (11.1% vs. 14.3%) and fewer adverse drug reactions leading to discontinuation (6.5% vs. 8.9%). The incidence of on-treatment serious adverse events is significantly lower in patients receiving anastrozole 1 mg relative to tamoxifen 20 mg (33.3% versus 36.0%).

Adverse events occurring with an incidence of at least 5% in either treatment group during treatment or within 14 days of the end of treatment are presented below in Table 1.

Table 1: Adverse events occurring with an incidence of at least 5% in any treatment group during or within 14 days of the end of treatment from the ATAC trial

Body system and adverse event by COSTART-preferred term	Number (%) of patients ^a			
	33-month analysis (data cut-off 29 June 2001)		5-year treatment completion analysis (data cut-off 31 March 2004)	
	Anastrozole 1 mg (N=3092)	Tamoxifen 20 mg (N=3094)	Anastrozole 1 mg (N=3092)	Tamoxifen 20 mg (N=3094)
Body as a whole				
Asthenia	483 (15.6)	466 (15.1)	575 (18.6)	544 (17.6)
Pain	432 (14.0)	413 (13.3)	533 (17.2)	485 (15.7)
Back pain	238 (7.7)	234 (7.6)	321 (10.4)	309 (10.0)
Headache	253 (8.2)	197 (6.4)	314 (10.2)	249 (8.0)
Accidental injury	195 (6.3)	189 (6.1)	311 (10.1)	303 (9.8)
Infection	197 (6.4)	205 (6.6)	285 (9.2)	276 (8.9)
Abdominal pain	202 (6.5)	211 (6.8)	271 (8.8)	276 (8.9)
Chest pain	145 (4.7)	115 (3.7)	200 (6.5)	150 (4.8)
Flu syndrome	146 (4.7)	164 (5.3)	175 (5.7)	195 (6.3)
Neoplasm	101 (3.3)	99 (3.2)	162 (5.2)	144 (4.7)
Cyst	96 (3.1)	110 (3.6)	138 (4.5)	162 (5.2)
Cardiovascular				
Vasodilation	1060 (34.3)	1229 (39.7)	1104 (35.7)	1264 (40.9)
Hypertension	255 (8.2)	218 (7.0)	402 (13.0)	349 (11.3)
Digestive				
Nausea	287 (9.3)	281 (9.1)	343 (11.1)	335 (10.8)
Diarrhea	206 (6.7)	168 (5.4)	265 (8.6)	216 (7.0)
Constipation	183 (5.9)	203 (6.6)	249 (8.1)	252 (8.1)
Gastrointestinal disorder	126 (4.1)	104 (3.4)	210 (6.8)	158 (5.1)
Dyspepsia	150 (4.9)	124 (4.0)	206 (6.7)	169 (5.5)
Haemic and lymphatic				
Lymphoedema	247 (8.0)	277 (9.0)	304 (9.8)	341 (11.0)
Anemia	73 (2.4)	102 (3.3)	113 (3.7)	159 (5.1)
Metabolic and nutritional				
Peripheral edema	236 (7.6)	246 (8.0)	311 (10.1)	343 (11.1)
Weight gain	234 (7.6)	236 (7.6)	285 (9.2)	274 (8.9)
Hypercholesterolemia	186 (6.0)	68 (2.2)	278 (9.0)	108 (3.5)
Musculoskeletal disorders				
Arthritis	380 (12.3)	296 (9.6)	512 (16.6)	445 (14.4)
Arthralgia	386 (12.5)	252 (8.1)	467 (15.1)	344 (11.1)
Osteoporosis	192 (6.2)	134 (4.3)	325 (10.5)	226 (7.3)
Fracture	183 (5.9)	115 (3.7)	315 (10.2)	209 (6.8)
Arthrosis	161 (5.2)	112 (3.6)	207 (6.7)	156 (5.0)
Bone pain	158 (5.1)	139 (4.5)	201 (6.5)	185 (6.0)
Joint disorder	102 (3.3)	95 (3.1)	184 (6.0)	160 (5.2)
Myalgia	114 (3.7)	103 (3.3)	179 (5.8)	160 (5.2)
Nervous system				

Body system and adverse event by COSTART-preferred term	Number (%) of patients ^a			
	33-month analysis (data cut-off 29 June 2001)		5-year treatment completion analysis (data cut-off 31 March 2004)	
	Anastrozole 1 mg (N=3092)	Tamoxifen 20 mg (N=3094)	Anastrozole 1 mg (N=3092)	Tamoxifen 20 mg (N=3094)
Depression	323 (10.4)	315 (10.2)	413 (13.4)	382 (12.3)
Insomnia	253 (8.2)	226 (7.3)	309 (10.0)	281 (9.1)
Dizziness	180 (5.8)	191 (6.2)	236 (7.6)	234 (7.6)
Paraesthesia	181 (5.9)	106 (3.4)	215 (7.0)	145 (4.7)
Anxiety	147 (4.8)	147 (4.8)	195 (6.3)	180 (5.8)
Respiratory				
Pharyngitis	335 (10.8)	327 (10.6)	443 (14.3)	422 (13.6)
Cough increased	194 (6.3)	216 (7.0)	261 (8.4)	287 (9.3)
Dyspnea	173 (5.6)	164 (5.3)	234 (7.6)	237 (7.7)
Sinusitis	137 (4.4)	118 (3.8)	184 (6.0)	159 (5.1)
Bronchitis	126 (4.1)	107 (3.5)	167 (5.4)	153 (4.9)
Skin and appendages				
Rash	281 (9.1)	314 (10.1)	333 (10.8)	387 (12.5)
Sweating	112 (3.6)	158 (5.1)	145 (4.7)	177 (5.7)
Special senses				
Cataract specified	107 (3.5)	116 (3.7)	182 (5.9)	213 (6.9)
Urogenital				
Breast pain	176 (5.7)	121 (3.9)	251 (8.1)	169 (5.5)
Urinary tract infection	169 (5.5)	224 (7.2)	244 (7.9)	313 (10.1)
Vulvovaginitis	169 (5.5)	119 (3.8)	194 (6.3)	150 (4.8)
Breast neoplasm	94 (3.0)	89 (2.9)	164 (5.3)	139 (4.5)
Vaginitis	79 (2.6)	122 (3.9)	125 (4.0)	158 (5.1)
Vaginal hemorrhage ^b	100 (3.2)	151 (4.9)	122 (3.9)	180 (5.8)
Leucorrhea	68 (2.2)	264 (8.5)	86 (2.8)	286 (9.2)

^a Patients with multiple events in the same category are counted only once in that category. Patients with events in more than 1 category are counted once in each of those categories.

^b Vaginal hemorrhage without further diagnosis.

COSTART: Coding Symbols for Thesaurus of Adverse Reaction Terms.

N: Number of patients treated.

Certain adverse events (irrespective of drug causality) and combinations of adverse events were prospectively specified for analysis, based on the known pharmacological properties and side effect profiles of anastrozole and tamoxifen. Tamoxifen was statistically superior to anastrozole for the adverse events of joint disorders and fractures (including fractures of spine, hip and wrist) while anastrozole was statistically superior to tamoxifen for the adverse events of hot flushes, vaginal bleeding, vaginal discharge, endometrial cancer, venous thromboembolic events (including deep thromboembolic events) and ischemic cerebrovascular events.

A fracture rate of 22 per 1000 patient years was observed on anastrozole and 15 per 1000 patient years with the tamoxifen group with a median follow-up of 68 months. The rate of hip fractures was similar for anastrozole and tamoxifen in the ATAC trial. After a median follow-up of 100 months, fractures were reported more frequently in patients treated with anastrozole in comparison to tamoxifen, both during and off-treatment (13.7% vs. 10.1%; see Table 2), but the rate of fracture remained stable between the two groups. During the post-treatment follow-up period, the annual fracture rates were similar in the anastrozole and tamoxifen arms and the increased fracture episode rate seen during treatment was not observed following treatment completion as shown in Figure 1.

Table 2: Incidence of Fractures (during or off-trial treatment)

Category	Number (%) of Patients ^a	
	2007 update analysis (data cut-off 31 March 2007)	
	Anastrozole 1 mg (N=3092)	Tamoxifen 20 mg (N=3094)
Non-serious or serious		
All Fractures	425 (13.7)	313 (10.1)
Wrist/Colles	95 (3.1)	84 (2.7)
Spine	61 (2.0)	38 (1.2)
Hip	49 (1.6)	42 (1.4)
Serious		
All Fractures	212 (6.9)	170 (5.5)
Wrist/Colles	49 (1.6)	45 (1.5)
Spine	23 (0.7)	18 (0.6)
Hip	46 (1.5)	40 (1.3)


^a Patients with multiple events in the same category are counted only once in that category. Patients with events in more than one category are counted once in each of those categories.

Note: Off-trial treatment AEs were SAEs and any fracture event reported as being serious or non-serious that occurred more than 14 days after stopping study treatment (but within 10 years of starting study treatment). AEs starting after the patient's first recurrence visit were not reported.

Note: Off-trial treatment AEs included all off-treatment reports regardless of whether a patient had had a similar report on treatment.

N: Number of patients treated.

Figure 1: Annual first event rates of all fractures on or off study therapy



In the ATAC trial, ischemic cardiovascular events were reported more frequently in patients treated with anastrozole compared to those treated with tamoxifen, although the difference was not statistically significant (see Table 3). A retrospective evaluation has shown that this numerical difference was associated with a sub-group of patients with pre-existing ischemic heart disease. A statistical analysis could not be performed on this subgroup evaluation. From the 33 month analysis to the 68 month analysis, the incidence of cardiovascular events also remains stable over time between the two treatment groups. The incidence of myocardial infarctions increased by 0.1% in the anastrozole treatment group and 0.2% in the tamoxifen treatment group; the incidence of cerebrovascular accidents increased by 0.3% in each treatment group. During the off-treatment follow-up, when serious adverse events continued to be collected, the incidence of myocardial infarctions and cerebrovascular accidents was similar in both treatment groups.

Table 3 provides a summary of the pre-specified adverse events that occurred in either treatment group during treatment and after cessation of trial therapy.

Table 3: Incidence of pre-specified adverse events occurring in either treatment group during treatment and after cessation of trial therapy from the ATAC trial*

Adverse Event	Number (%) of patients ^a			
	5 year treatment completion analysis (data cut-off 31 March 2004)			
	Anastrozole 1 mg (N=3092)	Tamoxifen 20 mg (N=3094)	Odds ratio ^b	p-value
Hot flushes	1104 (35.7)	1264 (40.9)	0.80	<0.0001
Mood disturbances	600 (19.4)	557 (18.0)	1.10	0.2
Fatigue/asthenia	577 (18.7)	544 (17.6)	1.08	0.3
Nausea and vomiting	396 (12.8)	385 (12.4)	1.03	0.7
Vaginal discharge	111 (3.6)	407 (13.2)	0.25	<0.0001
Vaginal bleeding	171 (5.5)	323 (10.4)	0.50	<0.0001
Joint pain/stiffness	1111 (35.9)	922 (29.8)	1.32	<0.0001
Fractures	340 (11.0)	238 (7.7)	1.48	<0.0001
Fractures of the spine, hip, or wrist/Colles	148 (4.8)	112 (3.6)	1.34	0.02
Hip ^c	37 (1.2)	31 (1.0)	NC	NC
Spine ^c	45 (1.5)	27 (0.9)	NC	NC
Wrist/Colles ^c	72 (2.3)	63 (2.0)	NC	NC
Cataracts	191 (6.2)	219 (7.1)	0.86	0.2
Ischemic cardiovascular disease	137 (4.4)	119 (3.8)	1.16	0.2
Angina Pectoris ^c	75 (2.4)	56 (1.8)	NC	NC
Myocardial infarct ^c	42 (1.4)	40 (1.3)	NC	NC
Coronary artery disorder ^c	26 (0.7)	27 (0.9)	NC	NC
Myocardial ischemia ^c	24 (0.8)	16 (0.5)	NC	NC
Venous thromboembolic events	95 (3.1)	151 (4.9)	0.62	0.0003
Deep venous thromboembolic events	57 (1.8)	83 (2.7)	0.68	0.03
Ischemic cerebrovascular events	67 (2.2)	94 (3.0)	0.71	0.03
Endometrial cancer ^d	5 (0.2)	17 (0.8)	0.29	0.02

* All adverse events occurring during treatment or within 14 days of the end of treatment; all serious adverse events and all non-serious fractures occurring after 14 days from the end of treatment and prior to the confirmation of recurrence of breast cancer.

^a Patients with multiple events in the same category are counted only once in that category. Patients with events in more than one category are counted once in each of those categories.

^b Odds ratios of <1.00 indicate that treatment with anastrozole 1 mg is associated with a lower incidence of a specific event than Tamoxifen 20 mg.

^c Individual COSTART-preferred terms for a particular category of event – the broader category was the ‘pre-specified adverse event’.

^d Percentages calculated based upon the numbers of patients with an intact uterus at baseline (N=2229 for anastrozole and N=2236 for tamoxifen).

N: Number of patients treated.

NC: Not calculated.

Patients with Advanced Breast Cancer:

Two controlled clinical trials involving postmenopausal women with advanced breast cancer, compared treatment with tamoxifen (20 mg daily) versus treatment with anastrozole (1 mg daily). Table 4 presents adverse events reported in these trials with an incidence of greater than 5% in either treatment group, regardless of causality.

Table 4: Number (%) of Patients with Adverse Events from Trials 0027 and 0030*

Adverse Event by Body System	Anastrozole 1 mg (n = 506)	Tamoxifen 20 mg (n=511)
Body as a whole		
Asthenia	83 (16.4)	81 (15.9)
Pain	70 (13.8)	73 (14.3)
Back Pain	60 (11.9)	68 (13.3)
Headache	47 (9.3)	40 (7.8)
Chest Pain	37 (7.3)	37 (7.2)
Flu Syndrome	35 (6.9)	30 (5.9)
Pelvic Pain	23 (4.5)	30 (5.9)
Cardiovascular		
Vasodilation	128 (25.3)	106 (20.7)
Hypertension	25 (4.9)	36 (7.0)
Digestive		
Nausea	94 (18.6)	106(20.7)
Constipation	47 (9.3)	66 (12.9)
Abdominal Pain	40 (7.9)	38 (7.4)
Diarrhea	40 (7.9)	33 (6.5)
Vomiting	38 (7.5)	36 (7.0)
Anorexia	26 (5.1)	46 (9.0)
Metabolic and Nutritional		
Peripheral Edema	51 (10.1)	41 (8.0)
Musculoskeletal Disorders		
Bone Pain	54 (10.7)	52 (10.2)
Nervous System		
Insomnia	30 (5.9)	28 (5.5)
Dizziness	30 (5.9)	22 (4.3)
Depression	23 (4.5)	32 (6.3)
Hypertonia	16 (3.2)	26 (5.1)
Respiratory		
Cough Increased	55 (10.9)	52 (10.2)
Dyspnea	51 (10.1)	47 (9.2)
Pharyngitis	49 (9.7)	68 (13.3)
Skin and Appendages		
Rash	38 (7.5)	34 (6.7)
Urogenital		
Leucorrhea	9 (1.8)	31 (6.1)

*A patient may have more than one adverse event

Based on results from the established safety profiles of anastrozole and tamoxifen, the incidences of nine pre-specified adverse event categories, potentially causally related to one or both therapies because of their pharmacology, were statistically analyzed. No statistically significant differences were seen between treatment groups. The results are shown in Table 5.

Table 5: Number (%) of Patients from Trials 0027 and 0030*

Adverse Event by Body System	Anastrozole 1 mg (n = 506)	Tamoxifen 20 mg (n=511)
Body as a Whole		
Tumour Flare	15 (3.0)	18 (3.5)
Cardiovascular		
Hot Flushes	134 (26.5)	118 (23.1)
Thromboembolic Disease	23 (4.5)	39 (7.6)
Digestive		
Gastrointestinal Disturbances	170 (33.6)	196 (38.4)
Metabolic and Nutritional		
Weight Gain	11 (2.2)	8 (1.6)
Nervous System		
Depression	23 (4.5)	32 (6.3)
Lethargy	6 (1.2)	15 (2.9)
Urogenital		
Vaginal Dryness	15 (3.0)	13 (2.5)
Vaginal Bleeding	5 (1.0)	11 (2.2)

*Patients may appear in more than one row

The low incidence of vaginal bleeding and vaginal discharge was consistent with the known pharmacology of anastrozole, which would be predicted to have no estrogenic effect, and no effect on the endometrium. Despite the lack of estrogenic activity, there was no increase in myocardial infarction or pathological fracture when compared with tamoxifen. There was a low incidence of thromboembolic disease.

Patients with Advanced Breast Cancer Who had Disease Progression Following Tamoxifen Therapy

For two controlled clinical trials comparing anastrozole (1 mg and 10 mg) versus megestrol acetate (160 mg), adverse events reported in greater than 5% of the patients in any of the treatment groups, regardless of causality, are presented in Table 6.

Table 6: Number (n) and Percentage of Patients with Adverse Events from Trials 0004 and 0005*

Adverse Event by Body System	Anastrozole 1 mg (n=262) n (%)	Anastrozole 10 mg (n=246) n (%)	Megestrol Acetate 160 mg (n=253)
Body as a Whole			
Asthenia	42 (16.0)	33 (13.4)	47 (18.6)
Headache	34 (13.0)	44 (17.9)	24 (9.5)
Pain	28 (10.7)	38 (15.4)	29 (11.5)
Back Pain	28 (10.7)	26 (10.6)	19 (7.5)
Pelvic Pain	14 (5.3)	17 (6.9)	13 (5.1)
Chest Pain	13 (5.0)	18 (7.3)	13 (5.1)
Cardiovascular			
Hot Flashes	32 (12.2)	29 (10.6)	21 (8.3)
Digestive			
Nausea	41(15.6)	48 (19.5)	28 (11.1)
Vomiting	24 (9.2)	26 (10.6)	16 (6.3)
Diarrhea	22 (8.4)	18 (7.3)	7 (2.8)
Constipation	18 (6.9)	18 (7.3)	21 (8.3)
Abdominal Pain	18 (6.9)	14 (5.7)	18 (7.1)
Anorexia	18 (6.9)	19 (7.7)	11 (4.3)
Dry Mouth	15 (5.7)	11(4.5)	13 (5.1)
Metabolic and Nutritional			
Peripheral Edema	14 (5.3)	21 (8.5)	28 (11.1)
Weight Gain	4 (1.5)	9 (3.7)	30 (11.9)
Increased Appetite	0 (0)	1 (0.4)	13 (5.1)
Musculoskeletal Disorders			
Bone Pain	17 (6.5)	26 (11.8)	19 (7.5)
Nervous System			
Dizziness	16 (6.1)	12 (4.9)	15 (5.9)
Depression	14 (5.3)	6 (2.4)	5 (2.0)
Paresthesia	12 (4.6)	15 (6.1)	9 (3.6)
Respiratory			
Dyspnea	24 (9.2)	27 (11.0)	53 (20.9)
Cough Increased	22 (8.4)	18 (7.3)	19 (7.5)
Pharyngitis	16 (6.1)	23 (9.3)	15 (5.9)
Skin and Appendages			
Rash	15 (5.7)	15 (6.1)	19 (7.5)
Sweating	4 (1.5)	3 (1.2)	16 (6.3)
Urogenital			
Vaginal Hemorrhage	6 (2.3)	4 (1.6)	13 (5.1)

* A patient may have more than one adverse event.

Other less frequent (2% to 5%) adverse experiences reported in patients receiving anastrozole 1 mg in the two pivotal clinical trials are listed below. These adverse experiences are listed by

body system and are in order of decreasing frequency within each body system regardless of assessed causality.

Body as a Whole: Flu syndrome; fever; neck pain; malaise; accidental injury; infection

Cardiovascular: Hypertension; thrombophlebitis

Hepatic: Gamma GT increased; SGOT increased; SGPT increased

Hematologic: Anemia; leukopenia

Metabolic and Nutritional: Alkaline phosphatase increased; weight loss

Mean serum total cholesterol levels increased by 0.5 mmol/L among patients receiving anastrozole. Increases in LDL cholesterol have been shown to contribute to these changes.

Musculoskeletal: Myalgia; arthralgia; pathological fracture

Nervous: Somnolence; confusion; insomnia; anxiety; nervousness

Respiratory: Sinusitis; bronchitis; rhinitis

Skin and Appendages: Hair thinning; pruritus

Urogenital: Urinary tract infection; breast pain

The incidence of the following adverse event groups, potentially causally related to one or both of the therapies because of their pharmacology, were statistically analyzed: weight gain, edema, thromboembolic disease, gastrointestinal disturbance, hot flushes, and vaginal dryness. These six groups, and the adverse events captured in the groups, were prospectively defined. The results are shown in Table 7.

Table 7: Number (n) and Percentage of Patients from Trials 004 and 005

Adverse Event by Body System	Anastrozole 1 mg (n=262) n (%)	Anastrozole 10 mg (n=246) n (%)	Megestrol Acetate 160 mg n=253 n (%)
Cardiovascular			
Hot Flushes	33 (12.6)	29 (11.8)	35 (13.8)
Thromboembolic Disease	9 (3.4)	4 (1.6)	12 (4.7)
Digestive			
Gastrointestinal Disturbance	77 (29.4)	81 (32.9)	54 (21.3)
Metabolic and Nutritional			
Edema	19 (7.3)	28 (11.4)	35 (13.8)
Weight Gain	4 (1.5)	10 (4.1)	30 (11.9)
Urogenital			
Vaginal Dryness	5 (1.9)	3 (1.2)	2 (0.8)

More patients treated with megestrol acetate reported weight gain as an adverse event compared to patients treated with anastrozole 1 mg ($p < 0.0001$). Other differences were not statistically significant.

An examination of the magnitude of change in weight in all patients was also conducted. Thirty-four percent (87/253) of the patients treated with megestrol acetate experienced weight gain of 5% or more and 11% (27/253) of the patients treated with megestrol acetate experienced weight gain of 10% or more. Among patients treated with anastrozole 1 mg, 13% (33/262) experienced weight gain of 5% or more and 3% (6/262) experienced weight gain of 10% or more. On average, this 5 to 10% weight gain represented between 6 and 12 pounds.

No patients receiving anastrozole or megestrol acetate discontinued treatment due to drug-related weight gain.

Abnormal Hematologic and Clinical Chemistry Findings

Systematic collection of laboratory results (including total cholesterol) was not performed as specific endpoints in the ATAC trial. Abnormal laboratory results in ATAC are reported as an adverse event. During the ATAC trial, more patients receiving anastrozole were reported to have elevated serum cholesterol levels compared to patients receiving tamoxifen (9% versus 3.5%, respectively). In the SABRE trial, which was designed to specifically evaluate lipid levels in patients on anastrozole, no difference was observed in levels of low density lipoprotein-cholesterol (LDL-C), total cholesterol or triglycerides in patients taking anastrozole for 12 months compared to levels prior to commencing anastrozole treatment. There was a statistically significant increase in high density lipoprotein-cholesterol (HDL-C) in patients taking anastrozole for 12 months compared to levels prior to commencing anastrozole treatment (see Clinical Trials, Adjuvant treatment of breast cancer in postmenopausal women – assessment of lipids). On the basis of the SABRE data, no specific requirements for lipid monitoring due to anastrozole therapy are recommended.

Post-Market Adverse Drug Reactions

A case of severe acute hepatitis has been reported. Although late onset hepatotoxicity due to previous chemotherapy could not be ruled out, the temporal evidence suggested anastrozole as a possible cause. Cases of toxic hepatitis have been reported in association with anastrozole administration.

Cases of cutaneous vasculitis (including some reports of Henoch-Schönlein purpura) have been associated with anastrozole administration and symptoms have been reported to resolve within 10 - 30 days of discontinuing the drug, either spontaneously or with additional treatments.

Severe hypercalcaemia with high serum parathyroid hormone (PTH) levels was reported in a 65-year old woman on anastrozole. All parathyroid glands were considered normal and the hypercalcaemia and high PTH levels resolved with one month of anastrozole withdrawal. Calcium and PTH values increased to high levels again within 6 weeks of resumption of anastrozole.

Cases of paraesthesia (pain, numbness, and tingling of skin) and dysgeusia (taste loss and perversion) have been associated with ARIMIDEX administration.

DRUG INTERACTIONS

Overview

Anastrozole inhibits reactions catalyzed by cytochrome P₄₅₀ 1A2, 2C8/9, and 3A4 *in vitro* with K_i values which are approximately 30 times higher than the mean plasma steady-state C_{max} values observed following a 1 mg daily dose. Anastrozole has no inhibitory effect on reactions catalyzed by cytochrome P₄₅₀ 2A6 or 2D6 *in vitro*. Administration of a single 30 mg or multiple 10 mg doses of anastrozole to subjects had no effect on the clearance of antipyrine or urinary recovery of antipyrine metabolites. Based on these *in vitro* and *in vivo* results, it is unlikely that the administration of anastrozole 1 mg will result in clinically significant inhibition of cytochrome P₄₅₀-mediated metabolism of co-administered drugs.

Antipyrine, cimetidine, tamoxifen and warfarin clinical interaction studies indicate that the co-administration of anastrozole with other drugs is unlikely to result in clinically significant drug interactions mediated by cytochrome P₄₅₀.

A review of a global clinical trial safety database did not reveal evidence of clinically significant interactions in patients treated with anastrozole who also received other commonly prescribed drugs.

Estrogen-containing therapies should not be used with Zinda-Anastrozole as they may counteract the goal of achieving estrogen suppression.

Drug-Drug Interactions

Warfarin

The pharmacokinetics and anticoagulant activity of warfarin (25 mg) co-administered with anastrozole (1 mg daily) have been studied in healthy male volunteers. The mean plasma concentrations of anastrozole achieved throughout the warfarin dosing and sampling period were within the range seen in postmenopausal women with advanced breast cancer taking the clinically recommended dose of the drug. Overall, there was no evidence to suggest that anastrozole has any clinically relevant effects on the pharmacokinetics or anti-coagulant activity of warfarin.

Bisphosphonates

A review of a global clinical trial safety database showed that there are no clinically significant interactions with bisphosphonates. Results from the SABRE trial demonstrate that anastrozole in combination with the bisphosphonate, risedronate, was well tolerated.

Tamoxifen

The effect of anastrozole on tamoxifen (20 mg daily) pharmacokinetics has been studied in postmenopausal women with early breast cancer, who were already receiving tamoxifen as adjuvant therapy. There was no evidence of anastrozole having any significant effect on blood levels of tamoxifen compared to placebo (p=0.919).

Co-administration of anastrozole and tamoxifen did not affect tamoxifen or N-desmethyltamoxifen plasma concentrations, however, anastrozole plasma concentrations were reduced by 27% compared to those achieved with anastrozole alone. Combination treatment of anastrozole with tamoxifen has shown that anastrozole does not have a significant effect on blood levels of tamoxifen; estradiol suppression is consistent with that seen in patients treated with anastrozole alone.

Results from the ATAC trial (median follow-up of 33 months) suggest that tamoxifen should not be co-administered with anastrozole. The combination did not demonstrate any efficacy or safety benefit when compared to anastrozole or tamoxifen treatment alone, subsequently resulting in the discontinuation of the combination arm from the ATAC trial.

Table 8: Established or potential drug-drug interactions

Anastrozole	Ref	Effect	Clinical comment
Tamoxifen	CT	Tamoxifen and metabolite N-desmethyltamoxifen concentrations not affected. Anastrozole concentrations are decreased.	ATAC results indicate that the anastrozole-tamoxifen combination does not demonstrate any efficacy or safety benefits compared to tamoxifen monotherapy.

Legend: C = Case Study; CT = Clinical Trial; T = Theoretical

Drug-Food Interactions

Interactions with particular food have not been established.

Drug-Herb Interactions

Interactions with herbal products have not been established. Estrogen-containing herb therapies should not be used with anastrozole as they may counteract the goal of achieving estrogen suppression.

Drug-Laboratory Interactions

Anastrozole has not been observed to interfere with routine clinical laboratory tests results.

DOSAGE AND ADMINISTRATION

Dosing Considerations

Age: Patients should be postmenopausal.

Recommended Dose and Dosage Adjustment

Anastrozole should be administered 1 mg orally, once a day.

In the adjuvant setting, it is currently recommended that treatment be given for 5 years.

Elderly: No changes in dose are necessary for elderly patients.

Hepatic Impairment: Although the apparent oral clearance of anastrozole was decreased in subjects with cirrhosis due to alcohol abuse, plasma anastrozole concentrations remained within the range seen across all clinical trials in subjects without liver disease. Therefore, no changes in dose are recommended for patients with mild-to-moderate hepatic impairment,

although patients should be monitored for side effects. Anastrozole has not been studied in patients with severe hepatic impairment. The potential risk/benefit to such patients should be carefully considered before administration of Zinda-Anastrozole.

Renal Impairment: No changes in dose are necessary for patients with renal impairment. The potential risk/benefit to patients with severe renal impairment should still be considered prior to the administration of Zinda-Anastrozole in these patients.

Missed Dose

A missed dose should be taken as soon as possible, as long as it is taken at least 12 hours before the next dose is due. A missed dose should not be taken within 12 hours of the next dose.

Administration

Patients should swallow Zinda-ANASTROZOLE with fluids.

Patients should try to take Zinda-ANASTROZOLE at the same time each day.

OVERDOSAGE

For management of suspected drug overdose, contact your regional Poison Control Centre.

There is limited clinical experience of accidental overdose. Acute toxicity was seen in animals at a dose greater than 45 mg /kg (equivalent to 2.7 g). Clinical trials have been conducted with various dosages of anastrozole, up to 60 mg in a single dose given to healthy male volunteers and up to 10 mg daily given to postmenopausal women with advanced breast cancer; these dosages were well tolerated. A single dose of anastrozole that results in life-threatening symptoms has not been established.

There is no specific antidote to overdose and treatment must be symptomatic. In the management of an overdose, consideration should be given to the possibility that multiple agents may have been taken. Vomiting may be induced if the patient is alert. Dialysis may be helpful because anastrozole is not highly protein bound. General supportive care, including frequent monitoring of vital signs and close observation of the patient, is indicated. For management of suspected drug overdose, contact your regional Poison Control Centre.

ACTION AND CLINICAL PHARMACOLOGY

Mechanism of Action

Many breast cancers have estrogen receptors and growth of these tumours can be stimulated by estrogens. In postmenopausal women, the principal source of circulating estrogen (primarily estrone) is conversion of adrenally-generated androstenedione to estrone by aromatase in peripheral tissues, such as adipose tissue, with further conversion of estrone to estradiol. Many breast cancers also contain aromatase; the importance of tumour-generated estrogens is uncertain.

Treatment of breast cancer has included efforts to decrease estrogen levels by ovariectomy pre-menopausally and by use of anti-estrogens and progestational agents both pre- and post-menopausally, and these interventions lead to decreased tumour mass or delayed progression of tumour growth in some women.

Anastrozole is a potent and selective non-steroidal aromatase inhibitor. It significantly lowers serum estradiol concentrations and has no detectable effect on formation of adrenal corticosteroids or aldosterone.

Pharmacodynamics

Inhibition of aromatase activity is primarily due to anastrozole, the parent drug

The relationship between dose and response, measured as suppression of serum estradiol, was studied in postmenopausal women. Daily doses of anastrozole at 1 mg for 14 days produced estradiol suppression of greater than 80%. Suppression of serum estradiol was maintained for up to 6 days after cessation of daily dosing with 1 mg anastrozole.

In a study of 14 postmenopausal women diagnosed with locally advanced (Stage T3-T4) breast cancer with non-inflammatory, estrogen-receptor positive tumours, anastrozole was shown to be a potent suppressor of intra-tumoural estrogen levels. Following use as a 15-week primary systemic treatment (prior to any local surgery and/or radiotherapy), anastrozole suppressed intra-tumoural concentrations of estradiol (E_2), estrone (E_1) and estrone sulfate (E_1S) to mean values of 11.1%, 16.7% and 26.6%, respectively, of baseline levels. Three patients had intra-tumoural levels of E_2 , E_1 and E_1S suppressed below assay detection limits.

The selectivity of anastrozole to the aromatase enzyme, rather than other cytochrome P₄₅₀ enzymes controlling glucocorticoid and mineralocorticoid synthesis in the adrenal gland, has been established. Furthermore, provocative stimulation of the adrenal glands by ACTH in subjects under treatment with anastrozole up to 10 mg, produced a normal response in terms of cortisol and aldosterone secretion. Therefore, patients treated with anastrozole do not require glucocorticoid or mineralocorticoid replacement therapy.

Anastrozole does not possess direct progestogenic, androgenic or estrogenic activity and does not interfere with secretion of thyroid stimulating hormone (TSH).

Because of its pharmacological action, patients with estrogen and/or progesterone receptor-positive disease are the most appropriate population for anastrozole therapy.

Pharmacokinetics

Absorption: Absorption of anastrozole is rapid and maximum plasma concentrations typically occur within 2 hours of dosing under fasted conditions. Studies with radiolabeled drug have demonstrated that orally administered anastrozole is well absorbed into the systemic circulation. Food reduces the rate, but not the overall extent of anastrozole absorption.

Distribution: The pharmacokinetics of anastrozole are linear over the dose range of 1 to 20 mg and do not change with repeated dosing. Consistent with the 50-hour plasma elimination

half-life, plasma concentrations of anastrozole approach steady-state concentrations after 7 days of once daily dosing and are approximately three- to four-fold higher than the concentrations observed after a single dose of anastrozole. The protein binding of anastrozole to plasma proteins is about 40% and independent of concentration over a range, which includes therapeutic concentrations.

Metabolism: Metabolism of anastrozole occurs by N-dealkylation, hydroxylation and glucuronidation. Three metabolites of anastrozole (triazole, a glucuronide conjugate of hydroxy-anastrozole, and a glucuronide conjugate of anastrozole itself) have been identified in human plasma or urine. Several minor (less than 5% of the radioactive dose) metabolites excreted in the urine have not been identified. The major metabolite of anastrozole in the circulation, triazole, lacks pharmacologic activity.

Excretion: Studies in postmenopausal women with radiolabeled anastrozole demonstrated that elimination occurs primarily via metabolism (approximately 85%) and to a lesser extent renal excretion of unchanged anastrozole (approximately 11%). Anastrozole is eliminated slowly with a plasma elimination half-life of approximately 50 hours in postmenopausal women.

Special Populations and Conditions

Geriatrics: Anastrozole pharmacokinetics have been investigated in postmenopausal female volunteers and patients with breast cancer. The pharmacokinetics were similar in volunteers and in patients and no age related effects were seen.

Race: Anastrozole pharmacodynamics and pharmacokinetics have been studied in healthy, postmenopausal women in Japan, dosed for 16 days. The pharmacodynamic effect and pharmacokinetics of anastrozole 1 mg daily were similar in Japanese and Caucasian volunteers, and there was no indication that there would be any clinically significant differences in therapeutic responses to anastrozole between Japanese and Caucasian patients with breast cancer.

Hepatic Insufficiency: Anastrozole pharmacokinetics have been investigated in subjects with stable hepatic cirrhosis related to alcohol abuse. The apparent oral clearance of anastrozole was approximately 30% lower in subjects with hepatic cirrhosis than in control subjects with normal liver function. However, plasma anastrozole concentrations in the subjects with hepatic cirrhosis are within the range of concentrations seen in normal subjects across all clinical trials. Dosage adjustment in patients with mild-to-moderate hepatic impairment is not necessary. Anastrozole has not been studied in patients with severe hepatic impairment. The potential risk/benefit to such patients should be carefully considered before administration of Zinda-ANASTROZOLE.

Renal Insufficiency: Anastrozole pharmacokinetics have been investigated in subjects with renal insufficiency. Anastrozole renal clearance decreased proportionately with creatinine clearance and was approximately 50% lower in volunteers with severe renal impairment (creatinine clearance less than 30 mL/min/1.73m² or 0.5 mL/sec/1.73m²) compared to controls. Because renal clearance is not a significant pathway of elimination, the apparent oral clearance of anastrozole is unchanged even in severe renal impairment. Dosage adjustment in

patients with renal dysfunction is not necessary. The potential risk/benefit to patients with severe renal impairment should still be considered prior to the administration of Zinda-Anastrozole in these patients.

STORAGE AND STABILITY

Zinda-ANASTROZOLE should be stored at room temperature (15°C to 30°C). Keep container tightly closed.

SPECIAL HANDLING INSTRUCTIONS

No special instructions for handling are required.

DOSAGE FORMS, COMPOSITION AND PACKAGING

Zinda-ANASTROZOLE (anastrozole) is a 1 mg, round, film coated, white to off white tablet with 'BGN' engraved on one side and "2033" on the other side.

In addition to the active ingredient anastrozole, each tablet contains the following inactive ingredients: croscarmellose sodium, hypromellose, lactose monohydrate, macrogol, magnesium stearate, maltodextrin, and povidone.


Packaging formats: Zinda-ANASTROZOLE is available in bottles of 30 tablets and 500 tablets.

PART II: SCIENTIFIC INFORMATION

PHARMACEUTICAL INFORMATION

Drug Substance

Proper name:	anastrozole
Chemical name:	2, 2'-[5-(1H-1,2,4-triazol-1 ylmethyl)-1, 3-phenylene] bis (2-methylpropionitrile) (IUPAC)
Molecular formula and molecular mass:	C ₁₇ H ₁₉ N ₅ ; 293.37 g/mol
Structural formula:	



Physicochemical properties:

Anastrozole is a fine white to off-white powder, which has moderate aqueous solubility (0.53 mg/mL at 25°C) which is dependent on pH, from pH 1 to 4, but independent of pH thereafter. The molecule has a pKa of 1.4 (base) and is therefore only ionized at low pH. The log P (octanol/water) is 1.58, indicating that it is a moderately lipophilic compound.

Recrystallization of anastrozole from various solvents/solvent mixtures has to date yielded only one morphological form, which has been characterized using Differential Scanning Calorimetry (DSC), X-Ray Powder Diffraction (XRPD) and Fourier Transform Infra-Red Spectroscopy (FTIR).

CLINICAL TRIALS

Comparative Bioavailability Studies

A single dose crossover comparative bioavailability study of Zinda-Anastrozole 1 mg tablets and Arimidex[®] (anastrozole) 1 mg tablets (AstraZeneca Canada Inc.) following a single dose of 1 mg anastrozole in 23 healthy male adult volunteers under fasting conditions was conducted. The results indicate that Zinda-Anastrozole 1 mg tablets are bioequivalent to Arimidex[®] 1 mg tablets. The summary of results is presented in the following table:

Anastrozole (1 x 1 mg) From measured data uncorrected for potency Geometric LS Mean Arithmetic Mean (CV %)				
Parameter	Zinda-Anastrozole Stason Pharmaceuticals	Arimidex ^{®†} AstraZeneca Canada Inc.	% Ratio of Geometric LS Means	90% Confidence Interval
AUC ₀₋₇₂ (ng·h/mL)	465.52 477.4 (22.1)	465.39 472.8 (17.6)	100.0	95.9 – 104.2
C _{max} (ng/mL)	14.5116 14.745 (17.8)	14.3586 14.605 (19.3)	100.9	96.6 – 105.3
T _{max} [§] (h)	1.500 (0.667-4.000)	1.500 (0.667-4.000)		

[†] Reference product purchased in Canada

[§] Expressed as the median (min-max) only

AUC₁ and T_{1/2} are not reported; these parameters could not be accurately estimated due to the long half-life of the active ingredient and the design of the study.

Adjuvant treatment of breast cancer in postmenopausal women

Study demographics and trial design

A multicentre phase III trial entitled ‘A Randomized, Double-Blind Trial Comparing anastrozole alone with tamoxifen alone, with anastrozole and tamoxifen in combination as Adjuvant Treatment in Postmenopausal Women with Breast Cancer: (ATAC)’ was conducted in 9,366 postmenopausal patients with operable breast cancer. The patients were randomized to receive anastrozole 1 mg daily, tamoxifen 20 mg daily, or a combination of the two treatments for five years or until recurrence of the disease. However, at the time of the primary analysis (at a median of 33 months follow-up), the combination of anastrozole and tamoxifen did not demonstrate any efficacy benefit when compared with tamoxifen, and this treatment arm was subsequently discontinued from the trial, leaving the 6,241 patients who had been randomized to the anastrozole and tamoxifen monotherapy arms of the study. These patients will be followed to 10 years post-randomization.

The primary endpoints were disease-free survival and safety. Disease-free survival includes loco-regional (including new primary ipsilateral breast cancer) and distant recurrences, new

contralateral primaries and death from any cause as a first event. The secondary endpoints were distant disease-free survival (time to a first event of distant recurrence or death from any cause), the incidence of new contralateral breast primaries and overall survival. The primary analysis for disease-free survival was to be carried out after 352 events per treatment arm and occurred after a median of 33 months of follow-up; the major analysis for survival was to be carried out after a total of 352 events per treatment arm and occurred after a median follow-up of 68 months.

Demographics and other baseline characteristics were similar between the two treatment groups and are summarized in Table 9.

Table 9: Summary of demographic and baseline characteristics for the ATAC trial

Demographic Characteristic	Anastrozole 1 mg (N=3125)	Tamoxifen 20 mg (N=3116)
Mean age (yrs)	64.1	64.1
Age Range (yrs)	38.1 - 92.8	32.8 - 94.9
<45 yrs	0.7	0.4
45-60 yrs	34.6	35.1
>60<70 yrs	38.0	37.1
>70 yrs	26.7	27.4
Mean Weight (kg)	70.8	71.1
Receptor Status (%)		
Positive ¹	83.8	83.4
Negative ²	7.5	8.0
Other ³	8.8	8.6
Other treatment prior to randomization (%)		
Mastectomy	47.8	47.3
Breast conservation ⁴	52.2	52.7
Axillary surgery	95.5	95.7
Radiotherapy	63.4	62.5
Chemotherapy	22.3	20.8
Neoadjuvant Tamoxifen	1.6	1.6
Primary tumour size (%)		
T1 (≤ 2 cm)	63.9	62.9
T2 (>2 cm and ≤ 5 cm)	32.5	34.2
T3 (>5 cm)	2.7	2.2
Nodal status (%)		
Node positive	34.9	33.6
1-3 (# of nodes)	24.5	24.5
4-9	7.5	6.4
>9	2.9	2.7
Tumour grade (%)		
Well-differentiated	20.8	20.5
Moderately differentiated	46.8	47.8
Poorly/undifferentiated	23.6	23.3

Demographic Characteristic	Anastrozole 1 mg (N=3125)	Tamoxifen 20 mg (N=3116)
Not assessed/recorded	8.7	8.4

- 1 includes patients who were estrogen receptor (ER) positive or progesterone receptor (PgR) positive, or both positive.
 - 2 includes patients with both ER negative and PgR negative receptor status.
 - 3 includes all other combinations of ER and PgR receptor status unknown.
 - 4 among the patients who had breast conservation, radiotherapy was administered to 95.0% of patients in the anastrozole arm and 94.1% in the tamoxifen arm
- N=Number of patients randomized to the treatment.


Study results

Patients in the ATAC trial have now been treated for a median of 60 months (5 years) and followed for a median of 100 months. The primary analysis was carried out after a median follow-up of 33 months; the most recent analyses were carried out after a median follow-up of 68 and 100 months.

In the assessment of disease-free survival, anastrozole was superior to tamoxifen in the intent-to-treat population with a statistically significant 17% reduction in the risk of disease recurrence or death from any cause ($p=0.01$) at the primary analysis (median follow-up of 33 months) and a 13% reduction in risk after a median follow-up of 68 months ($p=0.01$). At a median follow-up of 100 months, anastrozole maintained statistical superiority with a 10% reduction in the risk of disease recurrence or death from any cause ($p=0.0252$). In the hormone receptor positive subgroup (representing about 84% of trial patients), there was a significant 22% reduction in the risk of disease recurrence or death from any cause ($p=0.006$) at the primary analysis, a 17% reduction ($p=0.005$) at the 68 month analysis and a 15% reduction ($p=0.0027$) at the 100 month analysis. These results demonstrate a carryover effect of the efficacy benefit of anastrozole following treatment completion in both the ITT and the HR+ populations. The absolute difference in disease-free survival continues to increase from 2.4% at 68 months to 2.8% at 100 months in the intent-to-treat population and from 2.5% in the hormone receptor positive subgroup at 68 months to 4.1% at 100 months.

Figure 2 and 3 presents the Kaplan-Meier probability of the protocol-defined disease-free survival for the intent to treat population and the hormone receptor positive subgroup.

Figure 2: Kaplan-Meier probability of disease-free survival for the intention-to treat (ITT) population.




NUMBER OF PATIENTS AT RISK:

MONTHS	0	12	24	36	48	60	72	84	96	108
A	3125	3005	2879	2763	2656	2511	2316	2087	1722	708
T	3116	2992	2838	2713	2585	2426	2215	2005	1648	661

NB: THE VERTICAL AXIS HAS BEEN TRUNCATED TO IMPROVE THE CLARITY OF THE KAPLAN-MEIER CURVES

Figure 3: Kaplan-Meier probability of disease-free survival for the hormone- receptor-positive subgroup.



The post-treatment follow-up analysis continues to demonstrate a significant advantage of anastrozole over tamoxifen in time to recurrence in both the intent-to treat population (HR 0.81, 95% CI 0.73 to 0.91, $p=0.0004$) and the hormone receptor positive subgroup (HR 0.76, 95% CI 0.67 to 0.87, $p=0.0001$), as well as in the time to distant recurrence in both the intent-to-treat population (HR 0.86, 95% CI 0.75 to 0.98, $p=0.022$) and the hormone receptor positive subgroup (HR 0.84, 95% CI 0.72 to 0.97, $p=0.022$). Additionally, there is a significant advantage of anastrozole over tamoxifen in risk of invasive contralateral breast cancer in both the intent-to-treat population (HR 0.68, 95% CI 0.49 to 0.94, $p=0.02$) and the hormone receptor positive subgroup (HR 0.60, 95% CI 0.42 to 0.85, $p=0.004$).

Anastrozole 1 mg did not show a survival advantage over tamoxifen 20 mg in the primary analysis of survival, which was carried out after a median follow-up of 68 months. Overall survival was similar in the two arms of the trial for both the intent-to-treat population (HR 0.97, 95% CI 0.85 to 1.12, $p=0.71$) and the hormone receptor positive subgroup (HR 0.97, 95% CI 0.83 to 1.14, $p=0.73$). After a median follow-up of 100 months, overall survival continued to be similar in the two arms of the trial for both the intent-to-treat population (HR 1.00, 95% CI 0.89 to 1.12, $p=0.99$) and the hormone receptor positive subgroup (HR 0.97, 95% CI 0.86 to 1.11, $p=0.68$).

Overall, a similar number of deaths occurred in the tamoxifen group and the anastrozole arm although there were fewer deaths following breast cancer recurrence in the anastrozole arm (HR=0.91; A=11.2%, T=12.3%). There were more deaths due to causes other than breast cancer and fewer deaths related to breast cancer among patients receiving anastrozole therapy. The difference in the numbers of non-breast cancer deaths before recurrence in the intention to treat population between the two treatment groups is small (absolute difference of 1.1%; A=8.9%, T =7.8%). The largest imbalance is seen among deaths from secondary cancers (A=2.8%, T=2.1%) and, in particular, deaths from lung and colorectal cancer compared with tamoxifen.

The incidence of ovarian cancer, endometrial cancer and melanoma was lower with anastrozole than in the tamoxifen group (see Table 10).

Table 10: Incidences of new primary cancers in either treatment group prior to recurrence (during or off-trial treatment)

Body System and adverse event by COSTART-preferred term	Number (%) of Patients	
	2007 update analysis (data cut-off 31 March 2007)	
	Anastrozole 1 mg (N=3092)	Tamoxifen 20 mg (N=3094)
Skin – non melanoma ^{a, b}	94 (3.0)	100 (3.2)
Contralateral breast cancer ^c	62 (2.0)	87 (2.8)
Colorectal	56 (1.8)	36 (1.2)
Lung	42 (1.4)	24 (0.8)
Ovary	12 (0.4)	26 (0.8)
Head and Neck	12 (0.4)	5 (0.2)
Kidney	11 (0.4)	6 (0.2)
Lymphoma (non-Hodgkins)	10 (0.3)	8 (0.3)
Gastric ^d	10 (0.3)	6 (0.2)
Melanoma	8 (0.3)	18 (0.6)
Leukaemia	7 (0.2)	9 (0.3)
Bladder	6 (0.2)	9 (0.3)
Brain	4 (0.1)	6 (0.2)
Endometrium ^a	4 (0.1)	23 (0.7)
Cervix ^a	2 (0.1)	5 (0.2)
Pancreas	2 (0.1)	6 (0.2)
Other	34 (1.1)	21 (0.7)
TOTAL	351 (11.4)	365 (11.8)

a In addition to the new primary cancers tabulated here, the following new primary cancers were reported as SAEs: 4 skin cancers and 1 endometrial cancer in the anastrozole 1 mg group and 8 skin cancers, 1 cervix cancer and 1 endometrial cancer in the tamoxifen 20 mg group.

b These totals include 2 patients in the anastrozole 1 mg group and 1 patient in the tamoxifen 20 mg group with new primary skin cancers that were categorized as skin (non-Hodgkin's).

c Excludes any new primary (contralateral) breast cancer occurring after recurrence.

d These totals include 2 patients in the anastrozole 1 mg group with new primary gastric cancers that were categorized as stomach cancers.

COSTART Coding Symbols for Thesaurus of Adverse Reaction Terms.

N=Number of patients treated.

Deaths (both during and off-trial treatment) resulting from ischaemic cardiovascular events (A =1.3%, T =1.2%) or other cardiovascular events (A =0.9%, T =0.9%) occurred with similar frequency for both treatment groups.

The analyses of the study endpoints in the intent-to-treat population and hormone receptor positive subgroup at the time of the primary, the 5-year treatment completion and the 100 month analyses are summarized in Table 11. The frequency of individual events in the intent-to-treat population and the hormone receptor positive subgroup at the 100 month analyses are described in Table 12.

Table 11: ATAC endpoint summary

	33-month analysis (data cut-off 29 June 2001)				68-month analysis (data cut-off 31 March 2004)				100-month analysis (data cut-off 31 March 2007)			
	Intent to Treat Population		Hormone Receptor positive population		Intent to Treat Population		Hormone Receptor positive population		Intent to Treat Population		Hormone Receptor positive population	
	Anastrozole 1 mg (N=3125)	Tamoxifen 20 mg (N=3116)	Anastrozole 1 mg (N=2618)	Tamoxifen 20 mg (N=2598)	Anastrozole 1 mg (N=3125)	Tamoxifen 20 mg (N=3116)	Anastrozole 1 mg (N=2618)	Tamoxifen 20 mg (N=2598)	Anastrozole 1 mg (N=3125)	Tamoxifen 20 mg (N=3116)	Anastrozole 1 mg (N=2618)	Tamoxifen 20 mg (N=2598)
Disease free survival (# of events) Hazard ratio (2-sided 95% CI) p-value	318 0.83 (0.17 to 0.96) 0.01	379 0.83 (0.17 to 0.96) 0.01	217 0.78 (0.65 to 0.93) 0.005	272 0.78 (0.65 to 0.93) 0.005	575 0.87 (0.78 to 0.97) 0.01	651 0.87 (0.78 to 0.97) 0.01	424 0.83 (0.73 to 0.94) 0.005	497 0.83 (0.73 to 0.94) 0.005	817 0.90 (0.82 to 0.99) 0.0252	887 0.90 (0.82 to 0.99) 0.0252	619 0.85 (0.76 to 0.94) 0.0027	702 0.85 (0.76 to 0.94) 0.0027
Time to recurrence (# of events) Hazard ratio (2-sided 95% CI) p-value	240 0.80 (0.67 to 0.94) 0.009	298 0.80 (0.67 to 0.94) 0.009	153 0.73 (0.59 to 0.90) 0.004	204 0.73 (0.59 to 0.90) 0.004	402 0.79 (0.70 to 0.90) 0.0005	498 0.79 (0.70 to 0.90) 0.0005	282 0.74 (0.64 to 0.87) 0.0002	370 0.74 (0.64 to 0.87) 0.0002	538 0.81 (0.73 to 0.91) 0.0004	645 0.81 (0.73 to 0.91) 0.0004	391 0.76 (0.67 to 0.87) 0.0001	494 0.76 (0.67 to 0.87) 0.0001
Distant disease-free survival (# of events) Hazard ratio (2-sided 95% CI) p-value	267 0.89 (0.74 to 1.07) 0.2	299 0.89 (0.74 to 1.07) 0.2	185 0.86 (0.69 to 1.08) 0.1	212 0.86 (0.69 to 1.08) 0.1	500 0.94 (0.83 to 1.06) 0.3	530 0.94 (0.83 to 1.06) 0.3	370 0.93 (0.80 to 1.07) 0.3	394 0.93 (0.80 to 1.07) 0.3	N/A N/A N/A	N/A N/A N/A	N/A N/A N/A	N/A N/A N/A
Contralateral breast primary (# of events) Odds ratio (2-sided 95% CI) p-value	14 0.42 (0.22 to 0.79) 0.007	33 0.42 (0.22 to 0.79) 0.007	11 0.36 (0.18 to 0.72) 0.004	30 0.36 (0.18 to 0.72) 0.004	35 0.59 (0.39 to 0.89) 0.01	59 0.59 (0.39 to 0.89) 0.01	26 0.47 (0.30 to 0.76) 0.002	54 0.47 (0.30 to 0.76) 0.002	61 0.68 (0.49 to 0.94) 0.02	87 0.68 (0.49 to 0.94) 0.02	50 0.60 (0.42 to 0.85) 0.004	80 0.60 (0.42 to 0.85) 0.004
Overall Survival (# of events) Hazard ratio (2-sided 95% CI) p-value	202 Not Calculated (not enough events at cut-off to conduct analysis)	203 Not Calculated (not enough events at cut-off to conduct analysis)	131 Not Calculated (not enough events at cut-off to conduct analysis)	136 Not Calculated (not enough events at cut-off to conduct analysis)	411 0.97 (0.85 to 1.12) 0.7	420 0.97 (0.85 to 1.12) 0.7	296 0.97 (0.83 to 1.14) 0.7	301 0.97 (0.83 to 1.14) 0.7	629 1.0 (0.89 to 1.12) 0.99	624 1.0 (0.89 to 1.12) 0.99	472 0.97 (0.86 to 1.11) 0.68	477 0.97 (0.86 to 1.11) 0.68
Time to distant recurrence (# of events) Hazard ratio (2-sided 95% CI) p-value	Not analyzed at the 33 month analysis.		Not analyzed at the 33 month analysis.		324 0.86 (0.74 to 0.99) 0.0427	375 0.86 (0.74 to 0.99) 0.0427	226 0.84 (0.74 to 0.99) 0.0559	265 0.84 (0.74 to 0.99) 0.0559	424 0.86 (0.75 to 0.98) 0.022	487 0.86 (0.75 to 0.98) 0.022	305 0.84 (0.72 to 0.97) 0.022	357 0.84 (0.72 to 0.97) 0.022

N = Number of patients randomized to the treatment

N/A = Not Available

Table 12: All recurrence and death events (data cut-off 31 March 2007; median follow-up 100 months)

	Number (%) of patients			
	Intention-to-treat population		Hormone receptor positive population	
	Anastrozole 1 mg (N=3125)	Tamoxifen 20 mg (N=3116)	Anastrozole 1 mg (N=2618)	Tamoxifen 20 mg (N=2598)
Loco-regional recurrence^{a,b}	155 (5.0)	184 (5.9)	102 (3.9)	130 (5.0)
Chest Wall	56 (1.8)	66 (2.1)	37 (1.4)	47 (1.8)
Ipsilateral breast ^c	56 (1.8)	73 (2.3)	38 (1.5)	56 (2.2)
Axillary lymph nodes	27 (0.9)	39 (1.3)	20 (0.8)	28 (1.1)
Other regional nodes ^d	31 (1.0)	43 (1.4)	16 (0.6)	28 (1.1)
Contralateral recurrence^e	61 (2.0)	87 (2.8)	50 (1.9)	80 (3.1)
Invasive	42 (1.3)	68 (2.2)	36 (1.4)	63 (2.4)
Ductal Carcinoma in situ	15 (0.5)	9 (0.3)	10 (0.4)	8 (0.3)
Unknown	4 (0.1)	10 (0.3)	4 (0.2)	9 (0.3)
Distant Recurrence^a	333 (10.7)	389 (12.5)	250 (9.5)	294 (11.3)
Bone/soft tissue	213 (6.8)	226 (7.3)	170 (6.5)	182 (7.0)
Bone	208 (6.7)	223 (7.2)	166 (6.3)	180 (6.9)
Soft tissue	8 (0.3)	8 (0.3)	6 (0.2)	6 (0.2)
Visceral	239 (7.6)	290 (9.3)	165 (6.3)	215 (8.3)
Pulmonary	110 (3.5)	140 (4.5)	78 (3.0)	95 (3.7)
Hepatic	82 (2.6)	144 (4.6)	61 (2.3)	113 (4.3)
Other	74 (2.4)	81 (2.6)	49 (1.9)	63 (2.4)
Death from Any Cause	629 (20.1)	624 (20.0)	472 (18.0)	477 (18.4)
Deaths following recurrence	350 (11.2)	382 (12.3)	245 (9.4)	269 (10.4)
Deaths without recurrence	279 (8.9)	242 (7.8)	227 (8.7)	208 (8.0)

a Patients may fall into more than one category.

b Patients who presented with distant recurrence or new primary (contralateral) breast cancer on the same day as loco-regional recurrence are included in this table but were counted either as distant recurrence or new primary (contralateral) breast cancer, respectively, in the summary of DFS (protocol-specified definition).

c Includes ductal carcinoma in situ and ipsilateral new breast primaries.

d Includes supraclavicular and internal mammary.

e Any new primary breast cancers occurring after loco-regional recurrence or distant recurrence were not included in this variable.

N= Number of patients randomized to the treatment.

Adjuvant treatment of breast cancer in postmenopausal women – assessment of bone

In the phase III/IV SABRE trial, 234 postmenopausal women with hormone receptor positive early breast cancer scheduled for treatment with anastrozole were stratified to low (T-score in both lumbar spine and total hip of -1.0 or higher and with no personal history of fragility fracture), moderate (T-score < -1.0 in either lumbar spine or total hip, provided neither of these was less than -2.0, and with no personal history of a fragility fracture) and high-risk (T-score < -2.0 in either lumbar spine, or total hip, or with a personal history of fragility fracture) groups. All patients received treatment with vitamin D and calcium. Patients in the low-risk group received anastrozole alone, those in the moderate group were randomised to

anastrozole plus bisphosphonate (risedronate) or anastrozole plus placebo and those in the high-risk group received anastrozole plus bisphosphonate (risedronate). The primary variable of the SABRE trial was the change from baseline in lumbar spine (L1-L4) bone mineral density (BMD) following 12 months of treatment. Secondary variables were changes in total hip BMD at 12 and 24 months as well as lumbar spine BMD at 24 months.

In postmenopausal breast cancer patients with a high-risk of fragility fracture, treatment with anastrozole and risedronate was associated with a statistically significant increase from baseline in lumbar spine BMD at 12 months (estimate percentage change 3.36%; 95% CI: 2.05, 4.69; $p > 0.0001$) and 24 months (estimated percentage change 3.02%; 95% CI: 1.40, 4.67; $p = 0.0006$).

In postmenopausal breast cancer patients with a moderate-risk of fragility fracture, treatment with anastrozole and risedronate resulted in a statistically significant increase in lumbar spine BMD at 12 months compared with anastrozole and placebo treatment (estimated percentage change 1.20% versus -1.22%; treatment ratio 1.02; 95% CI: 1.01, 1.04; $p < 0.001$) and at 24 months (estimated percentage change 2.24% versus -1.76%; treatment ratio 1.04; 95% CI: 1.02, 1.06; $p < 0.0001$).

In postmenopausal breast cancer patients with a low-risk of fragility fracture, treatment with anastrozole monotherapy was associated with no statistically significant change in lumbar spine BMD at 12 months (estimated percentage change -0.62%; 95% CI: -1.93, 0.71; $p = 0.3511$). The change in lumbar spine BMD at 24 months was statistically significant (estimated percentage change -2.07%; 95% CI: -3.60, -0.53, $p = 0.0109$).

Table 13: Analysis of lumbar spine BMD (g/cm^2) at 12 & 24 months in the high-risk, moderate-risk and low-risk strata (PAP)

High-risk stratum: analysis of change from baseline							
Anastrozole + Risedronate							
N ^a	Baseline gmean (g/cm^2)	gmean at time point (g/cm^2)	Estimated % change ^b (95% CI)	Time effect ^c (95% CI)	p-value		
12 months							
36	0.84	0.87	3.36 (2.05, 4.69)	1.03 (1.02, 1.05)	< 0.0001		
24 months							
33	0.83	0.86	3.02 (1.40, 4.67)	1.03 (1.01, 1.05)	0.0006		
Moderate-risk stratum: randomized comparison							
	N ^a	Baseline gmean (g/cm^2)	gmean at time point (g/cm^2)	Estimated % change ^{b,d} (95% CI)	glsmean ^d (g/cm^2)	Treatment ratio ^e (95% CI)	p-value ^d
12 months							
Anastrozole + placebo	65	0.98	0.97	-1.22 (-2.19, -0.24)	0.99		
Anastrozole + risedronate	73	0.98	1.00	1.20 (0.22, 2.19)	1.01	1.02 (1.01, 1.04)	<0.0001
24 months							
Anastrozole + placebo	54	0.96	0.95	-1.76 (-3.25, -0.25)	0.98		
Anastrozole + risedronate	60	0.98	1.00	2.24 (0.73, 3.76)	1.02	1.04 (1.02, 1.06)	<0.0001

Low-risk stratum: analysis of change from baseline Anastrozole monotherapy					
N ^a	Baseline gmean (g/cm ²)	gmean at time point (g/cm ²)	Estimated % change ^b (95% CI)	Time effect ^c (95% CI)	p-value
12 months					
35	1.15	1.14	-0.62 (-1.93, 0.71)	0.99 (0.98, 1.01)	0.3511
24 months					
26	1.15	1.12	-2.07 (-3.60, -0.53)	0.98 (0.96, 0.99)	0.0109

a Patients with values at baseline and 12 month visit

b 100*(time effect)-1

c Ratio of post baseline value/baseline value

d Covariance analysis

e Anastrozole + risedronate/anastrozole + placebo

BMD: Bone Mineral Density; CI: Confidence Interval; glsmean: geometric least square mean; gmean: geometric mean; PAP: Primary Analysis Population

In summary, the 12- and 24- month main analyses have shown that patients already at moderate to high-risk of fragility had their bone health (assessed by bone mineral density and bone formation and resorption markers) successfully managed by using anastrozole in combination with a bisphosphonate (risedronate). These findings were mirrored in the secondary efficacy variable of change from baseline in total hip BMD at 12 and 24 months (see Table 14). In addition, no changes in lumbar spine BMD were seen in the low risk group following 12 months of treatment with anastrozole alone and given vitamin D and calcium but were seen following 24 months of treatment. No change in total hip BMD was seen at 12 and 24 months in the low-risk group.

This study provides evidence that postmenopausal women with early breast cancer scheduled to be treated with anastrozole should have their bone status managed according to treatment guidelines already available for postmenopausal women at similar risk of fragility fracture.

Table 14: Analysis of total hip BMD (g/cm²) at 12 & 24 months in the high-risk, moderate-risk and low-risk strata (PAP)

High-risk stratum: analysis of change from baseline Anastrozole + Risedronate							
N ^a	Baseline gmean (g/cm ²)	gmean at time point (g/cm ²)	Estimated % change ^b (95% CI)	Time effect ^c (95% CI)	p-value		
12 months							
37	0.79	0.81	1.53 (0.37, 2.71)	1.02 (1.00, 1.03)	0.0112		
24 months							
33	0.80	0.81	1.96 (0.49, 3.44)	1.02 (1.00, 1.03)	0.0104		
Moderate-risk stratum: randomized comparison							
	N ^a	Baseline gmean (g/cm ²)	gmean at time point (g/cm ²)	Estimated % change ^{b,d} (95% CI)	glsmea n ^d (g/cm ²)	Treatment ratio ^e (95% CI)	p-value ^d
12 months							
Anastrozole + placebo	65	0.87	0.87	-0.44 (-1.17, 0.31)	1.00		
Anastrozole + risedronate	73	0.89	0.90	0.86 (0.12, 1.61)	1.01	1.01 (1.00, 1.02)	0.0023
24 months							

Anastrozole + placebo	54	0.87	0.86	-1.12 (-2.14, -0.10)	0.99		
Anastrozole + risedronate	60	0.90	0.92	1.81 (0.78, 2.86)	1.02	1.03 (1.02, 1.04)	<0.0001
Low-risk stratum: analysis of change from baseline Anastrozole monotherapy							
N ^a	Baseline gmean (g/cm ²)	gmean at time point (g/cm ²)	Estimated % change ^b (95% CI)	Time effect ^c (95% CI)	p-value		
12 months							
35	1.00	1.01	-0.35 (-1.37, 0.68)	1.00 (0.99, 1.01)	0.4918		
24 months							
26	1.01	1.00	-0.44 (-2.10, 1.26)	1.00 (0.98, 1.01)	0.5988		

a Patients with values at baseline and 12 month visit

b $100 * (\text{time effect}) - 1$

c Ratio of post baseline value/baseline value

d Covariance analysis

e Anastrozole + risedronate/anastrozole + placebo

BMD: Bone Mineral Density; CI: Confidence Interval; glsmean: geometric least square mean; gmean: geometric mean; PAP: Primary Analysis Population

Adjuvant treatment of breast cancer in postmenopausal women – assessment of lipids

In postmenopausal women with early breast cancer in the SABRE trial who received anastrozole alone (the primary analysis population), there was no statistically significant change in low-density lipoprotein-cholesterol (LDL-C) from baseline to 12 months (mean percent change -2.25% (95% CI: -7.64, 3.13) p-value 0.2859), a statistically significant increase in high-density lipoprotein-cholesterol (HDL-C) from baseline to 12 months (mean percent change 6.85% (95% CI: 2.79, 10.91) p-value 0.0016) and no statistically significant changes in total cholesterol or triglycerides (see Table 15).

In addition, no statistically significant changes from baseline to 12 months were seen in LDL-C [(mean percent change (-2.91% (95% CI: -7.20, 1.38) p-value 0.0770)], HDL-C [(mean percent change (4.00% (95% CI: 0.21, 7.79) p-value 0.1070)], total cholesterol or triglycerides in patients who received anastrozole in combination with the bisphosphonate, risedronate (the secondary analysis population)

The mean TC:HDL-C ratio decreased from baseline to 12 months in both populations for lipids. In the primary analysis population, the TC:HDL-C ratio decreased from a mean of 3.30 mmol/L (SD=0.82) at baseline to 3.11 mmol/L (SD=0.86) at 12 months while the secondary analysis population decreased from a mean of 3.48 mmol/L (SD=0.90) at baseline to 3.28 mmol/L (SD=0.85) at 12 months.

Table 15: Summary of lipid profile changes from baseline in LDL-C, HDL-C, total cholesterol and serum triglycerides (mmol/L) at 12 months

	Anastrozole 1 mg Population (PAPL) (N=66)	Anastrozole 1 mg + risedronate 35 mg Population (SP) (N=65)
LDL-C		
N ^a	54	59
Mean (baseline)	2.97	2.99
Mean (12 months)	2.88	2.89
Difference in means	-0.09	-0.11
Mean % change (95% CI)	-2.25 (-7.64, 3.13)	-2.91 (-7.20, 1.38)
p-value ^b	0.2859	0.0770
HDL-C		
N ^a	54	60
Mean (baseline)	1.68	1.62
Mean (12 months)	1.79	1.67
Difference in means	0.11	0.05
Mean % change (95% CI)	6.85 (2.79, 10.91)	4.00 (0.21, 7.79)
p-value ^b	0.0016	0.1070
Total Cholesterol (TC)		
N ^a	54	60
Mean (baseline)	5.25	5.24
Mean (12 months)	5.27	5.19
Difference in means	0.02	-0.05
Mean % change (95% CI)	0.76 (-3.08, 4.60)	-0.44 (-3.27, 2.39)
p-value ^b	0.8647	0.4840
Serum triglycerides (TG)		
N ^a	54	60
Mean (baseline)	1.31	1.40
Mean (12 months)	1.31	1.50
Difference in means	0.00	0.11
Mean % change (95% CI)	-0.60 (-7.15, 5.94)	7.03 (-5.02, 19.09)
p-value ^b	0.9881	0.4313

a Patients with values at baseline and 6 or 12 months

b Paired t-test comparing the means at baseline and 12 months

CI: Confidence Interval; ITT: Intent-To-Treat; LDL-C: Low-Density Lipoprotein-Cholesterol; PAPL: Primary Analysis Population for Lipids; SP: Secondary analysis population for lipids.

LDL-C, HDL-C, TC, TG and TC:HDL-C ratio were analyzed independently of strata in patients who did not have elevated cholesterol at baseline, according to the ATP [Adult Treatment Panel] III criteria.

Treatment for 12 months with anastrozole alone or in combination treatment with anastrozole and risedronate had a neutral effect on lipid profile. Therefore, no specific requirements for lipid monitoring due to anastrozole therapy are recommended.

Treatment of Postmenopausal Women with Advanced Breast Cancer

Anastrozole was studied in two, double-blind, controlled trials of similar design (0030, a North American study; 0027, a predominantly European study) in 1021 postmenopausal women with advanced breast cancer. Eligible patients were randomized to receive a single


daily dose of either anastrozole 1 mg, or tamoxifen 20 mg. The trials were designed to allow data to be pooled.

Demographics and other baseline characteristics were similar for the two treatment groups, however there were differences in hormone receptor status between the two trials. In Trial 0030, 88.3% of anastrozole-treated patients and 89.0% of tamoxifen-treated patients were known to be estrogen and/or progesterone receptor positive, compared to 45.3% and 43.9% (respectively) of patients in Trial 0027.


Anastrozole was shown to be at least as effective as tamoxifen for the primary endpoints of time to progression and objective response rate. In Trial 0030, a non-protocolled analysis indicated that anastrozole had a statistically significant advantage over tamoxifen ($p=0.005$) for time to progression (11.1 months versus 5.6 months, respectively) (see Figure 4a). Trial 0027 showed anastrozole to be at least as effective as tamoxifen for time to progression (8.2 months versus 8.3 months, respectively) (see Figure 4b) and objective response rate. The combined data from the two trials showed anastrozole to be numerically superior to tamoxifen for time to progression (8.5 months versus 7.0 months, respectively) (see Figure 4c). In a retrospective data analysis, patients from Trial 0027 who were known to be estrogen and/or progesterone receptor positive were shown to have longer median times to progression (271 days) when treated with anastrozole, than those treated with tamoxifen (237 days) (see Figure 4d). In addition, combined data from both trials, for patients who were estrogen and/or progesterone receptor positive, showed median times to progression of 10.7 months versus 6.4 months for anastrozole versus tamoxifen treated patients (two sided, $p=0.022$, retrospective analysis). These subgroup analyses support the results of Trial 0030 in suggesting numerical superiority for anastrozole over tamoxifen in patients known to be estrogen and/or progesterone receptor positive. Furthermore, these analyses demonstrate that patients with estrogen and/or progesterone receptor positive tumours are clearly the most appropriate population for anastrozole therapy.

Figure 4: Kaplan-Meier plots of time to progression (intention-to-treat population).
 a) Trial 0030 all patients; b) Trial 0027 all patients; c) Trials 0030 and 0027 combined; d) Trial 0027 estrogen/progesterone receptor positive patients only.


a)




b)



c)



d)



Results from the secondary endpoints of time to treatment failure, duration of response, and duration of clinical benefit were supportive of the results of the primary efficacy endpoints. The number of patients who experienced clinical benefit (best objective response of complete response [CR], partial response [PR] or stable disease [SD] \geq 24 weeks is shown in Table 16.

Table 16: Analysis of secondary endpoints in Trials 0030, 0027 and combined

Clinical Benefit	Number (%) of Patients					
	Trial 0030		Trial 0027		Combined Trials	
	Anastrozole 1 mg (n=171)	Tamoxifen 20 mg (n=182)	Anastrozole 1 mg (n=340)	Tamoxifen 20 mg (n=328)	Anastrozole 1 mg (n=511)	Tamoxifen 20 mg (n=510)
CR	5 (2.9)	5 (2.7)	19 (5.6)	16 (4.9)	24 (4.7)	21 (4.1)
PR	31 (18.1)	26 (14.3)	93 (27.4)	91 (27.7)	124 (24.3)	117 (22.9)
SD \geq 24 weeks	65 (38.0)	52 (28.6)	79 (23.2)	75 (22.9)	144 (28.2)	127 (24.9)
Total	101 (59.1)*	83 (45.6)*	191 (56.2)	182 (55.5)	292 (57.1)	265 (52.0)

CR complete response

PR partial response

SD stable disease

*two-sided p=0.0098, retrospective analysis

There were too few deaths occurring across treatment groups of both trials to assess overall survival differences at the time of data analysis.

Treatment of Postmenopausal Women with Advanced Breast Cancer who had Disease Progression following Tamoxifen Therapy

Anastrozole was studied in two well-controlled clinical trials (0004, a North American study; 0005, a predominantly European study) in postmenopausal women with advanced breast cancer who had disease progression following tamoxifen therapy. Most patients were estrogen receptor-positive; a smaller fraction was estrogen receptor-unknown or estrogen receptor-negative. Eligible patients were randomized to receive either a single daily dose of 1 mg or 10 mg of anastrozole, or megestrol acetate 40 mg four times a day. The studies were double-blinded with respect to anastrozole. Approximately 1/3 of the patients in each treatment group in both studies had either an objective response or stabilization of their disease for greater than 24 weeks. Hazard ratios for time to progression and odds ratios for response rates were calculated for the pooled studies were shown to be similar. After analysis of mature data involving 473 patients among 764 randomized participants, the hazard ratios for survival demonstrated a significant

prolongation of survival in the 1 mg anastrozole group compared to hormonal treatment with megestrol acetate.

Table 17: Analysis of time to death for patients in Trails 0004 and 0005 combined

Time of Death	Trial Treatment			Hazard ratio*, (97.5% CI), and p-values [#]	
	Anastrozole 1 mg	Anastrozole 10 mg	MA	Anastrozole 1mg vs. MA	Anastrozole 10 mg vs. MA
Number of patients who died (%)	151 of 263 (57.4)	151 of 248 (60.9)	171 of 253 (67.6)		
2-year survival rate	56.1%	54.6%	46.3%		
Median time to death (months)	26.7	25.5	22.5	0.78 (0.6040 to 0.9996) p=0.0248 ⁺	0.83 (0.6452 to 1.0662) p=0.0951 ⁺

* Hazard ratio greater than 1.00 indicated that the first treatment is associated with shorter time to death than is the second treatment

The critical p-value for statistical significance is 0.025

+ Calculated using Cox's regression model

CI: Confidence interval

MA: Megestrol acetate

Patients with estrogen receptor-negative disease rarely responded to anastrozole, but there were too few patients in this group for a meaningful analysis.

DETAILED PHARMACOLOGY

Animal Pharmacology

Pharmacodynamics

in vitro: Anastrozole inhibited human placental aromatase with an IC₅₀ (concentration inhibiting enzyme activity by 50%) of 15 nM; K_i values could not be calculated. It is therefore an inherently potent aromatase inhibitor.

in vivo: In rats, aromatase is not found in peripheral adipose tissue but is confined to the ovaries and brain so chronic inhibition of aromatase in this species invariably leads to marked compensatory ovarian changes. Two assessments of aromatase inhibitory activity in rats were therefore made in circumstances in which ovarian-hypothalamo-pituitary feedback effects were minimized, namely acute (single dose) inhibition of ovulation in adult rats, and

inhibition of uterine hypertrophy in sexually immature rats given exogenous androstenedione. Anastrozole consistently inhibited ovulation at a dose of 0.1 mg/kg. In this respect it was comparable in activity to fadrozole and 200 times as potent as aminoglutethimide, and it completely prevented the uterotrophic response to exogenous androstenedione at the same dose (given daily for three days).

As in the human, in Macaque monkeys aromatase is present in peripheral tissues in both males and females and the male monkey therefore affords an opportunity to assess chronic aromatase inhibitory activity in circumstances entirely analogous to the human male and comparable to postmenopausal women. Measurement of plasma estradiol concentrations in male pigtailed Macaques after ascending doses, each given for periods of 7 days, showed that anastrozole achieved maximal suppression at a dose of 0.1 mg/kg b.i.d. and was again comparable in potency to fadrozole.

Pharmacokinetics

Although species and gender-dependent effects are noted in anastrozole pharmacokinetics, anastrozole is rapidly and completely absorbed in all species evaluated. Elimination half-life of anastrozole is longer in humans (approximately 50 hours) than in animals and is independent of the anastrozole dose. Following administration of a single 1 mg/kg anastrozole dose, elimination half-life of anastrozole is approximately 10 hours in male dogs, 9 hours in female dogs, 7 hours in female rats and 2 hours in male rats. Consistent with its elimination half-life and with once a day dosing, 3-4 fold accumulation of anastrozole is observed in patients, while the accumulation pattern of anastrozole in rat and dog varies in a time and dose-dependent manner at doses greater than 5 mg/kg/day in the rat and 3 mg/kg/day in the dog.

Anastrozole is widely distributed into the tissues and is eliminated in both urine and bile in rats and dogs. Metabolism was qualitatively similar in rats, dogs and man, although a glucuronide conjugate of anastrozole was detectable in humans but not in rats or dogs. While some metabolites possess intrinsic aromatase inhibitory activity, they were not detectable in the plasma or were quantitatively minor metabolites (<5%). The results show that anastrozole itself is responsible for the observed pharmacological activity *in vivo*.

Adequate exposure to anastrozole and all metabolites except for the anastrozole glucuronide was achieved in the rat and dog relative to man. The anastrozole glucuronide is unlikely to possess pharmacological or toxicological activity.

TOXICOLOGY

The preclinical safety evaluation of anastrozole has included acute studies, 1 and 6-month toxicity studies in rats and dogs, teratology, genetic toxicology, antigenicity and irritancy studies. Additional toxicology studies include a 2-year oncogenicity study in rats and a 2-year

oncogenicity study in mice. Two additional investigative studies have also been completed to assist interpretation of the neoplastic changes observed in the rat oncogenicity study.

Acute Toxicity

The majority of mice dosed orally with 250 mg/kg anastrozole and all mice dosed intraperitoneally with 50 mg/kg showed signs of non-specific toxicity following dosing, but all recovered by day 2 and appeared normal for the remainder of the 14 day observation periods. Rats did not tolerate doses of 250 mg/kg and above by either route. No atypical signs were seen in rats following 100 mg/kg orally. However, there were signs of non-specific toxicity, but no deaths, following 50 mg/kg intraperitoneally. Non-specific toxicity in the rodent comprised the following: subdued behaviour, hunched posture, trembling, decreased respiration rate, fully or partially closed eyes, pilo-erection, salivation, lacrimation, convulsions, loss of skin tone, and lying prone.

In dogs treated orally with 45 mg/kg anastrozole, only minimally toxic effects were observed consisting of emesis, loose stools, body weight loss and reduced food consumption.

Multiple Dose Toxicity Studies

Anastrozole was well tolerated at up to 50 mg/kg/day in multiple dosing studies in rats, but 12 mg/kg/day was not tolerated in dogs in the 1 month study. Consequently, the top dose in the 6 month dog study was set at 8 mg/kg/day.

Anastrozole is a potent inhibitor of the aromatase enzyme and as such may be expected to induce a variety of effects resulting from the long-term inhibition of estrogen production in multiple dosing studies. Such pharmacological effects were observed in the reproductive tract and endocrine organs at all dose levels in rat and dog in both 1 month and 6 month toxicity studies. These effects included increased ovarian weight with increased numbers of Graafian follicles and/or corpora lutea together with mammary gland/uterine/vaginal changes in rats and dogs and testicular Leydig cell changes in dogs. Other pharmacologically induced changes in rats were reduced pituitary and adrenal gland weights, while in dogs, thymic involution was seen in both sexes in all dose groups. Changes in blood parameters included reversible increases in platelet numbers in both species, a reversible increase in erythrocyte parameters in female rats at 1 month, with a reversible decrease in male rats and dogs at 6 months, and increased white blood cells in rats of both sexes.

Non-pharmacologically induced changes in rats included an increased incidence of chronic progressive glomerular nephropathy at high dose (50 mg/kg/day) in the 6 month study. This was of minimal to mild severity and is thought to represent an exacerbation of the spontaneously occurring condition, possibly due to a slightly increased protein load in these animals. In addition, liver enlargement (reversible on withdrawal) accompanied by centrilobular hypertrophy and reduced glycogen at doses of 5 mg/kg/day and above in both

the 1 month and 6 month studies, was considered indicative of induction of mixed function oxidases by anastrozole.

In the dog, liver enlargement (reversible on withdrawal), generally accompanied by centrilobular hypertrophy and increased plasma alkaline phosphatase, was seen at mid and high dose levels in both multiple dose toxicity studies. This finding was consistent with induction of mixed function oxidase enzymes. Reversible hepatotoxicity, characterised by multifocal degeneration/necrosis and accompanied by elevated plasma alanine aminotransferase, was seen at the high dose (8 mg/kg/day) in the 6 month dog study. No degenerative changes were seen at the mid dose (3 mg/kg/day) in dogs, implying at least a 150 fold margin for hepatotoxicity in the dog based on a human dosage of 1 mg/day (approximately 0.02 mg/kg and approximately a 40 fold margin based on comparable AUC data).

Changes in clinical chemistry parameters in the toxicology studies included a reduction in triglycerides (all doses) and an increase in cholesterol (5 and 25 mg/kg/day) in male rats after 1 month dosing, and changes in potassium levels at 25 mg/kg/day. In dogs, plasma cholesterol and urinary creatine were reduced after 1 month at 12 mg/kg/day. Cholesterol was increased in female dogs (no change in males) after 6 months at 8 mg/kg/day. However, no ocular effects were seen in either species.

A reversible reduction in R-wave amplitude was seen in the dog studies at the high doses of 12 and 8 mg/kg/day in the 1 and 6 month studies respectively. This effect was not accompanied by any waveform interval or histopathological changes and is of unknown etiology.

Reproductive Toxicology

Anastrozole has been found to cross the placenta following oral administration of 0.1 mg/kg in rats and rabbits. Studies in both rats and rabbits at doses equal to or greater than 0.1 and 0.02 mg/kg/day, respectively, administered during the period of organogenesis showed that anastrozole increased pregnancy loss (increased pre- and/or post-implantation loss, increased resorption and decreased numbers of live fetuses). Effects were dose related in rats. Placental weights were significantly increased in rats at doses of 0.1 mg/kg/day or more.

Evidence of fetotoxicity, including delayed fetal development (i.e. incomplete ossification and depressed fetal body weights), was observed in rats administered doses of 1 mg/kg/day. There was no evidence of teratogenicity in rats administered doses up to 1 mg/kg/day. In rabbits, anastrozole caused pregnancy failure at doses equal to or greater than 1 mg/kg/day. There was no evidence of teratogenicity in rabbits administered 0.2 mg/kg/day.

Oral administration of anastrozole to female rats produced a high incidence of infertility at 1 mg/kg/day and increased pre-implantation loss at 0.02 mg/kg/day. These effects were related

to the pharmacology of the compound and were completely reversed after a 5-week compound withdrawal period.

Oncogenicity

The oncogenicity study in rats at doses of 1.0 to 25 mg/kg/day, administered by oral gavage for up to 2 years, revealed increases in the incidence of hepatocellular adenoma and carcinoma in high dose females, uterine stromal polyps in the high dose females and thyroid adenoma in the high dose males. Dose related increases were observed in the incidences of ovarian and uterine hyperplasia in females. At 25 mg/kg/day, plasma AUC_(0-24hr) levels in rats were about 100 times higher than the level exhibited in postmenopausal volunteers at the recommended dose.

A separate oncogenicity study in mice at oral doses of 5 to 50 mg/kg/day for up to 2 years, produced increases in the incidence of benign ovarian epithelial and sex cord stromal granulosa cell tumours, at all dose levels. A dose related increase in the incidence of ovarian stromal hyperplasia was also observed in female mice. These ovarian changes are considered to be rodent-specific effects of aromatase inhibition and are of no significance to postmenopausal breast cancer patients. The incidence of lymphosarcoma was marginally increased in males and females at the high dose. At 50 mg/kg/day, plasma AUC_(0-t) levels in mice were about 30 times higher than the level exhibited in postmenopausal volunteers at the recommended dose.

Other Toxicology Studies

There were no significant findings in genetic toxicology or in special toxicity studies designed to assess the irritant or antigenic potential of anastrozole.

Additional studies to provide further reassurance of the mechanisms underlying the formation of liver and thyroid tumours in rats have been completed.

In the first study, female rats dosed with anastrozole at 25 mg/kg/day for up to 28 days showed a 27% increase in relative liver weight, an increase in hepatocyte replication, and centrilobular hepatocyte hypertrophy. It was concluded that anastrozole, a known hepatic cytochrome P₄₅₀ enzyme inducer in rats, elicited a spectrum of biological changes in the rat liver similar to those observed with the non-genotoxic hepatocarcinogen, phenobarbitone. The hepatic changes in this study, and the tumours seen in female rats at 25 mg/kg/day after two-years, are considered a result of this non-genotoxic process.

In the second study, male rats were dosed at 25 mg/kg/day for 30 days. Thyroid follicular epithelial cell hypertrophy, increased TSH activity and increased plasma clearance of 125I-T₄, in association with liver enlargement, centrilobular hepatocyte hypertrophy, increase in CYP2B (predominantly) activity and increase in T₄ UDP-glucuronyltransferase activity, are consistent with anastrozole being a liver enzyme inducer of the phenobarbitone type. Thus, the thyroid tumours that occurred in male rats dosed with 25 mg/kg/day anastrozole over two-

years can be considered to be mechanistically related to an increased clearance of thyroid hormone resulting from an induction of specific liver enzymes resulting in a TSH-mediated non-genotoxic response.

The spectrum of biological changes in the rat liver and thyroid are similar to those reported in the literature following the administration of the non-genotoxic carcinogen, phenobarbitone. It is, therefore, concluded that the hepatic and thyroid changes seen in these investigative studies confirm the non-genotoxic mechanism responsible for the formation of tumours in the two-year rat oncogenicity study. The results do not alter the risk benefit assessment for the clinical use of anastrozole.

In support of clinical investigations using the combination of anastrozole and tamoxifen, an investigative study was performed in the rat to determine whether anastrozole, when administered in combination with tamoxifen, alters the metabolism of tamoxifen and the level of tamoxifen-DNA adducts in the rat liver. In the high dose group, where anastrozole produced a major increase in liver P₄₅₀ enzyme activity (specifically CYP2B and CYP3A), there was a significant reduction in the number of tamoxifen-DNA liver adducts compared to the animals given tamoxifen alone or in combination with a non-inducing dose of anastrozole. The plasma concentration of anastrozole was not determined but, in combination with the high dose of anastrozole, there was a reduction in the concentration of tamoxifen in plasma and liver, and a reduction in the concentration of tamoxifen metabolites in the liver.

REFERENCES

1. ATAC (Arimidex, Tamoxifen Alone or in Combination) Trialists' Group.
Anastrozole alone or in combination with tamoxifen versus tamoxifen alone for adjuvant treatment of postmenopausal women with early breast cancer: first results for the ATAC randomised trial. *Lancet* 2002; 359: 2131-2139.
2. ATAC Trialists's Group.
Anastrozole alone or in combination with tamoxifen versus tamoxifen alone for adjuvant treatment of postmenopausal women with early breast cancer: Results of the ATAC (Arimidex, Tamoxifen Alone or in Combination) Trial Efficacy and Safety Update Analyses. *Cancer* 2003; 98: 1802-1810.
3. ATAC Trialists' Group.
Results of the ATAC (Arimidex, Tamoxifen, Alone or in Combination) trial after completion of 5 years' adjuvant treatment for breast cancer. *Lancet* 2005; 365: 60-62.
4. ATAC Trialists' Group.
ATAC (Arimidex, Tamoxifen, Alone or in Combination): A phase III adjuvant therapy trial in postmenopausal breast cancer – 100 month follow-up results. *Lancet Oncol* 2008; 9(1):45-53.
5. Bonneterre J, Buzdar A, Nabholz JM et al.
Anastrozole is Superior to Tamoxifen as First-Line Therapy in Hormone Receptor Positive Advanced Breast Carcinoma. *Cancer* 2001; 92(9):2247-2258.
6. Bonneterre J, Thürlimann BJK, Robertson JFR et al.
Anastrozole Versus Tamoxifen as First-Line Therapy for Advanced Breast Cancer in 668 Postmenopausal Women: Results of the Tamoxifen or Arimidex Randomized Group Efficacy and Tolerability Study. *Journal of Clin Oncology* 2000; 18 (issue 22):3748-3757.
7. Bonneterre J, Thürlimann BJK, Robertson JFR.
Preliminary results of a large comparative multi-centre clinical trial comparing the efficacy and tolerability of Arimidex (anastrozole) and Tamoxifen (TAM) in postmenopausal women with advanced breast cancer (ABC). [abstract]. *Euro Journal of Cancer* 1999; 35 (suppl. 4): 313.
8. Brodie AMH.
Aromatase inhibitors in the treatment of breast cancer. *Journal of Steroid Biochemistry and Molecular Biology* 1994; 49(4-6):281-287.

9. Buzdar A, Jonat W, Howell A, Jones SE et al.
Anastrozole, a potent and selective aromatase inhibitor, versus megestrol acetate in postmenopausal women with advanced breast cancer: Results of overview analysis of two phase III trials. *Journal Clin Oncology* 1996; 14:2000-2011
10. Buzdar A, Jonat W, Howell A, Jones SE, et al.
Anastrozole versus megestrol acetate in the treatment of postmenopausal women with advanced breast carcinoma: results of a survival update based on a combined analysis of data from two mature phase III trials. *Cancer* 1998; 83 (8): 1142 -1152.
11. De la Cruz L, Romero-Vazquez, J, Jimenez-Saenz M, Padron JR, Herrerias- Gutierrez JM.
Severe acute hepatitis in a patient treated with anastrozole. *Lancet* 2007; 369:23-4.
12. Dowsett M, Yates RA, Lindsay A, Dukes M.
Endocrine effects of 'Arimidex', a potent aromatase inhibitor, in men and postmenopausal women. *Breast Cancer Research and Treatment* 1993; 27(1/2):152 Abs.
13. Dowsett M, Yates R, Wong YWJ.
'Arimidex' (anastrozole): Lack of interactions with tamoxifen, antipyrine, cimetidine and warfarin. *European Journal of Cancer* 1998;34 Suppl 1:S39-S40 Abs P100. Proceedings of the 6th International Conference on Adjuvant Therapy of Primary Breast Cancer, St. Gallen, 25-28 Feb 1998.
14. Duffy SR, Greenwood M.
The endometrial cancer data from the ATAC (Arimidex, Tamoxifen, Alone or in Combination) trial indicates a protective effect of anastrozole (Arimidex) upon the endometrium. *Breast Cancer Research and Treatment* 2003;82(Suppl 1):S29, Abs 134.
15. Eastell R.
Effect of anastrozole on bone mineral density: 2-year results of the 'Arimidex' (anastrozole), tamoxifen, alone or in combination (ATAC) trial. *Journal of Bone and Mineral Research* 2003;18(Suppl 2):S312, Abs M070.
16. Fallowfield L, Cella D.
Assessing the quality of life (QOL) of postmenopausal (PM) women randomized into the ATAC (Arimidex, Tamoxifen, Alone or in Combination) adjuvant breast cancer (BC) trial. *Program/proceedings - American Society of Clinical Oncology* 2002;21(1):40A, Abs 159.
17. Geisler J, Berntsen H, Ottestad L, Lindtjorn B, Dowsett M, Lonning PE.
Neoadjuvant treatment with anastrozole ('ARIMIDEX') causes profound suppression of intra-tumor estrogen levels [abstract]. *Proc ASCO* 1999;18: 82a, (Abstract 311).

18. Goss PE, Gwyn KMEH.
Current perspectives on aromatase inhibitors in breast cancer. *Journal Clin Oncology* 1994; 12(11):2460-2470.
19. Järhult J.
Anastrozole can cause severe hypercalcaemia mimicking primary hyperparathyroidism. *Breast Cancer*; 2011 Feb 3, DOI 10.1007/s12282-011-0253-x.
20. Locker GY, Eastell R
The time course of bone fractures observed in the ATAC ('Arimidex', Tamoxifen, Alone or in Combination) trial. Program/proceedings - American Society of Clinical Oncology 2003;22:25, Abs 98.
21. Nabholz JM.
Results of a North American, first-line trial comparing 'Arimidex' with tamoxifen in postmenopausal women with advanced breast cancer [abstract]. Presented at ECCO 10, the European Cancer Conference, Vienna, Austria; September 15, 1999.
22. Nabholz JM, Buzdar A, Pollak, M et al.
Anastrozole is Superior to Tamoxifen as First-Line Therapy for Advanced Breast Cancer in Postmenopausal Women: Results of a North American Multicenter Randomized Trial. *Journal of Clinical Oncology* 2000; 18 (issue 22): 3758-3767.
23. Pellegrini F, Rossi V, Fassone F, Castracane E, Persemoli L et al. Henoch-Schonlein purpura and aromatase inhibitors. *European Journal of Dermatology* 2009 ;19(5) :519 -20
24. Plourde PV, Dyroff M, Dukes M.
'Arimidex': A potent and selective fourth-generation aromatase inhibitor. *Breast Cancer Research and Treatment* 1994; 30(1):103-111.
25. Thürlimann BJK, Nabholz JM, Bonnetterre J, Buzdar AU, et al.
Preliminary results of two comparative multi-centre clinical trials comparing the efficacy and tolerability of Arimidex(r) (anastrozole) and tamoxifen (TAM) in postmenopausal women with advanced breast cancer (ABC). *Breast* 1999; 8 (4):214 Abs 004. 6th Nottingham International Breast Cancer Conference, 22-24 Sep 1999.
26. Vanden-Bossche H, Moereels H, Koymans LMH.
Aromatase inhibitors - mechanisms for non-steroidal inhibitors. *Breast Cancer Research and Treatment* 1994; 30(1):43-55.
27. Walton PL, Yates RA, Dukes M.

'Arimidex': an overview of a new selective nonsteroidal aromatase inhibitor. In: Motta M, Serio M, editors. Sex hormones and antihormones in endocrine dependent pathology: basic and clinical aspects. Proceedings of an international symposium; 1994 April 10-14; Milan, Italy. Excerpta Medica International Congress Series 1994(1064); 311-316.

28. Zapata E, Zubiaurre L, Bujanda L, Pierola A.
Anastrozole-induced hepatotoxicity. Eur J Gastroenterol Hepatol 2006; 7:1233-4.
29. AstraZeneca Canada Inc. ARIMIDEX[®] (anastrozole) Product Monograph, Control #170828, March 26, 2014

IMPORTANT: PLEASE READ

PART III: CONSUMER INFORMATION

Pr **Zinda-ANASTROZOLE** anastrozole

This leaflet is part III of a three-part "Product Monograph" published when Zinda-ANASTROZOLE was approved for sale in Canada and is designed specifically for Consumers. This leaflet is a summary and will not tell you everything about Zinda-ANASTROZOLE. Contact your doctor or pharmacist if you have any questions about the drug.

ABOUT THIS MEDICATION

What Zinda-ANASTROZOLE is used for:

Zinda-ANASTROZOLE is used for the treatment of postmenopausal women with hormone receptor positive breast cancer in the following conditions:

- adjuvant treatment for early breast cancer
- advanced breast cancer

What Zinda-ANASTROZOLE does:

In hormone sensitive breast cancer, estrogens fuel tumour growth. Following menopause, estrogens are still produced in small amounts in other tissues of the body such as the breasts, muscle and fat. These estrogens are produced when androgens (hormones produced by the adrenal glands) interact with aromatase, a naturally occurring enzyme in the body.

Zinda-ANASTROZOLE belongs to a group of medicines called aromatase inhibitors and works by inhibiting the aromatase enzyme, thereby, suppressing the production of estrogens that can stimulate tumour growth. Suppressing the production of estrogens may help reduce the growth of breast cancer and delay the breast cancer from recurring.

Adjuvant means "in addition to." In early breast cancer, this means that additional treatment is required after primary treatment. The reason for this is that after surgery, a small number of cancer cells may remain in the body. These cells can continue to multiply and spread. Adjuvant therapy is given to prevent or delay these cells from multiplying and spreading. The purpose of adjuvant therapy with Zinda-ANASTROZOLE is to help to delay the breast cancer from recurring. Cytotoxic chemotherapy, radiation, and hormonal treatment are three common forms of adjuvant treatment.

When Zinda-ANASTROZOLE should not be used:

- If you are allergic to the active ingredient anastrozole or any nonmedicinal ingredients of Zinda-ANASTROZOLE. If you think you may be allergic, ask your doctor for advice.
- If you are pregnant or breast-feeding.

What the medicinal ingredient is:

Anastrozole

What the nonmedicinal ingredients are:

Croscarmellose sodium, hypromellose, lactose monohydrate, macrogol, magnesium stearate, maltodextrin, and povidone

What dosage forms Zinda-ANASTROZOLE comes in:

Each Zinda-ANASTROZOLE tablet contains 1 milligram of anastrozole.

WARNINGS AND PRECAUTIONS

Serious Warnings and Precautions

- Zinda-ANASTROZOLE should not be given to premenopausal women.
- Zinda-ANASTROZOLE should not be given to children.
- Patients with liver and/or kidney problems, and patients with osteoporosis or at risk for osteoporosis should be carefully monitored by the doctor.
- Zinda-ANASTROZOLE should be prescribed by a doctor experienced in the use of anti-cancer drugs.

BEFORE you use Zinda-ANASTROZOLE talk to your doctor or pharmacist:

- If you have any disorder or disease which affects your heart, liver or kidneys.
- Zinda-ANASTROZOLE lowers the level of female hormones and this may lead to a loss of mineral content of bones, which might decrease their strength and lead to a broken bone. You should talk to your doctor about your osteoporosis risk before using Zinda-ANASTROZOLE.

Zinda-ANASTROZOLE tablets are unlikely to affect your ability to drive a car or to operate machinery. However, some patients may occasionally feel weak or sleepy. If this happens, you should not drive or operate machinery.

INTERACTIONS WITH THIS MEDICATION

BEFORE you use Zinda-ANASTROZOLE talk to your doctor or pharmacist:

- If you take medicine containing estrogen (a female sex hormone). It may oppose the effect of Zinda-ANASTROZOLE. Some herbal products contain estrogen.
- If you are currently taking tamoxifen.
- If you are taking or have recently taken other medicines, even those not prescribed by a doctor.

Please note that these statements may also apply to medicine used some time ago.

PROPER USE OF THIS MEDICATION

Usual dose:

Follow your doctor's instructions about when and how to take your Zinda-ANASTROZOLE tablets. The usual dose is one tablet once a day. Swallow the tablet with fluids. Try to take your tablet at the same time each day.

For adjuvant treatment of early breast cancer, currently it is recommended that Zinda-ANASTROZOLE be taken for 5 years.

Overdose:

In case of drug overdose, contact a health care practitioner, hospital emergency department or regional Poison Control Centre immediately, even if there are no symptoms.

Missed Dose:

Take the last missed dose as soon as you remember, as long as it is at least 12 hours before the next dose is due. If it is less than 12 hours to the next dose, do not take the dose you have missed.

SIDE EFFECTS AND WHAT TO DO ABOUT THEM

Like all medicines, Zinda-ANASTROZOLE can have side effects.

Contact your doctor immediately if any of the following happens to you. You may need further examinations or treatment:

- Severe skin reactions (Stevens-Johnson syndrome) with lesions, ulcers or blisters. This type of skin reaction is very rare.
- Allergic reaction with swelling of the face, lips, tongue and/or throat which may cause difficulty in swallowing and/or breathing.
- Chest pain or angina, as a result of ischemic heart disease (reduced blood flow in the vessels of the heart).
- Inflammation of the liver (hepatitis). Symptoms may include a general feeling of being unwell, with or without jaundice (yellowing of the skin and eyes) and pain in the upper abdomen on the right side.
- If you experience nausea, vomiting and thirst, you should tell your doctor. These symptoms may indicate possible increased blood calcium levels.

SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM

Symptom / effect	Talk with your doctor or pharmacist In all cases	Stop taking drug and seek immediate emergency medical attention
Very Common (greater than or equal to 10 of every 100 patients are likely to have these events)		
Hot flushes	✓	
Joint pain, joint stiffness or broken bones	✓	
Common (greater than or equal to 1 of every 100 patients, but less than 10 of every 100 patients, are likely to have these events)		
Weakness	✓	
Carpal tunnel syndrome (tingling, pain, coldness, weakness in parts of the hand)	✓	
Tickling, tingling or numbness of skin, loss/ lack of taste	✓	
Vaginal dryness	✓	
Hair thinning (alopecia)	✓	
Rash	✓	
Nausea	✓	
Diarrhea	✓	
Headache	✓	
Changes in blood tests of liver function	✓	
Bone pain	✓	
Muscle pain	✓	
Uncommon (greater than or equal to 1 of every 1000 patients, but less than 10 of every 1000 patients, are likely to have these events)		
Vaginal bleeding	✓	
Loss of appetite	✓	
High blood cholesterol	✓	
Vomiting	✓	
Sleepiness/tiredness	✓	
Trigger finger	✓	
Hepatitis	✓	✓
Very Rare (less than 1 of every 10 000 patients are likely to have these events)		
Sever skin reactions	✓	✓
Allergic reactions	✓	✓

This is not a complete list of side effects. For any unexpected effects while taking Zinda-ANASTROZOLE, contact your doctor or pharmacist.

HOW TO STORE IT

- Keep out of reach and sight of children.
- Store at room temperature, 15°C to 30°C. Keep container tightly closed.
- Keep your Zinda-ANASTROZOLE tablets in the

- original container.
- Do not use Zinda-ANASTROZOLE after the expiry date on the packaging.

REPORTING SUSPECTED SIDE EFFECTS

You can report any suspected adverse reactions associated with the use of health products to the Canada Vigilance Program by one of the following 3 ways:

- Report online at www.healthcanada.gc.ca/medeffect
- Call toll-free at 1-866-234-2345
- Complete a Canada Vigilance Reporting Form and:
 - Fax toll-free to 1-866-678-6789, or
 - Mail to: Canada Vigilance Program
Health Canada
Postal Locator 0701E
Ottawa, ON K1A 0K9

Postage paid labels, Canada Vigilance Reporting Form and the adverse reactions reporting guidelines are available on the MedEffect™ Canada Website at www.healthcanada.gc.ca/medeffect.

NOTE: Should you require information related to the management of side effects, contact your health professional. The Canada Vigilance Program does not provide medical advice.

MORE INFORMATION

This document plus the full product monograph, prepared for health professionals can be found by contacting the distributor, Zinda Pharma Limited, at

8250 Decarie Suite 110
Montreal, QC H4P 2P5

This leaflet was prepared by Stason Pharmaceuticals, Inc.

Last revised: October 10, 2014