

PRODUCT MONOGRAPH

Pr MYLAN-VALPROIC

Valproic Acid Capsules, USP

250 mg

Antiepileptic

Mylan Pharmaceuticals ULC
85 Advance Road
Etobicoke Ontario
M8Z 2S6

Date of Revision: May 7, 2015

Submission Control No.: 184065

Table of Contents

PART I: HEALTH PROFESSIONAL INFORMATION.....	3
SUMMARY PRODUCT INFORMATION	3
INDICATIONS AND CLINICAL USE	3
CONTRAINDICATIONS.....	4
WARNINGS AND PRECAUTIONS	5
ADVERSE REACTIONS	20
DRUG INTERACTIONS	24
DOSAGE AND ADMINISTRATION	34
OVERDOSAGE	37
ACTION AND CLINICAL PHARMACOLOGY	37
STORAGE AND STABILITY.....	40
DOSAGE FORMS, COMPOSITION AND PACKAGING.....	40
PART II: SCIENTIFIC INFORMATION	41
PHARMACEUTICAL INFORMATION.....	41
CLINICAL TRIALS	42
DETAILED PHARMACOLOGY.....	43
TOXICOLOGY.....	44
REFERENCES.....	47
PART III: CONSUMER INFORMATION.....	50

Pr **MYLAN-VALPROIC**
valproic acid

PART I: HEALTH PROFESSIONAL INFORMATION

SUMMARY PRODUCT INFORMATION

Route of Administration	Dosage Form / Strength	All Nonmedicinal Ingredients
Oral	Capsule / 250 mg	Corn Oil, D&C Yellow No. 10, FD&C Red No. 40, Gelatin, Glycerine, Purified Water and Titanium Dioxide.

INDICATIONS AND CLINICAL USE

Mylan-Valproic (valproic acid) is indicated for:

- use as sole or adjunctive therapy in the treatment of simple or complex absence seizures, including petit mal, and is useful in primary generalized seizures with tonic-clonic manifestations.
- use adjunctively in patients with multiple seizure types which include either absence or tonic-clonic seizures.

In accordance with the International Classification of Seizures, simple absence is defined as a very brief clouding of the sensorium or loss of consciousness (lasting usually 2 to 15 seconds), accompanied by certain generalized epileptic discharges without other detectable clinical signs. Complex absence is the term used when other signs are also present.

See **CONTRAINDICATIONS** and **WARNINGS AND PRECAUTIONS, Hepatic/Biliary/Pancreatic, Serious or Fatal Hepatotoxicity** for statement regarding serious or fatal hepatic dysfunction.

Geriatrics (≥ 65 years of age):

The safety and efficacy of valproic acid in elderly patients with epilepsy has not been evaluated in clinical trials. Caution should thus be exercised in dose selection for an elderly patient, recognizing the more frequent hepatic and renal dysfunctions, and limited experience with valproic acid in this population. For a brief discussion, see **WARNINGS AND PRECAUTIONS, Special Populations, Geriatrics (≥ 65 years of age), DOSAGE AND**

ADMINISTRATION and ACTION AND CLINICAL PHARMACOLOGY, Special Populations and Conditions, Geriatrics.

Pediatrics (< 18 years of age):

When valproic acid is used in children under the age of 2 years, it should be used with extreme caution and as a sole agent. Above the age of 2 years, experience in epilepsy has indicated that the incidence of fatal hepatotoxicity decreases considerably in progressively older patient groups. For a brief discussion, see **WARNINGS AND PRECAUTIONS, Special Populations, Pediatrics (< 18 years of age).**

CONTRAINDICATIONS

Mylan-Valproic (valproic acid) is contraindicated:

- in patients with hepatic disease or significant hepatic dysfunction. See **WARNINGS AND PRECAUTIONS, Serious Warnings and Precautions, Hepatotoxicity and WARNINGS AND PRECAUTIONS, Hepatic/Biliary/Pancreatic, Serious or Fatal Hepatotoxicity.**
- in patients known to have mitochondrial disorders caused by mutations in mitochondrial DNA polymerase γ (POLG; e.g., Alpers-Huttenlocher Syndrome) and children under two years of age who are suspected of having a POLG-related disorder. See **WARNINGS AND PRECAUTIONS, Hepatic/Biliary/Pancreatic.**
- in patients with known hypersensitivity to the drug, any ingredient in the formulation or component of the container. For a complete listing of ingredients, see the **DOSAGE FORMS, COMPOSITION AND PACKAGING** section of the Product Monograph.
- in patients with known urea cycle disorders. See **WARNINGS AND PRECAUTIONS, Endocrine and Metabolism, Urea Cycle Disorders.**
- in patients with known porphyria.

WARNINGS AND PRECAUTIONS

Serious Warnings and Precautions

- **Hepatotoxicity:** Hepatic failure resulting in fatalities has occurred in patients receiving valproic acid and its derivatives. These incidences usually occurred during the first 6 months of treatment with valproic acid. Caution should be observed when administering Mylan-Valproic to patients with a prior history of hepatic disease. Patients on multiple anticonvulsants, children, those with congenital metabolic disorders, those with severe seizure disorders accompanied by mental retardation, and those with organic brain disease may be at particular risk. Experience has indicated that children under the age of 2 years are at a considerably increased risk of developing fatal hepatotoxicity, especially those on multiple anticonvulsants. See **WARNINGS AND PRECAUTIONS, Hepatic/Biliary/Pancreatic, Serious or Fatal Hepatotoxicity.**
- **Teratogenicity:** Valproic acid can produce teratogenic effects such as neural tube defects (e.g., spina bifida) in a dose-dependent manner. Accordingly, the use of the medication in women of childbearing potential requires that the benefits of its use be weighed against the risk of injury to the fetus. This is especially important when Mylan-Valproic use is considered for a condition not usually associated with permanent injury or death. Women of childbearing potential should use an effective method of contraception during Mylan-Valproic therapy. Mylan-Valproic should only be used to treat pregnant women if other medications have failed to control their symptoms or are otherwise unacceptable. See **WARNINGS AND PRECAUTIONS, Special Populations, Women of Childbearing Potential and WARNINGS AND PRECAUTIONS, Special Populations, Pregnant Women.**
- **Patients with Mitochondrial Disease:** There is an increased risk of valproate-induced acute liver failure and resultant deaths in patients with hereditary neurometabolic syndromes caused by DNA mutations of the mitochondrial DNA Polymerase γ (POLG) gene (e.g. Alpers Huttenlocher Syndrome). Valproic acid is contraindicated in patients known to have mitochondrial disorders caused by POLG mutations and children under two years of age who are clinically suspected of having a mitochondrial disorder. See **CONTRAINDICATIONS.** In patients over two years of age who are clinically suspected of having a hereditary mitochondrial disease. Mylan-Valproic should only be used after other anticonvulsants have failed. This older group of patients should be closely monitored during treatment with Mylan-Valproic for the development of acute liver injury with regular clinical assessments and serum liver testing. POLG mutation screening should be performed in accordance with current clinical practice. See **WARNINGS AND PRECAUTIONS, Hepatic /Biliary/ Pancreatic, Patients with Mitochondrial Disease.**
- **Pancreatitis:** Cases of life-threatening pancreatitis have been reported in both children and adults receiving valproic acid. Some of the cases have been described as hemorrhagic with a rapid progression from initial symptoms to death. Patients and guardians should be warned that abdominal pain, nausea, vomiting and/or anorexia can be symptoms of pancreatitis that require prompt medical attention. If pancreatitis is diagnosed, Mylan-Valproic should ordinarily be discontinued. Alternative treatment for the underlying medical condition should be initiated as clinically indicated. Some cases have occurred shortly after initial use as well as after several years of use. See **WARNINGS AND PRECAUTIONS, Hepatic/Biliary/Pancreatic, Pancreatitis.**

General

Interaction with Carbapenem Antibiotics

Carbapenem antibiotics (ertapenem, imipenem, meropenem, doripenem) can reduce serum valproic acid concentrations to sub-therapeutic levels. This can result in loss of seizure control in epileptic patients or loss of efficacy in non-epileptics. In some cases of co-administration in epileptic patients, breakthrough seizures have occurred. Increasing valproic acid dose may not be sufficient to overcome this interaction. If co-administration is essential, serum valproic acid concentrations should be monitored daily after initiating carbapenem therapy. Alternative antibacterial or anticonvulsant therapy should be considered if serum valproic acid concentrations drop significantly or seizure control deteriorates. See **DRUG INTERACTIONS, Drug-Drug Interactions, Table 1.**

Effects of Valproate on HIV and CMV Viruses Replication

There are in vitro studies that suggest valproate stimulates the replication of the HIV (Human Immunodeficiency Virus) and CMV (Cytomegalovirus) viruses under certain experimental conditions. The clinical relevance of these in vitro data is unknown. Additionally, the relevance of these in vitro findings is uncertain for patients receiving maximally suppressive antiretroviral therapy. Nevertheless, these data should be borne in mind when interpreting the results from regular monitoring of the viral load in HIV infected patients receiving valproate or when following CMV infected patients clinically.

Carcinogenesis and Mutagenesis

Long-term animal toxicity studies indicate that valproic acid is a weak carcinogen or promoter in rats and mice. The significance of these findings for humans is unknown at present. See **TOXICOLOGY, Mutagenicity and Carcinogenicity.**

Endocrine and Metabolism

Urea Cycle Disorders

Mylan-Valproic is contraindicated in patients with known urea cycle disorders. Hyperammonemic encephalopathy, sometimes fatal, has been reported following initiation of valproic acid therapy in patients with urea cycle disorders, a group of uncommon genetic abnormalities, particularly ornithine transcarbamylase deficiency. Prior to initiation of Mylan-Valproic therapy, evaluation for urea cycle disorders (UCD) should be considered in the following patients:

- 1) those with a history of unexplained encephalopathy or coma, encephalopathy associated with protein load, pregnancy-related or postpartum encephalopathy, unexplained mental retardation, or history of elevated plasma ammonia or glutamine;
- 2) those with signs and symptoms of UCD, for example, cyclical vomiting and lethargy, episodic extreme irritability, ataxia, low blood urea nitrogen (BUN), protein avoidance;

- 3) those with a family history of UCD or a family history of unexplained infant deaths (particularly males);
- 4) those with other signs or symptoms of UCD. Patients receiving Mylan-Valproic therapy who develop symptoms of unexplained hyperammonemic encephalopathy should receive prompt treatment (including discontinuation of Mylan-Valproic therapy) and be evaluated for underlying urea cycle disorders. See **CONTRAINDICATIONS** and **WARNINGS AND PRECAUTIONS, Endocrine and Metabolism, Hyperammonemia and Hyperammonemia and Encephalopathy Associated with Concomitant Topiramate Use**.

Hyperammonemia

Hyperammonemia has been reported in association with valproic acid therapy and may be present despite normal liver function tests. In patients who develop unexplained lethargy and vomiting or changes in mental status, hyperammonemic encephalopathy should be considered as a possible cause and serum ammonia level should be measured. Hyperammonemia should also be considered in patients with hypothermia. See **WARNINGS AND PRECAUTIONS, Endocrine and Metabolism, Hypothermia**. If serum ammonia is increased, Mylan-Valproic therapy should be discontinued. Appropriate interventions for treatment of hyperammonemia should be initiated, and such patients should undergo investigation for underlying urea cycle disorders. See **CONTRAINDICATIONS** and **WARNINGS AND PRECAUTIONS, Endocrine and Metabolism Urea Cycle Disorders and Hyperammonemia and Encephalopathy Associated with Concomitant Topiramate Use**.

Asymptomatic elevations of serum ammonia are more common and, when present, require close monitoring of serum ammonia levels. If the elevation persists, discontinuation of Mylan-Valproic therapy should be considered.

Hyperammonemia and Encephalopathy Associated with Concomitant Topiramate Use

Concomitant administration of topiramate and valproic acid has been associated with hyperammonemia with or without encephalopathy in patients who have tolerated either drug alone. Clinical symptoms of hyperammonemic encephalopathy often include acute alterations in level of consciousness and/or cognitive function with lethargy or vomiting. Hypothermia can also be a manifestation of hyperammonemia. See **WARNINGS AND PRECAUTIONS, Endocrine and Metabolism, Hypothermia**. In most cases, symptoms and signs abated with discontinuation of either drug. This adverse event is not due to a pharmacokinetic interaction.

It is not known if topiramate monotherapy is associated with hyperammonemia.

Patients with inborn errors of metabolism or reduced hepatic mitochondrial activity may be at an increased risk for hyperammonemia with or without encephalopathy. Although not studied, an interaction of topiramate and Mylan-Valproic may exacerbate existing

defects or unmask deficiencies in susceptible persons. See CONTRAINDICATIONS and WARNINGS AND PRECAUTIONS, Endocrine and Metabolism, Urea Cycle Disorders and Hyperammonemia.

Hypothermia

Hypothermia, defined as an unintentional drop in core body temperature to $< 35^{\circ}\text{C}$ (95°F), has been reported in association with valproic acid both in conjunction with and in the absence of hyperammonemia. This adverse reaction can also occur in patients using concomitant topiramate with valproic acid after starting topiramate treatment or after increasing the daily dose of topiramate. See **DRUG INTERACTIONS, Drug-Drug Interactions, Table 1.** Hypothermia may be manifested by a variety of clinical abnormalities including, lethargy, confusion, coma, and significant alterations in other major organ systems such as the cardiovascular and respiratory systems. Clinical management and assessment should include examination of blood ammonia levels. Consideration should be given to stopping Mylan-Valproic in patients who develop hypothermia. See **WARNINGS AND PRECAUTIONS, Endocrine and Metabolism, Hyperammonemia.**

Hematologic

Thrombocytopenia

Because of reports of thrombocytopenia and inhibition of the second phase of platelet aggregation, and abnormal coagulation parameters (*e.g.*, low fibrinogen), platelet counts and coagulation tests are recommended before initiating therapy and at periodic intervals. It is recommended that patients receiving Mylan-Valproic be monitored for platelet count and coagulation parameters prior to planned surgery. Clinical evidence of hemorrhage, bruising or a disorder of hemostasis/coagulation is an indication for reduction of Mylan-Valproic dosage or withdrawal of therapy. See also **WARNINGS AND PRECAUTIONS, Hematologic, Dose-related Adverse Reactions: Thrombocytopenia.**

Dose-related Adverse Reactions: Thrombocytopenia

The frequency of adverse effects thrombocytopenia (particularly elevated liver enzymes and thrombocytopenia) may be dose-related. In a clinical trial of divalproex sodium as monotherapy in patients with epilepsy, 34/126 patients (27%) receiving approximately 50 mg/kg/day on average, had at least one value of platelets $\leq 75 \times 10^9/\text{L}$. Approximately half of these patients had treatment discontinued with return of platelet counts to normal. In the remaining patients, platelet counts normalized with continued treatment. In this study, the probability of thrombocytopenia appeared to increase significantly at total valproate concentrations of ≥ 110 mcg/mL (females) or ≥ 135 mcg/mL (males). The therapeutic benefit which may accompany the higher doses should therefore be weighed against the possibility of a greater incidence of adverse events.

In addition, the findings from a crossover clinical trial conducted with divalproex sodium extended-release tablets, in 44 epilepsy patients, indicate that the frequency of treatment-emergent mild thrombocytopenia (platelet count between 100 to 150 x 10⁹/L) was significantly higher after 12 weeks of treatment with divalproex sodium extended-release than after 12 weeks of treatment with divalproex sodium (7 versus 3 low counts, respectively).

Hepatic/Biliary/Pancreatic

Serious or Fatal Hepatotoxicity

Hepatic failure resulting in fatalities has occurred in patients receiving valproic acid and its derivatives. These incidences usually have occurred during the first 6 months of treatment with valproic acid. Caution should be observed when administering Mylan-Valproic products to patients with a prior history of hepatic disease. Patients on multiple anticonvulsants, children, those with congenital metabolic disorders, those with severe seizure disorders accompanied by mental retardation, and those with organic brain disease may be at particular risk.

Experience has indicated that children under the age of 2 years are at a considerably increased risk of developing fatal hepatotoxicity, especially those on multiple anticonvulsants, those with congenital metabolic disorders, those with severe seizure disorders accompanied by mental retardation, and those with organic brain disease. The risk in this age group decreased considerably in patients receiving valproic acid as monotherapy. Similarly, patients aged 3 to 10 years were at somewhat greater risk if they received multiple anticonvulsants than those who received only valproic acid. Above the age of 2 years, experience in epilepsy has indicated that the incidence of fatal hepatotoxicity decreases considerably in progressively older patients. No deaths have been reported in patients over 10 years of age who received valproic acid alone.

If Mylan-Valproic is to be used in children 2 years old or younger, it should be used with extreme caution and as a sole agent. The benefits of therapy should be weighed against the risk. See **WARNINGS AND PRECAUTIONS, Special Populations, Pediatrics (<18 years of age)**.

Serious or fatal hepatotoxicity may be preceded by non-specific symptoms such as loss of seizure control, malaise, weakness, lethargy, facial edema, anorexia and vomiting. Patients should be monitored closely for appearance of these symptoms. Patients and parents should be instructed to report such symptoms. Because of the non-specific nature of some of the early signs, hepatotoxicity should be suspected in patients who become unwell, other than through obvious cause, while taking Mylan-Valproic.

Liver function tests should be performed prior to therapy and at frequent intervals thereafter especially during the first 6 months. However, physicians should not rely totally on serum biochemistry since these tests may not be abnormal in all instances, but should also consider the results of careful interim medical history and physical examination.

In high-risk patients, it might also be useful to monitor serum fibrinogen and albumin for decreases in concentration and serum ammonia for increases in concentration. If changes occur, Mylan-Valproic should be discontinued. Dosage should be titrated to and maintained at the lowest dose consistent with optimal seizure control.

The drug should be discontinued immediately in the presence of significant hepatic dysfunction, suspected or apparent. In some cases, hepatic dysfunction has progressed in spite of discontinuation of drug. The frequency of adverse effects (particularly elevated liver enzymes and thrombocytopenia) may increase with increasing dose. The therapeutic benefit which may accompany the higher doses should therefore be weighed against the possibility of a greater incidence of adverse effects. See **CONTRAINDICATIONS**.

Patients with Mitochondrial Disease

Valproate induced acute liver failure and liver-related deaths have been reported in patients with hereditary neurometabolic syndromes caused by mutations in the gene for mitochondrial DNA polymerase γ (POLG) (e.g., Alpers-Huttenlocher Syndrome) at a higher rate than those without these syndromes. See **CONTRAINDICATIONS**.

POLG-related disorders should be suspected in patients with a family history or suggestive symptoms of a POLG-related disorder, including but not limited to unexplained encephalopathy, refractory epilepsy (focal, myoclonic), status epilepticus at presentation, developmental delays, psychomotor regression, axonal sensorimotor neuropathy, myopathy cerebellar ataxia, ophthalmoplegia, or complicated migraine with occipital aura. POLG mutation testing should be performed in accordance with current clinical practice for the diagnostic evaluation of such disorders. The A467T and W748S mutations are present in approximately 2/3 of patients with autosomal recessive POLG-related disorders.

In patients over two years of age who are clinically suspected of having a hereditary mitochondrial disease, Mylan-Valproic should only be used after other anticonvulsants have failed. This older group of patients should be closely monitored during treatment with Mylan-Valproic for the development of acute liver injury with regular clinical assessments and liver function test monitoring.

In the presence of significant hepatic dysfunction, suspected or apparent, Mylan-Valproic should be discontinued and alternative therapy initiated. In some cases, hepatic dysfunction has progressed in spite of discontinuation of the drug. See **CONTRAINDICATIONS** and **WARNINGS AND PRECAUTIONS**.

Pancreatitis

Cases of life-threatening pancreatitis have been reported in both children and adults receiving valproic acid. Some of the cases have been described as hemorrhagic with a rapid progression from initial symptoms to death. Some cases have occurred shortly after initial use as well as after several years of use. The rate based upon the reported cases exceeds that expected in the general population and there have been cases in which pancreatitis recurred after rechallenge with valproic acid. In clinical trials, there were 2 cases of pancreatitis without alternative etiology in 2,416 patients, representing 1,044 patient-years experience. Patients and guardians should be warned that abdominal pain, nausea,

vomiting, and/or anorexia can be symptoms of pancreatitis that require prompt medical evaluation. If pancreatitis is diagnosed, Mylan-Valproic should ordinarily be discontinued. Alternative treatment for the underlying medical condition should be initiated as clinically indicated.

Muscle Effects /Rhabdomyolysis

Rare cases of rhabdomyolysis, independent of neuroleptic malignant syndrome, have been reported to occur in patients treated with Mylan-Valproic. Cases have included renal failure and fatalities.

Patients should be carefully monitored for muscle pain, tenderness or weakness, particularly if accompanied by malaise or fever or tea-coloured urine. Blood creatine phosphokinase (CPK) levels should be assessed in patients experiencing these symptoms and Mylan-Valproic therapy should be discontinued if markedly elevated CPK levels are measured or if the patient develops signs and symptoms indicative of rhabdomyolysis.

Caution should be exercised in prescribing Mylan-Valproic to patients with predisposing/risk factors, including: prior history of muscular disorders such as CPT II deficiency (carnitine palmitoyltransferase type II); uncontrolled hypothyroidism; hepatic or renal impairment; concomitant medications that are known to be associated with rhabdomyolysis (eg statins, antipsychotics, diuretics, some antidepressants).

Neurologic

Brain Atrophy

There have been postmarketing reports of reversible and irreversible cerebral and cerebellar atrophy with neurological symptoms, in children, adults, and the elderly, receiving valproate therapy. A temporal relationship between valproate therapy and the development of cerebral atrophy and associated signs and symptoms was also demonstrated. In some cases, symptoms disappeared after valproate discontinuation but patients recovered with permanent sequelae. See **ADVERSE REACTIONS**. The motor and cognitive functions of patients on valproate should be routinely monitored and drug should be discontinued in the presence of suspected or apparent signs of brain atrophy.

Neurological Problems in Children after In-utero Exposure to Valproate

Reports of cerebral atrophy with various forms of neurological problems including developmental delays, psychomotor impairment and decreased IQ scores have been reported in children who were exposed in-utero to valproate products. See **WARNINGS AND PRECAUTIONS, Special Populations, Pregnant Women**.

Driving and Hazardous Occupations

Mylan-Valproic may produce central nervous system (CNS) depression, especially when combined with another CNS depressant, such as alcohol. Therefore, patients should be advised not to engage in hazardous occupations, such as driving a car or operating dangerous machinery, until it is known that they do not become drowsy from the drug.

Psychiatric

Suicidal Behaviour and Ideation

Suicidal ideation and behaviour have been reported in patients treated with antiepileptic agents in several indications.

All patients treated with antiepileptic drugs (AEDs), irrespective of indication, should be monitored for signs of suicidal ideation and behaviour and appropriate treatment should be considered. Patients (and caregivers of patients) should be advised to seek medical advice should signs of suicidal ideation or behaviour emerge.

An FDA meta-analysis of randomized placebo controlled trials, in which AEDs were used for various indications, has shown a small increased risk of suicidal ideation and behaviour in patients treated with these drugs. The mechanism of this risk is not known.

There were 43,892 patients treated in the placebo controlled clinical trials that were included in the meta-analysis. Approximately 75% of patients in these clinical trials were treated for indications other than epilepsy and, for the majority of non-epilepsy indications the treatment (AED or placebo) was administered as monotherapy. Patients with epilepsy represented approximately 25% of the total number of patients treated in the placebo controlled clinical trials and, for the majority of epilepsy patients, treatment (AED or placebo) was administered as adjunct to other antiepileptic agents (i.e., patients in both treatment arms were being treated with one or more AED). Therefore, the small increased risk of suicidal ideation and behaviour reported from the meta-analysis (0.43% for patients on AEDs compared to 0.24% for patients on placebo) is based largely on patients that received monotherapy treatment (AED or placebo) for non-epilepsy indications. The study design does not allow an estimation of the risk of suicidal ideation and behaviour for patients with epilepsy that are taking AEDs, due both to this population being the minority in the study, and the drug-placebo comparison in this population being confounded by the presence of adjunct AED treatment in both arms.

Behavioural Disorders

There have been postmarketing reports of behavioural disorders, including aggression, agitation, abnormal behaviour, disturbance in attention, and learning disorders. Although patients of all ages were affected, including the elderly and those exposed to valproate products in-utero, a large number of cases were reported in children. There was no clear trend with respect to valproate dose. In some cases, patients improved or recovered following valproate discontinuation.

Renal

Renal Impairment

Renal impairment is associated with an increase in the unbound fraction of valproate. In several studies, the unbound fraction of valproate in plasma from renally impaired patients was approximately double that for subjects with normal renal function. Accordingly, monitoring of total concentrations in patients with renal impairment may be misleading since free concentrations may be substantially elevated whereas total concentrations may appear to be normal. Hemodialysis in renally impaired patients may remove up to 20% of the circulating valproate.

Sensitivity/Resistance

Multi-organ Hypersensitivity Reaction

Multi-organ hypersensitivity reactions have been rarely reported in close temporal association to the initiation of valproic acid therapy in adult and pediatric patients (median time to detection 21 days; range 1 to 40). Although there have been a limited number of reports, many of these cases resulted in hospitalization and at least one death has been reported. Signs and symptoms of this disorder were diverse; however, patients typically, although not exclusively, presented with fever and rash associated with other organ system involvement. Other associated manifestations may include lymphadenopathy, hepatitis, liver function test abnormalities, hematological abnormalities (e.g., eosinophilia, thrombocytopenia, neutropenia), pruritus, nephritis, oliguria, hepato-renal syndrome, arthralgia, and asthenia. Because the disorder is variable in its expression, other organ system symptoms and signs, not noted here may occur. If this reaction is suspected, Mylan-Valproic should be discontinued and an alternative treatment started. Although the existence of cross sensitivity with other drugs that produce this syndrome is unclear, the experience amongst drugs associated with multi-organ hypersensitivity would indicate this to be a possibility.

Sexual Function/Reproduction

Fertility

The effect of valproic acid on the development of the testis and on sperm production and fertility in humans is unknown. See **TOXICOLOGY, Reproduction and Teratology, Fertility** for results in animal studies.

Skin

Serious Skin Reactions

The dose of lamotrigine should be reduced when co-administered with Mylan-Valproic. Serious skin reactions (such as Stevens-Johnson syndrome and Toxic Epidermal Necrolysis) have

been reported with concomitant lamotrigine and valproic acid administration (see Lamotrigine Product Monograph for details on lamotrigine dosing with concomitant valproic acid administration).

Special Populations

Women of Childbearing Potential

Because of the risk to the fetus of major congenital malformations (including neural tube defects) Mylan-Valproic should be considered for women of childbearing potential only after the risks have been thoroughly discussed with the patient and weighed against the potential benefits of treatment. See **WARNINGS AND PRECAUTIONS, Special Populations, Pregnant Women**.

This is especially important when Mylan-Valproic use is considered for a condition not usually associated with permanent injury or death.

Women of childbearing potential must use an effective method of contraception while using Mylan-Valproic.

Pregnant Women

Mylan-Valproic can cause fetal harm when administered to pregnant women. In comparison with some other antiepileptics drugs (AEDs), Valproic acid use during pregnancy is associated with an increased risk of severe birth defects such as neural tube defects (e.g., spina bifida), craniofacial defects, cleft palate, cardiovascular malformations, hypospadias, etc. See WARNINGS AND PRECAUTIONS, Special Populations, Pregnant Women, Birth Defects.

Since some of the congenital malformations occur in the first trimester of pregnancy before many women know that they are pregnant, all women of childbearing potential should be informed of the potential hazard to the fetus from exposure to Mylan-Valproic.

If the decision is made to use Mylan-Valproic, women who are not planning a pregnancy should be advised to use an effective method of contraception. If the woman is planning a pregnancy or becomes pregnant, risks of using Mylan-Valproic should be thoroughly discussed with the patient, weighed against the potential benefits of treatment, and alternative therapeutic options should be considered for these patients.

Women who are pregnant or who plan to become pregnant should not be treated with Mylan-Valproic unless other treatments have failed to provide adequate symptom control or are otherwise unacceptable. In such women, the benefits of treatment with Mylan-Valproic during pregnancy should be carefully weighed against the risk of malformation and other fetal abnormalities.

Antiepileptic drugs should not be abruptly discontinued in patients to whom the drug is administered to prevent major seizures, because of the strong possibility of precipitating status epilepticus with attendant hypoxia and risks to both the mother and the unborn child. With regard to drugs given for minor seizures, the risks of discontinuing medication prior to or during

pregnancy should be weighed against the risk of congenital defects in each particular case and with the particular family history. In individual cases where the severity and frequency of the seizure disorder are such that the removal of medication does not pose a serious threat to the patient, discontinuation of the drug may be considered prior to and/or during pregnancy, and alternative therapy considered, although it cannot be said with any confidence that even minor seizures do not pose some hazard to the developing embryo or fetus.

Women of childbearing age should be encouraged to seek the counsel of their physician and should report a positive pregnancy test promptly. Where the necessity for continued use of antiepileptic medication is in doubt, appropriate consultation is indicated. Current best practice guidelines should be considered in order to provide the optimal counsel to patients regarding the teratogenic risks associated with Mylan-Valproic.

Tests to detect neural tube and other defects using current accepted procedures should be considered a part of routine prenatal care in pregnant women receiving Mylan-Valproic.

Pregnancy Registry

Pregnant patients taking Mylan-Valproic should be encouraged to enroll in the North American Antiepileptic Drug (NAAED) Pregnancy Registry. This can be done by calling the toll free number 1-888-233-2334, and must be done by patients themselves. Information on the registry can also be found at the following website: <http://www.aedpregnancyregistry.org/>.

Birth Defects

There are multiple reports in the clinical literature that indicate the use of antiepileptic drugs during pregnancy results in an increased incidence of birth defects in the offspring. The incidence of congenital malformations in the general population is regarded to be approximately 2%; in children of treated epileptic women, this incidence may be increased 2- to 3-fold. The increase is largely due to specific defects such as congenital malformations of the heart, cleft lip and/or palate, craniofacial abnormalities and neural tube defects. Nevertheless, the great majority of mothers receiving antiepileptic medications deliver normal infants.

The data described below were gained almost exclusively from women who received valproate to treat epilepsy. Data from Pregnancy Registries indicate an increased risk of congenital anomalies in infants exposed to valproic acid monotherapy during the first trimester of pregnancy as compared to other antiepileptic drugs. Based on Pregnancy Registry data and the United States Centers for Disease Control (CDC), the estimated risk of valproate-exposed women having children with spina bifida, oral clefts, neural tube defects, and hypospadias is approximately 1 to 2% as compared to the risk of spina bifida in the general population which is about 0.06 to 0.07%.

In a study using NAAED Pregnancy Registry data, 16 cases of major malformations following prenatal valproate exposure were reported among offspring of 149 enrolled women who used valproate during pregnancy. Three of the 16 cases were neural tube defects; the remaining cases included craniofacial defects, cardiovascular malformations and malformations of

varying severity involving other body systems. The NAAED Pregnancy Registry has reported a major malformation rate of 10.7% in the offspring of women exposed to valproate monotherapy during pregnancy (average daily dose: 1000 mg; range: 500 – 2000 mg/day) as compared to major malformation rate of 2.9% among 1,048 epileptic women who received any other antiepileptic drug monotherapy during pregnancy. These data show a four-fold increased risk for any major malformation following valproate exposure in utero compared to the risk following exposure in utero to any other antiepileptic drug monotherapy.

Data from a meta-analysis (including registries and cohort studies) has shown an increased incidence of congenital malformations in children born to epileptic women exposed to valproate monotherapy during pregnancy. Available data indicate dose-dependency of this effect.

Risk of Neurological Problems from In-Utero Exposure

Cerebral Atrophy

Exposure in-utero to valproate products has been associated with cerebral atrophy with varying degrees/manifestations of neurological compromise, including developmental delays and psychomotor impairment. See **ADVERSE REACTIONS** and **WARNINGS AND PRECAUTIONS, Neurologic, Brain Atrophy**.

Developmental Delay, Decreased IQ, Autism and/or Autism Spectrum Disorders

There have been post-marketing reports of developmental delay, autism and/or autism spectrum disorder in the offspring of women exposed to Mylan-Valproic during pregnancy.

Valproate can cause decreased IQ scores following in utero exposure in a dose-dependent fashion. Published epidemiological studies have indicated that children exposed to valproate in utero have lower cognitive test scores than children exposed in utero to either another antiepileptic drug or to no antiepileptic drugs. Although all of the available studies have methodological limitations, the weight of the evidence supports the conclusion that in utero exposure to Mylan-Valproic can cause decreased IQ in children.

Withdrawal Syndrome in the Neonate

Withdrawal syndrome (such as, in particular, agitation, irritability, hyperexcitability, jitteriness, hyperkinesia, tonic disorders, tremor, convulsions and feeding disorders) may occur, in the days following birth, in neonates whose mothers have taken valproate during the last trimester of pregnancy.

Coagulation Abnormalities

There have been reported postmarketing cases of coagulation abnormalities in patients of all ages receiving valproate therapy. These include thrombocytopenia, hypofibrinogenemia, and/or decrease in other coagulation factors, which can lead to bleeding and other complications, especially in the cases of decrease in factors VII, VIII and XIII. These abnormalities may not necessarily be dose-dependent. Some of the hemorrhage manifestations may include mucosal

bleeding (e.g., menorrhagia, epistaxis, hematuria, melena), easy-bruising, soft-tissue hematoma, hemarthrosis, and intracranial hemorrhage. Caution should be taken in patients taking valproate and anticoagulants and in cases of injury or surgery to avoid life-threatening or fatal bleeding. See **WARNINGS AND PRECAUTIONS, Hematologic, Thrombocytopenia**.

Pregnant women taking valproic acid may also develop coagulation abnormalities, which may result in hemorrhagic complications in the neonate including death. See **WARNINGS AND PRECAUTIONS, Hematologic, Thrombocytopenia**. If Mylan-Valproic is used in pregnancy, the coagulation parameters should be monitored carefully.

Hepatic Failure

Hepatic failure, resulting in the death of a newborn and of an infant has been reported following the use of valproate during pregnancy.

Hypoglycemia

Serious postmarketing reports of hypoglycaemia have been received for neonates whose mothers received valproic acid treatment during pregnancy. In most cases, valproic acid was the only reported antiepileptic drug (AED). Most of these neonates also displayed other congenital anomalies including hypospadias, complex facial dysmorphism, limb anomalies, severe cardiac anomalies, etc. Therefore, if a decision has been made to use Mylan-Valproic during pregnancy, or if the patient becomes pregnant while taking this drug, the patient should be made aware of the potential hazard to the fetus.

Thyroid Gland Abnormalities

Cases of hypothyroidism have been reported in neonates whose mothers have taken valproate during pregnancy. There have also been reported cases of increased serum thyroid stimulating hormone or decreased serum thyroxine levels in children receiving valproate therapy. In addition, there have been reported cases of hypothyroidism and hyperthyroidism in adults and children receiving valproate monotherapy.

Teratogenicity in Animals

Animal studies have demonstrated valproic acid induced teratogenicity, see **TOXICOLOGY, Reproduction and Teratology**, and studies in human females have demonstrated placental transfer of the drug. Increased frequencies of malformations, as well as intrauterine growth retardation and death, have been observed in mice, rats, rabbits, and monkeys following prenatal exposure to valproate. Malformations of the skeletal system are the most common structural abnormalities produced in experimental animals, but neural tube closure defects have been seen in mice exposed to maternal plasma valproate concentrations exceeding 230 mcg/mL (2.3 times the upper limit of the human therapeutic range for epilepsy) during susceptible periods of embryonic development.

Administration of an oral dose of 200 mg/kg/day or greater (50% of the maximum human daily dose or greater on a mg/m² basis) to pregnant rats during organogenesis produced malformations (skeletal, cardiac and urogenital) and growth retardation in the offspring. These doses resulted in peak maternal plasma valproate levels of approximately 340 mcg/mL or greater (3.4 times the upper limit of the human therapeutic range for epilepsy or greater). Behavioural deficits have been reported in the offspring of rats given a dose of 200 mg/kg/day throughout most of pregnancy.

An oral dose of 350 mg/kg/day (approximately 2 times the maximum human daily dose on a mg/m² basis) produced skeletal and visceral malformations in rabbits exposed during organogenesis. Skeletal malformations, growth retardation, and death were observed in rhesus monkeys following administration of an oral dose of 200 mg/kg/day (equal to the maximum human daily dose on a mg/m² basis) during organogenesis. This dose resulted in peak maternal plasma valproate levels of approximately 280 mcg/mL (2.8 times the upper limit of the human therapeutic range for epilepsy).

Nursing Women

Valproic acid is secreted in breast milk. Concentrations in breast milk have been reported to be 1 to 10% of serum concentrations. As a general rule, nursing should not be undertaken while a patient is receiving Mylan-Valproic. Based on literature and clinical experience, hematological disorders in the newborn may occur.

Pediatrics (< 18 years of age)

Experience has indicated that children under the age of 2 years are at a considerably increased risk of developing fatal hepatotoxicity, especially those with the aforementioned conditions. See **WARNINGS AND PRECAUTIONS Hepatic/Biliary/Pancreatic, Serious or Fatal Hepatotoxicity**. When valproic acid is used in this patient group, it should be used with extreme caution and as a sole agent. The benefits of therapy should be weighed against the risks. See **CONTRAINDICATIONS**.

Above the age of 2 years, experience in epilepsy has indicated that the incidence of fatal hepatotoxicity decreases considerably in progressively older patient groups.

In patients over two years of age who are clinically suspected of having a hereditary mitochondrial disease, Mylan-Valproic should only be used after other anticonvulsants have failed. See **WARNINGS AND PRECAUTIONS, Hepatic/Biliary/Pancreatic, Patients with Mitochondrial Disease**.

Younger children, especially those receiving enzyme-inducing drugs, will require larger maintenance doses to attain targeted total and unbound valproate concentrations. The variability in free fraction limits the clinical usefulness of monitoring total serum valproate concentrations. Interpretation of valproate concentrations in children should include consideration of factors that affect hepatic metabolism and protein binding.

Geriatrics (≥ 65 years of age)

Alterations in the kinetics of unbound valproate in the elderly indicate that the initial dosage should be reduced in this population. See **DOSAGE AND ADMINISTRATION** and **ACTION AND CLINICAL PHARMACOLOGY, Special Populations and Conditions, Geriatrics**.

The safety and efficacy of valproic acid in elderly patients with epilepsy has not been evaluated in clinical trials. Caution should thus be exercised in dose selection for an elderly patient, recognizing the more frequent hepatic and renal dysfunctions, and limited experience with valproic acid in this population.

A study of elderly patients revealed valproate-related somnolence and discontinuation of valproic acid therapy for this adverse event. See **WARNINGS AND PRECAUTIONS, Special Populations, Geriatrics, Somnolence in the Elderly**. The starting dose should be reduced in elderly patients, and dosage reductions or discontinuation should be considered in patients with excessive somnolence. See **DOSAGE AND ADMINISTRATION**.

Somnolence in the elderly

In a group of elderly patients (mean age 83 years old, n = 172), valproic acid doses were increased by 125 mg/day to a target dose of 20 mg/kg/day. Compared to placebo a significantly higher number of valproate-treated patients had somnolence, and although not statistically significant, a higher number of valproate-treated patients experienced dehydration. Discontinuations for somnolence were also significantly higher in valproate-treated patients compared to placebo. In approximately one-half of the patients with somnolence, there was also associated reduced nutritional intake and weight loss. In elderly patients, dosage should be increased more slowly and with regular monitoring for fluid intake, dehydration, somnolence, urinary tract infection and other adverse events. Dose reductions or discontinuation of Mylan-Valproic should be considered in patients with decreased food or fluid intake and in patients with excessive somnolence. See **DOSAGE AND ADMINISTRATION**.

Monitoring and Laboratory Tests

Since Mylan-Valproic may interact with concurrently administered drugs which are capable of enzyme induction, periodic plasma concentration determinations of valproate and concomitant drugs are recommended during the early course of therapy and whenever enzyme-inducing drugs are introduced or withdrawn. See **DRUG INTERACTIONS**.

Monitoring Valproate Concentrations

Protein binding of valproate is reduced in the elderly, in patients with renal impairment, and in the presence of other drugs (e.g., acetylsalicylic acid). Accordingly, measurements of plasma levels of valproate may be misleading in these patients, as actual drug exposure may be higher than measured values. See **WARNINGS AND PRECAUTIONS, Hepatic/Biliary/Pancreatic**.

WARNINGS AND PRECAUTIONS, Endocrine and Metabolism, Hyperammonemia, WARNINGS AND PRECAUTIONS, Hematologic, Thrombocytopenia and DRUG INTERACTIONS, Drug-Drug Interactions, Table 1.

ADVERSE REACTIONS

Adverse Drug Reaction Overview

The most commonly reported adverse reactions are nausea, vomiting and indigestion. Since valproic acid has usually been used with other antiepileptics, it is not possible in most cases to determine whether the adverse reactions mentioned in this section are due to valproic acid alone or to the combination of drugs.

Adverse events that have been reported with valproic acid from epilepsy trials, spontaneous reports, and other sources are listed below by system organ class.

Blood and Lymphatic System Disorders:	Thrombocytopenia and inhibition of the secondary phase of platelet aggregation may be reflected in altered bleeding time, petechiae, bruising, hematoma formation, epistaxis, and hemorrhage. See WARNINGS AND PRECAUTIONS, <u>Hematologic</u>, Thrombocytopenia . Relative lymphocytosis, macrocytosis and hypofibrinogenemia have been noted. Leukopenia and eosinophilia have also been reported. Anemia, including macrocytic with or without folate deficiency, aplastic anemia, pancytopenia, bone marrow suppression, agranulocytosis and acute intermittent porphyria have been reported.
Cardiac Disorders:	Bradycardia.
Ear and Labyrinth Disorders:	Hearing loss, either reversible or irreversible, has been reported; however, a cause and effect relationship has not been established. Ear pain has also been reported.

Gastrointestinal Disorders:	<p>Nausea, vomiting and indigestion are the most commonly reported side effects at the initiation of therapy. These effects are usually transient and rarely require discontinuation of therapy. Diarrhea, abdominal cramps, constipation and gingival disorder (mainly gingival hyperplasia) have also been reported.</p> <p>There have been reports of acute pancreatitis, including rare fatal cases, occurring in association with valproic acid therapy. See WARNINGS AND PRECAUTIONS, <u>Hepatic/Biliary/Pancreatic, Pancreatitis.</u></p> <p>Parotid gland swelling has also been reported in patients receiving valproic acid.</p>
General Disorders and Administration Site Conditions Hepatobiliary Disorders:	<p>Edema of the extremities, fever and hypothermia</p> <p>Minor elevations of transaminases [e.g., serum glutamic-oxaloacetic transaminase (SGOT) and serum glutamic-pyruvic transaminase (SGPT)] and lactate dehydrogenase (LDH) are frequent and appear to be dose-related. Occasionally, laboratory tests also show increases in serum bilirubin and abnormal changes in other liver function tests. These results may reflect potentially serious hepatotoxicity. See WARNINGS AND PRECAUTIONS, Hepatic/Biliary/Pancreatic, Serious or Fatal Hepatotoxicity.</p>
Immune System Disorder:	Allergic reaction, anaphylaxis
Infections and Infestations:	Pneumonia and otitis media
Investigations:	Abnormal thyroid function tests (including both hyperthyroidism and hypothyroidism). See WARNINGS AND PRECAUTIONS, Special Populations. Pregnant Women, Thyroid Gland Abnormalities and (DRUG INTERACTIONS, Drug-Laboratory Interactions.
Metabolism and Nutrition Disorders:	<p>Hyperammonemia, see WARNINGS AND PRECAUTIONS, Endocrine and Metabolism, Hyperammonemia, hyponatremia and inappropriate antidiuretic hormone (ADH) secretion. There have been rare reports of Fanconi syndrome (proximal renal tubular dysfunction) occurring primarily in children. Decreased carnitine concentrations have been reported although the clinical relevance is undetermined. Hyperglycinemia has been reported and associated with a fatal outcome in patient with pre-existing nonketotic hyperglycinemia.</p> <p>Anorexia with some weight loss and increased appetite with some weight gain have also been reported.</p>

Musculoskeletal and
Connective Tissue
Disorders:

Weakness and bone pain have been reported (see WARNINGS AND PRECAUTIONS, Muscle Effects Rhabdomyolysis).

Reports have been received of decreased bone mass, potentially leading to osteoporosis and osteopenia, during long-term therapy with some anticonvulsant medications, including valproic acid. Some studies have indicated that supplemental calcium and vitamin D may be of benefit to patients who are on chronic valproic acid therapy.

A lupus erythematosus-like syndrome has been reported rarely.

Neoplasms Benign,
Malignant and Unspecified
(including cysts and
polyps):
Nervous System Disorders:

Myelodysplastic syndrome in both adults and children (all children were on valproate monotherapy). In some cases in adults and/or children, myelodysplastic syndrome was reversible upon valproate discontinuation.

Sedative effects have been noted in patients receiving valproic acid alone but occur most often in patients on combination therapy. Sedation usually disappears upon reduction of other anti-epileptic medication.

Hallucination, ataxia, headache, nystagmus, diplopia, asterixis, "spots before the eyes", tremor (may be dose-related), confusion, dysarthria, dizziness, hypesthesia, vertigo, incoordination, memory impairment, cognitive disorder, and parkinsonism have been noted. Rare cases of coma have been reported in patients receiving valproic acid alone or in conjunction with phenobarbital.

Encephalopathy, with or without fever or hyperammonemia, has been reported without evidence of hepatic dysfunction or inappropriate valproate plasma levels. Most patients recovered, with noted improvement of symptoms, upon discontinuation of the drug.

There have been postmarketing reports of reversible and irreversible cerebral and cerebellar atrophy temporally associated with the use of valproate products. In some cases the patients recovered with permanent sequelae. See WARNINGS AND PRECAUTIONS, Neurologic, Brain Atrophy. Cerebral atrophy seen in children exposed to valproate in utero led to various forms of neurological events, including developmental delays and psychomotor impairment. See WARNINGS AND PRECAUTIONS, Special Populations, Pregnant Women.

Psychiatric Disorders:

Emotional upset, depression, psychosis, aggression, psychomotor hyperactivity, hostility, agitation, disturbance in attention, abnormal behaviour, learning disorder and behavioural deterioration.

See WARNINGS AND PRECAUTIONS, Psychiatric.

Renal and Urinary Disorders: Enuresis, acute renal failure, tubulointerstitial nephritis and urinary tract infection.

Reproductive System and Breast Disorders: There have been reports of irregular menses, secondary amenorrhea, breast enlargement and galactorrhea in patients receiving valproic acid. Hyperandrogenism (hirsutism, virilism, acnea, male pattern alopecia, and/or androgen increased).

There have been rare spontaneous reports of polycystic ovary disease. A cause and effect relationship has not been established.

Respiratory, Thoracic and Mediastinal Disorders: Increased cough, pleural effusion.

Skin and Subcutaneous Tissue Disorders: Transient and/or dose related alopecia (hair loss), hair disorders (such as hair texture abnormal, hair colour changes, increases hair growth abnormal), have been observed. Skin rash, photosensitivity, generalized pruritus, erythema multiforme, Stevens-Johnson syndrome, and petechiae have rarely been noted.

Rare cases of Toxic Epidermal Necrolysis (TEN) have been reported including a fatal case of a 6 month old infant taking valproic acid and several other concomitant medications. An additional case of Toxic Epidermal Necrolysis resulting in death was reported in a 35 year old patient with AIDS taking several concomitant medications and with a history of multiple cutaneous drug reactions.

Serious skin reactions have been reported with concomitant administration of lamotrigine and valproic acid. See DRUG INTERACTIONS, Drug-Drug Interactions, Table 1.

Cutaneous vasculitis has also been reported.

Adverse Events in Elderly Patients

In elderly patients (above 65 years of age), there were more frequent reports of accidental injury, infection, pain, and to a lesser degree, somnolence and tremor, when compared to patients 18 to 65 years of age. Somnolence and tremor tended to be associated with the discontinuation of valproic acid.

DRUG INTERACTIONS

Serious Drug Interactions

- Rare cases of coma have been reported in patients receiving valproic acid alone or in conjunction with phenobarbital. See **Drug-Drug Interactions, Table 1**.
- Serious skin reactions (such as Stevens-Johnson syndrome and Toxic Epidermal Necrolysis) have been reported with concomitant lamotrigine and valproic acid administration. See **Drug-Drug Interactions, Table 1**.

Overview

Valproic acid has been found to be a weak inhibitor of some P₄₅₀ isozymes, epoxide hydrase, and glucuronyl transferases.

Drugs that affect the level of expression of hepatic enzymes, particularly those that elevate levels of glucuronyl transferases (such as ritonavir; see **Table 1** below), may increase the clearance of valproate. For example, phenytoin, carbamazepine, and phenobarbital (or primidone) can double the clearance of valproate. Thus, patients on valproic acid monotherapy will generally have longer half-lives and higher concentrations than patients receiving polytherapy with antiepilepsy drugs.

In contrast, drugs that are inhibitors of cytochrome P₄₅₀ isozymes, such as antidepressants, may be expected to have little effect on valproate clearance because cytochrome P₄₅₀ microsomal mediated oxidation is a relatively minor secondary metabolic pathway compared to glucuronidation and beta-oxidation.

The concomitant administration of valproic acid with drugs that exhibit extensive protein binding (e.g., acetylsalicylic acid, carbamazepine, dicumarol, warfarin, tolbutamide, and phenytoin) may result in alteration of serum drug levels.

Since valproic acid may interact with concurrently administered drugs which are capable of enzyme induction, periodic plasma concentration determinations of valproate and concomitant drugs are recommended during the early course of therapy and whenever enzyme-inducing drugs are introduced or withdrawn.

Drug-Drug Interactions

Table 1 provides information about the potential influence of several commonly prescribed medications on valproic acid pharmacokinetics as well as the potential influence of valproic acid on the pharmacokinetics and pharmacodynamics of several commonly prescribed medications. The list is not exhaustive nor could it be, since new interactions are continuously being reported. Please note that drugs may be listed under specific name, family or pharmacologic class. Reading the entire section is recommended.

Table 1. Summary of Drug-Drug Interaction Studies Including Important Interactions, Non-clinically Important Interaction and No Observed Interactions			
Concomitant Drug	Ref	Effect	Clinical Comment
Acetaminophen	CT	↔ acetaminophen	Valproic acid had no effect on any of the pharmacokinetic parameters of acetaminophen when it was concurrently administered to three epileptic patients.
Acetylsalicylic Acid	CT	↑ valproate	A study involving the co-administration of acetylsalicylic acid at antipyretic doses (11 to 16 mg/kg) with valproic acid to pediatric patients (n=6) revealed a decrease in protein binding and an inhibition of metabolism of valproate. Valproate free fraction was increased 4-fold in the presence of acetylsalicylic acid compared to valproic acid alone. The beta-oxidation pathway consisting of 2-E-valproic acid, 3-OH-valproic acid, and 3-keto valproic acid was decreased from 25% of total metabolites excreted on valproic acid alone to 8.3% in the presence of acetylsalicylic acid. Caution should be observed when valproic acid is administered with drugs affecting coagulation, (e.g., acetylsalicylic acid and warfarin). See ADVERSE REACTIONS .
Alcohol	T	No pharmacokinetic (PK) interaction	Valproic acid may potentiate the CNS depressant action of alcohol.
Amitriptyline / Nortriptyline	CT	In general: ↓amitriptyline. ↓nortriptyline Rarely: ↑ amitriptyline ↑nortriptyline	Administration of a single oral 50 mg dose of amitriptyline to 15 normal volunteers (10 males and 5 females) who received valproic acid (500 mg twice daily) resulted in a 21% decrease in plasma clearance of amitriptyline and a 34% decrease in the net clearance of nortriptyline. Rare post-marketing reports of concurrent use of valproic acid and amitriptyline resulting in an increased amitriptyline and nortriptyline levels have been received. Concurrent use of valproic acid and amitriptyline has rarely been associated with toxicity. Monitoring of amitriptyline levels should be considered for patients taking Mylan-Valproic concomitantly with amitriptyline. Consideration should be given to lowering the dose of amitriptyline/nortriptyline in the presence of valproic acid.
Antacids	CT	↔ valproate	A study involving the co-administration of valproic acid 500 mg with commonly administered antacids (Maalox®, Trisogel, and Titalac™ - 160 milliequivalent doses) did not reveal any effect on the extent of absorption of valproic acid.

Table 1. Summary of Drug-Drug Interaction Studies Including Important Interactions, Non-clinically Important Interaction and No Observed Interactions			
Concomitant Drug	Ref	Effect	Clinical Comment
Other - Antipsychotics, Monoamine Oxidase Inhibitors MAO Inhibitors (MAOIs) and Tricyclic Antidepressants			In addition to enhancing (CNS) depression when used concurrently with valproic acid, antipsychotics, tricyclic antidepressants and MAOIs may lower the seizure threshold. Dosage adjustments may be necessary to control seizures.
Benzodiazepines			Valproic acid may decrease oxidative liver metabolism of some benzodiazepines, resulting in increased serum concentrations. See Table 1. Diazepam and Lorazepam.
Carbamazepine / carbamazepine-10,11-epoxide	CT	↓ CBZ ↑ CBZ-E ↓ valproate	Concomitant use of carbamazepine (CBZ) with valproic acid may result in decreased serum concentrations and half-life of valproate due to increased metabolism induced by hepatic microsomal enzyme activity. Monitoring of serum concentrations is recommended when either medication is added to or withdrawn from an existing regimen. Changes in the serum concentration of the 10,11-epoxide metabolite of carbamazepine, however, will not be detected by routine serum carbamazepine assay. Serum levels of carbamazepine decreased 17% while that of carbamazepine-10,11-epoxide (CBZ-E) increased by 45% upon co-administration of valproic acid and CBZ to epileptic patients.
Carbapenem Antibiotics		↓ valproate	Carbapenem antibiotics (ertapenem, imipenem, meropenem, doripenem) can reduce serum valproic acid concentrations to sub-therapeutic levels. This can result in loss of seizure control in epileptic patients or loss of efficacy in non-epileptics. In some cases of co-administration in epileptic patients, breakthrough seizures have occurred. Increasing valproic acid dose may not be sufficient to overcome this interaction. If co-administration is essential, serum valproic acid concentrations should be monitored daily. Alternative antibacterial or anticonvulsant therapy should be considered if serum valproic acid concentrations drop significantly or seizure control deteriorates. See WARNINGS AND PRECAUTIONS, General, Interaction with Carbapenem Antibiotics.

Table 1. Summary of Drug-Drug Interaction Studies Including Important Interactions, Non-clinically Important Interaction and No Observed Interactions			
Concomitant Drug	Ref	Effect	Clinical Comment
Chlorpromazine	CT	↑ valproate	A study involving the administration of 100 to 300 mg/day of chlorpromazine to schizophrenic patients already receiving valproic acid (200 mg twice daily) revealed a 15% increase in trough plasma levels of valproate. This increase is not considered clinically important.
Cimetidine	T	↑ valproate	Cimetidine may decrease the clearance and increase the half-life of valproic acid by altering its metabolism. In patients receiving valproic acid, serum valproic acid levels should be monitored when treatment with cimetidine is instituted, increased, decreased, or discontinued. The valproic acid dose should be adjusted accordingly.
Clonazepam	T	No PK interaction	The concomitant use of valproic acid and clonazepam may induce absence status in patients with a history of absence type seizures.
Clozapine	CT	No interaction	In psychotic patients (n=11), no interaction was observed when valproic acid was co-administered with clozapine.
Diazepam	CT	↑ diazepam	Valproate displaces diazepam from its plasma albumin binding sites and inhibits its metabolism. Co-administration of valproic acid (1500 mg daily) increased the free fraction of diazepam (10 mg) by 90% in healthy volunteers (n=6). Plasma clearance and volume of distribution for free diazepam were reduced by 25% and 20%, respectively, in the presence of valproate. The elimination half-life of diazepam remained unchanged upon addition of valproate.
Ethosuximide	CT	↑ ethosuximide	Valproate inhibits the metabolism of ethosuximide. Administration of a single ethosuximide dose of 500 mg with valproic acid (800 to 1,600 mg/day) to healthy volunteers (n=6) was accompanied by a 25% increase in elimination half-life of ethosuximide and a 15% decrease in its total clearance as compared to ethosuximide alone. Patients receiving valproic acid and ethosuximide, especially along with other anticonvulsants, should be monitored for alterations in serum concentrations of both drugs.

Table 1. Summary of Drug-Drug Interaction Studies Including Important Interactions, Non-clinically Important Interaction and No Observed Interactions

Concomitant Drug	Ref	Effect	Clinical Comment
Felbamate	CT	↑ valproate	A study involving the co-administration of 1,200 mg/day of felbamate with valproic acid to patients with epilepsy (n=10) revealed an increase in mean valproate peak concentration by 35% (from 86 to 115 mcg/mL) compared to valproic acid alone. Increasing the felbamate dose to 2,400 mg/day increased the mean valproate peak concentration to 133 mcg/mL (another 16% increase). A decrease in Mylan-Valproic dosage may be necessary when felbamate therapy is initiated. Lower doses of Mylan-Valproic may be necessary when used concomitantly with felbamate.
Haloperidol	CT	↔ valproate	A study involving the administration of 6 to 10 mg/day of haloperidol to schizophrenic patients already receiving valproic acid (200 mg twice daily) revealed no significant changes in valproate trough plasma levels.

Table 1. Summary of Drug-Drug Interaction Studies Including Important Interactions, Non-clinically Important Interaction and No Observed Interactions

Concomitant Drug	Ref	Effect	Clinical Comment
Lamotrigine	CT	<p>↑ lamotrigine ↓ valproate</p>	<p>The effects of valproic acid on lamotrigine were investigated in six healthy male subjects. Each subject received a single oral dose of lamotrigine alone and with valproic acid 200 mg every 8 hours for 6 doses starting 1 hour before the lamotrigine dose was given. Valproic acid administration reduced the total clearance of lamotrigine by 21% and increased the plasma elimination half-life from 37.4 hours to 48.3 hours ($p < 0.005$). Renal clearance of lamotrigine was unchanged. In a steady-state study involving 10 healthy volunteers, the elimination half-life of lamotrigine increased from 26 to 70 hours with valproic acid co-administration (a 165% increase).</p> <p>In a study involving 16 epileptic patients, valproic acid doubled the elimination half-life of lamotrigine. In an open labelled study, patients receiving enzyme inducing AEDs (<i>e.g.</i>, carbamazepine, phenytoin, phenobarbital, or primidone) demonstrated a mean lamotrigine plasma elimination half-life of 14 hours while the elimination half-life was 30 hours in patients taking valproic acid plus an enzyme inducing antiepileptic agent. The latter value is similar to the lamotrigine half-life during monotherapy indicating that valproic acid may counteract the effect of the enzyme inducer. If valproic acid is discontinued in a patient receiving lamotrigine and an enzyme inducing antiepileptic serum lamotrigine concentrations may decrease. Patients receiving combined antiepileptic therapy require careful monitoring when another agent is started, stopped or when the dose is altered.</p> <p>Serious skin reactions (such as Stevens-Johnson Syndrome and toxic epidermal necrolysis) have been reported with concomitant lamotrigine and valproic acid administration.</p>

Table 1. Summary of Drug-Drug Interaction Studies Including Important Interactions, Non-clinically Important Interaction and No Observed Interactions

Concomitant Drug	Ref	Effect	Clinical Comment
Lithium	CT	↔ lithium	<p>In a double-blind placebo-controlled multiple dose crossover study in 16 healthy male volunteers, pharmacokinetic parameters of lithium were not altered by the presence or absence of valproate. The presence of lithium, however, resulted in an 11 to 12% increase in the AUC and C_{max} of valproate. T_{max} was also reduced. Although these changes were statistically significant, they are not likely to have clinical importance.</p> <p>Co-administration of valproic acid (500 mg-twice daily) and lithium carbonate (300 mg three times daily) to normal male volunteers (n=16) had no effect on the steady-state kinetics of lithium.</p>
Lorazepam	CT	↑ lorazepam	<p>Concomitant administration of valproic acid (500 mg twice daily) and lorazepam (1 mg twice daily) in normal male volunteers (n=9) was accompanied by a 17% decrease in the plasma clearance of lorazepam. This decrease is not considered clinically important.</p>
Olanzapine	CT	↓ olanzapine	<p>Administration of a single 5 mg dose of olanzapine to 10 healthy, non-epileptic volunteers with divalproex sodium extended-release tablets 1000 mg QD did not affect olanzapine C_{max} and elimination half-life. However, olanzapine AUC was 35% lower in the presence of divalproex sodium extended-release tablets. The clinical significance of these observations is unknown.</p>

Table 1. Summary of Drug-Drug Interaction Studies Including Important Interactions, Non-clinically Important Interaction and No Observed Interactions

Concomitant Drug	Ref	Effect	Clinical Comment
Oral contraceptive Steroids	CT	No PK interaction	<p>Evidence suggests that there is an association between the use of certain antiepileptic drugs (AEDs) capable of enzyme induction and failure of oral contraceptives. One explanation for this interaction is that enzyme-inducing drugs effectively lower plasma concentrations of the relevant steroid hormones, resulting in unimpaired ovulation. However, other mechanisms, not related to enzyme induction, may contribute to the failure of oral contraceptives. Valproic acid is not a significant enzyme inducer and would not be expected to decrease concentrations of steroid hormones. However, clinical data about the interaction of valproic acid with oral contraceptives are minimal.</p> <p>Administration of a single-dose of ethinyloestradiol (50 mcg)/levonorgestrel (250 mcg) to 6 women on valproic acid (200 mg twice daily) therapy for 2 months did not reveal any pharmacokinetic interaction.</p>
Phenobarbital	CT	↑ phenobarbital	<p>Valproate was found to inhibit the metabolism of phenobarbital. Co-administration of valproic acid (250 mg twice daily for 14 days) with phenobarbital to normal subjects (n=6) resulted in a 50% increase in half-life and a 30% decrease in plasma clearance of phenobarbital (60 mg single-dose). The fraction of phenobarbital dose excreted unchanged increased by 50% in the presence of valproate.</p> <p>There is evidence for severe CNS depression, with or without significant elevations of barbiturate or valproate serum concentrations. All patients receiving concomitant barbiturate therapy should be closely monitored for neurological toxicity.</p> <p>Serum barbiturate concentrations should be obtained, if possible, and the barbiturate dosage decreased, if appropriate.</p>

Table 1. Summary of Drug-Drug Interaction Studies Including Important Interactions, Non-clinically Important Interaction and No Observed Interactions

Concomitant Drug	Ref	Effect	Clinical Comment
Phenytoin	CT	↑ phenytoin	Valproate displaces phenytoin from its plasma albumin binding sites and inhibits its hepatic metabolism. Co-administration of valproic acid (400 mg three times daily) with phenytoin (250 mg) in normal volunteers (n=7) was associated with a 60% increase in the free fraction of phenytoin. Total plasma clearance and apparent volume of distribution of phenytoin increased 30% in the presence of valproate. Both the clearance and apparent volume of distribution of free phenytoin were reduced by 25%. In patients with epilepsy, there have been reports of breakthrough seizures occurring with the combination of valproic acid and phenytoin. The dosage of phenytoin should be adjusted as required by the clinical situation.
Primidone	T	↑ phenobarbital	Primidone is metabolized into a barbiturate (phenobarbital), and therefore, may also be involved in a similar or identical interaction with valproic acid as phenobarbital.
Quetiapine			Co-administration of valproate and quetiapine may increase the risk of neutropenia/leucopenia.
Rifampin	CT	↓ valproate	A study involving the administration of a single dose of valproic acid (7 mg/kg) 36 hours after 5 nights of daily dosing with rifampin (600 mg) revealed a 40% increase in the oral clearance of valproate. Mylan-Valproic dosage adjustment may be necessary when it is co-administered with rifampin.
Ritonavir	C	↓ valproate	Reduction of therapeutic effect of valproate was observed in a patient with bipolar disorder with the initiation of HIV treatment with lopinavir/ritonavir, zidovudine, and lamivudine.
Selective Serotonin Re-Uptake Inhibitors (SSRIs)	C	↑ valproate	Some evidence suggests that SSRIs inhibit the metabolism of valproic acid, resulting in higher than expected levels of valproate.
Tolbutamide	T	↑ tolbutamide	From <i>in vitro</i> experiments, the unbound fraction of tolbutamide was increased from 20 to 50% when added to plasma samples taken from patients treated with valproic acid. The clinical relevance of this displacement is unknown.

Table 1. Summary of Drug-Drug Interaction Studies Including Important Interactions, Non-clinically Important Interaction and No Observed Interactions			
Concomitant Drug	Ref	Effect	Clinical Comment
Topiramate	CT	Effect unknown	<p><u>Hyperammonemia</u></p> <p>Concomitant administration of valproic acid and topiramate has been associated with hyperammonemia with and without encephalopathy. See CONTRAINDICATIONS and WARNINGS AND PRECAUTIONS), <u>Endocrine and Metabolism</u>, Urea Cycle Disorders, Hyperammonemia and Hyperammonemia and Encephalopathy Associated with Concomitant Topiramate Use.</p> <p><u>Hypothermia</u></p> <p>Concomitant administration of topiramate with valproic acid has also been associated with hypothermia in patients who have tolerated either drug alone. Blood ammonia levels should be measured in patients with reported onset of hypothermia. See WARNINGS AND PRECAUTIONS, <u>Endocrine and Metabolism</u>, Hypothermia.</p>
Warfarin	T	Effect unknown	<p>In an <i>in vitro</i> study, valproate increased the unbound fraction of warfarin by up to 32.6%. The therapeutic relevance of this is unknown, however, coagulation tests should be monitored if valproic acid therapy is instituted in patients taking anticoagulants.</p> <p>Caution is recommended when valproic acid is administered with drugs affecting coagulation. See ADVERSE REACTIONS.</p>
Zidovudine	CT	↑ zidovudine	<p>In 6 patients who were seropositive for HIV, the clearance of zidovudine (100 mg every 8 hours) was decreased by 38% after administration of valproate (250 or 500 mg every 8 hours); the half-life of zidovudine was unaffected.</p>
Legend: C = Case Study; CT = Clinical Trial; T = Theoretical			

Drug-Food Interactions

Co-administration of Mylan-Valproic with food should cause no clinical problems in the management of patients with epilepsy.

Drug-Herb Interactions

Interactions with herbal products have not been established.

Drug-Laboratory Interactions

Valproic acid is partially eliminated in the urine as a ketone-containing metabolite which may lead to a false interpretation of the urine ketone test.

There have been reports of altered thyroid function tests associated with valproic acid; the clinical significance of these is unknown. See **WARNINGS AND PRECAUTIONS, Special Populations, Pregnant Women, Thyroid Gland Abnormalities**.

Drug-Lifestyle Interactions

Refer to **WARNINGS AND PRECAUTIONS, Neurologic, Driving and Hazardous Occupations** for details.

DOSAGE AND ADMINISTRATION

Dosing Considerations

Patients receiving combined antiepileptic therapy require careful monitoring when another agent is started, stopped or when the dose is altered. See **DRUG INTERACTIONS**.

As the dosage of Mylan-Valproic is titrated upward, blood concentrations of phenobarbital and/or phenytoin may be affected. See **DRUG INTERACTIONS**.

AEDs should not be abruptly discontinued in patients in whom the drug is administered to prevent major seizures because of the strong possibility of precipitating status epilepticus with attendant hypoxia and threat to life.

Any changes in dosage and administration, or the addition or discontinuance of concomitant drugs, should ordinarily be accompanied by close monitoring of clinical status and valproate plasma concentrations.

When changing therapy involving drugs known to induce hepatic microsomal enzymes (e.g., carbamazepine) or other drugs with valproate interactions, see **DRUG INTERACTIONS**, it is advisable to monitor serum valproate concentrations.

Dosing in Elderly Patients

Due to a decrease in unbound clearance of valproate and possibly a greater sensitivity to somnolence in the elderly, the starting dose should be reduced. Dosage should be increased more slowly and with regular monitoring for fluid and nutritional intake, dehydration, somnolence, urinary tract infection, and other adverse events. Dose reductions or discontinuation of Mylan-Valproic should be considered in patients with decreased food or fluid intake and in patients with excessive somnolence. The ultimate therapeutic dose should be achieved on the basis of clinical response. See **WARNINGS AND PRECAUTIONS, Special Populations, Geriatrics (≥ 65 years of age)**.

Dose-Related Adverse Events

The frequency of adverse events (particularly elevated liver enzymes and thrombocytopenia) may be dose related. The probability of thrombocytopenia appears to increase significantly at total valproate concentration of ≥ 110 mcg/mL (females) or ≥ 135 mcg/mL (males). See **WARNINGS AND PRECAUTIONS, Hematologic, Dose-related Adverse Reactions: Thrombocytopenia**. Therefore, the benefit of improved therapeutic effect with higher doses should be weighed against the possibility of a greater incidence of adverse effects.

Recommended Dose and Dosage Adjustment

Mylan-Valproic (valproic acid) is administered orally. The recommended initial dosage is 15 mg/kg/day, increasing at one-week intervals by 5 to 10 mg/kg/day until seizures are controlled or side effects preclude further increases.

The maximum recommended dosage is 60 mg/kg/day. When the total daily dose exceeds 250 mg, it should be given in a divided regimen (**Table 2**).

Weight		Total Daily Dose (mg)	Number of 250 mg Capsules		
kg	lb		Dose 1	Dose 2	Dose 3
10 to 24.9	22 to 54.9	250	0	0	1
25 to 39.9	55 to 87.9	500	1	0	1
40 to 59.9	88 to 131.9	750	1	1	1
60 to 74.9	132 to 164.9	1,000	1	1	2
75 to 89.9	165 to 197.9	1,250	2	1	2

Therapeutic Blood Levels

A good correlation has not been established between daily dose, total serum valproate concentration and therapeutic effect. However, therapeutic valproate serum concentrations for most patients with epilepsy will range from 50 to 100 mcg/mL (350 to 700 micromole/L). Some patients may be controlled with lower or higher serum concentrations. See **WARNINGS AND PRECAUTIONS**.

Conversion from valproic acid to divalproex sodium

Divalproex sodium enteric-coated tablets dissociate to the valproate ion in the gastrointestinal tract. Divalproex sodium tablets are uniformly and reliably absorbed, however, because of the enteric coating, absorption is delayed by an hour when compared to valproic acid.

The bioavailability of both types of divalproex sodium tablets (divalproex sodium and divalproex sodium extended release) is equivalent to that of valproic acid capsules.

In patients previously receiving valproic acid therapy, divalproex sodium should be initiated at the same daily dosing schedule. After the patient is stabilized on divalproex sodium, a dosing schedule of two or three times a day may be elected in selected patients. Changes in dosage administration of valproate or concomitant medications should be accompanied by increased monitoring of plasma concentrations of valproate and other medications, as well as the patient's clinical status.

Missed Dose

The patient should not abruptly stop taking their medication because of the risk of increasing their seizures.

If the patient misses a dose, they should not try to make up for it by doubling up on their next dose. They should take their next regularly scheduled dose and try not to miss any more doses.

Administration

Mylan-Valproic may be taken with or without food.

Patients who experience gastrointestinal irritation may benefit from administration of the drug with food or by a progressive increase of the dose from an initial low level. The capsules should be swallowed without chewing to avoid local irritation of the mouth and throat. Co-administration of oral Mylan-Valproic with food should cause no clinical problems in the management of patients with epilepsy.

OVERDOSAGE

For management of a suspected drug overdose, contact your Poison Control Centre Immediately.

Overdosage with valproic acid may result in somnolence, muscular hypotonia, hyporeflexia, miosis, impaired respiratory function, hypotension, metabolic acidosis, heart block, deep coma and circulatory collapse/shock. Cases of intracranial hypertension related to cerebral oedema have been reported. Fatalities have been reported; however, patients have recovered from valproate levels as high as 2,120 mcg/mL.

The presence of sodium content in the valproate formulations may lead to hypernatremia when taken in overdose.

In a reported case of overdosage with valproic acid after ingesting 36 g in combination with phenobarbital and phenytoin, the patient presented in deep coma. An electroencephalogram (EEG) recorded diffuse slowing, compatible with the state of consciousness. The patient made an uneventful recovery.

In overdose situations, the fraction of drug not bound to protein is high and hemodialysis or tandem hemodialysis plus hemoperfusion may result in significant removal of drug. The benefit of gastric lavage or emesis will vary with the time since ingestion. As valproic acid is absorbed very rapidly, gastric lavage may be of limited value. General supportive measures should be applied with particular attention to the prevention of hypovolemia and the maintenance of adequate urinary output.

Naloxone has been reported to reverse the CNS depressant effects of valproic acid overdosage. Because naloxone could theoretically also reverse the antiepileptic effects of valproic acid, it should be used with caution.

ACTION AND CLINICAL PHARMACOLOGY

Pharmacotherapeutic group: Anticonvulsant and mood-stabilizing drug; ATC-Code: N03AG01.

Mechanism of Action

Valproic acid has anticonvulsant properties. Although its mechanism of action has not yet been established, it has been suggested that its activity is related to increased brain levels of gamma-aminobutyric acid (GABA). The effect on the neuronal membrane is unknown.

Pharmacodynamics

A good correlation has not been established between daily dose, serum level and therapeutic effect. In epilepsy, the therapeutic plasma concentrations range is believed to be from 50 to 100 mcg/mL (350 to 700 micromole/L) of total valproate. Occasional patients may be controlled with serum levels lower or higher than this range. See **DOSAGE AND ADMINISTRATION**.

Pharmacokinetics

Absorption

Valproic acid is rapidly absorbed after oral administration. Peak serum levels occur approximately 1 to 4 hours after a single oral dose. A slight delay in absorption occurs when the drug is administered with meals but this does not affect the total absorption.

Distribution

Valproic acid is rapidly distributed throughout the body and the drug is strongly bound (90%) to human plasma proteins. Increases in dose may result in decreases in the extent of protein binding and variable changes in valproic acid clearance and elimination.

Protein Binding

The plasma protein binding of valproate is concentration dependent and the free fraction increases from approximately 10% at 40 mcg/mL to 18.5% at 130 mcg/mL. Protein binding of valproate is reduced in the elderly, in patients with chronic hepatic diseases, in patients with renal impairment, in hyperlipidemic patients, and in the presence of other drugs (*e.g.*, acetylsalicylic acid). Conversely, valproate may displace certain protein-bound drugs (*e.g.*, phenytoin, carbamazepine, warfarin, and tolbutamide). See **DRUG INTERACTIONS** for more detailed information on the pharmacokinetic interactions of valproate with other drugs.

CNS Distribution

Valproate concentrations in cerebrospinal fluid (CSF) approximate unbound concentrations in plasma (ranging from 7 to 25% of total concentration).

Metabolism

Valproate is metabolized almost entirely by the liver. In adult patients on monotherapy, 30 to 50% of an administered dose appears in urine as a glucuronide conjugate. Mitochondrial (beta)-oxidation is the other major metabolic pathway, typically accounting for over 40% of the dose. Usually, less than 15 to 20% of the dose is eliminated by other oxidative mechanisms. Less than 3% of an administered dose is excreted unchanged in urine.

Due to the saturable plasma protein binding, the relationship between dose and total valproate concentration is nonlinear; concentration does not increase proportionally with the dose, but rather increases to a lesser extent. The kinetics of unbound drug are linear.

Excretion

Mean plasma clearance and volume of distribution for total valproate are 0.56 L/hr/1.73 m² and 11 L/1.73 m², respectively. Mean plasma clearance and volume of distribution for free valproate are 4.6 L/hr/1.73 m² and 92 L/1.73 m², respectively. These estimates cited apply primarily to patients who are not taking drugs that affect hepatic metabolizing enzyme systems. For example, patients taking enzyme-inducing AEDs (carbamazepine, phenytoin, and phenobarbital) will clear valproate more rapidly. Because of these changes in valproic acid clearance, monitoring of valproate and concomitant drug concentrations should be intensified whenever enzyme-inducing drugs are introduced or withdrawn.

Elimination of valproic acid and its metabolites occurs principally in the urine, with minor amounts in the feces and expired air. Very little unmetabolized parent drug is excreted in the urine.

The serum half-life ($t_{1/2}$) of valproic acid is typically in the range of 6 to 16 hours. Half-lives in the lower part of the above range are usually found in patients taking other AEDs capable of enzyme induction.

Special Populations and Conditions

Neonates/ Infants:

Within the first 2 months of life, infants have a markedly decreased ability to eliminate valproate compared to children and adults. This is a result of reduced clearance (perhaps due to delay in development of glucuronosyltransferase and other enzyme systems involved in valproate elimination) as well as increased volume of distribution (in part due to decreased plasma protein binding). For example, in one study, the half-life in neonates under 10 days ranged from 10 to 67 hours, compared to a range of 7 to 13 hours in children greater than 2 months.

Pediatrics

Patients between 3 months and 10 years have 50% higher clearances expressed on weight (i.e., L/min/kg) than do adults. Over the age of 10 years, children have pharmacokinetic parameters that approximate those of adults.

Geriatrics

The capacity of elderly patients (age range: 68 to 89 years) to eliminate valproate has been shown to be reduced compared to younger adults (age range: 22 to 26 years). Intrinsic clearance is reduced by 39%; the free fraction is increased by 44%. See **DOSAGE AND ADMINISTRATION**.

Gender

There are no differences in unbound clearance (adjusted for body surface area) between males and females (4.8 ± 0.17 and 4.7 ± 0.07 L/hr per 1.73 m^2 , respectively).

Race

The effects of race on the kinetics of valproate have not been studied.

Hepatic Insufficiency

See **CONTRAINDICATIONS and WARNINGS AND PRECAUTIONS, Hepatic/Biliary/Pancreatic, Serious or Fatal Hepatotoxicity** for statements regarding hepatic dysfunction and associated fatalities.

Renal Insufficiency

See **WARNINGS AND PRECAUTIONS, Renal, Renal Impairment**.

Genetic Polymorphism

No data available on genetic polymorphism.

STORAGE AND STABILITY

Store Mylan-Valproic capsules between 15°C and 30°C .

DOSAGE FORMS, COMPOSITION AND PACKAGING

Composition

Mylan-Valproic (valproic acid) Capsules contain 250 mg of Valproic Acid. The non-medical ingredients are: Corn Oil, D&C Yellow' No. 10, FD&C Red No. 40, Gelatin, Glycerine, Purified Water and Titanium Dioxide.

Availability of Dosage Forms

Mylan-Valproic (valproic acid): available as oblong, orange soft gelatin capsules with clear fill liquid, imprinted in black ink with "VA250" on one side. These capsules are available in bottles of 100 and 500.

PART II: SCIENTIFIC INFORMATION

PHARMACEUTICAL INFORMATION


Drug Substance

Proper name: valproic acid

Chemical name: 2-propylpentanoic acid or dipropylacetic acid

Molecular formula and molecular mass: $C_8H_{16}O_2$ 144.21 g/mol

Structural formula:



Physicochemical properties: Valproic acid is chemically a carboxylic acid designated as 2-propylpentanoic acid, also known as dipropylacetic acid. It is a colourless liquid, slightly soluble in water and very soluble in organic solvents.

CLINICAL TRIALS

Comparative Bioavailability Studies

A single-dose, crossover, comparative bioavailability study of Mylan-Valproic (Valproic Acid) Capsules 250 mg and Depakene[®] (Valproic Acid) Capsules 250 mg (Abbott, Canada) following a single-dose of 250 mg valproic acid in 21 subjects under fed conditions was conducted. The results indicate that Mylan-Valproic Capsules 250 mg are bioequivalent to Depakene[®] Capsules 250 mg. The summary of results is presented in the following table:

Valproic Acid (1 x 250 mg) From measured data				
Geometric Mean Arithmetic Mean (CV %)				

Parameter	Test [‡]	Reference ^{⊕†}	% Ratio of Geometric Means	95% Confidence Interval
AUC _T μg/ml•h	592.29 620 (30.5)	598.24 624 (30.2)	99	97 - 102
C _{max} μg/ml•h	26.0 26.3 (17.2)	26.2 26.5 (17.1)	99	95 - 103
T _{max} [§] (h)	4.48 (30.5)	4.64 (24.8)		
T _{1/2} [§] (h)	19.7 (32.2)	19.8 (29.4)		

[‡] Mylan-Valproic Capsules 250 mg were manufactured by Mylan Pharmaceuticals ULC

[†] Depakene[®] Capsules 250 mg was manufactured by Abbott Canada

[§] Expressed as the arithmetic mean (CV%) only

Comparative Bioavailability Studies

A single-dose, crossover, comparative bioavailability study of Mylan-Valproic (Valproic Acid) Capsules 250 mg and Depakene[®] (Valproic Acid) Capsules 250 mg (Abbott, Canada) following a single-dose of 250 mg valproic acid in 20 subjects under fasted conditions was conducted. The results indicate that Mylan-Valproic Capsules 250 mg are bioequivalent to Depakene[®] Capsules 250 mg. The summary of results is presented in the following table:

**Valproic Acid
(1 x 250 mg)
From measured data**

**Geometric Mean
Arithmetic Mean (CV %)**

Parameter	Test [‡]	Reference ^{⊕†}	% Ratio of Geometric Means	95% Confidence Interval
AUC _T μg/ml•h	566.33 579.10 (22.1)	566.88 579.39 (21.5)	99.90	96 - 104
AUC _I μg/ml•h	605.70 620.73 (23.3)	605.23 620.08 (22.9)	100.08	-
C _{max} μg/ml•h	31.72.0 32.28 (17.8)	31.73 32.26 (17.2)	99.96	95 - 105
T _{max} [§] (h)	1.40	1.63		
T _{1/2} [§] (h)	17.7	17.4		

[‡] Mylan-Valproic Capsules 250 mg were manufactured by Mylan Pharmaceuticals ULC

[†] Depakene[®] Capsules 250 mg was manufactured by Abbott Canada

[§] Expressed as the arithmetic mean (CV%) only

DETAILED PHARMACOLOGY

Animal

Valproic acid has been shown to be effective against several types of chemically and electrically induced convulsions in a variety of animal species. These included maximal electroshock, low frequency electroshock, CO₂ withdrawal, pentylene tetrazole, cobalt, bemegride, bicuculline and l-glutamate. Many forms of photic and auditory induced seizures are also effectively blocked by valproic acid.

In animal studies, valproic acid at doses of 175 mg/kg or less had no effect on locomotor activity and conditioned responses to positive reinforcement.

Doses greater than 175 mg/kg inhibited spontaneous and conditioned behaviour in mice and rats and interfered with coordination of hind limbs in rats. Suppression of spontaneous and evoked brain potentials was also demonstrated at these higher dose levels.

Valproic acid at doses of 175 mg/kg or less had little or no effect on the autonomic nervous system, cardiovascular system, respiration, body temperature, inflammatory responses, smooth muscle contraction or renal activity. Intravenous doses of 22, 43 and 86 mg/kg in animals caused very transient decreases followed by compensatory increases in blood pressure.

Sodium valproate injectable caused decreased activity, ataxia, dyspnea, prostration and death in rats and mice acutely exposed to dosages exceeding 200 mg/kg.

TOXICOLOGY

The initial animal testing was done with sodium valproate, whereas most of the recent research has been with valproic acid. The conversion factor is such that 100 mg of the sodium salt is equivalent to 87 mg of the acid. References to dosage are in terms of valproic acid activity.

Acute Toxicity

Acute toxicity has been determined in several animal species using oral, intravenous, intraperitoneal and subcutaneous routes. The oral median lethal dose in adult rats and dogs was about 1 to 2 g/kg. Toxicity was similar for both sexes; however, it tended to be greater in newborn and 14-day old rats and in young adult rats. The signs of toxicity were those of central nervous system depression. Specific organ damage was limited to cellular debris in reticuloendothelial tissue and slight fatty degeneration of the liver.

Large oral doses (more than 500 mg/kg) produced irritation of the gastrointestinal tract of rats.

In adult male mice, the oral medial lethal dose of divalproex sodium was 1.66 g/kg (equal to approximately 1.54 g/kg valproic acid).

Pulverized divalproex sodium enteric-coated tablets (equivalent to 250 mg valproic acid), suspended in 0.2% methylcellulose, were administered orally to mice and rats of both sexes (10/sex/species/group) in dosages ranging from 1.74 to 4.07 g/kg. The oral median lethal dose (LD₅₀) ranged from 2.06 to 2.71 g/kg. No consistent sex-related or species-related differences were observed.

Signs of central nervous system depression, such as decreased activity, ataxia, and sleep, were observed. At necropsy, discolouration and/or thickening of the glandular mucosa were observed in only 2 female rats treated with 2.71 g/kg that died acutely.

When mature rats and dogs were administered up to 240 mg/kg/day or 120 mg/kg/day, respectively, for at least four consecutive weeks, no significant toxicologic effects were reported. However, significant reductions in testicular weights and total white cell counts in rats given 240 mg/kg/day were considered as evidence of subtle toxicity from sodium valproate injectable. Therefore, 90 mg/kg/day in rats and 120 mg/kg/day in dogs were considered the highest non-toxic doses.

The acute intravenous toxicity of sodium valproate injectable formulation containing the equivalent of 100 mg valproic acid/mL was evaluated in both sexes of mice and rats. Groups of mice and rats (five/sex/species/group) were treated at dosages ranging from 0.5 to 9.0 mL/kg (50 to 900 mg valproate/kg). No overt signs of toxicity were present in rats and mice given 0.5 mL/kg (50 mg valproate/kg). LD₅₀ values for the test solution in mice and rats (data combined for both sexes) were 7.3 and 7.0 mL/mg (730 and 700 mg valproate/kg), respectively.

Subacute and Chronic Toxicity

Subacute and chronic toxicity studies consisted of 1, 3, 6 and 18 months studies in rats and 3, 6 and 12 months studies in dogs. Pathologic changes included suppression of the hematopoietic system, depletion of lymphocytes from lymphoid tissues and the loss of germinal epithelial cells from seminiferous tubules. Reduced spermatogenesis and testicular atrophy occurred in dogs at doses greater than 90 mg/kg/day and in rats at doses greater than 350 mg/kg/day. In rats, the first indication of toxicity at 350 mg/kg/day was decreased food consumption and growth.

Mutagenicity and Carcinogenicity

Mutagenicity

Valproate was not mutagenic in an *in vitro* bacterial assay (Ames test), did not produce dominant lethal effects in mice, and did not increase chromosome aberration frequency in an *in vivo* cytogenetic study in rats. Increased frequencies of sister chromatid exchange (SCE) have been reported in a study of epileptic children taking valproate, but this association was not observed in another study conducted in adults. There is some evidence that increased SCE frequencies may be associated with epilepsy. The biological significance of increase in SCE frequency is not known.

Carcinogenicity

Two hundred rats were given valproic acid in the diet for 107 weeks. Mean doses consumed in the treatment period were: 81 mg/kg/day (males) and 85 mg/kg/day (females), in the low dose group; 161 mg/kg/day (males) and 172 mg/kg/day (females) in the high dose group (approximately 10 to 50% of the maximum human daily dose on a mg/m² basis). Control animals received corn oil in the diet. The chief finding in the study was an increased incidence of skin fibrosarcomas in treated males of the high-dose group. There were 2 such neoplasms in the low dose group, 5 in the high dose group and none in control males. Fibrosarcomas in rats are relatively infrequent, usually occurring in less than 3% of animals.

Valproic acid was also administered in the diet to female mice for nearly 19 months at doses of 81 and 163 mg/kg/day and to male mice for nearly 23 months at doses of 80 and 159 mg/kg/day. A significant dose related trend occurred in male mice in the incidence of bronchoalveolar adenomas, and when the data were adjusted for the times of death, the incidence in the high dose group was significantly increased.

Depending on the method of statistical analysis, the incidence of hepatocellular carcinomas and/or adenomas also showed significant or almost significant increases for the corresponding observations. The results of these two studies indicate that valproic acid is a weak carcinogen or promoter in rats and mice. The significance of these findings for humans is unknown at present.

Reproduction and Teratology

Studies in rats have shown placental transfer of the drug. Doses greater than 65 mg/kg/day given to rats, mice and rabbits produced an increased incidence of skeletal abnormalities of the ribs, vertebrae and palate.

Doses greater than 150 mg/kg/day given to pregnant rabbits produced fetal resorptions and (primarily) soft-tissue abnormalities in the offspring.

In rats, there was a dose related delay in onset of parturition. Post-natal growth and survival of the progeny were adversely affected, particularly when drug administration spanned the entire gestation and early lactation period. Embryo lethality or major developmental abnormalities occurred in rats and rabbits at doses of 350 mg/kg/day.

Survival among pups born to the high dose females was very poor but was improved when pups were transferred to control dams shortly after birth.

Fertility

Chronic toxicity studies in juvenile and adult rats and dogs demonstrated reduced spermatogenesis and testicular atrophy at oral doses of valproic acid of 400 mg/kg/day or greater in rats (approximately equivalent to or greater than the maximum human daily dose on a mg/m² basis) and 150 mg/kg/day or greater in dogs (approximately 1.4 times the maximum human daily dose or greater on a mg/m² basis). Segment I fertility studies in rats have shown that oral doses up to 350 mg/kg/day (approximately equal to the maximum human daily dose on a mg/m² basis) for 60 days have no effect on fertility.

The effect of valproate on testicular development and on sperm production and fertility in humans is unknown.

REFERENCES

1. Godin Y, Heiner L, Mark J, Mandel P. Effects of di-n-propyl-acetate, An Anticonvulsive Compound on GABA Metabolism. *J. Neurochem.* 1969;16:869.
2. Simler S, Ciesielski L, Maitre M. Effects of Sodium n-dipropyl-acetate on Audiogenic Seizures and Brain Aminobutyric Acid Level. *Biochem. Pharmacol.* 1973;22:1701.
3. Gugler R, Schell A, Froscher W, Schulz HU. Oral Disposition of Valproic Acid Following Single and Multiple Doses. *Arch. Pharmacol.* 1977;297(Suppl. 2):R62.
4. Eadie MJ. Plasma Level Monitoring of Anticonvulsants. *Clin Pharmacokinet.* 1976;1:52.
5. Gram L, Wulff K, Kasmussen KE, Flachs H, Wuertz-Jorgensen A, Somerbeck KW, Lohren V. Valproate Sodium: A Controlled Clinical Trial Including Monitoring of Drug Levels. *Epilepsia.* 1977; 18:141.
6. Benoit JM, Besson JM, Bivot JB, Aleonard P. Effects of Na Dipropylacetate (DPA) on Dorsal Root Potentials Induced by Cortical Seizures. *J. Pharmacol.* 1971;2:23.
7. Frey HH, Loscher W. Di-n-propyl-acetic Acid. Profile of Anticonvulsant Activity in Mice. *Arzneim-Forsch.* 1976;26:299.
8. Kupferberg H, Lust WD, Penry JK. Anticonvulsant Activity of Dipropylacetic acid (DPA) in Relation to GABA and CGMP Brain Levels in Mice. *Fed. Proc.* 1975;34:283.
9. Misslin R, Ropartz PH, Mandel P. The Effects of n-dipropylacetate on the Acquisition of Conditioned Behaviour with Negative Reinforcement in Mice. *Psychopharmacologie.* 1975;44:263.
10. Pinder PM, Brogden RN, Speight TM, Avery GS. Sodium Valproate: A Review of its Pharmacological Properties and Therapeutic Efficacy in Epilepsy. *Drugs.* 1977;13:81.
11. Shearer DE, Fleming DE, Bigler ED, Wilson CE. Suppression of Photically Evoked After-discharge Bursting Following Administration of Anticonvulsants in Waking Rats, *Pharmacol. Biochem. Behav.* 1974;2:839.
12. Shuto K, Nishigaki T. The Pharmacological Studies on Sodium Dipropylacetate Anticonvulsant Activities and General Pharmacological Actions. *Oyo Yakuri (Pharmacometrics).* 1970;4:937.
13. Simon D, Perry JK. Sodium di-n-propylacetate (DPA) in the Treatment of Epilepsy. A Review. *Epilepsia.* 1975;16:549.

14. Robert E, Guibaud P. Maternal Valproic Acid and Congenital Neural Tube Defects. *The Lancet* 2(8304):October 23, 1982;937.
15. Shakir RA. Comparison of Sodium Valproate and Phenytoin as Single Drug Treatment in Epilepsy. *Epilepsia*. 1981 ;22 :27-33.
16. Shakir RA. Comparative Trial of Valproate Sodium and Clonazepam in Chronic Epilepsy. *Arch Neurol*. 1979;36:301 –304.
17. Turnbull DM, Rawlins MD, Weightman D, Chadwick DW. A Comparison of Phenytoin and Valproate in Previously Untreated Adult Epileptic Patients. *J. Neurol Neurosurg. Psychiatr.* 1982;45:55-59.
18. Gugler R, vonUnruh GE. Clinical Pharmacokinetics of Valproic Acid. *Clinical Pharmacokinetics*. 1980;5:67-83.
19. Cramer JA, Mattson RH. Valproic Acid: In Vitro Plasma Protein Binding and Interaction with Phenytoin. *Ther. Drug. Monit.* 1979;1(1):105-116.
20. Bowdle TA, Patel JH, Levy RH. Valproic Acid Dosage and Plasma Protein Binding and Clearance. *Clin. Pharmacol. Ther.* 1980;28(4) :487-492.
21. Gram L, Flachs H. Sodium Valproate, Serum Level and Clinical Effect in Epilepsy: A Controlled Study. *Epilepsia*. 1979;20: 303-312.
22. Kapetamovic IM, Kupferberg HJ. Inhibition of Microsomal Phenobarbital Metabolism of Valproic Acid. *Biochem. Pharmacol.* 1981 ;30(1 1): 1361-1363.
23. Patel IH, Levy RH. Phenobarbital - Valproic Acid Interaction. *Clin. Pharmacol. Ther.* 1979;27(4):515-521.
24. Bruni J, Wilder BJ. Valproic Acid and Plasma Levels of Phenobarbital. *Neurol.* 1980;30:94-97.
25. Pinder PM, Brogden RN. Sodium Valproate: A Review of Its Pharmacological Properties and Therapeutic Efficacy in Epilepsy. *Drugs*. 1977;B:81-123.
26. Millet Y, Sinty JM. Problems Created by Combining Phenobarbital and Sodium Dipropylacetate Therapeutically: Case Report. *Europ. Toxicol.* 1976;9(6)6: 381-383.
27. Rai PV. Acute Intoxication During a Combined Treatment of Sodium Valproate and Phenobarbitone. In: Meinardi H and Rowan AJ, editors. *Advances in epileptology*, Swets Zertlinger, Amsterdam, pp. 366-369, 1978.

28. Valproic acid (Systemic). USPDI 1990; 10th Ed(1 B): 2720-25. Data on file at Mylan Pharmaceuticals.
29. Product Monograph DEPAKENE[®] (valproic acid capsules USP). BGP Pharma ULC, 8401 Trans-Canada Highway, Saint-Laurent (QC) Canada. January 13, 2015, Control No. 178659.

PART III: CONSUMER INFORMATION

Pr MYLAN-VALPROIC (Valproic Acid Capsules, USP)

This leaflet is part III of a three-part "Product Monograph" published when Mylan-Valproic was approved for sale in Canada and is designed specifically for consumers. This leaflet is a summary and will not tell you everything about Mylan-Valproic. Contact your doctor or pharmacist if you have any questions about the drug.

ABOUT THIS MEDICATION

What the medication is used for:

- Mylan-Valproic has been prescribed to you to control your epilepsy. Please follow your doctor's recommendations carefully.

What it does:

Mylan-Valproic has anticonvulsant properties. The mechanism of action has not yet been established. It has been suggested that its activity in epilepsy is related to increased brain concentrations of gamma-aminobutyric acid (GABA).

When it should not be used:

Mylan-Valproic should not be taken by:

- patients with liver disease or significant liver dysfunction
- patients with mitochondrial diseases (e.g., Alpers or Alpers-Huttenlocher disease)
- patients who are allergic to the drug
- patients with known urea cycle disorders (a genetic disorder)
- patients with known porphyria (a genetic disorder)

What the medicinal ingredient is:

Valproic acid.

What the non-medicinal ingredients are:

Mylan-Valproic 250 mg capsules contain the following non-medicinal ingredients: Corn Oil, D&C Yellow' No. 10, FD&C Red No. 40, Gelatin, Glycerine, Purified Water and Titanium Dioxide.

What dosage forms it comes in:

Mylan-Valproic is available as a capsule containing 250 mg of valproic acid.

WARNINGS AND PRECAUTIONS

Serious Warnings and Precautions

- **Hepatotoxicity:** liver failure resulting in death has occurred in patients receiving valproic acid. These incidents usually occurred during the first 6 months of treatment with valproic acid. Patients taking several anticonvulsant drugs, children, those with a history of liver disease, metabolic disorders, severe seizure disorders accompanied by mental retardation, and those with brain disease may be at particular risk. Experience has indicated that children under the age of 2 years are at a considerably increased risk of developing fatal hepatotoxicity, especially those on multiple anticonvulsants.
- **Teratogenicity:** Valproic acid can produce birth defects to an unborn baby. Accordingly, the use of Mylan-Valproic in women of childbearing potential requires that the benefits of its use be weighed against the risk of injury to the fetus.
- **Pancreatitis:** cases of life threatening pancreas disorder have been reported in both children and adults receiving valproic acid. Some cases have occurred shortly after first use as well as after several years of use. Abdominal pain, nausea, vomiting and/or anorexia can be symptoms of pancreatitis that require immediate medical evaluation.

BEFORE you use Mylan-Valproic talk to your doctor or pharmacist if:

- you have a history of, or suffer from a liver disease, such as jaundice (yellowing of the skin and eyes);
- you have ever had an unusual or allergic reaction to valproic acid (including fever or rash);
- you are allergic to any component of Mylan-Valproic capsules;
- you are pregnant or are planning to become pregnant;
- you are breast-feeding (nursing); valproic acid passes into breast milk;
- you are taking any other prescription or over the counter medicine;
- you have kidney disease;
- you have other medical conditions including a history of unexplained coma, intellectual disability or any type of brain dysfunction;
- you have a psychiatric disorder or have thoughts of suicide;
- you consume alcohol on a regular basis.

Precautions while taking Mylan-Valproic:

- Your doctor will monitor your response to Mylan-Valproic on a regular basis. However, if your seizures get worse, you should tell your doctor immediately.

- Since Mylan-Valproic may cause poor coordination and/or drowsiness, you should not engage in hazardous activities, such as driving and operating machinery, until you know that you don't become drowsy from the drug.
- You should not stop taking your medication unless directed by your doctor. You should always check that you have an adequate supply of Mylan-Valproic. You should remember that this medicine was prescribed only for you; it should never be given to anyone else.

Women of Childbearing Potential

- All women of childbearing age who are being treated with Mylan-Valproic should talk to their healthcare providers about using other possible treatments instead of Mylan-Valproic. If the decision is made to use Mylan-Valproic, you must use an effective method of birth control (contraception). You should talk to your doctor about the best kind of birth control to use while you are taking Mylan-Valproic.

Pregnant Women

- If you take Mylan-Valproic during pregnancy, your child has a higher risk for birth defects and adverse effects on IQ and brain development, which can begin early in the pregnancy, even before you know that you are pregnant.
- There may be other medications to treat your condition that have a lower chance of birth defects.
- If you are planning to become pregnant, or if you become pregnant while taking Mylan-Valproic, you should promptly inform your doctor. Do not suddenly stop taking the drug. Appropriate treatment options will need to be discussed with your physician to ensure the benefits outweigh the risks.
- Pregnancy Registry: If you become pregnant while taking Mylan-Valproic, talk to your doctor about registering with the North American Antiepileptic Drug Pregnancy Registry. You can enrol in this registry by calling 1-888-233-2334. The purpose of this registry is to collect information about the safety of antiepileptic medicines during pregnancy. Information on the registry can also be found at the following website:
<http://www.aedpregnancyregistry.org/>.

INTERACTIONS WITH THIS MEDICATION

Serious Drug Interactions

- Rare cases of coma have been reported in patients receiving valproic acid alone or when taken with phenobarbital.
- Serious skin reactions (such as conditions called Stevens-Johnson syndrome and Toxic Epidermal Necrolysis) have been reported when valproic acid and lamotrigine were taken together.

Drugs that may interact with Mylan-Valproic include:

- anticonvulsants such as carbamazepine, lamotrigine, primidone, topiramate, felbamate, phenytoin, ethosuximide, phenobarbital;
- anticoagulants such as acetylsalicylic acid, warfarin, dicumarol;
- benzodiazepines such as diazepam, lorazepam, clonazepam;
- some medicines used to treat infections such as rifampin;
- some medicines used to treat diabetes such as tolbutamide;
- some HIV-antiviral medication such as zidovudine;
- any of the group of antibiotics in the carbapenem class such as doripenem, ertapenem, imipenem, meropenem;
- some medicines used to treat heartburn and peptic ulcers such as cimetidine;
- medicines used to treat depression such as Selective Serotonin Re-Uptake Inhibitors (SSRIs), Monoamine Oxidase Inhibitors (MAOIs), Tricyclic antidepressants such as amitriptyline, nortriptyline;
- antipsychotics.

PROPER USE OF THIS MEDICATION

Please consult your doctor before taking any other medication, including over-the-counter medicines. Some drugs can produce various side effects when they are used in combination with Mylan-Valproic.

It is important to keep your appointments for medical checkups.

The doctor may need to take blood tests to measure the amount of Mylan-Valproic in your blood when adjusting your medications.

IMPORTANT: PLEASE READ

Usual dose:

It is very important to take Mylan-Valproic exactly as instructed by your doctor.

The recommended starting dose of Mylan-Valproic will be decided by your doctor based on your weight, your seizures or manic episodes and your concomitant medications. Be sure to tell your doctor all the prescription and over the counter medications that you are currently taking. Your doctor will gradually increase the dosage until your condition is well controlled without experiencing side effects. You should carefully follow the instructions that were given to you and not change your dose without consulting with your doctor.

Mylan-Valproic may be taken with or without food.

Overdose:

In case of drug overdose, contact a health care practitioner, hospital emergency department or regional Poison Control Centre immediately, even if there are no symptoms.

Missed Dose:

Do not suddenly stop taking your medicine because of the risk of increasing your epileptic seizures.

If you miss a dose, you should not try to make up for it by doubling up on your next dose. You should take your next regularly scheduled dose and try not to miss any more doses.

SIDE EFFECTS AND WHAT TO DO ABOUT THEM

You should check with your doctor or pharmacist right away if you notice any bothersome or unusual effects while taking Mylan-Valproic.

The most commonly reported adverse reactions include nausea, vomiting, indigestion sleepiness, headache, diarrhea, weakness, tremor and dizziness. Changes in hair are also reported, such as hair loss or in increase in hair on face, chest and back. If any of these affect you severely, contact your doctor or pharmacist.

You should know that this does not mean that you will experience such effects, because people can react in different ways to the same medicine.

SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM				
Symptom / effect		Talk with your doctor or pharmacist right away		Stop taking drug and seek immediate emergency medical attention
		Only if severe	In all cases	
Common	Hallucinations: seeing or hearing something that is not there	√		
Uncommon	Brain dysfunction from high ammonia levels in the blood (tiredness, vomiting, abnormal walking, extreme irritability†, combative/bizarre behaviour††, refusal to eat meat or high protein products††)		√	
	Decreased number of platelets in the blood (may result in easy bruising and bleeding from the skin or other areas)		√	
	Liver disorder (weakness, tiredness, abdominal pain, diarrhea, facial swelling, loss of appetite, yellowing of the skin or eyes, dark urine, nausea, and vomiting)		√	
	Pancreas Disorder (abdominal pain, nausea, vomiting and /or loss of appetite)		√	
	Thoughts of suicide or hurting yourself, symptoms of depression or unusual changes in mood or		√	

SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM

Symptom / effect	Talk with your doctor or pharmacist right away		Stop taking drug and seek immediate emergency medical attention
	Only if severe	In all cases	
behaviour			
Rare Muscle disorder (unexplained muscle pain or tenderness, with a fever or “tea-coloured” urine, or reduced urination)		√	
† In young children †† In older children or adults			

This is not a complete list of side effects. For any unexpected effects while taking Mylan-Valproic, contact your doctor or pharmacist.

HOW TO STORE IT

Mylan-Valproic should be stored between 15°C and 30°C.

Mylan-Valproic should be kept out of reach of children.

Reporting Side Effects

You can help improve the safe use of health products for Canadians by reporting serious and unexpected side effects to Health Canada. Your report may help to identify new side effects and change the product safety information.

3 ways to report:

- Online at MedEffect (<http://hc-sc.gc.ca/dhp/mps/medeff/index-eng.php>);
- By calling 1-866-234-2345 (toll-free);
- By completing a Consumer Side Effect Reporting Form and sending it by:
 - Fax to 1-866-678-6789 (toll-free), or
 - Mail to: Canada Vigilance Program
Health Canada, Postal Locator
0701E
Ottawa, ON
K1A 0K9

Postage paid labels and the Consumer Side Effect Reporting Form are available at MedEffect (<http://hc-sc.gc.ca/dhp/mps/medeff/index-eng.php>).

NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

MORE INFORMATION

This document plus the full Product Monograph, prepared for health professionals can be requested by contacting the sponsor, Mylan Pharmaceuticals ULC at: 1-800-575-1379

This leaflet was prepared by Mylan Pharmaceuticals ULC.

Last revised: May 7, 2015



Mylan Pharmaceuticals ULC
85 Advance Road
Etobicoke, Ontario
M8Z 2S6