PRESEPT Tablets 5.0g Outer label (50 Tablets container)

READ ENCLOSED DIRECTIONS BEFORE USE

DESCRIPTION

PRESEPT® Tablets are effervescent disinfectant tablets containing the active ingredient **Sodium Dichloroisocyanurate**. They dissolve rapidly in water to produce a rapid acting, wide spectrum disinfectant solution effective against vegetative bacteria, fungi, viruses and bacterial spores.

USES

Solutions prepared from PRESEPT® Tablets are suitable for the disinfection of hard surfaces. In addition, efficacy has been demonstrated in the following organisms:

- Clostridium difficile
- Methicillin Resistant Staphylococcus aureus (MRSA)
- Canine Parvovirus
- Feline Panleukopenia virus

PREPARATION

To prepare a disinfectant solution of 5000ppm, two 5.0g PRESEPT® Tablet to 1L of water.

For more specific dilution instructions refer to enclosed sheet.

DANGER

Contact with acids liberates toxic gas. Causes serious eye irritation. May cause respiratory irritation. Very toxic to aquatic life with long lasting effects. Avoid breathing dust/fume/gas/mist/vapors/spray. Avoid release to the environment. Wear protective gloves/protective clothing/eye protection/face protection. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do. Continue rinsing. Call a POISON CENTER or doctor if

5.0 g

PRESEPT® DISINFECTANT TABLETS

50 TABLETS each containing
5.0g SODIUM DICHLOROISOCYANURATE

French equivalent

DIN 01

Johnson & Johnson Medical Products Markham, Ontario L3R 0T5 © Ethicon, Inc. 2001- 2015 949-581-5799



DIN 01946404

MPR031

5.0g/50

French

left panel

equivalent of



Danger

you feel unwell. If eye irritation persists: Get medical attention.		
CAUTION		
Harmful if swallowed. Avoid skin and eye contact. If eye contact		
occurs, irrigate with copious amounts of water. Do not use broken or		
chipped tablets. Store in a cool dry place. Do not mix with acids or		
cationic detergents. Do not breathe dust. Do not dissolve tablets in a		
closed container.		

PRESEPT Tablets 5.0g Outer label (50 Tablets container)

READ ENCLOSED DIRECTIONS BEFORE USE

DESCRIPTION

PRESEPT® Tablets are effervescent disinfectant tablets containing the active ingredient **Sodium Dichloroisocyanurate**. They dissolve rapidly in water to produce a rapid acting, wide spectrum disinfectant solution effective against vegetative bacteria, fungi, viruses and bacterial spores.

USES

Solutions prepared from PRESEPT® Tablets are suitable for the disinfection of hard surfaces. In addition, efficacy has been demonstrated in the following organisms:

- Clostridium difficile
- Methicillin Resistant Staphylococcus aureus (MRSA)
- Canine Parvovirus
- Feline Panleukopenia virus

PREPARATION

To prepare a disinfectant solution of 5000ppm, two 5.0g PRESEPT® Tablet to 1L of water.

For more specific dilution instructions refer to enclosed sheet.

DANGER

Contact with acids liberates toxic gas. Causes serious eye irritation. May cause respiratory irritation. Very toxic to aquatic life with long lasting effects. Avoid breathing dust/fume/gas/mist/vapors/spray. Avoid release to the environment. Wear protective gloves/protective clothing/eye protection/face protection. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do. Continue rinsing. Call a POISON CENTER or doctor if

5.0 g

PRESEPT® DISINFECTANT TABLETS

50 TABLETS each containing 5.0g SODIUM DICHLOROISOCYANURATE

French equivalent

DIN 01946404

MPR031

5.0g/50

French

left panel

equivalent of

Danger

Johnson & Johnson Medical Products Markham, Ontario L3R 0T5 © Ethicon, Inc. 2001- 2015 949-581-5799

Draft Outerlabel - 5.0g/tab (non-annotated)

you feel unwell. If eye irritation persists: Get medical attention.		
CAUTION		
Harmful if swallowed. Avoid skin and eye contact. If eye contact		
occurs, irrigate with copious amounts of water. Do not use broken or		
chipped tablets. Store in a cool dry place. Do not mix with acids or		
cationic detergents. Do not breathe dust. Do not dissolve tablets in a		
closed container.		