

**PRODUCT MONOGRAPH**

**<sup>PR</sup>MYLAN-TOLTERODINE ER**

**(tolterodine L-tartrate extended release capsules)**

**2 mg and 4 mg Capsules**

**Anticholinergic - Antispasmodic Agent**

**Mylan Pharmaceuticals ULC  
85 Advance Road,  
Etobicoke ON,  
M8Z 2S6**

**Date of Preparation:  
April 28, 2015**

**Submission Control No: 147332, 183925**

## TABLE OF CONTENTS

<b>PART I: HEALTH PROFESSIONAL INFORMATION.....</b>	<b>3</b>
SUMMARY PRODUCT INFORMATION .....	3
INDICATIONS AND CLINICAL USE.....	3
CONTRAINDICATIONS .....	4
WARNINGS AND PRECAUTIONS.....	4
ADVERSE REACTIONS .....	7
DRUG INTERACTIONS.....	9
DOSAGE AND ADMINISTRATION .....	11
OVERDOSAGE .....	12
ACTION AND CLINICAL PHARMACOLOGY .....	13
STORAGE AND STABILITY.....	15
SPECIAL HANDLING INSTRUCTIONS .....	15
DOSAGE FORMS, COMPOSITION AND PACKAGING.....	15
<b>PART II: SCIENTIFIC INFORMATION.....</b>	<b>16</b>
PHARMACEUTICAL INFORMATION .....	16
CLINICAL TRIALS.....	17
COMPARATIVE BIOAVAILABILITY .....	17
DETAILED PHARMACOLOGY.....	20
TOXICOLOGY .....	26
REFERENCES .....	29
<b>PART III: CONSUMER INFORMATION .....</b>	<b>33</b>

# PR MYLAN-TOLTERODINE ER

(tolterodine L-tartrate extended release capsules)

## PART I: HEALTH PROFESSIONAL INFORMATION

### SUMMARY PRODUCT INFORMATION

Route of Administration	Dosage Form / Strength	All Nonmedicinal Ingredients
Oral	Extended release capsules 2 mg and 4 mg	Ethylcellulose, FD&C Blue #2, gelatin, hypromellose, iron oxide yellow (2 mg capsules only), starch, sucrose, talc, titanium dioxide; Ink: povidone, propylene glycol, shellac and sodium hydroxide.

### INDICATIONS AND CLINICAL USE

MYLAN-TOLTERODINE ER (tolterodine L-tartrate extended release capsules) is indicated for:

- the symptomatic management of patients with an overactive bladder with symptoms of urinary frequency, urgency, or urge incontinence, or any combination of these symptoms. (see **WARNINGS AND PRECAUTIONS and DETAILED PHARMACOLOGY, Electrophysiology**).

**Geriatrics (≥ 65 years of age):** No overall differences were observed in safety between older (patients ≥ 65 years) and younger patients (patients < 65 years) on tolterodine extended release capsules; and therefore, no dosage adjustment for elderly patients is recommended (see **WARNINGS AND PRECAUTIONS, Special Populations, DETAILED PHARMACOLOGY and CLINICAL TRIALS**).

## CONTRAINDICATIONS

**MYLAN-TOLTERODINE ER** (tolterodine L-tartrate extended release capsules) is contraindicated in patients with:

- urinary retention,
- gastric retention,
- uncontrolled narrow angle glaucoma,
- a known hypersensitivity to this drug or to any ingredient in the formulation or component of the container (see **PHARMACEUTICAL INFORMATION**).

## WARNINGS AND PRECAUTIONS

### **Gastrointestinal and Genitourinary**

#### **Patients at Risk of Urinary Retention and Gastric Retention**

**MYLAN-TOLTERODINE ER** (tolterodine L-tartrate extended release capsules) should be administered with caution to patients with clinically significant bladder outflow obstruction because of the risk of urinary retention, to patients at risk of decreased gastrointestinal motility, and to patients with gastrointestinal obstructive disorders, such as pyloric stenosis, because of the risk of gastric retention (see **CONTRAINDICATIONS**).

### **Cardiovascular**

#### **Patients with Congenital or Acquired QT Prolongation:**

In a clinical QT study, the QT prolonging effect of two times the highest labeled dose of tolterodine (8 mg/per day in divided doses, given as tolterodine immediate release tablets) was 50% to 60% less than that of the active control moxifloxacin (400 mg) at its labeled dose. At the recommended therapeutic dose (4 mg daily) of tolterodine L-tartrate tablets, the effect was lower. Since the QT prolongation effect is in linear relationship with exposure, any QT effect of tolterodine L-tartrate extended release capsules would also be expected to be similarly lower. This study, however, was not designed to make direct statistical comparisons between drugs, tolterodine formulations, or dose levels.

The clinical relevance of these findings will depend on individual patient risk factors and susceptibilities present. Particular care should be exercised in patients who are at an increased risk of experiencing torsade de pointes during treatment with QT/QTc-prolonging drugs. This especially holds true in patients with abnormally long baseline QT/QTc intervals or when taking potent CYP3A4 inhibitors (see **DRUG INTERACTIONS, Drug-Drug Interactions, DOSAGE AND ADMINISTRATION, DETAILED PHARMACOLOGY, Electrophysiology**).

In the general population, the risk factors for torsade de pointes include, but are not limited to, the following:

- female;
- elderly (65 years);
- genetic variants affecting cardiac ion channels or regulatory proteins, especially congenital long QT syndrome;

- family history of sudden cardiac death at <50 years;
- cardiac disease (e.g., myocardial ischemia or infarction, congestive heart failure, left ventricular hypertrophy, cardiomyopathy);
- demonstrated history of arrhythmias (especially ventricular arrhythmias, atrial fibrillation, or recent conversion from atrial fibrillation);
- bradycardia (<50 beats per minute);
- acute neurological events (e.g., intracranial or subarachnoid haemorrhage, stroke, intracranial trauma);
- electrolyte disturbances (e.g., hypokalemia, hypomagnesemia, hypocalcemia);
- nutritional deficits (e.g., eating disorders, extreme diets);
- diabetes mellitus;
- autonomic neuropathy;
- hepatic or renal dysfunction if relevant to the elimination of the drug.

Approximately 7% of Caucasians are poor metabolizers of CYP2D6 substrates. A pharmacokinetic/pharmacodynamic model estimated that QTc interval increases in poor metabolizers treated with tolterodine 2 mg BID are comparable to those observed in extensive metabolizers receiving 4 mg BID.

Discontinuation of the drug should be considered if symptoms suggestive of arrhythmia occur.

### **Aggravation with Pre-existing Cardiac Conditions**

Although there are no clinical trial or post-marketing data to confirm the potential for tolterodine L-tartrate extended release capsules to aggravate certain pre-existing cardiac conditions, this product is in the class anticholinergic medications which are known to have cardiac effects. Prescribers should therefore use caution when prescribing MYLAN-TOLTERODINE ER to patients with ischemic heart disease, congestive heart failure, cardiac arrhythmias, or tachycardia.

### **Neurologic**

MYLAN-TOLTERODINE ER should be used with caution in patients with myasthenia gravis.

### **Ophthalmologic**

#### **Controlled Narrow Angle Glaucoma**

MYLAN-TOLTERODINE ER should be used with caution in patients being treated for narrow angle glaucoma.

### **Hepatic/Biliary/Pancreatic/Renal**

Patients with impaired hepatic function and patients with renal impairment should not receive doses of MYLAN-TOLTERODINE ER greater than 2 mg daily (see **DETAILED PHARMACOLOGY, Pharmacokinetics in Special Populations**).

### **Special Populations**

**Pregnant Women:** Studies in mice have shown that at doses of 30 to 40 mg/kg/day, tolterodine caused embryoletality, reduced fetal weight, and increased incidence of fetal abnormalities (cleft palate, digital abnormalities, intra-abdominal hemorrhage, various skeletal abnormalities, primarily

reduced ossification in mice). At these doses, AUC values were about 20 to 25-fold higher than in humans. At doses of 20 mg/kg/day (AUC value was about 15-fold higher than in humans), no anomalies or malformations were seen in mice. There are no studies of tolterodine in pregnant women. Therefore, **MYLAN-TOLTERODINE ER** should be used during pregnancy only if the potential benefit for the mother justifies the potential risk for the fetus. Women of childbearing potential should be considered for treatment only if using adequate contraception (see **TOXICOLOGY**).

**Nursing Women:** Tolterodine is excreted into the milk in mice. It is not known whether tolterodine is excreted in human milk. Because many drugs are excreted into human milk, administration of **MYLAN-TOLTERODINE ER** should be avoided during nursing.

**Pediatrics:** The safety and effectiveness of tolterodine extended release capsules in pediatric patients have not been established.

**Geriatrics (65 – 93 years of age):** Of the 1120 patients who were treated in the four, phase III, 12- week clinical studies of tolterodine, 474 (42%) were 65 to 91 years of age. No overall differences in safety were observed between the older and younger patients.

Of the 1526 patients who were treated in the 12-week clinical study comparing tolterodine extended release capsules and tolterodine immediate release tablets versus placebo, 642 (42%) were 65 to 93 years of age. No overall differences in safety were observed between the older and younger patients (see **DETAILED PHARMACOLOGY, Pharmacokinetics in Special Populations**).

### **Monitoring and Laboratory Tests**

Monitoring of the QT/QTc interval and/or serum electrolyte levels may be appropriate in high risk patients who are being treated with **MYLAN-TOLTERODINE ER**, such as:

- patients with known congenital or acquired QT/QTc prolongation or electrolyte disturbances;
- patients with impaired hepatic or renal function or other comorbid conditions that may increase tolterodine exposure or cause QT/QTc prolongation;
- patients who are taking drugs that have been associated with QT/QTc prolongation and/or torsade de pointes such as Class IA (e.g., quinidine, procainamide) or Class III (e.g., amiodarone, sotalol) antiarrhythmic medications, or those taking potent CYP3A4 inhibitors.

(see **WARNINGS AND PRECAUTIONS, Cardiovascular, DRUG INTERACTIONS, Drug- Drug Interactions, DOSAGE AND ADMINISTRATION, DETAILED PHARMACOLOGY, Electrophysiology**).

Discontinuation of the drug should be considered if symptoms suggestive of arrhythmia occur or if the QT/QTc interval becomes markedly prolonged.

## Information For Patients

The ability to drive and use machinery may be negatively affected. Patients should be advised to exercise caution.

## ADVERSE REACTIONS

### Adverse Drug Reaction Overview

In a large randomized, multicenter, double-blind, 12-week study, patients treated with tolterodine L-tartrate extended release capsules, 4 mg once daily (N=505), tolterodine immediate release tablets, 2 mg twice daily (N=512), or placebo (N=507), were evaluated for safety.

Tolterodine L-tartrate extended release capsules, 4 mg once daily, was generally well tolerated, with an overall incidence of adverse events comparable to tolterodine immediate release tablets, 2 mg twice daily, and placebo. Dry mouth was the most frequently reported adverse event for patients treated with tolterodine L-tartrate extended release capsules occurring in 23.4% of patients treated with tolterodine L-tartrate extended release capsules, 30.5% in patients treated with tolterodine immediate release tablets and 7.7% of placebo-treated patients. The overall dry mouth rate for patients taking tolterodine L-tartrate extended release capsules, in this single pivotal trial, was 23% lower than for tolterodine immediate release tablets ( $P<0.02$ ).

Adverse events considered to be related to treatment with tolterodine L-tartrate extended release capsules, tolterodine immediate release tablets, versus placebo			
	Tolterodine extended release capsules	Tolterodine immediate release tablets	Placebo
Dry Mouth	23.4%	30.5%	7.7%
Abdominal Pain	3.8%	2.5%	1.6%
Dyspepsia	3.0%	3.1%	1.4%
Dizziness/Vertigo	2.2%	1.8%	1.0%
Fatigue	2.2%	1.2%	0.8%
Sinusitis	1.8%	0.6%	0.6%
Abnormal Vision	1.2%	0.8%	0.4%
Dysuria	1.0%	1.6%	0.2%

Dry mouth, constipation, abnormal vision (accommodation abnormalities), urinary retention, and dry eyes are expected side effects of antimuscarinic agents.

The frequency of discontinuation due to adverse events was highest during the first 4 weeks of treatment. Similar percentages of patients treated with tolterodine extended release capsules, tolterodine immediate release tablets or placebo, discontinued treatment due to adverse events; the most common adverse events associated with discontinuation were dry mouth (1.6%), headache (1.0%), and constipation (0.7%).

### **Clinical Trial Adverse Drug Reactions**

*Because clinical trials are conducted under very specific conditions the adverse reaction rates observed in the clinical trials may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse drug reaction information from clinical trials is useful for identifying drug-related adverse events and for approximating rates.*

The following table lists the adverse events reported in  $\geq 5\%$  or more of patients treated with tolterodine L-tartrate extended release capsules, 4 mg once daily, in the 12-week study. The adverse events were reported regardless of causality.

<b>Incidence (%) of Adverse Events that Occurred in <math>\geq 5\%</math> of Patients Treated with tolterodine L-tartrate extended release capsules and tolterodine immediate release tablets in a 12-week Controlled Clinical Trial</b>			
	<b>tolterodine extended release capsules 4 mg Once Daily N=505</b>	<b>Placebo N=507</b>	<b>Tolterodine immediate release tablets 2 mg twice daily N=512</b>
% Patients Reporting Serious Events	1.4	3.6	2.3
% Patients Discontinuing due to Adverse Events	5.3	6.5	5.4
Dry mouth	23.4	7.7	30.5
Headache	6.3	4.5	3.7
Constipation	5.9	4.3	6.6

### **Less Common Clinical Trial Adverse Drug Reactions (1% to $< 5\%$ )**

Other events reported by 1% to  $< 5\%$  of patients treated with tolterodine L-tartrate extended release capsules and numerically greater than those reported for patients receiving placebo are listed in order of descending frequency: abdominal pain, dry eyes, urinary tract infection, dyspepsia, upper respiratory tract infection, somnolence, dizziness, fatigue, flatulence, sinusitis, edema, pain, abnormal vision, and dysuria.

Over 400 patients treated for up to 6 months with tolterodine L-tartrate extended release capsules 4 mg once daily, had an overall incidence and adverse event profile similar to those patients treated with tolterodine L-tartrate extended release capsules for 12 weeks.



## **Post-Market Adverse Drug Reactions**

The following events have been reported in association with tolterodine use in clinical practice: anaphylactoid reactions (including angioedema), tachycardia, palpitations, peripheral edema, hallucinations, disorientation, memory impairment, and diarrhea.

**Cholinesterase Inhibitors:** Worsening of symptoms of dementia (e.g. confusion, disorientation, delusion) have been reported after tolterodine therapy was initiated in patients taking cholinesterase inhibitors for the treatment of dementia.

## **DRUG INTERACTIONS**

### **Overview**

Concomitant medication with other drugs that possess antimuscarinic properties may result in more pronounced therapeutic and/or adverse effects. Conversely, the therapeutic effect of tolterodine may be reduced by concomitant administration of muscarinic receptor agonists.

### **Drug-Drug Interactions**

#### **Effects of Other Drugs on MYLAN-TOLTERODINE ER**

Drugs Which Prolong the QT/QTc Interval: Drugs that have been associated with QT/QTc interval prolongation and/or torsade de pointes include, but are not limited to, the examples in the following list. Chemical/pharmacological classes are listed if some, although not necessarily all, class members have been implicated in QT/QTc prolongation and/or torsade de pointes:

- Antiarrhythmics (Class IA, e.g., quinidine, procainamide, disopyramide; Class III, e.g., amiodarone, sotalol, ibutilide; Class IC, e.g., flecainide, propafenone);
- Antipsychotics (e.g., thioridazine, chlorpromazine, pimozide, haloperidol, droperidol);
- Antidepressants (e.g., amitriptyline, imipramine, maprotiline, fluoxetine, venlafaxine);
- Opioids (e.g., methadone);
- Antibacterials (e.g., erythromycin, clarithromycin, telithromycin, moxifloxacin, gatifloxacin);
- Antimalarials (e.g., quinine);
- Pentamidine;
- Azole antifungals (e.g., ketoconazole, fluconazole, voriconazole);
- Gastrointestinal drugs (e.g., domperidone, dolasetron, ondasetron);
- B2-adrenoreceptor agonist (salmeterol, formoterol);
- Tacrolimus

This list of potentially interacting drugs is not comprehensive. Prior to initiating drug treatment in the presence of concomitant medications, physicians should consult current scientific literature for

information on the ability of newly approved drugs to prolong the QT/QTc interval, inhibit the metabolizing enzyme or transporter, or cause electrolyte disturbances, as well for older drugs for which these effects have recently been established (see **WARNINGS AND PRECAUTIONS**).

Cytochrome P450 3A4 inhibitors: Patients treated with ketoconazole or other potent CYP3A4 inhibitors such as other azole antifungals (e.g., itraconazole, miconazole) or macrolide antibiotics (e.g., erythromycin, clarithromycin) or cyclosporine or vinblastine, should not receive doses of **MYLAN-TOLTERODINE ER** (tolterodine L-tartrate extended release capsules) greater than 2 mg daily (see **DETAILED PHARMACOLOGY, Drug Interactions**).

Fluoxetine: Fluoxetine, a potent inhibitor of P450 2D6, inhibits significantly the metabolism of tolterodine in extensive metabolizers. The sum of unbound serum concentrations of tolterodine and the 5-hydroxymethyl derivative (DD 01) is 25% higher when the two drugs are administered concomitantly. No dose adjustment is required (see **DETAILED PHARMACOLOGY, Drug Interactions**).

### **Effects of MYLAN-TOLTERODINE ER on Other Drugs**

Other Drugs Metabolized by P450 2D6: The potential effect of tolterodine on the pharmacokinetics of drugs that are metabolized by P450 2D6 (such as flecainide, vinblastine, carbamazepine, tricyclic antidepressants) has not been formally evaluated (see **DETAILED PHARMACOLOGY, Drug Interactions**).

Diuretics: Coadministration of diuretics (such as indapamide, hydrochlorothiazide, triamterene, bendroflumethiazide, chlorothiazide, methylchlorothiazide, or furosemide) with tolterodine (2 mg, twice daily) did not cause any adverse ECG effects, however, in the presence of diuretics causing hypokalemia, and, concomitant medications known or suspected to cause adverse ECG effects (such as QT/QTc prolongation), the physician is advised to exercise caution and advise the patient about the signs and symptoms of cardiac arrhythmia (see **DETAILED PHARMACOLOGY, Drug Interactions**).

Oral Contraceptives: Clinical drug interaction studies have shown that there are no known interactions between tolterodine immediate release tablets and oral contraceptives (ethinyl estradiol/levonorgestrel).

Warfarin: Clinical drug interaction studies have shown that there are no known interactions between tolterodine immediate release tablets and warfarin.

### **Concomitant use of MYLAN-TOLTERODINE ER with alpha-blockers in men**

Tolterodine L-tartrate extended release capsules efficacy has not been established in studies of men on an alpha-blocker therapy. The two trials conducted in men with overactive bladder symptoms (OAB) with and without benign prostatic hyperplasia or in men with persistent OAB symptoms on alpha-blocker therapy demonstrated that tolterodine, in addition to an alpha-blocker therapy, showed no excess incidence of acute urinary retention. However, in these studies, the incidence of adverse events such as dry mouth, constipation, nasal congestion, ejaculation failure, headache, and dysuria was increased in

patients treated with an alpha-blocker in combination with Tolterodine L-tartrate extended release capsules (n=554) compared to those patients treated with an alpha-blocker alone (n=538). An increase in discontinuation due to adverse events was also observed in patients treated with an alpha-blocker in combination with Tolterodine L-tartrate extended release capsules (7%) compared to those treated with an alpha-blocker alone (3%). MYLAN-TOLTERODINE ER should be administered with caution in men who are suspicious of having bladder outlet obstruction (see WARNINGS AND PRECAUTIONS, Gastrointestinal and Genitourinary).

### **Drug-Food Interactions**

Food intake does not result in clinically relevant changes in the pharmacokinetic profile of either the tolterodine immediate release tablets or extended release capsules.

### **Drug-Herb Interactions**

Interaction with herbal products has not been established.

### **Drug-Laboratory Interactions**

Interactions between tolterodine and laboratory tests have not been studied.

### **Patient Counselling**

Patients should be informed that antimuscarinic agents such as **MYLAN-TOLTERODINE ER** (tolterodine L- tartrate extended release capsules) may produce blurred vision or dizziness.

## **DOSAGE AND ADMINISTRATION**

### **Dosing Considerations**

Dosing of **MYLAN-TOLTERODINE ER** (tolterodine L- tartrate extended release capsules) may be affected by the following:

- individual response and tolerability
- impaired hepatic function and renal impairment
- potent CYP3A4 inhibitors

(see **WARNINGS AND PRECAUTIONS** and **DOSAGE AND ADMINISTRATION, Recommended Dose and Dosage Adjustment**)

### **Recommended Dose and Dosage Adjustment**

The initial recommended maximum dose of **MYLAN-TOLTERODINE ER** (tolterodine L- tartrate extended release capsules) is 4 mg once daily. The dose may be reduced to 2 mg once daily based on

individual response and tolerability. However, limited efficacy data are available for tolterodine L-tartrate extended release capsules 2 mg once daily. For patients with impaired hepatic function and patients with renal impairment, the recommended dose is 2 mg once daily (see **WARNINGS AND PRECAUTIONS**). No dosage adjustment for elderly patients ( $\geq 65$  years of age) is recommended (see **WARNINGS AND PRECAUTIONS, Special Populations and DETAILED PHARMACOLOGY**).

Patients treated with potent CYP3A4 inhibitors should not receive doses of **MYLAN-TOLTERODINE ER** greater than 2 mg once daily (see **WARNINGS AND PRECAUTIONS**).

The maximum recommended daily dose of 4 mg should not be exceeded.

### **Administration**

**MYLAN-TOLTERODINE ER** can be taken with food. It should be swallowed whole.

### **OVERDOSAGE**

**For management of a suspected drug overdose, contact your regional Poison Control Centre Immediately.**

The highest dose of tolterodine tartrate given to human volunteers was 12.8 mg as single dose. The most severe adverse events observed were accommodation disturbances and micturition difficulties. One case of overdose has been reported prior to the marketing of the tolterodine immediate release tablets that involved a 27-month-old child who ingested 5 to 7 tablets of tolterodine immediate release 2 mg. He was hospitalized overnight with symptoms of dry mouth and was treated with a suspension of activated charcoal. The child recovered fully.

### **Management of Overdosage**

Treatment of overdosage with **MYLAN-TOLTERODINE ER** (tolterodine L-tartrate extended release capsules) should consist of gastric lavage and activated charcoal. Treatments for symptoms are recommended as follows. For severe central anticholinergic effects (hallucinations, severe excitation), an anticholinesterase agent, such as physostigmine, may be used. If excitation and convulsions occur, administer an anticonvulsant, such as diazepam. Patients with respiratory insufficiency should be given respiratory assistance. If respiratory arrest occurs, patients should be given artificial respiration. Patients with tachycardia may be treated with a beta-blocker, and those with urinary retention may be catheterized. Patients with troublesome mydriasis may be placed in a dark room or treated with pilocarpine eye drops, or both. ECG should be monitored.

In clinical trials of normal volunteers, QT interval prolongation was observed with tolterodine immediate release at doses of 8 mg (4 mg BID). The risk of torsade de pointes with a QT/QTc prolonging drug is usually dose-dependent. It is recommended that continuous ECG monitoring may be appropriate in cases of overdose with **MYLAN-TOLTERODINE ER**. Concomitant therapy should be immediately reviewed and stopped if potential for drug-drug interaction and exacerbation of the QT prolongation effect is possible (see **WARNINGS AND PRECAUTIONS, DRUG INTERACTIONS, Drug-Drug Interactions, DETAILED PHARMACOLOGY, Electrophysiology**).

## ACTION AND CLINICAL PHARMACOLOGY

### Mechanism of Action

Tolterodine L-tartrate, is a competitive muscarinic receptor antagonist, which has been shown to inhibit carbachol-induced contraction of isolated bladder preparations from rats, guinea pigs, and man. Tolterodine L-tartrate (henceforth referred to as tolterodine) inhibits contractions of the detrusor muscle from the guinea pig, and electrically induced contractions of human detrusor muscle from stable and overactive bladders *ex vivo*. Tolterodine is significantly more active in inhibiting acetylcholine-induced urinary bladder contractions than electrically induced salivation in the anesthetized cat.

### Pharmacodynamics

Tolterodine has a pronounced effect on bladder function in healthy volunteers. The main effects following a 6.4 mg single dose of tolterodine were an increase in residual urine, reflecting an incomplete emptying of the bladder, and a decrease in detrusor pressure. These findings are consistent with antimuscarinic action on the lower urinary tract.

In patients with an overactive bladder who received recommended therapeutic doses of the tolterodine immediate release tablets, urodynamic measurements have shown that tolterodine increased the volume at first contraction and maximum cystometric capacity.

Tolterodine is converted to a pharmacologically active 5-hydroxymethyl metabolite (DD 01) by the isozyme cytochrome P450 2D6 (debrisoquine hydroxylase). This metabolite exhibits an antimuscarinic profile similar to that of tolterodine, both *in vitro* and *in vivo*. In view of the antimuscarinic activity of DD 01 and pharmacokinetic data from both humans and animals, it has been concluded that this metabolite contributes significantly to the therapeutic effect in extensive metabolizers (see **Metabolism below, and DETAILED PHARMACOLOGY**).

A dose-effect relationship was established in a Phase II study for the tolterodine extended release capsule (002) for mean residual volume per micturition during 12 hours. The dose of the tolterodine extended release capsule that has the same effect as the tolterodine immediate release tablets, 2 mg twice daily, was estimated to be 4.7 mg (3.7 mg after correction for relative exposure to the active moiety). A dose-effect relationship was also observed for the inhibition of salivation (see **DETAILED PHARMACOLOGY**).

### Pharmacokinetics

**Absorption:** In a study of <sup>14</sup>C-tolterodine in healthy volunteers who received a 5 mg oral dose, at least 77% of the radiolabeled dose was absorbed. Tolterodine immediate release tablets are rapidly absorbed, and maximum serum concentrations ( $C_{max}$ ) occur within 1 to 2 hours after dose administration. The pharmacokinetics of tolterodine immediate release tablets, based on  $C_{max}$  and area under the concentration-time curve (AUC) determinations, are dose-proportional over the range of 1 to 4 mg. Based on the sum of unbound serum concentrations of tolterodine and DD 01, the AUC of tolterodine

extended release capsules, 4 mg once daily, is equivalent to tolterodine immediate release tablets, 2 mg twice daily.  $C_{max}$  and  $C_{min}$  levels of the extended release capsule are about 75% and 150% of the immediate release tablet, respectively, with maximum serum concentrations observed 2 to 6 hours after dose administration. Food intake does not result in clinically relevant changes in the pharmacokinetic profile of either the tolterodine immediate release tablets or extended release capsules (see **DETAILED PHARMACOLOGY**).

**Metabolism:** Tolterodine is extensively metabolized by the liver following oral dosing, and is converted to DD 01 by the isozyme cytochrome P450 2D6. Further metabolism leads to formation of the 5-carboxylic acid and N-dealkylated 5-carboxylic acid metabolites which account for 51% + 14% and 29% + 6.3% of the metabolites recovered in the urine respectively. (see **DETAILED PHARMACOLOGY**).

The potential effect of tolterodine on the pharmacokinetics of other drugs also metabolized by P450 2D6, such as tricyclic antidepressants, some antiarrhythmics and selective serotonin reuptake inhibitors, and neuroleptics has not been formally evaluated.

**Variability in Metabolism:** A subset (about 7%) of the population is devoid of the drug-metabolizing isoenzyme cytochrome P450 2D6, the enzyme responsible for the formation of DD 01. The identified pathway of metabolism for these individuals, referred to as “poor metabolizers” (PMs), is dealkylation via cytochrome P450 3A4 to N-dealkylated tolterodine. The remainder of the population is referred to as “extensive metabolizers” (EMs). Pharmacokinetic studies revealed that tolterodine is metabolized at a slower rate in PMs than in EMs. Since tolterodine and DD 01 have similar antimuscarinic effects, the net activity of **MYLAN-TOLTERODINE ER** is expected to be similar in EMs and PMs (see **DETAILED PHARMACOLOGY**).

**Distribution:** Tolterodine is highly bound to plasma proteins, primarily  $\alpha$ 1-acid glycoprotein. Unbound concentrations of tolterodine average  $3.7\% \pm 0.13\%$  over the concentration range achieved in clinical studies. The 5-hydroxymethyl metabolite (DD 01) is not extensively protein bound, with unbound fraction concentrations averaging  $36\% \pm 4.0\%$ . The blood to serum ratio of tolterodine and DD 01 averages 0.6 and 0.8, respectively, indicating that these compounds do not distribute extensively into erythrocytes. The volume of distribution of tolterodine following administration of a 1.28 mg intravenous dose is  $113 \pm 26.7$  L.

**Excretion:** Following administration of a 5 mg oral dose of  $^{14}\text{C}$ -tolterodine solution to healthy volunteers, 77% of radioactivity was recovered in urine and 17% was recovered in feces in 7 days. Less than 1% (<2.5% in poor metabolizers) of the dose was recovered in urine and feces as intact tolterodine; 5% to 14% (<1% in poor metabolizers) was recovered as DD 01 within the first 24 hours. This is consistent with the apparent half-life of tolterodine: 1.9 to 3.7 hours.

The levels of the serum metabolites other than DD 01 determined in four poor metabolizers and four extensive metabolizers, were comparable for the tolterodine extended release capsule and immediate release tablet.

### **Special Populations and Conditions**

**Age:** No overall differences were observed in safety between older and younger patients on

tolterodine immediate release tablets in Phase III, 12 week, controlled clinical studies; and therefore, no dosage adjustment for elderly patients is recommended (see **DETAILED PHARMACOLOGY**).

**Gender:** There are no sex dependent differences in the pharmacokinetic profile of tolterodine or DD 01.

**Race:** Pharmacokinetic differences due to race have not been identified.

**Hepatic Insufficiency:** Subjects with hepatic cirrhosis exhibit higher serum concentrations and longer half-lives of tolterodine and DD 01 compared to young healthy subjects given the same dose (see **DETAILED PHARMACOLOGY**).

**Renal Insufficiency:** Potential pharmacologic effects and also the toxicological significance of metabolite levels should be taken into account if exposing subjects with renal impairment (GFR < 30 mL/min) to repeated doses of tolterodine (see **DETAILED PHARMACOLOGY**).

## **STORAGE AND STABILITY**

Store at room temperature 15°C to 30°C. Protect from light.

## **SPECIAL HANDLING INSTRUCTIONS**

Not applicable.

## **DOSAGE FORMS, COMPOSITION AND PACKAGING**

**MYLAN-TOLTERODINE ER** (tolterodine L-tartrate extended release capsules) is available as:

**2 mg capsules:** No. 4, blue-green opaque cap and blue-green opaque body, hard-shell gelatin capsule filled with white to off-white beads. The capsule is axially printed with MYLAN over TT 2 in white ink on both the cap and body.

**4 mg capsules:** No. 3, blue opaque cap and blue opaque body, hard-shell gelatin capsule filled with white to off-white beads. The capsule is axially printed with MYLAN over TT 4 in white ink on both the cap and body.

Both strengths are supplied in bottles of 30's and 100's

MYLAN-TOLTERODINE ER (tolterodine L-tartrate extended release capsules) contain the following non-medicinal ingredients: Ethylcellulose, FD&C Blue #2, gelatin, hypromellose, iron oxide yellow (2 mg capsules only), starch, sucrose, talc, titanium dioxide; Ink: povidone, propylene glycol, shellac and sodium hydroxide.

## PART II: SCIENTIFIC INFORMATION

### PHARMACEUTICAL INFORMATION


#### Drug Substance

Proper name: Tolterodine L-Tartrate

Chemical name: (1) (R)-2-[3[bis(1-methylethyl)amino]-1-phenylpropyl]-4-methylphenol  
[R-(R\*,R\*)]-2,3-dihydroxybutanedioate (1:1) (salt)  
(2) (+)-(R)-2-[ $\alpha$ -[2-(diisopropylamino)ethyl]benzyl]-p-cresol L-tartrate  
(1:1) (salt)

Molecular formula and molecular mass: C<sub>26</sub>H<sub>37</sub>NO<sub>7</sub>; 475.6

Structural formula:



Physicochemical properties:

Physical form: White to off-white crystalline powder

Solubility: Soluble in water at room temperature, soluble in methanol, slightly soluble in ethanol and practically insoluble in toluene.

PH: 3.75

pKa: 9.9

Melting point: 213°C



## CLINICAL TRIALS

### COMPARATIVE BIOAVAILABILITY

Comparative bioavailability studies were conducted on MYLAN-TOLTERODINE ER (tolterodine L-tartrate extended-release Capsules) against the reference listed product, <sup>pr</sup>Detrol LA<sup>®</sup> as follows:

- A blinded, single-dose, randomized, two-period, two-treatment, crossover study to investigate the bioequivalence of MYLAN-TOLTERODINE ER 1x4 mg (Mylan Pharmaceuticals ULC) to Detrol LA<sup>®</sup> capsules 1x4 mg (Pfizer Canada Inc) in twenty-seven (27) healthy, adult Asian male volunteers under fasting conditions.
- A blinded, single-dose, randomized, two-period, two-treatment, crossover study to investigate the bioequivalence of MYLAN-TOLTERODINE ER 1x4 mg (Mylan Pharmaceuticals ULC) to Detrol LA<sup>®</sup> capsules, 1x4 mg (Pfizer Canada Inc) in twenty-five (25) healthy, adult Asian male volunteers under fed conditions.
- A blinded, multiple-dose, randomized, two-period, two-treatment, crossover study to investigate the bioequivalence of MYLAN-TOLTERODINE ER 1x4 mg (Mylan Pharmaceuticals ULC) to Detrol LA<sup>®</sup> capsules, 1x4 mg (Pfizer Canada Inc) in thirty-seven (37) healthy, adult Asian male volunteers at steady-state conditions.

#### Summary Table of Comparative Bioavailability Data (from measured data) fasting conditions.

Tolterodine (1 x 4 mg) From measured data Geometric Mean Arithmetic Mean (CV %)				
Parameter	Test*	Reference †	% Ratio of Geometric Means	90% Confidence Interval
<b>AUC<sub>t</sub> (ng•hr/mL)</b>	19.76	21.25	93%	85.14%-101.64%
	28.42 (114.77%)	29.78 (110.31%)		
<b>AUC<sub>inf</sub> (ng•hr/mL)</b>	23.16	24.22	96%	87.86%-104.07%
	31.47 (108.00%)	32.44 (105.39%)		
<b>C<sub>MAX</sub> (ng/mL)</b>	1.50	1.45	103%	92.49%-115.29%
	1.99 (96.86%)	2.06 (109.97%)		
<b>T<sub>1/2</sub> (hr) §</b>	9.00 (49.35%)	8.15 (45.01%)	--	--
<b>T<sub>MAX</sub> (h) §</b>	5.11 (14.70%)	6.70 (42.56%)	--	--

\*Mylan-Tolterodine ER (Tolterodine L-Tartrate Extended-release Capsules), 4 mg (Mylan Pharmaceuticals ULC)

†Detrol<sup>®</sup> LA Capsules, 4 mg (Pfizer Canada Inc)

‡ Geometric mean followed by arithmetic mean (CV%)

§ Expressed as the arithmetic mean (CV%)

#### Summary Table of Comparative Bioavailability Data (from measured data) fed conditions.

Tolterodine (1 x 4 mg) From measured data Geometric Mean Arithmetic Mean (CV %)				
Parameter	Test *	Reference †	% Ratio of Geometric Means	90% Confidence Interval
<b>AUC<sub>t</sub> (ng•hr/mL)</b>	22.70 28.93 (71.87%)	22.65 29.71 (78.97%)	100%	94.18%-106.68%
<b>AUC<sub>inf</sub> (ng•hr/mL)</b>	24.80 31.07 (69.73%)	25.69 35.22 (67.29%)	97%	89.14%-104.59%
<b>C<sub>MAX</sub> (ng/mL)</b>	2.28 2.65 (53.18%)	1.90 2.38 (72.21%)	120%	105.25%-136.42%
<b>T<sub>½</sub> (hr) §</b>	6.61 (43.23%)	7.14 (48.60%)	--	--
<b>T<sub>MAX</sub> (h) §</b>	5.80 (53.14%)	5.92 (9.66%)	--	--

\*Mylan-Tolterodine ER (Tolterodine L-Tartrate Extended-release Capsules), 4 mg (Mylan Pharmaceuticals ULC)

†Detrol® LA Capsules, 4 mg (Pfizer Canada Inc)

\* Geometric mean followed by arithmetic mean (CV%)

§ Expressed as the arithmetic mean (CV%)

### Summary Table of Comparative Bioavailability Data (from measured data) steady state conditions.

Tolterodine (1 x 4 mg) From measured data Geometric Mean Arithmetic Mean (CV %)				
Parameter	Test *	Reference †	% Ratio of Geometric Means	90% Confidence Interval
<b>AUC<sub>TAU</sub> (ng•hr/mL)</b>	16.84 30.60 (138.77%)	17.88 29.08 (120.56%)	94%	83.39%-106.38%
<b>C<sub>MAX</sub> (ng/mL)</b>	1.48 2.46 (122.15%)	1.40 2.15 (107.26%)	106%	91.97%-121.42%
<b>C<sub>MIN</sub> (ng/mL)</b>	0.29 0.49 (185.53%)	0.33 0.49 (154.23%)	90%	77.06%-105.77%
<b>T<sub>MAX</sub> (hr) §</b>	5.16 (17.40%)	5.19 (39.56%)	--	--
<b>FLUCT(%) §</b>	190.89 (24.82%)	163.53 (40.80%)	--	--

\*Mylan-Tolterodine ER (Tolterodine L-Tartrate Extended-release Capsules), 4 mg (Mylan Pharmaceuticals ULC)

†Detrol® LA Capsules, 4 mg (Pfizer Canada Inc)

\* Geometric mean followed by arithmetic mean (CV%)

§ Expressed as the arithmetic mean (CV%)

### Study demographics and trial design

Tolterodine L-tartrate extended release capsules was evaluated in patients with symptoms of

overactive bladder with urinary urge incontinence, frequency, and/or urgency in a large randomized, placebo-controlled, multicenter, double-blind, 12-week study. A total of 507 patients were treated with tolterodine L-tartrate extended release capsules, 4 mg once daily, 514 were treated with tolterodine immediate release tablets, 2 mg twice daily, and 508 were treated with placebo. The majority of patients were Caucasian (95%), with a mean age of 61 years (range, 20 to 93 years). Women (81%) and men (19%) participated in the study; 53% of patients had prior pharmacotherapy for overactive bladder (included responders and nonresponders). At study entry, 97% of patients had at least 5 urge incontinence episodes per week and 91% of patients had 8 or more micturitions per day. The primary efficacy endpoint was change in mean number of incontinence episodes per week at week 12 from baseline. Secondary efficacy endpoints included change in mean number of micturitions per day and mean volume voided per micturition at week 12 from baseline. As shown below, the efficacy results for tolterodine L-tartrate extended release capsules and tolterodine immediate release tablets were significantly better than placebo for all efficacy parameters.

### Study results

<b>Difference between tolterodine L-tartrate extended release capsules (4 mg once daily), tolterodine immediate release tablets (2 mg twice daily) and Placebo for Mean Change at Week 12 from Baseline*</b>			
	<b>Tolterodine extended release capsules (N=507)</b>	<b>Tolterodine immediate release tablets (n=514)</b>	<b>Placebo (N=508)†</b>
Number of incontinence episodes/week			
Mean Baseline	22.1	23.2	23.3
Mean Change from Baseline (%)	-11.8 (53)	-10.6 (46)	-6.9 (30)
p-value‡	0.0001	0.0005	
Number of micturitions/day			
Mean Baseline	10.9	11.1	11.3
Mean Change from Baseline (%)	-1.8 (17)	-1.7 (15)	-1.2 (11)
p-value‡	0.0047	0.0079	
Volume per micturition (mL)			
Mean Baseline	141	137	136
Mean Change from Baseline (%)	34 (24)	29 (21)	14 (10)
p-value‡	0.0001	0.0001	

\*Intent-to-treat analysis

† 1 to 2 patients missing in placebo group for each efficacy parameter

‡Mean change versus placebo

## DETAILED PHARMACOLOGY

### Preclinical Pharmacology

Tolterodine is a competitive muscarinic receptor antagonist, which has been shown to inhibit carbachol-induced contraction of isolated bladder preparations from rats, guinea pigs, and man. Tolterodine is significantly more active in inhibiting acetylcholine-induced urinary bladder contractions ( $ID_{50} = 101 \text{ nmol/kg}$ ) than electrically induced salivation ( $ID_{50} = 257 \text{ nmol/kg}$ ) in the anesthetized cat; whereas oxybutynin exhibits the opposite selectivity profile (urinary bladder contraction  $ID_{50} = 200 \text{ nmol/kg}$ ; salivation  $ID_{50} = 104 \text{ nmol/kg}$ ). At unbound serum concentrations relevant to those observed clinically, tolterodine has no effects on central nervous system (CNS) or intestinal motility in mice. Tolterodine has high affinity for muscarinic receptors and has a very weak affinity for  $\alpha$  - adrenoreceptors, histamine receptors, the neuromuscular junction, and calcium channels.

Preclinical studies have shown that tolterodine is as active as oxybutynin in inhibiting contractions of the detrusor muscle from the guinea pig. Tolterodine also has similar activity to oxybutynin in inhibiting electrically induced contractions of human detrusor muscle from stable and overactive bladders *ex vivo*. These electrically induced contractions are completely blocked by tolterodine.

Effects on the cardiovascular system in conscious dogs, treated orally with tolterodine for 10 days, have been investigated using telemetry technique. Heart rate and diastolic blood pressure were increased at 1 mg/kg (tolterodine 103  $\mu\text{g/L}$ , 5-hydroxymethyl metabolite (DD 01) 25  $\mu\text{g/L}$ ). Except for a prolongation of the QT-interval (10-20 %) observed at 4.5 mg/kg (tolterodine >600  $\mu\text{g/L}$ , DD 01 100  $\mu\text{g/L}$ ), there were no abnormalities of the ECG pattern and no signs of arrhythmias were observed.

In anaesthetised dogs, tolterodine had little or no effect on the cardiovascular and respiratory systems when administered as a continuous i.v. infusion. Marked effects (20-40% prolongation of the QT-interval and T-wave duration) occurred only at tolterodine concentrations 500  $\mu\text{g/L}$ . Heart rate, blood pressure and respiration remained virtually unaffected (1000  $\mu\text{g/L}$ ).

Effects of tolterodine (p.o.) on the central nervous system, gastrointestinal tract and renal function have been evaluated in the mouse. The strict no observed effect level for these effects is 1.5 mg/kg (tolterodine 2.1  $\mu\text{g/L}$ , DD 01 2.4  $\mu\text{g/L}$ ). However, the dose at which effects were observed (15 mg/kg) was in some other studies a no effect dose. The true no observed effect level may therefore be closer to 15 mg/kg than to 1.5 mg/kg. A dose of 15 mg/kg can be expected to result in high serum levels of both tolterodine (83  $\mu\text{g/L}$ ) and DD 01 (63  $\mu\text{g/L}$ ).

Most of the effects observed at high doses in the mouse ( $\geq 15 \text{ mg/kg}$ ) and dog ( $\geq 1 \text{ mg/kg}$ ) were antimuscarinic in nature. Increased locomotor activity, mydriasis, decreased intestinal motility, increased residual urine and increased heart rate can all be attributed to the primary action of tolterodine and DD 01 on muscarinic receptors. Preclinical studies have shown that DD 01 exhibits a similar antimuscarinic profile to that of tolterodine, and a greater antimuscarinic activity on the bladder relative to the salivary gland *in vivo*.

The degree of serum protein binding differs between species and this must be taken into account when comparisons to humans are made. Thus, the unbound concentrations of tolterodine (2.2 µg/L) and DD 01 (8 µg/L) at which an increased heart rate was observed in the dog, are 17 and 8 times higher than the unbound serum concentrations achieved in most patients treated with tolterodine 2 mg bid (tolterodine: 0.13 µg/L; DD 01: 1.04 µg/L). The unbound concentrations at which effects on the central nervous system, intestinal motility and renal function were observed in the mouse (tolterodine: 13 µg/L; DD 01: 45 µg/L) are approximately 100 and 40 times, respectively, higher than those expected to be achieved in patients. Almost the same factors (100 and 30 times) were calculated for the unbound concentrations at which a slight QT-prolongation was recorded in the conscious dog (tolterodine: 13 µg/L; DD 01: 32 µg/L).

## **Clinical Pharmacology**

### ***Pharmacodynamics***

After oral administration, tolterodine is metabolized in the liver, resulting in the formation of the 5-hydroxymethyl derivative, a major pharmacologically active metabolite. The 5-hydroxymethyl metabolite (DD 01), which exhibits an antimuscarinic activity similar to that of tolterodine, contributes significantly to the therapeutic effect. Both tolterodine and DD 01 exhibit a high affinity for muscarinic receptors and have a very weak affinity for  $\alpha$ -adrenoreceptors, histamine receptors, neuromuscular junction, and calcium channels.

Preclinical studies have shown that tolterodine is as active as oxybutynin in inhibiting contractions of the detrusor muscle from the guinea pig; it has a potency similar to that of oxybutynin in inhibiting electrically induced contractions of human detrusor muscle from stable and overactive bladders *ex vivo*.

### ***Bioavailability***

The absolute bioavailability of the tolterodine immediate release tablet was determined using a 1.28 mg intravenous dose as reference. Reported values in the oral dose interval 3.2 - 12.8 mg were 29- 39%. In selected extensive metabolizers (EMs) and poor metabolizers (PMs) the bioavailability was 17±9% and 65±26%. This difference is explained by a higher degree of first-pass metabolism in EMs. The bioavailability estimate as such is, however, not an informative parameter with respect to clinical effect, since DD 01 is found in pharmacologically active concentrations in the majority of the population (EMs).

In a phase II study (002), serum concentrations of tolterodine and DD 01 were assessed in 25 to 29 patients per dose group. At the end of the 7-day treatment using the prototype tolterodine extended release capsule, blood samples were taken periodically during the 24 hours following the last dose of capsules (2, 4, 6 or 8 mg), and 12 hours following tolterodine immediate release tablets, 2 mg bid.

The relative bioavailability (dose-normalized AUC<sub>t</sub> estimated over all doses) of the prototype extended release capsule used compared with the immediate release tablet averaged 71% for tolterodine and 73% based on DD 01. The relative exposure to the active moiety from the prototype capsule used compared with the tablet averaged 79% based on dose-normalized AUC<sub>12</sub>.

### ***Pharmacokinetics***

**Absorption:** In a study of <sup>14</sup>C-tolterodine in healthy volunteers who received a 5 mg oral dose, at least 77% of the radiolabeled dose was absorbed. Tolterodine is rapidly absorbed, and maximum serum concentrations (C<sub>max</sub>) typically occur within 1 to 2 hours after dose administration. The pharmacokinetics of tolterodine immediate release tablets, based on C<sub>max</sub> and area under the concentration-time curve (AUC) determinations, are dose-proportional over the range of 1 to 4 mg.

After single-dose administration, dose-normalized AUC (corrected for differences in dose) of both tolterodine and DD 01 showed equivalence after administration of the final extended release capsule (2x4 mg) and the immediate release tablets (2x2mg). C<sub>max</sub> for the capsule was markedly lower than for the tablet.

After multiple-dose administration, the final extended release capsule was equivalent to the immediate release tablet based on AUC for the active moiety. The extended release capsule showed the desired extended release properties with C<sub>max</sub> levels of the active moiety, which were approximately 75% of the tablet C<sub>max</sub>. C<sub>min</sub> levels (active moiety) were about 1.4 times higher for the capsule. The fluctuation index was consequently lower for the capsule than for the tablet.

Food intake does not result in clinically relevant changes in the pharmacokinetic profile of either the tolterodine immediate release tablets or extended release capsules.

**Metabolism:** Tolterodine is extensively metabolized by the liver following oral dosing. The primary metabolic route involves the oxidation of the 5-methyl group and is mediated by the isoenzyme cytochrome P450 2D6 and leads to the formation of a major pharmacologically active 5-hydroxymethyl metabolite. Further metabolism leads to formation of the 5-carboxylic acid and N-dealkylated 5-carboxylic acid metabolites, which account for 51% ± 14% and 29% ± 6.3%, respectively, of the metabolites recovered in the urine.

Variability in Metabolism: A subset (about 7%) of the population is devoid of the drug-metabolizing isoenzyme cytochrome P450 2D6, the enzyme responsible for the formation of DD 01. The identified pathway of metabolism for these individuals, referred to as “poor metabolizers” (PMs), is dealkylation via cytochrome P450 3A4 to N-dealkylated tolterodine. The remainder of the population is referred to as “extensive metabolizers” (EMs). Pharmacokinetic studies revealed that tolterodine is metabolized at a slower rate in PMs than in EMs. This results in significantly higher serum concentrations of tolterodine and in negligible concentrations of DD 01. Because of differences in the protein-binding characteristics of tolterodine and DD 01, the sum of unbound serum concentrations of tolterodine and DD 01 is similar in EMs and PMs at steady state. Since tolterodine and DD 01 have similar antimuscarinic effects, the net activity of **MYLAN-TOLTERODINE ER** (tolterodine L-tartrate extended release capsules) is expected to be similar in EMs and PMs.

The  $AUC_{24}$  data for serum metabolites measured in 4 PMs and 4 EMs after multiple-dose administration indicated that the levels of the known serum metabolites are similar for the tolterodine extended release capsules and immediate release tablets. As for the tablet, the exposure to the active moiety is within the same range after administration to PMs and EMs.

The PMs in this study had levels of the active moiety that are within the range observed in the EMs, both for the extended release capsules and immediate release tablets. For the PMs, the effect of the sustained release and slower absorption of tolterodine is less pronounced than for the EMs.

**Excretion:** Following administration of a 5 mg oral dose of  $^{14}C$ -tolterodine to healthy volunteers, about 77% of radioactivity was recovered in urine and 17% was recovered in feces. Less than 1% (<2.5% in PMs) of the dose was recovered as intact tolterodine, and 5% to 14% was recovered as the active DD 01 metabolite. Most of the radioactivity was recovered within the first 24 hours, which is consistent with the apparent half-life of tolterodine: 1.9 to 3.7 hours in pharmacokinetic studies.

### ***Pharmacokinetics in Special Populations***

**Age:** In phase I multiple-dose studies in which tolterodine immediate release tablets 2 mg were administered twice daily, serum concentrations of tolterodine and of DD 01 were similar in healthy elderly volunteers (aged 64 through 80 years) and healthy young volunteers (aged less than 40 years). In another phase I study, elderly volunteers (aged 71 through 81 years) were given tolterodine immediate release tablets, 1 or 2 mg twice daily. Mean serum concentrations of tolterodine and DD 01 in these elderly volunteers were approximately 20% and 50% higher, respectively, than reported in young healthy volunteers. However, no overall differences were observed in safety between older and younger patients in phase III, 12-week, controlled clinical studies; and therefore, no dosage adjustment is recommended (see **WARNINGS AND PRECAUTIONS, Special Populations**).

**Pediatric:** The pharmacokinetics of the extended release capsules have not been established in pediatric patients.

**Gender:** Pharmacokinetic data from three Phase I clinical studies (Studies 022, 024, and 028) in which a tolterodine immediate release tablet dose of 2 mg was administered in the fasting state were analyzed with respect to gender. The pharmacokinetics of tolterodine and DD 01 are not influenced by gender. Mean  $C_{max}$  of tolterodine (1.6  $\mu\text{g/L}$  in males versus 2.2  $\mu\text{g/L}$  in females) and DD 01 (2.2  $\mu\text{g/L}$  in males versus 2.5  $\mu\text{g/L}$  in females) are similar in males and females who were administered tolterodine immediate release tablets 2 mg. Mean AUC values of tolterodine (6.7  $\mu\text{g/h/L}$  in males versus 7.8  $\mu\text{g/h/L}$  in females) and DD 01 (10  $\mu\text{g/h/L}$  in males versus 11  $\mu\text{g/h/L}$  in females) are also similar. The elimination half-life of tolterodine immediate release tablets for both males and females is 2.4 hours, and the half-life of DD 01 is 3.3 hours in males and 3.0 hours in females.

**Race:** Differences among races regarding metabolic capacity can be assumed to be of quantitative nature and are probably less than the thoroughly documented difference between extensive and poor

metabolizers. The few non-Caucasians included do not show a different pharmacokinetic profile of tolterodine or DD 01.

**Renal Impairment:** A study was conducted to evaluate the pharmacokinetics of tolterodine immediate release tablets in 12 subjects with renal impairment compared to 12 healthy volunteers. The exposure to unbound tolterodine and DD 01 was on average 2-3 fold higher in patients with renal impairment compared with healthy volunteers. AUC of N-dealkylated tolterodine was in an extreme case, about 60-fold higher in a poor metabolizer (PM) in the renal impairment group than in the only healthy extensive metabolizer (EM) with quantifiable AUC. However, the corresponding ratio for what is generally observed in healthy PMs is about 10. Tolterodine acid levels and N-dealkylated tolterodine acid were on average 5 times and 11 times higher, respectively, in the renal impairment group with respect to AUC (extreme case 9-fold and 31-fold higher than most exposed healthy subjects). Potential pharmacologic effects and also the toxicological significance of metabolite levels should be taken into account if exposing subjects with renal impairment (GFR < 30 mL/min) to repeated doses of tolterodine (see **WARNINGS AND PRECAUTIONS**).

**Hepatic Insufficiency:** As might be predicted from a drug in which hepatic metabolism is the primary route of elimination, liver impairment can significantly alter the disposition of tolterodine. In a study of cirrhotic patients, elimination half-life of tolterodine immediate release tablets was longer in cirrhotic patients (mean, 8.7 hours) than in healthy, young and elderly volunteers (mean, 2 to 4 hours). The clearance of orally administered tolterodine immediate release tablets was substantially lower in cirrhotic patients ( $1.1 \pm 1.7$  L/h/kg) than in the healthy volunteers ( $5.7 \pm 3.8$  L/h/kg). Patients with significantly reduced hepatic function should not receive doses of **MYLAN-TOLTERODINE ER** (tolterodine L-tartrate extended release capsules) greater than 2 mg daily (see **WARNINGS AND PRECAUTIONS**).

### ***Drug Interactions***

**Fluoxetine:** Fluoxetine is a selective serotonin reuptake inhibitor and a potent inhibitor of cytochrome P450 2D6 activity. In a study to assess the effect of fluoxetine on the pharmacokinetics of tolterodine and its metabolites, it was observed that fluoxetine significantly inhibited the metabolism of tolterodine immediate release tablets in extensive metabolizers, resulting in a 4.8-fold increase in tolterodine AUC. However, DD 01 showed a 52% decrease in  $C_{max}$  and a 20% decrease in AUC. Fluoxetine thus alters the pharmacokinetics in patients who would otherwise be extensive metabolizers of tolterodine to resemble the pharmacokinetic profile in poor metabolizers. The sums of unbound serum concentrations of tolterodine and DD 01 are 25% higher during the interaction. However, no dose adjustment is required when **MYLAN-TOLTERODINE ER** and fluoxetine are coadministered (see **DRUG INTERACTIONS**).

**Other Drugs Metabolized by P450 2D6:** The potential effect of **MYLAN-TOLTERODINE ER** (tolterodine L-tartrate extended release capsules) on the pharmacokinetics of drugs that are metabolized by P450 2D6 (such as flecainide, vinblastine, carbamazepine, tricyclic antidepressants) has not been formally evaluated (see **DRUG INTERACTIONS**).



**Warfarin:** In healthy volunteers, coadministration of tolterodine immediate release tablets 2 mg twice daily for 7 days and a single dose of warfarin 25 mg on day 4 had no effect on prothrombin time, Factor VII suppression, or on the pharmacokinetics of warfarin.

**Oral Contraceptives:** Tolterodine immediate release tablets, 2 mg twice daily, has no effect on the pharmacokinetics of an oral contraceptive (ethinyl estradiol 30 µg; levonorgestrel 150 µg) as evidenced by the monitoring of ethinyl estradiol and levonorgestrel over a 2-month cycle in healthy female volunteers.

**Diuretics:** Coadministration of tolterodine immediate release tablets up to 4 mg twice daily for up to 12 weeks with diuretic agents, such as indapamide, hydrochlorothiazide, triamterene, bendroflumethiazide, chlorothiazide, methylchlorothiazide, or furosemide, did not cause any adverse electrocardiographic (ECG) effects in patients with overactive bladder.

**Cytochrome P450 3A4 inhibitors:** The use of tolterodine immediate release tablets in combination with ketoconazole, a potent CYP3A4 inhibitor, was studied in 8 healthy subjects, all of whom were poor metabolizers of CYP2D6. Concomitant treatment with ketoconazole resulted in a 2.2 fold increase in tolterodine AUC at steady state. Based on these findings, potent CYP3A4 inhibitors such as macrolide antibiotics (erythromycin and clarithromycin) or azole antifungal agents (ketoconazole, itraconazole and miconazole), or cyclosporin or vinblastine may also lead to increases of tolterodine plasma concentrations (see **DRUG INTERACTIONS**).

A clinical explorative study with marker drugs for the major P450 isoenzymes suggests that metabolic activity of CYP2D6, 2C9, 2C19, 3A4 or 1A2 is unlikely to be inhibited by tolterodine immediate release tablets.

### ***Electrophysiology***

The QT effect of 2 mg BID and 4 mg BID doses of tolterodine L-tartrate immediate release tablets was evaluated in a steady-state, 4-way crossover, double-blind, placebo- and active-controlled (moxifloxacin 400 mg QD) study in 48 healthy volunteers (18-55 yrs age, with approximately equal representations of males and females and of CYP2D6 poor and extensive metabolizers). The QT interval was measured over a 12-hour period including peak times at steady state. This evaluation was done at up to two times the highest dose of tolterodine L-tartrate immediate release tablets at peak exposures equivalent to three times the highest dose of tolterodine L-tartrate extended release capsules.

The following table summarizes the largest time-matched, placebo and baseline-adjusted mean effects on Fridericia-corrected QTc (QTcF) at steady-state. The mean increase of heart rate associated with a 4 mg/day dose of tolterodine in this study was 2.0 beats/minute and 6.3 beats/minute with 8 mg/day tolterodine. The change in heart rate with moxifloxacin was 0.5 beats/minute.

### **Largest Time-Matched, Placebo and Baseline-Adjusted Mean Effects on Fridericia-corrected QTc (QTcF) at Steady-State**

Treatment Dose	Multiple of Maximum Recommend Dose	Machine-Read QTcF(msec) **			Manually-Read QTcF(msec) **		
		Time of Max Increase	Point Estimate*	90% Confidence Interval	Time of Max Increase	Point Estimate*	90% Confidence Interval
Tolterodine 2 mg BID	1 X	3	1.4	-2.8, 5.6	1	5.0	1.0, 9.0
Tolterodine 4 mg BID	2 X	1	5.6	2.2, 9.1	1	11.8	7.9, 15.8
Moxifloxacin 400 mg QD	1 X	4	13.5	9.9, 17.1	4	22.4***	19.3, 27.1

\* The point estimate is the difference between arithmetic means for pair-wise comparisons of the drug versus placebo treatments. QTc values are corrected for heart rate using Fridericia's formula ( $QTc=QT/RR^{0.33}$ )

\*\* The machine-read methodology is based on earliest Q onset to latest T offset in 12 simultaneous recorded leads, while the manual over-read method is based on lead II only.

The reason for the difference between machine and manual read of QT interval is unclear.

\*\*\* The effect on QTc interval with 4 days of moxifloxacin dosing in this QT trial may be greater than typically observed in QT trials of other drugs.

The QT effect appeared greater for 8 mg/day compared with 4 mg/day tolterodine immediate release tablets. The effect of the highest tolterodine dose (two times the therapeutic dose) was 50-60% less than that of the active control moxifloxacin (400 mg) at its therapeutic dose. Tolterodine's effect on QT interval was found to correlate with plasma concentration of tolterodine. The peak exposures of tolterodine and its 5-hydroxymethyl metabolite following dosing with tolterodine L-tartrate extended release capsules are about 61% and 67%, respectively, compared with tolterodine L-tartrate tablets. Since the QT prolongation effect is in linear relationship with exposure, any QT effect of tolterodine L-tartrate extended release capsules would also be expected to be similarly lower. The effect on QTc interval appeared to be greater in CYP2D6 poor metabolizers than in CYP2D6 extensive metabolizers. In this study, the point estimates of manual-read QTc interval increase were 2.1 msec in extensive metabolizers and 8.7 msec in poor metabolizers receiving tolterodine 2 mg BID treatment. However, this study was not designed to make direct statistical comparisons by CYP2D6 metabolizer status nor between drugs or dose levels. At both doses of tolterodine, no subject, irrespective of their metabolic profile (ie. poor/extensive metabolizers), exceeded 500 msec for absolute QTcF or 60 msec for change from baseline. The clinical relevance of these findings will depend on individual patient risk factors and susceptibilities present (see **WARNINGS AND PRECAUTIONS, Cardiovascular**).

## TOXICOLOGY

### Acute toxicity

The single oral dose administration studies in mice, rats and dogs showed species differences. At 300 mg/kg in mice, a 10-60% mortality was recorded, whereas 375 mg/kg was non-lethal in rats. In mice, a dose of 200 mg/kg caused no lethality. In the dog, at 40 mg/kg (the highest dose tested) no mortality occurred, but pronounced clinical signs were seen such as decreased locomotor activity, clouding of consciousness and stupor. Following a single intravenous dose, 8 mg/kg was a no

observed effect level in both rats and mice. At 24 mg/kg, 30% mortality was recorded in rats, and 80% mortality in mice.

### **Long-term toxicity**

The metabolic profiles in urine from the mouse, rat, dog and man given an oral dose of radioactively labeled tolterodine show that the mouse, dog and man have a similar metabolic pattern including the formation of the pharmacologically active 5-hydroxymethyl metabolite, DD 01. In contrast, the metabolism of tolterodine in the rat is more extensive and occurs also via other pathways involving mono- and dihydroxylation of the unsubstituted benzene ring. The mouse is considered to be a more appropriate species than the rat for the safety evaluation of tolterodine in man.

Mouse In the 2 week study, dose levels of 4, 12, 40 or 80 mg/kg/day were used, and in the 13 week study, the dose levels were 4, 12 or 40 mg/kg/day. In the 26 week study dose levels of 3, 10 or 30 mg/kg/day were used. In the 2 week study, no toxicity was found after doses up to 80 mg/kg/day. During the 13 week study, 7 males and 8 females receiving 40 mg/kg/day died shortly after dosing. Treatment related deaths also occurred in the 26 week study, where 12 males and 15 females treated at 30 mg/kg/day died within one hour of dosing. In both studies, the deaths were distributed throughout the treatment period starting from the second week of treatment. Although the mechanism of the unexpected deaths is unknown, it is most likely related to exaggerated pharmacological effects (circulatory and/or respiratory failure) occurring at serum peak levels.

Rat In the 13 week repeated dose study in rats, doses of 4, 12 or 40 mg/kg/day were given. In females given 40 mg/kg/day depressed body weight gain and reduced food consumption were recorded. Also, ten female rats died approximately 20 hours after dosing. The deaths occurred from week 3. Cause of death could not be established, but is most likely related to exaggerated pharmacologic effects (circulatory and/or respiratory failure) following the accumulation of tolterodine with time.

Dog The clinical signs that were associated with tolterodine treatment in the 13 week, 26 week and 52 week (0.5, 1.5 or 4.5 mg/kg/day) studies were characterized mainly by dose related peripheral antimuscarinic effects, i.e. dry mouth, mydriasis and dryness of the eye. In some dogs receiving 1.5 or 4.5 mg/kg/day, diminished lacrimation caused conjunctivitis and/or corneal changes especially at the high dose level.

Tolterodine, as well as its active human metabolites prolong action potential duration (90% repolarization) in canine Purkinje fibers (23 - 123 times therapeutic levels) and block the K<sup>+</sup>-current in cloned human ether-a-go-go-related gene (hERG) channels (0.8 – 14.7 times therapeutic levels). In dogs prolongation of the QT interval has been observed after application of tolterodine and its human metabolites (5.1 - 62.7 times therapeutic levels).

Central antimuscarinic effects, i.e. locomotor disturbances and drowsiness, were seen in all three studies on day 1, in a few dogs receiving 4.5 or 8 mg/kg/day. These symptoms occurred in dogs

with high serum concentrations of tolterodine ( $C_{\max}$  800-1250  $\mu\text{g/L}$ ), and DD 01. Ataxia and tremor were also observed occasionally in high dose animals during the 26 week study.

### **Carcinogenicity**

Carcinogenicity studies with tolterodine were conducted in mice and rats. At the maximum- tolerated dose in mice (30 mg/kg/day [123 mg/m<sup>2</sup>/day]), female rats (20 mg/kg/day), and male rats (30 mg/kg/day), AUC values obtained for tolterodine were 355, 291, and 462  $\mu\text{g}\cdot\text{h/L}$ , respectively. In comparison, the human AUC value for a 2-mg dose administered twice daily is estimated at 34  $\mu\text{g}\cdot\text{h/L}$ . Thus, tolterodine exposure in the carcinogenicity studies was 9- to 14-fold higher than expected in humans. No increase in tumors was found in either mice or rats.

### **Mutagenicity**

No mutagenic effects of tolterodine were detected in a battery of *in vitro* tests, including bacterial mutation assays (Ames test) in four strains of *Salmonella typhimurium* and in two strains of *Escherichia coli*, a gene mutation assay in L5178Y mouse lymphoma cells, and chromosomal aberration tests in human lymphocytes. Tolterodine was also negative *in vivo* in the bone marrow micronucleus test in the mouse.

### **Reproduction and Teratology**

In female mice treated for 2 weeks before mating and during gestation with 20 mg/kg/day (corresponding to AUC value of about 500  $\mu\text{g}\cdot\text{h/L}$ ), neither effects on reproductive performance or fertility nor any anomalies or malformations were seen. Based on AUC values, the systemic exposure was about 15-fold higher in animals than in humans. At doses of 30 to 40 mg/kg/day, tolterodine caused a dose-related increase in embryoletality, reduced fetal weight, and increased incidence of fetal abnormalities. At these doses, AUC values were about 20- to 25-fold higher than in humans. In male mice, a dose of 30 mg/kg/day did not induce any adverse effects on fertility.

## REFERENCES

1. Abrams P, Jackson S, Mattiasson A, et. al. A randomised, double-blind, placebo controlled, dose-ranging study of the safety and efficacy of tolterodine in patients with hyperreflexia [abstract]. 26th Annual Meeting of International Continence Society; 27-30 August 1996; Athens, Greece: International Continence Society, 1996. Publication Citation.
2. Bertilsson L. Geographical/interracial differences in polymorphic drug oxidation. Current state of knowledge of cytochromes P450 (CYP) 2D6 and 2C19. Clin Pharmacokinet 1995;29:192-209.
3. Brøsen K, Gram LF, Haghfelt T, Bertilsson L. Extensive metabolizers of debrisoquine become poor metabolizers during quinidine treatment. Pharmacology & Toxicology 1987;60:312-4.
4. Brynne N, Olofsson S, Hallén B, Grälls M. Pharmacokinetics of tolterodine in subjects with renal failure compared to healthy volunteers. An open controlled non-randomized parallel group study. CTN 97-OATA-040. Document c-0013132 (1999) - Data on file.
5. Brynne N, Stahl M, Hallén B, Edlund PO, Palmér L. Clinical Pharmacokinetics of Tolterodine: A New Drug in the Treatment of Urge Incontinence [abstract]. Therapie 1995; 50 (Suppl): abstr 353. Publication Citation.
6. Chapple C, Herschorn S, Abrams P, Sun F, Brodsky M, Guan Z. Tolterodine Treatment Improves Storage Symptoms Suggestive of Overactive Bladder in Men Treated With alpha-Blockers. Eur Urol 2009;56(3):534-41.
7. Eichelbaum M, Gross AS. The genetic polymorphism of debrisoquine/sparteine metabolism-clinical aspects. Pharmacol Ther 1990;46:377-94.
8. Ekström B, Stahl M, Mattiasson A, Andersson K-E. Effects of Tolterodine on Bladder Function in Healthy Volunteers [abstract]. AUA; J Urol 1995;153 (4 Suppl): 394A. Publication Citation.
9. Fantl JA, Newman DK, Co-Chair. Treatment of urinary incontinence. In: Urinary incontinence in adults: acute and chronic management, clinical practice guideline; Number 2, 1996 update. US Department of Health and Human Services, Agency for Health Care Policy and Research, 1996: 31-73.
10. Freeman R, Hill S, Millard R, Slack M, Sutherst J. Tolterodine study group. Reduced perception of urgency in treatment of overactive bladder with extended-release tolterodine. Obstet Gynecol 2003;102(3):605-11.
11. Frewen W. Role of bladder training in the treatment of the unstable bladder in the female. Urol Clin North Am 1979;6:273-7.
12. Griebing TL, Kraus SR, Richter HE, Glasser DB, Carlsson M. Tolterodine extended release is well tolerated in older subjects. Int J Clin Pract. 2009; 63(8):1198-204.

13. Gustafsson L. The influence of ketoconazole on the pharmacokinetics and safety of tolterodine. An open, single group study in healthy volunteers. CTN 97-OATA-036 Pharmacia & Upjohn Document 9810186 (1998) - Data on file.
14. Kaplan SA, Roehrborn CG, Rovner ES, Carlsson M, Bavendam T, Guan Z. Tolterodine and tamsulosin for treatment of men with lower urinary tract symptoms and overactive bladder: a randomized controlled trial. *JAMA* 2006;296(19):2319-28.
15. Keam SJ, Perry CM. Management of overactive bladder: Defining the role of extended release tolterodine. *Dis Manage Health Outcomes* 2004;12 (2): 121-142.
16. Khullar V, Hill S, Laval KU, Schiøtz HA, Jonas U, Versi E. Treatment of urge-predominant mixed urinary incontinence with tolterodine extended release: A randomized, placebocontrolled trial. *urology*. 2004;64(2):269-75.
17. Landis JR, Kaplan S, Swift S, Versi E. Efficacy of antimuscarinic therapy for overactive bladder with varying degrees of incontinence severity. *J Urol* 2004;171(2 Pt 1):752-6.
18. Larsson G, Hallén B, Nilvebrant L. Tolterodine in the Treatment of Urge Incontinence. Analysis of the Pooled Phase II Efficacy and Safety Data [abstract]. 26th Annual Meeting of International Continence Society; 27-30 August 1996; Athens, Greece: International Continence Society, 1996. Publication Citation.
19. Layton D, Pearce GL, Shakir SAW. Safety profile of tolterodine as used in general practice in England – Results of prescription-event monitoring. *Drug Saf* 2001; 24(9):703-13.
20. Malone-Lee J, Lubel D, Szonyi G. Low dose oxybutynin for the unstable bladder [abstract]. *BMJ* 1992;304:1053.
21. Marinac JS, Foxworth JW, Willsie SK. Dextromethrophan polymorphic hepatic oxidation (CYP2D6) in healthy black American adult subjects. *Therapeutic Drug Monitoring* 1995;17,120-4.
22. Maurice M, Pichard L, Daujat M, Fabre I, Joyeux H, Domergue J, Maurel P. Effects of imidazole derivates of cytochromes P450 from human hepatocytes in primary culture. *The FASEB Journal* 1992; 6:752-8.
23. Messelink EJ, Soler JM, Madersbacher H, et al. Urodynamic Aspects of the Efficacy of Tolterodine, A New Antimuscarinic Drug in the Treatment of Detrusor Hyperreflexia [abstract]. 25th Annual Meeting of International Continence Society; 17-20 October 1995; Sydney, Australia: International Continence Society, 1995. Publication Citation.
24. Naerger H, Fry CH, Nilvebrant L. Effect of tolterodine on electrically induced contractions of isolated human detrusor muscle from stable and unstable bladders [abstract]. *Neurology and Urodynamics* 1995;14:524-6.

25. Olsson B, Szamosi J. Multiple dose pharmacokinetics and pharmacodynamics of tolterodine prolonged release capsules in comparison with tolterodine immediate release tablets. An open-randomized, cross-over trial in healthy volunteers. Final report of the trial CTN 98-TOCR-006 Pharmacia & Upjohn Document No. c0008272 (13 October 1999) – data on file.
26. Olsson B, Szamosi J. The effect of food on the bioavailability of tolterodine prolonged release capsules. An open, randomized, cross-over trial in healthy volunteers. Final report of the trial CTN 98-TOCR-010. Pharmacia & Upjohn Document No. c0003212 (25 February 1999) – data on file.
27. Otton SV, Wu D, Joffe RT, Cheung SW and Sellers EM. Inhibition by fluoxetine of cytochrome P450 2D6 activity. *Clin Pharmacol Ther* 1992;53:401-9.
28. Ouslander JG, Blaustein J, Coonor A, Orzeck S, Yong CL. Pharmacokinetics and clinical effects of oxybutynin in geriatric patients. *Journal of Urology* 1988;140: 47-50.
29. Ouslander JG. Management of overactive bladder. *N Engl J Med*. 2004;350(8): 786-99.
30. Rentzhog L, Abrams P, Cardozo L, et al. Tolterodine - A New Bladder Selective Drug for the Treatment of Detrusor Instability [abstract]. 25th Annual Meeting of International Continence Society; 17-20 October 1995; Sydney, Australia: International Continence Society, 1995. Publication Citation.
31. Riva D, Casolati E. Oxybutynin chloride in the treatment of female idiopathic bladder instability. Results from double blind treatment. *Clin Exp Obstet Gynecol* 1984;11:37-42.
32. Salvatore S, Khullar V, Cardozo L, Kelleher CJ, Abbott D, Hill S. Long term outcome of women with detrusor instability treated with oxybutynin [abstract]. *Neurourol Urodyn* 1995;14:460-1.
33. Söderström J, Szamosi J. Clinical efficacy and tolerability/safety of tolterodine prolonged release capsules and tolterodine immediate release tablets vs. placebo. A randomized, double-blind, placebo-controlled, multinational study in patients with symptoms of overactive bladder. Final report of the trial 98-TOCR-007. Pharmacia & Upjohn Document No. c0013194 (13 December 1999) – data on file.
34. Stahl M, Brynne N, Ekström, et al. Pharmacokinetics of Tolterodine in Relation to Effects on the Bladder in Healthy Volunteers [abstract]. *Therapie* 1995; 50 (Suppl): abstr 355. Publication Citation.
35. Stahl MMS, Ekström B, Sparf B, Mattiasson A, Andersson K-E. Urodynamic and Other Effects of Tolterodine: A Novel Antimuscarinic Drug for the Treatment of Detrusor Overactivity [abstract]. *Neurourol Urodyn* 1995; 14:647-55. Publication Citation.
36. Strömberg J, Vågerö M, Olsson B. Dose effect trial of tolterodine prolonged release capsules. A double-blind, double-dummy, cross-over trial in patients with overactive bladder. Final report of the study 97-TOCR-002. Pharmacia & Upjohn Document No. c0003471 (29 November 1999) – data on file.

37. Tapp AJS, Cardozo LD, Versi E, Cooper D. The treatment of detrusor instability in postmenopausal women with oxybutynin chloride: a double blind placebo controlled study. *Br J Obstet Gynaecol* 1990; 97: 521-6.
38. Thomas TM, Plymat KR, Blannin J, Meade TW. Prevalence of urinary incontinence. *BMJ* 1980;281:1243-5.
39. Thüroff JW, Bunke B, Ebner A, Faber P, de Geeter P, Hannappel J, Heidler H, Madersbacher H, Melchoir H, Schafer W, Schwenzer T, Stockle M. Randomized, doubleblind, multicenter trial on treatment of frequency, urgency and incontinence related to detrusor hyperactivity: Oxybutynin versus propantheline versus placebo. *J Urol* 1991;145:813-7.
40. Urinary incontinence in adults - National Institute of Health consensus statement. *JAMA* 1989;261:2685-90.
41. Van Kerrebroeck P, Kreder K, Jonas U, Zinner N, Wein A. Tolterodine Once-Daily: Superior Efficacy and Tolerability in the Treatment of the Overactive Bladder. *Urology* 2001;57(3):414-21.
42. Winter S. Tolterodine: a new drug for urinary incontinence. *Inpharma* 1996;1040. Publication Citation.
43. Yarker YE, Goa KL, Fitton A. Oxybutynin. A review of its pharmacodynamic and pharmacokinetic properties, and its therapeutic use in detrusor instability. *Drugs and Aging* 1995;6:243-62.
44. Zinner NR, Mattiasson A, Stanton S.L. Efficacy, Safety, and Tolerability of Extended-Release Once-Daily Tolterodine Treatment for Overactive Bladder in Older versus Younger Patients. *J Am Geriatr Soc* 2002; 50: 799-807.
45. DETROL LA<sup>®</sup> Product Monograph, Pfizer Canada Inc. Submission Control No. 180260. Last Revised: February 2, 2015.



### PART III: CONSUMER INFORMATION

#### Pr MYLAN-TOLTERODINE ER

(tolterodine L-tartrate extended release capsules)

This leaflet is part III of a three-part "Product Monograph" published when MYLAN-TOLTERODINE ER (tolterodine L-tartrate extended release capsules) was approved for sale in Canada and is designed specifically for Consumers. This leaflet is a summary and will not tell you everything about MYLAN-TOLTERODINE ER. Contact your doctor or pharmacist if you have any questions about the drug.

#### ABOUT THIS MEDICATION

##### What the medication is used for:

The name of this medication is MYLAN-TOLTERODINE ER. It is used for the treatment of the symptoms of overactive bladder which include frequency, urgency, and urge incontinence.

**REMEMBER: This medication is for YOU. Never give it to others. It may harm them even if their symptoms are the same as yours.**

##### What it does:

Tolterodine works to prevent bladder contractions or spasms. This results in more bladder capacity and less frequency, urgency and involuntary loss of urine.

##### When it should not be used:

You should not take MYLAN-TOLTERODINE ER if you have:

- urinary retention,
- gastric retention,
- uncontrolled narrow angle glaucoma,
- known hypersensitivity to tolterodine L-tartrate or any of the other ingredients in MYLAN-TOLTERODINE ER.

##### What the medicinal ingredient is:

Each capsule contains film-coated beads of 2 mg or 4 mg of the active ingredient, tolterodine L-tartrate. The film-coated beads dissolve over time, releasing the active ingredient over 24 hours.

##### What the important nonmedicinal ingredients are:

The capsules also contain the following inactive ingredients: Ethylcellulose, FD&C Blue #2, gelatin, hypromellose, iron oxide yellow (2 mg capsules only), starch, sucrose, talc, titanium dioxide; Ink: povidone, propylene glycol, shellac and sodium hydroxide.

##### What dosage forms it comes in:

MYLAN-TOLTERODINE ER 2 mg is a blue-green capsule printed with "MYLAN" over "TT 2" in white ink on both the cap and body.

MYLAN-TOLTERODINE ER 4 mg is a blue capsule printed with "MYLAN" over "TT 4" in white ink on both the cap and body.

#### WARNINGS AND PRECAUTIONS

MYLAN-TOLTERODINE ER may have an effect on the electrical activity of the heart. This effect can be measured as a change in the electrocardiogram (ECG). It is important to follow the instructions of your doctor with regard to dosing or any special tests. In very rare cases, drugs with an effect on the ECG can lead to disturbances in heart rhythm (arrhythmias/dysrhythmias). These heart rhythm disturbances are more likely in patients with risk factors, such as heart disease, or in the presence of certain interacting drugs. If you experience any symptoms of a possible heart rhythm disturbance, such as dizziness, palpitations (sensation of rapid pounding, or irregular heart beat), fainting, or seizures, you should stop taking MYLAN-TOLTERODINE ER and seek immediate medical attention.

##### **BEFORE you use MYLAN-TOLTERODINE ER talk to your doctor or pharmacist if:**

- you are pregnant, or trying to become pregnant
- you are breastfeeding your child
- you have myasthenia gravis (a chronic autoimmune neuromuscular disease which cause muscle weakness)
- you have stomach problems affecting passage and digestion of food
- you have liver problems
- you have kidney problems
- you are taking medication bought without a prescription. They may affect your condition, or how MYLAN-TOLTERODINE ER works for you.
- you are a female or are over 65 years in age; you have a disorder known as Long QT Syndrome; a heart disease; a history of stroke or brain hemorrhage; a personal history of fainting spells; a family history of sudden cardiac death at <50 years; electrolyte disturbances (e.g., low blood potassium levels); an eating disorder or are following an extreme diet; diabetes, especially with associated nerve disorders

#### INTERACTIONS WITH THIS MEDICATION

The following list includes some, but not all, of the drugs that may increase the risk of side effects while receiving MYLAN-TOLTERODINE ER. You should check with your doctor or pharmacist before taking any other medication with MYLAN-TOLTERODINE ER.

##### **Drugs that may interact with MYLAN-TOLTERODINE ER include:**

- other drugs that possess antimuscarinic/anticholinergic properties (drugs that cause blurred vision, constipation, dry

mouth, etc.)

- antifungals (drugs to treat fungal infections, such as, fluconazole, ketoconazole, or itraconazole)
- antibiotics (ie. erythromycin, clarithromycin)
- cyclosporine (a drug to prevent rejection of organ transplants)
- vinblastine (a drug to treat some types of cancer)
- antiarrhythmics (drugs that stabilize the heart rhythm function, such as procainamide, quinidine, amiodarone, sotalol, etc.)
- antidepressants (mood disorder drugs)
- antipsychotics (drugs to stabilize thinking and behavior)
- anti-asthmatic (salmeterol)

- palpitations (sensation of rapid, pounding, or irregular heart beat)
- fainting
- difficulty in urination (passing water)

The most common side effect is dry mouth. Less commonly reported side effects are headache, constipation, dizziness, fatigue, abdominal pain and dry eyes.

If you experience any symptoms of a possible heart rhythm disturbance, such as dizziness, palpitations (sensation of rapid, pounding, or irregular heart beat), fainting, or seizures, you should stop taking **MYLAN-TOLTERODINE ER** and seek immediate medical attention.

**PROPER USE OF THIS MEDICATION**

Take **MYLAN-TOLTERODINE ER** as instructed by your doctor. Do not increase, decrease or stop taking **MYLAN-TOLTERODINE ER** without first talking to your doctor.

**Usual dose:**

The usual starting dose is 4 mg once daily, but may be decreased to 2 mg once daily. The capsule should be swallowed whole. **MYLAN-TOLTERODINE ER** can be taken with food.

**Overdose:**

In case of drug overdose, contact a health care practitioner, hospital emergency department or regional Poison Control Centre immediately, even if there are no symptoms.

Do not take more capsules than your doctor has told you to. If you take too many tablets by accident, call your doctor or pharmacist or a poison control centre immediately.

**Missed Dose:**

If you miss taking your capsule, take it as soon as you remember. But if it is almost time for the next dose, skip the missed dose and just take the next dose. Do not take more than one dose at a time.

Check with your doctor or pharmacist right away if you have *any* bothersome or unusual effects while taking **MYLAN-TOLTERODINE ER**.

Use caution while driving or using machinery until you know how **MYLAN-TOLTERODINE ER** affects you.

**SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM**

Symptom / effect	Talk with your doctor or pharmacist		Stop taking drug and seek immediate emergency medical attention
	Only if severe	In all cases	
Uncommon Allergic reaction			√

*This is not a complete list of side effects. For any unexpected effects while taking **MYLAN-TOLTERODINE ER**, stop taking the drug and contact your doctor or pharmacist.*

Store at room temperature 15°C to 30°C. Protect from light.

You should not use your medication after the expiration date printed on the carton and label.

**Keep all medications out of the reach of children. This medication could harm them.**

**SIDE EFFECTS AND WHAT TO DO ABOUT THEM**

As with most drugs, **MYLAN-TOLTERODINE ER** can cause some side effects.

Tell your doctor or pharmacist right away if you suffer from any of the following side effects while taking this medication:

- dry mouth
- stomach pain
- decreased tear production (dry irritable eye)
- heartburn
- blurred vision
- dizziness

**REPORTING SUSPECTED SIDE EFFECTS**

You can help improve the safe use of health products for Canadians by reporting serious and unexpected side effects to Health Canada. Your report may help to identify new side effects and change the product safety information.

**3 ways to report:**

- Online at MedEffect (<http://hc-sc.gc.ca/dhp-mps/medeff/index-eng.php>);
- By calling 1-866-234-2345 (toll-free);
- By completing a Consumer Side Effect Reporting Form and sending it by:
  - Fax to 1-866-678-6789 (toll-free), or
  - Mail to: Canada Vigilance Program  
Health Canada, Postal Locator  
0701E  
Ottawa, ON  
K1A 0K9

Postage paid labels and the Consumer Side Effect Reporting Form are available at MedEffect. (<http://hc-sc.gc.ca/dhp-mps/medeff/index-eng.php>)

*NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.*

**MORE INFORMATION**

For more information, please contact your healthcare professionals or pharmacist first, or Mylan Pharmaceuticals ULC Canada at: 1-800-668-3174, or visit the website at [www.mylan.ca](http://www.mylan.ca)

This leaflet was prepared by Mylan Pharmaceuticals ULC, Etobicoke, Ontario, Canada, M8Z 2S6.

**Last prepared: April 28, 2015**