

PRODUCT MONOGRAPH

**Pr IPG-RIZATRIPTAN**

rizatriptan benzoate tablets

5 mg and 10 mg rizatriptan

5-HT<sub>1</sub> Receptor Agonist

**MIGRAINE THERAPY**

Marcan Pharmaceuticals Inc.  
4-77 Auriga Drive  
Ottawa, ON, K2E 7Z7  
[www.marcanpharma.com](http://www.marcanpharma.com)

Date of Preparation:  
January 8, 2016

Control # 190185

## Table of Contents

PART I: HEALTH PROFESSIONAL INFORMATION.....	3
SUMMARY PRODUCT INFORMATION .....	3
INDICATIONS AND CLINICAL USE .....	3
CONTRAINDICATIONS .....	3
WARNINGS AND PRECAUTIONS .....	4
ADVERSE REACTIONS.....	10
DRUG INTERACTIONS .....	15
DOSAGE AND ADMINISTRATION .....	17
OVERDOSAGE .....	18
ACTION AND CLINICAL PHARMACOLOGY.....	19
STORAGE AND STABILITY .....	21
DOSAGE FORMS, COMPOSITION AND PACKAGING .....	22
PART II: SCIENTIFIC INFORMATION .....	23
PHARMACEUTICAL INFORMATION.....	23
CLINICAL TRIALS .....	24
DETAILED PHARMACOLOGY .....	26
TOXICOLOGY .....	29
REFERENCES .....	31
PART III: CONSUMER INFORMATION .....	32

**Pr** **IPG-RIZATRIPTAN**  
as rizatriptan benzoate

**PART I: HEALTH PROFESSIONAL INFORMATION**

**SUMMARY PRODUCT INFORMATION**

<b>Route of Administration</b>	<b>Dosage Form / Strength</b>	<b>Nonmedicinal Ingredients</b>
Oral	tablets 5 mg , 10 mg	<i>Lactose monohydrate</i> <i>For a complete listing see Dosage Forms, Composition and Packaging section.</i>

**INDICATIONS AND CLINICAL USE**

**Adults**

IPG-RIZATRIPTAN (rizatriptan benzoate) is indicated for:

- acute treatment of migraine attacks with or without aura in adults

IPG-RIZATRIPTAN is not intended for the prophylactic therapy of migraine or for use in the management of hemiplegic, ophthalmoplegic or basilar migraine (see CONTRAINDICATIONS). Safety and effectiveness of IPG-RIZATRIPTAN have not been established for cluster headache, which is present in an older, predominantly male population.

**Pediatrics (<18 years of age)**

The safety and efficacy of IPG-RIZATRIPTAN has not been established in patients under 18 years of age and its use in this age group is not recommended (see WARNINGS AND PRECAUTIONS, Special Populations and Conditions).

**Geriatrics (> 65 years of age)**

The safety and effectiveness of IPG-RIZATRIPTAN has not been adequately studied in individuals over 65 years of age. Its use in this age group is, therefore, not recommended (see WARNINGS AND PRECAUTIONS).

**CONTRAINDICATIONS**

**IPG-RIZATRIPTAN is contraindicated in patients with history, symptoms, or signs of ischemic cardiac, cerebrovascular or peripheral vascular syndromes, valvular heart disease or cardiac arrhythmias (especially tachycardias). In addition, patients with other significant underlying cardiovascular diseases (e.g., atherosclerotic disease, congenital**

heart disease) should not receive IPG-RIZATRIPTAN. Ischemic cardiac syndromes include, but are not restricted to, angina pectoris of any type (e.g., stable angina of effort and vasospastic forms of angina such as the Prinzmetal's variant), all forms of myocardial infarction, and silent myocardial ischemia. Cerebrovascular syndromes include, but are not limited to, strokes of any type as well as transient ischemic attacks (TIAs). Peripheral vascular disease includes, but is not limited to, ischemic bowel disease or Raynaud's syndrome (see WARNINGS AND PRECAUTIONS).

Because IPG-RIZATRIPTAN may increase blood pressure, it is contraindicated in patients with uncontrolled or severe hypertension (see WARNINGS AND PRECAUTIONS).

IPG-RIZATRIPTAN is contraindicated within 24 hours of treatment with another 5-HT<sub>1</sub> agonist, or an ergotamine-containing or ergot-type medication like dihydroergotamine or methysergide.

IPG-RIZATRIPTAN is contraindicated in patients with hemiplegic, ophthalmoplegic or basilar migraine.

Concurrent administration of MAO inhibitors or use of rizatriptan within 2 weeks of discontinuation of MAO inhibitor therapy is contraindicated (see DRUG INTERACTIONS).

Because there are no data available, IPG-RIZATRIPTAN is contraindicated in patients with severe hepatic impairment.

IPG-RIZATRIPTAN is contraindicated in patients who are hypersensitive to rizatriptan or any component of the formulation.

## **WARNINGS AND PRECAUTIONS**

### **General**

IPG-RIZATRIPTAN should only be used where a clear diagnosis of migraine has been established.

For a given attack, if a patient has no response to the first dose of rizatriptan, the diagnosis of migraine should be reconsidered before administration of a second dose.

### **Psychomotor Effect**

Dizziness, somnolence and asthenia/fatigue were experienced by some patients in clinical trials with IPG-RIZATRIPTAN (see ADVERSE EVENTS). Patients should be advised to avoid driving a car or operating hazardous machinery until they are reasonably certain that IPG-RIZATRIPTAN does not adversely affect them.

## Cardiovascular

**Risk of Myocardial Ischemia and/or Infarction and Other Adverse Cardiac Events** IPG-RIZATRIPTAN has been associated with transient chest and/or neck pain and tightness which may resemble angina pectoris. Following the use of other 5-HT<sub>1</sub> agonists, in rare cases these symptoms have been identified as being the likely result of coronary vasospasm or myocardial ischemia. Rare cases of serious coronary events or arrhythmia have occurred following use of other 5-HT<sub>1</sub> agonists, and may therefore also occur with IPG-RIZATRIPTAN. Because of the potential of this class of compounds (5-HT<sub>1B/1D</sub> agonists) to cause coronary vasospasm, IPG-RIZATRIPTAN should not be given to patients with documented ischemic or vasospastic coronary artery disease (see CONTRAINDICATIONS). It is strongly recommended that IPG-RIZATRIPTAN not be given to patients in whom unrecognized coronary artery disease (CAD) is predicted by the presence of risk factors (e.g., hypertension, hypercholesterolemia, smoker, obesity, diabetes, strong family history of CAD, female with surgical or physiological menopause, or male over 40 years of age) unless a cardiovascular evaluation provides satisfactory clinical evidence that the patient is reasonably free of coronary artery and ischemic myocardial disease or other significant underlying cardiovascular disease. The sensitivity of cardiac diagnostic procedures to detect cardiovascular disease or predisposition to coronary artery vasospasm is unknown. If, during the cardiovascular evaluation, the patient's medical history, electrocardiographic or other investigations reveal findings indicative of, or consistent with, coronary artery vasospasm or myocardial ischemia, IPG-RIZATRIPTAN should not be administered (see CONTRAINDICATIONS).

For patients with risk factors predictive of CAD, who are considered to have a satisfactory cardiovascular evaluation, the first dose of rizatriptan should be administered in the setting of a physician's office or similar medically staffed and equipped facility. Because cardiac ischemia can occur in the absence of clinical symptoms, consideration should be given to obtaining on the first occasion of use an electrocardiogram (ECG) during the interval immediately following IPG-RIZATRIPTAN, in these patients with risk factors. However, an absence of drug-induced cardiovascular effects on the occasion of the initial dose does not preclude the possibility of such effects occurring with subsequent administrations.

Intermittent long-term users of IPG-RIZATRIPTAN who have or acquire risk factors predictive of CAD, as described above, should receive periodic interval cardiovascular evaluation as they continue to use IPG-RIZATRIPTAN.

If symptoms consistent with angina occur after the use of IPG-RIZATRIPTAN, ECG evaluation should be carried out to look for ischemic changes.

*The systematic approach described above is intended to reduce the likelihood that patients with unrecognized cardiovascular disease will be inadvertently exposed to IPG-RIZATRIPTAN.*

Discomfort in the chest, neck, throat and jaw (including pain, pressure, heaviness and

tightness) has been reported after administration of rizatriptan. Because drugs in this class may cause coronary artery vasospasm, patients who experience signs or symptoms suggestive of angina following dosing should be evaluated for the presence of CAD or a predisposition to Prinzmetal's variant angina before receiving additional doses of medication, and should be monitored electrocardiographically if dosing is resumed and similar symptoms recur. Similarly, patients who experience other symptoms or signs suggestive of decreased arterial flow, such as ischemic bowel syndrome or Raynaud's syndrome following IPG-RIZATRIPTAN administration should be evaluated for atherosclerosis or predisposition to vasospasm (see CONTRAINDICATIONS and WARNINGS AND PRECAUTIONS).

### **Cardiac Events and Fatalities Associated with 5-HT<sub>1</sub> Agonists**

IPG-RIZATRIPTAN may cause coronary artery vasospasm. Serious adverse cardiac events, including acute myocardial infarction, life-threatening disturbances of cardiac rhythm, and death have been reported within a few hours following the administration of 5-HT<sub>1</sub> agonists. Considering the extent of use of 5-HT<sub>1</sub> agonists in patients with migraine, the incidence of these events is extremely low.

#### ***Premarketing Experience with Rizatriptan***

Among the approximately 4200 patients who were treated with at least a single oral dose of either 5 or 10 mg rizatriptan in premarketing clinical trials of rizatriptan, electrocardiac adverse experiences were observed in 33 patients. One patient was reported to have chest pain with possible ischemic ECG changes following a single dose of 10 mg.

#### ***Post marketing Experience with Rizatriptan***

Serious cardiovascular events have been reported in association with the use of rizatriptan. The uncontrolled nature of post marketing surveillance, however, makes it impossible to determine definitively the proportion of reported cases that were actually caused by rizatriptan or to reliably assess causation in individual cases.

### **Cerebrovascular Events and Fatalities Associated with 5-HT<sub>1</sub> Agonists**

Cerebral hemorrhage, subarachnoid hemorrhage, stroke, and other cerebrovascular events have been reported in patients treated with 5-HT<sub>1</sub> agonists; and some have resulted in fatalities. In a number of cases, it appears possible that the cerebrovascular events were primary, the agonist having been administered in the incorrect belief that the symptoms experienced were a consequence of migraine, when they were not. Before treating migraine headaches with rizatriptan in patients not previously diagnosed as migraineurs, and in migraineurs who present with atypical symptoms, care should be taken to exclude other potentially serious neurological conditions. If a patient does not respond to the first dose, the opportunity should be taken to review the diagnosis before a second dose is given. It should be noted that patients with migraine may be at increased risk of certain cerebrovascular events (e.g., stroke, hemorrhage, transient ischemic attack).

## **Special Cardiovascular Pharmacology Studies with another 5-HT<sub>1</sub> Agonist**

In subjects (n=10) with suspected coronary artery disease undergoing angiography, a 5-HT<sub>1</sub> agonist at a subcutaneous dose of 1.5 mg produced an 8% increase in aortic blood pressure, an 18% increase in pulmonary artery blood pressure, and an 8% increase in systemic vascular resistance. In addition, mild chest pain or tightness was reported by four subjects. Clinically significant increases in blood pressure were experienced by three of the subjects (two of whom also had chest pain/discomfort). Diagnostic angiogram results revealed that 9 subjects had normal coronary arteries and one had insignificant coronary artery disease.

In an additional study with this same drug, migraine patients (n=35) free of cardiovascular disease were subjected to assessments of myocardial perfusion by positron emission tomography while receiving a subcutaneous 1.5 mg dose in the absence of a migraine attack. Reduced coronary vasodilatory reserve (~10%), increased coronary resistance (~20%), and decreased hyperemic myocardial blood flow (~10%) were noted. The relevance of these findings to the use of the recommended oral dose of this 5-HT<sub>1</sub> agonist is not known.

Similar studies have not been done with rizatriptan. However, owing to the common pharmacodynamic actions of 5-HT<sub>1</sub> agonists, the possibility of cardiovascular effects of the nature described above should be considered for any agent of this pharmacological class.

## **Other Vasospasm-Related Events**

5-HT<sub>1</sub> agonists may cause vasospastic reactions other than coronary artery vasospasm. Extensive post-market experience has shown the use of another 5-HT<sub>1</sub> agonist to be associated with rare occurrences of peripheral vascular ischemia and colonic ischemia with abdominal pain and bloody diarrhea.

## **Increase in Blood Pressure**

Significant elevation in blood pressure, including hypertensive crisis, has been reported on rare occasions in patients receiving 5-HT<sub>1</sub> agonists with and without a history of hypertension. In healthy young male and female subjects who received maximal doses of rizatriptan (10 mg every 2 hours for 3 doses), slight increases in blood pressure (approximately 2-3 mmHg) were observed. Rizatriptan is contraindicated in patients with uncontrolled or severe hypertension (see CONTRAINDICATIONS). In patients with controlled hypertension, rizatriptan should be administered with caution, as transient increases in blood pressure and peripheral vascular resistance have been observed in a small portion of patients.

### **Hepatic/Biliary/Pancreatic**

Rizatriptan should be used with caution in patients with moderate hepatic insufficiency due to an increase in plasma concentrations of approximately 30% (see ACTION AND CLINICAL PHARMACOLOGY, Special Populations and Conditions, and DOSAGE AND ADMINISTRATION). Since there are no data in patients with severe hepatic impairment, rizatriptan is contraindicated in this population (see CONTRAINDICATIONS and DOSAGE AND ADMINISTRATION).

### **Immune**

Rare hypersensitivity (anaphylaxis/anaphylactoid) reactions may occur in patients receiving 5-HT<sub>1</sub> agonists such as IPG-RIZATRIPTAN. Such reactions can be life threatening or fatal. In general, hypersensitivity reactions to drugs are more likely to occur in individuals with a history of sensitivity to multiple allergens. Owing to the possibility of cross-reactive hypersensitivity reactions, IPG-RIZATRIPTAN should not be used in patients having a history of hypersensitivity to chemically-related 5-HT<sub>1</sub> receptor agonists.

### **Neurologic**

Care should be taken to exclude other potentially serious neurologic conditions before treating headache in patients not previously diagnosed with migraine or who experience a headache that is atypical for them. There have been rare reports where patients received 5-HT<sub>1</sub> agonists for severe headache that were subsequently shown to have been secondary to an evolving neurological lesion. For newly diagnosed patients or patients presenting with atypical symptoms, the diagnosis of migraine should be reconsidered if no response is seen after the first dose of IPG-RIZATRIPTAN.

### **Seizures**

Caution should be observed if IPG-RIZATRIPTAN is to be used in patients with a history of epilepsy or structural brain lesions which lower the convulsion threshold. There have been very rare reports of seizures following administration of IPG-RIZATRIPTAN in patients with or without risk factors or previous history of seizures (see ADVERSE REACTIONS, Post-Marketing Adverse Reactions, Nervous System).

### **Ophthalmologic**

#### **Binding to Melanin-Containing Tissues**

The propensity for rizatriptan to bind melanin has not been investigated. Based on its chemical properties, rizatriptan may bind to melanin and accumulate in melanin-rich tissue (e.g., eye) over time. This raises the possibility that rizatriptan could cause toxicity in these tissues after extended use. There were, however, no adverse ophthalmologic changes related to treatment with rizatriptan in the one-year dog toxicity study. Although no systematic monitoring of ophthalmologic function was undertaken in clinical trials, and no specific recommendations for ophthalmologic monitoring are offered, prescribers should be aware of the possibility of long-term ophthalmologic effects.



## **Renal**

Rizatriptan should be used with caution in dialysis patients due to a decrease in the clearance of rizatriptan, resulting in approximately 44% increase in plasma concentrations (see ACTION AND CLINICAL PHARMACOLOGY, Special Populations and Conditions, and DOSAGE AND ADMINISTRATION).

## **Selective Serotonin Reuptake Inhibitors/Serotonin Norepinephrine Reuptake Inhibitors and Serotonin Syndrome**

Cases of life-threatening serotonin syndrome have been reported during combined use of selective serotonin reuptake inhibitors (SSRIs)/serotonin norepinephrine reuptake inhibitors (SNRIs) and triptans. If concomitant treatment with IPG-RIZATRIPTAN and SSRIs (e.g., sertraline, escitalopram oxalate, and fluoxetine) or SNRIs (e.g., venlafaxine, duloxetine) is clinically warranted, careful observation of the patient is advised, particularly during treatment initiation and dose increases. Serotonin syndrome symptoms may include mental status changes (e.g., agitation, hallucinations, coma), autonomic instability (e.g., tachycardia, labile blood pressure, hyperthermia), neuromuscular aberrations (e.g., hyperreflexia, incoordination) and/or gastrointestinal symptoms (e.g., nausea, vomiting, diarrhea) (see DRUG INTERACTIONS).

**Medication Overuse Headache:** Overuse of acute migraine treatments has been associated with the exacerbation of headache (medication overuse headache) in susceptible patients. Medication overuse headache may present as migraine-like daily headaches or as a marked increase in frequency of migraine attacks. Detoxification of patients, including withdrawal of the overused drugs, and treatment of withdrawal symptoms (which often includes a transient worsening of headache) may be necessary.

## **Special Populations and Conditions**

**Pregnant Women:** In a reproduction study in rats, birth weights and pre- and post-weaning weight gain were reduced in the offspring of females treated prior to and during mating and throughout gestation and lactation. These effects occurred in the absence of any apparent maternal toxicity (maternal plasma drug exposures were 22 and 337 times, respectively, the exposure in humans receiving the maximum recommended daily dose (MRDD) of 20 mg). The developmental no-effect dose was equivalent to 2.25 times human exposure at the MRDD.

In embryofetal development studies, no teratogenic effects were observed when pregnant rats and rabbits were administered doses at the equivalent of 337 times and 168 times, respectively, the human MRDD, during organogenesis. However, fetal weights were decreased in conjunction with decreased maternal weight gain at these same doses. The developmental no-effect dose in both rats and rabbits was 22 times the human MRDD. Toxicokinetic studies demonstrated placental transfer of drug in both species.

There are no adequate and well-controlled studies in pregnant women; therefore, rizatriptan should be used during pregnancy only if the potential benefit justifies the potential risk to the

fetus.

### **Impairment of Fertility**

In a fertility study in rats, altered estrus cyclicity and delays in time to mating were observed in females treated orally with an equivalent of 337 times the maximum recommended daily dose (MRDD) of 20 mg in humans. The no-effect dose was 22 times the MRDD. There was no impairment of fertility or reproductive performance in male rats treated with up to 825 times the MRDD.

**Nursing Women:** It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when rizatriptan is administered to women who are breast-feeding. Rizatriptan is extensively excreted in rat milk, at a level of 5-fold or greater than maternal plasma levels.

**Pediatrics (<18 years of age):** The safety and efficacy of Rizatriptan have not been established in patients under 18 years of age and its use in this age group is not recommended (see ADVERSE REACTIONS, Special Populations).

**Geriatrics (> 65 years of age):** The safety and effectiveness of rizatriptan has not been adequately studied in individuals over 65 years of age. The risk of adverse reactions to this drug may be greater in elderly patients, as they are more likely to have decreased hepatic function, be at higher risk for CAD, and experience blood pressure increases that may be more pronounced. Clinical studies with rizatriptan did not include a substantial number of patients over 65 years of age (n=17). Its use in this age group is, therefore, not recommended.

### **Special Disease Conditions:**

IPG-RIZATRIPTAN (rizatriptan benzoate) should be administered with caution to patients with diseases that may alter the absorption, metabolism, or excretion of drugs (see ACTION AND CLINICAL PHARMACOLOGY, Special Populations and Conditions).

### **Monitoring and Laboratory Tests**

No specific laboratory tests are recommended for monitoring patients prior to and/or after treatment with rizatriptan.

## **ADVERSE REACTIONS**

### **Adverse Drug Reaction Overview**

**Serious cardiac events, including some that have been fatal, have occurred following use of 5-HT<sub>1</sub> agonists. These events are extremely rare and most have been reported in patients with risk factors predictive of CAD. Events reported have included coronary artery vasospasm, transient myocardial ischemia, myocardial infarction, ventricular tachycardia,**

**and ventricular fibrillation (see CONTRAINDICATIONS, WARNINGS AND PRECAUTIONS).**

### **Clinical Trial Adverse Drug Reactions**

*Because clinical trials are conducted under very specific conditions the adverse reaction rates observed in the clinical trials may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse drug reaction information from clinical trials is useful for identifying drug-related adverse events and for approximating rates.*

### **Experience in Controlled Clinical Trials with Rizatriptan**

#### **Typical 5-HT<sub>1</sub> Agonist Adverse Reactions**

As with other 5-HT<sub>1</sub> agonists, rizatriptan has been associated with sensations of heaviness, pressure, tightness or pain which may be intense. These may occur in any part of the body including the chest, throat, neck, jaw and upper limb.

#### **Acute Safety**

Adverse experiences to rizatriptan were assessed in controlled clinical trials that included over 3700 adult patients who received single or multiple doses of rizatriptan tablets. The most common adverse events during treatment with rizatriptan were asthenia/fatigue, somnolence, pain/pressure sensation and dizziness. These events appeared to be dose-related. In long-term extension studies where patients were allowed to treat multiple attacks for up to 1 year, 4% (59 out of 1525 patients) withdrew because of adverse experiences.

Table 1 list the adverse events regardless of drug relationship (incidence  $\geq$  1% and greater than placebo) after a single dose of rizatriptan tablets. Most of the adverse events appear to be dose-related. The events cited reflect experience gained under closely monitored conditions of clinical trials in a highly selected patient population. In actual clinical practice or in other clinical trials, these frequency estimates may not apply, as the conditions of use, reporting behavior, and the kinds of patients treated may differ.

### **Table 1 Incidence ( $\geq$ 1% and Greater than Placebo) of Adverse Experiences in Adults After a Single Dose of rizatriptan tablets or Placebo (Prior to Subsequent Dose) in Phase III Controlled Clinical Trials<sup>†</sup>**

	% of Patients		
	Placebo	Rizatriptan 5 mg	Rizatriptan 10 mg
<b>Number of Patients</b>	627	977	1167
<b>Symptoms of Potentially Cardiac Origin</b>			
Upper Limb Sensations*	1.3	1.7	1.8
Chest Sensations*	1.0	1.6	3.1
Neck/Throat/Jaw Sensations*	0.6	1.4	2.5
Palpitations	0.2	0.9	1.0
<b>Body as a Whole</b>			
Asthenia/Fatigue	2.1	4.2	6.9
Abdominal Pain	1.0	1.7	2.2
<b>Digestive System</b>			
Nausea	3.5	4.1	5.7
Dry Mouth	1.3	2.6	3.0
Vomiting	2.1	1.6	2.3
<b>Nervous System</b>			
Dizziness	4.5	4.2	8.9
Somnolence	3.5	4.2	8.4
Headache	0.8	1.8	2.1
Paresthesia	1.0	1.5	2.9
Tremor	1.0	1.3	0.3
Insomnia	0.3	1.0	0.3
<b>Skin and Skin Appendage</b>			
Flushing	1.0	0.6	1.1

\* The term “sensations” encompasses adverse events described as pain, discomfort, pressure, heaviness, constriction, tightness, heat/burning sensation, paresthesia, numbness, tingling, weakness and strange sensations.

† Data from Studies 022, 025, 029 and 030.

Rizatriptan was generally well-tolerated. Adverse experiences were typically mild in intensity and were transient. The frequencies of adverse experiences in clinical trials did not increase when up to three doses were taken within 24 hours. The incidences of adverse experiences were not affected by age, gender or use of prophylactic medications. There were insufficient data to assess the impact of race on the incidence of adverse events.

### **Long-Term Safety**

In long-term extension studies, a total of 1854 patients treated 16,150 migraine attacks with rizatriptan 5 mg Tablets and 24,043 attacks with rizatriptan 10 mg Tablets over a period of up to 1 year. In general, the types of clinical adverse experiences observed in the extension studies were similar to those observed in the acute studies. However, the incidences of most clinical adverse events were approximately 3-fold higher in extension, as expected, based on increased observation time. The most common adverse events per attack (defined as occurring at an incidence of at least 1%) for rizatriptan 5 mg and 10 mg, respectively, were as follows: nausea (3%, 4%), dizziness (2%, 2%), somnolence (2%, 4%), asthenia/fatigue (2%, 2%), headache (1%, 2%), vomiting (1%, <1%), chest pain (<1%, 1%) and paresthesia (<1%, 2%). Due to the lack of placebo controls in the extension studies, the role of rizatriptan in causation cannot be reliably determined.

### **Other Events Observed in Association with the Administration of Rizatriptan**

In the section that follows, the frequencies of less commonly reported adverse clinical events are presented. Because the reports include events observed in open studies, the role of rizatriptan in their causation cannot be reliably determined. Furthermore, variability associated with adverse event reporting, the terminology used to describe adverse events, etc. limit the value of the quantitative frequency estimates provided. Event frequencies are calculated as the number of patients who used rizatriptan 5 mg and 10 mg tablets in Phase II and III studies (n=3716) and reported an event divided by the total number of patients exposed to rizatriptan. All reported events are included, except those already listed in the previous table, those too general to be informative, and those not reasonably associated with the use of the drug. Events are further classified within body system categories and enumerated in order of decreasing frequency using the following definitions: frequent adverse events are those defined as those occurring in at least 1/100 patients; infrequent adverse experiences are those occurring in 1/100 to 1/1000 patients; and rare adverse experiences are those occurring in fewer than 1/1000 patients.

### **Body as a Whole**

Frequent were warm sensations, chest pain and chills/cold sensations. Infrequent were heat sensitivity, facial edema, hangover effect, abdominal distention, edema/swelling and malaise. Rare were fever, orthostatic effects, and syncope.

### **Cardiovascular**

Frequent was palpitation. Infrequent were tachycardia, cold extremities, hypertension, arrhythmia, and bradycardia. Rare were angina pectoris and blood pressure increased.

**Digestive**

Frequent was diarrhea. Infrequent were dyspepsia, thirst, acid regurgitation, dysphagia, constipation, flatulence, and tongue edema. Rare were anorexia, appetite increase, gastritis, paralysis (tongue), eructation and glossodynia.

**Metabolic**

Infrequent was dehydration.

**Musculoskeletal**

Infrequent were muscle weakness, stiffness, myalgia, muscle cramp, musculoskeletal pain, and arthralgia.

**Neurological/Psychiatric**

Frequent were hypoesthesia and mental acuity decreased. Infrequent were nervousness, vertigo, insomnia, anxiety, depression, euphoria, disorientation, ataxia, dysarthria, confusion, dream abnormality, gait abnormality, irritability, memory impairment, agitation, hyperesthesia, sleep disorder, speech disorder, migraine and spasm. Rare were dysesthesia, depersonalization, akinesia/bradykinesia, apprehension, hyperkinesia, hypersomnia, and hyporeflexia.

**Respiratory**

Frequent were dyspnea and pharyngeal discomfort. Infrequent were pharyngitis, irritation (nasal), congestion (nasal), dry throat, upper respiratory infection, yawning, respiratory congestion, dry nose, epistaxis, and sinus disorder. Rare were cough, hiccups, hoarseness, rhinorrhea, sneezing, tachypnea, and pharyngeal edema.

**Special Senses**

Frequent was taste perversion. Infrequent were blurred vision, tinnitus, dry eyes, burning eye, eye pain, eye irritation, ear pain, and tearing. Rare were hyperacusis, smell perversion, photophobia, photopsia, itching eye, and eye swelling.

**Skin and Skin Appendage**

Infrequent were sweating, pruritus, rash, and urticaria. Rare were erythema, acne, and photosensitivity.

**Urogenital System**

Frequent was hot flashes. Infrequent were urinary frequency, polyuria, and menstruation disorder. Rare was dysuria.

**Post-Market Adverse Drug Reactions**

The following additional adverse reactions have been reported very rarely and most have been reported in patients with risk factors predictive of CAD: Myocardial ischemia or infarction, cerebrovascular accident.

The following adverse reactions have also been reported:

**Hypersensitivity:** Hypersensitivity reaction, anaphylaxis/anaphylactoid reaction, angioedema (e.g., facial edema, tongue swelling, pharyngeal edema), wheezing, urticaria, rash, toxic epidermal necrolysis.

**Nervous System:** serotonin syndrome.

**Seizures:** There have been very rare reports of seizures following administration of rizatriptan in patients with or without risk factors or previous history of seizures (see WARNINGS AND PRECAUTIONS).

**Musculoskeletal:** facial pain.

**Special Senses:** Dysgeusia.

**Vascular disorders:** Peripheral vascular ischemia

**Gastrointestinal Disorders:** Ischemic colitis.

### **Drug Abuse and Dependence**

Although the abuse potential of rizatriptan has not been specifically assessed, no abuse of, tolerance to, withdrawal from, or drug-seeking behavior was observed in patients who received rizatriptan in clinical trials or their extensions. The 5-HT<sub>1B/1D</sub> agonists, as a class, have not been associated with drug abuse.

## **DRUG INTERACTIONS**

### **Drug-Drug Interactions**

#### **Ergot-Containing Drugs**

Ergot-containing drugs have been reported to cause prolonged vasospastic reactions. Because there is a theoretical basis that these effects may be additive, use of ergotamine-containing or ergot-type medications (like dihydroergotamine or methysergide) and rizatriptan within 24 hours is contraindicated (see CONTRAINDICATIONS).

#### **Monoamine Oxidase Inhibitors**

Rizatriptan is principally metabolized via monoamine oxidase, 'A' subtype (MAO-A). In a drug interaction study, when rizatriptan 10 mg was administered to subjects (n=12) receiving concomitant therapy with the selective, reversible MAO-A inhibitor, moclobemide 150 mg t.i.d., there were mean increases in rizatriptan AUC and C<sub>max</sub> of 119% and 41%, respectively; and the AUC of the active N-monodesmethyl metabolite of rizatriptan was increased more than 400%. The interaction would be expected to be greater with irreversible MAO inhibitors. Drug interaction studies were not conducted with selective MAO-B inhibitors. The specificity of MAO-B inhibitors diminishes with higher doses and varies among patients. Therefore, co-administration of rizatriptan in patients taking MAO-A or MAO-B inhibitors is contraindicated

(see CONTRAINDICATIONS).

### **Nadolol/Metoprolol**

In a drug interactions study, effects of multiple doses of nadolol 80 mg or metoprolol 100 mg every 12 hours on the pharmacokinetics of a single dose of 10 mg rizatriptan were evaluated in healthy subjects (n=12). No pharmacokinetic interactions were observed.

### **Oral Contraceptives**

In a study of concurrent administration of an oral contraceptive during 6 days of administration of rizatriptan (10-30 mg/day) in healthy female volunteers (n=18), rizatriptan did not affect plasma concentrations of ethinyl estradiol or norethindrone.

### **Other 5-HT<sub>1</sub> Agonists**

The administration of rizatriptan with other 5-HT<sub>1</sub> agonists has not been evaluated in migraine patients. Because their vasospastic effects may be additive, co-administration of rizatriptan and other 5-HT<sub>1</sub> agonists within 24 hours of each other is contraindicated (see CONTRAINDICATIONS).

### **Propranolol**

IPG-RIZATRIPTAN should be used with caution in patients receiving propranolol, since the pharmacokinetic behavior of rizatriptan during co-administration with propranolol may be unpredictable. In a study of concurrent administration of propranolol 240 mg/day and a single dose of rizatriptan 10 mg in healthy adult subjects (n=11), mean plasma AUC and C<sub>max</sub> for rizatriptan were increased by 70% and 75%, respectively, during propranolol administration. In one subject, a 4-fold increase in AUC and 5-fold increase in C<sub>max</sub> was observed. This subject was not distinguishable from the others based on demographic characteristics. The AUC of the active N-monodesmethyl metabolite of rizatriptan was not affected by propranolol. (See DOSAGE AND ADMINISTRATION).

### **Selective Serotonin Reuptake Inhibitors / Serotonin Norepinephrine Reuptake Inhibitors and Serotonin Syndrome**

Cases of life-threatening serotonin syndrome have been reported in post-marketing experience during combined use of selective serotonin reuptake inhibitors (SSRIs) or serotonin norepinephrine reuptake inhibitors (SNRIs) and triptans. (See WARNINGS AND PRECAUTIONS.)

In a pharmacokinetic study with paroxetine and rizatriptan, paroxetine had no influence on the plasma levels of rizatriptan.

### **Drug-Food Interactions**

Interactions with food have not been studied. Food has no significant effect on the bioavailability of rizatriptan but delays the time to reach peak concentration by an hour. In clinical trials, rizatriptan was administered without regard to food.



### **Drug-Herb Interactions**

Interactions with herbal products have not been studied.

### **Drug-Laboratory Interactions**

Rizatriptan is not known to interfere with commonly employed clinical laboratory tests.

### **Drug-Lifestyle Interactions**

Lifestyle interactions have not been established.

## **DOSAGE AND ADMINISTRATION**

### **Dosing Considerations**

IPG-RIZATRIPTAN is recommended only for the acute treatment of migraine attacks. IPG-RIZATRIPTAN should not be used prophylactically.

Controlled trials have not established the effectiveness of a second dose if the initial dose is ineffective.

The safety of treating, on average, more than four headaches in a 30-day period has not been established.

### **Recommended Dose and Dosage Adjustment**

#### **ADULTS**

##### **IPG-RIZATRIPTAN Tablets**

The recommended single adult dose is 5 mg. The maximum recommended single dose is 10 mg. There is evidence that the 10 mg dose may provide a greater effect than the 5 mg dose (see ACTION AND CLINICAL PHARMACOLOGY, Clinical Studies). The choice of dose should therefore be made on an individual basis, weighing the possible benefit of the 10 mg dose with the potential risk for increased adverse events.

##### **Redosing**

Doses should be separated by at least 2 hours; no more than a total of 20 mg (Tablets) should be taken in any 24-hour period.

##### **Patients receiving propranolol**

A single 5 mg dose of IPG-RIZATRIPTAN should be used. In no instances should the total daily dose exceed 10 mg per day, given in two doses, separated by at least two hours (see DRUG INTERACTIONS).

##### **Renal Impairment**

In hemodialysis patients with severe renal impairment (creatinine clearance  $<2$  mL/min/1.73 m<sup>2</sup>), the AUC of rizatriptan was approximately 44% greater than in patients with normal renal

function (see ACTION AND CLINICAL PHARMACOLOGY, Special Populations and Conditions). Consequently, if treatment is deemed advisable in these patients, the 5 mg IPG-RIZATRIPTAN tablet should be administered. No more than a total of 10 mg should be taken in any 24-hour period. Repeated dosing in renally impaired patients has not been evaluated.

### **Hepatic Impairment**

IPG-RIZATRIPTAN is contraindicated in patients with severe hepatic impairment (Child-Pugh grade C) due to the absence of safety data. Plasma concentrations of rizatriptan were approximately 30% greater in patients with moderate hepatic insufficiency (see ACTION AND CLINICAL PHARMACOLOGY, Special Populations and Conditions). Consequently, if treatment is deemed advisable in the presence of moderate hepatic impairment, the 5 mg IPG-RIZATRIPTAN tablet should be administered. No more than a total of 10 mg should be taken in any 24-hour period. Repeated dosing in hepatically impaired patients has not been evaluated.

### **Patients with Hypertension**

IPG-RIZATRIPTAN should not be used in patients with uncontrolled or severe hypertension. In patients with mild to moderate controlled hypertension, patients should be treated cautiously at the lowest effective dose.

## **OVERDOSAGE**

For management of a suspected drug overdose, contact your regional Poison Control Centre immediately.

No overdoses of rizatriptan (rizatriptan benzoate) were reported during clinical trials in adults.

Rizatriptan 40 mg (administered as either a single dose or as two doses with a 2-hour interdose interval) was generally well tolerated in over 300 adult patients; dizziness and somnolence were the most common drug-related adverse effects.

In a clinical pharmacology study in which 12 adult subjects received rizatriptan, at total cumulative doses of 80 mg (given within four hours), two subjects experienced syncope and/or bradycardia. One subject, a female aged 29 years, developed vomiting, bradycardia, and dizziness beginning three hours after receiving a total of 80 mg rizatriptan (administered over two hours); a third degree AV block, responsive to atropine, was observed an hour after the onset of the other symptoms. The second subject, a 25-year-old male, experienced transient dizziness, syncope, incontinence, and a 5-second systolic pause (on ECG monitor) immediately after a painful venipuncture. The venipuncture occurred two hours after the subject had received a total of 80 mg rizatriptan (administered over four hours).

In addition, based on the pharmacology of rizatriptan, hypertension or other more serious cardiovascular symptoms could occur after overdose. Gastrointestinal decontamination (i.e., gastric lavage followed by activated charcoal) should be considered in patients suspected of an overdose with rizatriptan. The elimination half-life of rizatriptan is 2 to 3 hours (see ACTION

AND CLINICAL PHARMACOLOGY). Clinical and electrocardiographic monitoring should be continued for at least 12 hours, even if clinical symptoms are not observed.

There is no specific antidote to rizatriptan. In cases of severe intoxication, intensive care procedures are recommended, including establishing and maintaining a patent airway, ensuring adequate oxygenation and ventilation, and monitoring and support of the cardiovascular system.

The effects of hemo- or peritoneal dialysis on serum concentrations of rizatriptan are unknown.

## **ACTION AND CLINICAL PHARMACOLOGY**

### **Mechanism of Action**

Rizatriptan is a selective 5-hydroxytryptamine<sub>1B/1D</sub> (5-HT<sub>1B/1D</sub>) receptor agonist. Rizatriptan binds with high affinity to human cloned 5-HT<sub>1B</sub> and 5-HT<sub>1D</sub> receptors. It has weak affinity for other 5-HT<sub>1</sub> receptor subtypes (5-HT<sub>1A</sub>, 5-HT<sub>1E</sub>, 5-HT<sub>1F</sub>) and the 5-HT<sub>7</sub> receptor, but has no significant activity at 5-HT<sub>2</sub>, 5-HT<sub>3</sub>, alpha- and beta-adrenergic, dopaminergic, histaminergic, muscarinic or benzodiazepine receptors.

Current theories on the etiology of migraine headache suggest that symptoms are due to local cranial vasodilatation and/or to the release of vasoactive and pro-inflammatory peptides from sensory nerve endings in an activated trigeminal system. The therapeutic activity of rizatriptan in migraine can most likely be attributed to agonist effects at 5-HT<sub>1B/1D</sub> receptors on the extracerebral, intracranial blood vessels that become dilated during a migraine attack and on nerve terminals in the trigeminal system. Activation of these receptors results in cranial vessel constriction, inhibition of neuropeptide release and reduced transmission in trigeminal pain pathways.

### **Pharmacokinetics**

**Absorption:** Rizatriptan is completely absorbed following oral administration. The mean oral absolute bioavailability of the rizatriptan tablet is about 45%, and mean peak plasma concentrations ( $C_{max}$ ) are reached in approximately 1-1.5 hours ( $T_{max}$ ). The presence of a migraine headache did not appear to affect the absorption or pharmacokinetics of rizatriptan. Food has no significant effect on the bioavailability of rizatriptan but delays the time to reach peak concentration by an hour. In clinical trials, rizatriptan was administered without regard to food. The plasma half-life of rizatriptan in males and females averages 2-3 hours.

When rizatriptan 10 mg was administered every 2 hours for three doses on four consecutive days, the plasma concentrations of rizatriptan within each day were approximately 3-fold greater than those seen with a single 10 mg dose and no plasma accumulation of the drug occurred from day to day.

In a pharmacokinetic study in adults, median  $T_{max}$  was 0.67 hours for the 10-mg tablet. AUC of rizatriptan is approximately 30% higher in females than in males.

**Distribution:** The mean volume of distribution is approximately 140 liters in male subjects and 110 liters in female subjects. Rizatriptan is minimally bound (14%) to plasma proteins.

**Metabolism:** The primary route of rizatriptan metabolism is via oxidative deamination by monoamine oxidase-A (MAO-A) to the indole acetic acid metabolite, which is not active at the 5-HT<sub>1B/1D</sub> receptor. N-monodesmethyl-rizatriptan, a metabolite with activity similar to that of parent compound at the 5-HT<sub>1B/1D</sub> receptor, is formed to a minor degree. Plasma concentrations of N-monodesmethyl-rizatriptan are approximately 14% of those of parent compound, and it is eliminated at a similar rate. Other minor metabolites, the N-oxide, the 6-hydroxy compound, and the sulfate conjugate of the 6-hydroxy metabolite are not active at the 5-HT<sub>1B/1D</sub> receptor.

Rizatriptan is not an inhibitor of the activities of human liver cytochrome P450 isoforms 3A4/5, 1A2, 2C9, 2C19, or 2E1; rizatriptan is a competitive inhibitor (K<sub>i</sub>=1400 nM) of cytochrome P450 2D6, but only at high, clinically irrelevant concentrations.

**Excretion:** The total radioactivity of the administered dose recovered over 120 hours in urine and feces was 82% and 12%, respectively, following a single 10 mg oral administration of <sup>14</sup>C-rizatriptan. Following oral administration of <sup>14</sup>C-rizatriptan, rizatriptan accounted for about 17% of circulating plasma radioactivity. Approximately 14% of an oral dose is excreted in urine as unchanged rizatriptan while 51% is excreted as indole acetic acid metabolite, indicating substantial first pass metabolism.

**Table 2**

**Pharmacokinetic Parameters of a Single Dose of Rizatriptan in Females (n = 12)**

	Arithmetic Mean (±SD)	
	RIZATRIPTAN 5mg Tablet	RIZATRIPTAN 10mg Tablet
AUC <sub>(0,4)</sub> (ng.hr/ml) <sup>a</sup>	34.5±13.0	73.9±23.4
C <sub>max</sub> (ng/ml) <sup>a</sup>	10.4±3.9	21.3±6.9
T <sub>max</sub> (hr)	1.0±0.6	1.5±0.8
t <sub>1/2</sub> (hr) <sup>b</sup>	1.7	1.7
Plasma Clearance (ml/min) <sup>c</sup>	1050.5±224.5	1081.6±239.4

<sup>a</sup>Potency-normalized

<sup>b</sup>Harmonic mean

<sup>c</sup>Plasma clearance of 1 mg stable, heavy-labeled I.V dose of rizatriptan given concomitantly with oral dose

**Special Populations and Conditions**

**Pediatrics:** The mean AUC<sub>0-∞</sub> and C<sub>max</sub> of rizatriptan (10 mg orally) were about 12% and 19% higher in adolescents (n=12) as compared to historical data in adults, respectively.

The safety and efficacy of Rizatriptan have not been established in patients under 18 years of age and its use in this age group is not recommended (see WARNINGS AND PRECAUTIONS, Special Populations and Conditions).

In an earlier randomized placebo-controlled trial with 291 adolescents migraineurs (dosed without regard to body weight), aged 12-17 years, the efficacy of Rizatriptan Tablets (5 mg) was not different from that of placebo.

**Geriatrics:** Rizatriptan pharmacokinetics in healthy elderly non-migraineur volunteers (age 65-77 years) were similar to those in younger non-migraineur volunteers (age 18-45 years).

**Gender:** The mean  $AUC_{0-\infty}$  and  $C_{max}$  of rizatriptan (10 mg orally) were about 30% and 11% higher in females as compared to males, respectively, while  $T_{max}$  occurred at approximately the same time.

**Race:** Pharmacokinetic data revealed no significant differences between African American and Caucasian subjects. The effect of race on the pharmacokinetics of rizatriptan has not been systematically evaluated.

**Hepatic Insufficiency:** Following oral administration in patients with hepatic impairment caused by mild to moderate alcoholic cirrhosis of the liver, plasma concentrations of rizatriptan were similar in patients with mild hepatic insufficiency compared to a control group of healthy subjects; plasma concentrations of rizatriptan were approximately 30% greater in patients with moderate hepatic insufficiency (see WARNINGS AND PRECAUTIONS). Since there are no data in patients with severe hepatic impairment (Child-Pugh grade C), rizatriptan is contraindicated in this population (see CONTRAINDICATIONS).

**Renal Insufficiency:** In patients with renal impairment (creatinine clearance 10-60 mL/min/1.73 m<sup>2</sup>), the  $AUC_{0-\infty}$  of rizatriptan was not significantly different from that in healthy subjects. In hemodialysis patients (creatinine clearance < 2 mL/min/1.73 m<sup>2</sup>), however, the AUC for rizatriptan was approximately 44% greater than that in patients with normal renal function (see WARNINGS AND PRECAUTIONS).

## **STORAGE AND STABILITY**

Store the tablets at room temperature (15°C - 30°C).

## **DOSAGE FORMS, COMPOSITION AND PACKAGING**

Each compressed tablet contains either 5 mg or 10 mg of rizatriptan (corresponding to 7.265 mg or 14.53 mg of the benzoate salt, respectively) and the following non-medicinal ingredients: Lactose Monohydrate, Microcrystalline Cellulose, Pregelatinized Starch, Ferric Oxide (Red) and Magnesium Stearate,

IPG-RIZATRIPTAN 5 mg is Light pink colored, capsule-shaped, biconvex uncoated tablets, with “NAT” on one side and “123” on the other side. Available in blister packages of 6 tablets and bottles of 100’s & 500’s.

IPG-RIZATRIPTAN 10 mg is Light pink colored, capsule-shaped, biconvex uncoated tablets, with “NAT” on one side and “121” on the other side. Available in blister packages of 6 tablets and bottles of 100’s & 500’s.

## PART II: SCIENTIFIC INFORMATION

### PHARMACEUTICAL INFORMATION

#### Drug Substance


Proper name: rizatriptan benzoate

Chemical name: *N,N*-dimethyl-5-(1*H*-1,2,4-triazol-1-ylmethyl)-1*H*-indole-3-ethanamine monobenzoate

Molecular formula: C<sub>15</sub>H<sub>19</sub>N<sub>5</sub>. C<sub>7</sub>H<sub>6</sub>O<sub>2</sub>

Molecular mass: 391.47 g/mol;

Structural formula:



Physicochemical properties:

**Description:** Rizatriptan benzoate is a white to almost white crystalline powder. Rizatriptan benzoate is soluble in water at about 42 mg per mL (expressed as free base) at 25°C.

## CLINICAL TRIALS

### Comparative Bioavailability Studies

#### Study results in adults

An open label, balanced, randomized, two-treatment, two-sequence, two-period, single-dose, cross-over, bioequivalence study comparing Rizatriptan Benzoate 10mg tablets of Marcan Pharmaceuticals Inc., with Maxalt 10mg tablets (containing Rizatriptan Benzoate 10mg) of Merck & Co. Inc., USA in healthy, adult, human subjects, under fasting condition. Twenty nine (29) subjects were included in the calculation of the presented pharmacokinetic parameters. The Comparative Bioavailability Data for rizatriptan is summarized below.

SUMMARY TABLE OF THE COMPARATIVE BIOAVAILABILITY DATA

Rizatriptan (1 x 10 mg) From measured data Geometric Mean Arithmetic Mean (CV %)				
Parameter	IPG- Rizatriptan*	Maxalt †	% Ratio of Least- Square Means	Confidence Interval, 90%
AUC <sub>T</sub> (ng*h/mL)	67.92 70.71 (28)	71.76 74.07 (25)	94.65	89.18 – 100.45
AUC <sub>inf</sub> (ng*h/mL)	70.22 73.13 (28)	73.79 76.16 (25)	95.17	89.72 – 100.94
C <sub>max</sub> (ng/mL)	16.43	17.88	91.89	82.49 – 102.36
T <sub>max</sub> <sup>§</sup> (h)	1.71 (61)	1.47 (42)		
T <sub>1/2</sub> <sup>§</sup> (h)	2.34 (79)	2.08 (21)		

\* IPG-Rizatriptan, Manufactured by Marcan Pharmaceuticals Inc.,

† Maxalt 10mg Tablets (Merck & Co. Inc., USA)

§ Expressed as the arithmetic mean (CV%) only

### Rizatriptan Tablets

The efficacy of rizatriptan tablets was established in four multicenter, randomized, placebo-controlled trials. Patients enrolled in these studies were primarily female (84%) and Caucasian (88%), with a mean age of 40 years (range of 18 to 71). Patients were instructed to treat a moderate to severe headache. Headache response (defined as a reduction of moderate or severe headache pain to no or mild headache pain), was assessed for up to 2 hours (Study 1) or up to 4 hours after dosing (Studies 2, 3 and 4). Associated symptoms of nausea, photophobia, and phonophobia and maintenance of response up to 24 hours postdose were evaluated. A second dose of rizatriptan tablets was allowed 2 to 24 hours after dosing for treatment of recurrent headache in Studies 1 and 2. Additional analgesics and/or antiemetics were allowed 2 hours after



initial treatment for rescue in all four studies.

In all studies, the percentage of patients achieving headache response 2 hours after treatment was significantly greater in patients who received either rizatriptan 5 or 10 mg compared to those who received placebo. In a separate study, doses of 2.5 mg were not different from placebo. Doses greater than 10 mg were associated with an increased incidence of adverse effects. The results from the 4 controlled studies using the marketed formulation are summarized in Table 3.

**Table 3**

**Response Rates<sup>1</sup> 2 Hours Following Treatment of Initial Headache**

<b>Study</b>	<b>Placebo</b>	<b>Rizatriptan Tablets 5 mg</b>	<b>Rizatriptan Tablets 10mg</b>
1	35% (n=304)	62%* (n=458)	71%*,** (n=456)
2 <sup>†</sup>	37% (n= 82)	-	77%* (n=320)
3	23% (n= 80)	63%* (n=352)	-
4	40% (n=159)	60%* (n=164)	67%* (n=385)

<sup>1</sup> Pain response is defined as a reduction in headache severity from grade 3 or 2 (severe or moderate pain) to grade 1 or 0 (mild or no pain)

\* p value < 0.05 in comparison with placebo

\*\* p value < 0.05 in comparison with 5 mg

† Results for initial headache only

**Comparisons of drug performance based upon results obtained in different clinical trials are never reliable. Because studies are conducted at different times, with different samples of patients, by different investigators, employing different criteria and/or different interpretations of the same criteria, under different conditions (dose, dosing regimen, etc.), quantitative estimates of treatment response and the timing of response may be expected to vary considerably from study to study.**

For patients with migraine -associated photophobia, phonophobia, and nausea at baseline, there was a decreased incidence of these symptoms following administration of rizatriptan compared to placebo.

Efficacy was unaffected by the presence of aura; by the gender, or age of the patient; or by concomitant use of common migraine prophylactic drugs (e.g., beta-blockers, calcium channel blockers, tricyclic antidepressants) or oral contraceptives. There were insufficient data to assess the impact of race on efficacy.

The long-term efficacy of rizatriptan 5 mg and 10 mg was investigated in a total of 1854 patients

in optional extension phases of three Phase III studies. The extension phases were single blind (except in one study where only 10 mg rizatriptan was used) and patients were randomized to either 5 mg or 10 mg rizatriptan or standard care. Approximately 16,150 attacks were treated with rizatriptan 5 mg and approximately 24,043 with rizatriptan 10 mg over a period of up to 12 months (median number of attacks treated per patient was approximately 17). Headache response was sustained (as judged by the proportion of attacks treated with rizatriptan per patient resulting in headache relief).

## **DETAILED PHARMACOLOGY**

### **Nonclinical Pharmacodynamics**

#### ***In Vitro***

Rizatriptan has been shown to be a selective agonist at human 5-HT<sub>1B</sub> (IC<sub>50</sub> 11 nM) and 5-HT<sub>1D</sub> (IC<sub>50</sub> 41nM) receptors with 20-40 fold reduced activity at 5-HT<sub>1A</sub> receptors using radioligand binding and functional pharmacological bioassays.

Rizatriptan has no significant activity at 5-HT<sub>2</sub> or 5-HT<sub>3</sub> receptor subtypes or at alpha- and beta-adrenergic receptors, dopaminergic, histaminergic, muscarinic or benzodiazepine receptors.

Rizatriptan contracted human isolated cerebral (middle meningeal) arteries (EC<sub>50</sub> 0.09 micromolar) yet was significantly less potent in contracting human isolated coronary arteries (EC<sub>50</sub> 1.0 micromolar) with a 5.6 fold lower maximum response.

Rizatriptan's primary metabolite, indole acetic acid derivative, was devoid of activity at human 5-HT receptors. The minor metabolite, monodesmethyl rizatriptan retained activity at human 5-HT<sub>1B</sub> (IC<sub>50</sub> 19 nM) and 5-HT<sub>1D</sub> (IC<sub>50</sub> 9 nM) receptors. The 6-hydroxytryptamine metabolite of rizatriptan that is found only at low levels in humans had an affinity of IC<sub>50</sub> 600nM at  $\alpha_{2a}$  and of IC<sub>50</sub> 140 nM at  $\alpha_{2c}$  adrenergic receptors.

#### ***In vivo***

Rizatriptan caused dose-dependent vasoconstriction within the carotid artery vascular bed in anesthetized dogs (ED<sub>50</sub> 54  $\mu$ g/kg I.V.; plasma concentration 16 ng/mL) with selectivity for the craniovasculature since effects were not seen in the coronary artery until high (>137 ng/mL) plasma concentrations were reached. In ferrets rizatriptan caused vasoconstriction in the carotid artery (ED<sub>50</sub> 20  $\mu$ g/kg I.V.) with no evidence for tachyphylaxis to 3 sequential doses of 15  $\mu$ g/kg I.V.

Rizatriptan produced a dose-dependent inhibition of dural plasma protein extravasation, dural vasodilatation, and central trigeminal pain signal transmission evoked by electrical stimulation of the trigeminal ganglion or dura mater in anesthetized rats.

Rizatriptan (1.0 mg/kg P.O.; plasma level 317 ng/mL) was free of cardiovascular effects in conscious dogs but at higher doses (5 mg/kg P.O. plasma level 1642 ng/mL) produced increases in blood pressure (~ 40 mmHg) and heart rate (~ 80 beats/min) and behavioural activation that were sustained for at least 120 minutes.

Rizatriptan caused a 12% increase in mean arterial blood pressure (mmHg) and a 16% decrease in heart rate (beats/min) with lengthening of the P-R interval (90-103 msec) in anesthetized dogs. QT intervals were unchanged.

Rizatriptan prevented the increases in blood pressure evoked in pithed rats and guinea pigs by stimulation of vascular pre-ganglionic sympathetic nerves.

Rizatriptan caused increases in urine flow (0.46 to 0.83 mL/min), sodium (56-110 microEq/min) and potassium (28- 40 microEq/min) excretion in conscious dogs. Average glomerular filtration and effective renal plasma flow rates were unchanged.

Rizatriptan caused behavioural activation and mydriasis at doses >2 mg/kg P.O.

Rizatriptan, in conscious rhesus monkeys, caused behavioral activation and emesis at plasma levels of 260 ng/mL. Rizatriptan (7 mg/kg S.C.) caused mild hypothermia and emesis in squirrel monkeys at high plasma levels of 1546 ng/mL.

### **Animal Pharmacokinetics**

The absorption of oral solutions of rizatriptan in rats and dogs was rapid and high. The time to reach the maximum plasma concentration was 71 minutes in rats and 26 minutes in dogs. The absorption, as measured by the relative urinary excretion of total radioactivity (drug and metabolites) after oral and intravenous doses, was 78% in rats and essentially complete in dogs. The oral bioavailability, as estimated by comparison of plasma AUC<sub>0-∞</sub> of rizatriptan after oral and intravenous doses, was 64% in rats and 47% in dogs. The apparent discrepancy in the absorption and bioavailability of rizatriptan is consistent with moderate first-pass metabolism of the drug.

The pharmacokinetics of rizatriptan were generally similar in rats and dogs. Plasma concentrations of rizatriptan declined biexponentially with terminal half-lives of 65 minutes in rats and 72 minutes in dogs. The plasma clearance was rapid in rats (71 mL/min/kg) and dogs (46 mL/min/kg). The steady state volumes of distribution were 4.3 L/kg in rats and 3.2 L/kg in dogs. The systemic exposure of rizatriptan increased slightly greater than proportionally with increases in oral doses in rats (3 and 10 mg/kg) and dogs (1, 2 and 5 mg/kg).

Radioactivity was distributed widely throughout the body after a single 3 mg/kg oral or intravenous dose of <sup>14</sup>C rizatriptan in rats. In general, the pattern of distribution was similar after both routes of administration. Tissues with significant percentages of dose were the liver, kidney, small intestine and stomach, while the brain contained only a trace amount of radioactivity. At 24 hours, tissue concentrations of rizatriptan equivalents had declined to trace levels except in organs involved in elimination (i.e., liver, kidney and the G.I. tract). Over a 24-hour period, the

recoveries of radioactivity in the G.I. contents and excreta were 85% and 96% of I.V. and oral doses, respectively.

Toxicokinetic studies demonstrated high placental transfer of rizatriptan in pregnant rats and rabbits and high milk transfer in lactating rats (see TOXICOLOGY, Reproduction).

Rizatriptan was poorly bound to plasma proteins of mouse, rat, rabbit, dog and human and the binding was independent of concentration in the range 50- 5000 ng/mL. The unbound fraction was 81% in mouse, 82% in rat, 73% in rabbit, 88% in dog and 86% in human.

The pathways of rizatriptan metabolism in laboratory animals (rat, mouse, rabbit, dog) are similar to those in humans, and consist of oxidative deamination, aromatic hydroxylation followed by sulfation, N-oxidation and N-demethylation.

In general, there were no major differences in the qualitative metabolism of rizatriptan in rodents and humans with respect to dosing frequency (single or multiple oral doses), the size of the dose (therapeutically relevant or toxicological doses) or sex. The extent of metabolism, however, was species dependent. Rizatriptan and its metabolites were eliminated primarily via renal excretion, although biliary excretion also played a role in rats. Rizatriptan underwent more extensive metabolism in dogs and humans when compared to two rodent species, less than 15% of oral doses being excreted in urine as the unchanged drug. The indole acetic acid derivative of rizatriptan was the most prominent metabolite in all species studied although the N-oxide analog also was a significant metabolite in rats and dogs.

In rats, both urinary and biliary routes contributed to the elimination of rizatriptan and its metabolites, while in dogs and human, renal excretion was the primary route of elimination. Accordingly, the excretion of an intravenous dose of rizatriptan in rats was 58% in urine and 23% in feces, while that of the oral dose was balanced evenly between urine (45%) and feces (42%). Approximately one-half (I.V.) and one-third (p.o.) of the urinary radioactivity represented rizatriptan while the remainder was divided among several metabolites. In dogs, 75% of oral and intravenous doses was excreted in urine and 12- 13% in feces. The majority of urinary radioactivity after either route of administration represented metabolites while the unchanged drug accounted for approximately 7% (p.o.) and 21% (I.V.) of the radioactivity.

### **Human Pharmacodynamics**

In healthy young male and female subjects who received maximal doses of rizatriptan (10 mg every 2 hours for three doses), slight increases in blood pressure (approximately 2-3 mmHg) were observed. These small, transient increases in blood pressure were not clinically significant. During long-term monitoring of migraine patients in controlled studies, no consistent effects on blood pressure or heart rate were observed.

In a study in healthy male subjects, rizatriptan 10 mg produced slight, transient peripheral vasoconstriction (measured as a 5.1 mmHg increase in toe-arm systolic blood pressure gradient). In contrast, intravenous ergotamine (0.25 mg) produced a 14.6 mmHg increase in toe-arm systolic blood pressure gradient. When ergotamine and rizatriptan were given together, the

increase in toe-arm systolic blood pressure gradient was similar to that when ergotamine was given alone.

No change in the ECG parameters measured was observed when two 10-mg doses of rizatriptan, separated by 2 hours, were studied in 157 migraine patients (age range 18 to 72 years) during a migraine attack.

## **TOXICOLOGY**

### **Acute Toxicity**

The approximate oral LD<sub>50</sub> of rizatriptan was 700 mg/kg and 2227 mg/kg in mice and rats, respectively. The approximate intravenous LD<sub>50</sub> values were 89 and 141 mg/kg in mice and rats, respectively.

### **Chronic Toxicity**

The toxicity potential of rizatriptan was evaluated in a series of repeat dose oral toxicity studies up to one year in dogs and rats and up to 14 weeks in mice. There were no adverse findings that would preclude administration of rizatriptan (rizatriptan benzoate) at the recommended therapeutic dosages to humans.

### **Carcinogenesis**

The carcinogenic potential of rizatriptan was evaluated in a 106-week study in rats and a 100-week study in mice at oral doses of up to 125 mg/kg/day (625-fold the human dose of 10 mg or 0.2 mg/kg). These doses produced exposure margins (AUC ratios) of up to 600- and 400-fold in rats and mice, respectively, over the human systemic exposure at a therapeutic dose of 10 mg (0.2 mg/kg). For both rats and mice, no evidence of carcinogenicity was seen with increasing doses of rizatriptan.

### **Mutagenesis**

Rizatriptan, with and without metabolic activation, was neither genotoxic, mutagenic, nor clastogenic in all *in vitro* and *in vivo* genetic toxicity studies, including: microbial mutagenesis, *in vitro* chromosome aberration assays, *in vitro* V-79 mammalian cell mutagenesis assays, an *in vitro* alkaline elution/rat hepatocyte assay, and an *in vivo* chromosome aberration assay in mouse bone marrow.

### **Reproduction**

No adverse effects on fertility or reproductive performance, and no fetal toxicity or malformations were observed in female and male rats (except slightly decreased body weight at the high dose) given oral doses of rizatriptan up to 100 and 250 mg/kg/day, respectively (500- and 1250-fold the human dose of 10 mg or 0.2 mg/kg). In addition, no adverse effects on reproductive parameters were detected during early or late gestation, or during the lactation period. These doses provided exposure margins more than 900-fold over the human systemic exposure, based on the AUC ratio derived from rat maternal drug levels compared to humans treated with 10 mg (0.2 mg/kg). High placental transfer occurred, as evidenced by rat fetal plasma levels of 20 to 40% of the maternal plasma levels. High milk transfer occurred, and

resulted in rat milk levels that were 5- fold, or greater, the maternal plasma levels. Although high maternal, fetal and neonatal exposure to rizatriptan occurred in these studies, no adverse treatment-related effects were observed on F<sub>1</sub> survival, development, behavior, reproductive performance, or testicular histology, nor were there any effects seen in the F<sub>2</sub> offspring.

No adverse effects on development, and no fetal toxicity or malformations were observed in pregnant rabbits (except slightly decreased body weight at the high dose) given oral doses of rizatriptan up to 50 mg/kg/day (250- fold the human dose of 10 mg or 0.2 mg/kg). These doses produced high maternal drug levels, resulting in a 475-fold exposure margin, based on the AUC ratio derived from rabbit maternal drug levels compared to humans treated with 10 mg (0.2 mg/kg). High placental transfer occurred with rabbit fetal tissue levels reaching 42 to 49% of the maternal plasma levels.

### **Development**

There were no adverse effects on fetal development in rats or rabbits exposed to large multiples of the human therapeutic dose of rizatriptan during early and late gestation. High placental transfer of rizatriptan was documented by fetal plasma and tissue levels.

## REFERENCES

1. Goldberg MR, Lowry RC, Musson DG, Birk KL, Fisher A, DePuy ME, Shadle CR. Lack of pharmacokinetic and pharmacodynamic interaction between rizatriptan and paroxetine. *J Clin Pharmacol* 1999;39:192-9.
2. Ho TW, Pearlman E, Lewis D, Hamalainen M, Connor K, Michelson D, Zhang Y, Assaid C, Mozley LH, Strickler N, Bachman R, Mahoney E, Lines C and Hewitt DJ. Efficacy and tolerability of rizatriptan in pediatric migraineurs: Results from a randomized, double-blind, placebo-controlled trial using a novel adaptive enrichment design.
3. Longmore J, Boulanger CM, Desta B, Hill RG, Schofield WN, Taylor AA. 5-HT<sub>1D</sub> receptor agonists and human coronary artery reactivity *in vitro*: crossover comparisons of 5-HT and sumatriptan with rizatriptan and L-741,519. *Br J Clin Pharmacol* 1996;42:431-41.
4. Longmore J, Razzaque Z, Shaw D, Davenport AP, Maguire J, Pickard JD, Schofield WN, Hill RG. Comparison of the vasoconstrictor effects of rizatriptan and sumatriptan in human isolated cranial arteries: immunohistological demonstration of the involvement of 5-HT<sub>1B</sub>-receptors. *Br J Clin Pharmacol* 1998;46:577–582.
5. Visser HW, Terwindt GM, Reines SA, Jiang K, Lines CR, Ferrari MD. Rizatriptan vs. sumatriptan in the acute treatment of migraine. *Arch of Neurol* 1996;53:1132-7.
6. Product Monograph for Maxalt® (Rizatriptan benzoate tablets) 5mg and 10mg, submission control No. 151004, date of revision: May 08, 2013

## PART III: CONSUMER INFORMATION

### Pr **IPG-RIZATRIPTAN** rizatriptan benzoate tablets

This leaflet is part III of a three-part "Product Monograph" published when IPG-RIZATRIPTAN was approved for sale in Canada and is designed specifically for Consumers. This leaflet is a summary and will not tell you everything about IPG-RIZATRIPTAN. Contact your physician or pharmacist if you have any questions about the drug.

Please read this leaflet carefully before you take IPG-RIZATRIPTAN. It provides a summary of the information available on your medication. Please do not throw it away. Keep it for reference until you have finished your medicine. This leaflet does not contain all the information available on IPG-RIZATRIPTAN. For further information or advice, ask your physician or pharmacist.

#### ABOUT THIS MEDICATION

The name of your medicine is IPG-RIZATRIPTAN. It can only be obtained by prescription from your physician. The decision to use IPG-RIZATRIPTAN is one that you and your physician should make jointly, taking into account your individual preferences and medical circumstances. If you have risk factors for heart disease (such as high blood pressure, high cholesterol, obesity, diabetes, smoking, strong family history of heart disease, or you are a post-menopausal female, or a male over 40), tell your physician. Your physician should evaluate you for heart disease in order to determine whether IPG-RIZATRIPTAN is appropriate for you.

#### What the medication is used for

IPG-RIZATRIPTAN is intended to relieve your migraine headache and other associated symptoms of a migraine attack in adults. **IPG-RIZATRIPTAN should not be used continuously to prevent or reduce the number of attacks you experience. Use IPG-RIZATRIPTAN only to treat an actual migraine headache attack.**

#### What it does:

Migraine headache is believed to be caused by a widening of the blood vessels in the head. IPG-RIZATRIPTAN narrows the vessels and relieves the pain and other symptoms of migraine headache.

#### When it should not be used

Do not take IPG-RIZATRIPTAN if you:

- are allergic to any of the ingredients (see **What the nonmedicinal ingredients are** section)
- have uncontrolled or severe high blood pressure
- have heart disease or history of heart disease
- have severe liver disease
- have had a stroke or transient ischemic attack (TIA)

- have or have had blood vessels problems, including ischemic bowel disease
- take monoamine oxidase (MAO) inhibitors such as moclobemide, phenelzine, tranylcypromine, or pargyline, or have taken MAO inhibitors within the last two weeks.

IPG-RIZATRIPTAN should not be used to relieve pain other than that associated with migraine headache.

IPG-RIZATRIPTAN should not be used within 24 hours of treatment with another 5-HT<sub>1</sub> agonist such as sumatriptan (Imitrex<sup>\*</sup>), naratriptan (Amerge<sup>\*</sup>) or zolmitriptan (Zomig<sup>\*</sup>), or ergotamine-type medications such as ergotamine, dihydroergotamine or methysergide.

Please also see WARNINGS AND PRECAUTIONS.

#### What the medicinal ingredient is:

rizatriptan benzoate

#### What the nonmedicinal ingredients are:

Ferric oxide (red), lactose monohydrate, magnesium stearate, microcrystalline cellulose and pregelatinized starch.

#### What dosage forms it comes in:

IPG-RIZATRIPTAN (rizatriptan benzoate) is available as a 5 mg or 10 mg pale pink tablet containing either 5 mg or 10 mg of rizatriptan.

#### WARNINGS AND PRECAUTIONS

If the answer to any of the following questions is YES or if you do not know the answer, then please speak with your physician before you take any IPG-RIZATRIPTAN.

- Are you pregnant? Do you think you might be pregnant? Are you trying to become pregnant? Are you using inadequate contraception? Are you breast feeding?
- **Do you or have you ever experienced any pain or tightness in the chest, (which may or may not spread to your neck, jaw, or upper arm), shortness of breath, rapid heartbeats or irregular heartbeats? Do you have angina?**
- Have you ever had heart or blood vessel disease? Have you had a heart attack or stroke?
- **Do you have risk factors for heart disease, such as: high blood pressure, high cholesterol, smoking, obesity, diabetes, or strong family history of heart disease? Are you post-menopausal or a male over 40?**
- Have you ever had to stop taking this or any other medication because of an allergy or other problems?
- Are you taking any other migraine 5-HT<sub>1</sub> agonist medications such as Imitrex<sup>\*</sup> (sumatriptan succinate/sumatriptan) or Amerge<sup>\*</sup> (naratriptan as naratriptan hydrochloride), Zomig<sup>\*</sup> (zolmitriptan) or migraine medications containing ergotamine, dihydroergotamine, or methysergide?



- Are you taking any medications for depression such as selective serotonin reuptake inhibitors (SSRIs) such as sertraline, escitalopram oxalate, and fluoxetine or serotonin norepinephrine reuptake inhibitors (SNRIs) such as venlafaxine, duloxetine, or monoamine oxidase inhibitors (MOIs)?
- Have you ever experienced numbness on one side of your body when you have a headache?
- Have you ever had, or do you have epilepsy or seizures?
- Have you ever had, or do you have liver disease or kidney disease?
- Is this headache different from your usual migraine attacks?
- Are you over 65 years of age?

#### **Continuous use of IPG-Rizatriptan:**

If you take IPG-Rizatriptan too often, this may result in you getting chronic headaches or marked increases in the frequency of your migraine attacks. In such cases, you should contact your doctor, as you may have to stop taking IPG-Rizatriptan.

If you are not sure whether you should take IPG-RIZATRIPTAN, contact your physician or pharmacist.

#### **IPG-RIZATRIPTAN use during pregnancy:**

Do not use IPG-RIZATRIPTAN if you are pregnant, think you might be pregnant, are trying to become pregnant or are using inadequate contraception, unless you have discussed this with your physician.

### **INTERACTIONS WITH THIS MEDICATION**

If you are taking any other migraine medication, check with your physician first before taking IPG-RIZATRIPTAN and inform your physician of any other medications you may be taking, including those obtained without a prescription.

Do not take IPG-RIZATRIPTAN if you take monoamine oxidase (MAO) inhibitors such as moclobemide, phenelzine, tranylcypromine, or pargyline, or have taken MAO inhibitors within the last two weeks.

Ask your physician for instructions about taking IPG-RIZATRIPTAN if you are taking selective serotonin reuptake inhibitors (SSRIs) such as sertraline, escitalopram oxalate, and fluoxetine or serotonin norepinephrine reuptake inhibitors (SNRIs) such as venlafaxine, duloxetine for depression. A life-threatening condition called serotonin syndrome can happen when medicines called triptans, such as IPG-RIZATRIPTAN, and medicines used to treat depression and mood disorders called SSRIs and SNRIs are used together. Signs and symptoms of serotonin syndrome include the following: restlessness, diarrhea, hallucinations, coma, loss of coordination, nausea, fast heart beat, vomiting, increased body temperature, changes in blood pressure and overactive reflexes.

### **PROPER USE OF THIS MEDICATION**

**REMEMBER:** this medicine was prescribed only for YOU. Only a physician knows who can use it safely. Never give it to someone else. It may harm them, even if their symptoms are the same as yours.

#### **Usual dose**

For adults, the usual dosage is 5 mg or 10 mg, as recommended by your physician. The dose should be taken as soon as your migraine appears, but it may be taken at any time during your migraine headache.

If you are taking a tablet, swallow your dose with water.

A second dose may be taken if your headache returns. Repeat doses cannot be taken any sooner than 2 hours following the first dose. Do not take more than 20 mg in any 24-hour period.

For any attack where you have no response to the first dose, do not take a second dose without first consulting your physician.

#### **Overdose:**

**In case of drug overdose, contact a health care practitioner, hospital emergency department or regional Poison Control Centre immediately, even if there are no symptoms.**

### **SIDE EFFECTS AND WHAT TO DO ABOUT THEM**

Although the vast majority of IPG-RIZATRIPTAN users have not experienced any significant problems, you should be aware of the following side effects:

#### **Common side effects**

- **Sensations of pain, pressure or tightness in the chest, neck, throat, jaw or arms. If this happens to you, then discuss it with your physician before using any more IPG-RIZATRIPTAN. If the chest pain is severe (may resemble an angina attack) or does not go away, stop taking drug and call your physician immediately.**
- Tingling, heat, heaviness or pressure, dizziness, tiredness, or feeling sick. Tell your physician if you have any of these symptoms.
- Drowsiness. Do not drive or operate machinery until you are sure that you are not drowsy.
- Irregular or rapid heartbeats. If this happens to you, stop taking drug and then discuss it with your physician before using any more IPG-RIZATRIPTAN.

#### **Uncommon side effects**

- Rapid onset of numbness or weakness on one side of the body (often beginning in one arm). If this occurs, stop taking drug and call your physician immediately.
- Muscle pain. If this occurs, stop taking drug and call your physician immediately.

#### **Rare side effects**

- shortness of breath, wheeziness, heart throbbing, swelling of the eyelids, face, tongue or lips; or a skin rash or skin lumps

or hives or spasms of blood vessels of the extremities including coldness and numbness of the hands and feet. These effects happen rarely, but if any do occur, do not take any more IPG-RIZATRIPTAN and contact your physician immediately.

- Sudden or severe abdominal pain. If this occurs, stop taking drug and call your physician immediately.

**Very rare side effects**

- There have been very rare reports of seizures. If this occurs, stop taking drug and call your physician immediately.
- Spasm of the blood vessels of the colon (large bowel). If this occurs, stop taking drug and call your physician immediately.

If you feel unwell in any other way or have symptoms that you do not understand or find distressing, you should contact your physician immediately.

SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM			
Symptom/ effect	Talk to your physician or pharmacist		Stop taking drug and seek immediate emergency medical attention.
	Only if severe	In all cases	
	Sudden or severe abdominal pain		√
Very rare	Seizures		√
	Spasm of the blood vessels of the colon (large bowel)		√

*This is not a complete list of side effects. For any unexpected effects while taking IPG-RIZATRIPTAN, contact your physician or pharmacist.*

SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM			
Symptom/ effect	Talk to your physician or pharmacist		Stop taking drug and seek immediate emergency medical attention.
	Only if severe	In all cases	
Common	Sensation of pain, pressure or tightness in the chest, neck, throat, jaw or arms		√
	Tingling, heat, heaviness or pressure in any part of the body	√	
	Headache, dizziness, tiredness, nausea, vomiting, dry mouth, drowsiness	√	
	Irregular or rapid heartbeats		√
Uncommon	Rapid onset of numbness or weakness on one side of the body (often beginning in one arm)		√
	Muscle pain		√
Rare	Shortness of breath, wheeziness, heart throbbing, swelling of the eyelids, face, tongue or lips; or a skin rash or skin lumps or hives or spasms of blood vessels of the extremities		√

**HOW TO STORE IT**

Keep your medicine in a safe place where children cannot reach it. IPG-RIZATRIPTAN could be harmful to children. Store your medication between 15°C and 30°C.

If your physician decides to stop your treatment, do not keep any leftover medicine unless your physician tells you to do so.

If your medicine has expired, do not use it. It is recommended to take leftover medications to a pharmacy for proper disposal.

**REPORTING SUSPECTED SIDE EFFECTS**

You can report any suspected adverse reactions associated with the use of health products to the Canada Vigilance Program by one of the following 3 ways:

- Report online at [www.healthcanada.gc.ca/medeffect](http://www.healthcanada.gc.ca/medeffect)
- Call toll-free at 1-866-234-2345
- Complete a Canada Vigilance Reporting Form and:  
Fax toll-free to 1-866-678-6789, or  
Mail to:  
Canada Vigilance Program  
Health Canada  
Postal Locator 0701E  
Ottawa, Ontario  
K1A 0K9

Postage paid labels, Canada Vigilance Reporting Form and the adverse reaction reporting guidelines are available on the MedEffect™ Canada Web site at [www.healthcanada.gc.ca/medeffect](http://www.healthcanada.gc.ca/medeffect).

Note: Should you require information related to the management of side effects, contact **IMPORTANT: PLEASE READ** your health professional. The Canada Vigilance Program does not provide medical advice.

## **MORE INFORMATION**

This document plus the full product monograph, prepared for health professionals can be found at: <http://www.marcanpharma.com>. Or by contacting the sponsor, Marcan Pharmaceuticals Inc. at 1-613-228-2600.

Date of Preparation: January 8, 2016.