

PRODUCT MONOGRAPH

^{Pr}**pms-ENALAPRIL**

Enalapril Maleate Tablets, House Standard

2.5 mg, 5 mg, 10 mg and 20 mg

Each tablet is made with 2.5 mg, 5 mg, 10 mg or 20 mg of enalapril maleate that appears as 2 mg, 4 mg, 8 mg or 16 mg of enalapril sodium in the tablets

Angiotensin Converting Enzyme Inhibitor

PHARMASCIENCE INC.
6111 Royalmount Ave., Suite 100
Montréal, QC, Canada
H4P 2T4

Date of Revision:
April 7, 2016

www.pharmascience.com

Submission Control No.: 193659

Table of contents

PART I: HEALTH PROFESSIONAL INFORMATION	3
SUMMARY PRODUCT INFORMATION	3
INDICATIONS AND CLINICAL USE	3
CONTRAINDICATIONS.....	4
WARNINGS AND PRECAUTIONS.....	4
ADVERSE REACTIONS	9
DRUG INTERACTIONS.....	13
DOSAGE AND ADMINISTRATION	14
OVERDOSAGE.....	16
ACTION AND CLINICAL PHARMACOLOGY	17
STORAGE AND STABILITY	19
DOSAGE FORMS, COMPOSITION AND PACKAGING.....	19
PART II: SCIENTIFIC INFORMATION.....	20
PHARMACEUTICAL INFORMATION	20
CLINICAL TRIALS.....	21
DETAILED PHARMACOLOGY.....	25
TOXICOLOGY	27
REFERENCES.....	33
PART III: CONSUMER INFORMATION	36

Pr pms-ENALAPRIL

Enalapril Maleate Tablets, House Standard

PART I: HEALTH PROFESSIONAL INFORMATION

SUMMARY PRODUCT INFORMATION

Route of Administration	Dosage Form/ Strength	All Non Medicinal Ingredients
Oral	Tablet* 2.5 mg, 5 mg, 10 mg, 20 mg	Copovidone, lactose, pregelatinized starch, sodium bicarbonate and zinc stearate. 10 mg tablets also contain: red iron oxide 20 mg tablets also contain: red iron oxide and yellow iron oxide

*The splitting of pms-ENALAPRIL tablets is not advised. Each tablet is made with 2.5 mg, 5 mg, 10 mg or 20 mg enalapril maleate that appears as 2 mg, 4 mg, 8 mg or 16 mg enalapril sodium in the tablet.

INDICATIONS AND CLINICAL USE

pms-ENALAPRIL (enalapril) is indicated for:

- essential or renovascular hypertension
- treatment of symptomatic congestive heart failure

Hypertension

pms-ENALAPRIL is indicated in the treatment of essential or renovascular hypertension. It is usually administered in association with other drugs, particularly thiazide diuretics.

In using pms-ENALAPRIL consideration should be given to the risk of angioedema (see WARNINGS AND PRECAUTIONS).

Geriatrics (>65 years of age): See DOSAGE AND ADMINISTRATION.

Pediatrics (<16 years of age): pms-ENALAPRIL may be used in children (see DOSAGE AND ADMINISTRATION).

Congestive Heart Failure

pms-ENALAPRIL is indicated in the treatment of symptomatic congestive heart failure usually in combination with diuretics and/or digitalis. In these patients, enalapril improves symptoms, increases survival, and decreases the frequency of hospitalization (see CLINICAL TRIALS for details and limitations of survival trials). Treatment with enalapril should be initiated under close medical supervision.

In clinically stable asymptomatic patients with left ventricular dysfunction (ejection fraction $\leq 35\%$), enalapril decreases the rate of development of overt heart failure and decreases the

incidence of hospitalization for heart failure (see CLINICAL TRIALS for details and limitations of survival trials).

CONTRAINDICATIONS

pms-ENALAPRIL (enalapril) is contraindicated in:

- Patients who are hypersensitive to these products or to any ingredient in their formulation. For a complete listing, see the DOSAGE FORMS, COMPOSITION AND PACKAGING section of the product monograph.
- Patients with a history of angioneurotic edema relating to previous treatment with an angiotensin converting enzyme inhibitor.
- Patients with hereditary or idiopathic angioedema.

Concomitant use of angiotensin converting enzyme inhibitors (ACEIs) – including enalapril – with aliskiren-containing drugs in patients with diabetes mellitus (type 1 or type 2) or moderate to severe renal impairment (GFR < 60 ml/min/1.73m²) is contraindicated (see WARNINGS AND PRECAUTIONS, Cardiovascular, Dual Blockade of the Renin-Angiotensin System (RAS) and Renal, and DRUG INTERACTIONS, Dual Blockade of the Renin-Angiotensin System (RAS) with ACEIs, ARBs or aliskiren-containing drugs).

WARNINGS AND PRECAUTIONS

Serious Warnings and Precautions

When used in pregnancy, angiotensin converting enzyme (ACE) inhibitors can cause injury or even death of the developing fetus. When pregnancy is detected, pms-ENALAPRIL should be discontinued as soon as possible.

General

Angioedema: Angioedema of the face, extremities, lips, tongue, glottis and/or larynx has been reported rarely in patients treated with enalapril. This may occur at any time during treatment and may be life threatening.

Very rarely, fatalities have been reported due to angioedema associated with laryngeal edema or tongue edema. Patients with involvement of the tongue, glottis or larynx are likely to experience airway obstruction, especially those with a history of airway surgery. However, where there is involvement of the tongue, glottis or larynx, likely to cause airway obstruction, appropriate therapy which may include subcutaneous adrenaline solution 1:1000 (0.3 mL to 0.5 mL) and/or measures to ensure a patent airway should be administered promptly when indicated.

If angioedema occurs, pms-ENALAPRIL should be discontinued promptly and appropriate monitoring should be instituted to ensure complete resolution of symptoms prior to dismissing the patient. Even in those instances where swelling of only the tongue is involved, without respiratory distress, patients may require prolonged observation since this may be life threatening and treatment with antihistamines and corticosteroids may not be sufficient.

In patients who experience angioedema, future administration is contraindicated (see CONTRAINDICATIONS).

The incidence of angioedema during ACE inhibitor therapy has been reported to be higher in black than in non-black patients.

Patients with a history of angioedema unrelated to ACE inhibitor therapy may be at increased risk of angioedema while receiving an ACE inhibitor (see CONTRAINDICATIONS).

Patients receiving coadministration of ACE inhibitor and mTOR (mammalian target of rapamycin) inhibitor (e.g., temsirolimus, sirolimus, everolimus) therapy may be at increased risk for angioedema. Caution should be used when these drugs are used concomitantly (see DRUG INTERACTIONS).

Anaphylactoid Reactions during Membrane Exposure: Anaphylactoid reactions have been reported in patients dialysed with high-flux membranes (e.g., polyacrylonitrile [PAN]) and treated concomitantly with an ACE inhibitor. Dialysis should be stopped immediately if symptoms such as nausea, abdominal cramps, burning, angioedema, shortness of breath and severe hypotension occur. Symptoms are not relieved by antihistamines. In these patients consideration should be given to using a different type of dialysis membrane or a different class of antihypertensive agent.

Anaphylactoid Reactions during Desensitization: There have been isolated reports of patients experiencing sustained life-threatening anaphylactoid reactions while receiving ACE inhibitors during desensitizing treatment with hymenoptera (bees, wasp) venom. In the same patients, these reactions have been avoided when ACE inhibitors were temporarily withheld for at least 24 hours, but they have reappeared upon inadvertent rechallenge.

Anaphylactoid Reactions during LDL Apheresis: Rarely, patients receiving ACE inhibitors during low-density lipoprotein (LDL)-apheresis with dextran sulfate have experienced life-threatening anaphylactoid reactions. These reactions were avoided by temporarily withholding ACE inhibitor therapy prior to each apheresis.

Cardiovascular

Hypotension: Symptomatic hypotension has occurred after administration of enalapril, usually after the first or second dose or when the dose was increased. It is more likely to occur in patients who are volume depleted by diuretic therapy, dietary salt restriction, dialysis, diarrhea, or vomiting. In patients with severe congestive heart failure, with or without associated renal insufficiency, excessive hypotension has been observed and may be associated with oliguria and/or progressive azotemia, and rarely with acute renal failure and/or death. Because of the potential fall in blood pressure in these patients, therapy should be started under very close medical supervision, usually in a hospital. Such patients should be followed closely for the first two weeks of treatment and whenever the dose of enalapril and/or diuretic is increased. Similar considerations may apply to patients with ischemic heart or cerebrovascular disease in whom an excessive fall in blood pressure could result in a myocardial infarction or cerebrovascular accident (see ADVERSE REACTIONS).

If hypotension occurs, the patient should be placed in supine position and, if necessary, receive an intravenous infusion of normal saline. A transient hypotensive response is not a contraindication to further doses which usually can be given without difficulty once the blood pressure has increased after volume expansion.

Valvular Stenosis: There is concern on theoretical grounds that patients with aortic stenosis might be at particular risk of decreased coronary perfusion when treated with vasodilators because they do not develop as much afterload reduction.

Dual Blockade of the Renin-Angiotensin System (RAS)

There is evidence that co-administration of angiotensin converting enzyme inhibitors (ACEIs), such as enalapril, or of angiotensin receptor antagonists (ARBs) with aliskiren increases the risk of hypotension, syncope, stroke, hyperkalemia and deterioration of renal function, including renal failure, in patients with diabetes mellitus (type 1 or type 2) and/or moderate to severe renal impairment (GFR < 60 ml/min/1.73m²). Therefore, the use of pms-ENALAPRIL, in combination with aliskiren-containing drugs is contraindicated in these patients (see CONTRAINDICATIONS). Further, co-administration of ACEIs, including pms-ENALAPRIL, with other agents blocking the RAS, such as ARBs or aliskiren-containing drugs, is generally not recommended in other patients, since such treatment has been associated with an increased incidence of severe hypotension, renal failure, and hyperkalemia.

Ear/Nose/Throat

Cough: A dry, persistent cough, which usually disappears only after withdrawal or lowering of the dose of enalapril has been reported. Such possibility should be considered as part of the differential diagnosis of the cough.

Endocrine and Metabolism

Hypoglycemia: Rare cases of hypoglycemia in diabetic patients on oral antidiabetic agents or insulin have been reported. Diabetic patients treated with oral antidiabetic agents or insulin starting an ACE inhibitor should be told to closely monitor for hypoglycemia, especially during the first month of combined use. In addition, hypoglycemia appeared to be more likely to occur during the first weeks of combined treatment and in patients with renal impairment (see ADVERSE REACTIONS).

Hematologic

Neutropenia/Agranulocytosis: Agranulocytosis and bone marrow depression have been caused by angiotensin converting enzyme inhibitors. Several cases of agranulocytosis and neutropenia have been reported in which a causal relationship to enalapril cannot be excluded. Current experience with the drug shows the incidence to be rare. Periodic monitoring of white blood cell counts should be considered, especially in patients with collagen vascular disease and renal disease.

Hepatic/Biliary/Pancreatic

Patients with Impaired Liver Function: Hepatitis, jaundice (hepatocellular and/or cholestatic), elevations of liver enzymes and/or serum bilirubin have occurred during therapy with enalapril in patients with or without pre-existing liver abnormalities (see ADVERSE REACTIONS). In most cases the changes were reversed on discontinuation of the drug.

Should the patient receiving pms-ENALAPRIL experience any unexplained symptoms (see CONSUMER INFORMATION), particularly during the first weeks or months of treatment, it is recommended that a full set of liver function tests and any other necessary investigation be carried out. Discontinuation of enalapril should be considered when appropriate.

There are no adequate studies in patients with cirrhosis and/or liver dysfunction. pms-ENALAPRIL should be used with particular caution in patients with pre-existing liver abnormalities. In such patients baseline liver function tests should be obtained before administration of the drug and close monitoring of response and metabolic effects should apply.

Nitritoid Reactions – Gold: Nitritoid reactions (symptoms include facial flushing, nausea, vomiting and symptomatic hypotension) have been reported rarely in patients on therapy with injectable gold (sodium aurothiomalate) and concomitant ACE inhibitor therapy including enalapril (see DRUG INTERACTIONS).

Peri-Operative Considerations

Surgery/Anesthesia: In patients undergoing major surgery or during anesthesia with agents that produce hypotension, enalapril blocks angiotensin II formation, secondary to compensatory renin release. If hypotension occurs and is considered to be due to this mechanism, it can be corrected by volume expansion.

Renal

Renal Impairment: As a consequence of inhibiting the renin-angiotensin-aldosterone system, changes in renal function have been seen in susceptible individuals. In patients whose renal function may depend on the activity of the renin-angiotensin-aldosterone system, such as patients with bilateral renal artery stenosis, unilateral renal artery stenosis to a solitary kidney, or severe congestive heart failure, treatment with agents that inhibit this system has been associated with oliguria, progressive azotemia, and rarely, acute renal failure and/or death. In susceptible patients, concomitant diuretic use may further increase risk.

The use of ACEIs – including of pms-ENALAPRIL – or ARBs with aliskiren-containing drugs is contraindicated in patients with moderate to severe renal impairment (GFR < 60 ml/min/1.73m²). (See CONTRAINDICATIONS and DRUG INTERACTIONS, Dual Blockade of the Renin-Angiotensin-System (RAS) with ACEIs, ARBs or aliskiren-containing drugs).

Use of pms-ENALAPRIL should include appropriate assessment of renal function.

Hyperkalemia: Elevated serum potassium (greater than 5.7 mEq/L) was observed in approximately one percent of hypertensive patients in clinical trials with enalapril. In most cases these were isolated values which resolved despite continued therapy. Hyperkalemia was a cause of discontinuation of therapy in 0.28% of hypertensive patients. Risk factors for the development of hyperkalemia include renal insufficiency, diabetes mellitus, and concomitant use of potassium-sparing diuretics (e.g., spironolactone, eplerenone, triamterene, or amiloride), potassium supplements, or potassium-containing salt substitutes. The use of potassium supplements, potassium-sparing diuretics, or potassium-containing salt substitutes particularly in patients with impaired renal function should be given only for documented hypokalemia and with caution and frequent monitoring of serum potassium since they may lead to a significant increase in serum potassium. Hyperkalemia can cause serious, sometimes fatal, arrhythmias. If concomitant use of

pms-ENALAPRIL and any of the above-mentioned agents is deemed appropriate, they should be used with caution and with frequent monitoring of serum potassium (see DRUG INTERACTIONS, Agents Increasing Serum Potassium).

Special Populations

Pregnant Women: ACE inhibitors can cause fetal and neonatal morbidity and mortality when administered to pregnant women. When pregnancy is detected, pms-ENALAPRIL should be discontinued as soon as possible.

The use of ACE inhibitors during the second and third trimesters of pregnancy has been associated with fetal and neonatal injury including hypotension, neonatal skull hypoplasia, anuria, reversible or irreversible renal failure, and death. Oligohydramnios has also been reported, presumably resulting from decreased fetal renal function, associated with fetal limb contractures, craniofacial deformation, and hypoplastic lung development.

Prematurity, and patent ductus arteriosus and other structural cardiac malformations, as well as neurologic malformations, have also been reported following exposure in the first trimester of pregnancy.

Infants with a history of *in utero* exposure to ACE inhibitors should be closely observed for hypotension, oliguria, and hyperkalemia. If oliguria occurs, attention should be directed toward support of blood pressure and renal perfusion. Exchange transfusion or dialysis may be required as a means of reversing hypotension and/or substituting for impaired renal function; however, limited experience with those procedures has not been associated with significant clinical benefit.

Enalapril has been removed from the neonatal circulation by peritoneal dialysis with some clinical benefit and may, theoretically, be removed by exchange transfusion, although there is no experience with the latter procedure.

Animal Data

Maternal and fetal toxicity occurred in some rabbits given enalapril at doses of 1 mg/kg/day or more. Saline supplementation prevented the maternal and fetal toxicity seen at doses of 3 and 10 mg/kg/day, but not at 30 mg/kg/day (50 times the maximum human dose). Enalapril was not teratogenic in rabbits.

There was no fetotoxicity or teratogenicity in rats treated with enalapril at doses up to 200 mg/kg/day (333 times the maximum human dose). Fetotoxicity expressed as a decrease in average fetal weight, occurred in rats given 1200 mg/kg/day of enalapril, but did not occur when these animals were supplemented with saline. The drug crosses the placental barrier in rats and hamsters.

Nursing Women: Enalapril and enalaprilat are secreted in human milk in trace amounts. Use of ACE inhibitors (pms-ENALAPRIL) is not recommended during breast-feeding.

Pediatrics (<16 years of age): The safety and antihypertensive effect have been studied short-term (one month) in patients aged 6 to 16 years (see DOSAGE AND ADMINISTRATION and ACTION AND CLINICAL PHARMACOLOGY, Pharmacokinetics).

pms-ENALAPRIL is not recommended in neonates and in children with glomerular filtration rate <30 mL/min/1.73 m², as no data are available.

ADVERSE REACTIONS

Adverse Drug Reaction Overview

In controlled clinical trials involving 2314 hypertensive patients and 363 patients with congestive heart failure, the most severe adverse reactions were: angioedema (0.2%), hypotension (2.3%) and renal failure (5 cases).

In hypertensive patients, hypotension occurred in 0.9% and syncope in 0.5%, with a discontinuation rate of 0.1%.

In congestive heart failure patients, hypotension occurred in 4.4% and syncope in 0.8%, with a discontinuation rate of 2.5%.

The most frequent clinical adverse reactions in controlled clinical trials were: headache (4.8%), dizziness (4.6%) and fatigue (2.8%). Discontinuation of therapy was required in 6.0% of the 2677 patients.

Clinical Trial Adverse Drug Reactions – Hypertension

Because clinical trials are conducted under very specific conditions the adverse reaction rates observed in the clinical trials may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse drug reaction information from clinical trials is useful for identifying drug-related adverse events and for approximating rates.

Adverse experiences occurring in greater than one percent of patients with hypertension treated with enalapril in controlled clinical trials are shown below. In patients treated with enalapril, the maximum duration of therapy was three years; in placebo treated patients the maximum duration of therapy was 12 weeks.

Table 1 – Hypertension

	Enalapril n=2314	Placebo n=230
Body As A Whole		
Fatigue	3.0	2.6
Orthostatic Effects	1.2	0.0
Asthenia	1.1	0.9
Digestive		
Diarrhea	1.4	1.7
Nausea	1.4	1.7
Nervous/Psychiatric		
Headache	5.2	9.1
Dizziness	4.3	4.3
Respiratory		
Cough	1.3	0.9
Skin		
Rash	1.4	0.4

Less Common Clinical Trial Adverse Drug Reactions (<1%) – Hypertension

Cardiovascular: Hypotension, chest pain, palpitations, acute myocardial infarction

Digestive: Vomiting, dysphagia, abdominal pain

Hematologic: Anemia, Leukopenia

Hypersensitivity: Angioedema

Musculoskeletal: Muscle cramps

Nervous System/Psychiatric: Insomnia, nervousness, somnolence, paresthesia

Respiratory: Dyspnea

Skin: Pruritus, hyperhidrosis

Special Senses: Taste disturbance

Urogenital: Renal failure, proteinuria, oliguria, impotence

Clinical Trial Adverse Drug Reactions – Heart Failure

Because clinical trials are conducted under very specific conditions the adverse reaction rates observed in the clinical trials may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse drug reaction information from clinical trials is useful for identifying drug-related adverse events and for approximating rates.

Adverse experiences occurring in greater than one percent of patients with heart failure treated with enalapril are shown below. The incidences represent the experiences from both controlled and uncontrolled clinical trials (maximum duration of therapy was approximately one year). In the placebo treated patients, the incidences reported are from the controlled trials (maximum duration of therapy is 12 weeks). The percentage of patients with severe heart failure (New York Heart Association [NYHA] Class IV) was 29 percent and 43 percent for patients treated with enalapril and placebo, respectively.

Table 2 – Congestive Heart Failure

	Enalapril n=673	Placebo n=339
Body As A Whole		
Orthostatic Effects	2.2	0.3
Syncope	2.2	0.9
Chest Pain	2.1	2.1
Fatigue	1.8	1.8
Abdominal Pain	1.6	2.1
Asthenia	1.6	0.3
Cardiovascular		
Hypotension	6.7	0.6
Orthostatic Hypotension	1.6	0.3
Angina Pectoris	1.5	1.8
Myocardial Infarction	1.2	1.8
Digestive		
Diarrhea	2.1	1.2
Nausea	1.3	0.6
Vomiting	1.3	0.9
Nervous/Psychiatric		
Dizziness	7.9	0.6
Headache	1.8	0.9
Vertigo	1.6	1.2

	Enalapril n=673	Placebo n=339
Respiratory		
Cough	2.2	0.6
Bronchitis	1.3	0.9
Dyspnea	1.3	0.4
Pneumonia	1.0	2.4
Skin		
Rash	1.3	2.4
Urogenital		
Urinary Tract Infection	1.3	2.4

Less Common Clinical Trial Adverse Drug Reactions (<1%) – Heart Failure

Cardiovascular: Palpitations

Musculoskeletal: Muscle cramps

Nervous System/Psychiatric: Insomnia

Skin: Pruritus

Special Senses: Taste disturbance

Urogenital: Renal failure, impotence

Abnormal Hematologic and Clinical Chemistry Findings

Hyperkalemia: (See WARNINGS AND PRECAUTIONS, Renal)

Creatinine, Blood Urea Nitrogen (BUN): Increases in serum creatinine and BUN were reported in about 20% of patients with renovascular hypertension and in about 0.2% of patients with essential hypertension treated with enalapril alone.

In patients with congestive heart failure, who were also receiving diuretics and/or digitalis, increases in BUN and serum creatinine, usually reversible upon discontinuation of enalapril and/or concomitant therapy, were observed in about 9.7% of patients.

Hemoglobin and Hematocrit: Decreases in hemoglobin and hematocrit (mean approximately 0.34 g% and 1.0 vol%, respectively) occurred frequently in either hypertensive or congestive heart failure patients treated with enalapril, but were rarely of clinical importance. In clinical trials, less than 0.1% of patients discontinued therapy due to anemia.

Hepatic: Elevations of liver enzymes and/or serum bilirubin have occurred (see WARNINGS AND PRECAUTIONS).

Pediatric Patients: In a four-week placebo-controlled clinical trial, 110 hypertensive pediatric patients (6-16 years of age) received medication for 14 days including 51 patients for a four-week period. The adverse experience profile was no different from that seen in adult patients.

Post-Market Adverse Drug Reactions

Adverse Reactions Reported in Uncontrolled Trials and/or Marketing Experience:

Other serious clinical adverse experiences occurring since the drug was marketed or adverse experiences occurring in 0.5 to 1.0 percent of patients with hypertension or heart failure in clinical trials are listed below and, within each category, are in order of decreasing severity.

Body as a Whole

Anaphylactoid reactions (see WARNINGS AND PRECAUTIONS).

Cardiovascular

Cardiac arrest; myocardial infarction or cerebrovascular accident, possibly secondary to excessive hypotension in high-risk patients (see WARNINGS AND PRECAUTIONS); pulmonary embolism and infarction; pulmonary edema; angina pectoris; arrhythmia including atrial tachycardia and bradycardia; atrial fibrillation; palpitation, Raynaud's phenomenon.

Endocrine

Syndrome of inappropriate antidiuretic hormone secretion (SIADH).

Digestive

Ileus, pancreatitis, hepatic failure, hepatitis (hepatocellular or cholestatic jaundice), liver function abnormalities (see WARNINGS AND PRECAUTIONS), melena, anorexia, dyspepsia, constipation, glossitis, stomatitis, dry mouth.

Hematologic

Rare cases of neutropenia, thrombocytopenia, hemolytic anemia and bone marrow depression.

Metabolic

Rare cases of hypoglycemia in diabetic patients on oral antidiabetic agents or insulin have been reported (see WARNINGS AND PRECAUTIONS).

Musculoskeletal

Muscle cramps.

Nervous System/Psychiatric

Vertigo, depression, confusion, ataxia, somnolence, insomnia, nervousness, peripheral neuropathy (e.g., paresthesia, dysesthesia), dream abnormality.

Respiratory

Bronchospasm, rhinorrhea, sore throat and hoarseness, asthma, upper respiratory infection, pulmonary infiltrates, eosinophilic pneumonitis.

Skin

Exfoliative dermatitis, toxic epidermal necrolysis, Stevens-Johnson syndrome, pemphigus, herpes zoster, erythema multiforme, urticaria, pruritus, alopecia, flushing, diaphoresis, photosensitivity.

Special Senses

Blurred vision, taste alteration, anosmia, tinnitus, conjunctivitis, dry eyes, tearing, hearing impairment.

Urogenital

Renal failure, oliguria, renal dysfunction (see WARNINGS AND PRECAUTIONS and DOSAGE AND ADMINISTRATION), flank pain, gynecomastia, impotence.

A symptom complex has been reported which may include some or all of the following: fever, serositis, vasculitis, myalgia/myositis, arthralgia/arthritis, a positive antinuclear antibody (ANA), elevated erythrocyte sedimentation rate, eosinophilia and leukocytosis. Rash, photosensitivity or other dermatologic manifestations may occur. These symptoms may be reversible upon discontinuation of therapy.

In very rare cases, intestinal angioedema has been reported with angiotensin converting enzyme inhibitors including enalapril.

Laboratory Test Findings: Hyponatremia

DRUG INTERACTIONS

Drug-Drug Interactions

Hypotension – Patients on Diuretic Therapy: Patients on diuretics and especially those in whom diuretic therapy was recently instituted, may occasionally experience an excessive reduction of blood pressure after initiation of therapy with enalapril. The possibility of hypotensive effects with enalapril can be minimized by either discontinuing the diuretic or increasing the salt intake prior to initiation of treatment with enalapril (see WARNINGS AND PRECAUTIONS and DOSAGE AND ADMINISTRATION).

Agents Increasing Serum Potassium: Since enalapril decreases aldosterone production, elevation of serum potassium may occur. Potassium-sparing diuretics such as spironolactone, eplerenone, triamterene or amiloride, or potassium supplements should be given only for documented hypokalemia and with caution and frequent monitoring of serum potassium particularly in patients with impaired renal function since they may lead to a significant increase in serum potassium. Salt substitutes which contain potassium should also be used with caution (see WARNINGS AND PRECAUTIONS, Hyperkalemia).

Agents Causing Renin Release: The antihypertensive effect of enalapril is augmented by antihypertensive agents that cause renin release (e.g., diuretics).

Agents Affecting Sympathetic Activity: Agents affecting sympathetic activity (e.g., ganglionic blocking agents or adrenergic neuron blocking agents) may be used with caution. Beta-adrenergic blocking drugs add some further antihypertensive effect to enalapril.

Lithium Salts: As with other drugs which eliminate sodium, lithium clearance may be reduced. Therefore, the serum lithium levels should be monitored carefully if lithium salts are to be administered.

Non-Steroidal Anti-Inflammatory Drugs (NSAIDs) Including Selective Cyclooxygenase-2 Inhibitors: The antihypertensive effect of enalapril may be diminished with concomitant non-steroidal anti-inflammatory drug use including selective cyclooxygenase-2 inhibitors (COX-2 inhibitors). In some patients with compromised renal function (e.g., elderly patients or patients who are volume-depleted including those on diuretic therapy) who are being treated with NSAIDs including selective COX-2 inhibitors, the co-administration of ACE inhibitors or angiotensin II receptor antagonists may result in further deterioration of renal function. Cases of

acute renal failure, usually reversible, have also been reported. This combination should therefore be administered with caution in this patient population.

Dual Blockade of the Renin-Angiotensin System (RAS) with ACEIs, ARBs or aliskiren-containing drugs: Dual Blockade of the Renin-Angiotensin System (RAS) with ACEIs, ARBs or aliskiren-containing drugs is contraindicated in patients with diabetes and/or renal impairment, and is generally not recommended in other patients, since such treatment has been associated with an increased incidence of severe hypotension, renal failure, and hyperkalemia. See CONTRAINDICATIONS, WARNINGS AND PRECAUTIONS, Cardiovascular, Dual Blockade of the Renin-Angiotensin System (RAS) and Renal.

Gold: Nitritoid reactions (symptoms include facial flushing, nausea, vomiting and symptomatic hypotension) have been reported rarely in patients on therapy with injectable gold (sodium aurothiomalate) and concomitant ACE inhibitor therapy including enalapril (see WARNINGS AND PRECAUTIONS).

Mammalian Target of Rapamycin (mTOR) Inhibitors: Patients taking concomitant mTOR inhibitor (e.g., temsirolimus, sirolimus, everolimus) therapy may be at increased risk for angioedema. Caution should be used when these drugs are used concomitantly (see WARNINGS AND PRECAUTIONS).

Drug-Food Interactions

The absorption of enalapril is not influenced by the presence of food in the gastrointestinal tract.

DOSAGE AND ADMINISTRATION

pms-ENALAPRIL for Oral Administration Only

Dosing Considerations

- The absorption of pms-ENALAPRIL is not affected by food.
- Dosage must be individualized.
- Special attention for dialysis patients.
- The splitting of pms-ENALAPRIL tablets is not advised.

Recommended Dose and Dosage Adjustment

Hypertension: Initiation of therapy requires consideration of recent antihypertensive drug treatment, the extent of blood pressure elevation and salt restriction; the dosage of other antihypertensive agents being used with pms-ENALAPRIL may need to be adjusted.

The recommended initial dose in patients not on diuretics is 5 mg once a day. Dosage should be adjusted according to blood pressure response. The usual dosage range is 10 to 40 mg per day administered in a single dose or two divided doses. In some patients treated once daily, the antihypertensive effect may diminish toward the end of the dosing interval. In such patients, an increase in dosage or twice-daily administration should be considered. If blood pressure is not controlled, a diuretic may be added.

The maximum daily dose is 40 mg. Raising the dose above that level is not recommended because of the possibility of increased adverse reactions.

Symptomatic hypotension occasionally may occur following the initial dose of pms-ENALAPRIL and is more likely in patients who are currently being treated with a diuretic. The diuretic should, if possible, be discontinued for two to three days before beginning therapy with pms-ENALAPRIL to reduce the likelihood of hypotension (see WARNINGS AND PRECAUTIONS).

If the diuretic cannot be discontinued, an initial dose of 2.5 mg should be used to determine whether excessive hypotension occurs.

To date there is insufficient experience with pms-ENALAPRIL in the treatment of accelerated or malignant hypertension. pms-ENALAPRIL, therefore, is not recommended in such situations.

Pediatrics (<16 years of age): The usual recommended starting dose is 0.08 mg/kg (up to 5 mg) once daily. Dosage should be adjusted according to blood pressure response. Doses above 0.58 mg/kg (or in excess of 40 mg) have not been studied in pediatric patients (see ACTION AND CLINICAL PHARMACOLOGY). pms-ENALAPRIL is not recommended in neonates and in pediatric patients with glomerular filtration rate <30 mL/min/1.73 m², as no data are available.

Geriatrics (>65 years of age): The starting dose should be 2.5 mg. Some elderly patients may be more responsive to pms-ENALAPRIL than younger patients.

Dosing Adjustment in Renal Impairment: (See WARNINGS AND PRECAUTIONS, Anaphylactoid Reactions during Membrane Exposure)

The doses should be reduced in patients with hypertension according to the following guidelines:

Renal Status	Creatinine Clearance mL/min (mL/s)	Initial Dose mg/day
Normal Renal Function	>80 mL/min (>1.33 mL/s)	5 mg
Mild Impairment	≤80 >30 mL/min (≤1.33 > 0.50 mL/s)	5 mg
Moderate to Severe Impairment	≤30 mL/min (≤0.50 mL/s)	2.5 mg
Dialysis Patients	-	2.5 mg on dialysis days*

* Enalaprilat is dialyzable. Dosage on nondialysis days should be adjusted depending on the blood pressure response.

Congestive Heart Failure: Enalapril is generally used in conjunction with a diuretic and/or digitalis. Blood pressure and renal function should be monitored, both before and during treatment with pms-ENALAPRIL, because severe hypotension and, more rarely, consequent renal failure have been reported (see WARNINGS AND PRECAUTIONS).

Initiation of therapy requires consideration of recent diuretic therapy and the possibility of severe salt/volume depletion. If possible, the dose of diuretic should be reduced before beginning treatment to reduce the likelihood of hypotension. Serum potassium also should be monitored (see DRUG INTERACTIONS, Drug-Drug Interactions).

The recommended initial dose in patients with symptomatic heart failure or asymptomatic left ventricular dysfunction (ejection fraction $\leq 35\%$) is 2.5 mg once a day, to be administered under close medical supervision to determine the initial effect on blood pressure. After the initial dose, the patient should be observed for at least two hours or until the pressure has stabilized for at least another additional hour (see WARNINGS AND PRECAUTIONS, Hypotension).

In the absence of, or after effective management of symptomatic hypotension following initiation of therapy, the dose should be increased gradually depending on the patient's response. The usual therapeutic dosing range is 5 to 20 mg daily, given as a single dose or two divided doses.

This dose titration may be performed over a 2 to 4 week period, or more rapidly if indicated by the presence of residual signs and symptoms of heart failure. The dosage regimen, in patients with symptomatic heart failure, which was effective in reducing mortality and the need for hospitalization in multicentre studies ranged between 16.4 and 18.8 mg/day. The majority of patient experience in clinical studies has been with twice-daily dosage.

The maximum daily dose is 40 mg.

Dosage Adjustment in Patients with Congestive Heart Failure and Renal Impairment or Hyponatremia: In patients with heart failure who have hyponatremia (serum sodium less than 130 mEq/L) or with serum creatinine greater than 1.6 mg/dL, therapy should be initiated at 2.5 mg daily under close medical supervision (see DOSAGE AND ADMINISTRATION, Congestive Heart Failure and DRUG INTERACTIONS, Drug-Drug Interactions).

The dose may be increased to 2.5 mg b.i.d. then 5 mg b.i.d. and higher as needed, usually at intervals of four days or more if at the time of dosage adjustment there is not excessive hypotension or significant deterioration of renal function.

The maximum daily dose is 40 mg.

OVERDOSAGE

Limited data are available for overdosage in humans. The most prominent features of overdosage reported to date are marked hypotension, beginning some six hours after ingestion of tablets, concomitant with blockade of the renin-angiotensin system, and stupor. Serum enalaprilat levels 100- and 200-fold higher than usually seen after therapeutic doses have been reported after ingestion of 300 mg and 440 mg of enalapril, respectively.

The recommended treatment of overdosage is intravenous infusion of normal saline solution. If ingestion is recent, induce emesis. Enalaprilat may be removed from the general circulation by hemodialysis (see WARNINGS AND PRECAUTIONS, Anaphylactoid Reactions during Membrane Exposure).

For management of a suspected drug overdose, contact your regional
Poison Control Centre immediately.

ACTION AND CLINICAL PHARMACOLOGY

Mechanism of Action

Enalapril is an ACE inhibitor which is used in the treatment of hypertension and heart failure.

Angiotensin converting enzyme (ACE) is a peptidyl dipeptidase which catalyzes the conversion of angiotensin I to the pressor substance, angiotensin II. After absorption, enalapril, a pro-drug, is hydrolyzed to enalaprilat, its active metabolite, which inhibits ACE. Inhibition of ACE results in decreased plasma angiotensin II, which leads to increased plasma renin activity (due to removal of negative feedback of renin release) and decreased aldosterone secretion. Although the latter decrease is small, it results in a small increase in serum potassium. In patients treated with enalapril and a thiazide diuretic there was essentially no change in serum potassium (see WARNINGS AND PRECAUTIONS).

ACE is identical to kininase II. Thus, enalapril may also block the degradation of bradykinin, a potent vasodepressor peptide. However, the role that this plays in the therapeutic effects of either drug is unknown.

While the mechanism through which enalapril lowers blood pressure is believed to be primarily the suppression of the renin-angiotensin-aldosterone system, enalapril also lowers blood pressure in patients with low-renin hypertension.

Pharmacodynamics

Administration of enalapril to patients with hypertension results in a reduction of both supine and standing blood pressure. Abrupt withdrawal of enalapril has not been associated with a rapid increase in blood pressure. In most patients studied, after oral administration of an individual dose of enalapril, the onset of antihypertensive activity is seen at one hour with peak reduction of blood pressure achieved by 4-6 hours. At recommended doses, the antihypertensive effect has been shown to be maintained for at least 24 hours. In some patients the effect may diminish towards the end of the dosing interval (see DOSAGE AND ADMINISTRATION). On occasion, achievement of optimal blood pressure reduction may require several weeks of therapy.

In hemodynamic studies in patients with essential hypertension, blood pressure reduction was accompanied by a reduction in peripheral arterial resistance with an increase in cardiac output and little or no change in heart rate. Following administration of enalapril, there was an increase in renal blood flow; glomerular filtration rate was usually unchanged.

When enalapril is given together with thiazide-type diuretics, its blood pressure lowering effect is approximately additive.

Administration of enalapril to patients with congestive heart failure reduces afterload and preload of the heart, resulting in an increase in cardiac output, without reflex tachycardia.

When used in hypertensive, normolipidemic patients, enalapril had no effect on plasma lipoprotein fractions.

Studies in dogs indicate that enalapril crosses the blood brain barrier poorly, if at all; enalaprilat does not enter the brain.

Pharmacokinetics

Table 3 – Summary of Enalaprilat’s Pharmacokinetic Parameters in Healthy Volunteers Further to a 10 mg Oral Dose of Enalapril

	C_{max} ng/mL	t_{1/2} (h)*	AUC_{0-∞} ng.h/mL
Single dose mean	32.3	11	423

* Effective half-life of accumulation.

Absorption: Enalapril is rapidly absorbed with peak serum concentrations of enalapril occurring within one hour. Based on urinary recovery the extent of absorption of enalapril is approximately 60%. The absorption of enalapril is not influenced by the presence of food in the gastrointestinal tract.

Metabolism: Following absorption, enalapril is rapidly and extensively hydrolyzed to enalaprilat, a potent angiotensin converting enzyme inhibitor (which itself is poorly absorbed). Peak serum concentrations of enalaprilat occur 3 to 4 hours after an oral dose of enalapril. Except for conversion to enalaprilat, there is no evidence of significant metabolism of enalapril.

Excretion: Excretion of enalapril is primarily renal. Approximately 94% of the dose is recovered in the urine and feces as enalaprilat or enalapril. The principal components in urine are enalaprilat, accounting for about 40% of the dose, and intact enalapril.

The serum concentration profile of enalaprilat exhibits a prolonged terminal phase, apparently associated with binding to ACE. The effective half-life for accumulation of enalaprilat following multiple doses of enalapril is 11 hours.

In hypertensive children aged 2 months to 15 years the kinetics of enalapril were approximately similar to adults (see DOSAGE AND ADMINISTRATION).

Special Populations and Conditions

Pediatrics: In pediatric patients the antihypertensive effect of enalapril has been studied in hypertensive children aged 6–16 years (see DOSAGE AND ADMINISTRATION and ACTION AND CLINICAL PHARMACOLOGY, Pharmacokinetics).

Race: The antihypertensive effect of angiotensin converting enzyme inhibitors is generally lower in black than in non-black patients.

Renal Insufficiency: The disposition of enalapril and enalaprilat in patients with renal insufficiency is similar to that in patients with normal renal function until the glomerular filtration rate is 30 mL/min (0.50 mL/s) or less. With renal function ≤ 30 mL/min (≤ 0.50 mL/s), peak and trough enalaprilat levels increase, time to peak concentration increases and time to steady state

may be delayed. The effective half-life of enalaprilat following multiple doses of enalapril is prolonged at this level of renal insufficiency (see DOSAGE AND ADMINISTRATION). Enalaprilat is dialyzable at the rate of 62 mL/min (1.03 mL/s).

STORAGE AND STABILITY

Store between 15°C and 30°C. Protect from moisture.

NOTE: 100 and 500 tablet bottle: discard remaining tablets six months after opening bottle.

DOSAGE FORMS, COMPOSITION AND PACKAGING

Tablets

2.5 mg Each white, oval-shaped tablet, debossed with “P” letter on one side of the score line and “2.5” on the other side of the score line and nothing on the other side, contains 2.5 mg of enalapril maleate, which appears as 2 mg of enalapril sodium, and the following non medicinal ingredients: copovidone, lactose, pregelatinized starch, sodium bicarbonate and zinc stearate. Available in HDPE bottles of 100 tablets and 500 tablets.

5 mg Each white, shield-shaped tablet, debossed with “P” letter and “5” on one side and scored on the other side, contains 5 mg of enalapril maleate, which appears as 4 mg of enalapril sodium, and the following non medicinal ingredients: copovidone, lactose, pregelatinized starch, sodium bicarbonate and zinc stearate. Available in HDPE bottles of 100 tablets and 500 tablets.

10 mg Each salmon, shield-shaped tablet, debossed with “P” letter and “10” on one side and scored on the other side, contains 10 mg enalapril maleate, which appears as 8 mg of enalapril sodium, and the following non medicinal ingredients: copovidone, lactose, pregelatinized starch, red iron oxide, sodium bicarbonate and zinc stearate. Available in HDPE bottles of 100 tablets and 500 tablets.

20 mg Each peach, shield-shaped tablet debossed with “P” letter and “20” on one side and scored on the other side, contains 20 mg of enalapril maleate, which appears as 16 mg of enalapril sodium, and the following non medicinal ingredients: copovidone, lactose, red iron oxide, pregelatinized starch, sodium bicarbonate, yellow iron oxide and zinc stearate. Available in HDPE bottles of 100 tablets and 500 tablets.

The splitting of pms-ENALAPRIL tablets is not advised.

PART II: SCIENTIFIC INFORMATION

PHARMACEUTICAL INFORMATION



Drug Substance

Proper Name:	Enalapril maleate	Enalapril sodium
Chemical Name:	L-Proline, 1-[N-[1-(ethoxycarbonyl)-3-phenylpropyl]-L-alanyl]-, (s)-, (Z)-2-butenedioate (1:1)	L-Proline, 1-[N-[1-(ethoxycarbonyl)-3-phenylpropyl]-L-alanyl]-, (s)-, Sodium (1:1)

Molecular Formula:	$C_{20}H_{28}N_2O_5 \cdot C_4H_2O$	$C_{20}H_{27}N_2O_5Na$
---------------------------	------------------------------------	------------------------

Molecular Mass:	492.53 g/mol	398.43 g/mol
------------------------	--------------	--------------

Structural Formula:



Physicochemical Properties:	Enalapril maleate is a white to off-white crystalline powder which melts at $\sim 143^{\circ}C$ to $144^{\circ}C$. It is sparingly soluble in water (pH 3.4), soluble in ethanol, and freely soluble in methanol and dimethylformamide. The pK_a^1 and pK_a^2 of the base moiety are 3.0 and 5.4, respectively.
------------------------------------	--

CLINICAL TRIALS

Comparative Bioavailability Studies

A randomized two-way crossover, blinded, single-dose comparative bioavailability study was performed in normal healthy non-smoking male volunteers (n=24) under fasting conditions on Enalapril tablets using Pharmascience Inc. pms-ENALAPRIL 5 mg tablets versus the reference product, VASOTEC[®] 5 mg Tablets, by Merck Frosst Canada Ltd. The pharmacokinetic data calculated for the pms-ENALAPRIL 5 mg tablets and VASOTEC[®] 5 mg tablets formulation are tabulated below. The number of subjects that were included in the PK analysis was 23.

SUMMARY TABLE OF THE COMPARATIVE BIOAVAILABILITY DATA

Enalapril (1 x 5 mg) From measured data				
Geometric Mean Arithmetic Mean (CV %)				
Parameter	Test*	Reference†	% Ratio of Geometric Means‡	90% Confidence Interval
AUC _T (ng·h/mL)	50.82 53.66 (35.39)	51.20 55.46 (39.60)	98.95	92.72 – 105.60
AUC _I (ng·h/mL)	51.43 54.22 (34.96)	50.68 54.74 (39.35)	98.73	92.33 – 105.57
C _{max} (ng/mL)	33.28 35.26 (35.78)	33.19 36.67 (45.92)	99.81	92.63 – 107.54
T _{max} [§] (h)	0.75 (0.75-1.25)	0.75 (0.75-1.50)		
T _½ [€] (h)	1.36 (88.08)	1.26 (48.57)		

* pms-ENALAPRIL

† Vasotec[®] Merck Frosst, Canada; 5 mg tablets were purchased in Canada.

§ Expressed as the median (range) only

€ Expressed as the arithmetic mean (CV%) only

‡ The ratios were calculated using the least square mean differences

A randomized two-way crossover, blinded, single-dose comparative bioavailability study was performed in normal healthy non-smoking male volunteers (n=24) under fasting conditions on Enalapril tablets using Pharmascience Inc. pms-ENALAPRIL 20 mg tablets versus the reference product, VASOTEC[®] 20 mg Tablets, by Merck Frosst Canada Ltd. The pharmacokinetic data calculated for the pms-ENALAPRIL 20 mg tablets and VASOTEC[®] 20 mg tablets formulation are tabulated below. The number of subjects that were included in the PK analysis was 22.

SUMMARY TABLE OF THE COMPARATIVE BIOAVAILABILITY DATA

Enalapril (1 x 20 mg) From measured data Geometric Mean Arithmetic Mean (CV %)				
Parameter	Test[*]	Reference[†]	% Ratio of Geometric Means[‡]	90% Confidence Interval
AUC _T [‡] (ng*h/mL)	162.40 169.4 (31.51)	156.58 161.60 (27.45)	103.64	98.06 %-109.53 %
AUC _I (ng*h/mL)	163.22 170.31 (31.37)	158.01 163.26 (27.45)	102.68	97.25 %-108.41 %
C _{max} (ng/mL)	112.48 117.40 (30.00)	105.98 109.65 (26.92)	105.96	98.38 %-114.13 %
T _{max} [§] (h)	0.75 (0.5-1.0)	0.75 (0.5-1.25)		
T _{1/2} [€] (h)	1.67 (70.42)	1.59 (58.67)		

* pms-ENALAPRIL

† Vasotec[®] Merck Frosst, Canada; 20 mg tablets were purchased in Canada.

§ Expressed as the median (range) only

€ Expressed as the arithmetic mean (CV%) only

‡ The ratios were calculated using the least square mean differences

Study demographics and trial design

Table 4 – Summary of patient demographics for clinical trials

Study	Trial design	Dosage, route of administration and duration	Study subjects (n=number)	Mean age (Range)	Gender
SOLVD – Treatment	Multicentre, randomized, double-blind, placebo-controlled	2.5 or 5 mg twice daily Titrated up to 10 mg twice daily up to 55 months	2569 Placebo: 1284 Enalapril: 1285	60.2 years (20-80)	Male: 2065 Female: 504

Study results

Table 5 – Results of SOLVD-Treatment study

Primary Endpoints	Enalapril N (%)	Placebo N (%)	p-Value
Overall mortality	451 (35.1%)	510 (39.7%)	0.008

In a multicentre, placebo-controlled, double-blind study (SOLVD, see REFERENCES, no. 23), 2569 patients with symptomatic heart failure (primarily New York Heart Association Class II and III and ejection fraction $\leq 35\%$), were randomized to placebo or enalapril given as an adjunct to conventional therapy. Diseases that excluded patients from enrolment in the study included severe stable angina, hemodynamically significant valvular or outflow tract obstruction, renal failure, cerebral vascular disease (e.g., significant carotid artery diseases), advanced pulmonary disease, malignancies, active myocarditis and constrictive pericarditis. The use of enalapril was associated with an 11% reduction in all-cause mortality (which corresponds to a 16% risk reduction in all-cause mortality) and a 30% reduction in hospitalization for heart failure (which corresponds to a 36% risk reduction in hospitalization for heart failure). The chief difference in mortality was in deaths due to progressive heart failure. There was no significant difference in the number of deaths classified as due to arrhythmia without worsening congestive heart failure.

Study demographics and trial design

Table 6 – Summary of patient demographics for clinical trials

Study	Trial design	Dosage, route of administration and duration	Study subjects (n=number)	Mean age (Range)	Gender
SOLVD – Prevention	Multicentre, randomized, double-blind, placebo-controlled	2.5 mg twice daily gradually increased to 10 mg twice daily Follow-up for a minimum of 46 months to a maximum of 62 months	4228 Placebo: 2117 Enalapril: 2111	58.7 years (20-80)	Male: 3752 Female: 476

Study results

Table 7 – Results of SOLVD-Prevention study

Primary Endpoints	Enalapril N (%)	Placebo N (%)	p-Value
Overall mortality	306 (14.5%)	332 (15.7%)	0.211

A second multicentre trial used the SOLVD protocol for a study of asymptomatic or minimally symptomatic patients. SOLVD-Prevention patients, who had left ventricular ejection fraction $\leq 35\%$ and no history of symptomatic heart failure, were randomized to placebo (n=2117) or enalapril (n=2111) and followed for up to 5 years. The majority of patients in the SOLVD-Prevention trial had a history of ischemic heart disease. A history of myocardial infarction was present in 80% of patients, current angina pectoris in 34%, and a history of hypertension in 37%. No statistically significant mortality effect was demonstrated in this population. Enalapril-treated subjects had 32% fewer first hospitalizations for heart failure, and 32% fewer total heart failure hospitalizations. Compared to placebo, 32% fewer patients receiving enalapril developed symptoms of overt heart failure.

Hospitalizations for cardiovascular reasons were also reduced. There was an insignificant reduction in hospitalizations for any cause in the enalapril treatment group (for enalapril vs. placebo, respectively, 1166 vs. 1201 first hospitalizations, 2649 vs. 2840 total hospitalizations), although the study was not powered to look for such an effect.

The SOLVD-Prevention trial was not designed to determine whether treatment of asymptomatic patients with low ejection fraction would be superior, with respect to preventing hospitalization, to closer follow-up and use of enalapril at the earliest sign of heart failure. However, under the conditions of follow-up in the SOLVD-Prevention trial (every four months at the study clinic; personal physician as needed), 68% of patients on placebo who were hospitalized for heart failure had no prior symptoms recorded which would have signaled initiation of treatment.

The SOLVD-Prevention trial was also not designed to show whether enalapril modified the progression of underlying heart disease.

Study demographics and trial design

Table 8 – Summary of patient demographics for clinical trials

Study	Trial design	Dosage, route of administration and duration	Study subjects (n=number)	Mean age (Range)	Gender
CONSENSUS	Multicentre, randomized, double-blind, placebo-controlled	5 mg twice daily increased to a maximum of 20 mg twice daily	253 Placebo: 126 Enalapril: 127	70 years (36-91)	Male: 179 Female: 74

Study results

Table 9 – Results of CONSENSUS study

Primary Endpoints		Enalapril N (%)	Placebo N (%)	p-Value
Overall mortality	6 months	33 (26.0%)	55 (43.6%)	0.004
	12 months	46 (36.2%)	66 (52.4%)	0.011
	End of study	50 (39.4%)	68 (54.0%)	0.003

In another multicentre, placebo-controlled trial (CONSENSUS, see BIBLIOGRAPHY, no. 22), 253 patients with severe congestive heart failure (New York Heart Association Class IV) were randomized to placebo or enalapril given as an adjunct to conventional therapy. The use of enalapril was associated with an improvement of symptoms and a reduction in mortality from the progression of heart failure. No difference was seen in the incidence of sudden cardiac death.

DETAILED PHARMACOLOGY

Mechanism of Action

Study	Species/Strain	No. of Animals/Group	Route	Dose	Results
Effect of enalapril maleate on total serum ACE in rats and dogs	Male Sprague/Dawley Rats	12 experimental 6 placebo	P.O.	10 mg/kg/day for 7 or 14 days	79% increase in ACE after 7 days & 140% after 14 days
	Male beagle hounds	3 dogs	P.O.	10 mg/kg (free base) for 7 or 14 days	30% increase in ACE after 7 days & 48% after 14 days
		3 dogs	P.O.	30 mg/kg/day for 3 days	1.5-fold increase in ACE
<i>In vivo</i> ACE inhibition in anesthetized and unanesthetized rats and dogs	Male Sprague/Dawley rats (Blue Spruce)	6 rats	I.V. P.O.	3, 10, 30 mcg/kg 0.1, 0.3, 1.0 and 3.0 mg/kg	The ED ₅₀ is 14.0 mcg/kg I.V. and 0.29 mg/kg p.o.

Study	Species/Strain	No. of Animals/Group	Route	Dose	Results
	Mongrel or beagle dogs (male & female)	6 dogs per dose	I.V.	30, 130, 430, 1430 mcg/kg	Dose-related inhibition of pressor response to angiotensin
					ED ₅₀ : Enalaprilat: 6.4 mcg/kg. Enalapril maleate: 278 mcg/kg.
Effect of enalaprilat on canine hind limb vasodilator response to bradykinin and vasoconstrictor response to angiotensins	Anesthetized dogs male or female	4 dogs	I.V.	0.3-100 mcg/kg	Local inhibition of ACE: (enalaprilat) ED ₅₀ = 4.8 (4.4 to 5.2 mcg/kg) I.V.

Effects on Blood Pressure

Study	Species/Strain	No. of Animals/Group	Route	Dose	Results
Antihypertensive activity in sodium-deficient rats	Male Sprague/Dawley rats	6 rats/group and at least 8 treatment groups	P.O.	Enalapril 1 to 10 mg/kg	Enalapril produced a dose-dependent decrease in systolic BP for 3 or more hours
Effect on renal hypertensive rats (Grollman technique)	Male Sprague/Dawley rats	Most groups=6 to 8 rats/treatment group	P.O.	Enalapril 3.0 mg/kg	Enalapril produced a mean decrease in systolic pressure of ≈20 mmHg and a slight tachycardia
Relationship between angiotensin I blockade and blood pressure lowering in spontaneous hypertensive rats, renal hypertensive rats, and renal hypertensive dogs and normotensive sodium depleted dogs	Sprague/Dawley rats Normotensive dogs (mongrel)	At least 4 to 5 rats/group and at least 3 dogs per group	P.O.	Enalapril 0.1 to 3 mg/kg	Time course of blood pressure decrease did not coincide with time course for maximal inhibition of angiotension I pressor response

Other Effects

Study	Species/Strain	No. of Animals/Group	Route	Dose	Results
Effects in acute renal failure in dogs	Mongrel dogs	4/group	P.O.	1.0 mg/kg b.i.d for 3 days	No further deterioration of acute renal failure occurred
Whole body autoradiography	Golden hamsters	Min. 16	P.O.	5 mg/kg	No radioactivity was found in the spinal cord or brain of either male or female hamsters

TOXICOLOGY

Acute Toxicity

LD₅₀ Values:

Route	Species	Sex	MSDRL ^a	NMB/RL ^b
Oral	Mouse	Male	2 g/kg	3.5 g/kg
		Female	2 g/kg	3.5 g/kg
	Rat	Male	2 g/kg	3.5 g/kg
		Female	2 g/kg	3.0 g/kg
Intravenous	Mouse	Male	-	900 mg/kg
		Female	750 mg/kg	900 mg/kg
	Rat	Male	-	950 mg/kg
		Female	-	850 mg/kg
Subcutaneous	Mouse	Male	-	1150 mg/kg
		Female	-	1500 mg/kg
	Rat	Male	-	1750 mg/kg
		Female	-	1400 mg/kg

^a Merck Sharp and Dohme Research Laboratories, West Point, PA, USA

^b Nippon Merck-Banyu Co., Menuuma, Japan

Signs of toxicity: ptosis, decreased activity, bradypnea, loss of righting, ataxia, dyspnea, and clonic convulsions.

Sub-Acute and Chronic Toxicity

Species	Duration	No. of Animals/Group	Route	Dose mg/kg/day	Effects
Rat	1 Month	10 M + 10 F	Oral	0, 10, 30, 90	At all doses: Slight decrease in body weight gain.

Species	Duration	No. of Animals/Group	Route	Dose mg/kg/day	Effects
					At 30 & 90 mg/kg/day: Dose-related increase in BUN in males.
Rat	3 Months	15 M + 15 F	Oral	0, 10, 30, 90	At all doses: Slight decrease in body weight gain and in serum sodium, slight increase in serum potassium. Small increase in kidney weight and decrease in heart weight. At 30 & 90 mg/kg/day: Dose-related increase in BUN.
Rat	1 Year	25 M + 25 F	Oral	0, 10, 30, 90	6-month interim kill: Males given 90 mg/kg/day had a significantly ($P \leq 0.05$) greater kidney weight than controls. 1 year: Dose-related decrease in weight gain (7 to 19%). Dose-related increase in serum urea nitrogen in males given 30 and 90 mg/kg/day (values up to 52.9 and 89.2 mg/100 mL, respectively). Three high dose females showed elevated serum urea nitrogen levels. Serum potassium values were increased (0.1 to 0.8 mEq/L) in male rats on the high dose. Males given 90 mg/kg/day had a significantly ($P \leq 0.05$) greater kidney weight than controls.
Rat	1 Month	20 M + 20 F	Oral	0, 90 & 90 with physiologic saline for drinking	Unsupplemented: Less weight gain (8 to 19%), increase in serum urea nitrogen (up to 62.8 mg%). Supplemented: Body weight gain and serum urea nitrogen levels similar to controls.
Rat (sodium depleted)	3 Weeks	30 M + 30 F	Oral	0, 90	A marked potentiation in toxicity included: death, weight loss, marked increases in serum urea nitrogen, creatinine and potassium, renal tubular degeneration.
Dog Beagle	1 Month	3M + 3F	Oral	0, 10, 30, 90 (4 doses only) reduced to 60 (4 doses only)	At 30 mg: One dog showed increase in BUN and renal tubular degeneration. At high doses: 6/6: deaths (7-12 days)

Species	Duration	No. of Animals/Group	Route	Dose mg/kg/day	Effects
					Increase in serum urea nitrogen, glucose, SGOT, SGPT, and potassium; decrease in serum sodium and chloride; renal tubular degeneration and increased hepatocellular fat.
Dog Beagle	3 Months	3 M + 3 F	Oral	0, 10, 30, 90 (7 doses only)	<p>At All Doses: Slight decrease in serum sodium.</p> <p>At 30 mg: 2/6: deaths Increase in BUN and serum glucose; renal tubular degeneration.</p> <p>At 90 mg: 5/6: deaths Increase in BUN, serum glucose, SGOT, SGPT, alkaline phosphatase and potassium. Decrease in serum chloride; renal tubular degeneration, increased hepatocellular fat; hepatocellular necrosis.</p>
Dog Beagle	1 Year	5 M + 5 F	Oral	0, 3, 5, 15	No drug-induced changes were seen.
Dog Beagle	15 Days	3 M + 3 F	Oral	0, 60 with and without saline supplementation	<p>Unsupplemented treated dogs: 3/6: deaths 4/6: increase in serum urea nitrogen 3/6: decrease in serum chloride increase in SGOT, SGPT and potassium 1/6: increase in alkaline phosphatase 1/6: hepatocellular lesions (in 1st animal which died) 5/6: renal lesions (3 moderate, 2 slight renal tubular necrosis).</p> <p>Saline supplemented treated dogs: 0/6: deaths 3/6: increase in serum urea nitrogen 1/6: very slight renal tubular necrosis and moderate renal tubular cell vacuolation</p>
Dog Beagle	15 Days	3 M + 3 F	Oral	0, 90 with and without saline supplementation	<p>Unsupplemented treated dogs: 6/6: deaths 6/6: increase in serum urea nitrogen, creatinine and SGPT 5/6: increase in SGOT 2/6: increase in serum potassium 5/6: marked renal tubular degeneration 1/6: moderate renal tubular degeneration</p>

Species	Duration	No. of Animals/Group	Route	Dose mg/kg/day	Effects
					6/6: slight to marked thymic atrophy 3/6: ulceration of distal esophagus 2/6: oral mucosal lesions Supplemented treated dogs: 2/6: deaths 6/6: increase in serum urea nitrogen, creatinine 3/6: increase in SGOT and SGPT 0/6: increase in potassium 2/6: moderate renal tubular degeneration 4/6: slight renal tubular degeneration 4/6: slight to moderate thymic atrophy 3/6: liver degeneration

Teratology Studies

Species	No. of Animals/Group	Dose mg/kg/day	Duration of Dosing	Results
Rat (Charles River CD)	20 F	0, 10, 30, 90	Day 15 of gestation through Day 20 of lactation	At all dosage levels: <ul style="list-style-type: none"> - Decreased maternal weight gain during days 15 –20 - Dose-related retardation in growth of F1 offspring during lactation At 90 mg/kg/day: <ul style="list-style-type: none"> - Mean Day 1 pup weight/litter was significantly less than that of controls
Rat (Charles River CD)	25 F	0, 10, 100, 200 100 + saline, 200 + saline	Days 6 through Day 17 of gestation	Decreased maternal weight gain at 100 and 200 mg/kg/day in unsupplemented rats. No treatment-related effects on reproductive status or teratogenic effects in any of the groups.
Rat (CLEA Japan Inc-JCL :SD)	25F	0, 12, 120, 1200, 1200 + saline	Days 6 through Day 17 of gestation	Unsupplemented treated rats: <ul style="list-style-type: none"> - Average maternal body weight gain significantly reduced at all doses. At 1200 mg/kg/day: <ul style="list-style-type: none"> - Slight but significant decrease in fetal weight - Increase in the number of fetuses with the 14th rib skeletal variation - Decrease in the number of fetuses with ossified caudal vertebrae Supplemented treated rats:

Species	No. of Animals/Group	Dose mg/kg/day	Duration of Dosing	Results
				- No evidence of maternotoxicity or fetotoxicity
Rabbit (New Zealand albino)	18 F	0, 3, 10, 30 (with saline)	Days 6 through Day 18 of gestation	<p>At 3 and 10 mg/kg/day:</p> <ul style="list-style-type: none"> - No treatment-related effects on reproductive status or teratogenicity was observed <p>At 30 mg/kg/day:</p> <ul style="list-style-type: none"> - 4 deaths - Reduced food and water intake - Significant increase in the mean number of resorptions per litter - 2 abortions - No evidence of teratogenicity was observed

Fertility and Postnatal Evaluation Studies

Species	No. of Animals/Group	Dose mg/kg/day	Duration of Dosing	Results
Rat (Charles River CD)	15 M + 30 F	0, 10, 30, 90	<p>Males 70 days prior to mating to termination of females.</p> <p>Females 15 days prior to mating and throughout gestation.</p>	<p>No effects on reproductive status were observed at any dose.</p> <p>Males at 30 & 90 mg/kg/day:</p> <ul style="list-style-type: none"> - At approximately 14 weeks of age, and after 6 weeks of dosing, the F0 males started producing an increased number of seminal plugs and lacerated genitalia - At termination of treatment, weight gain was significantly reduced in F0 males - A slight treatment-related reduction in mean postweaning weight gain among F1 males of the 30 and 90 mg/kg/day groups. <p>Females at 30 & 90 mg/kg/day:</p> <ul style="list-style-type: none"> - Decrease weight gain during gestation. <p>Pups: Reduced body weights in F1 pups at 90 mg/kg/day on Day 1 postpartum and secondarily a delay in postnatal development. Increased incidence of deaths of F1 pups at 30 and 90 mg/kg/day during lactation.</p>

Mutagenicity Studies

Enalapril was not mutagenic in the Ames microbial mutagen test with or without metabolic activation, in the Rec-Assay, sister chromatid exchange with cultured chinese hamster cells, (up to 20 mg/mL) and the micro-nucleus test with mice.

In vitro chromosomal aberration test = enalapril was clastogenic at 10 and 20 mg/mL but not at 5 mg/mL.

Carcinogenicity Studies

There was no evidence of a carcinogenic effect when enalapril was administered for 106 weeks to rats (Charles River CD-1) at doses up to 90 mg/kg/day (150 times the maximum daily human dose).

Enalapril has also been administered for 94 weeks to male and female mice (Charles River CD-1) at doses up to 90 and 180 mg/kg/day, respectively (150 and 300 times the maximum daily dose for humans), and no evidence of carcinogenicity was noted.

REFERENCES

1. Biollaz J, Burnier M, Turini GA, Brunner DB, Porchet M, Gomez HJ, Jones KH. Three new long-acting converting-enzyme inhibitors: relationship between plasma converting-enzyme activity and response to angiotensin I. *Clin Pharmacol Ther* 1981;29:665-70.
2. Biollaz J, Schelling JL, Jacot des Combes B, Brunner DB, Desponds G, Brunner HR. Enalapril maleate and a lysine analogue (MK-521) in normal volunteers; relationship between plasma drug levels and the renin angiotensin system. *Br J Clin Pharmacol* 1982;14:363-8.
3. Biollaz J, Brunner HR, Gavras I, Waeber B, Gavras H. Antihypertensive therapy with MK-421: Angiotensin II - Renin relationships to evaluate efficacy of converting enzyme blockade. *J Cardiovasc Pharmacol* 1982;4:966-72.
4. Brunner DB, Desponds G, Biollaz J, Keller I, Ferber F, Gavras H, Brunner HR, Schelling JL. Effect of a new angiotensin converting enzyme inhibitor MK-421 and its lysine analogue on the components of the renin system in healthy subjects. *Br J Clin Pharmacol* 1981;11:461-7.
5. Cleary JP, Taylor JW. Enalapril: a new angiotensin converting enzyme inhibitor. *Drug Intell Clin Pharm* 1986;20(3):177-86.
6. Cleland JGF, Dargie HJ, McAlpine H, Ball SG, Morton JJ, Robertson JIS, Ford I. Severe hypotension after first dose of enalapril in heart failure. *Br Med J* 1985;291:1309-12.
7. Cody RJ, Covit AB, Schaer GL, Laragh JH. Evaluation of a long-acting converting enzyme inhibitor (enalapril) for the treatment of chronic congestive heart failure. *J Am Coll Cardiol* 1983;1(4):1154-9.
8. Cohn JN, Johnson G, Ziesche S, Cobb F, Francis G, Tristani F, Smith R, Dunkman WB, Loeb H, Wong M, Bhat G, Goldman S, Fletcher RD, Doherty J, Hughes CV, Carson P, Cintron G, Shabetai R, Haakenson C. A comparison of enalapril with hydralazine-isosorbide dinitrate in the treatment of chronic congestive heart failure. *N Engl J Med* 1991;325:303-10.
9. Gavras H, Biollaz J, Waeber B, Brunner HR, Gavras I, Davies RO. Antihypertensive effect of the new oral angiotensin converting enzyme inhibitor "MK-421". *Lancet* 1981;ii:543-7.
10. Gomez HJ, Cirillo VJ. Angiotensin converting enzyme inhibitors: The clinical pharmacology of captopril and enalapril maleate. *Clinical pharmacology and therapeutics (International Congress Series No. 604)*, Velasco, M. ed., Excerpta Medica, Amsterdam. 1983;52-7.
11. Hodsman GP, Brown JJ, Davies DL, Fraser R, Lever AF, Morton JJ, Murray GD, Robertson JIS. Converting-enzyme inhibitor enalapril (MK-421) in treatment of hypertension with renal artery stenosis. *Br Med J* 1982;285:1697-9.
12. Hodsman GP, Brown JJ, Cumming AMM, Davies DL, East BW, Lever AF, Morton JJ, Murray GD, Robertson JIS. Enalapril (MK-421) in the treatment of hypertension with renal artery stenosis, (Symposium: on the renin-angiotensin-aldosterone system: treatment of

- hypertension and heart failure, Milan, Italy, 28 May 1983). *J Hypertension* 1(Suppl. 1)1983;109-17.
13. Ibsen H, Egan B, Osterziel K, Vander A, Julius S. Reflex-hemodynamic adjustments and baroreflex sensitivity during converting enzyme inhibition with MK-421 in normal humans. *Hypertension* 1983;5 (Suppl. D):184-91.
 14. Larochelle P, Carruthers SG, Krip G. Comparison of monotherapy with enalapril and atenolol in mild to moderate hypertension. *Can Med Assoc J* 1987;(37):803-8.
 15. McNabb WR, Brooks BA, Noormohamed F, Lant AF. The effect of enalapril on serum prolactin. *Br J Clin Pharmacol* 1983;15:752-4.
 16. Millar JA, Derx FHM, McLean K, Reid JL. Pharmacodynamics of converting enzyme inhibition: The cardiovascular, endocrine and autonomic effects of MK-421 (enalapril) and MK-521. *Br J Clin Pharmacol* 1982;14:347-55.
 17. Oberman A, Cutter GR, Rogers WJ, Mantle JA. Clinical Endpoints. 1982;341-55 in *Congestive Heart Failure*; ed. Braunwald E, Mock M, Watson J. Grune & Stratton.
 18. Packer M, Hung Lee W, Yushak M, Medina N. Comparison of captopril and enalapril in patients with severe chronic heart failure. *N Engl J Med* 1986;315:847-53.
 19. Packer M, Hung Lee W, Medina N, Yushak M, Kessler PD. Functional renal insufficiency during long-term therapy with captopril and enalapril in severe chronic heart failure. *Ann Intern Med* 1987;106(3):346-54.
 20. Packer M, ed. Do vasodilators prolong life in heart failure? *N Engl J Med* 1987;316:1471-73.
 21. Turini GA, Waeber B, Brunner HR. The renin-angiotensin system in refractory heart failure: Clinical, hemodynamic and hormonal effects of captopril and enalapril. *Eur Heart J* 1983;4:189-97.
 22. The CONSENSUS Trial Study Group. Effects of enalapril on mortality in severe congestive heart failure. Results of the Cooperative North Scandinavian Enalapril Survival Study (CONSENSUS). *N Engl J Med* 1987;316:1429-35.
 23. The SOLVD Investigators. Effect of enalapril on survival in patients with reduced left ventricular ejection fractions and congestive heart failure. *N Engl J Med* 1991;325:293-302.
 24. The SOLVD Investigators. Effect of enalapril on mortality and the development of heart failure in asymptomatic patients with reduced left ventricular ejection fractions. *N Engl J Med* 1992;327:685-91.
 25. Wilkins LH, Dustan HP, Walker JF, Oparil S. Enalapril in low-renin essential hypertension, *Clin Pharmacol Ther* 1983;34(3):297-302.

26. Product Monograph: VASOTEC[®], Merck Canada Inc., Control No. 188571, Revision dated January 25, 2016.

PART III: CONSUMER INFORMATION

Pr pms-ENALAPRIL

Enalapril Maleate Tablets, House Standard

This leaflet is part III of a three-part “Product Monograph” published when pms-ENALAPRIL was approved for sale in Canada and is designed specifically for Consumers. This leaflet is a summary and will not tell you everything about pms-ENALAPRIL. Contact your physician or pharmacist if you have any questions about the drug.

ABOUT THIS MEDICATION

pms-ENALAPRIL is available **only on prescription** from your physician.

What the medication is used for:

pms-ENALAPRIL is used for:

- reducing high blood pressure

When **blood pressure** is high, the workload of the heart and arteries increases so that over time, these organs may not function as they should. In turn, this could lead to damage of the “vital organs”: brain – heart – kidneys, and result in stroke, heart failure, heart attack, blood vessel disease or kidney disease.

pms-ENALAPRIL is used for:

- treating patients with heart failure

pms-ENALAPRIL may also be used to treat patients with **heart failure**. This is a condition where the heart cannot pump adequate amounts of blood to satisfy the needs of the body.

If your physician has recommended a particular diet, for instance less salt – follow the diet carefully. This could help your medicine to better control your blood pressure. Your physician may also recommend weight loss. Do follow these suggestions.

What it does:

pms-ENALAPRIL is part of a class of medicines known as angiotensin converting enzyme (ACE) inhibitors. They lower blood pressure by specifically blocking a naturally occurring substance called angiotensin II. Angiotensin II normally tightens your

blood vessels. pms-ENALAPRIL allows them to relax and therefore help lower high blood pressure.

This medicine does not cure high blood pressure, **but does help control it**. So, it is important to continue taking the tablets regularly to keep your blood pressure down. You may have to take high blood pressure medicine for life.

Keep your regular appointments with your physician, even if you feel well. High blood pressure may not be easily recognized by you, because you may not “feel any symptoms”; but your physician can measure your blood pressure very easily, and check how the medicine is controlling it.

Read the following information carefully. **If you need any explanations, or further information, ask your physician or pharmacist.**

When it should not be used:

Do not take pms-ENALAPRIL if you:

- are allergic to enalapril or any other component of pms-ENALAPRIL (see What the non-medicinal ingredients are).
- have a history of swelling of the face, lips, tongue, throat, or sudden difficulty breathing or swallowing.
- have been diagnosed with swelling of the face, lips, tongue, throat, or sudden difficulty breathing or swallowing due to genetic factors or unknown reasons (please refer to Side Effects and What to Do About Them).
- Are already taking a blood pressure-lowering medicine that contains aliskiren and you have diabetes or kidney disease.

What the medicinal ingredient is:

Each tablet is made with enalapril maleate that appears as enalapril sodium in the tablet.

What the non medicinal ingredients are:

Copovidone, lactose, pregelatinized starch, sodium bicarbonate and zinc stearate. 10 mg tablets also contain: red iron oxide. 20 mg tablets also contain: red iron oxide and yellow iron oxide.

What dosage forms it comes in:

Tablets: 2.5 mg, 5 mg, 10 mg, and 20 mg.

WARNINGS AND PRECAUTIONS

Serious Warnings and Precautions

pms-ENALAPRIL should not be used during pregnancy. If you discover that you are pregnant while taking pms-ENALAPRIL, stop the medication and please contact your physician as soon as possible.

This medicine may not be suitable for certain people. Tell your physician or pharmacist if you think **any** of the following applies to you:

- You have previously taken pms-ENALAPRIL or other medication of the same type – Angiotensin Converting Enzyme (ACE) inhibitors such as enalapril, lisinopril, captopril, and you were allergic or reacted badly to it, particularly if you experienced swelling of the face, lips, tongue, or throat, or had sudden difficulty breathing or swallowing. These are symptoms of conditions called hereditary angioedema or idiopathic angioedema.
- Are taking a medicine that contains aliskiren, used to lower high blood pressure. The combination with pms-ENALAPRIL is not recommended.
- Are taking an angiotensin receptor blocker (ARB). You can recognize an ARB because its medicinal ingredient ends in “-SARTAN”.
- Are taking anti-cancer (temsirolimus, everolimus) or anti-rejection (sirolimus) medications. Use of ACE inhibitors, such as pms-ENALAPRIL, with these drugs may increase the chance of having an allergic reaction (angioedema).
- **You should not take this medicine if you have been diagnosed with hereditary angioedema or idiopathic angioedema (angioedema of unknown cause).**
- Dizziness or drowsiness may occasionally occur when taking medication to lower blood pressure. Therefore, before you perform tasks which may require special attention (driving a car or operating dangerous machinery), wait until you know how you respond to your medicine.
- You should be aware that black patients are at increased risk of these types of reactions to ACE inhibitors.
- **You are pregnant, breast-feeding or thinking of becoming pregnant.** Taking pms-ENALAPRIL during pregnancy can cause injury and even death

to your developing baby. This medicine should **not** be used during pregnancy. If you become pregnant while taking pms-ENALAPRIL, stop the medication and report to your physician as soon as possible. It is possible that pms-ENALAPRIL passes into breast milk. You should not breast-feed while taking pms-ENALAPRIL.

- You suffer from low blood pressure (you may notice this as faintness or dizziness, especially when standing).
- You are undergoing dialysis.
- You have any of these conditions:
 - diabetes
 - heart or blood vessel disease
 - liver disease
 - kidney disease
- You are receiving gold (sodium aurothiomalate) injections.
- You are taking “water pills” or potassium supplements.
- You use potassium containing salt substitutes with your food.

You should also inform your physician or pharmacist if you have recently suffered from excessive vomiting or diarrhea.

If you have diabetes and are taking oral medicines to treat diabetes or insulin, you should closely monitor for low blood glucose levels, especially during the first month of treatment with pms-ENALAPRIL.

If you have to undergo any dental or other surgery, inform the dentist or the physician in charge that you are taking this medicine.

Remember – This medicine is prescribed for the particular condition that you have. **Do not give this medicine to other people, nor use it for any other condition.**

Do not use outdated medicine.

INTERACTIONS WITH THIS MEDICATION

Do not take any other medicines unless you have discussed the matter with your physician or pharmacist. Certain medications tend to increase your blood pressure, for example, non-prescription

IMPORTANT: PLEASE READ

preparations for appetite control, asthma, colds, coughs, hay fever and sinus problems, or may also react badly with pms-ENALAPRIL.

Your physician or pharmacist also needs to know if you are taking any other medication, whether on prescription or otherwise. It is particularly important to inform your physician or pharmacist if you are taking:

- Diuretics or “water pills”; any other medicines to reduce blood pressure.
- Diabetes medicine and/or insulin.
- Blood pressure-lowering drugs, including diuretics (“water pills”), aliskiren-containing products, or angiotensin receptor blockers (ARBs).
- Potassium-containing medicines, potassium supplements.
- Salt substitutes that contain potassium, as these may lead to increased levels of potassium in the blood which can be serious. In these cases, your physician may need to adjust the dosage of pms-ENALAPRIL or monitor your blood level of potassium.
- Lithium (a drug used to treat a certain kind of depression).
- Certain pain and arthritis medicines, including gold therapy and non-steroidal anti-inflammatory drugs.
- mTOR inhibitor (e.g., temsirolimus, sirolimus, everolimus) therapy. Taking these drugs together could increase the risk for an allergic reaction called angioedema.

PROPER USE OF THIS MEDICATION

Usual Dose:

- Take this medicine exactly as your physician ordered.
- The absorption of this medicine is not affected by food; so it can be taken with or without a meal.
- Try to take your medicine every day at the same time. This way it becomes easy to remember your doses.
- The splitting of pms-ENALAPRIL tablets is not advised.

For patients with high blood pressure:

Adults:

Your physician may adjust the dose according to your blood pressure response. The usual dose range is 10 mg to 40 mg per day given in a single dose or two divided doses. The maximum daily dose for pms-ENALAPRIL oral administration is 40 mg.

Children (less than 16 years):

The usual starting dose is up to 5 mg once daily. Doses above 40 mg have not been studied.

For patients with congestive heart failure:

The initial dose is 2.5 mg taken once a day. Your physician will closely monitor you to determine the initial effect on your blood pressure. The usual daily dose is 5 mg to 20 mg given in a single dose or two divided doses. The maximum daily dose is 40 mg.

Overdose:

If you think you have taken too much pms-ENALAPRIL, contact your healthcare professional, hospital emergency department or regional Poison Control Centre immediately, even if there are no symptoms.

The most likely symptom would be a feeling of lightheadedness or dizziness due to a sudden or excessive drop in blood pressure.

Missed Dose:

- If you miss a dose of this medicine, take it as soon as possible. However, if no more than six hours have elapsed since the missed dose, you may take that day’s dose of medication and then go back to your regular dosing schedule. **Do not take a double dose.**

SIDE EFFECTS AND WHAT TO DO ABOUT THEM

Along with its intended action, any medication, including pms-ENALAPRIL, may cause side effects. Most people do not experience any problem when taking these medicines; but if you notice any of the following, have other side effects or if the condition persists or worsens, seek medical attention.

- Dry cough, sore throat.
- The initial dose may cause a greater fall in blood

IMPORTANT: PLEASE READ

pressure than will occur following continued treatment. You may notice this as faintness or dizziness and it may help to lie down. If concerned, please consult your physician or pharmacist.

SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM				
Symptom / effect		Talk with your physician or pharmacist		Stop taking drug and seek immediate emergency medical attention
		Only if severe	In all cases	
Common	Fatigue	✓		
	Dizziness/Fainting/Lightheadedness, especially following exercise, and/or when it is hot and you have lost a lot of water by sweating			✓
	Low blood pressure		✓	
	Headache	✓		
	Rash/Itching		✓	
	Nausea/Vomiting/Diarrhea	✓		
	Lasting cough		✓	
	Chest pain		✓	
	Shortness of breath		✓	
Uncommon	Allergic reactions/ Angioedema, (sudden difficulty in breathing or swallowing, swelling of face, eyes, lips, tongue and/or throat, hands or feet)			✓

SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM			
Symptom / effect	Talk with your physician or pharmacist		Stop taking drug and seek immediate emergency medical attention
	Only if severe	In all cases	
Flu-like symptoms (fever, malaise, muscle pain, rash, itching, abdominal pain, nausea, vomiting, diarrhea, jaundice, loss of appetite)			✓
Liver impairment such as jaundice, dark/brown urine		✓	
Abdominal pain	✓		
Low blood sugars in diabetic patients	✓		
Loss of appetite	✓		

This is not a complete list of side effects. For any unexpected effects while taking pms-ENALAPRIL contact your physician or pharmacist.

HOW TO STORE IT

Store between 15°C and 30°C. Protect from moisture.

Discard remaining tablets six months after opening the bottle.

Keep all medicines out of the reach and sight of children.

Reporting Side Effects

You can help improve the safe use of health products for Canadians by reporting serious and unexpected side effects to Health Canada. Your report may help to identify new side effects and change the product safety information.

3 ways to report:

- Online at [MedEffet](http://www.healthcanada.gc.ca/medeffect) (www.healthcanada.gc.ca/medeffect);
- By calling 1-866-234-2345 (toll-free);
- By completing a Patient Side Effect Reporting Form and sending it by:
 - Fax to 1-866-678-6789 (toll-free), or
 - Mail to: Canada Vigilance Program
Health Canada, Postal Locator
0701E
Ottawa, ON
K1A 0K9Postage paid labels and the Consumer Side Effect Reporting Form are available at [MedEffet](http://www.healthcanada.gc.ca/medeffect) (www.healthcanada.gc.ca/medeffect).

NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

MORE INFORMATION

This document plus the full product monograph, prepared for health professionals can be obtained by contacting the sponsor Pharmascience Inc., at 1-888-550-6060.

This leaflet was prepared by

Pharmascience Inc.
Montréal, Québec
H4P 2T4

www.pharmascience.com

Last revised: April 7, 2016