

PRODUCT MONOGRAPH

Pr ^{phl}-LAMOTRIGINE

Lamotrigine Tablets, USP
25 mg, 100 mg and 150 mg

Antiepileptic

PHARMEL INC.
6111 Royalmount Ave., Suite 100
Montréal, Québec
H4P 2T4

Date of Revision:
June 15, 2016

Submission Control No.: 194956

Table of Contents

PART I: HEALTH PROFESSIONAL INFORMATION	3
SUMMARY PRODUCT INFORMATION.....	3
INDICATIONS AND CLINICAL USE	3
CONTRAINDICATIONS.....	3
WARNINGS AND PRECAUTIONS	4
ADVERSE REACTIONS	13
DRUG INTERACTIONS	19
DOSAGE AND ADMINISTRATION	24
OVERDOSAGE.....	28
ACTION AND CLINICAL PHARMACOLOGY	29
STORAGE AND STABILITY	33
DOSAGE FORMS, COMPOSITION AND PACKAGING.....	33
PART II: SCIENTIFIC INFORMATION	34
PHARMACEUTICAL INFORMATION	34
CLINICAL TRIALS	35
DETAILED PHARMACOLOGY	37
TOXICOLOGY.....	39
REFERENCES.....	42
PART III: CONSUMER INFORMATION.....	45

Pr **phl-LAMOTRIGINE**
Lamotrigine Tablets, USP
25 mg, 100 mg and 150 mg

PART I: HEALTH PROFESSIONAL INFORMATION

SUMMARY PRODUCT INFORMATION

Route of Administration	Dosage Form / Strength	Non-medicinal Ingredients
Oral	Tablets, 25 mg, 100 mg and 150 mg	Lactose monohydrate, magnesium stearate, microcrystalline cellulose, povidone, sodium starch glycolate, and the following: 100 mg tablets also include: FD&C yellow no. 6 150 mg tablets also include: yellow iron oxide

INDICATIONS AND CLINICAL USE

Adults (≥18 years of age)

phl-LAMOTRIGINE (lamotrigine) is indicated:

- as adjunctive therapy for the management of epilepsy who are not satisfactorily controlled by conventional therapy;
- for use as monotherapy following withdrawal of concomitant antiepileptic drugs;
- as adjunctive therapy for the management of the seizures associated with Lennox-Gastaut syndrome.

Geriatrics (> 65 years of age):

No dosage adjustment is required in patients over 65 years of age.

Pediatrics (<18 years of age)

phl-LAMOTRIGINE is indicated as adjunctive therapy for the management of the seizures associated with Lennox-Gastaut. Safety and efficacy in patients below the age of 16 years, other than those with Lennox-Gastaut Syndrome, have not been established.

CONTRAINDICATIONS

- Patients who are hypersensitive to this drug or to any ingredient in the formulation or component of the container. For a complete listing, see DOSAGE FORMS,

WARNINGS AND PRECAUTIONS

SERIOUS WARNING AND PRECAUTION: SERIOUS SKIN RASHES

Serious rashes associated with hospitalization have occurred with the use of lamotrigine. The incidence of these rashes in clinical trials was 1% (1/100) in pediatric patients (age < 16 years) and 0.3% (3/1000) in adults. The incidence of serious rash reported as Stevens-Johnson Syndrome (SJS) in clinical trials was 0.5% (1/200) in pediatric patients and 0.1% (1/1000) in adults. In worldwide post-marketing experience, rare cases of toxic epidermal necrolysis and/or death associated with rash have been reported, but their numbers are too few to permit a precise estimate of the rate.

In children, the initial presentation of a rash can be mistaken for an infection. Physicians should consider the possibility of a drug reaction in children that develop symptoms of rash and fever during the first eight weeks of therapy.

SERIOUS RASH ASSOCIATED WITH RAPID TITRATION

Do not exceed the recommended initial dose and subsequent dose escalations of *phl*-LAMOTRIGINE. A higher incidence of serious dermatologic events (see WARNINGS AND PRECAUTIONS, Skin-related Events, Table 1 and Table 2; see also DOSAGE AND ADMINISTRATION) has been associated with more rapid initial titration (exceeding the recommended initial dose or exceeding the recommended dose escalation), and use of concomitant valproic acid.

RASH ASSOCIATED WITH A HISTORY OF RASH TO OTHER ANTI-EPILEPTIC DRUGS

In two studies (n = 767 and n = 988), the frequency of rash with lamotrigine treatment was approximately 3-4 times higher in patients with a history of allergy or rash to other anti-epileptics, compared to those without such history.

Nearly all cases of rash associated with lamotrigine have occurred within 2 to 8 weeks of treatment initiation. However, isolated cases have been reported after prolonged treatment (e.g. 6 months). Accordingly, duration of therapy cannot be relied upon as a means to predict the potential risk signalled by the first appearance of a rash.

Although benign rashes also occur with lamotrigine, it is not possible to predict reliably which rashes will prove to be life threatening. Accordingly, all patients who develop rash should be promptly evaluated and lamotrigine withdrawn immediately, unless the rash is clearly not drug related.

General

Acute Multi-organ Failure

Multi-organ failure, which in some cases has been fatal or irreversible, has been observed in patients receiving lamotrigine. Fatalities associated with multi-organ failure and various degrees of hepatic failure have been reported in 2 of 3,796 adult patients and 4 of 2,435 pediatric patients who received lamotrigine in epilepsy clinical trials. No such fatalities have been reported in bipolar patients in clinical trials. Rare fatalities from multi-organ failure have also been reported in compassionate plea and post-marketing use. The majority of these deaths occurred in association with other serious medical events, including status epilepticus and overwhelming sepsis, and hantavirus, making it difficult to identify the initial cause.

Additionally, 3 patients (a 45-year-old woman, a 3.5-year-old boy, and an 11-year-old girl) developed multi-organ dysfunction and disseminated intravascular coagulation 9 to 14 days after lamotrigine was added to their AED regimens. Rash and elevated transaminases were also present in all patients and rhabdomyolysis was noted in 2 patients. Both pediatric patients were receiving concomitant therapy with valproate, while the adult patient was being treated with carbamazepine and clonazepam. All patients subsequently recovered with supportive care after treatment with lamotrigine was discontinued.

Drug Discontinuation

Abrupt discontinuation of any antiepileptic drug (AED) in a responsive patient with epilepsy may provoke rebound seizures. In general, withdrawal of an AED should be gradual to minimize this risk. Unless safety concerns (i.e. rash) require a more rapid withdrawal, the dose of phl-LAMOTRIGINE (lamotrigine) should be tapered over a period of at least two weeks (see DOSAGE AND ADMINISTRATION).

Hormonal Contraceptives

Patients taking phl-LAMOTRIGINE should be advised not to start or stop their oral contraceptives without consulting their physician. Significant adjustments in the maintenance dose of phl-LAMOTRIGINE may be required in some patients (see DRUG INTERACTIONS, Drug-Drug Interactions, Oral Contraceptives; ADVERSE REACTIONS, PRECAUTIONS, Post-Market Adverse Drug Reactions, Oral Contraceptives and DOSAGE AND ADMINISTRATION, Women and Oral Contraceptives).

Interactions with Antiepileptic Drugs

In a multidrug regimen, antiepileptic drugs may affect lamotrigine clearance, and dosage adjustment of phl-LAMOTRIGINE might be required (see DOSAGE AND ADMINISTRATION and DRUG INTERACTIONS).

Occupational Hazards

Patients with uncontrolled epilepsy should not drive or handle potentially dangerous machinery. During clinical trials common adverse effects included dizziness, ataxia, drowsiness, diplopia, and blurred vision. Patients should be advised to refrain from activities requiring mental

alertness or physical coordination until they are sure that phl-LAMOTRIGINE does not affect them adversely.

Other Products Containing Lamotrigine

Lamotrigine tablets should not be administered to patients currently being treated with any other preparation containing lamotrigine without consulting a doctor.

Potential Medication Errors

Medication errors involving lamotrigine have occurred. In particular, the names phl-LAMOTRIGINE or lamotrigine can be confused with the names of other commonly used medications. Medication errors may also occur between the different formulations of lamotrigine. To reduce the potential of medication errors, write and say phl-LAMOTRIGINE clearly. To avoid the medication error of using the wrong drug or formulation, patients should be strongly advised to visually inspect their tablets to verify that they are phl-LAMOTRIGINE, as well as the correct formulation of lamotrigine, each time they fill their prescription.

Hematologic

Blood Dyscrasias

There have been reports of blood dyscrasias that may or may not be associated with the hypersensitivity syndrome. These have included neutropenia, leukopenia, anemia, thrombocytopenia, pancytopenia, and, rarely, aplastic anemia and pure red cell aplasia.

Hypersensitivity Reactions

Drug Reaction with Eosinophilia and Systemic Symptoms

Multi-organ hypersensitivity reactions, also known as Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS), have occurred with lamotrigine. Some have been fatal or life threatening. DRESS typically, although not exclusively, presents with fever, rash, and/or lymphadenopathy in association with other organ system involvement, such as hepatitis, nephritis, hematologic abnormalities, aseptic meningitis, myocarditis, or myositis, sometimes resembling an acute viral infection (see ADVERSE REACTIONS). Eosinophilia is often present. This disorder is variable in its expression and other organ systems not noted here may be involved. The syndrome shows a wide spectrum of clinical severity and may rarely lead to disseminated intravascular coagulation (DIC) and multi-organ failure.

It is important to note that early manifestations of hypersensitivity (e.g. fever, lymphadenopathy) may be present even though a rash is not evident. If such signs or symptoms are present, the patient should be evaluated immediately. phl-LAMOTRIGINE should be discontinued if an alternative etiology for the signs or symptoms cannot be established.

Prior to initiation of treatment with phl-LAMOTRIGINE, the patient should be instructed that a rash or other signs or symptoms of hypersensitivity (e.g. fever, lymphadenopathy) may herald a serious medical event and that the patient should report any such occurrence to a physician immediately.

Neurologic

Aseptic Meningitis

Therapy with lamotrigine increases the risk of developing aseptic meningitis. Because of the potential for serious outcomes of untreated meningitis due to other causes, patients should also be evaluated for other causes of meningitis and treated as appropriate.

Post-marketing cases of aseptic meningitis have been reported in pediatric and adult patients taking lamotrigine for various indications. Symptoms upon presentation have included headache, fever, nausea, vomiting, and nuchal rigidity. Rash photophobia, myalgia, chills, altered consciousness, and somnolence were also noted in some cases. Symptoms have been reported to occur within 1 to 40 days following the initiation of treatment. In most cases, symptoms were reported to resolve after discontinuation of lamotrigine. Re-exposure resulted in a rapid return of symptoms (from within 30 minutes to 1 day following re-initiation of treatment) that were frequently more severe. Some of the patients treated with lamotrigine who developed aseptic meningitis had underlying diagnoses of systemic lupus erythematosus or other autoimmune diseases. Lamotrigine should not be restarted in patients who have discontinued due to aseptic meningitis associated with prior treatment of lamotrigine.

Cerebrospinal fluid (CSF) analysed at the time of clinical presentation in reported cases was characterized by a mild to moderate pleocytosis, normal glucose levels, and mild to moderate increase in protein. CSF white blood cell count differentials showed a predominance of neutrophils in a majority of the cases, although a predominance of lymphocytes was reported in approximately one third of the cases. Some patients also had new onset of signs and symptoms of involvement of other organs (predominantly hepatic and renal involvement) which may suggest that in these cases the aseptic meningitis observed was part of a hypersensitivity reaction.

Status Epilepticus

There are reports in the literature that severe convulsive seizures including status epilepticus may lead to rhabdomyolysis, multi-organ dysfunction and disseminated intravascular coagulation, sometimes with fatal outcome. Similar cases have occurred in association with the use of lamotrigine.

Psychiatric

Suicidal Behaviour and Ideation

Symptoms of depression and/or bipolar disorder may occur in patients with epilepsy, and there is evidence that patients with epilepsy and bipolar disorder have an elevated risk for suicidality.

Twenty-five to 50% of patients with bipolar disorder attempt suicide at least once, and may experience worsening of their depressive symptoms and/or the emergence of suicidal ideation and behaviours (suicidality) whether or not they are taking medications for bipolar disorder, including lamotrigine.

Suicidal ideation and behaviour have been reported in patients treated with antiepileptic agents in several indications.

All patients treated with antiepileptic drugs, irrespective of indication, should be monitored for signs of suicidal ideation and behaviour and appropriate treatment should be considered. Patients (and caregivers of patients) should be advised to seek medical advice should signs of suicidal ideation or behaviour emerge.

An FDA meta-analysis of randomized placebo controlled trials, in which antiepileptic drugs were used for various indications, has shown a small increased risk of suicidal ideation and behaviour in patients treated with these drugs. The mechanism of this risk is not known.

There were 43,892 patients treated in the placebo controlled clinical trials that were included in the meta-analysis. Approximately 75% of patients in these clinical trials were treated for indications other than epilepsy and, for the majority of non-epilepsy indications the treatment (antiepileptic drug or placebo) was administered as monotherapy. Patients with epilepsy represented approximately 25% of the total number of patients treated in the placebo controlled clinical trials and, for the majority of epilepsy patients, treatment (antiepileptic drug or placebo) was administered as adjunct to other antiepileptic agents (i.e. patients in both treatment arms were being treated with one or more antiepileptic drug). Therefore, the small increased risk of suicidal ideation and behaviour reported from the meta-analysis (0.43% for patients on antiepileptic drugs compared to 0.24% for patients on placebo) is based largely on patients that received monotherapy treatment (antiepileptic drug or placebo) for non-epilepsy indications. The study design does not allow an estimation of the risk of suicidal ideation and behaviour for patients with epilepsy that are taking antiepileptic drugs, due both to this population being the minority in the study, and the drug-placebo comparison in this population being confounded by the presence of adjunct antiepileptic drug treatment in both arms.

Skin

Skin-Related Events

In adult controlled studies of adjunctive lamotrigine therapy, the incidence of rash (usually maculopapular and/or erythematous) in patients receiving lamotrigine was 10% compared with 5% in placebo patients. The rash usually occurred within the first six weeks of therapy and resolved during continued administration of lamotrigine. Lamotrigine was discontinued because of rash in 1.1% of adult patients in controlled studies and 3.8% of all patients in all studies. The rate of rash-related withdrawal in clinical studies was higher with more rapid initial titration dosing, and in patients receiving concomitant valproic acid (VPA), particularly in the absence of AEDs that induce lamotrigine glucuronidation. See Table 1 and Table 2; see also DOSAGE AND ADMINISTRATION.

Table 1 Effect of Concomitant AEDs on Rash Associated with Lamotrigine in All Adult Controlled and Uncontrolled Clinical Trials Regardless of Dosing Escalation Scheme

AED Group	Total Patient Number	All Rashes	Withdrawal Due to Rash	Hospitalization in Association with Rash
AEDs that induce lamotrigine glucuronidation ¹	1788	9.2%	1.8%	0.1%
AEDs that induce lamotrigine glucuronidation ¹ + VPA	318	8.8%	3.5%	0.9%
VPA ± AEDs that neither inhibit nor induce lamotrigine glucuronidation ²	159	20.8%	11.9%	2.5%
AEDs that neither inhibit nor induce lamotrigine glucuronidation ²	27	18.5%	0.0%	0.0%

VPA = Valproic Acid

1. AEDs that induce lamotrigine glucuronidation include carbamazepine, phenobarbital, phenytoin, and primidone
2. AEDs that neither inhibit nor induce lamotrigine glucuronidation include clonazepam, clobazam, ethosuximide, methsuximide, vigabatrin, and gabapentin

Table 2 Effect of the Initial Daily Dose¹ of Lamotrigine in the Presence of Concomitant AEDs, on the Incidence of Rash Leading to Withdrawal of Treatment in Adult Add-On Clinical Trials

AED Group	AEDs that induce lamotrigine glucuronidation ²		AEDs that induce lamotrigine glucuronidation ² + VPA		VPA ± AEDs that neither inhibit nor induce lamotrigine glucuronidation ³	
	Total Patient Number	Percentage of Patients Withdrawn	Total Patient Number	Percentage of Patients Withdrawn	Total Patient Number	Percentage of Patients Withdrawn
Lamotrigine Average Daily Dose (mg)						
12.5	9	0.0	10	0.0	51	7.8
25	3	0.0	7	0.0	58	12.1
50	182	1.1	111	0.9	35	5.7
100	993	1.4	179	4.5	15	40.0
≥125	601	2.8	11	18.2	0	0.0

VPA = valproic acid

¹Average daily dose in week 1

²AEDs that induce lamotrigine glucuronidation include carbamazepine, phenobarbital, phenytoin, and primidone

³AEDs that neither inhibit nor induce lamotrigine glucuronidation include clonazepam, clobazam, ethosuximide, methsuximide, vigabatrin, and gabapentin

Increased incidence of rash-related withdrawal was seen when initial doses were higher and titration more rapid than recommended under DOSAGE AND ADMINISTRATION.

Patients with a history of allergy or rash to other antiepileptic drugs

Caution is also required when treating patients with a history of allergy or rash to other antiepileptic drugs, as it was found in two studies (n = 767 and n = 988) on the frequency of rash after treatment with lamotrigine that the rate of rash was approximately three to four times higher in patients with such a history than those without.

Restarting phl-LAMOTRIGINE Therapy

It is recommended that phl-LAMOTRIGINE not be restarted in patients who discontinued due to rash associated with prior treatment with phl-LAMOTRIGINE unless the potential benefits clearly outweigh the risks. If the decision is made to restart a patient who has discontinued phl-LAMOTRIGINE for any reason, the need to restart with the initial dosing recommendations should be assessed. The greater the interval of time since the previous dose, the greater consideration should be given to restarting with the initial dosing recommendations. If a patient has discontinued phl-LAMOTRIGINE for a period of more than 5 half-lives, it is recommended that initial dosing recommendations and guidelines be followed. The half-life of lamotrigine is affected by other concomitant medications (see ACTION AND CLINICAL PHARMACOLOGY, Pharmacokinetics and DOSAGE AND ADMINISTRATION).

Special Populations

Pregnant Women: phl-LAMOTRIGINE should only be used during pregnancy if the expected benefits outweigh the potential risks. If therapy with phl-LAMOTRIGINE is considered necessary during pregnancy, the lowest possible therapeutic dose is recommended.

Physiological changes during pregnancy may affect lamotrigine levels and/or therapeutic effect. There have been reports of decreased lamotrigine levels during pregnancy. Appropriate clinical management of pregnant women during lamotrigine therapy should be ensured.

Studies in mice, rats and rabbits given lamotrigine orally or intravenously revealed no evidence of teratogenicity, however, maternal and secondary fetal toxicity producing reduced fetal weight and/or delayed ossification in mice and rats were observed.

In rats, enhanced fetal as well as post-natal mortality was observed when lamotrigine was administered during late gestation and through the early post-natal period. These effects were observed at the expected clinical exposure. Lamotrigine reduced fetal folic acid levels in rats. Folic acid deficiency is assumed to be associated with an enhanced risk of congenital malformations in animals as well as in humans. Intake of folic acid when planning pregnancy and during early pregnancy may be considered.

Studies in rats and rabbits indicate that lamotrigine crosses the placenta; placental and fetal plasma concentrations of lamotrigine are comparable to levels in maternal plasma. In studies where simultaneous maternal and umbilical cord samples were taken, the ratio of umbilical cord/maternal lamotrigine plasma concentrations were generally close to unity (range of 0.4 - 1.4).

Pregnancy Registries

Post-marketing data from six prospective pregnancy registries have documented outcomes in approximately 8700 women exposed to lamotrigine monotherapy during the first trimester of pregnancy. Two registries have reported an increase in the risk of isolated oral cleft (isolated) malformations with exposure to lamotrigine in the first trimester, over both the study reference population and reported background rates from the literature. The observed risk of malformations following first trimester monotherapy exposure ranged from 2.0% in the North American Registry to 4.9% in the Swedish Registry as compared to unexposed groups.

North American Registry

The North American Anti-Epileptic Drug Registry reported rates of: 7.3/1000 for lamotrigine exposures in the registry with n = 684 vs. 0.70/1000 for the reference population and 0.50-2.16/1000 for the background rates.

Swedish Registry

The Swedish Medical Birth Register reported rates of 9.9/1000 for lamotrigine exposures in the registry with n = 403 vs. 2.0/1000 for the registry reference population and 0.50- 2.16/1000 for the background rates. In the remaining registries with approximately 3000 women, 4 cases of isolated oral cleft malformation were found (1.3/1000 vs. 0.50-2.16/1000 for the background rates).

EUROCAT

The European Network of Congenital Anomaly and Twin Registries (EUROCAT) is a network of 40 registries across 20 European countries. The results of a case control design using the EUROCAT dataset demonstrated that the risk of development of oral clefts in the newborn infants was similar to other malformations (e.g. club foot, limb, heart and respiratory defects) following exposure to lamotrigine in the first trimester. Within the EUROCAT network, rates of exposure to lamotrigine in the first trimester of pregnancy were similar among 4571 non-chromosomal, isolated oral cleft cases and 80,052 non-chromosomal, non-oral cleft defect controls.

The data on use of lamotrigine in polytherapy combinations are insufficient to assess whether the risk of malformation associated with other agents is affected by concomitant lamotrigine use.

To provide information regarding the effects of *in utero* exposure to lamotrigine, physicians are advised to recommend that pregnant patients taking lamotrigine enroll in the North American Antiepileptic Drug (NAAED) Pregnancy Registry. This can be done by calling the toll-free number 1-888-233-2334, and must be done by patients themselves. Information on the registry can also be found at the website <http://www.aedpregnancyregistry.org/>.

Labour and Delivery: The effect of lamotrigine on labour and delivery in humans is unknown.

Nursing Women: Lamotrigine has been reported to pass into breast milk resulting in total lamotrigine levels in infant's plasma, including neonates as well as older infants, of up to approximately 50% of the mother's plasma. Therefore, in some breast-fed infants, plasma

concentrations of lamotrigine may reach levels at which pharmacological effects do occur.

Because of the potential for adverse reactions from lamotrigine in nursing infants, breast-feeding while taking this medication is not recommended. Should a woman breast-feed while on therapy with lamotrigine, the infant should be monitored for adverse effects. Events including apnea, drowsiness, and poor sucking have been reported in infants who have been human milk-fed by mothers using lamotrigine; whether or not these events were caused by lamotrigine is unknown.

Geriatrics: As the pharmacokinetics in this age group do not differ significantly from a non-elderly adult population, no dosage adjustment from the recommended adult schedule is required (see also DOSAGE AND ADMINISTRATION, ADVERSE REACTIONS, and ACTION AND CLINICAL PHARMACOLOGY).

Patients with Special Diseases and Conditions: Clinical experience with lamotrigine in patients with concomitant illness is limited. Caution is advised when using phl-LAMOTRIGINE in patients with diseases or conditions that could affect the metabolism or elimination of the drug.

Renal Impairment: A study in individuals with chronic renal failure (not receiving other AEDs) indicated that the elimination half-life of unchanged lamotrigine is prolonged relative to individuals with normal renal function (see ACTION AND CLINICAL PHARMACOLOGY). phl-LAMOTRIGINE should be used with caution in patients with severe renal impairment.

Hepatic Impairment: Results from a single dose pharmacokinetic study indicate that the apparent clearance of lamotrigine decreased in subjects with Grades A, B or C hepatic impairment. A reduced dosage should be used for all hepatically impaired patients, and phl-LAMOTRIGINE should be administered with caution particularly in those patients with severe hepatic impairment (see also DOSAGE AND ADMINISTRATION and ACTION AND CLINICAL PHARMACOLOGY).

Cardiac Conduction Abnormalities: One placebo-controlled trial that compared electrocardiograms at baseline and during treatment, demonstrated a mild prolongation of the P-R interval associated with lamotrigine administration. The prolongation was statistically significant but clinically insignificant. Patients with significant cardiovascular disease or electrocardiographic abnormalities were, however, systematically excluded from clinical trials. Thus, phl-LAMOTRIGINE should be used with caution in patients with cardiac conduction abnormalities, and in patients taking concomitant medications which depress AV conduction.

Monitoring and Laboratory Tests

The relationship between clinical efficacy and plasma concentrations has not been clearly established. Based on the possible pharmacokinetic interactions between lamotrigine and other drugs including AEDs, monitoring of the plasma levels of lamotrigine and concomitant drugs may be indicated, particularly during dosage adjustments. In general, clinical judgment should be exercised regarding monitoring of plasma levels of lamotrigine and other drugs and whether or

not dosage adjustments are necessary.

ADVERSE REACTIONS

Adverse Drug Reaction Overview

Rarely, serious skin rashes, including Stevens Johnson Syndrome and Toxic Epidermal Necrolysis (Lyell Syndrome) have been reported. Although the majority recover following drug withdrawal, some patients experience irreversible scarring and there have been rare cases of associated death (see WARNINGS AND PRECAUTIONS).

Adverse experiences in patients receiving lamotrigine were generally mild, occurred within the first two weeks of therapy, and resolved without discontinuation of the drug.

Clinical Trial Adverse Drug Reactions

Because clinical trials are conducted under very specific conditions the adverse reaction rates observed in the clinical trials may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse drug reaction information from clinical trials is useful for identifying drug-related adverse events and for approximating rates.

Commonly Observed

The most commonly observed adverse experiences associated with the use of adjunctive therapy with lamotrigine (incidence of at least 10%) were dizziness, headache, diplopia, somnolence, ataxia, nausea, and asthenia.

Dizziness, diplopia, ataxia, and blurred vision were dose-related and occurred more commonly in patients receiving carbamazepine in combination with lamotrigine than in patients receiving other AEDs that induce lamotrigine glucuronidation in combination with lamotrigine. Reduction of the daily dose and/or alteration of the timing of doses of concomitant antiepileptic drugs and/or lamotrigine may reduce or eliminate these symptoms. Clinical data suggest a higher incidence of rash in patients who are receiving concomitant valproic acid, or non-inducing AEDs (see WARNINGS AND PRECAUTIONS, Skin-Related Events, Table 1).

Adverse Events Associated with Discontinuation of Treatment

Across all adult add-on studies, the most common adverse experiences associated with discontinuation of lamotrigine were rash, dizziness, headache, ataxia, nausea, diplopia, somnolence, seizure exacerbation, asthenia, and blurred vision. In controlled clinical trials, 6.9% of the 711 patients receiving lamotrigine discontinued therapy due to an adverse experience, versus 2.9% of the 419 patients receiving placebo. Of 3501 patients and volunteers who received lamotrigine in pre-marketing clinical studies, 358 (10.2%) discontinued therapy due to an adverse experience.

Serious Adverse Events Associated with Discontinuation of Treatment

Discontinuation due to an adverse experience classified as serious occurred in 2.3% of adult patients and volunteers who received lamotrigine in the pre-market studies. Rash accounted for almost half of the discontinuations due to serious adverse experiences. More rapid initial titration of lamotrigine and concomitant use of valproic acid were associated with higher incidences of rash-related withdrawal in clinical studies (see WARNINGS AND PRECAUTIONS, Skin-Related Events, Table 2).

Adult Controlled Add-on Clinical Studies

Table 3 enumerates adverse experiences that occurred with an incidence of 2% or greater among refractory patients with epilepsy treated with lamotrigine.

Table 3 Percentage of Treatment - Emergent Adverse Experiences in Adult Placebo or Comparator-Controlled Clinical Studies¹

	ADULTS (ADJUNCTIVE THERAPY) ²		ELDERLY (MONOTHERAPY) ³
	LAMOTRIGINE (and other AEDs)	Placebo (and other AEDs)	LAMOTRIGINE
Total Number of Patients	(n = 711)	(n = 419)	(n = 102)
Body System/Adverse Experience⁴			
BODY AS A WHOLE			
Headache	29.1	19.1	8.8
Accidental Injury	9.1	8.6	8.8
Asthenia	8.6	8.8	4.9
Flu Syndrome	7.0	5.5	4.9
Pain	6.2	2.9	5.9
Back Pain	5.8	6.2	3.9
Fever	5.5	3.6	0.9
Abdominal Pain	5.2	3.6	3.9
Infection	4.4	4.1	5.9
Neck Pain	2.4	1.2	0
Malaise	2.3	1.9	4.9
Seizure Exacerbation	2.3	0.5	n/a
CARDIOVASCULAR			
Chest pain	n/a	n/a	2.9
Syncope	n/a	n/a	2.9
Cerebrovascular accident	n/a	n/a	3.9
DIGESTIVE			
Nausea	18.6	9.5	8.8
Vomiting	9.4	4.3	8.8
Diarrhea	6.3	4.1	6.9
Dyspepsia	5.3	2.1	5.9
Constipation	4.1	3.1	8.9
Tooth Disorder	3.2	1.7	0

	ADULTS (ADJUNCTIVE THERAPY) ²		ELDERLY (MONOTHERAPY) ³
	LAMOTRIGINE (and other AEDs)	Placebo (and other AEDs)	LAMOTRIGINE
Total Number of Patients	(n = 711)	(n = 419)	(n = 102)
Body System/Adverse Experience⁴			
MUSCULOSKELETAL			
Myalgia	2.8	3.1	0.9
Arthralgia	2.0	0.2	2.9
NERVOUS			
Dizziness	38.4	13.4	9.8
Ataxia	21.7	5.5	0
Somnolence	14.2	6.9	11.8
Incoordination	6.0	2.1	12.7
Insomnia	5.6	1.9	3.9
Tremor	4.4	1.4	0.9
Depression	4.2	2.6	4.9
Anxiety	3.8	2.6	0.9
Convulsion	3.2	1.2	1.9
Irritability	3.0	1.9	0
Speech Disorder	2.5	0.2	0.9
Memory Decreased	2.4	1.9	n/a
Memory Decreased (Memory Rating Question)	n/a	n/a	19.6
RESPIRATORY			
Rhinitis	13.6	9.3	0.9
Pharyngitis	9.8	8.8	1.9
Cough Increased	7.5	5.7	2.9
Respiratory Disorder	5.3	5.5	0.9
Asthma			3.0
SKIN AND APPENDAGES			
Rash	10.0	5.0	8.8
Pruritus	3.1	1.7	5.9
Herpes Zoster	n/a	n/a	3.0
Eczema	n/a	n/a	2.0
Ulcer Skin	n/a	n/a	2.0
SPECIAL SENSES			
Diplopia	27.6	6.7	0
Blurred Vision	15.5	4.5	0
Vision Abnormality	3.4	1.0	0
UROGENITAL			
Female Patients	(n = 365)	(n = 207)	(n = 47)
Dysmenorrhea	6.6	6.3	n/a
Menstrual Disorder	5.2	5.8	n/a
Vaginitis	4.1	0.5	0

¹ Patients from the studies summarized in the first two columns were receiving 1 to 3 concomitant enzyme-inducing antiepileptic drugs in addition to lamotrigine or placebo. Patients from the single study summarized in the last column were compared to n = 48 patients receiving carbamazepine. Patients may have reported multiple adverse experiences during the

study or at discontinuation. Thus, patients may be included in more than one category.

² Studies 05, 06, 16 (US) & 16, 21, 35 & 37 (UK)

³ Study 105 - 124 - C93

⁴ All Adverse Experiences reported by at least 2% of patients treated with either lamotrigine add-on or monotherapy are included.

Other Events Observed During Clinical Studies

During clinical testing, multiple doses of lamotrigine were administered to 3501 patients and volunteers. The conditions and duration of exposure to lamotrigine during these clinical studies varied greatly. Studies included monotherapy and pediatric trials. A substantial proportion of the exposure was gained in open, uncontrolled clinical studies. Adverse experiences associated with exposure to lamotrigine were recorded by clinical investigators using terminology of their own choosing. Consequently, it is not possible to provide a meaningful estimate of the proportion of individuals experiencing adverse events without first grouping similar types of adverse experiences into a smaller number of standardized event categories.

Since the adverse experiences reported occurred during treatment with lamotrigine in combination with other antiepileptic drugs, they were not necessarily caused by lamotrigine.

The following adverse events have been reported on one or more occasions by at least 1% of patients and volunteers exposed to lamotrigine: anorexia, weight gain, amnesia, concentration disturbance, confusion, emotional lability, nervousness, nystagmus, paresthesia, thinking abnormality and vertigo (all types of events are included except those already listed in Table 3).

Adult Monotherapy Clinical Studies

Withdrawals due to adverse events were reported in 42 (9.5%) of newly diagnosed patients treated with lamotrigine monotherapy. The most common adverse experiences associated with discontinuation of lamotrigine were rash (6.1%), asthenia (1.1%), headache (1.1%), nausea (0.7%) and vomiting (0.7%).

Geriatric Monotherapy Clinical Studies

A study with elderly newly-diagnosed epilepsy patients yielded rates of adverse events which were generally similar to those reported in adults (see Table 3). The rate of withdrawal due to adverse events was 21.6 %, with rash (3%), nausea (3%) and coordination abnormalities (3%) representing the most common events associated with withdrawal, followed by somnolence (2%), depression (2%), accidental injury (2%) and malaise (2%) (see also DOSAGE AND ADMINISTRATION, and ACTION AND CLINICAL PHARMACOLOGY).

Adjunctive Therapy in Lennox-Gastaut Syndrome

In 169 adult and pediatric patients with Lennox-Gastaut syndrome, 3.8% of patients on lamotrigine and 7.8% of patients on placebo discontinued due to adverse experiences. The most commonly reported adverse experiences that led to discontinuation were rash for patients treated with lamotrigine and deterioration of seizure control for patients treated with placebo. Fever and infection occurred at least 10% more frequently in patients ≤ 12 years of age than in patients > 12 years of age on lamotrigine. Rash occurred at least 10% more frequently in female patients than male patients on lamotrigine. Table 4 lists adverse events that occurred in at least 1% of 79 adult

and pediatric patients who received lamotrigine up to 15 mg/kg per day or a maximum of 400 mg per day.

Table 4 Treatment-Emergent Adverse Experience Incidence in Placebo-Controlled Add-On Trial in Adult and Pediatric Patients with Lennox-Gastaut Syndrome¹

Body System/ Adverse Experience	Percent of Patients Receiving Lamotrigine (n = 79)	Percent of Patients Receiving Placebo (n = 90)
Body as a whole:		
Infection	13	8
Accidental injury	9	7
Flu Syndrome	5	0
Asthenia	3	1
Abdominal Pain	3	0
Back Pain	1	0
Edema of the face	1	0
Lab test abnormal	1	0
Pain	1	0
Cardiovascular:		
Hemorrhage	3	0
Digestive:		
Vomiting	9	7
Constipation	5	2
Diarrhea	4	2
Nausea	4	1
Anorexia	3	1
Stomatitis aphtha	1	0
Tooth disorder	1	0
Endocrine:		
Cushing's syndrome	1	0
Hypothyroidism	1	0
Hemic and lymphatic:		
Lymphadenopathy (enlarged cervical nodes)	1	0
Nervous system:		
Ataxia	4	1
Convulsions	4	1
Tremor	3	0
Agitation	1	0
Coordination	1	0
Dizziness	1	0
Emotional lability	1	0
Nervousness	1	0
Vertigo	1	0
Respiratory:		
Pharyngitis	14	10
Bronchitis	9	7
Pneumonia	3	0
Dyspnea	1	0
Skin:		
Rash	9	7
Eczema	4	0
Nail disorder	1	0
Special senses:		
Blepharitis	1	0
Conjunctivitis	1	0
Keratitis	1	0
Ear pain	1	0
Eye pain	1	0
Urogenital:		
Urinary tract infection	3	0
Balanitis	2	0

Body System/ Adverse Experience	Percent of Patients Receiving Lamotrigine (n = 79)	Percent of Patients Receiving Placebo (n = 90)
Penis disorder	2	0

¹ = the most frequently reported adverse reactions in children ≤ 12 years of age in both treatment groups were pharyngitis, fever and infection.

Less Common Clinical Trial Adverse Drug Reactions (≥1% and <2%)

Body as a Whole: chest pain, chills

Cardiovascular: cerebral infarction, hot flashes, palpitations

Gastrointestinal: dry mouth

Musculoskeletal: joint disorder, myasthenia

Nervous System: dream abnormality, dysarthria, grand mal convulsion, hostility, hypertension, hypesthesia, mind racing, muscle spasm, sleep disorder

Metabolic: hyperkalemia, peripheral edema

Skin: acne, alopecia

Special Senses: eye disorder

Urogenital: amenorrhea, dysuria

Post-Market Adverse Drug Reactions

In addition to the adverse experiences reported during clinical testing of lamotrigine, the following adverse experiences have been reported in patients receiving marketed lamotrigine and from worldwide investigational use. These adverse experiences have not been listed above, and data are insufficient to establish causation.

Body as a Whole: hypersensitivity reaction, multi-organ failure, progressive immunosuppression, tiredness

Gastrointestinal: Esophagitis

Hematologic and Lymphatic: Hematological abnormalities (anemia, agranulocytosis, aplastic anemia, disseminated intravascular coagulation, hemolytic anemia, leucopenia, neutropenia, pancytopenia, red cell aplasia thrombocytopenia), lymphadenopathy*

*There have been reported cases of lymphadenopathy in the absence of hypersensitivity reactions, in patients with or without previous history, and taking lamotrigine alone or in combination with other medications (including other AEDs). Discontinuation of lamotrigine or, in some cases, dose reduction resolved the reaction.

Hepatobiliary Tract and Pancreas: elevations of liver function tests, hepatic dysfunction, including hepatic failure*, pancreatitis

*Hepatic dysfunction usually occurs in association with hypersensitivity reactions but isolated cases have been reported without overt signs of hypersensitivity.

Immunologic: lupus-like reaction, vasculitis

Lower Respiratory: apnea

Musculoskeletal: rhabdomyolysis (observed in patients experiencing hypersensitivity reactions)

Neurologic: aggression, aseptic meningitis*, exacerbation of parkinsonian symptoms**, extrapyramidal effects, choreoathetosis, hallucinations, movement disorders (such as tics and unsteadiness), nightmares

*There have been rare reports of aseptic meningitis in patients taking lamotrigine alone or in combination with other AEDs. In some cases, discontinuation of lamotrigine resolved this reaction (see WARNINGS AND PRECAUTIONS, Hypersensitivity Reactions).

**There have been reports that lamotrigine may worsen parkinsonian symptoms in patients with pre-existing Parkinson's disease, and isolated reports of extrapyramidal effects and choreoathetosis in patients without this underlying condition.

Oral Contraceptives: There have been post-marketing reports of breakthrough seizures in patients taking lamotrigine and hormonal contraceptives. In some cases, reduced serum lamotrigine levels during co-administration with a hormonal contraceptive have been documented. Most cases have responded to lamotrigine dose increase (see DOSAGE AND ADMINISTRATION, Women and Oral Contraceptives and DRUG INTERACTIONS, Oral Contraceptives).

DRUG INTERACTIONS

Drug-Drug Interactions

Antiepileptic Drugs (AEDs)

The net effects of co-administration of lamotrigine with other antiepileptic drugs are summarized in Table 5. Additional details of these drug interaction studies are provided below.

Table 5 Summary of AED Interactions with Lamotrigine

AED	AED Plasma Concentration with Adjunctive Lamotrigine ¹	Lamotrigine Plasma Concentration with Adjunctive AEDs ²
Carbamazepine (CBZ) CBZ epoxide ³	No significant effect Conflicting data	↓40%
Felbamate	Not evaluated	No significant effect
Gabapentin	Not evaluated	No significant effect
Levetiracetam	No significant effect	No significant effect
Oxcarbazepine	No significant effect	No significant effect
10-monohydroxy oxcarbazepine metabolite	No significant effect	Not evaluated
Phenobarbital	No significant effect	↓40%

AED	AED Plasma Concentration with Adjunctive Lamotrigine ¹	Lamotrigine Plasma Concentration with Adjunctive AEDs ²
Phenytoin (PHT)	No significant effect	↓40 %
Pregabalin	No significant effect	No significant effect
Primidone	No significant effect	↓40%
Topiramate	No significant effect ⁴	No significant effect
Valproic Acid (VPA)	Decreased ⁵	↑200%
VPA + PHT and/or CBZ	Not evaluated	No significant effect
Zonisamide	Not evaluated	No significant effect

1 From adjunctive clinical trials and volunteer studies.

2 Net effects were estimated by comparing the mean clearance values obtained in adjunctive clinical trials and volunteer studies.

3 Not administered, but an active metabolite of carbamazepine.

4 Slight increase, not expected to be clinically relevant

5 Refer to Valproic Acid paragraph below for additional information

↓ Decreased (induces lamotrigine glucuronidation)

↑ Increased (inhibits lamotrigine glucuronidation)

Lamotrigine does not affect the plasma concentrations of concomitantly administered AEDs that induce lamotrigine glucuronidation. Antiepileptic drugs that induce lamotrigine glucuronidation (such as phenytoin, carbamazepine, phenobarbital, and primidone) increase the plasma clearance and reduce the elimination half-life of lamotrigine (see ACTION AND CLINICAL PHARMACOLOGY).

Felbamate (not available in Canada)

In a cross-over study of 21 healthy male volunteers, co-administration of felbamate (1200 mg twice daily) with lamotrigine (100 mg twice daily for 10 days) appeared to have no clinically relevant effects on the pharmacokinetics of lamotrigine. Furthermore, felbamate had no significant effect on urinary excretion of total lamotrigine, unconjugated lamotrigine or N-glucuronide metabolite.

Gabapentin

Based on a retrospective analysis of plasma levels of 34 epileptic patients who received lamotrigine both with and without gabapentin, gabapentin did not appear to change the apparent clearance of lamotrigine. No information is available about the effects of gabapentin on C_{max} or AUC of lamotrigine.

Levetiracetam

Potential drug interactions between levetiracetam and lamotrigine were assessed by evaluating serum concentrations of both agents during placebo-controlled clinical trials. In 48 refractory epileptic patients receiving stable doses of lamotrigine, concomitant levetiracetam (1000-4000 mg/day) did not influence mean steady serum concentrations of lamotrigine. Similarly, lamotrigine did not influence the pharmacokinetics of levetiracetam.

Oxcarbazepine

The AUC and C_{max} of oxcarbazepine and its active 10-monohydroxy oxcarbazepine metabolite were not significantly different following the addition of oxcarbazepine (600 mg twice daily) to

lamotrigine (200 mg once daily) in healthy male volunteers (n = 13) compared to healthy male volunteers receiving oxcarbazepine alone (n = 13). Limited clinical data suggest a higher incidence of headache, dizziness, nausea, and somnolence with co-administration of lamotrigine and oxcarbazepine compared to lamotrigine alone or oxcarbazepine alone.

Pregabalin

In 12 patients with partial-onset seizures receiving combination lamotrigine (mean lamotrigine dose 287.5 mg/day, range 100-600 mg/day) and pregabalin 600 mg/day (200 mg three times daily) therapy, steady-state trough plasma concentrations of lamotrigine were not affected by pregabalin. The pharmacokinetics of pregabalin when co-administered with lamotrigine in this study appeared similar to historical values for pregabalin in healthy volunteers.

Topiramate

In three studies in patients with epilepsy (n=52), topiramate (dose range of topiramate in the 2 studies where this was recorded was 75-800 mg/day) resulted in no change in plasma concentrations of lamotrigine. Administration of lamotrigine resulted in a 15% increase in plasma concentrations of topiramate.

Valproic Acid

Valproic acid reduces the plasma clearance and prolongs the elimination half-life of lamotrigine (see ACTION AND CLINICAL PHARMACOLOGY). When lamotrigine was administered to 18 healthy volunteers already receiving valproic acid, a modest decrease (25% on average) in the trough steady-state valproic acid plasma concentrations was observed over a 3-week period, followed by stabilization. However, the addition of lamotrigine did not affect the plasma concentration of valproic acid in patients receiving AEDs that induce lamotrigine glucuronidation in combination with valproic acid. See also WARNINGS AND PRECAUTIONS, Skin-Related Events.

Zonisamide (not available in Canada)

In a study of 18 patients with epilepsy, co-administration of zonisamide (200 to 400 mg/day) with lamotrigine (150 to 500 mg/day) for 35 days had no significant effect on the pharmacokinetics of lamotrigine. Although the effects of lamotrigine on pharmacokinetics of zonisamide have not been studied, two post-marketing cases have reported zonisamide toxicity when lamotrigine was administered (zonisamide dose: 600-800 mg daily; lamotrigine dose: 400 mg daily). In both cases, positive dechallenge and rechallenge were observed.

Other Drug Interactions

Aripiprazole

In a study of 18 adult patients with bipolar I disorder, receiving an established regimen of lamotrigine (≥ 100 mg/day), doses of aripiprazole were increased from 10 mg/day to a target of 30 mg/day over a 7 day period and continued once daily for a further 7 days. An average reduction of approximately 10% in C_{max} and AUC of lamotrigine was observed. An effect of this magnitude is not expected to be of clinical consequence.

Atazanavir/ritonavir

In a study of 21 healthy adult volunteers, atazanavir/ritonavir combination (300 mg/100 mg) reduced the plasma AUC, C_{max} and elimination half-life of lamotrigine (single 100 mg dose) by an average of 32%, 6% and 27%, respectively. However, atazanavir alone did not induce lamotrigine glucuronidation (did not change pharmacokinetics of lamotrigine) to a clinically significant extent.

To initiate lamotrigine in patients already on the combination therapy of atazanavir/ritonavir, dose titration should follow the recommended guidelines based on whether lamotrigine is added to valproate (an inhibitor of lamotrigine glucuronidation), or to an inducer of lamotrigine glucuronidation, or whether lamotrigine is added in the absence of valproate or an inducer of lamotrigine glucuronidation (see DOSAGE AND ADMINISTRATION, Table 6).

To initiate atazanavir/ritonavir in patients already on maintenance doses of lamotrigine, the lamotrigine dose may need to be increased if another glucuronidation inducer is taken, but its dose may remain unchanged in the absence of another inducer. The lamotrigine dose needs to be decreased if atazanavir/ritonavir are discontinued.

Lopinavir/ritonavir

In a study in healthy volunteers, lopinavir/ritonavir approximately halved the plasma concentrations of lamotrigine. At least 14 out of 22 healthy adult volunteers reported diarrhea and 5 out of 22 subjects reported rash following the addition of lopinavir/ritonavir (400 mg/100 mg b.i.d.) to lamotrigine (up to 200 mg b.i.d.), compared to 1 out of 24 reporting diarrhea or rash, when lamotrigine was administered alone (up to 100 mg b.i.d.). Diarrhea is a common adverse event reported for lopinavir/ritonavir. The titration of lamotrigine in this study was rapid, which is a known risk factor for the development of a rash. The rashes necessitated the discontinuation of the medications. One subject also discontinued the medications due to elevated AST and ALT levels. Caution is advised for the co-administration of lopinavir/ritonavir with lamotrigine. In patients receiving concomitant therapy with lopinavir/ritonavir, the treatment regimen recommended for concomitant medications that induce lamotrigine glucuronidation (without VPA) should be used (see DOSAGE AND ADMINISTRATION, Table 6).

Olanzapine

The AUC and C_{max} of lamotrigine was reduced on average by 24% and 20%, respectively, following the addition of olanzapine (15 mg once daily) to lamotrigine (200 mg once daily) in healthy male volunteers (n = 16) compared to healthy male volunteers receiving lamotrigine alone (n = 12). This reduction in lamotrigine plasma concentrations is not expected to be clinically relevant.

Organic Cationic Transporter 2 (OCT 2)

Lamotrigine is an inhibitor of renal tubular secretion via OCT 2 proteins. This may result in increased plasma levels of certain drugs that are substantially excreted via this route. Co-administration of lamotrigine with OCT 2 substrates (i.e. procainamide and metformin) with a

narrow therapeutic index is not recommended.

Data from *in vitro* assessment of the effect of lamotrigine at OCT 2 demonstrate that lamotrigine, but not the N(2)-glucuronide metabolite, is an inhibitor of OCT 2 at potentially clinically relevant concentrations. These data demonstrate that lamotrigine is a more potent inhibitor of OCT 2 than cimetidine, with IC₅₀ values of 53.8 mcM and 186 mcM, respectively.

Oral Contraceptives

Effect of Oral Contraceptives on Lamotrigine

In a study in 16 female volunteers, an oral contraceptive preparation containing 30 mcg ethinylestradiol and 150 mcg levonorgestrel increased the apparent clearance of lamotrigine (300 mg/day) by approximately 2-fold with a mean decrease in AUC of 52% and in C_{max} of 39%. In this study, trough serum lamotrigine concentrations gradually increased and were approximately 2-fold higher on average at the end of the week of the inactive preparation compared to trough lamotrigine concentrations at the end of the active hormone cycle.

Gradual transient increases in lamotrigine levels will occur during the week of no active hormone preparation (pill-free week) for women not also taking a drug that increases the clearance of lamotrigine (carbamazepine, phenytoin, phenobarbital, primidone, or rifampin). The increase in lamotrigine levels will be greater if the dose of lamotrigine is increased in the few days before or during the pill-free week.

There have been post-marketing reports of breakthrough seizures in women taking lamotrigine and hormonal contraceptives. Dosage adjustments will be necessary in most cases for women receiving oral contraceptive preparations (see DOSAGE AND ADMINISTRATION, Women and Oral Contraceptives and ADVERSE REACTIONS, Post-Market Adverse Drug Reactions).

Effect of Lamotrigine on Oral Contraceptives

Co-administration of lamotrigine (300 mg/day) in 16 female volunteers did not affect the pharmacokinetics of the ethinylestradiol component of an oral contraceptive preparation containing 30 mcg ethinylestradiol and 150 mcg levonorgestrel. There was a mean decrease in the AUC and C_{max} of the levonorgestrel component of 19% and 12%, respectively. Measurement of serum progesterone indicated that there was no hormonal evidence of ovulation in any of the 16 volunteers, although measurement of serum FSH, LH, and estradiol indicated that there was some loss of suppression of the hypothalamic-pituitary-ovarian axis.

The effects of doses of lamotrigine other than 300 mg/day have not been studied.

The clinical significance of the observed hormonal changes on ovulatory activity is unknown. However, the possibility of decreased contraceptive efficacy in some patients cannot be excluded. Therefore, patients should be instructed to promptly report changes in their menstrual pattern (e.g. break-through bleeding).

Interactions with Other Hormonal Contraceptives or Hormone Replacement Therapy

The effect of other hormonal contraceptive preparations or hormone replacement therapy on the pharmacokinetics of lamotrigine has not been evaluated, although the effect may be similar to oral contraceptive preparations. Therefore, as for oral contraceptives, dosage adjustments will be necessary in most cases (see DOSAGE AND ADMINISTRATION: Women and Oral Contraceptives and ADVERSE REACTIONS, Post-Market Adverse Drug Reactions).

Rifampin

In a study in 10 male volunteers, rifampin (600 mg/day for 5 days) significantly increased the apparent clearance of a single 25 mg dose of lamotrigine by approximately 2-fold (AUC decreased by approximately 40%). For patients taking rifampin with phl-LAMOTRIGINE, follow the titration schedule for concomitant medications that induce lamotrigine glucuronidation (without VPA) (see DOSAGE AND ADMINISTRATION, Table 6).

Risperidone

Following the co-administration of 2 mg risperidone with lamotrigine (up to 400 mg daily), 12 out of the 14 healthy adult volunteers reported somnolence compared to 1 out of 20 when risperidone was given alone, and 0 out of 20 when lamotrigine was administered alone.

Drugs Depressing Cardiac Conduction

See WARNINGS AND PRECAUTIONS, Special Populations, Patients with Special Diseases and Conditions, Cardiac Conduction Abnormalities.

Drug-Laboratory Interactions

Lamotrigine has been reported to interfere with the assay used in some rapid urine drug screens, which can result in false positive readings, particularly for phencyclidine (PCP). A more specific alternative chemical method should be used to confirm a positive result.

Drug-Lifestyle Interactions

Dependence Liability

No evidence of abuse potential has been associated with lamotrigine, nor is there evidence of psychological or physical dependence in humans.

DOSAGE AND ADMINISTRATION

Dosing Considerations

Potential for Serious Dermatological Reactions

Do not exceed the recommended initial dose and subsequent dose escalations of phl-LAMOTRIGINE. More rapid initial titration has been associated with an increased incidence

of serious dermatological reactions (see WARNINGS AND PRECAUTIONS), as has concomitant use of valproic acid, particularly in the absence of AEDs that induce lamotrigine glucuronidation (see DRUG INTERACTIONS). Patients with a history of rash or allergy to other AEDs are more at risk for phl-LAMOTRIGINE-associated rash than those without such history (see WARNINGS AND PRECAUTIONS).

Restarting phl-LAMOTRIGINE Therapy

It is recommended that phl-LAMOTRIGINE not be restarted in patients who discontinued due to rash associated with prior treatment with phl-LAMOTRIGINE, unless the potential benefits clearly outweigh the risks. If the decision is made to restart a patient who has discontinued phl-LAMOTRIGINE for any reason, the need to restart with the initial dosing recommendations should be assessed. The greater the interval of time since the previous dose, the greater consideration should be given to restarting with the initial dosing recommendations. If a patient has discontinued phl-LAMOTRIGINE for a period of more than 5 half-lives, it is recommended that initial dosing recommendations and guidelines be followed. The half-life of lamotrigine is affected by other concomitant medications (see ACTION AND CLINICAL PHARMACOLOGY).

Withdrawal of Concomitant AEDs in Adults

Concomitant AEDs may be decreased over a 5-week period, by approximately 20% of the original dose every week. However, a slower taper may be used if clinically indicated. During this period, the dose of phl-LAMOTRIGINE administered will be dependent upon the effect of the drug being withdrawn on the pharmacokinetics of lamotrigine, together with the overall clinical response of the patient. The withdrawal of enzyme inducing AEDs (i.e. phenytoin, phenobarbital, primidone, and carbamazepine) will result in an approximate doubling of the $t_{1/2}$ of lamotrigine. Under these conditions, it may be necessary to reduce the dose of phl-LAMOTRIGINE. In contrast, the withdrawal of enzyme inhibiting AEDs (i.e. valproic acid) will result in a decrease in the $t_{1/2}$ of lamotrigine and may require an increase in the dose of phl-LAMOTRIGINE.

Recommended Dose and Dosage Adjustment

phl-LAMOTRIGINE should be added to the patient's current antiepileptic therapy.

Valproic acid more than doubles the elimination half-life of lamotrigine and reduces the plasma clearance by 50%; conversely, drugs that induce lamotrigine glucuronidation such as carbamazepine, phenytoin, phenobarbital, and primidone reduce the elimination half-life of lamotrigine by 50% and double the plasma clearance (see ACTION AND CLINICAL PHARMACOLOGY). These clinically important interactions require dosage schedules of phl-LAMOTRIGINE as summarized in Table 6 through Table 7.

Lamotrigine does not alter plasma concentrations of concomitantly administered AEDs that induce lamotrigine glucuronidation and therefore they do not usually require dose adjustment to maintain therapeutic plasma concentrations. For patients receiving phl-LAMOTRIGINE in

combination with other AEDs, an evaluation of all AEDs in the regimen should be considered if a change in seizure control or an appearance or worsening of adverse experiences is observed. If there is a need to discontinue therapy with phl-LAMOTRIGINE, a step-wise reduction of dose over at least two weeks (approximately 50% per week) is recommended unless safety concerns (i.e. rash) require a more rapid withdrawal (see WARNINGS AND PRECAUTIONS).

The relationship of plasma concentration to clinical response has not been established for lamotrigine. Dosing of phl-LAMOTRIGINE should be based on therapeutic response. In controlled clinical studies, doses of lamotrigine that were efficacious generally produced steady-state trough plasma lamotrigine concentrations of 1 to 4 mcg/mL in patients receiving one or more concomitant AEDs. Doses of lamotrigine producing this plasma concentration range were well tolerated. As with any antiepileptic drug, the oral dose of phl-LAMOTRIGINE should be adjusted to the needs of the individual patient, taking into consideration the concomitant AED therapy the patient is receiving.

Adults and Children Over 12 Years of Age

For patients taking AEDs whose pharmacokinetic interactions with phl-LAMOTRIGINE are currently unknown, follow the titration schedule for concomitant VPA (regardless of any concomitant mediation).

Table 6 Escalation regimen for phl-LAMOTRIGINE in Patients over 12 Years of Age

	Patients Taking Medications that induce lamotrigine glucuronidation ¹		Patients Taking Medications that neither induce nor inhibit lamotrigine glucuronidation ²
	with valproate ³	without valproate ³	
Weeks 1 and 2	25 mg once a day	50 mg once a day	25 mg once a day
Weeks 3 and 4	25 mg twice a day	50 mg twice a day	25 mg twice a day
Weeks 5 onwards to maintenance	Increase by 25-50 mg every 1 to 2 weeks	Increase by 100 mg every 1 to 2 weeks	Increase by 25-50 mg every 1 to 2 weeks
Usual Maintenance Dose	50-100 mg twice a day	150-250 mg twice a day	50-100 mg twice a day

^{1.} Medications that induce lamotrigine glucuronidation include carbamazepine, phenobarbital, phenytoin, primidone, rifampin, lopinavir/ritonavir and atazanavir/ritonavir

^{2.} Medications that neither inhibit nor induce lamotrigine glucuronidation include olanzapine, oxcarbazepine, felbamate, gabapentin, levetiracetam, pregabalin, topiramate and zonisamide

^{3.} Valproic acid is an inhibitor of lamotrigine glucuronidation

For patients taking valproic acid regardless of any concomitant medication, a more cautious titration schedule is available than that detailed in Table 6. During weeks 1 and 2, a dose of 25 mg every other day may be given instead of 25 mg once daily dose. During weeks 3 and 4, a dose of 25 mg once a day may be given instead of the 25 mg twice daily dose (total daily dose of 50 mg).

There have been no controlled studies to establish the effectiveness or optimal dosing regimen of add-on lamotrigine therapy in patients receiving only AEDs that neither inhibit nor induce lamotrigine glucuronidation or valproic acid. However, available data from open clinical trials indicate that the addition of lamotrigine under these conditions is associated with a higher incidence of serious rash or rash-related withdrawal, even at an initial titration dose of 12.5 mg daily (see WARNINGS AND PRECAUTIONS, Skin Related Events, Table 1 and Table 2).

The potential medical benefits of the addition of phl-LAMOTRIGINE under these conditions must be weighed against the increased risk of serious rash. If use of phl-LAMOTRIGINE under these conditions is considered clinically indicated, titration should proceed with extreme caution, especially during the first six weeks of treatment.

Starting phl-LAMOTRIGINE in Women Taking Oral Contraceptives:

Although oral contraceptives have been shown to increase the clearance of lamotrigine (see DRUG INTERACTIONS, Drug-Drug Interactions, Other Drug Interactions), no adjustments to the recommended dose escalation guidelines for phl-LAMOTRIGINE should be necessary solely based on the use of oral contraceptives. Therefore, dose escalation should follow the recommended guidelines based on whether lamotrigine is added to valproate (an inhibitor of lamotrigine glucuronidation) or an inducer of lamotrigine glucuronidation, or whether lamotrigine is added in the absence of valproate or an inducer of lamotrigine glucuronidation.

Adjustments to the Maintenance Dose of phl-LAMOTRIGINE:

Taking or Starting Oral Contraceptives: For women not taking inducers of lamotrigine glucuronidation, the maintenance dose of phl-LAMOTRIGINE will in most cases need to be increased, by as much as 2-fold over the recommended target maintenance dose (see DRUG INTERACTIONS, Drug-Drug Interactions, Other Drug Interactions). It is recommended that from the time that the hormonal contraceptive is started, the phl-LAMOTRIGINE dose is increased by 50 to 100 mg/day every week, according to the individual clinical response. Dose increases should not exceed this rate, unless the clinical response supports larger increases.

Stopping Oral Contraceptives: For women not taking inducers of lamotrigine glucuronidation (e.g., carbamazepine, phenytoin, phenobarbital, primidone, or rifampin), the maintenance dose of phl-LAMOTRIGINE will in most cases need to be decreased by as much as 50% of the maintenance dose with concurrent oral contraceptives, (see DRUG INTERACTIONS, Drug-Drug Interactions, Other Drug Interactions). It is recommended to gradually decrease the daily dose of lamotrigine by 50 to 100 mg each week (at a rate not exceeding 25% of the total daily dose per week) over a period of 3 weeks, unless the clinical response indicates otherwise.

Women and Other Hormonal Contraceptive Preparations or Hormone:

Replacement Therapy: Although the effect of other hormonal contraceptive preparations or hormone replacement therapy on the pharmacokinetics of lamotrigine has not been evaluated, the effect may be similar to oral contraceptives (see DRUG INTERACTIONS, Drug-Drug Interactions, Other Drug Interactions). Therefore, similar adjustments to the dosage of phl-LAMOTRIGINE may be needed, based on clinical response.

Pediatrics

Safety and efficacy in patients below the age of 16 years, other than those with Lennox-Gastaut Syndrome, have not been established. phl-LAMOTRIGINE is not to be administered to children under the age of 12 years due to limitations of dosage availability.

Geriatrics

No dosage adjustment from the recommended adult schedule is required. The pharmacokinetics of lamotrigine in this age group does not differ significantly from a non-elderly population (see also ACTION AND CLINICAL PHARMACOLOGY and ADVERSE REACTIONS).

Renal Impairment

The elimination half-life of lamotrigine is prolonged in patients with impaired renal function (see ACTION AND CLINICAL PHARMACOLOGY). Caution should be exercised in dose selection for patients with impaired renal function.

Hepatic Impairment

phl-LAMOTRIGINE is not to be administered to patients with hepatic impairment due to limitations of dosage availability.

Missed Dose

If a dose is missed, the patient should be instructed to take the next dose as soon as they remember unless the next dose is due in less than 4 hours, in which case they should skip the missed dose and take the next dose when it is due. The patient should be instructed to not make up for a missed dose by taking a double dose next time.

Administration

phl-LAMOTRIGINE (lamotrigine) is intended for oral administration and may be taken with or without food. phl-LAMOTRIGINE tablets should be swallowed whole and should not be chewed or crushed.

OVERDOSAGE

Adults

Acute overdoses involving quantities up to 15 g have been reported for lamotrigine, some of which have been fatal. In general, overdose has resulted in symptoms including nystagmus,

ataxia, grand mal convulsions, impaired consciousness coma and intraventricular conduction delay (QRS broadening).

Pediatrics

Among patients ≤ 16 years of age, reports of single doses of lamotrigine have included ingestion of 3 g by a 14 year old female and approximately 1 g by a 4 year old male. The 14 year old female was taking marketed lamotrigine; after the dose, she lost consciousness and was admitted to the hospital for supportive therapy, where she recovered fully (time to recovery not reported). The 4 year old male was drowsy and agitated when found, and his condition worsened to coma level II after hospitalization. He was given supportive therapy, and his condition improved rapidly with full recovery in 3 days.

There are no specific antidotes for lamotrigine. Following a suspected overdose, immediate hospitalization of the patient is advised. In the event of a very recent overdose/ingestion of a potentially life-threatening amount of the drug, emesis may be induced if indicated. General supportive care is indicated, including frequent monitoring of vital signs and close observation of the patient. It is uncertain whether hemodialysis is an effective means of removing lamotrigine from the blood. In six renal failure patients, about 20% of the amount of lamotrigine in the body was removed during 4 hours of hemodialysis.

For management of a suspected drug overdose,
contact your regional Poison Control Centre immediately.

ACTION AND CLINICAL PHARMACOLOGY

Mechanism of Action

phl-LAMOTRIGINE (lamotrigine) is a drug of the phenyltriazine class chemically unrelated to existing antiepileptic drugs (AEDs).

Lamotrigine is thought to act at voltage-sensitive sodium channels to stabilize neuronal membranes and inhibit the release of excitatory amino acid neurotransmitters (e.g. glutamate, aspartate) that are thought to play a role in the generation and spread of epileptic seizures.

Pharmacokinetics

Adults

Absorption: Lamotrigine is rapidly and completely absorbed following oral administration, reaching peak plasma concentrations 1.4 to 4.8 hours (T_{max}) post-dosing. When administered with food, the rate of absorption is slightly reduced, but the extent remains unchanged. Following single lamotrigine doses of 50–400 mg, peak plasma concentration (C_{max} =0.6–4.6 mcg/mL) and the area under the plasma concentration-versus-time curve (AUC =29.9–211 h mcg/mL) increase linearly with dose. The time-to-peak concentration,

elimination half-life ($t_{1/2}$) and volume of distribution (Vd/F) are independent of dose. The $t_{1/2}$ averages 33 hours after single doses and Vd/F ranges from 0.9 to 1.4 L/kg. Following repeated dosing in healthy volunteers for 14 days, the $t_{1/2}$ decreased by an average of 26% (mean steady state $t_{1/2}$ of 26.4 hours) and plasma clearance increased by an average of 33%. In a single-dose study where healthy volunteers were administered both oral and intravenous doses of lamotrigine, the absolute bioavailability of oral lamotrigine was 98%.

Distribution: Lamotrigine is approximately 55% bound to human plasma proteins. This binding is unaffected by therapeutic concentrations of phenytoin, phenobarbital or valproic acid. Lamotrigine does not displace other antiepileptic drugs (carbamazepine, phenytoin, phenobarbital) from protein binding sites.

Metabolism: Lamotrigine is metabolized predominantly in the liver by glucuronic acid conjugation. The major metabolite is an inactive 2-N-glucuronide conjugate that can be hydrolyzed by β -glucuronidase.

Elimination: Approximately 70% of an oral lamotrigine dose is recovered in urine as this metabolite.

Special Populations and Conditions

Pediatrics: Lamotrigine was rapidly absorbed in children, with a T_{max} ranging from 1 to 6 hours. The mean Vd/F of lamotrigine in children aged 5 to 11 years (1.3 to 1.4 L/kg) was similar to those seen in adults (0.9 to 1.4 L/kg) but was larger in younger children (1.8 to 2.3 L/kg). As with adults, the elimination of lamotrigine in pediatric patients was similarly affected by concomitant AEDs. While the CL/F was higher and $t_{1/2}$ was shorter in younger children than in older children, the mean CL/F was higher and mean $t_{1/2}$ was shorter in both pediatric groups than in adults. Population analysis results showed that the estimated apparent plasma clearances in patients aged 13 to 18 years were similar to those found in adult patients.

Geriatrics: Results of a population pharmacokinetic analysis, based on individual trials in which both adult (n = 138) and elderly (n = 13) patients with epilepsy were enrolled, indicated that the clearance of lamotrigine in elderly patients did not change to a clinically relevant extent. After single doses, apparent clearance was lower in the elderly by 12% (31 mL/min at age 70 vs. 35 mL/min at age 20). After 48 weeks of treatment, the difference in clearance was 10% (37 mL/min at age 70 vs. 41 mL/min at age 20). In addition, the pharmacokinetics of lamotrigine was studied in 12 healthy elderly volunteers who each received a single oral dose of 150 mg. The mean clearance in the elderly (0.39 mL/min) lies within the range of mean clearance values (0.31 to 0.65 mL/min) obtained in 9 studies with non-elderly adults after single doses of 30 to 450 mg (see also DOSAGE AND ADMINISTRATION and ADVERSE REACTIONS).

Renal Impairment: The pharmacokinetics of a single oral dose of lamotrigine (100 mg) was evaluated in 12 individuals with chronic renal failure (with mean creatinine clearance of 13 mL/min) who were not receiving other antiepileptic drugs. In this study, the elimination half-life of unchanged lamotrigine was prolonged (by an average of 63%) relative to individuals with

normal renal function (see WARNINGS AND PRECAUTIONS, Renal Impairment and DOSAGE AND ADMINISTRATION).

Hemodialysis: In six hemodialysis patients, the elimination half-life of unchanged lamotrigine was doubled off dialysis, and reduced by 50% on dialysis, relative to individuals with normal renal function.

Hepatic Impairment: A single dose pharmacokinetic study was performed in 24 subjects with hepatic impairment (n = 12 mild/Grade A; n = 5 moderate/Grade B and n = 7 severe/Grade C), versus 12 healthy controls. For the moderate and severe subgroups, the mean values for AUC and plasma half-life were increased approximately 2-fold and 3-fold respectively over control values, with clearance decreased proportionately. For the mild group, while mean values were not statistically different from those of controls, a subgroup of 1-to 4 subjects (dependant on pharmacokinetic parameter examined) showed abnormal individual values which were in the range of the moderately impaired subjects (see also DOSAGE AND ADMINISTRATION and WARNINGS AND PRECAUTIONS).

Gilbert's Syndrome: Gilbert's syndrome (idiopathic unconjugated hyperbilirubinemia) does not appear to affect the pharmacokinetic profile of lamotrigine.

Concomitant Antiepileptic Drugs: In patients with epilepsy, concomitant administration of lamotrigine and AEDs that induce lamotrigine glucuronidation (phenytoin, carbamazepine, primidone or phenobarbital) decreases the mean lamotrigine $t_{1/2}$ to 13 hours. Concomitant administration of lamotrigine with valproic acid significantly increases $t_{1/2}$ and decreases the clearance of lamotrigine, whereas concomitant administration of lamotrigine with valproic acid plus AEDs that induce lamotrigine glucuronidation can prolong $t_{1/2}$ up to approximately 27 hours. The key lamotrigine parameters for adult patients and healthy volunteers are summarized in Table 7, and for pediatric patients in Table 8.

Table 7 Mean Pharmacokinetic Parameters in Adult Patients with Epilepsy or Healthy Volunteers

		Healthy Young Volunteers		Patients with Epilepsy		
		Lamotrigine	Lamotrigine + Valproic Acid ²	Lamotrigine + AEDs that induce lamotrigine glucuronidation	Lamotrigine + Valproic Acid	Lamotrigine + AEDs that induce lamotrigine glucuronidation + Valproic Acid
Lamotrigine Administered						
T_{max} (hrs)	Single Dose	2.2 (0.25-12.0) ¹	1.8 (1.0-4.0)	2.3 (0.5-5.0)	4.8 (1.8-8.4)	3.8 (1.0-10.0)
	Multiple Dose	1.7 (0.5-4.0)	1.9 (0.5-3.5)	2.0 (0.75-5.93)	ND	ND
$t_{1/2}$	Single Dose	32.8 (14.0-103.0)	48.3 (31.5-88.6)	14.4 (6.4-30.4)	58.8 (30.5-88.8)	27.2 (11.2-51.6)
	Multiple Dose	25.4 (11.6-61.6)	70.3 (41.9-113.5)	12.6 (7.5-23.1)	ND	ND

		Healthy Young Volunteers		Patients with Epilepsy		
		Lamotrigine	Lamotrigine + Valproic Acid ²	Lamotrigine + AEDs that induce lamotrigine glucuronidation	Lamotrigine + Valproic Acid	Lamotrigine + AEDs that induce lamotrigine glucuronidation + Valproic Acid
Lamotrigine Administered						
Plasma Clearance (mL/min/kg)	Single Dose	0.44 (0.12-1.10)	0.30 (0.14-0.42)	1.10 (0.51-2.22)	0.28 (0.16-0.40)	0.53 (0.27-1.04)
	Multiple Dose	0.58 (0.25-1.15)	0.18 (0.12-0.33)	1.21 (0.66-1.82)	ND	ND

ND=Not done

1. Range of individual values across studies
2. Valproic Acid administered chronically (Multiple Dose Study) or for 2 days (Single Dose Study)

Table 8 Mean Pharmacokinetic Parameters in Pediatric Patients with Epilepsy

Pediatric Study Population	Number of Subjects	T _{max} (h)	t _{1/2} (h)	CL/F (mL/min/kg)
Ages 10 months-5.3 years				
Patients taking AEDs that induce lamotrigine glucuronidation	10	3.0 (1.0-5.9)	7.7 (5.7-11.4)	3.62 (2.44-5.28)
Patients taking AEDs with no known effect on drug-metabolizing enzymes	7	5.2 (2.9-6.1)	19.0 (12.9-27.1)	1.2 (0.75-2.42)
Patients taking VPA only	8	2.9 (1.0-6.0)	44.9 (29.5-52.5)	0.47 (0.23-0.77)
Ages 5 to 11 years				
Patients taking AEDs that induce lamotrigine glucuronidation	7	1.6 (1.0-3.0)	7.0 (3.8-9.8)	2.54 (1.35-5.58)
Patients taking AEDs that induce lamotrigine glucuronidation plus VPA	8	3.3 (1.0-6.4)	19.1 (7.0-31.2)	0.89 (0.39-1.93)
Patients taking VPA only*	3	4.5 (3.0-6.0)	55.4 (24.3-73.7)	0.31 (0.20-0.54)
Ages 13 to 18 years				
Patients taking AEDs that induce lamotrigine glucuronidation	11	†	†	1.3
Patients taking AEDs that induce lamotrigine glucuronidation plus VPA	8	†	†	0.5
Patients taking VPA only	4	†	†	0.3

VPA = valproic acid

* Two subjects were included in the calculation for mean t_{max}

† Parameter not estimated

Oxcarbazepine, gabapentin, levetiracetam, pregabalin, felbamate, zonisamide and topiramate did not affect the plasma concentrations of lamotrigine (see DRUG INTERACTIONS, Drug-Drug Interactions, Antiepileptic Drugs (AEDs)).

Other Drug Interactions

Chronic administration of acetaminophen was shown to slightly decrease the $t_{1/2}$ and increase the clearance of a single dose of lamotrigine. Oral contraceptives and rifampin have also been shown to increase the apparent clearance of lamotrigine (see DRUG INTERACTIONS). Co-administration of olanzapine did not have a clinically relevant effect on lamotrigine pharmacokinetics (see DRUG INTERACTIONS).

STORAGE AND STABILITY

Store between 15°C and 30°C, in a dry place. Protect from light.

DOSAGE FORMS, COMPOSITION AND PACKAGING

Tablets

- 25 mg:** Each white to off-white, shield shaped tablet debossed “P” logo over “25” on one side and scored on the other side contains 25 mg of lamotrigine, and the following non-medicinal ingredients: lactose monohydrate, magnesium stearate, microcrystalline cellulose, povidone and sodium starch glycolate. Available in bottles of 100 and 500 tablets.
- 100 mg:** Each peach coloured, shield shaped tablet debossed “P” logo over “100” on one side and scored on the other side contains 100 mg of lamotrigine, and the following non-medicinal ingredients: FD&C yellow no. 6, lactose monohydrate, magnesium stearate, microcrystalline cellulose, povidone and sodium starch glycolate. Available in bottles of 100 and 500 tablets.
- 150 mg:** Each cream coloured, shield shaped tablet debossed “P” logo over “150” on one side and scored on the other side contains 150 mg of lamotrigine, and the following non-medicinal ingredients: lactose monohydrate, magnesium stearate, microcrystalline cellulose, povidone, sodium starch glycolate and yellow iron oxide. Available in bottles of 60, 100 and 500 tablets.

PART II: SCIENTIFIC INFORMATION


PHARMACEUTICAL INFORMATION

Drug Substance

Proper Name: Lamotrigine

Chemical Name: 1,2,4-Triazine-3,5-diamine, 6-(2,3-dichlorophenyl)- [USAN]
6-(2, 3-dichlorophenyl)-1, 2, 4-triazine-3, 5-diamine [Chem. Abstr.]

Structural Formula:



Molecular Formula: C₉H₇Cl₂N₅

Molecular mass: 256.09 g/mol

Description: Lamotrigine is a white to pale cream powder. The pK_a at 25°C is 5.7. It is practically insoluble in water (0.017% w/v); slightly soluble in ethanol (0.41% w/v), chloroform (0.11% w/v) and octanol (0.28% w/v).

CLINICAL TRIALS

A comparative bioavailability study of phl-LAMOTRIGINE 150 mg tablets was performed. Pharmacokinetic and bioavailability data were measured in 24 volunteers in the *fasting* and *fed* states. The results are summarized in the following tables:

SUMMARY TABLE OF THE COMPARATIVE BIOAVAILABILITY DATA phl-LAMOTRIGINE 150 mg Tablets (Pharmel Inc., Québec, Canada) [A single 150 mg dose- 1 x 150 mg) oral administration of lamotrigine in the <i>fasting</i> state] From measured data Geometric Mean Arithmetic Mean (CV%) TEST versus REFERENCE Pharmacokinetic Parameters				
Parameter	Test phl-LAMOTRIGINE [†]	Reference*	% Ratio of Geometric Means	95% Confidence Interval
AUC _{0-72h} (ng.h/mL)	65226.2 66407.4 (19.2)	64022.0 65261.4 (20.0)	101.88	99.53-104.29
AUC ₁ (ng.h/mL)	93598.1 99490.7 (39.1)	92171.4 99980.7 (48.2)	101.55	95.76-107.68
C _{max} (ng/mL)	1862.3 1881.8 (14.3)	1906.9 1923.6 (13.5)	97.66	93.16-102.38
T _{max} ** (h)	2.00 (59.1)	1.58 (64.6)		
t _{1/2} ** (h)	42.98 (39.2)	44.13 (51.7)		

[†]phl-LAMOTRIGINE Tablets, Pharmel Inc.

*LAMICTAL[®] 150 mg Tablets (GlaxoSmithKline Inc., Mississauga, Ontario)

** For T_{max} and T_{1/2} only the arithmetic mean (CV%) is presented

SUMMARY TABLE OF THE COMPARATIVE BIOAVAILABILITY DATA
 phl-LAMOTRIGINE 150 mg Tablets (Pharmel Inc., Québec, Canada)
 [A single 150 mg dose- 1 x 150 mg) oral administration of lamotrigine in the *fed* state]
 From measured data
 Geometric Mean
 Arithmetic Mean (CV%)

TEST versus REFERENCE
Pharmacokinetic Parameters

Parameter	Test phl-LAMOTRIGINE[†]	Reference*	% Ratio of Geometric Means	95% Confidence Interval
AUC _{0-72 h} (ng.h/mL)	65459.4 66344.0 (18.3)	64305.9 65307.3 (19.3)	101.79	99.83-103.80
AUC _I (ng.h/mL)	87521.3 90442.9 (27.1)	85888.1 89053.3 (28.6)	101.90	97.80-106.18
C _{max} (ng/mL)	1904.4 1912.4 (11.0)	1865.2 1869.9 (9.7)	102.10	99.20-105.09
T _{max} ^{**} (h)	3.12 (35.0)	3.23 (32.3)		
t _{1/2} ^{**} (h)	36.27 (25.3)	36.04 (24.8)		

[†]phl-LAMOTRIGINE Tablets, Pharmel Inc.

*LAMICTAL[®] 150 mg Tablets (GlaxoSmithKline Inc., Mississauga, Ontario)

**For T_{max} and t_{1/2}, only the arithmetic mean (CV%) is presented

In adult placebo-controlled clinical studies, lamotrigine has been shown to be effective in reducing seizure frequency and the number of days with seizures when added to existing antiepileptic drug therapy in adult patients with partial seizures, with or without generalized tonic-clonic seizures that are not satisfactorily controlled.

The effectiveness of lamotrigine adjunctive therapy has also been shown in pediatric and adult patients with Lennox-Gastaut syndrome. A significant reduction in major motor seizures, drop attacks, and tonic-clonic seizures were seen following lamotrigine treatment compared with placebo treated patients. Improvements in cognitive skills (speech, nonverbal communication, alertness, attention, intellectual capacity), behaviour, and fine co-ordination have been seen with lamotrigine treatment in these patients.

Studies have also been conducted using lamotrigine monotherapy in adult patients (n = 443) newly diagnosed with epilepsy (partial seizures, with or without secondary generalization or primary generalized tonic-clonic). Results have shown comparable efficacy (time to first seizure, seizure frequency, percentage of patients seizure-free) with fewer side effects than currently approved therapies.

Clinical trials have also demonstrated that adult patients (any seizure type) can be converted to lamotrigine monotherapy from polytherapy with significant numbers of patients maintaining or improving seizure control. Efficacy was maintained during long-term treatment (up to 152 weeks).

A 24 week monotherapy trial was conducted in elderly newly diagnosed patients (102 patients received lamotrigine and 48 received carbamazepine). The findings indicate comparable efficacy, and demonstrate that lamotrigine was well tolerated in the elderly. However, the small and unbalanced number of patients in the study precludes any firm conclusions on the relative safety of the two drugs.

DETAILED PHARMACOLOGY

Animal Pharmacology

In vivo studies

In *in vivo* studies in animal models, lamotrigine has an antiepileptic profile suggesting utility in the treatment of partial seizures and generalized tonic-clonic seizures. Lamotrigine was effective in the maximal electroshock (MES), maximum pentylenetetrazol, electrically evoked after discharge (EEAD) tests and in visually evoked after discharge (VEAD) tests. In mice and rats, lamotrigine has a longer duration of action than phenytoin, carbamazepine, diazepam or valproate. Lamotrigine's potency is similar to that of phenytoin (mouse), phenobarbital (rat), carbamazepine (rat) and diazepam (mouse) (see Table 9).

Table 9 Potency and Duration of Lamotrigine Following Maximal Electroshock-Induced Seizures

Drug	ED ₅₀ for Abolition of Hind Leg Extension (mg/kg p.o.)		Duration Maintenance of Peak Activity (hours)	
	Mouse	Rat	Mouse	Rat
Lamotrigine	2.6-3.8	1.9-3.3	1-8	1-8
Phenytoin	3.5	19.7	8	1
Phenobarbital	9.1	4.7	1	1
Carbamazepine	6.9	2.5	1	0.25
Valproate	332.4	238	0.25	1
Diazepam	3.2	16.9	1	1

In single-dose mouse and rat studies, the anticonvulsant ED₅₀s of orally administered lamotrigine ranged from 1.9 to 3.8 mg/kg. Signs of CNS toxicity did not occur until high multiples of the lamotrigine ED₅₀s were reached, and consisted of ataxia and tremors (at 140 mg/kg), and convulsions (at 300 and 675 mg/kg in mice and rats, respectively). In multiple-dose studies in mice and rats, the anticonvulsant ED₅₀s were unchanged. In mice, lamotrigine was well tolerated at chronic doses up to 30 mg/kg/day. In rats, convulsions possibly related to drug administration were rarely observed (in no more than 1 of 46 to 49 animals per dose group) and did not occur until week 24 of chronic oral dosing with 15 mg/kg/day.

Preclinical Pharmacokinetics

Lamotrigine was found to accumulate in the kidney of the male rat, bind to melanin-containing ocular tissue of the pigmented rat and cynomolgus monkey, and prolong gastric emptying time in rats. In dogs, lamotrigine was extensively metabolized to a 2-N-methyl metabolite which produced dose-dependent prolongations of the P-R interval and QRS duration in ECG tracings in this species. Only trace amounts of this metabolite (< 0.6% of lamotrigine dose) were found in human urine. Clinical studies showed no evidence in humans for manifestations of these preclinical observations regarding accumulation in the kidney, melanin binding, gastric emptying, or cardiac effects.

In vitro Studies

In vitro pharmacological studies suggest that lamotrigine acts at voltage-sensitive sodium channels to stabilize neuronal membranes and inhibit the release of excitatory amino acid neurotransmitters (e.g. glutamate, aspartate) that are thought to play a role in the generation and spread of epileptic seizures.

Lamotrigine was shown to be a weak inhibitor of dihydrofolate reductase *in vitro*, hence there is a possibility of interference with folate metabolism during long-term therapy (see WARNINGS AND PRECAUTIONS, Special Populations, Pregnant Women). In clinical studies, lamotrigine did not affect blood folate concentrations or associated hematologic parameters.

TOXICOLOGY

Acute Toxicity

Single-dose lethality values were calculated in male and female mice and rats receiving lamotrigine by the oral and intravenous routes of administration. The calculated LD₅₀ values are listed in Table 10.

Table 10 Acute Toxicity Studies

Route	LD ₅₀ Dose (mg/kg)			
	Mouse		Rat	
	Male	Female	Male	Female
Oral	245	292	205	163
I.V.	141	134	107	112

The lowest doses (mg/kg) at which any deaths occurred were 300 (oral) and 125 (I.V.) in mice and 140 (oral) and 100 (I.V.) in rats. Deaths were seen as early as 1 minute following I.V. dosing and 30 minutes following oral dosing. The most severe sign noted was clonic convulsions (rats only). Other signs, including tremors, ataxia, hypoactivity, decreased respiration, and hypothermia were also observed. Survivors generally recovered within 24 hours, but hypoactivity lasted for several days in some animals.

Long-Term Toxicity

Subacute to subchronic (14–30 days) studies were conducted in rats (oral and intravenous), marmosets (oral) and cynomolgus monkeys (intravenous).

Effects seen in rats which were considered to be consistently associated with oral lamotrigine administration included specific nephropathy (males, 1 mg/kg/day), increased weight of stomach contents (6.25 mg/kg/day), increased water consumption and urine output (10 mg/kg/day), reduced weight gain and food consumption (22.5 mg/kg/day), and convulsions (50 mg/kg/day).

Marmosets were given either single daily lamotrigine doses of 10-100 mg/kg/day or 3 daily doses ranging from 10 to 50 mg/kg/dose. The following effects were observed at the following lowest doses given: slight decreases in WBC, RBC and related values (100 mg/kg/day or 22.5 mg/kg/dose t.i.d.); post-dose incoordination, slight body weight loss, decreased food consumption (50 mg/kg/day or 22.5 mg/kg/dose t.i.d.); post-dose salivation and vomiting (10 mg/kg/day or 22.5 mg/kg/dose t.i.d.). Blood, brain and liver folate levels were not affected.

In a 14-day intravenous study in cynomolgus monkeys, daily lamotrigine doses ranged from 5 to 20 mg/kg. The following effects were observed (shortly after dosing) at the following lowest doses given: ataxia, vomiting, and decreased food consumption (10 mg/kg); nystagmus (15 mg/kg); agitation, slight body weight loss (20 mg/kg).

In chronic oral toxicity studies, mice were given daily lamotrigine doses of up to 60 mg/kg for 90 days. Rats were given doses of up to 30 mg/kg/day for 90 days, up to 25 mg/kg/day for 6

months, and up to 15 mg/kg/day for one year. The only effect seen in mice was increased stomach content weight, likely related to delay gastric emptying (30 and 60 mg/kg/day). The most prominent drug-related effects in rats were renal histopathological changes, seen at all dose levels; other effects seen were similar to those documented in the subacute and subchronic rat studies. Other species were also evaluated by light microscopic examination of kidney sections from animals given lamotrigine and by *in vitro* studies of kidney cortex slice uptake of radiolabelled lamotrigine. The studies demonstrated that the renal effects seen are limited to male rats and represent exacerbation of spontaneous hyaline droplet changes related to the renal proximal tubular handling of α_{2U} -globulin. This globulin is a specific protein synthesized by the liver in male rats only. The hyaline droplets are secondary lysosomes and contain α_{2U} -globulin. Lamotrigine is also accumulated in the renal proximal tubular epithelium in the form of rectilinear and crystalline inclusions which likely contain both α_{2U} -globulin and lamotrigine. Time-and-dose dependent loading of these cells leads to cellular degeneration and eventual regeneration. Changes are consistent with those seen in "light hydrocarbon nephropathy," a male-rat-specific condition, which is reversible following termination of treatment and has no human counterpart.

Cynomolgus monkeys were treated with daily oral lamotrigine doses of up to 20 mg/kg in 13-week, 26-week, and 52-week studies. Effects noted were limited to animals given 20 mg/kg/day and included postdose ataxia, lethargy, trembling, locomotor incoordination and convulsions in some animals in the 26-week study. A reduced rate of body weight gain and transient weight loss was seen at doses as low as 5 mg/kg/day, but only in the 52-week study.

Carcinogenicity

There was no evidence of carcinogenicity in mice treated orally with lamotrigine (10-30 mg/kg/day) for up to 106 weeks. Lamotrigine was not carcinogenic in two rat studies where animals received doses of 1–10 mg/kg/day for up to 104 weeks (females) or 112 weeks (males). In both studies, survival was not affected by treatment.

Mutagenicity

In vitro, lamotrigine was not mutagenic in microbial (Ames) or mammalian (mouse lymphoma) mutagenicity tests with or without metabolic activation. Lamotrigine had no effect on the incidence of chromosomal abnormalities in cultured human lymphocytes exposed to concentrations of up to 1000 mcg/mL in the presence or absence of S9 metabolic activation. Concentrations of 500 and 1000 mcg/mL without activation were cytotoxic.

In vivo, there was no increase in the incidence of chromosomal abnormalities in bone marrow cells of rats given doses of lamotrigine of up to 200 mg/kg.

Reproduction and Teratology

No evidence of teratogenicity was observed in mice, rats, and rabbits given oral doses of lamotrigine at up to 14, 4, and 4 times, respectively, the currently recommended upper human maintenance dose (500 mg/day or 7 mg/kg/day). This was true when lamotrigine was given during the period of major organogenesis or was started prior to and continued throughout the period of organogenesis. In these same oral dosing studies maternal toxicity and secondary fetal

toxicity, resulting in reduced fetal weight and/or delayed ossification were seen. Teratology studies were also conducted using bolus intravenous (I.V.) administration of the isethionate salt of lamotrigine in multiples of the projected human oral dose. Intravenous lamotrigine resulted in convulsions or impaired coordination in rat and rabbit dams at 30 mg/kg and 15 mg/kg, respectively. The 30 mg/kg dose also produced an increased incidence of intrauterine death without signs of teratogenicity in rats only.

When rats were dosed prior to and during mating, and throughout gestation and lactation at daily oral doses of 5, 10, and 20 mg/kg, gestation was slightly longer in the dams allowed to deliver in the 20 mg/kg/group (22.0 ± 0.0 versus 21.5 ± 0.5 days in non-treated controls). These doses are approximately 1, 1.5, and 3 times the currently recommended upper human maintenance dose (500 mg/day or 7 mg/kg/day). In this same study, body weight gain and food consumption of the parent generation (F₀) dams dosed at 20 mg/kg/day were less than the control dams and were indicative of maternal toxicity. There was no evidence of teratogenic effects in the litters of the dams designated for cesarean section. Effects secondary to maternal toxicity consisted of a decrease in mean fetal weight and length of 20 mg/kg pups; the incidence of two skeletal variants was increased and the incidence of one skeletal variant was decreased. When pregnant rats of the same strain were dosed only on days 15–20 of gestation at the same daily doses of 5, 10, and 20 mg/kg, more pronounced maternal toxicity, than noted in the previously described study at the same doses, was seen in dams given 10 and 20 mg/kg doses and consisted of dehydration, hypothermia, and decreased weight gain and food consumption. A smaller decrease in body weight gain was seen in the 5 mg/kg group. Gestation was prolonged in the 20 mg/kg group (22.6 vs. 22.0 days in non-treated controls) and secondary to maternal toxicity there were increased numbers of stillborn pups (partial to entire litters) in the 10 and 20 mg/kg groups and increased early neonatal mortality.

Even at maternally toxic levels leading to fetal death, there was no evidence of teratogenicity. Because animal reproduction studies are not always predictive of human response, this drug should be used during pregnancy only if clearly needed. Clinical data indicate that lamotrigine has no effect on blood folate concentrations in adults; however, the effects of lamotrigine on fetal blood folate levels *in utero* are unknown.

REFERENCES

1. Anon. Lamotrigine. *Drugs Future* 1987; 12(6):595-596.
2. Betts T, Goodwin G, Withers RM, Yuen AW. Human safety of lamotrigine. *Epilepsia* 1991; 32 Suppl 2:S17-S21.
3. Binnie CD, van Emde BW, Kasteleijn-Nolste-Trenite DG, de Korte RA, Meijer JW, Meinardi H et al. Acute effects of lamotrigine (BW430C) in persons with epilepsy. *Epilepsia* 1986; 27(3):248-254.
4. Binnie CD, Debets RM, Engelsman M, Meijer JW, Meinardi H, Overweg J et al. Double-blind crossover trial of lamotrigine (Lamictal) as add-on therapy in intractable epilepsy. *Epilepsy Res* 1989; 4(3):222-229.
5. Binnie CD, Van Wieringen A, Meijer JW, Peck AW, de Vries J. Comparison of the effects of lamotrigine and phenytoin on the EEG power spectrum and cortical and brainstem-evoked responses of normal human volunteers. *Neuropsychobiology* 1989; 21(3):157-169.
6. Binnie CD. Lamotrigine. *Comprehensive Epileptology* Ed M Dam and L Gram. New York, Raven Press, 1991;665-670.
7. Brodie MJ, Richens A, Yuen AW. Double-blind comparison of lamotrigine and carbamazepine in newly diagnosed epilepsy. UK Lamotrigine/Carbamazepine Monotherapy Trial Group. *Lancet* 1995; 345(8948):476-479.
8. Brodie MJ. Lamotrigine monotherapy: An overview. *R Soc Med: Int Congr Symp Ser* 1996;(214):43-49.
9. Brodie MJ, Overstall PW, Giorgi L. Multicentre, double-blind, randomised comparison between lamotrigine and carbamazepine in elderly patients with newly diagnosed epilepsy. The UK Lamotrigine Elderly Study Group. *Epilepsy Res* 1999; 37(1):81-87.
10. Cohen AF, Ashby L, Crowley D, Land G, Peck AW, Miller AA. Lamotrigine (BW430C), a potential anticonvulsant. Effects on the central nervous system in comparison with phenytoin and diazepam. *Br J Clin Pharmacol* 1985; 20(6):619-629.
11. Cohen AF, Land GS, Breimer DD, Yuen WC, Winton C, Peck AW. Lamotrigine, a new anticonvulsant: pharmacokinetics in normal humans. *Clin Pharmacol Ther.* 1987; 42(5):535-541.
12. Depot M, Powell JR, Messenheimer JA, Jr., Cloutier G, Dalton MJ. Kinetic effects of multiple oral doses of acetaminophen on a single oral dose of lamotrigine. *Clin Pharmacol Ther* 1990; 48(4):346-355.

13. Donaldson JA, Glauser TA, Olberding LS. Lamotrigine adjunctive therapy in childhood epileptic encephalopathy (the Lennox Gastaut syndrome). *Epilepsia* 1997; 38(1):68-73.
14. Dulac O, Kaminska A. Use of lamotrigine in Lennox-Gastaut and related epilepsy syndromes. *J Child Neurol* 1997; 12 Suppl 1:S23-S28.
15. Fitton A, Goa KL. Lamotrigine. An update of its pharmacology and therapeutic use in epilepsy. *Drugs* 1995; 50(4):691-713.
16. Goa KL, Ross SR, Chrisp P. Lamotrigine. A review of its pharmacological properties and clinical efficacy in epilepsy. *Drugs* 1993; 46(1):152-176.
17. Jawad S, Oxley J, Yuen WC, Richens A. The effect of lamotrigine, a novel anticonvulsant, on interictal spikes in patients with epilepsy. *Br J Clin Pharmacol* 1986; 22(2):191-193.
18. Jawad S, Richens A, Goodwin G, Yuen WC. Controlled trial of lamotrigine (Lamictal) for refractory partial seizures. *Epilepsia* 1989; 30(3):356-363.
19. Leach MJ, Baxter MG, Critchley MA. Neurochemical and behavioral aspects of lamotrigine. *Epilepsia* 1991; 32 Suppl 2:S4-S8.
20. Loiseau P, Yuen AW, Duche B, Menager T, Arne-Bes MC. A randomised double-blind placebo-controlled crossover add-on trial of lamotrigine in patients with treatment-resistant partial seizures. *Epilepsy Res* 1990; 7(2):136-145.
21. Matsuo F, Bergen D, Faught E, Messenheimer JA, Dren AT, Rudd GD et al. Placebo-controlled study of the efficacy and safety of lamotrigine in patients with partial seizures. U.S. Lamotrigine Protocol 0.5 Clinical Trial Group. *Neurology* 1993; 43(11):2284-2291.
22. Messenheimer J, Ramsay RE, Willmore LJ, Leroy RF, Zielinski JJ, Mattson R et al. Lamotrigine therapy for partial seizures: a multicenter, placebo-controlled, double-blind, cross-over trial. *Epilepsia* 1994; 35(1):113-121.
23. Mikati MA, Schachter SC, Schomer DL, Keally M, Osborne-Shafer P, Seaman CA et al. Long-term tolerability, pharmacokinetic and preliminary efficacy study of lamotrigine in patients with resistant partial seizures. *Clin Neuropharmacol* 1989; 12(4):312-321.
24. Miller AA, Sawyer DA, Roth B, Peck AW, Leach MJ, Wheatley PL et al. Lamotrigine. *New Anticonvulsant Drugs* Meldrum BS, Porter RJ, Eds London: Libbey, 1986;165-177.
25. Motte J, Trevathan E, Arvidsson JF, Barrera MN, Mullens EL, Manasco P. Lamotrigine for generalized seizures associated with the Lennox-Gastaut syndrome. Lamictal Lennox-Gastaut Study Group. *N Engl J Med* 1997; 337(25):1807-1812.
26. Peck AW. Clinical pharmacology of lamotrigine. *Epilepsia* 1991; 32 Suppl 2:S9-12.

27. Posner J, Cohen AF, Land G, Winton C, Peck AW. The pharmacokinetics of lamotrigine (BW430C) in healthy subjects with unconjugated hyperbilirubinaemia (Gilbert's syndrome). *Br J Clin Pharmacol* 1989; 28(1):117-120.
28. Posner J, Holdich T, Crome P. Comparison of lamotrigine pharmacokinetics in young and elderly healthy volunteers. *J Pharm Med* 1991; 1(3):121-128.
29. Richens A, Yuen AW. Overview of the clinical efficacy of lamotrigine. *Epilepsia* 1991; 32 Suppl 2:S13-S16.
30. Richens A. Pharmacokinetics of lamotrigine. In: Richens A., ed. *Clinical update on lamotrigine: a novel antiepileptic agent*. Wells Medical Limited, Kent, 1992;21-27.
31. Sander JW, Patsalos PN, Oxley JR, Hamilton MJ, Yuen WC. A randomised double-blind placebo-controlled add-on trial of lamotrigine in patients with severe epilepsy. *Epilepsy Res* 1990; 6(3):221-226.
32. Schlumberger E, Chavez F, Palacios L, Rey E, Pajot N, Dulac O. Lamotrigine in treatment of 120 children with epilepsy. *Epilepsia* 1994; 35(2):359-367.
33. Smith D, Baker G, Davies G, Dewey M, Chadwick DW. Outcomes of add-on treatment with lamotrigine in partial epilepsy. *Epilepsia* 1993; 34(2):312-322.
34. Timmings PL, Richens A. Lamotrigine as an add-on drug in the management of Lennox-Gastaut syndrome. *Eur Neurol* 1992; 32(6):305-307.
35. Uvebrant P, Bauziene R. Intractable epilepsy in children. The efficacy of lamotrigine treatment, including non-seizure-related benefits. *Neuropediatrics* 1994; 25(6):284-289.
36. Yuen AW. Lamotrigine: a review of antiepileptic efficacy. *Epilepsia* 1994; 35 Suppl 5:S33-S36.
37. Product Monograph: LAMICTAL[®] (lamotrigine) by GlaxoSmithKline Inc. Date of Revision: March 23, 2015. Control No.: 179706

PART III: CONSUMER INFORMATION

Pr **phl-LAMOTRIGINE**
Lamotrigine Tablets, USP

This leaflet is part III of a "Product Monograph" published when phl-LAMOTRIGINE was approved for sale in Canada and is designed specifically for Consumers. This leaflet is a summary and will not tell you everything about phl-LAMOTRIGINE. Contact your doctor or pharmacist if you have any questions about the drug.

The information provided below is for patients, or parents of patients, who will be receiving phl-LAMOTRIGINE. Please read the following information carefully before you start to take phl-LAMOTRIGINE, even if you have taken this drug before. Please do not discard this leaflet; you may need to read it again.

ABOUT THIS MEDICATION

What the medication is used for:

phl-LAMOTRIGINE has been prescribed to you/your child to control your/their epilepsy. Please follow your doctor's recommendations carefully.

What it does:

phl-LAMOTRIGINE affects chemicals in the brain that are involved in sending signals to the nerves. phl-LAMOTRIGINE helps to control epileptic seizures.

When it should not be used:

Do NOT take or use phl-LAMOTRIGINE if you are hypersensitive (allergic) to lamotrigine or any of the other ingredients of phl-LAMOTRIGINE tablets (see What the non-medicinal ingredients are).

What the medicinal ingredient is:

Lamotrigine

What the non-medicinal ingredients are:

Lactose monohydrate, magnesium stearate, microcrystalline cellulose, povidone, sodium starch glycolate, and the following:

100 mg tablets also include: FD&C yellow no. 6

150 mg tablets also include: yellow iron oxide.

What dosage forms it comes in:

Tablets: 25 mg, 100 mg and 150 mg.

WARNINGS AND PRECAUTIONS

Serious Warnings and Precautions – Serious Skin Rashes

Serious skin rashes requiring hospitalization have occurred with the use of lamotrigine. Symptoms include blistering, peeling of the lips, eyes, mouth or genitals, swelling of the face and/or tongue, hives, swollen lymph nodes and fever. If you or your child develops these symptoms seek immediate medical help.

In children serious skin rashes can be mistaken for an infection. If your child develops a rash and fever while taking phl-LAMOTRIGINE contact your doctor immediately.

Do NOT take more tablets than has been prescribed by your doctor. There is a higher chance of serious skin reactions occurring when more than the recommended number of tablets is taken.

There is a higher chance of developing a serious skin rash if you have a history of allergy or rash to other anti-epileptic drugs. If you or your child has had a serious skin reaction or allergy to another anti-epileptic drug you must tell your doctor.

BEFORE you use phl-LAMOTRIGINE talk to your doctor or pharmacist if you or your child:

- Ever had an unusual or allergic reaction to phl-LAMOTRIGINE.
- Are/is allergic to any component of phl-LAMOTRIGINE tablets.
- Have ever had a rash during previous treatment with lamotrigine or with any other antiepileptic drug.
- Have ever developed meningitis after taking lamotrigine
- Are/is pregnant or are planning to become pregnant. There is a risk of malformations (e.g. cleft lip or cleft palate) associated with the use of lamotrigine in the first few months of pregnancy. Taking folic acid when planning pregnancy and during early pregnancy may be considered. If you take phl-LAMOTRIGINE while pregnant, you may register yourself at the North American Antiepileptic Drug Pregnancy Registry (see contact number at the end of this information)
- Are/is breast-feeding (nursing). Lamotrigine passes into breast milk and may cause side-effects in a breast-fed baby. If the decision by you and your healthcare provider is to breast-feed while taking phl-LAMOTRIGINE, watch your baby closely for symptoms such as trouble breathing, episodes of temporarily stopping breathing, sleepiness or poor sucking. Call your baby's healthcare provider right away if you see any of these problems
- Are/is taking any other prescription or over-the-counter medicine, including birth control pills or other female hormonal products.
- Have/has liver, kidney disease, heart problems or other medical conditions.
- Consume alcohol on a regular basis.

A small number of people taking lamotrigine get an allergic reaction or potentially serious skin reaction, which may develop into more serious problems such as organ failure if not treated. You need to know the symptoms to look out for while you are taking phl-LAMOTRIGINE. Please read the Serious Side Effects, How Often Do They Happen and What to do

About Them section of this leaflet for more information.

There have been a small number of reports of suicidal behaviour (including suicidal thoughts and suicide attempts) in patients being treated with anti-epileptic medicines such as lamotrigine. If at any time you have these thoughts, contact your doctor immediately. **Do not discontinue phl-LAMOTRIGINE on your own.**

The seizures in some types of epilepsy may occasionally become worse or happen more often while you're taking phl-LAMOTRIGINE. Some patients may experience severe seizures, which may cause serious health problems. If your seizures happen more often or if you experience a more severe seizure while you're taking phl-LAMOTRIGINE, contact a doctor as soon as possible.

Do not stop taking/giving phl-LAMOTRIGINE unless directed by your doctor. Always check that you have an adequate supply of phl-LAMOTRIGINE. Remember that this medicine was prescribed only for you/your child; never give it to anyone else.

Taking the wrong medication can cause serious health problems. When your healthcare provider gives you a prescription for phl-LAMOTRIGINE, make sure you can read it clearly and talk to your pharmacist to check that you are given the correct medicine.

Driving and using machines: Patients with uncontrolled epilepsy should not drive or operate machinery. Before doing tasks which require special attention, wait until you know how you respond to phl-LAMOTRIGINE. Dizziness, poor coordination, drowsiness, double vision and blurred vision have been reported in people taking lamotrigine.

INTERACTIONS WITH THIS MEDICATION

Consult your doctor before taking/giving your child any other medication, including over-the-counter medicines, herbal and alternative medicines. Some drugs can produce various side-effects when they are used in combination with phl-LAMOTRIGINE.

Do not start or stop using birth control pills or other female hormonal products, without consulting with your doctor.

Tell your doctor as soon as possible if you experience changes in your menstrual pattern (e.g. break-through bleeding) while taking phl-LAMOTRIGINE and birth control pills or other female hormonal products.

phl-LAMOTRIGINE may interfere with some laboratory tests to detect other drugs. If you require a laboratory test, tell your doctor or hospital that you are taking phl-LAMOTRIGINE.

Drug-Drug Interactions

Drugs that may interact with phl-LAMOTRIGINE include:

- Valproic acid (valproate)
- Carbamazepine
- Phenytoin
- Phenobarbital

- Primidone
- Risperidone
- Rifampin
- Procainamide
- Metformin
- Combination of lopinavir and ritonavir, or atazanavir and ritonavir
- Birth control pills or other female hormonal products.

PROPER USE OF THIS MEDICATION

Usual dose:

It is very important that you/your child take phl-LAMOTRIGINE exactly as your doctor instructed.

Your doctor may increase or decrease your/your child's medication according to your/their specific needs. Carefully follow the instructions you were given. Do not change the dose yourself.

If you plan to start or stop taking hormonal contraceptives or other female hormonal products, your doctor will give you specific instructions on the dose of phl-LAMOTRIGINE.

Do not stop taking your medication abruptly, because your/your child's seizures may increase. Speak to your doctor before stopping phl-LAMOTRIGINE.

It is important to keep your/your child's appointments for medical checkups.

phl-LAMOTRIGINE may be taken with or without food.

phl-LAMOTRIGINE tablets should be swallowed whole and should not be chewed or crushed.

Overdose:

If you think you/your child has taken too much phl-LAMOTRIGINE, contact your healthcare professional, hospital emergency department or regional Poison Control Centre immediately, even if there are no symptoms.

Missed Dose:

If you/your child happen to miss a dose, take it as soon as you remember. If your next dose is less than 4 hours away, skip the missed dose and take the next dose at the normal time. Do not try to make up for it by doubling up on the next dose. Try not to miss any more doses. Ask your doctor for advice on how to start taking it again, even if you only stop for a few days.

SIDE EFFECTS AND WHAT TO DO ABOUT THEM

Side effects include:

- double vision, blurred vision
- headache
- tremor, poor coordination
- dizziness
- nausea, vomiting, stomach upset

- pain in your neck, abdomen, or joints
- feeling of weakness or tiredness (fatigue)
- sleepiness/drowsiness
- difficulty sleeping (insomnia)
- nasal congestion
- skin rash
- aggression, agitation or irritability

SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM

Symptom/ effect		Talk with your doctor or pharmacist right away		Seek immediate emergency medical assistance
		Only if severe	In all cases	
Very common	Skin rashes or redness		√	
	Severe Skin Reactions: skin rashes or redness developing into widespread rash with blisters and peeling skin, particularly around the mouth, nose, eyes and genitals, sore mouth or eyes, a high temperature (fever), flu-like symptoms or drowsiness			√
Rare	Choreoathetosis: abnormal uncontrollable muscle movements that may involve the face, eyes (nystagmus), neck, trunk, arms or legs	√		
	Conjunctivitis: itchy eyes with discharge and crusty eyelids		√	
	Aseptic meningitis: fever, nausea, vomiting, headache, stiff neck and extreme sensitivity to bright light			√
	Hallucinations: seeing or hearing things that aren't really there		√	
Very rare	Nightmares	√		
	Liver and blood problems: yellowing of the skin, itching, abdominal pain and/or tenderness, feeling very tired, unexpected bleeding or bruising or the fingers turning blue, a sore throat or more infections (such as colds) than usual		√	

	Lymphadenopathy: swelling of the face or swollen glands in the neck, armpits or groin		√	
	Serious blood clotting disorder: unexpected or prolonged bleeding, including from your gums or nose; blood in your urine; and unexpected bruising, including in the form of small dots on the skin.			√
Not known	Thoughts of suicide or hurting yourself		√	
	Seizures happening more often in people who already have epilepsy			√

This is not a complete list of side effects. For any unexpected effects while taking phl-LAMOTRIGINE contact your doctor or pharmacist.

HOW TO STORE IT

Store between 15°C and 30°C, in a dry place. Protect from light. Cap the bottle tightly immediately after use. **Keep out of sight and reach of children.**

REPORTING SIDE EFFECTS

You can help improve the safe use of health products for Canadians by reporting serious and unexpected side effects to Health Canada. Your report may help to identify new side effects and change the product safety information.

3 ways to report:

- Online at [MedEffect](#);
- By calling 1-866-234-2345 (toll-free);
- By completing a Consumer Side Effect Reporting Form and sending it by:
 - Fax to 1-866-678-6789 (toll-free), or
 - Mail to: Canada Vigilance Program
Health Canada, Postal Locator 0701E
Ottawa, ON
K1A 0K9
 Postage paid labels and the Consumer Side Effect Reporting Form are available at [MedEffect](#).

NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

MORE INFORMATION

This document plus the full product monograph, prepared for health professionals, can be obtained by contacting Pharmel Inc. at 1-888-550-6060.

This leaflet was prepared by

Pharmel Inc.
Montréal Québec
H4P 2T4

Last revised: June 15, 2016