

## PRODUCT MONOGRAPH

### <sup>Pr</sup>**pendo-PRAVASTATIN**

Pravastatin Sodium Tablets, House Standard

10 mg and 20 mg

Lipid Metabolism Regulator

PENDOPHARM, division of Pharmascience Inc.  
6111 Royalmount Ave., Suite 100  
Montréal, Québec  
H4P 2T4

Date of Revision:  
July 25, 2016

[www.pendopharm.com](http://www.pendopharm.com)

Submission Control No: 196360

## Table of Contents

<b>PART I: HEALTH PROFESSIONAL INFORMATION .....</b>	<b>3</b>
SUMMARY PRODUCT INFORMATION .....	3
INDICATIONS AND CLINICAL USE .....	3
CONTRAINDICATIONS .....	5
WARNINGS AND PRECAUTIONS .....	5
ADVERSE REACTIONS .....	10
DRUG INTERACTIONS .....	14
DOSAGE AND ADMINISTRATION .....	16
OVERDOSAGE .....	17
ACTION AND CLINICAL PHARMACOLOGY .....	17
STORAGE AND STABILITY .....	19
DOSAGE FORMS, COMPOSITION AND PACKAGING .....	19
<b>PART II: SCIENTIFIC INFORMATION .....</b>	<b>21</b>
PHARMACEUTICAL INFORMATION .....	21
CLINICAL TRIALS .....	22
DETAILED PHARMACOLOGY .....	28
TOXICOLOGY .....	30
REFERENCES .....	34
<b>PART III: CONSUMER INFORMATION .....</b>	<b>39</b>

# Pr<sup>r</sup>pendo-PRAVASTATIN

Pravastatin Sodium Tablets, House Standard

## PART I: HEALTH PROFESSIONAL INFORMATION

### SUMMARY PRODUCT INFORMATION

Route of Administration	Dosage Form / Strength	All non-medicinal Ingredients
oral	Tablets 10 mg and 20 mg	Colloidal Silicon Dioxide, Copovidone, Croscarmellose Sodium, Dibasic Calcium Phosphate, Iron Oxide Red (10 mg only), Iron Oxide Yellow (20 mg only), Lactose, Magnesium Stearate, Microcrystalline Cellulose, Polyethylene Glycol.

### INDICATIONS AND CLINICAL USE

Therapy with lipid-altering agents should be considered a component of multiple risk factor intervention in those individuals at increased risk for atherosclerotic vascular disease due to dyslipidemia. pendo-PRAVASTATIN (pravastatin sodium) should be used in addition to a diet restricted in saturated fat and cholesterol when the response to diet and other non-pharmacological measures alone has been inadequate.

#### Hypercholesterolemia

pendo-PRAVASTATIN is indicated as an adjunct to diet (at least an equivalent of the Adult Treatment Panel III [ATP III TLC diet]) for the reduction of elevated Total and Low Density Lipoprotein Cholesterol (LDL-C) levels in patients with primary hypercholesterolemia (Types IIa and IIb), when the response to diet and other non-pharmacologic measures alone has been inadequate.

Prior to initiating therapy with pendo-PRAVASTATIN, secondary causes for hypercholesterolemia, such as obesity, poorly controlled diabetes mellitus, hypothyroidism, nephrotic syndrome, dysproteinemias, obstructive liver disease, other drug therapy or alcoholism, should be excluded and it should be determined that patients for whom treatment with pendo-PRAVASTATIN is being considered have an elevated LDL-C level as the cause for an elevated total serum cholesterol. A lipid profile should be performed to measure Total Cholesterol, High Density Lipoprotein Cholesterol (HDL-C) and Triglycerides (TG).

For patients with total triglycerides less than 4.52 mmol/L (400 mg/dL), LDL-C can be estimated using the following equation:

$$\text{LDL-C (mmol/L)} = \text{Total Cholesterol} - [(0.37 \times \text{triglycerides}) + \text{HDL-C}]$$

$$\text{LDL-C (mg/dL)} = \text{Total Cholesterol} - [(0.16 \times \text{triglycerides}) + \text{HDL-C}]$$

When total triglyceride levels exceed 4.52 mmol/L (400 mg/dL), this equation is less accurate and LDL-C concentrations should be determined by ultracentrifugation.

As with other lipid-lowering therapy, pendo-PRAVASTATIN is not indicated when hypercholesterolemia is due to hyperalphalipoproteinemia (elevated HDL-C). The efficacy of pravastatin has not been evaluated in conditions where the major abnormality is elevation of chylomicrons, VLDL or LDL (i.e. hyperlipoproteinemia or dyslipoproteinemia types I, III, IV or V).

### **Primary Prevention of Coronary Events**

In hypercholesterolemic patients without clinically evident coronary heart disease, pendo-PRAVASTATIN is indicated to:

- Reduce the risk of myocardial infarction;
- Reduce the risk for undergoing myocardial revascularization procedures;
- Reduce the risk of total mortality by reducing cardiovascular deaths.

### **Secondary Prevention of Cardiovascular Events**

In patients with total cholesterol in the normal to moderately elevated range who have clinically evident coronary heart disease, pendo-PRAVASTATIN is indicated to:

- Reduce the risk of total mortality
- Reduce the risk of death due to coronary heart disease
- Reduce the risk of myocardial infarction
- Reduce the risk of undergoing myocardial revascularization procedures
- Reduce the risk of stroke and transient ischemic attack (TIA).
- Reduce total hospitalization

Pravastatin sodium was also found to reduce the rate of progression of atherosclerosis in patients with coronary heart disease as part of a treatment strategy to lower Total and LDL-cholesterol to desired levels. In two trials including this type of patients<sup>1</sup> (i.e. in a secondary prevention intervention), pravastatin sodium monotherapy was shown to reduce the rate of progression of atherosclerosis as evaluated by quantitative angiography and B-mode ultrasound. This effect may be associated with an improvement in the coronary endpoints (fatal or non fatal myocardial infarction). In these trials, however, no effect was observed in all cause mortality (see CLINICAL TRIALS - Atherosclerotic disease Progression).

### **Pediatric Use (< 16 years of age)**

There is no experience to date with the use of pravastatin sodium in such patients. Treatment in these patients is not recommended at this time.

### **Elderly (≥ 65 years of age)**

Pharmacokinetic evaluation of pravastatin in patients over the age of 65 years indicates an increased AUC. As a precautionary measure, the lowest dose should be administered initially in

---

<sup>1</sup> Pravastatin Limitation of Atherosclerosis in the Coronary/Carotid Arteries (PLAC I and II)

these patients (see REFERENCES).

## CONTRAINDICATIONS

Patients who are hypersensitive to this drug or to any ingredient in the formulation.

Active liver disease or unexplained persistent elevations of serum transaminases exceeding 3 times the upper limit of normal (ULN) (see WARNINGS AND PRECAUTIONS).

### **In Pregnant and Nursing Women**

Cholesterol and other products of cholesterol biosynthesis are essential components for fetal development (including synthesis of steroids and cell membranes). Since HMG-CoA reductase inhibitors such as pendo-PRAVASTATIN (pravastatin sodium) decrease cholesterol synthesis and possibly the synthesis of other biologically active substances derived from cholesterol, they may cause fetal harm when administered to pregnant women. pendo-PRAVASTATIN should be administered to women of childbearing age only when such patients are highly unlikely to conceive and have been informed of the possible harm. If the patient becomes pregnant while taking pendo-PRAVASTATIN, the drug should be discontinued immediately and the patient apprised of the potential harm to the fetus. Atherosclerosis being a chronic process, discontinuation of lipid metabolism regulating drugs during pregnancy should have little impact on the outcome of long-term therapy of primary hypercholesterolemia (see WARNINGS AND PRECAUTIONS - Use in Pregnancy, Use in Nursing Mothers).

## WARNINGS AND PRECAUTIONS

### **Muscle Effects**

Elevations of creatinine phosphokinase levels (CK [MM fraction]), have been reported with the use of HMG-CoA reductase inhibitors, including pravastatin sodium.

Effects on skeletal muscle such as myalgia, myopathy and, rarely, rhabdomyolysis have been reported in patients treated with pravastatin sodium.

Muscle weakness and rhabdomyolysis have been reported in patients receiving other HMG-CoA reductase inhibitors concomitantly with itraconazole and cyclosporine.

The benefits and risks of using HMG-CoA reductase inhibitors concomitantly with immunosuppressive drugs, fibrates, erythromycin, systemic azole derivative antifungal agents or lipid-lowering doses of niacin should be carefully considered.

Rare cases of rhabdomyolysis with acute renal failure secondary to myoglobinuria, have been reported with pravastatin sodium and with other HMG-CoA reductase inhibitors.

Myopathy, defined as muscle pain or muscle weakness in conjunction with increases in creatine phosphokinase (CK) values to greater than ten times the upper limit of normal, should be considered in any patient with diffuse myalgias, muscle tenderness or weakness, and/or marked

elevation of CK. Patients should be advised to report promptly any unexplained muscle pain, tenderness or weakness, particularly if associated with malaise or fever. Patients who develop any signs or symptoms suggestive of myopathy should have their CK levels measured. Pravastatin sodium therapy should be discontinued if markedly elevated CK levels are measured or myopathy is diagnosed or suspected.

As with other statins, the risk of myopathy including rhabdomyolysis may be substantially increased by concomitant immunosuppressive therapy including cyclosporines, and by concomitant therapy with gemfibrozil, erythromycin or niacin (see WARNINGS AND PRECAUTIONS).

Myopathy has not been observed in clinical trials involving small numbers of patients who were treated with pravastatin sodium together with immunosuppressants, fibric acid derivatives or niacin (see CLINICAL TRIALS).

The use of fibrates alone is occasionally associated with myopathy. In a limited size clinical trial of combined therapy with pravastatin (40 mg/day) and gemfibrozil (1200 mg/day), myopathy was not reported, although a trend towards CK elevations and musculoskeletal symptoms was seen. The combined use of pravastatin and fibrates should generally be avoided.

No information is available on the combined therapy of pravastatin with erythromycin.

Pre-disposing Factors for Myopathy/Rhabdomyolysis: pravastatin sodium, as with other HMG-CoA reductase inhibitors, should be prescribed with caution in patients with pre-disposing factors for myopathy/rhabdomyolysis. Such factors include:

- Personal or family history of hereditary muscular disorders
- Previous history of muscle toxicity with another HMG-CoA reductase inhibitor
- Concomitant use of a fibrate or niacin
- Uncontrolled hypothyroidism
- Alcohol abuse
- Excessive physical exercise
- Advance age >65 years
- Renal impairment
- Hepatic impairment
- Diabetes with hepatic fatty change
- Surgery and trauma
- Frailty
- Situation where an increase in plasma levels of active ingredient may occur.

Pravastatin sodium therapy should be temporarily withheld or discontinued in any patient with an acute serious condition suggestive of myopathy or predisposing to the development of rhabdomyolysis (e.g. sepsis, hypotension, major surgery, trauma, severe metabolic endocrine and electrolyte disorders, or uncontrolled seizures).

There have been very rare reports of an immune-mediated necrotizing myopathy (IMNM) during or after treatment with some statins including pravastatin. IMNM is clinically characterized by persistent proximal muscle weakness and elevated serum creatine kinase, which persist despite discontinuation of statin treatment.

### **Liver Dysfunction**

HMG-CoA reductase inhibitors have been associated with biochemical abnormalities of liver function. As with other lipid-lowering agents, including non-absorbable bile acid-binding resins, increases in liver enzymes to less than three times the upper limit of normal have occurred during therapy with pravastatin. The significance of these changes, which usually appear during the first few months of treatment initiation, is not known. In the majority of patients treated with pravastatin, in clinical trials, these increased values declined to pretreatment levels despite continuation of therapy at the same dose.

**Marked persistent increases (greater than three times the upper limit of normal) in serum transaminases were seen in 6 out of 1142 (0.5%) patients treated with pravastatin in clinical trials** (see ADVERSE REACTIONS). The increases usually appeared 3 to 12 months after the start of therapy with pravastatin sodium. These elevations were not associated with clinical signs and symptoms of liver disease and usually declined to pretreatment levels upon discontinuation of therapy. Patients rarely had persistent marked abnormalities possibly attributable to therapy. In the largest long-term placebo-controlled trial with pravastatin (Pravastatin Primary Prevention Study/WOSCOPS), no patient with normal liver function after 12 weeks of treatment (N = 2875 pravastatin-treated patients) had subsequent ALT elevations greater than three times the upper limit of normal on two consecutive measurements. Two of these 2875 patients treated with pravastatin (0.07%) and one of 2 919 placebo patients (0.03%) had elevations of AST greater than three times the upper limit of normal on two consecutive measurements during the 4.8 years (median treatment) of the study.

Liver function tests should be performed at baseline and at 12 weeks following initiation of therapy or the elevation of dose. Special attention should be given to patients who develop increased transaminase levels. Liver function tests should be repeated to confirm an elevation and subsequently monitored at more frequent intervals. **If increases in alanine aminotransferase (ALAT) and aspartate aminotransferase (ASAT) equal or exceed three times the upper limit of normal and persist, therapy should be discontinued.**

There have been rare post-marketing reports of fatal and non-fatal hepatic failure in patients taking statins, including pravastatin sodium. If serious liver injury with clinical symptoms and/or hyperbilirubinemia or jaundice occurs during treatment with pravastatin sodium, promptly interrupt therapy. If an alternate etiology is not found, do not restart pravastatin sodium.

Pravastatin sodium, as well as other HMG-CoA reductase inhibitors should be used with caution in patients who consume substantial quantities of alcohol and/or have a past history of liver disease. Active liver disease or unexplained serum transaminase elevations are contraindications to the use of pravastatin sodium; if such condition develops during therapy, the drug should be discontinued.

## **General**

Before instituting therapy with pravastatin sodium, an attempt should be made to control hypercholesterolemia with appropriate diet, exercise, weight reduction in overweight and obese patients, and to treat other underlying medical problems (see INDICATIONS AND CLINICAL USE). The patient should be advised to inform subsequent physicians of the prior use of pravastatin sodium.

Pravastatin may elevate creatine phosphokinase and transaminase levels. This should be considered in the differential diagnosis of chest pain in a patient on therapy with pravastatin.

## **Effect on the Lens**

Current data from clinical trials do not indicate an adverse effect of pravastatin on the human lens.

## **Interstitial Lung Disease**

Exceptional cases of interstitial lung disease have been reported with some statins, including pravastatin, especially with long term therapy (see ADVERSE REACTIONS, Post-Market Adverse Drug Reactions). Presenting features can include dyspnoea, non-productive cough and deterioration in general health (fatigue, weight loss, and fever). If it is suspected a patient has developed interstitial lung disease, statin therapy should be discontinued.

## **Homozygous Familial Hypercholesterolemia**

Pravastatin has not been evaluated in patients with rare homozygous familial hypercholesterolemia. Most HMG-CoA reductase inhibitors are less or not effective in this subgroup of hypercholesterolemic patients (see REFERENCES).

## **Effect on Lipoprotein (a)**

In some patients, the beneficial effect of lowered total cholesterol and LDL-C levels may be partly blunted by a concomitant increase in the Lipoprotein (a)[Lp(a)] level. Further research is ongoing to elucidate the significance of Lp(a) variations. Therefore, until further experience is obtained, where feasible, it is suggested that measurements of serum Lp(a) be followed up in patients placed on pravastatin therapy (see REFERENCES).

## **Effect on CoQ10 Levels (Ubiquinone)**

A significant short-term decrease in plasma CoQ10 levels in patients treated with pravastatin sodium has been observed. Longer clinical trials have also shown reduced serum ubiquinone levels during treatment with pravastatin and other HMG CoA reductase inhibitors. The clinical significance of a potential long-term statin-induced deficiency of CoQ10 has not yet been established. It has been reported that a decrease in myocardial ubiquinone levels could lead to impaired cardiac function in patients with borderline congestive heart failure (see REFERENCES).

## **Carcinogenesis**

A 21-month oral study in mice, with doses of 10 to 100 mg/kg daily of pravastatin did not demonstrate any carcinogenic potential. In a 2-year oral study in rats, a statistically significant increase in the incidence of hepatocellular carcinoma was observed in male rats given 100 mg/kg daily (60 times the maximum human dose) of pravastatin. This change was not seen in male rats given 40 mg/kg daily (25 times the recommended human dose) or less, or in female rats at any dose level.



### **Hypersensitivity**

With lovastatin an apparent hypersensitivity syndrome has been reported rarely which has included one or more of the following features: anaphylaxis, angioedema, lupus-like syndrome, polymyalgia rheumatica, thrombocytopenia, leukopenia, hemolytic anemia, positive antinuclear antibody (ANA), erythrocytes sedimentation rate (ESR) increase, arthritis, arthralgia, urticaria, asthenia, photosensitivity, fever and malaise.

Although to date hypersensitivity syndrome has not been described as such, in few instances eosinophilia and skin eruptions appear to be associated with pravastatin sodium treatment. If hypersensitivity is suspected pravastatin sodium should be discontinued. Patients should be advised to report promptly any signs of hypersensitivity such as angioedema, urticaria, photosensitivity, polyarthralgia, fever, malaise.

### **Endocrine Function**

HMG-CoA reductase inhibitors interfere with cholesterol synthesis and as such could theoretically blunt adrenal and/or gonadal steroid production.

In one long-term study investigating the endocrine function in hypercholesterolemic patients, pravastatin sodium exhibited no effect upon basal and stimulated cortisol levels, as well as on aldosterone secretion. Although no change was reported in the testicular function, conflicting results were observed in the analysis of sperm motility after administration of pravastatin sodium. A case of reversible impotence has been reported in a 57-year old man administered pravastatin 20 mg/day and metoprolol (see REFERENCES). A causal relationship to therapy with pravastatin sodium has not been established. Further studies are needed to clarify the effects of HMG CoA reductase inhibitors on male fertility. Furthermore, the effects, if any, on the pituitary-gonadal axis in premenopausal women are unknown.

Patients treated with pravastatin sodium who develop clinical evidence of endocrine dysfunction should be evaluated appropriately. Caution should be exercised if an HMG-CoA reductase inhibitor or other agent used to lower cholesterol levels is administered to patients receiving other drugs (e.g. ketoconazole, spironolactone, or cimetidine) that may decrease the levels of endogenous steroid hormones.

Increases in fasting glucose and HbA1c levels have been reported with inhibitors of HMG-CoA reductase as a class. For some patients at high risk of diabetes mellitus, hyperglycemia was sufficient to shift them to diabetes status. The benefit of treatment continues to outweigh the small increased risk. Periodic monitoring of these patients is recommended.

### **Patients with Severe Hypercholesterolemia**

Higher doses ( $\geq 40$  mg/day) required for some patients with severe hypercholesterolemia are associated with increased plasma levels of pravastatin. **Caution should be exercised in such patients who are also significantly renally impaired or elderly** (see WARNINGS AND PRECAUTIONS - Muscle Effects)

### **Special Populations**

**Use in Pregnancy: pendo-PRAVSTATIN is contraindicated during pregnancy (see CONTRAINDICATIONS).**

Safety in pregnant women has not been established. Although pravastatin was not teratogenic in rats at doses as high as 1000 mg/kg daily nor in rabbits at doses of up to 50 mg/kg daily, pendo-PRAVASTATIN should be administered to women of childbearing age only when such patients are highly unlikely to conceive and have been informed of potential hazards. If a woman becomes pregnant while taking pravastatin sodium, pravastatin sodium should be discontinued and the patient advised again as to the potential hazards to the fetus.

**Use in Nursing Mothers:** A negligible amount of pravastatin is excreted in human breast milk. Because of the potential for adverse reactions in nursing infants, if the mother is being treated with pravastatin sodium, nursing should be discontinued or treatment with pravastatin sodium stopped.

**Pediatric Use (< 16 years of age):** Only limited experience with the use of statins in children is available (see REFERENCES). There is no experience to date with the use of pravastatin sodium in such patients. Treatment in these patients is not recommended at this time.

**Elderly ( $\geq$  65 years of age):** Pharmacokinetic evaluation of pravastatin in patients over the age of 65 years indicates an increased AUC. There were no reported increases in the incidence of adverse effects in these or other studies involving patients in that age group. As a precautionary measure, the lowest dose should be administered initially (see REFERENCES).

Elderly patients may be more susceptible to myopathy (see WARNINGS AND PRECAUTIONS - Muscle Effects - Pre-disposing Factors for Myopathy/Rhabdomyolysis).

**Use in Patients with Impaired Renal Function:** There have been no studies on the use of pravastatin in patients with renal insufficiency. As a precautionary measure, the lowest dose should be used in these patients (see WARNINGS AND PRECAUTIONS - Muscle Effects).

## ADVERSE REACTIONS

### Adverse Drug Reaction Overview

Pravastatin is generally well tolerated. Adverse events have been usually mild to moderate and transient. Adverse events observed or reported in short- and long-term trials are as follows.

### **Clinical Trial Adverse Drug Reactions**

*Because clinical trials are conducted under very specific conditions the adverse reaction rates observed in the clinical trials may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse drug reaction information from clinical trials is useful for identifying drug-related adverse events and for approximating rates.*

### Short-term Controlled Trials

All adverse clinical events (regardless of attribution) reported in more than 2% of pravastatin treated patients in placebo-controlled trials of up to four months duration are identified in the

following table; also shown are the percentages of patients in whom these medical events were believed to be related or possibly related to the drug.

**Adverse Events in > 2 Percent of Patients Treated with  
Pravastatin 10-40 mg in Short-Term Placebo-Controlled Trials**

Body System/ Event	All Events		Events Attributed to Study Drug	
	Pravastatin (N = 900) % of patients	Placebo (N = 411) % of patients	Pravastatin (N = 900) % of patients	Placebo (N = 411) % of patients
<b>Cardiovascular</b>				
Cardiac Chest Pain	4.0	3.4	0.1	0.0
<b>Dermatologic</b>				
Rash	4.0*	1.1	1.3	0.9
<b>Gastrointestinal</b>				
Nausea/Vomiting	7.3	7.1	2.9	3.4
Diarrhea	6.2	5.6	2.0	1.9
Abdominal Pain	5.4	6.9	2.0	3.9
Constipation	4.0	7.1	2.4	5.1
Flatulence	3.3	3.6	2.7	3.4
Heartburn	2.9	1.9	2.0	0.7
<b>General</b>				
Fatigue	3.8	3.4	1.9	1.0
Chest Pain	3.7	1.9	0.3	0.2
Influenza	2.4*	0.7	0.0	0.0
<b>Musculoskeletal</b>				
Localized Pain	10.0	9.0	1.4	1.5
Myalgia	2.7	1.0	0.6	0.0
<b>Nervous System</b>				
Headache	6.2	3.9	1.7*	0.2
Dizziness	3.3	3.2	1.0	0.5
<b>Renal/Genitourinary</b>				
Urinary Abnormality	2.4	2.9	0.7	1.2
<b>Respiratory</b>				
Common Cold	7.0	6.3	0.0	0.0
Rhinitis	4.0	4.1	0.1	0.0
Cough	2.6	1.7	0.1	0.0

\* Statistically significantly different from placebo

The safety and tolerability of pravastatin sodium at a dose of 80 mg in two controlled trials with a mean exposure of 8.6 months was similar to that of pravastatin sodium at lower doses except that 4 out of 464 patients taking 80 mg of pravastatin had a single elevation of CK > 10X ULN compared to 0 out of 115 patients taking 40 mg of pravastatin.

**Long-term Controlled Morbidity and Mortality Trials**

In seven randomized double blind placebo-controlled trials involving over 21,500 patients treated with pravastatin (N= 10,784) or placebo (N= 10,719), the safety and tolerability in the pravastatin group was comparable to that of the placebo group. Over 19,000 patients were followed for a median of 4.8 - 5.9 years, while the remaining patients were followed for two years or more.

Clinical adverse events probably or possibly related, or of uncertain relationship to therapy, occurring in at least 0.5% of patients treated with pravastatin or placebo in these long-term morbidity/mortality trials are shown in the table below:

	<b>Pravastatin sodium (N = 10,784) %</b>	<b>Placebo (N = 10,719) %</b>
<b>Cardiovascular</b>		
Angina pectoris	3.1	3.4
Disturbance rhythm subjective	0.8	0.7
Hypertension	0.7	0.9
Edema	0.6	0.6
Myocardial infarction	0.5	0.7
<b>Gastrointestinal</b>		
Dyspepsia / heartburn	3.5	3.7
Nausea/ vomiting	1.4	1.6
Flatulence	1.2	1.1
Constipation	1.2	1.3
Diarrhea	0.9	1.1
Abdominal pain	0.9	1.0
Distention abdomen	0.5	0.5
<b>Musculoskeletal</b>		
Musculoskeletal Pain (includes arthralgia)	5.9	5.7
Muscle cramp	2.0	1.8
Myalgia	1.4	1.4
Musculoskeletal trauma	0.5	0.3
<b>Nervous System</b>		
Dizziness	2.2	2.1
Headache	1.9	1.8
Sleep disturbance	1.0	0.9
Depression	1.0	1.0
Anxiety / nervousness	1.0	1.2
Paresthesia	0.9	0.9
Numbness	0.5	0.4
<b>General</b>		
Fatigue	3.4	3.3
Chest pain	2.6	2.6
Weight gain	0.6	0.7
Influenza	0.6	0.5
<b>Special senses</b>		
Vision disturbance (includes blurred vision)	1.5	1.3
Disturbance eye (includes eye inflammation)	0.8	0.9
Hearing abnormality (includes tinnitus and hearing loss)	0.6	0.5
Lens opacity	0.5	0.4
<b>Dermatologic</b>		
Rash	2.1	2.2
Pruritis	0.9	1.0
<b>Renal / Genitourinary</b>		
Urinary abnormality (includes dysuria and nocturia)	1	0.8

	<b>Pravastatin sodium (N = 10,784) %</b>	<b>Placebo (N = 10,719) %</b>
<b>Respiratory</b>		
Dyspnea	1.6	1.6
Upper respiratory infection	1.3	1.3
Cough	1.0	1.0
Sinus abnormality (includes sinusitis)	0.8	0.8
Pharyngitis	0.5	0.6

### **Abnormal Hematologic and Clinical Chemistry Findings**

Increases in serum transaminases and in creatine phosphokinase (CK) in patients treated with pravastatin sodium have been discussed (see WARNINGS AND PRECAUTIONS).

### **Post-Market Adverse Drug Reactions**

The following adverse events have also been rarely reported during post-marketing experience with pravastatin sodium, regardless of causality assessment:

- Cardiovascular:* angioedema
- Dermatologic:* a variety of skin changes (pruritis, scalp hair abnormalities, skin dryness and dermatitis)
- Endocrine:* increase in fasting glucose and HbA1C
- Gastrointestinal:* pancreatitis, hepatitis and fulminant hepatic necrosis, fatal and non-fatal hepatic failure, jaundice (including cholestatic), fatty change in liver, cirrhosis, thrombocytopenia, hepatoma, abnormal stool and appetite change. Liver Function Test (LFT) abnormalities have also been reported.
- General:* chest pain (non cardiovascular), weakness, excess sweating hot flashes and fever
- Hypersensitivity:* anaphylaxis, lupus erythematosus-like syndrome, polymyalgia, rheumatica, dermatomyositis, vasculitis, purpura, hemolytic anemia, positive ANA, ESR increase, arthritis, arthralgia, asthenia, photosensitivity reaction, chills, malaise, toxic epidermal necrolysis, erythema multiforme, including Stevens-Johnson syndrome
- Immunologic:* allergy
- Musculoskeletal:* myopathy, rhabdomyolysis, tendon disorder (specifically tendonitis and tendon rupture), polymyositis and immune mediated necrotizing myopathy
- Nervous System:* dysfunction of certain cranial nerves (including alteration of taste, impairment of extra-ocular movement, facial paresis), peripheral

nerve palsy, paresthesia equilibrium disturbance, vertigo, memory impairment, tremor, mood change, mood related disorders including depression, sleep disturbances including insomnia and nightmares

*Pulmonary:* Very rare cases of interstitial lung disease, especially with long term therapy. If it is suspected a patient has developed interstitial lung disease, statin therapy should be discontinued.

*Reproductive:* gynecomastia, impotence (see Endocrine Function), urticaria, sexual dysfunction, libido change.

*Special Senses:* eye symptoms (including soreness, dryness or itching), tinnitus, taste disturbance.

There have been rare post-marketing reports of cognitive impairment (e.g., memory loss, forgetfulness, amnesia, memory impairment, confusion) associated with statin use. These cognitive issues have been reported for all statins. The reports are generally non-serious and reversible upon statin discontinuation, with variable times to symptom onset (1 day to years) and symptom resolution (median of 3 weeks).

The following have also been reported with other statins: hepatitis, cholestatic jaundice, anorexia, psychic disturbances including anxiety, hypospermia, hypersensitivity, and increase in fasting glucose and HbA1C (see WARNINGS AND PRECAUTIONS).

#### **Lens**

Current data from clinical trials do not indicate an adverse effect of pravastatin on the human lens.

## **DRUG INTERACTIONS**

### **Drug-Drug Interactions**

#### **Concomitant Therapy with Other Lipid Metabolism Regulators**

Based on post-marketing surveillance, gemfibrozil, fenofibrate, other fibrates and lipid lowering doses of niacin (nicotinic acid) may increase the risk of myopathy when given concomitantly with HMG-CoA reductase inhibitors, probably because they can produce myopathy when given alone (see WARNINGS AND PRECAUTIONS - Muscle Effects). Therefore, combined drug therapy should be approached with caution.

#### **Bile Acid Sequestrants**

Preliminary evidence suggests that the cholesterol-lowering effects of pravastatin sodium and the bile acid sequestrants, cholestyramine/colestipol are additive.

When pravastatin was administered one hour before or four hours after cholestyramine or one hour before colestipol and a standard meal, there was no clinically significant decrease in bioavailability or therapeutic effect. Concomitant administration resulted in an approximately 40 to 50% decrease in the mean AUC of pravastatin (see DOSAGE AND ADMINISTRATION - Concomitant Therapy).

### **Gemfibrozil and nicotinic acid**

Gemfibrozil and nicotinic acid do not statistically significantly affect the bioavailability of pravastatin. However, in a limited size clinical trial, a trend toward CK elevations and musculoskeletal symptoms was seen in patients treated concurrently with pravastatin and gemfibrozil.

Myopathy, including rhabdomyolysis, has occurred in patients who were receiving coadministration of HMG-CoA reductase inhibitors with fibric acid derivatives and niacin, particularly in subjects with pre-existing renal insufficiency (see WARNINGS AND PRECAUTIONS - Muscle Effects).

### **Other Concomitant Therapy**

The use of HMG CoA reductase inhibitors has been associated with severe myopathy, including rhabdomyolysis, which may be more frequent when they are administered with drugs that inhibit the cytochrome P-450 enzyme system. *In vitro* and *in vivo* data indicate that pravastatin is not metabolized by cytochrome P450 3A4 to a clinically significant extent. This has been shown in studies with known cytochrome P450 3A4 inhibitors.

### **Digoxin**

Coadministration of digoxin and other HMG CoA reductase inhibitors has been shown to increase the steady state digoxin concentrations. The potential effects of coadministration of digoxin and pravastatin sodium are not known. As a precautionary measure, patients taking digoxin should be closely monitored.

### **Antipyrine**

Antipyrine was used as a model for drugs metabolized by the microsomal hepatic enzyme system (cytochrome P450 system). Pravastatin had no effect on the pharmacokinetics of antipyrine.

### **Coumarin Anticoagulants**

Pravastatin had no clinically significant effect on prothrombin time when administered in a study to normal elderly subjects who were stabilized on warfarin.

### **Antacids and Cimetidine**

On the average, antacids (one hour prior to pravastatin sodium) reduce and cimetidine increases the bioavailability of pravastatin. These changes were not statistically significant. The clinical significance of these interactions is not known but is probably minimal as judged from the interaction with food (see ACTION AND CLINICAL PHARMACOLOGY – Human Pharmacology).

No information is available regarding interactions with erythromycin (see WARNINGS AND PRECAUTIONS - Muscle Effects).

Although specific interaction studies were not performed during clinical trials, no noticeable drug interactions were reported when pravastatin sodium was added to diuretics, antihypertensives, angiotensin converting-enzyme (ACE) inhibitors, calcium channel blockers, or nitroglycerin.

### **Propranolol**

Co-administration of propranolol and pravastatin reduced the AUC values by 23% and 16% respectively.

### **Cyclosporine**

Several studies show that cyclosporine appears to increase pravastatin plasma concentration by several folds. In a multicentre study, the AUC values of pravastatin were shown to be five-fold higher in the presence of cyclosporine. There was no accumulation of pravastatin after multiple doses (see DOSAGE AND ADMINISTRATION and REFERENCES).

### **Macrolides**

Macrolides have the potential to increase statin exposure while used in combination. Pravastatin should be used cautiously with macrolide antibiotics due to potential increased risk of myopathies.

### **Drug-Laboratory Interactions**

Pravastatin may elevate creatine phosphokinase and transaminase levels. This should be considered in the differential diagnosis of chest pain in a patient on therapy with pravastatin.

## **DOSAGE AND ADMINISTRATION**

### **Dosing Considerations**

Patients should be placed on a standard cholesterol-lowering diet (at least equivalent to the Adult Treatment Panel III [ATP III TLC diet]) before receiving pendo-PRAVASTATIN (pravastatin sodium), and should continue on this diet during treatment with pendo-PRAVASTATIN. If appropriate, a program of weight control and physical exercise should be implemented.

Prior to initiating therapy with pendo-PRAVASTATIN, secondary causes for elevations in plasma lipid levels should be excluded. A lipid profile should also be performed.

### **Recommended Dose and Dosage Adjustment**

#### **Hypercholesterolemia and Coronary Heart Disease**

The recommended starting dose is 20 mg once daily at bed time. Patients who require a large dose reduction in LDL-C may be started at 40 mg once daily. The dose of 80 mg once daily should be reserved for patients who do not achieve their treatment goal with lower doses. pendo-PRAVASTATIN may be taken without regard to meals (see ACTION AND CLINICAL PHARMACOLOGY).

In patients with a history of significant renal or hepatic dysfunction, a starting dose of 10 mg daily is recommended.

#### **Concomitant Therapy**

Some patients may require combination therapy with one or more lipid-lowering agents. Pharmacokinetic interaction with pravastatin administered concurrently with nicotinic acid, or gemfibrozil did not statistically significantly affect the bioavailability of pravastatin. The combined use of pravastatin and fibrates should however generally be avoided (see WARNINGS AND PRECAUTIONS - Muscle Effects).



The lipid-lowering effects of pravastatin sodium on Total and Low Density Lipoprotein Cholesterol are additive when combined with a bile acid-binding resin. However, when administering a bile acid-binding resin (e.g. cholestyramine, colestipol) and pravastatin, pravastatin should not be administered concomitantly, but should be given either one hour or more before or at least four hours following the resin (see DRUG INTERACTIONS - Concomittant Therapy with Other Lipid Metabolism Regulators).

In patients taking cyclosporine, with or without other immunosuppressive drugs, concomitantly with pravastatin, therapy should be initiated with 10 mg per day and titration to higher doses should be performed with caution. Most patients treated with this combination received a maximum pravastatin dose of 20 mg/day (see WARNINGS AND PRECAUTIONS and DRUG INTERACTIONS - Other Concomitant Therapy, Cyclosporine).

The dosage of pendo-PRAVASTATIN should be individualized according to baseline LDL-C, total- C/HDL-C ratio and/or TG levels to achieve the desired lipid values at the lowest possible dose.

## OVERDOSAGE

There have been two reports of overdose with pravastatin, both of which were asymptomatic and not associated with clinical laboratory abnormalities.

In the event of overdose, treatment should be symptomatic and supportive, and appropriate therapy instituted. Until further experience is obtained, no specific therapy of overdose can be recommended. The dialyzability of pravastatin and its metabolites is not known.

For management of a suspected drug overdose, contact your regional Poison Control Centre immediately.

## ACTION AND CLINICAL PHARMACOLOGY

### **Mechanism of Action**

Pravastatin sodium is one of a new class of lipid-lowering compounds known as HMG-CoA reductase inhibitors (statins) that reduce cholesterol biosynthesis. These agents are competitive inhibitors of 3-hydroxy- 3-methylglutaryl-coenzyme A (HMG-CoA) reductase, the enzyme catalyzing the early rate-limiting step in cholesterol biosynthesis, conversion of HMG-CoA to mevalonate. Pravastatin is isolated from a strain of *Penicillium citrinum*. The active drug substance is the hydroxyacid form.

Pravastatin sodium produces its lipid-lowering effect in two ways. First, as a consequence of its reversible inhibition of HMG-CoA reductase activity, it effects modest reductions in intracellular pools of cholesterol. This results in an increase in the number of Low Density Lipoproteins (LDL) - receptors on cell surfaces and enhanced receptor-mediated catabolism and clearance of

circulating LDL. Second, pravastatin inhibits LDL production by inhibiting hepatic synthesis of Very Low Density Lipoproteins (VLDL), the LDL precursor.

Epidemiologic and clinical investigations have associated the risk of coronary artery disease (CAD) with elevated levels of Total-C, LDL-C and decreased levels of HDL-C. These abnormalities of lipoprotein metabolism are considered as major contributors to the development of the disease. Other factors, e.g. interactions between lipids/ lipoproteins and endothelium, platelets and macrophages, have also been incriminated in the development of human atherosclerosis and of its complications.

In long-term, prospective clinical trials effective treatment of hypercholesterolemia/ dyslipidemia has consistently been associated with a reduction in the risk of CAD.

Treatment with pravastatin sodium has been shown to reduce circulating Total-C, LDL-C, and apolipoprotein B, modestly reduce VLDL-C and triglycerides (TG) while producing increases of variable magnitude in HDL-C and apolipoprotein A. Clinical trials suggest that pravastatin sodium's effect on reducing clinical events appears to incorporate both cholesterol modification and some ancillary mechanism.

Pravastatin has complex pharmacokinetic characteristics.

### **Human Pharmacology**

In both normal volunteers and patients with hypercholesterolemia, treatment with pravastatin sodium reduced total-C, LDL-C, apolipoprotein B, VLDL-C and TG while increasing HDL-C and apolipoprotein A. The mechanism of action of pravastatin sodium is complex. Inhibition of hepatic VLDL synthesis and/or secretion occurs, leading to a decrease in LDL precursor formation. The reduction in hepatic cellular pools of cholesterol, resulting from the specific and reversible inhibition of HMG-CoA reductase activity, leads to an increase in the fractional catabolic rate of IDL and LDL via increased expression of LDL receptors on the surface of hepatic cells. Through a combination of these and possibly other unknown metabolic effects, a decline in the serum level of cholesterol results.

### **Pharmacokinetics**

**Absorption:** Pravastatin sodium is administered orally in the active form. Following oral ingestion, pravastatin is rapidly absorbed with peak plasma levels attained at about 1 to 1.5 hours. Average oral absorption of pravastatin, based on urinary recovery of radiolabelled drug after oral and intravenous dosing, is 34%; average absolute bioavailability of the parent drug is 17%. The therapeutic response to pravastatin sodium is similar, whether taken with meals or one hour prior to meals, even though the presence of food in the gastrointestinal tract causes a reduction in systemic bioavailability.

**Percent Decrease in LDL-C**

<b>Pravastatin</b>	<b>10 mg bid</b>	<b>20 mg bid</b>
With meals	- 25%	- 37%
Before meals *	- 26%	- 36%

\* administered one hour or more prior to eating.

**Distribution:** Pravastatin undergoes extensive first pass extraction in the liver (estimated hepatic extraction ratio, 66%), its primary site of action, and is excreted in the bile. Therefore, plasma levels of the drug are probably of limited value in predicting therapeutic effectiveness. Nevertheless, measurement of plasma pravastatin concentrations by gas chromatography and mass-spectrometry showed dose proportionality for area under the concentration-time curve (AUC) and maximum and steady-state plasma levels. Steady-state areas under the plasma concentration-time-curves and maximum ( $C_{MAX}$ ) or minimum ( $C_{MIN}$ ) plasma concentrations showed no accumulation following once or twice-daily administration of pravastatin sodium tablets.

**Metabolism:** Pravastatin is extensively metabolized. The major metabolite is the 3  $\alpha$ -hydroxy isomer, which has one-tenth to one-fortieth of the inhibitory activity of the parent compound on HMG-CoA reductase.

**Excretion:** Protein binding of pravastatin is approximately 50%. The plasma elimination half-life of pravastatin is between 1.5 and 2 hours (2.5 - 3 hours in hypercholesterolemic subjects). Approximately 20% of a radiolabelled oral dose is excreted in the urine and 70% in the feces.

After intravenous administration to healthy subjects, approximately 47% of the total drug clearance occurs via renal excretion of intact pravastatin, and about 53% is cleared by non-renal routes, i.e. biliary excretion and biotransformation.

### **Special Populations and Conditions**

**Pediatrics:** Only limited experience with the use of statins in children is available (see REFERENCES). There is no experience to date with the use of pravastatin sodium in such patients. Treatment in these patients is not recommended at this time.

**Geriatrics:** Studies of pravastatin sodium administered as a single dose to healthy elderly male and female subjects (age 65 to 78 years) indicated a 30 - 50% increase in plasma levels.

**Renal Insufficiency:** No studies have been carried out in patients with renal insufficiency.

### **STORAGE AND STABILITY**

Store between 15°C and 30°C. Protect from moisture and light.

### **DOSAGE FORMS, COMPOSITION AND PACKAGING**

#### **Dosage Forms**

**10 mg:** Each pink to peach rounded, rectangular-shaped biconvex tablet, is debossed with “P” over “10” on one side and nothing on the other side. Each tablet contains 10 mg of pravastatin sodium and the following non-medicinal ingredients: Colloidal Silicon Dioxide, Copovidone, Croscarmellose Sodium, Dibasic Calcium Phosphate, Iron Oxide

Red, Lactose, Magnesium Stearate, Microcrystalline Cellulose, Polyethylene Glycol.  
Available in bottles of 100 and in blister packages of 30.

**20 mg:** Each yellow, rounded, rectangular-shaped biconvex tablet is debossed with “P” over “20” on one side and nothing on the other side. Each tablet contains 20 mg of pravastatin sodium and the following non-medicinal ingredients: Colloidal Silicon Dioxide, Copovidone, Croscarmellose Sodium, Dibasic Calcium Phosphate, Iron Oxide Yellow, Lactose, Magnesium Stearate, Microcrystalline Cellulose, Polyethylene Glycol. Available in bottles of 100 and 500 and in blister packages of 30.

## PART II: SCIENTIFIC INFORMATION

### PHARMACEUTICAL INFORMATION

#### Drug Substance


Proper Name: Pravastatin Sodium

Chemical Name: Pravastatin sodium is designated chemically as [1S-[1 $\alpha$  ( $\beta$ S\*, $\delta$ S\*)2 $\alpha$ ,6 $\alpha$ ,8 $\beta$ (R\*),8 $\alpha\alpha$ ]]-1,2,6,7,8,8a-hexahydro- $\beta$ , $\delta$ ,6-trihydroxy-2-methyl-8-(2-methyl-1-oxobutoxy)-1-naphthaleneheptanoic acid, monosodium salt.

Molecular Formula: C<sub>23</sub>H<sub>35</sub>NaO<sub>7</sub>

Molecular Weight: 446.52 g/mol

Structural Formula:



Physiochemical properties: Pravastatin is a white to off white, hygroscopic crystalline powder that is soluble in water, in methanol, ethanol, slightly soluble in isopropyl alcohol and practically insoluble in acetone, acetonitrile, chloroform, ethyl-acetate and ether.

## CLINICAL TRIALS

### Comparative Bioavailability Studies

A single dose cross-over comparative bioavailability study to evaluate the pharmacokinetic profile and estimate the bioequivalence of 40 mg tablets of Pravastatin Sodium compared to the Reference formulation, i.e. PRAVACHOL<sup>®</sup> 40 mg tablets (Squibb Canada Division Bristol-Myers Squibb Canada Inc.) was performed in 18 healthy male volunteers under fasting conditions. The results are summarized below.

#### SUMMARY TABLE OF THE COMPARATIVE BIOAVAILABILITY DATA

<b>Pravastatin ( 1 x 40 mg) From measured data Geometric Mean Arithmetic Mean (CV %)</b>				
<b>Parameter</b>	<b>Pravastatin Sodium Tablets</b>	<b>PRAVACHOL**</b>	<b>% Ratio of Geometric Means</b>	<b>90% Confidence Interval</b>
AUC <sub>1</sub> (ng.h/mL)	128.16 141.27 (51.12)	143.28 156.57 (47.14)	91.61	81.56 to 102.90
AUC <sub>1</sub> (ng.h/mL)	130.14 143.52 (51.10)	144.84 158.41 (46.53)	91.69	81.86 to 102.69
C <sub>MAX</sub> (ng/mL)	58.97 61.56 (49.10)	61.70 74.86 (63.38)	90.39	77.95 to 104.82
T <sub>MAX</sub> * (h)	1.06 (33.08)	1.07 (47.98)		
T <sub>1/2</sub> * (h)	2.97 (60.91)	3.09 (64.15)		

\* T<sub>max</sub> and the T<sub>1/2</sub> parameter are expressed as the arithmetic mean (CV%)

\*\* PRAVACHOL<sup>®</sup> tablets are manufactured in Canada by Squibb Canada Inc.

## **Study Results**

### **Hypercholesterolemia**

Pravastatin sodium is highly effective in reducing total and LDL cholesterol in patients with primary hypercholesterolemia. A marked response is seen within one week, and the maximum therapeutic response usually occurs within four weeks. The response is maintained during extended periods of therapy. In addition, pravastatin sodium is effective in reducing the progressive cause of atherosclerosis and risk of coronary events, decreasing total mortality, decreasing death due to coronary heart disease, and decreasing the incidence of stroke, in hypercholesterolemic patients with atherosclerotic cardiovascular disease. Pravastatin sodium is also effective in reducing the risk of CHD death (fatal MI and sudden death) plus non-fatal MI with no increase in deaths from non-cardiovascular causes in hypercholesterolemic patients without previous myocardial infarction. Risk reduction is evident within 6 months of the initiation of treatment (see Figure 1).

Single daily doses of pravastatin sodium are effective. As shown in the table which follows, the Total-C and LDL-C lowering effects are the same whether pravastatin sodium is administered in single or divided (bid) doses. Once-daily administration in the evening appears to be marginally more effective than once-daily administration in the morning, perhaps because hepatic cholesterol is synthesized mainly at night.

The results of a multicenter, double-blind regimen response comparative study of placebo and pravastatin, given to parallel groups of patients for 8 weeks are as follows:

**Single-Daily Versus Twice-Daily Dosing\***

<b>Pravastatin</b>	<b>N</b>	<b>Total-C</b>	<b>LDL-C</b>	<b>HDL-C</b>	<b>TG</b>
40 mg qam	41	- 23%	- 30%	+ 4%	- 11%
40 mg qpm	33	- 26%	- 33%	+ 8%	- 24%
20 mg bid	44	- 27%	- 34%	+ 8%	- 25%

\* Evening doses were administered at least 3 hours after the evening meal. Morning doses were administered at least 1 hour prior to breakfast.

Patients with primary hypercholesterolemia (71% Familial or Familial Combined, 29% Non-Familial). Baseline mean LDL-C = 6.34 mmol/L (245.4 mg/dL).

In multicenter, double-blind studies of patients with primary hypercholesterolemia, pravastatin sodium administered in daily doses ranging from 5 mg to 80 mg to over 1100 patients was compared with placebo. Pravastatin sodium significantly decreased Total-C and LDL-C levels, and Total-C/HDL-C and LDL-C/HDL-C ratios. In addition, pravastatin slightly increased HDL-C and decreased VLDL-C and plasma TG levels.

Dose-response effects on lipids from two studies evaluated after 8 weeks of administering pravastatin sodium once or twice-daily are illustrated in the tables below.

**Dose-Response Results\***  
**(Once-Daily Administration at Bedtime)**

<b>Pravastatin</b>	<b>N</b>	<b>Total-C</b>	<b>LDL-C</b>	<b>HDL-C</b>	<b>TG</b>
5 mg qd	16	-14%	- 19%	+ 5%	- 14%
10 mg qd	18	- 16%	- 22%	+ 7%	- 15%
20 mg qd	19	- 24%	- 32%	+ 2%	- 11%
40 mg qd	18	- 25%	- 34%	+ 12%	- 24%

\* Patients with primary hypercholesterolemia (28% Familial or Familial Combined, 72% Non-Familial). Baseline mean LDL-C = 5.68 mmol/L (219.6 mg/dL)

**Dose-Response Results \***  
**(BID Administration)**

<b>Pravastatin</b>	<b>N</b>	<b>Total-C</b>	<b>LDL-C</b>	<b>HDL-C</b>	<b>TG</b>
5 mg bid	59	-15%	- 20%	+ 7%	- 14%
10 mg bid	53	- 18%	- 24%	+ 6%	- 17%
20 mg bid	56	- 24%	- 31%	+ 5%	- 17%

\* Patients with primary hypercholesterolemia (70% Familial or Familial Combined, 30% Non-Familial). Baseline mean LDL-C = 6.06 mmol/L (234.5 mg/dL)

Pravastatin sodium is also effective when given with a bile acid-binding resin. In a study of pravastatin sodium administered alone or in combination with cholestyramine, marked reductions in the level of LDL-C were observed. In addition, pravastatin sodium attenuated the increase in TG levels observed with cholestyramine alone (The results of the study cited in the table which follows should be interpreted in the context of the exceptionally high rate of patient compliance with the bile acid-binding resin [70% of patients were taking 20 or 24 g daily]).

**Comparison With Cholestyramine Resin\***

	<b>N</b>	<b>Total-C</b>	<b>LDL-C</b>	<b>HDL-C</b>	<b>TG</b>
<b>Pravastatin</b>					
20 mg bid	49	- 24%	- 32%	+ 6%	- 10%
40 mg bid	52	- 30%	- 39%	+ 5%	- 15%
Resin Alone**	41	- 22%	- 31%	+ 2%	+ 16%
<b>Combination</b>					
20 mg bid & Resin**	49	- 38%	- 52%	+ 5%	- 1%

\* Patients with primary hypercholesterolemia (68% Familial or Familial Combined; 32% Non-Familial). Baseline mean LDL-C = 6.09 mmol/L (235.3 mg/dL)

\*\* The dose of resin used in this study was 24 g.

**Primary Prevention of Coronary Events**

Pravastatin sodium has been shown to be effective in reducing the risk of coronary heart disease (CHD) death plus non-fatal MI in hypercholesterolemic patients without previous myocardial infarction.

In the West of Scotland Study (WOS), the effect of pravastatin sodium on fatal and non-fatal coronary heart disease (CHD) was assessed in 6595 patients. The patient population consisted of men 45-64 years of age, without a previous MI, and with LDL-C levels between 4 - 6.7 mmol/L (156 - 254 mg/dL). In this randomized, double-blind, placebo-controlled study, patients were



treated with standard care, including dietary advice, and either pravastatin sodium 40 mg daily (n = 3302) or placebo (n = 3293) for a median duration of 4.8 years.

Pravastatin sodium significantly reduced the risk of CHD death plus non-fatal MI by 31% (248 patients in the placebo group [CHD death = 44, non-fatal MI = 204] vs 174 patients in the pravastatin sodium group [CHD death = 31, non-fatal MI = 143],  $p = 0.0001$ ). As shown in the figure below, divergence in the cumulative event rate curves for this endpoint begins within 6 months of treatment. This reduction was similar and significant throughout the entire range of baseline LDL cholesterol levels with a 37% risk reduction for LDL cholesterol 4 - 4.8 mmol/L (156 - 188 mg/dL) ( $p = 0.003$ ) and a 27% risk reduction for LDL cholesterol 4.9 - 6.7 mmol/L (189 - 254 mg/dL) ( $p = 0.03$ ). This reduction was also similar and significant for all age groups studied with a 40% risk reduction for patients younger than 55 years ( $p = 0.002$ ) and 27% risk reduction for patients 55 years and older ( $p = 0.009$ ).




Figure 1.

Total cardiovascular deaths were reduced by 32% (73 vs 50,  $p = 0.03$ ) and overall mortality by 22% (135 vs 106,  $p = 0.051$ ). There was no statistically significant difference between treatment groups in non-cardiovascular mortality, including cancer deaths. Pravastatin sodium also decreased the risk for undergoing myocardial revascularization procedures (coronary artery bypass graft surgery or coronary angioplasty) by 37% (80 vs 51 patients,  $p = 0.009$ ) and coronary angiography by 31% (128 vs 90,  $p = 0.007$ ).

**The West of Scotland Study excluded female patients, elderly subjects and most patients with familial hypercholesterolemia (FH). Consequently it has not been established to what extent the findings of the WOS study can be extrapolated to these subpopulations of hypercholesterolemic patients.**

- In patients with heterozygous FH, optimal reduction in total and LDL cholesterol necessitates a combination drug therapy in the majority of patients (see REFERENCES), (For homozygous FH see WARNINGS AND PRECAUTIONS - Use in Homozygous Familial

Hypercholesterolemia).

- Because information on familial combined hyperlipidemic (FCH) patients is not available from the WOS study, the effect of pravastatin sodium in this subgroup of high risk dyslipidemic patients could not be assessed.

### **Secondary Prevention of Cardiovascular Events**

Pravastatin sodium has been shown to be effective in reducing the risk for total mortality, CHD death, recurrent coronary events (including myocardial infarction), frequency of stroke or transient ischemic attacks (TIA), need for myocardial revascularization procedures, and need for hospitalization in patients with a history of either myocardial infarction or unstable angina pectoris.

In the Long-Term Intervention with Pravastatin in Ischemic Disease (LIPID) study, the effect of pravastatin sodium 40 mg daily was assessed in 9014 men and women with normal to elevated serum cholesterol levels (baseline Total-C=155-271 mg/dL [4.0-7.0 mmol/L]; median Total-C=218 mg/dL [5.66 mmol/L]; median LDL-C =150 mg/dL [3.88 mmol/L]), and who had experienced either a myocardial infarction or had been hospitalized for unstable angina pectoris in the preceding 3-36 months. Patients with a wide range of baseline levels of triglycerides were included ( $\leq 443$  mg/dL [5.0 mmol/L]) and enrollment was not restricted by baseline levels of HDL cholesterol. At baseline, 82% of patients were receiving aspirin, 47% were receiving beta blockers, and 76% were receiving antihypertensive medication. Patients in this multicenter, double-blind, placebo-controlled study participated for a mean of 5.6 years (median=5.9 years).

Treatment with pravastatin sodium significantly reduced the risk for CHD death by 24% ( $p=0.0004$ ). The risk for coronary events (either CHD death or nonfatal MI) was significantly reduced by 24% ( $p<0.0001$ ) in the pravastatin sodium treated patients. The risk for fatal or nonfatal myocardial infarction was reduced by 29% ( $p<0.0001$ ). Pravastatin sodium reduced both the risk for total mortality by 23% ( $p<0.0001$ ) and cardiovascular mortality by 25% ( $p<0.0001$ ). The risk for undergoing myocardial revascularization procedures (coronary artery bypass grafting or percutaneous transluminal coronary angioplasty) was significantly reduced by 20% ( $p<0.0001$ ) in the pravastatin sodium treated patients. Pravastatin sodium also significantly reduced the risk for stroke by 19% ( $p=0.0477$ ). Treatment with pravastatin sodium significantly reduced the number of days of hospitalization per 100 person-years of follow-up by 15% ( $p<0.001$ ). The effect of pravastatin sodium on reducing CHD events was consistent regardless of age, gender, or diabetic status. Among patients who qualified with a history of myocardial infarction, pravastatin sodium significantly reduced the risk for total mortality and for fatal or non-fatal MI (risk reduction for total mortality = 21%,  $p = 0.0016$ ; risk reduction for fatal or non-fatal MI = 25%,  $p = 0.0008$ ). Among patients who qualified with a history of hospitalization for unstable angina pectoris, pravastatin sodium significantly reduced the risk for total mortality and for fatal or non-fatal MI (risk reduction for total mortality = 26%,  $p = 0.0035$ ; risk reduction for fatal or non-fatal MI = 37%,  $p = 0.0003$ ).

In the Cholesterol and Recurrent Events (CARE) study the effect of pravastatin sodium 40 mg daily on coronary heart disease death and nonfatal MI was assessed in 4159 men and women with normal serum cholesterol levels (baseline mean Total-C=209 mg/dL [5.4 mmol/L]), and who had experienced a myocardial infarction in the preceding 3-20 months. At baseline, 83% of patients were receiving aspirin, 55% had undergone PTCA/CABG, 40% were receiving beta blockers, and 82% were receiving antihypertensive medication. Patients in this double-blind, placebo-controlled

study participated for an average of 4.9 years. Treatment with pravastatin sodium significantly reduced the rate of a recurrent coronary event (either CHD death or nonfatal MI) by 24% (274 patients with events [13.3%] in the placebo group vs. 212 patients [10.4%] in the pravastatin sodium group,  $p=0.003$ ). The reduction in risk for this combined endpoint was significant for both men and women; in women, the reduction in risk was 43% ( $p=0.033$ ). The risk of undergoing revascularization procedures (coronary artery bypass grafting or percutaneous transluminal coronary angioplasty) was significantly reduced by 27% ( $p<0.001$ ) in the pravastatin sodium treated patients (391 [19.6%] vs 294 [14.2%] patients). Pravastatin sodium also significantly reduced the risk for stroke by 32% ( $p=0.032$ ), and stroke or transient ischemic attack (TIA) combined by 26% (124 [6.3%] vs 93 [4.7%] patients,  $p=0.025$ ).

### **Atherosclerotic Disease Progression**

In two controlled trials [PLAC I, PLAC II] in patients with moderate hypercholesterolemia and atherosclerotic cardiovascular disease, pravastatin was effective in reducing the progressive course of atherosclerosis as evaluated by quantitative angiography and B-mode ultrasound. This effect may be associated with an improvement in the coronary endpoints (fatal or non fatal MI). No difference in total mortality was detected during the 3 years of double-blind therapy.

In PLAC I (Pravastatin Limitation of Atherosclerosis in the Coronary Arteries), a 3-year, placebo-controlled, multicentre, randomized trial, of 408 patients with moderate hypercholesterolemia (baseline LDL-C range = 3.37 - 4.92 mmol/L (130-190 mg/dL)) and coronary artery disease, treatment with pravastatin reduced the rate of narrowing of the coronary artery lumen diameter as determined by quantitative angiography. The analyses of clinical cardiovascular events showed a favorable effect of pravastatin therapy on events that occurred > 90 days after randomization, as well as for events from the time of randomization. This effect was not accompanied by an improvement in the total mortality endpoint. In PLAC II (Pravastatin Lipids and Atherosclerosis in the Carotid Arteries), a 3-year, placebo-controlled trial, in 151 patients with moderate hypercholesterolemia (baseline LDL-C range = 3.76 - 4.92 mmol/L (145-190 mg/dL)) and coronary and carotid atherosclerosis, treatment with pravastatin significantly reduced the rate of progression of atherosclerosis in common carotid artery, as measured by B-mode ultrasound. The rate of progression of the mean-maximum intimal-medial thickness (IMT) was not significantly reduced. There was a decrease in the incidence of coronary events of borderline significance. No difference in total mortality was observed during the 3 years of double-blind therapy.

### **Solid Organ Transplant**

Myopathy has not been observed in clinical trials involving a total of 100 post-transplant patients (76 cardiac and 24 renal) treated concurrently for two years with pravastatin (10 - 40 mg) and cyclosporine some of whom also received other immunosuppressants. Further, in clinical trials involving small numbers of patients treated with pravastatin, together with niacin, there were no reports of myopathy.

## DETAILED PHARMACOLOGY

### Cell/Tissue Selective Inhibition of Cholesterol Synthesis

*In vitro* and animal studies have shown that pravastatin, a hydrophilic HMG-CoA reductase inhibitor, is tissue selective such that inhibitory activity is highest in those tissues with the highest rates of cholesterol synthesis, such as the liver and ileum.

In suspensions of freshly isolated rat hepatocytes and in one-day cultures of rat hepatocytes, pravastatin sodium showed potent inhibition of <sup>14</sup>C incorporation into cholesterol. In cultured human skin fibroblasts and other non hepatic cell types, pravastatin inhibited cholesterol synthesis 400 times less than in hepatocytes.

The accumulation of <sup>14</sup>C-pravastatin was concentration and time dependent in hepatocytes and barely detectable in fibroblasts.

In tissue slices from rats given oral doses of pravastatin sodium, cholesterol synthesis was inhibited by more than 90% in liver and ileum slices and was substantially lower or not detectable in other tissue slices such as prostate, testes and adrenal.

In the intact rat lens, pravastatin sodium inhibited cholesterol synthesis 10 times less than in liver from the same animals. The inhibition of sterol synthesis in lens epithelial lines derived from the mouse and the rabbit was 400 to 1500 times less than in rat hepatocytes.

### Specificity as an Inhibitor of HMG-CoA Reductase

The incorporation of <sup>14</sup>C-mevalonate, the product of HMG-CoA reductase reaction into sterols, was not affected in hepatocytes, fibroblasts, or CHO cells at concentrations of pravastatin sodium at least 20 times greater than those that inhibited <sup>14</sup>C-acetate incorporation into cholesterol.

At concentrations 500 times greater than those that inhibited acetate incorporation into cholesterol, pravastatin sodium did not alter the rate of incorporation of <sup>14</sup>C-acetate into total cell phospholipids in hepatocytes and the distribution of the radiolabel into the separate classes of phospholipids. Pravastatin sodium did not reduce the rate of incorporation of <sup>14</sup>C-acetate into triglycerides. These results demonstrate that pravastatin does not act in the sterol pathway at any step beyond the synthesis of mevalonate nor does it inhibit the enzymes required for the biosynthesis of two other major classes of lipids.

The inhibitory activity of pravastatin on the enzyme HMG-CoA reductase was 106 times greater than that demonstrated by pravastatin for HMG-CoA lyase. The active site of this enzyme, which also employs HMG-CoA as substrate, does not recognize pravastatin.

### General pharmacology

The effect of pravastatin sodium on major physiologic systems and isolated tissue and its agonist and antagonist effects towards principal neurohumoral transmitters or histamine, behavioral effects, convulsive threshold and tissue- or activity-specific effects were evaluated in animals or in vitro tissue preparations. With the exception of a moderate inhibition of gastric secretion at a dose of 300 mg/kg in rats, pravastatin sodium had no effect in any of these pharmacologic tests at doses of 1000 mg/kg in some species.

### **Pharmacokinetics**

Studies in rats, dogs and humans demonstrated that pravastatin sodium given orally has low bioavailability because of extensive first-pass hepatic extraction. Therefore, most of an oral dose of pravastatin sodium is delivered directly to the liver, the primary site of pharmacologic activity.

A relatively low extent of binding of pravastatin to plasma proteins was found in rats, dogs, monkeys and humans. The highest concentrations of <sup>14</sup>C-pravastatin were found in the excretory organs and the GI tract in rats (N=3-5), one dog and one monkey. Similar metabolic patterns and appreciable fecal excretion in rats, dogs, monkeys and man were also evident in these studies.

Dogs are unique as compared to all other species tested, including man, in that they have a much greater systemic exposure to pravastatin. Pharmacokinetic data from a study in dogs at a dose of 1.1 mg/kg (comparable to a 40 mg dose in humans) showed that the elimination of pravastatin is slower in dogs than in humans. Absolute bioavailability is two times greater in dogs compared to humans and estimated renal and hepatic extraction of pravastatin are about one-tenth and one-half, respectively, than those in humans. When concentrations of pravastatin in plasma or serum of rats, dogs, rabbits, monkeys and humans were compared, the exposure in dogs was dramatically higher, based on both C<sub>MAX</sub> and AUC. The mean AUC value in man at a therapeutic dose of 40 mg is approximately 100 times less than that in the dog at the no-effect dose of 12.5 mg/kg, and approximately 180 times lower than that in dogs at the threshold dose of 25 mg/kg for cerebral hemorrhage.

### **Placental Transfer**

Low levels of radioactivity were found in the fetuses of rats dosed orally with radiolabeled pravastatin sodium. Pravastatin sodium was also found to be secreted in the milk of rats.

## TOXICOLOGY

### Acute Toxicity

Species	Sex (N)	Route	LD50 (mg/kg)
Mouse	M (50)	Oral	10590
	F (50)		8939
Mouse	M (50)	i.v.	2114
	F (50)		2011
Mouse	M (50)	s.c.	2975
	F (50)		3667
Rat	M (20)	Oral	> 12000
	F (20)		> 12000
Rat	M (50)	i.v.	443
	F (50)		440
Rat	M (50)	s.c.	3172
	F (50)		4455
Dog	M (4)	Oral	> 800

Signs of toxicity in mice were decreased activity, irregular respiration, ptosis, lacrimation, soft stool, diarrhea, urine-stained abdomen, ataxia, creeping behavior, loss of righting reflex, hypothermia, urinary incontinence, pilo-erection convulsion and/or prostration.

Signs of toxicity in rats were soft stool, diarrhea, decreased activity, irregular respiration, waddling gait, ataxia, loss of righting reflex and/or weight loss.

### Subacute and Chronic Toxicity

The spectrum of effects produced by pravastatin in mice, rats, rabbits, dogs and monkeys shown on the following table is not unexpected in view of the magnitude of the dosage levels employed and the potency of pravastatin against the HMG-CoA reductase.

#### Target Organs Observed in Animal Studies

Organ	Mouse	Rat	Rabbit	Dog	Monkey
Liver, neoplastic effect	-	+	-	-	-
Liver, non-neoplastic effect	+	+	+	-	+
Kidney	-	-	+	-	+
Skeletal muscle	-	+	+	-	-
Brain	-	-	-	+	-

+ = Organ affected in some way by drug treatment

- = No effect observed in this organ in these species

On a mg/kg basis, rabbits appear to be more sensitive to the nephrotoxic effects of pravastatin sodium than monkeys, the only other species that exhibited renal toxicity. In rabbits, renal dysfunction and hepatic effects were observed at doses  $\geq 25$  mg/kg/day. In monkeys, hepatotoxicity

and nephrotoxicity occurred at doses of 100 mg/kg/day. The threshold dose for renal toxicity in rabbits is 31 times greater than the maximum human dose.

### Significant Adverse Changes

	Pravastatin	
	Minimal Toxic Dose (mg/kg/day)	No-Effect Dose (mg/kg/day)
<b>Mice</b>		
Single-cell necrosis in the liver	40	20
Elevated serum transaminase activity	20	10
<b>Rats</b>		
Hepatic tumors	100	40
Foci of hepato-cellular alteration	30	12
Elevated transaminase activity	100	50
Skeletal-muscle myolysis	400	250
<b>Rabbits</b>		
Death	400	100
Hepatocellular necrosis	100	25
Renal tubular degeneration	25	6.25
Skeletal-muscle myolysis	100	25
Elevated serum transaminase activity	100	25
<b>Dogs</b>		
Death	25	12.5
Cerebral hemorrhage	25	12.5
<b>Monkeys</b>		
Death	200	100
Hepatocellular necrosis	100	50
Renal tubular degeneration	100	50
Elevated serum transaminase activity	100	50

Noteworthy findings in these studies included varying degrees of hepatotoxicity in all species tested, renal toxicity in rabbits and monkeys, skeletal-muscle lesions in rabbits, CNS symptoms and death secondary to cerebral hemorrhage in dogs, and an increased incidence of hepatic lesions and evidence of hepatocarcinoma (the latter at 100 mg/kg) in rats treated for 2 years. In all cases, these changes occurred only at daily doses of 20 mg/kg or more (more than 25 times the maximum human dose).

The findings from the chronic toxicity in dogs are detailed on the following pages.

Species/ Strain	Sex	N/Dose	Dose (mg/kg/day)	Route	Time	Effects
<b>Subacute Toxicity</b>						
Dog Beagle	M F	3 3	0, 12.5, 50 or 200	Oral (capsule)	5 weeks	200 mg/kg: One dog died and 4 dogs sacrificed on days 11 to 22 after exhibiting ataxia and/or convulsions, salivation, urinary incontinence and/or defecation. Ecchymotic lesions (hemorrhagic foci) in the brain.
Dog Beagle	M F	6 6	0 or 100 (2M, 2F controls) (4M, 4F treated)	Oral (capsule)	13 weeks	100 mg/kg: One death (F) on day 42 preceded by marked decrease in activity, serous salivation and vomiting. Diapedetic hemorrhage and degeneration of venular endothelial cells in one F and the F that died.
<b>Chronic Toxicity</b>						
Dog Beagle	M F	4M, 4F at 12.5 and 25 - 6M, 6F at 0, 50 & 100	0, 12.5, 25, 50 or 100	Oral (capsule)	2 years	25 mg/kg: Two F sacrificed during weeks 60 and 61. One had lesions consistent with idiopathic coagulopathy. The other showed clinical signs of CNS toxicity prior to sacrifice and had brain lesions. <sup>1</sup>  50 mg/kg: All dogs showed clinical signs of CNS toxicity; 5/6 dogs had brain lesions. <sup>1</sup>  100 mg/kg: Three M and 5 F died or sacrificed between weeks 2 and 24. One M died in week 76. All dogs showed clinical signs of CNS toxicity prior to death/sacrifice. Nine/nine dogs had brain lesions.

<sup>1</sup> Brain lesions (primarily in the piriform lobes) were characterized by discrete multifocal perivascular capillary and venular hemorrhages. In more severe lesions, there was an increased number of focal perivascular hemorrhages and associated early degenerative neutrophil changes including vacuolization, edema, and mild neutrophil infiltration. Larger vascular elements were not involved. No vascular endothelial changes were present, based on light- and electron-microscopic studies.

### Chronic Toxicity (cont'd.)

In dogs, pravastatin sodium was toxic at high doses and caused cerebral hemorrhage with clinical evidence of acute CNS toxicity (e.g. ataxia, convulsions). A dose-response relationship with respect to the incidence of CNS toxicity was clearly evident. In dogs, the threshold dose for CNS toxicity is 25 mg/kg. The high systemic exposure to orally administered pravastatin in dogs (refer to DETAILED PHARMACOLOGY - pharmacokinetics) may be related to a greater bioavailability and slower elimination of pravastatin and likely plays an important role in the development of CNS lesions that occur in the dog.

Cerebral hemorrhages have not been observed to date in any other laboratory species and the CNS toxicity in dogs may represent a species-specific effect.



## **Reproduction and Teratology**

Aside from a slight maternal toxicity in rabbits at 50 mg/kg and in rats at 1000 mg/kg, there were no treatment-related findings.

In rabbits and rats at doses greater than 60 and 600 times respectively the maximum human dose, pravastatin sodium exerted no untoward effects on reproduction through the F1 generation in rats and did not cause any fetal or anatomic abnormalities through the F1 generation in rabbits and the F2 generation rats.

## **Carcinogenicity and Mutagenicity**

In mice and rats, treated for 21 months with oral doses approximately 12 and 25 times the maximum human dose respectively (i.e. 20 mg/kg daily and 40 mg/kg daily), pravastatin sodium was found to be non carcinogenic. After 86 and 104 weeks of dosing in mice and rats respectively, at oral doses approximately 60 times the maximum human dose (i.e. 100 mg/kg daily), statistically significant increases in the incidence of hepatocellular carcinoma were observed in male rats only.

In *in vivo* mutagenicity tests with i.p. doses up to 1400 mg/kg and in *in vitro* mutagenicity tests at concentrations up to 10 000 mcg/mL or plate, pravastatin sodium was found to be nonmutagenic.

Pravastatin was found to be non-genotoxic.

## REFERENCES

1. Brown W.V., Goldberg I.J., Ginsberg H.N.  
Treatment of Common Lipoprotein Disorders  
Prog Cardiovasc Dis 27 (1): 1-20, 1984
2. Byington R.P., Jukema J.W., Salonen J.T., Pitt B., Bruschke A.V., Hoen H., Furberg C.D., Mancini J.  
Reduction in cardiovascular events during pravastatin therapy. Pooled analysis of clinical events of the pravastatin atherosclerosis intervention program.  
Circulation 92: 2419 - 2425, 1995
3. Crouse J.R. et al.  
Pravastatin, Lipids and Atherosclerosis in the Carotid Arteries (PLAC II) Am J Cardiol 75: 455-459, 1995
4. Crouse J.R., Byington R.P., Bond M.G., Espeland M.A., Sprinkle J.W., McGovern M., Furberg C.D.  
Pravastatin, Lipids and Atherosclerosis in the Carotid Arteries: Design Features of a Clinical Trial with Carotid Atherosclerosis Outcome  
Controlled Clinical Trials 13: 495-506, 1992
5. Decoulx E., Millaire A., DeGroot P., Mahieux G., Ducloux G.  
Rhabdomyolysis Due to Pravastatin and Type I Macrocreatine Kinase  
Ann. Cardiol. Angeiol. 42(5): 267-269, 1993
6. Dobs A.S., Sarma P.S., Schteingart D.  
Long-Term Endocrine Function in Hypercholesterolemic Patients Treated With Pravastatin, a New 3-Hydroxy-3-Methylglutaryl Coenzyme A Reductase Inhibitor Metabolism 42: 1146-1152, 1993
7. East C, Bilheimer DW, Grundy SM.  
Combination Drug Therapy for Familial Combined Hyperlipidemia  
Ann Int. Med 109(1):25-32, 1988
8. Edelman S., Witztum J.L.  
Hyperkalemia during treatment with HMG-CoA reductase inhibitor  
New Eng J Med 320: 1219, 1989
9. Endo A.  
Compactin (ML-236B) and Related Compounds as Potential Cholesterol-Lowering Agents that Inhibit HMG-CoA Reductase  
J Med Chem 28 (4): 401-405, 1985
10. Eptastatin Sodium  
Drugs Future 12 (5): 437-442, 1987
11. Folkers K., Langsjoen P., Willis R., Richardson P., Xia L.J., Ye C.Q., Tamagawa H.

- Lovastatin decreases coenzyme Q levels in humans. Proc. nat. Acad. Sci. 87(22): 8931-8934, 1990
12. Furberg C.D. et al.  
Pravastatin, Lipids and Major Coronary Events  
Am J Cardiol 73: 1133-1134, 1994
  13. Ghirlanda G., Oradei A., Manto A., Lippa S., Liccioli L., et al.  
Evidence of Plasma CoQ10 Lowering Effect by HMG CoA Reductase Inhibitor. Double-Blind Placebo Control Study  
J Clin Pharmacol 33: 226-229, 1993
  14. Goto Y.  
The Profile of an HMG-CoA Reductase Inhibitor, CS-514 (SQ 31,000) In: Drugs Affecting Lipid Metabolism; R. Paoletti et al (eds)  
Springer-Verlag Berlin Heidelberg, pp 247-250, 1987
  15. Grundy S.M.  
HMG-CoA Reductase Inhibitors for Treatment of Hypercholesterolemia  
New Eng J Med 319 (1): 24-33, 1988
  16. Halkin A.  
HMG-CoA Reductase Inhibitor-Induced Impotence  
Ann Pharmacother 30:192, 1996
  17. Hoeg J.M., Brewer H.B.  
3-Hydroxy-3-Methylglutaryl-Coenzyme A Reductase Inhibitors in the Treatment of Hypercholesterolemia  
JAMA 258 (24): 3532-3536, 1987
  18. Hunninghake D.B., Stein E.A., Mellies M.J.  
Effects of One Year Treatment with Pravastatin, an HMG-CoA Reductase Inhibitor, on Lipoprotein A  
J Clin Pharmacol 33: 574-580, 1993
  19. Hunninghake D.B., Goldberg A.C., Insull W., Juo P., Mellies M., Pan H.Y.  
Pravastatin: A Tissue-Selective Once Daily HMG-CoA Reductase Inhibitor in the Treatment of Primary Hypercholesterolemia  
J Am Coll Cardiol 11 (2) (Suppl. A): 8A, 1988
  20. Illingworth D.R. Drug Therapy in Heterozygous Familial Hypercholesterolemia Am. J. Cardiol. 1993; 72: 54D - 58D
  21. Kazumi T., Yoshino G., Kasama T., Iwatani I., Iwai M., Morita S., Baba S.  
Effects of CS-514, a New Inhibitor of HMG-CoA Reductase, on Plasma Lipids, Lipoproteins and Apoproteins in Patients with Primary Hypercholesterolemia Horm Metabol Res 18: 654-655, 1986
  22. Kliem V., Wanner C., Eisenhauer T., Obricht C.J., Doll R., Boddaert M., O'Grady P., Krekler M., Mangold B., Christians U.

Comparison of pravastatin and lovastatin in renal transplant patients receiving cyclosporin.  
Transplant Proc. 28(6): 3126-3128, 1996

23. Lipid Research Clinics Coronary Primary Prevention Trial Results: I. Reduction in Incidence of Coronary Heart Disease. II. The Relationship of Reduction in Incidence of Coronary Heart Disease to Cholesterol Lowering  
JAMA 251: 351-374, 1984
24. Mabuchi H., Takeda R.  
Inhibitors of 3-Hydroxy-3-Methylglutaryl Coenzyme A Reductase: Compactin and its Analogues  
In: Pharmacological Control of Hyperlipidemia: Proceedings of an International Telesymposium on Hyperlipidaemia. J.R. Prous Science Publishers; Barcelona, Spain, pp 251-266, 1986
25. Mabuchi H., Kamon N., Fujita H., Michishita I., Takeda M., Kojinami K., Hoh H., Wakasugi T., Takeda R.  
Effects of CS-514 on Serum Lipoprotein Lipid and Apolipoprotein Levels in Patients with Familial Hypercholesterolemia  
Metabolism 36 (5): 475-479, 1987
26. Mabuchi H., Kamon N., Fujita H., Michishita I., Takeda M., Kajinami K., Itoh H., Wakasugi T., Takeda R.  
Long-Term Effects of CS-514 on Serum Lipoprotein Lipid and Apolipoprotein Levels in Patients with Familial Hypercholesterolemia  
In: Drugs Affecting Lipid Metabolism; R. Paoletti et al (eds) Springer-Verlag Berling Heidelberg, pp 261-268, 1987
27. Mabuchi H., Fujita H., Michishita I. et al  
Effects of CS-514 (eptastatin), an Inhibitor of 3-Hydroxy-3-Methylglutaryl-Coenzyme A- (HMG-CoA) Reductase, on Serum Lipid and Apolipoprotein Levels in Heterozygous Familial Hypercholesterolemia Patients Treated by Low Density Lipoprotein (LDL)- Apheresis  
Atherosclerosis 72: 183-188, 1988
28. Mellies M.J., DeVault A.R., Kassler-Taub K., McGovern M.E., Pan H.Y.  
Pravastatin Experience in Elderly and Non-Elderly Patients  
Atherosclerosis 101: 97-110, 1993
29. Mishkel M.  
Drug Treatment of Hypercholesterolemia  
Drug Protocol 3 (1): 9-17, 1988
30. Nakaya N., Homma Y., Tamachi H., Goto Y.  
The Effect of CS-514, an Inhibitor of HMG-CoA Reductase, on Serum Lipids on Healthy Volunteers  
Atherosclerosis 61: 125-128, 1986
31. Nakaya N., Homma Y., Tamachi H., Shigematsu H., Hata Y., Goto Y.  
The Effect of CS-514 on Serum Lipids and Apolipoproteins in Hypercholesterolemic

Subjects

JAMA 257 (22): 3088-3093, 1987

32. Nakaya N. and Goto Y.  
Effect of CS-514 on Hypercholesterolemic Patients  
In: Drugs Affecting Lipid Metabolism; R. Paoletti et al (eds) Springer-Verlag Berlin Heidelberg, pp 274-277, 1987
33. Pan H.Y., Willard D.A., Funke P.T., McKinstry D.  
The Clinical Pharmacology of SQ 31,000 (CS-514) in Healthy Subjects  
In: Drugs Affecting Lipid Metabolism; R. Paoletti et al (eds) Springer-Verlag Berlin Heidelberg, pp 255-259, 1987
34. Perault M.C., Ladouch-Bures L., Dejean C., Delauney C., Pouget Abadie J.F., Vandel B.  
Rhabdomyolyse associée à la prise de pravastatine (Vasten) *Thérapie* 48: 483-501, 1993
35. Saku K. et al  
Long-Term Effects of CS-514 (HMG-CoA Reductase Inhibitor) on Serum Lipids, Lipoproteins, and Apolipoproteins in Patients with Hypercholesterolemia  
*Curr Ther Res* 42 (3): 491-500, 1987
36. Saito Y., Goto Y., Nakaya N. et al  
Dose-Dependent Hypolipidemic Effect of an Inhibitor of HMG-CoA Reductase, Pravastatin (CS-514) in Hypercholesterolemic Subjects. A Double-Blind Test *Atherosclerosis* 7: 205-211, 1988
37. Scanu A.M.  
Lipoprotein(a) as a Cardiovascular Risk Factor  
*Trends Cardiovasc Med* 1: 294-299, 1991
38. Shephard J., Cobbe S.M., Ford I., Isles C.G., Lorimer A.R., Macfarlane P.W., McKillop J.H., Packard C.J.  
Prevention of Coronary Heart Disease Using Pravastatin in Hypercholesterolemic Men  
*NEJM* 333(20): 1301 - 1307, 1995
39. Smith G.D., Song F., Sheldon T.A.  
Cholesterol Lowering and Mortality: The Importance of Considering Initial Level of Risk  
*Br Med J* 306: 1367-1371, 1993
40. Stein E.A.  
Treatment of Familial Hypercholesterolemia with Drugs in Children  
*Arteriosclerosis Suppl. I*, 9: I-145 - I-151, 1989
41. Tobert J.A.  
New Developments in Lipid-Lowering Therapy: The Role of Inhibitors of Hydroxymethylglutaryl-Coenzyme A Reductase  
*Circulation* 76 (3): 534-538, 1987

42. Tsujita Y., Kuroda M., Shimada Y., Tanzawa K., Arai M., Kaneko I., Tanaka M., Masuda H., Tarumi C., Watanabe Y., Fujii S.  
CS-514, a Competitive Inhibitor of 3-Hydroxy-3-Methylglutaryl Coenzyme A Reductase: Tissue-Selective Inhibition of Sterol Synthesis and Hypolipidemic Effect on Various Animal Species  
*Biochim Biophys Acta* 877: 50-60, 1986
43. The West of Scotland Coronary Prevention Study Group  
A Coronary Primary Prevention Study of Scottish Men Aged 45-55 Years: Trial Design  
*J Clin Epidemiol* 45, No. 8, 849-860, 1992
44. The WOSCOPS Study Group  
Screening Experience and Baseline Characteristics in the West of Scotland Coronary Prevention Study  
*Am J Cardiol* 76: 485-491, 1995
45. Yamamoto A., Yokoyama S., Yamamura T.  
Combined Drug Treatment and Plasmapheresis for Familial Hypercholesterolemia In: Pharmacological Control of Hyperlipidemia: Proceedings of an International Telesymposium on Hyperlipidaemia. J.R. Prous Science Publishers; Barcelona, Spain, pp 333-342, 1986
46. Yamamoto A., Yokoyama S., Yamamura T.  
Intensive Drug Treatment for Familial Hypercholesterolemia In: Drugs Affecting Lipid Metabolism; R. Paoletti et al (eds) Springer-Verlag Berlin Heidelberg, pp 269-273, 1987
47. Yoshimura N., Takahiro O. et al  
The Effects of Pravastatin on Hyperlipidemia in Renal Transplant Recipients  
*Transplantation* 53: 94-99, 1992
48. Yoshino G., Kazumi T., Iwai M., Kasama T., Iwatani I., Matsuba K., Inui A., Venoyama R., Yokono K., Otsuki M., Baba S.  
CS-514 Suppresses Plasma Triglyceride in Hypertriglyceridemic Subjects Without Modifying a Lipoprotein Structural Model  
*Horm Metabol Res* 19: 513-514, 1987
49. Yoshino G., Kazumi T., Iwai M., Iwatani I., Matsuba K., Kasama T., Matsushita M., Otsuki M., Baba S.  
Effects of CS-514 on Plasma Lipids and Lipoprotein Composition in Hypercholesterolemic Subjects  
*Atherosclerosis* 71: 95-101, 1986.
50. PRAVACHOL<sup>®</sup> (pravastatin sodium) Product Monograph, Squibb Canada Division  
Bristol-Myers Squibb Canada Inc., Date of Revision: November 30, 2015,  
Control number 187002.

**PART III: CONSUMER INFORMATION****Pr pendo-PRAVASTATIN  
Pravastatin Sodium, House Standard**

**This leaflet is part III of a three-part "Product Monograph" published when pendo-PRAVASTATIN was approved for sale in Canada and is designed specifically for Consumers. This leaflet is a summary and will not tell you everything about pendo-PRAVASTATIN. Contact your doctor or pharmacist if you have any questions about the drug.**

**ABOUT THIS MEDICATION****What the medication is used for:**

pendo-PRAVASTATIN is available only with your physician's prescription. It is to be used as an adjunct to a medically recommended and carefully supervised diet for the long-term treatment of hypercholesterolemia and is not a substitute for such a diet. This has been shown to decrease the chances of experiencing a first or second heart attack, or stroke, and may help prevent heart disease if caused by cholesterol clogging the blood vessels, or slow the progression of atherosclerosis (hardening) of the arteries that nourish your heart, so-called coronary heart disease (CHD). In addition, depending on your condition, your physician may recommend an appropriate regimen of exercise, weight control and other measures.

**What it Does:**

pendo-PRAVASTATIN lowers the level of cholesterol, particularly Low Density Lipoprotein (LDL) cholesterol, in the blood. pendo-PRAVASTATIN reduces cholesterol production by the liver and induces some changes of cholesterol transport and disposition in the blood and tissues.

**When it should not be used:**

Do not use pendo-PRAVASTATIN if you:

- are pregnant, since its use in the event of pregnancy may harm the unborn. Only female patients who are highly unlikely to conceive can be candidates for pendo-PRAVASTATIN treatment. In the event of pregnancy during treatment, pendo-PRAVASTATIN should be discontinued and the physician should be informed.
- know that you are allergic to pravastatin or any of the non-medicinal ingredients of pendo-PRAVASTATIN.
- have liver disease.
- are less than 18 years of age. The safety of pendo-PRAVASTATIN in this age group has not been established.

**What the medicinal ingredient is:**

Pravastatin sodium

**What the non-medicinal ingredients are:**

Colloidal Silicon Dioxide, Copovidone, Croscarmellose Sodium, Dibasic Calcium Phosphate, Iron Oxide Red (10 mg only), Iron Oxide Yellow (20 mg only), Lactose, Magnesium Stearate, Microcrystalline Cellulose, Polyethylene Glycol.

**What dosage forms it comes in:**

**Tablets:** 10 mg and 20 mg

**WARNINGS AND PRECAUTIONS**

Use only as specifically directed. Do not alter the dosage unless ordered to do so by your physician. Check with your physician before discontinuing medication since this may result in an increase of your blood lipids.

Before taking pendo-PRAVASTATIN, tell your doctor or pharmacist if you:

- are breast-feeding or intend to breast-feed,
- have thyroid problems,
- have a family history of muscular disorders,
- have had any past problems with the muscles (pain, tenderness), after using an HMG-CoA reductase inhibitor ("statin") such as atorvastatin , fluvastatin , lovastatin , pravastatin , rosuvastatin or simvastatin , or have developed an allergy or intolerance to them,
- have kidney or liver problems,
- have diabetes,
- have undergone surgery or other tissue injury,
- do excessive physical exercise.
- suffer from alcohol abuse
- are 65 years of age or older.

**Pregnancy**

Before using this medication, discuss the following with your doctor:

- Cholesterol compounds are essential elements for the development of a fetus.
- Cholesterol-lowering drugs can harm the fetus. If you are of child-bearing age, discuss with your doctor the potential hazards to the fetus and the importance of birth control methods.
- pendo-PRAVASTATIN should not be used by pregnant women. If you become pregnant, discontinue use immediately and discuss with your doctor.

Slightly increased blood sugar can occur when you take HMG-CoA reductase inhibitor ("statin"). Discuss with the doctor your risk of developing diabetes

## INTERACTIONS WITH THIS MEDICATION

pendo-PRAVASTATIN may interact with other drugs, including those you take without a prescription. You must tell your doctor or pharmacist about all drugs, including prescription and non-prescription, herbal products and supplements, you are taking or planning to take before you take pendo-PRAVASTATIN.

You should tell your doctor if you are taking other cholesterol lowering medication such as fibrates (gemfibrozil, fenofibrate), niacin, or ezetimibe. If you take these drugs and pendo-PRAVASTATIN together, you may be at an increased risk of myopathy (muscle disease with aching or weakness).

If you are taking a bile acid-binding resin (such as cholestyramine, colestipol), your doctor will recommend that you take pendo-PRAVASTATIN either one hour or more before or at least four hours following the resin. Taking them together causes lower amounts of pendo-PRAVASTATIN in the blood, making it less effective.

If you are taking cyclosporine, your doctor may need to adjust the dose of pendo-PRAVASTATIN.

If you are taking certain antibiotics (such as azithromycin or clarithromycin) to treat an infection your doctor may need to adjust the dose of pendo-PRAVASTATIN.

Excessive alcohol intake should be avoided when taking pendo-PRAVASTATIN. Tell your doctor if you regularly drink *three or more* alcoholic drinks daily.

## PROPER USE OF THIS MEDICATION

- Do not change the dose unless directed by a doctor.
- pendo-PRAVASTATIN should be taken as a single dose at bedtime, as prescribed by your physician.
- Your physician will monitor your clinical condition and your blood tests at regular intervals. It is important to have these check-ups done on schedule. Please keep your appointments accurately.
- Notify your physician about any illness which may develop during your treatment with pendo-PRAVASTATIN and about any new prescription or non-prescription medication you may take. If you require medical help for other reasons, inform the attending physician that you are taking pendo-PRAVASTATIN,
- Notify your physician if you are going to have major surgery or have sustained a severe injury,
- Notify your physician of any muscle pain, tenderness or weakness developing during treatment with pendo-PRAVASTATIN (see SIDE EFFECTS AND WHAT TO DO ABOUT THEM).

### Usual adult dose:

The recommended starting dose is 20 mg once daily at bed time.

Patients who require a large dose reduction in cholesterol may be started at 40 mg once daily. The dose of 80 mg once daily should be reserved for patients who do not achieve their treatment goal with lower doses. pendo-PRAVASTATIN may be taken without regard to meals.

### Overdose:

If you think you have taken too much pendo-PRAVASTATIN contact your doctor, pharmacist, hospital emergency department or regional Poison Control Centre immediately, even if there are no symptoms.

## SIDE EFFECTS AND WHAT TO DO ABOUT THEM

Side effects may include:

- Abdominal pain, constipation, diarrhea, nausea, indigestion
- Headache, dizziness
- Skin rashes
- Persistent cough
- Trouble sleeping, insomnia, nightmares
- Sexual problems
- Depression, anxiety, nervousness

Muscle effects

Side effects such as muscle pain, muscle disease with aching or weakness, rhabdomyolysis (a muscle wasting disease), associated tenderness, and rare cases of rhabdomyolysis leading to kidney failure have been reported with other drugs of this class, known as HMG-CoA reductase inhibitors ("statins"), including pendo-PRAVASTATIN.

As these muscle problems are on rare occasions serious, you should contact your physician promptly if you experience any of the side effects listed in the Table below.

pendo-PRAVASTATIN can also cause abnormal blood test results. Your doctor will decide when to perform blood tests and will interpret the results.



**SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM**

Symptom / effect		Talk with your doctor or pharmacist		Stop taking drug and seek immediate emergency medical attention
		Only if severe	In all cases	
Rare	muscle pain that you cannot explain		√	
	muscle tenderness or muscle weakness or muscle cramping		√	
	generalized weakness, especially if you do not feel well (ie. fever or fatigue)		√	
	brownish or discoloured urine		√	
	Symptoms of liver problems (upper belly pain, dark urine, itchy skin, nausea or vomiting, loss of appetite, pale stools, yellowing of skin or the whites of your eyes)			√
	Cough and/or shortness of breath		√	
	Severe Skin reaction with fever, sore mouth or throat, blistering and/or peeling of your skin or mucous membranes			√
Uncommon	Allergic Reaction: rash, hives, swelling of the face, lips, tongue or throat, difficulty swallowing or breathing			√
	Memory problems: poor memory, forgetfulness, confusion, memory loss	√		
Common	Vision problems, blurred vision		√	

**SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM**

Symptom / effect		Talk with your doctor or pharmacist		Stop taking drug and seek
Unknown	Increased Blood Sugar: Frequent urination, thirst and hunger.	√		
	Chest pain		√	

*This is not a complete list of side effects. For any unexpected effects while taking pendo-PRAVASTATIN, contact your doctor or pharmacist.*

**HOW TO STORE IT**

Store between 15°C and 30°C. Protect from moisture and light.

**KEEP OUT OF THE REACH AND SIGHT OF CHILDREN.**

**REPORTING SUSPECTED SIDE EFFECTS**

You can report any suspected adverse reactions associated with the use of health products to the Canada Vigilance Program by one of the following 3 ways:

- Report online at [www.healthcanada.gc.ca/medeffect](http://www.healthcanada.gc.ca/medeffect)
- Call toll-free at 1-866-234-2345
- Complete a Canada Vigilance Reporting Form and:
  - Fax toll-free to 1-866-678-6789, or
  - Mail to: **Canada Vigilance Program  
Health Canada  
Postal Locator 0701E  
Ottawa, Ontario  
K1A 0K9**

Postage paid labels, Canada Vigilance Reporting Form and the adverse reaction reporting guidelines are available on the MedEffect™ Canada Web site at [www.healthcanada.gc.ca/medeffect](http://www.healthcanada.gc.ca/medeffect).

*NOTE: Should you require information related to the management of side effects, contact your health professional. The Canada Vigilance Program does not provide medical advice.*

**MORE INFORMATION**

This document plus the full product monograph, prepared for health professionals, can be obtained by contacting PENDOPHARM, division of Pharmascience Inc. at 1-888-550-6060.

This leaflet was prepared by  
**PENDOPHARM, division of Pharmascience Inc.**  
Montreal Canada  
H4P 2T4

[www.pendopharm.com](http://www.pendopharm.com)

Last revised: July 25, 2016