

PRODUCT MONOGRAPH

 **JANUVIA[®]**

sitagliptin tablets
(as sitagliptin phosphate monohydrate)

25, 50 and 100 mg

Oral Antihyperglycemic Agent
DPP-4 inhibitor
Incretin Enhancer

Merck Canada Inc.

16750 route Transcanadienne
Kirkland, QC Canada H9H 4M7
www.merck.ca

Date of Revision:
March 13, 2017

Submission Control No: 201760

Table of Contents

PART I: HEALTH PROFESSIONAL INFORMATION	3
SUMMARY PRODUCT INFORMATION.....	3
INDICATIONS AND CLINICAL USE	3
CONTRAINDICATIONS.....	4
WARNINGS AND PRECAUTIONS	4
ADVERSE REACTIONS	7
DRUG INTERACTIONS	22
DOSAGE AND ADMINISTRATION	24
OVERDOSAGE.....	25
ACTION AND CLINICAL PHARMACOLOGY	26
STORAGE AND STABILITY	30
DOSAGE FORMS, COMPOSITION AND PACKAGING.....	30
PART II: SCIENTIFIC INFORMATION	31
PHARMACEUTICAL INFORMATION	31
CLINICAL TRIALS	32
DETAILED PHARMACOLOGY	45
TOXICOLOGY	46
REFERENCES.....	48
PART III: CONSUMER INFORMATION	49

JANUVIA[®]

sitagliptin tablets
(as sitagliptin phosphate monohydrate)

PART I: HEALTH PROFESSIONAL INFORMATION

SUMMARY PRODUCT INFORMATION

Route of Administration	Dosage Form / Strength	Clinically Relevant Non-medicinal Ingredients
Oral	Tablet 25, 50 and 100 mg	<i>For a complete listing see DOSAGE FORMS, COMPOSITION AND PACKAGING section.</i>

INDICATIONS AND CLINICAL USE

Monotherapy

JANUVIA[®] (sitagliptin) is indicated as an adjunct to diet and exercise to improve glycemic control in adult patients with type 2 diabetes mellitus and for whom metformin is inappropriate due to contraindications or intolerance.

Combination with Metformin

JANUVIA[®] is indicated in combination with metformin in adult patients with type 2 diabetes mellitus to improve glycemic control when diet and exercise, plus metformin do not provide adequate glycemic control.

Combination with Metformin and a Sulfonylurea

JANUVIA[®] is indicated in combination with metformin and a sulfonylurea in adult patients with type 2 diabetes mellitus to improve glycemic control when diet and exercise, and dual therapy with these agents, do not provide adequate glycemic control.

Combination with Insulin

JANUVIA[®] is indicated as add-on combination therapy with premixed or long/intermediate acting insulin (with or without metformin) in adult patients with type 2 diabetes mellitus as an adjunct to diet and exercise to improve glycemic control when diet and exercise, and therapy with premixed or long/intermediate acting insulin (with or without metformin) do not provide adequate glycemic control (see CLINICAL TRIALS).

Combination with Pioglitazone

JANUVIA[®] is indicated in combination with pioglitazone in adult patients with type 2 diabetes mellitus to improve glycemic control when diet and exercise, plus pioglitazone do not provide adequate glycemic control.

Combination with Metformin and Pioglitazone

JANUVIA[®] is indicated in combination with metformin and pioglitazone in adult patients with type 2 diabetes mellitus to improve glycemic control when diet and exercise, and dual therapy with these agents, do not provide adequate glycemic control.

Geriatrics (≥65 years of age): No dosage adjustment is required based on age however, greater sensitivity of some older individuals cannot be ruled out (see WARNINGS AND PRECAUTIONS, DOSAGE AND ADMINISTRATION and ACTION AND CLINICAL PHARMACOLOGY).

Pediatrics (<18 years of age): Safety and effectiveness of JANUVIA[®] in pediatric patients have not been established. Therefore, JANUVIA[®] should not be used in this population.

CONTRAINDICATIONS

Patients who are hypersensitive to this drug or to any ingredient in the formulation (see WARNINGS AND PRECAUTIONS, Hypersensitivity Reactions and ADVERSE REACTIONS, Post-Marketing Adverse Drug Reactions). For a complete listing, see the DOSAGE FORMS, COMPOSITION AND PACKAGING section of the product monograph.

WARNINGS AND PRECAUTIONS

General

JANUVIA[®] should not be used in patients with type 1 diabetes or for the treatment of diabetic ketoacidosis.

Pancreatitis

There have been reports of acute pancreatitis, including fatal and non-fatal hemorrhagic or necrotizing pancreatitis, in patients taking JANUVIA[®]. In a long-term cardiovascular outcomes trial (see ADVERSE REACTIONS and CLINICAL TRIALS, TECOS Cardiovascular Safety Study), there were two adjudication-confirmed deaths due to acute pancreatitis in patients treated with JANUVIA[®] compared to none in the placebo group. After initiation of JANUVIA[®], patients should be observed carefully for signs and symptoms of pancreatitis. If pancreatitis is suspected, JANUVIA[®] should promptly be discontinued and appropriate management should be initiated. Risk factors for pancreatitis include a history of: pancreatitis, gallstones, alcoholism, or hypertriglyceridemia.

Hypoglycemia

When JANUVIA[®] was used in combination with metformin and a sulfonylurea, or with a stable dose of insulin (with or without metformin), the incidence of hypoglycemia was increased over that of placebo in combination with metformin and a sulfonylurea or that of placebo in combination with a stable dose of insulin (with or without metformin) (see ADVERSE REACTIONS). To reduce the risk of hypoglycemia associated with these indications, a lower dose of sulfonylurea or insulin may be considered (see DOSAGE AND ADMINISTRATION).

Hypersensitivity Reactions

There have been post-marketing reports of serious hypersensitivity reactions in patients treated with JANUVIA[®]. These reactions include anaphylaxis, angioedema, and exfoliative skin conditions including Stevens-Johnson syndrome. Onset of these reactions occurred within the first 3 months after initiation of treatment with JANUVIA[®], with some reports occurring after the first dose. If a hypersensitivity reaction is suspected, discontinue JANUVIA[®], assess for other potential causes for the event, and institute alternative treatment for diabetes (see CONTRAINDICATIONS and ADVERSE REACTIONS, Post-Marketing Adverse Drug Reactions).

Immune

Immunocompromised patients: A dose-related mean decrease in absolute lymphocyte count was observed with other members of this class. When clinically indicated, such as in settings of unusual or prolonged infection, lymphocyte count should be measured. The effect of sitagliptin on lymphocyte counts in patients with lymphocyte abnormalities (e.g. human immunodeficiency virus) is unknown. Immunocompromised patients, such as patients who have undergone organ transplantation or patients diagnosed with human immunodeficiency syndrome have not been studied in the sitagliptin clinical program. Therefore, the efficacy and safety profile of sitagliptin in these patients has not been established.

Skin

With other members of this class, ulcerative and necrotic skin lesions have been reported in monkeys in non-clinical toxicology studies. There is limited experience in patients with diabetic skin complications. In keeping with routine care of the diabetic patient, monitoring for skin disorders is recommended.

Bullous Pemphigoid

Postmarketing cases of bullous pemphigoid requiring hospitalization have been reported with the use of DPP-4 inhibitors, including JANUVIA[®]. In reported cases, patients typically recovered with topical or systemic immunosuppressive treatment and discontinuation of the DPP-4 inhibitor. Tell patients to report development of blisters or erosions while receiving JANUVIA[®]. If bullous pemphigoid is suspected, JANUVIA[®] should be discontinued and referral to a dermatologist should be considered for diagnosis and appropriate treatment.

Renal

Renal adverse events, including acute renal failure, have been observed during clinical trials and post-marketing use of sitagliptin in patients with and without known risk factors (see ADVERSE REACTIONS). Since JANUVIA[®] is renally excreted, dose adjustment is required in patients with moderate to severe renal impairment as well as end-stage renal disease patients on dialysis (see WARNINGS AND PRECAUTIONS, Monitoring and Laboratory Tests; and ACTION AND CLINICAL PHARMACOLOGY and DOSAGE AND ADMINISTRATION).

Hepatic

There are limited clinical experiences in patients with moderate hepatic insufficiency and no clinical experience in patients with severe hepatic insufficiency. Use in patients with severe hepatic insufficiency is not recommended (see ACTION AND CLINICAL PHARMACOLOGY).

Endocrine and Metabolism

The effectiveness of oral antidiabetic drugs in lowering blood glucose to a targeted level decreases in many patients over a period of time. This phenomenon, which may be due to progression of the underlying disease or to diminished responsiveness to the drug, is known as secondary failure, to distinguish it from primary failure in which the drug is ineffective during initial therapy. Should secondary failure occur with JANUVIA[®], therapeutic alternatives should be considered.

Special Populations

Pregnant Women: There are no adequate and well-controlled studies in pregnant women; therefore, the safety of JANUVIA[®] in pregnant women is not known. JANUVIA[®] is not recommended for use in pregnancy (see also TOXICOLOGY).

Nursing Women: Sitagliptin is secreted in the milk of lactating rats. It is not known whether sitagliptin is secreted in human milk. Therefore, JANUVIA[®] should not be used by a woman who is nursing.

Pediatrics (<18 years of age): Safety and effectiveness of JANUVIA[®] in pediatric patients have not been established. Therefore, JANUVIA[®] should not be used in this population.

Geriatrics (≥65 years of age): In clinical studies, no overall differences in safety or effectiveness were observed between subjects 65 years and over and younger subjects. While this and other reported clinical experience have not identified differences in responses between the geriatric and younger patients, greater sensitivity of some older individuals cannot be ruled out.

This drug is known to be substantially excreted by the kidney. Renal function should be assessed prior to initiating dosing and periodically thereafter in geriatric patients because they are more likely to have decreased renal function (see DOSAGE AND ADMINISTRATION and ACTION AND CLINICAL PHARMACOLOGY).

Monitoring and Laboratory Tests

Response to all diabetic therapies should be monitored by periodic measurements of blood glucose and HbA_{1c} levels, with a goal of decreasing these levels towards the normal range. HbA_{1c} is especially useful for evaluating long-term glycemic control.¹ Sitagliptin is substantially excreted by the kidney. Renal function should be assessed prior to initiating dosing and periodically thereafter.

ADVERSE REACTIONS

Adverse Drug Reaction Overview

JANUVIA[®] was generally well tolerated in controlled clinical studies as monotherapy and as part of combination therapy with metformin or combination therapy with metformin and a sulfonylurea agent, or add-on combination therapy with insulin (with or without metformin) or as add-on combination therapy with pioglitazone (with or without metformin).

The incidences of serious adverse reactions and discontinuation of therapy due to clinical adverse reactions were generally similar to placebo. The most frequent adverse events in trials of JANUVIA[®] as monotherapy (placebo-controlled) and as add-on combination therapy with metformin (reported regardless of causality, and more common with JANUVIA[®] than other treatments) was nasopharyngitis. The most frequent adverse events with JANUVIA[®] as add-on combination therapy with metformin and a sulfonylurea agent, or as add-on combination therapy with insulin (with or without metformin), was hypoglycemia.

Clinical Trial Adverse Drug Reactions

Because clinical trials are conducted under very specific conditions the adverse reaction rates observed in the clinical trials may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse drug reaction information from clinical trials is useful for identifying drug-related adverse events and for approximating rates.

Monotherapy:

Two placebo-controlled monotherapy studies, one of 18- and one of 24-week duration, included patients treated with JANUVIA[®] 100 mg once daily and patients given placebo. Adverse events, reported regardless of causality assessment, in $\geq 1\%$ of patients in these two studies pooled are shown in Table 1.

Table 1 – Adverse events $\geq 1\%$ in any treatment group (regardless of causality) reported in patients treated with JANUVIA[®] 100 mg or placebo in pooled 18 and 24-week placebo-controlled, double-blind clinical trials of JANUVIA[®] as monotherapy

Body system/Organ class Adverse event	Number of patients (%)	
	Sitagliptin 100 mg n=443	Placebo n=363
Eye disorders		
Conjunctivitis	3 (0.7)	4 (1.1)
Gastrointestinal disorders		
Abdominal pain	5 (1.1)	6 (1.7)
Constipation	13 (2.9)	5 (1.4)
Diarrhea	19 (4.3)	10 (2.8)
Gastritis	2 (0.5)	4 (1.1)
Nausea	7 (1.6)	3 (0.8)
Vomiting	3 (0.7)	4 (1.1)

Table 1 – Adverse events $\geq 1\%$ in any treatment group (regardless of causality) reported in patients treated with JANUVIA[®] 100 mg or placebo in pooled 18 and 24-week placebo-controlled, double-blind clinical trials of JANUVIA[®] as monotherapy

Body system/Organ class Adverse event	Number of patients (%)	
	Sitagliptin 100 mg n=443	Placebo n=363
General disorders and administration site conditions		
Fatigue	5 (1.1)	9 (2.5)
Edema peripheral	7 (1.6)	4 (1.1)
Pain	0 (0.0)	4 (1.1)
Infections and infestations		
Bronchitis	5 (1.1)	6 (1.7)
Gastroenteritis	6 (1.4)	4 (1.1)
Influenza	19 (4.3)	16 (4.4)
Nasopharyngitis	23 (5.2)	12 (3.3)
Pharyngitis	5 (1.1)	1 (0.3)
Sinusitis	6 (1.4)	9 (2.5)
Upper respiratory tract infection	29 (6.5)	24 (6.6)
Urinary tract infection	8 (1.8)	9 (2.5)
Viral infection	2 (0.5)	4 (1.1)
Viral upper respiratory tract infection	5 (1.1)	1 (0.3)
Injury, poisoning and procedural complications		
Limb injury	3 (0.7)	4 (1.1)
Investigations		
Blood glucose increased	7 (1.6)	13 (3.6)
Metabolism and nutrition disorders		
Hyperglycemia	5 (1.1)	7 (1.9)
Hypoglycemia	5 (1.1)	2 (0.6)
Musculoskeletal and connective tissue disorders		
Arthralgia	4 (0.9)	9 (2.5)
Back pain	14 (3.2)	12 (3.3)
Muscle spasm	6 (1.4)	4 (1.1)
Myalgia	6 (1.4)	4 (1.1)
Neck pain	1 (0.2)	4 (1.1)
Osteoarthritis	5 (1.1)	1 (0.3)
Pain in extremity	7 (1.6)	6 (1.7)
Nervous system disorders		
Dizziness	7 (1.6)	8 (2.2)
Headache	18 (4.1)	14 (3.9)
Paresthesia	4 (0.9)	4 (1.1)
Psychiatric disorders		
Anxiety	3 (0.7)	4 (1.1)
Insomnia	4 (0.9)	6 (1.7)

Table 1 – Adverse events $\geq 1\%$ in any treatment group (regardless of causality) reported in patients treated with JANUVIA[®] 100 mg or placebo in pooled 18 and 24-week placebo-controlled, double-blind clinical trials of JANUVIA[®] as monotherapy

Body system/Organ class Adverse event	Number of patients (%)	
	Sitagliptin 100 mg n=443	Placebo n=363
Respiratory, thoracic and mediastinal disorders		
Cough	8 (1.8)	10 (2.8)
Vascular disorders		
Hypertension	8 (1.8)	7 (1.9)

In a 24-week study which compared sitagliptin and metformin, adverse events, reported regardless of causality assessment, in $\geq 1\%$ of patients are shown in Table 2.

Table 2 – Adverse events $\geq 1\%$ in any treatment group (regardless of causality) reported in patients in 24-week active-controlled, double-blind clinical trial of JANUVIA[®] as monotherapy

Body system/Organ class Adverse event	Number of patients (%)	
	Sitagliptin 100 mg n=528	Metformin n=522
Gastrointestinal disorders		
Abdominal pain	4 (0.8)	6 (1.1)
Abdominal pain upper	5 (0.9)	12 (2.3)
Constipation	9 (1.7)	5 (1.0)
Diarrhea	19 (3.6)	57 (10.9)
Dyspepsia	1 (0.2)	7 (1.3)
Gastritis	6 (1.1)	11 (2.1)
Nausea	6 (1.1)	16 (3.1)
Vomiting	2 (0.4)	7 (1.3)
General disorders and administration site conditions		
Fatigue	6 (1.1)	6 (1.1)
Infections and infestations		
Bronchitis	4 (0.8)	7 (1.3)
Influenza	12 (2.3)	11 (2.1)
Nasopharyngitis	10 (1.9)	17 (3.3)
Upper respiratory tract infection	5 (0.9)	11 (2.1)
Urinary tract infection	3 (0.6)	13 (2.5)
Metabolism and nutrition disorders		
Hypoglycemia	9 (1.7)	18 (3.4)
Musculoskeletal and connective tissue disorders		
Back pain	9 (1.7)	9 (1.7)
Pain in extremity	7 (1.3)	2 (0.4)

Table 2 – Adverse events $\geq 1\%$ in any treatment group (regardless of causality) reported in patients in 24-week active-controlled, double-blind clinical trial of JANUVIA[®] as monotherapy

Body system/Organ class Adverse event	Number of patients (%)	
	Sitagliptin 100 mg n=528	Metformin n=522
Nervous system disorders		
Dizziness	9 (1.7)	5 (1.0)
Headache	17 (3.2)	17 (3.3)
Respiratory, thoracic and mediastinal disorders		
Cough	1 (0.2)	8 (1.5)
Vascular disorders		
Hypertension	12 (2.3)	4 (0.8)

In two monotherapy studies, diarrhea was the only drug-related adverse reaction reported by the investigator that occurred with an incidence $\geq 1\%$ in patients receiving JANUVIA[®] 100 mg (1.1%) and greater than in patients receiving placebo (0.3%).

Combination Therapy – Sitagliptin add-on to metformin:

In a 24-week placebo-controlled clinical study of patients receiving sitagliptin (100 mg daily) as add-on combination therapy with metformin the incidence of adverse events reported regardless of causality assessment, in $\geq 1\%$ of patients are shown in Table 3.

Table 3 – Adverse events $\geq 1\%$ in any treatment group (regardless of causality) reported in patients in a 24-week placebo-controlled, double-blind clinical trial of JANUVIA[®] in add-on combination use with metformin

Body system/Organ class Adverse event	Number of patients (%)	
	Sitagliptin 100 mg + Metformin n=464	Placebo + Metformin n=237
Ear and labyrinth disorders		
Vertigo	5 (1.1)	4 (1.7)
Eye disorders		
Vision blurred	1 (0.2)	3 (1.3)
Gastrointestinal disorders		
Abdominal pain	2 (0.4)	6 (2.5)
Abdominal pain upper	6 (1.3)	2 (0.8)
Constipation	5 (1.1)	1 (0.4)
Diarrhea	11 (2.4)	6 (2.5)
Nausea	6 (1.3)	2 (0.8)
Vomiting	5 (1.1)	2 (0.8)
General disorders and administration site conditions		
Fatigue	2 (0.4)	4 (1.7)
Edema peripheral	4 (0.9)	3 (1.3)
Infections and infestations		
Bronchitis	12 (2.6)	6 (2.5)
Bronchitis acute	2 (0.4)	3 (1.3)
Gastroenteritis	4 (0.9)	5 (2.1)
Influenza	19 (4.1)	12 (5.1)

Table 3 – Adverse events $\geq 1\%$ in any treatment group (regardless of causality) reported in patients in a 24-week placebo-controlled, double-blind clinical trial of JANUVIA[®] in add-on combination use with metformin

Body system/Organ class Adverse event	Number of patients (%)	
	Sitagliptin 100 mg + Metformin n=464	Placebo + Metformin n=237
Nasopharyngitis	19 (4.1)	7 (3.0)
Pharyngitis	6 (1.3)	1 (0.4)
Infections and infestations		
Pneumonia	5 (1.1)	0 (0.0)
Sinusitis	7 (1.5)	2 (0.8)
Tooth infection	5 (1.1)	2 (0.8)
Upper respiratory tract infection	34 (7.3)	22 (9.3)
Urinary tract infection	9 (1.9)	2 (0.8)
Injury, poisoning and procedural complications		
Contusion	5 (1.1)	1 (0.4)
Investigations		
Blood glucose increased	3 (0.6)	6 (2.5)
Metabolism and nutrition disorders		
Hyperglycemia	2 (0.4)	7 (3.0)
Hypoglycemia	6 (1.3)	5 (2.1)
Musculoskeletal and connective tissue disorders		
Arthralgia	14 (3.0)	1 (0.4)
Back pain	15 (3.2)	6 (2.5)
Muscle spasm	1 (0.2)	3 (1.3)
Myalgia	1 (0.2)	3 (1.3)
Pain in extremity	5 (1.1)	4 (1.7)
Shoulder pain	3 (0.6)	3 (1.3)
Nervous system disorders		
Dizziness	7 (1.5)	2 (0.8)
Headache	12 (2.6)	7 (3.0)
Sciatica	1 (0.2)	3 (1.3)
Sinus headache	0 (0.0)	3 (1.3)
Psychiatric disorders		
Insomnia	5 (1.1)	3 (1.3)
Renal and urinary disorders		
Nephrolithiasis	3 (0.6)	3 (1.3)
Respiratory, thoracic and mediastinal disorders		
Cough	14 (3.0)	4 (1.7)
Vascular disorders		
Hypertension	7 (1.5)	6 (2.5)

In a combination therapy study with metformin, nausea was the only drug-related adverse reaction reported by the investigator that occurred with an incidence $\geq 1\%$ in patients receiving JANUVIA[®] (1.1%) and greater than in patients receiving placebo (0.4%).

In pooled studies of up to one year duration which compared sitagliptin added to metformin or a sulfonylurea agent (glipizide) added to metformin, adverse events, reported regardless of causality assessment, in $\geq 1\%$ of patients are shown in Table 4.

Table 4 – Adverse events $\geq 1\%$ in any treatment group (regardless of causality) reported in patients from double-blind clinical trials of JANUVIA® in add-on combination use with metformin in studies up to one year compared to a sulfonylurea agent (glipizide)

Body system/Organ class Adverse event	Number of patients (%)	
	Sitagliptin 100 mg + Metformin n=979	Glipizide + Metformin n=748
Gastrointestinal disorders		
Abdominal pain	10 (1.0)	6 (0.8)
Abdominal pain upper	13 (1.3)	7 (0.9)
Constipation	17 (1.7)	13 (1.7)
Diarrhea	42 (4.3)	36 (4.8)
Dyspepsia	14 (1.4)	12 (1.6)
Nausea	19 (1.9)	16 (2.1)
Toothache	2 (0.2)	13 (1.7)
Vomiting	11 (1.1)	9 (1.2)
General disorders and administration site conditions		
Fatigue	20 (2.0)	8 (1.1)
Non-cardiac chest pain	10 (1.0)	6 (0.8)
Edema peripheral	16 (1.6)	14 (1.9)
Infections and infestations		
Bronchitis	27 (2.8)	22 (2.9)
Cellulitis	7 (0.7)	10 (1.3)
Gastroenteritis	19 (1.9)	13 (1.7)
Gastroenteritis viral	8 (0.8)	9 (1.2)
Herpes zoster	4 (0.4)	8 (1.1)
Influenza	35 (3.6)	32 (4.3)
Nasopharyngitis	75 (7.7)	49 (6.6)
Sinusitis	20 (2.0)	12 (1.6)
Upper respiratory tract infection	78 (8.0)	70 (9.4)
Urinary tract infection	41 (4.2)	21 (2.8)
Investigations		
Blood glucose decreased	5 (0.5)	16 (2.1)
Blood glucose increased	13 (1.3)	5 (0.7)
Weight increased	1 (0.1)	8 (1.1)
Metabolism and nutrition disorders		
Hyperglycemia	10 (1.0)	6 (0.8)
Hypoglycemia	32 (3.3)	217 (29.0)
Musculoskeletal and connective tissue disorders		
Arthralgia	34 (3.5)	29 (3.9)
Back pain	39 (4.0)	32 (4.3)
Muscle spasms	9 (0.9)	8 (1.1)
Neck pain	4 (0.4)	8 (1.1)
Osteoarthritis	18 (1.8)	5 (0.7)
Pain in extremity	23 (2.3)	9 (1.2)
Shoulder pain	7 (0.7)	14 (1.9)

Table 4 – Adverse events $\geq 1\%$ in any treatment group (regardless of causality) reported in patients from double-blind clinical trials of JANUVIA[®] in add-on combination use with metformin in studies up to one year compared to a sulfonylurea agent (glipizide)

Body system/Organ class Adverse event	Number of patients (%)	
	Sitagliptin 100 mg + Metformin n=979	Glipizide + Metformin n=748
Nervous system disorders		
Dizziness	26 (2.7)	14 (1.9)
Headache	34 (3.5)	31 (4.1)
Hypoaesthesia	3 (0.3)	11 (1.5)
Psychiatric disorders		
Anxiety	13 (1.3)	7 (0.9)
Depression	10 (1.0)	7 (0.9)
Insomnia	12 (1.2)	11 (1.5)
Reproductive system and breast disorders		
Erectile dysfunction	6 (0.6)	8 (1.1)
Respiratory, thoracic and mediastinal disorders		
Cough	19 (1.9)	23 (3.1)
Pharyngolaryngeal pain	10 (1.0)	9 (1.2)
Sinus congestion	5 (0.5)	8 (1.1)
Eczema	4 (0.4)	12 (1.6)
Vascular disorders		
Hypertension	33 (3.4)	29 (3.9)

Combination Therapy: Sitagliptin add-on to Metformin and a Sulfonylurea

In a 24-week placebo-controlled study of JANUVIA[®] 100 mg in combination with metformin and glimepiride (JANUVIA[®], N=116; placebo, N=113), the incidence of adverse events, reported regardless of causality assessment, in $\geq 1\%$ of patients are shown in Table 5. The overall incidence of adverse events with JANUVIA[®] was higher than with placebo, in part related to higher incidence of hypoglycemia (see Table 5).

Table 5 – Adverse events $\geq 1\%$ in any treatment group (regardless of causality) reported in patients in a 24-week placebo-controlled, double-blind clinical trial of JANUVIA[®] in add-on combination use with metformin and a sulfonylurea agent (glimepiride)

Body system/Organ class Adverse event	Number of patients (%)	
	Sitagliptin 100 mg + Metformin + Glimepiride n=116	Placebo + Metformin + Glimepiride n=113
Ear and Labyrinth Disorders		
Vertigo	2 (1.7)	0 (0.0)
Eye Disorders		
Diabetic retinopathy	0 (0.0)	2 (1.8)
Vision blurred	0 (0.0)	2 (1.8)
Gastrointestinal disorders		
Abdominal pain upper	2 (1.7)	2 (1.8)
Constipation	4 (3.4)	0 (0.0)
Diarrhea	1 (0.9)	4 (3.5)
Dyspepsia	3 (2.6)	2 (1.8)
Gastritis	0 (0.0)	4 (3.5)

Table 5 – Adverse events $\geq 1\%$ in any treatment group (regardless of causality) reported in patients in a 24-week placebo-controlled, double-blind clinical trial of JANUVIA® in add-on combination use with metformin and a sulfonylurea agent (glimepiride)

Body system/Organ class Adverse event	Number of patients (%)	
	Sitagliptin 100 mg + Metformin + Glimepiride n=116	Placebo + Metformin + Glimepiride n=113
Toothache	2 (1.7)	2 (1.8)
Vomiting	2 (1.7)	1 (0.9)
General disorders and administration site conditions		
Fatigue	0 (0.0)	3 (2.7)
Non-Cardiac chest pain	2 (1.7)	1 (0.9)
Pyrexia	0 (0.0)	2 (1.8)
Hepatobiliary disorders		
Cholelithiasis	0 (0.0)	2 (1.8)
Infections and infestations		
Bronchitis	2 (1.7)	2 (1.8)
Gastroenteritis	3 (2.6)	0 (0.0)
Gastroenteritis viral	2 (1.7)	2 (1.8)
Influenza	3 (2.6)	2 (1.8)
Nasopharyngitis	7 (6.0)	9 (8.0)
Pharyngitis	1 (0.9)	3 (2.7)
Pneumonia	3 (2.6)	0 (0.0)
Rhinitis	2 (1.7)	0 (0.0)
Sinusitis	1 (0.9)	2 (1.8)
Tooth abscess	2 (1.7)	1 (0.9)
Upper respiratory tract infection	8 (6.9)	9 (8.0)
Urinary tract infection	2 (1.7)	1 (0.9)
Injury, poisoning and procedural complications		
Fall	0 (0.0)	3 (2.7)
Polytraumatism	1 (0.9)	2 (1.8)
Investigations		
Blood glucose decreased	0 (0.0)	2 (1.8)
Metabolism and nutrition disorders		
Hypoglycemia	19 (16.4)	1 (0.9)
Musculoskeletal and connective tissue disorders		
Arthralgia	5 (4.3)	1 (0.9)
Back pain	1 (0.9)	2 (1.8)
Muscle spasms	2 (1.7)	1 (0.9)
Osteoarthritis	2 (1.7)	0 (0.0)
Pain in extremity	4 (3.4)	1 (0.9)
Shoulder pain	0 (0.0)	2 (1.8)
Nervous system disorders		
Dizziness	3 (2.6)	1 (0.9)
Headache	8 (6.9)	3 (2.7)
Hypoaesthesia	2 (1.7)	0 (0.0)
Somnolence	0 (0.0)	2 (1.8)

Table 5 – Adverse events $\geq 1\%$ in any treatment group (regardless of causality) reported in patients in a 24-week placebo-controlled, double-blind clinical trial of JANUVIA[®] in add-on combination use with metformin and a sulfonylurea agent (glimepiride)

Body system/Organ class Adverse event	Number of patients (%)	
	Sitagliptin 100 mg + Metformin + Glimepiride n=116	Placebo + Metformin + Glimepiride n=113
Respiratory, thoracic and mediastinal disorders		
Asthma	2 (1.7)	1 (0.9)
Skin and subcutaneous tissue disorders		
Pruritus	2 (1.7)	1 (0.9)
Rash	2 (1.7)	1 (0.9)
Vascular disorders		
Hypertension	2 (1.7)	0 (0.0)

In a combination therapy study with metformin and a sulfonylurea, hypoglycemia (JANUVIA[®] 13.8%; placebo 0.9%) and constipation (JANUVIA[®] 1.7%; placebo 0.0%) were the only drug-related adverse reactions reported by the investigator that occurred with an incidence $\geq 1\%$ in patients receiving JANUVIA[®] and metformin and a sulfonylurea and greater than in patients receiving placebo and metformin and a sulfonylurea.

Combination Therapy: Add-on with Insulin (with or without metformin)

In a 24-week placebo-controlled study of JANUVIA[®] 100 mg in combination with stable-dose insulin (with or without metformin) (JANUVIA[®], N=322; placebo, N=319), the incidence of adverse reactions, reported regardless of causality assessment, in $\geq 1\%$ of patients are shown in Table 6. The overall incidence of adverse events with JANUVIA[®] was higher than with placebo, in part related to higher incidence of hypoglycemia (see Table 6)

Table 6 – Adverse events $\geq 1\%$ in any treatment group (regardless of causality) reported in patients in a 24-week placebo-controlled, double-blind clinical trial of JANUVIA[®] in add-on combination use with stable dose insulin (with or without metformin)

Body system/Organ class Adverse event	Number of patients (%)	
	Sitagliptin 100 mg + Insulin (+/- Metformin) n=322	Placebo + Insulin (+/- Metformin) n=319
Gastrointestinal Disorders		
Constipation	6 (1.9)	1 (0.3)
Diarrhea	6 (1.9)	5 (1.6)
Nausea	4 (1.2)	5 (1.6)
Vomiting	5 (1.6)	2 (0.6)
Infections and infestations		
Bronchitis	6 (1.9)	5 (1.6)
Gastroenteritis	3 (0.9)	5 (1.6)
Influenza	13 (4.0)	12 (3.8)
Nasopharyngitis	10 (3.1)	8 (2.5)
Sinusitis	4 (1.2)	4 (1.3)
Upper respiratory tract infection	10 (3.1)	11 (3.4)
Urinary tract infection	9 (2.8)	6 (1.9)

Table 6 – Adverse events $\geq 1\%$ in any treatment group (regardless of causality) reported in patients in a 24-week placebo-controlled, double-blind clinical trial of JANUVIA[®] in add-on combination use with stable dose insulin (with or without metformin)

Body system/Organ class Adverse event	Number of patients (%)	
	Sitagliptin 100 mg + Insulin (+/- Metformin) n=322	Placebo + Insulin (+/- Metformin) n=319
Investigations		
Alanine aminotransferase increased	4 (1.2)	1 (0.3)
Creatinine renal clearance decreased	5 (1.6)	0 (0.0)
Metabolism and nutrition disorders		
Hyperglycemia	5 (1.6)	2 (0.6)
Hypoglycemia	50 (15.5)	25 (7.8)
Musculoskeletal and connective tissue disorders		
Arthralgia	4 (1.2)	6 (1.9)
Back pain	6 (1.9)	2 (0.6)
Muscle spasms	3 (0.9)	5 (1.6)
Pain in extremity	6 (1.9)	3 (0.9)
Nervous system disorders		
Dizziness	5 (1.6)	3 (0.9)
Headache	9 (2.8)	3 (0.9)
Respiratory, thoracic and mediastinal disorders		
Cough	5 (1.6)	3 (0.9)

In a combination therapy study with stable dose insulin (with or without metformin), hypoglycemia (JANUVIA[®], 9.6%; placebo, 5.3%), influenza (JANUVIA[®], 1.2%, placebo, 0.3%), and headache (JANUVIA[®], 1.2%, placebo, 0.0%) were the only drug-related adverse reactions reported by the investigator that occurred with an incidence $\geq 1\%$ in patients receiving JANUVIA[®] and greater than in patients receiving placebo.

Combination Therapy: Sitagliptin add-on to Pioglitazone (with or without Metformin)

In a 24-week placebo-controlled clinical study of patients receiving sitagliptin (100 mg daily) as add-on combination therapy with pioglitazone, the incidence of adverse events reported regardless of causality assessment, in $\geq 1\%$ of patients are shown in Table 7.

Table 7 – Adverse events $\geq 1\%$ in any treatment group (regardless of causality) reported in patients in a 24-week placebo-controlled, double-blind clinical trial of JANUVIA[®] in add-on combination with pioglitazone

Body system/Organ class Adverse event	Number of patients (%)	
	Sitagliptin 100 mg + Pioglitazone n=175	Placebo + Pioglitazone n=178
Ear and Labyrinth Disorders		
Vertigo	0 (0.0)	3 (1.7)
Eye Disorders		
Cataract	0 (0.0)	3 (1.7)
Vision Blurred	2 (1.1)	1 (0.6)

Table 7 – Adverse events $\geq 1\%$ in any treatment group (regardless of causality) reported in patients in a 24-week placebo-controlled, double-blind clinical trial of JANUVIA® in add-on combination with pioglitazone

Body system/Organ class Adverse event	Number of patients (%)	
	Sitagliptin 100 mg + Pioglitazone n=175	Placebo + Pioglitazone n=178
Gastrointestinal disorders		
Abdominal pain	2 (1.1)	0 (0.0)
Abdominal pain lower	2 (1.1)	0 (0.0)
Abdominal pain upper	2 (1.1)	0 (0.0)
Constipation	2 (1.1)	2 (1.1)
Diarrhea	3 (1.7)	2 (1.1)
Dyspepsia	2 (1.1)	1 (0.6)
Flatulence	2 (1.1)	0 (0.0)
Nausea	2 (1.1)	0 (0.0)
General disorders and administration site conditions		
Chest pain	2 (1.1)	0 (0.0)
Fatigue	1 (0.6)	3 (1.7)
Feeling abnormal	2 (1.1)	0 (0.0)
Oedema	2 (1.1)	1 (0.6)
Oedema peripheral	7 (4.0)	5 (2.8)
Hepatobiliary Disorders		
Cholelithiasis	0 (0.0)	2 (1.1)
Infections and infestations		
Bronchitis	3 (1.7)	1 (0.6)
Cellulitis	2 (1.1)	1 (0.6)
Influenza	6 (3.4)	5 (2.8)
Nasopharyngitis	7 (4.0)	7 (3.9)
Pharyngitis	2 (1.1)	2 (1.1)
Pneumonia	0 (0.0)	3 (1.7)
Pyoderma	2 (1.1)	0 (0.0)
Sinusitis	2 (1.1)	2 (1.1)
Tinea Pedis	2 (1.1)	0 (0.0)
Upper respiratory tract infection	11 (6.3)	6 (3.4)
Urinary tract infection	1 (0.6)	2 (1.1)
Viral infection	2 (1.1)	1 (0.6)
Injury, poisoning and procedural complications		
Joint sprain	2 (1.1)	2 (1.1)
Investigations		
Blood glucose increased	1 (0.6)	2 (1.1)
Weight increased	5 (2.9)	5 (2.8)
Metabolism and nutrition disorders		
Hypoglycemia	2 (1.1)	0 (0.0)
Musculoskeletal and connective tissue disorders		
Arthralgia	5 (2.9)	4 (2.2)
Back pain	3 (1.7)	5 (2.8)
Musculoskeletal stiffness	2 (1.1)	0 (0.0)
Myalgia	0 (0.0)	2 (1.1)
Neck pain	0 (0.0)	2 (1.1)

Table 7 – Adverse events $\geq 1\%$ in any treatment group (regardless of causality) reported in patients in a 24-week placebo-controlled, double-blind clinical trial of JANUVIA[®] in add-on combination with pioglitazone

Body system/Organ class Adverse event	Number of patients (%)	
	Sitagliptin 100 mg + Pioglitazone n=175	Placebo + Pioglitazone n=178
Osteoarthritis	3 (1.7)	3 (1.7)
Pain in extremity	4 (2.3)	3 (1.7)
Tendonitis	0 (0.0)	2 (1.1)
Nervous system disorders		
Dizziness	3 (1.7)	2 (1.1)
Headache	9 (5.1)	7 (3.9)
Psychiatric disorders		
Anxiety	1 (0.6)	2 (1.1)
Depression	4 (2.3)	2 (1.1)
Libido decreased	2 (1.1)	0 (0.0)
Respiratory, thoracic and mediastinal disorders		
Cough	3 (1.7)	3 (1.7)
Skin and subcutaneous tissue disorders		
Dermatitis allergic	0 (0.0)	2 (1.1)

In a combination therapy study with pioglitazone, hypoglycemia (JANUVIA[®], 1.1%, placebo, 0.0%), flatulence (JANUVIA[®], 1.1%, placebo, 0.0%), weight increase (JANUVIA[®], 2.3%, placebo, 1.7%), and headache (JANUVIA[®], 1.7%, placebo, 1.1%) were the only drug-related adverse reactions reported by the investigator that occurred with an incidence $\geq 1\%$ in patients receiving JANUVIA[®] and greater than in patients receiving placebo.

In a 26-week placebo-controlled clinical study of patients receiving sitagliptin (100 mg daily) as add-on combination therapy with metformin and pioglitazone, the incidence of adverse events reported regardless of causality assessment, in $\geq 1\%$ of patients are shown in Table 8.

Table 8 – Adverse events $\geq 1\%$ in any treatment group (regardless of causality) reported in patients in a 26-week placebo-controlled, double-blind clinical trial of JANUVIA[®] in add-on combination use with metformin and pioglitazone

Body system/Organ class Adverse event	Number of patients (%)	
	Sitagliptin 100 mg + Metformin + Pioglitazone n=157	Placebo + Metformin + Pioglitazone n=156
Ear and Labyrinth Disorders		
Cerumen impaction	2 (1.3)	1 (0.6)
Eye Disorders		
Conjunctivitis	3 (1.9)	1 (0.6)
Ocular hyperaemia	0 (0.0)	2 (1.3)
Gastrointestinal disorders		
Abdominal pain upper	1 (0.6)	2 (1.3)
Constipation	2 (1.3)	1 (0.6)
Dental Caries	2 (1.3)	1 (0.6)
Diarrhea	3 (1.9)	4 (2.6)

Table 8 – Adverse events $\geq 1\%$ in any treatment group (regardless of causality) reported in patients in a 26-week placebo-controlled, double-blind clinical trial of JANUVIA® in add-on combination use with metformin and pioglitazone

Body system/Organ class Adverse event	Number of patients (%)	
	Sitagliptin 100 mg + Metformin + Pioglitazone n=157	Placebo + Metformin + Pioglitazone n=156
Dyspepsia	1 (0.6)	2 (1.3)
Gastritis	0 (0.0)	2 (1.3)
Toothache	2 (1.3)	0 (0.0)
Vomiting	2 (1.3)	0 (0.0)
General disorders and administration site conditions		
Fatigue	0 (0.0)	2 (1.3)
Oedema peripheral	3 (1.9)	7 (4.5)
Infections and infestations		
Bronchitis	3 (1.9)	1 (0.6)
Cellulitis	2 (1.3)	0 (0.0)
Gastroenteritis	2 (1.3)	0 (0.0)
Gastroenteritis viral	2 (1.3)	0 (0.0)
Herpes zoster	2 (1.3)	0 (0.0)
Influenza	2 (1.3)	3 (1.9)
Nasopharyngitis	5 (3.2)	5 (3.2)
Tooth abscess	0 (0.0)	2 (1.3)
Upper respiratory tract infection	13 (8.3)	14 (9.0)
Urinary tract infection	5 (3.2)	6 (3.8)
Injury, poisoning and procedural complications		
Muscle strain	2 (1.3)	0 (0.0)
Investigations		
Blood creatine phosphokinase increased	1 (0.6)	3 (1.9)
Glomerular filtration rate decreased	2 (1.3)	0 (0.0)
Lymphocyte count increased	2 (1.3)	1 (0.6)
Neutrophil count decreased	2 (1.3)	1 (0.6)
Metabolism and nutrition disorders		
Hyperglycemia	2 (1.3)	2 (1.3)
Hypoglycemia	10 (6.4)	7 (4.5)
Musculoskeletal and connective tissue disorders		
Arthralgia	2 (1.3)	3 (1.9)
Back pain	7 (4.5)	4 (2.6)
Muscle spasms	2 (1.3)	0 (0.0)
Musculoskeletal pain	3 (1.9)	4 (2.6)
Pain in extremity	5 (3.2)	2 (1.3)
Nervous system disorders		
Headache	1 (0.6)	2 (1.3)
Psychiatric disorders		
Depression	4 (2.5)	1 (0.6)
Stress	2 (1.3)	0 (0.0)
Respiratory, thoracic and mediastinal disorders		

Table 8 – Adverse events $\geq 1\%$ in any treatment group (regardless of causality) reported in patients in a 26-week placebo-controlled, double-blind clinical trial of JANUVIA[®] in add-on combination use with metformin and pioglitazone

Body system/Organ class Adverse event	Number of patients (%)	
	Sitagliptin 100 mg + Metformin + Pioglitazone n=157	Placebo + Metformin + Pioglitazone n=156
Cough	2 (1.3)	2 (1.3)
Oropharyngeal pain	2 (1.3)	0 (0.0)
Rhinitis allergic	2 (1.3)	0 (0.0)

In a combination therapy study with pioglitazone and metformin, hypoglycemia (JANUVIA[®], 3.2%; placebo, 1.9%) was the only drug-related adverse reaction reported by the investigator that occurred with an incidence $\geq 1\%$ in patients receiving JANUVIA[®] and greater than in patients receiving placebo.

Less Common Clinical Trial Adverse Drug Reactions $>0.1\%$ and $<1\%$ (Drug-Related and Greater than Placebo in Pooled Monotherapy and in Individual Placebo-Controlled Studies)

Blood and Lymphatic System Disorders: anemia

Cardiac Disorders: bundle branch block, palpitations

Eye Disorders: vision blurred

Gastrointestinal Disorders: abdominal discomfort, abdominal pain upper, abdominal tenderness, constipation, diarrhea, dry mouth, dyspepsia, flatulence, reflux esophagitis disease, frequent bowel movements, gastroesophageal reflux disease, irritable bowel syndrome, retching, salivary hypersecretion

General Disorders and Administration Site Conditions: asthenia, chest discomfort, face edema, fatigue, feeling abnormal, hunger, irritability, malaise, peripheral edema, edema, pain, pyrexia, thirst, xerosis

Hepatobiliary Disorders: hepatic steatosis

Infections and Infestations: gastric ulcer helicobacter, genital abscess, helicobacter gastritis, localized infection, oropharyngeal candidiasis, sinusitis, upper respiratory tract infection, urinary tract infection

Investigations: alanine aminotransferase increased, aspartate aminotransferase increased, blood glucose decreased, blood glucose increased, blood pressure decreased, blood pressure increased, creatinine renal clearance decreased, glomerular filtration rate decreased, white blood cell count increased

Metabolism and Nutrition Disorders: decreased appetite, hypoglycemia

Musculoskeletal and Connective Tissue Disorders: muscle fatigue, muscle tightness

Nervous System Disorders: coordination abnormal, dizziness, headache, migraine, neuropathy peripheral, parosmia, somnolence

Psychiatric Disorders: anxiety, depression, insomnia, libido decreased

Renal and Urinary Disorders: renal disorders

Reproductive System and Breast Disorders: balanoposthitis, dysmenorrhea, erectile dysfunction

Respiratory, Thoracic and Mediastinal Disorders: cough

Skin and Subcutaneous Tissue Disorders: angioneurotic oedema, dermatitis acneiform, dry skin, erythema, exanthem, hyperhidrosis, leukocytoclastic vasculitis, nail disorder, prurigo, pruritus generalized, rash, rash macular, rosacea, urticaria

Vascular Disorders: orthostatic hypotension

Atrial fibrillation/atrial flutter: In a pooled analysis of randomized clinical trials, the pooled terms atrial fibrillation/atrial flutter were observed at an incidence rate of 0.45 events per 100 patient-years in the sitagliptin-exposed group compared to 0.28 events per 100 patient-years in the non-exposed group.

TECOS Cardiovascular Safety Study:

For details pertaining to study design and patient population, see CLINICAL TRIALS, TECOS Cardiovascular Safety Study.

The incidence of adjudication-confirmed pancreatitis events was higher in the JANUVIA[®] group (0.3%) compared to the placebo group (0.2%). The JANUVIA[®] group experienced a greater number of severe cases of pancreatitis including two confirmed deaths due to pancreatitis, compared to none in the placebo group.

Among patients who were using insulin and/or a sulfonylurea at baseline, the incidence of severe hypoglycemia was 2.7% in patients treated with JANUVIA[®] and 2.5% in patients treated with placebo; among patients who were not using insulin and/or a sulfonylurea at baseline, the incidence of severe hypoglycemia was 1.0% in patients treated with JANUVIA[®] and 0.7% in placebo-treated patients.

Abnormal Hematologic and Clinical Chemistry Findings

The incidence of laboratory adverse experiences was similar in patients treated with JANUVIA[®] 100 mg compared to patients treated with placebo. In most clinical studies, a slight decrease in alkaline phosphatase and small increases in uric acid and white blood cell count (due to an increase in neutrophils) were observed. In active comparator studies versus metformin or versus a sulfonylurea agent (glipizide) similar changes were seen in alkaline phosphatase and uric acid.

Mean Change from Baseline (Standard Error)				
Study	Treatment Group	Alkaline Phosphatase (IU/L)	Uric Acid (mg/dL)	WBC (cell/microl)
Placebo-controlled (monotherapy) ¹	Sitagliptin	-5.3 (0.5)	0.26 (0.04)	320.2 (71.7)
	Placebo	-0.8 (0.5)	-0.05 (0.05)	58.6 (80.0)
Active-controlled (monotherapy) ²	Sitagliptin	-3.9 (0.5)	-0.0 (0.0)	220.4 (77.7)
	Metformin	-4.7 (0.5)	0.1 (0.0)	184.7 (66.6)
Placebo-controlled (add-on to metformin) ³	Sitagliptin	-3.1 (0.4)	0.17 (0.04)	346.0 (64.3)
	Placebo	-1.3 (0.7)	0.05 (0.06)	142.4 (98.8)
Active-controlled (add-on to metformin) ⁴	Sitagliptin	-5.7 (0.5)	0.21 (0.05)	207.8 (67.4)
	Glipizide	-3.4 (0.5)	0.20 (0.05)	86.0 (62.5)

¹ pooled data from studies 3 and 4; see CLINICAL TRIALS, Table 10

² study 5; see CLINICAL TRIALS, Table 10

³ study 1; see CLINICAL TRIALS, Table 10

⁴ study 2; see CLINICAL TRIALS, Table 10

In a combination therapy study with stable dose insulin (with or without metformin), a greater proportion of patients was observed to have a decrease in hemoglobin ≥ 1.5 g/dL in the sitagliptin group (6.0%) compared with the placebo group (2.1%). No adverse experiences of anemia or hemoglobin decreased were reported in the sitagliptin group.

Post-Marketing Adverse Drug Reactions

The following adverse reactions have been identified during post-marketing use of JANUVIA[®] as monotherapy and/or in combination with other antihyperglycemic agents. Because these reactions are reported voluntarily from a population of uncertain size, it is generally not possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Immune system disorders: Hypersensitivity reactions including anaphylaxis, angioedema, rash, urticaria, cutaneous vasculitis, and exfoliative skin conditions, including Stevens-Johnson syndrome (see CONTRAINDICATIONS and WARNINGS AND PRECAUTIONS, Hypersensitivity Reactions)

Renal and urinary disorders: worsening renal function, including acute renal failure (sometimes requiring dialysis) (see WARNINGS AND PRECAUTIONS)

Gastrointestinal disorders: acute pancreatitis, including fatal and non-fatal hemorrhagic and necrotizing pancreatitis (see WARNINGS AND PRECAUTIONS); vomiting

Musculoskeletal and connective tissue disorders: arthralgia; myalgia; pain in extremity; back pain

Skin and subcutaneous tissue disorders: pruritus, bullous pemphigoid (see WARNINGS AND PRECAUTIONS, Skin)

DRUG INTERACTIONS

Overview

Sitagliptin is not an inhibitor of CYP isozymes CYP3A4, 2C8, 2C9, 2D6, 1A2, 2C19 or 2B6, and is not an inducer of CYP3A4. Sitagliptin is a p-glycoprotein substrate, but does not inhibit

p-glycoprotein mediated transport of digoxin. Based on these results, sitagliptin is considered unlikely to cause interactions with other drugs that utilize these pathways.

Sitagliptin is not extensively bound to plasma proteins. Therefore, the propensity of sitagliptin to be involved in clinically meaningful drug-drug interactions mediated by plasma protein binding displacement is very low.

Drug-Drug Interactions

In clinical studies, as described below, sitagliptin did not meaningfully alter the pharmacokinetics of metformin, glyburide, simvastatin, rosiglitazone, warfarin, or oral contraceptives, providing *in vivo* evidence of a low propensity for causing drug interactions with substrates of CYP3A4, CYP2C8, CYP2C9, and organic cationic transporter (OCT).

Metformin: Co-administration of multiple twice-daily doses of sitagliptin with metformin, an OCT substrate, did not meaningfully alter the pharmacokinetics of metformin or JANUVIA[®] in patients with type 2 diabetes. Therefore, sitagliptin is not an inhibitor of OCT-mediated transport.

Sulfonylureas: Single-dose pharmacokinetics of glyburide, a CYP2C9 substrate, were not meaningfully altered in subjects receiving multiple doses of sitagliptin. Clinically meaningful interactions would not be expected with other sulfonylureas (e.g., glipizide, tolbutamide, and glimepiride) which, like glyburide, are primarily eliminated by CYP2C9. The effect of sulfonylureas on the pharmacokinetics of sitagliptin was not assessed.

Simvastatin: Single-dose pharmacokinetics of simvastatin, a CYP3A4 substrate, were not meaningfully altered in subjects receiving multiple daily doses of sitagliptin. Therefore, sitagliptin is not an inhibitor of CYP3A4-mediated metabolism.

Thiazolidinediones: Single-dose pharmacokinetics of rosiglitazone were not meaningfully altered in subjects receiving multiple daily doses of sitagliptin. Therefore, sitagliptin is not an inhibitor of CYP2C8-mediated metabolism. Clinically meaningful interactions with pioglitazone are not expected because pioglitazone predominantly undergoes CYP2C8- or CYP3A4-mediated metabolism. The effect of thiazolidinediones on the pharmacokinetics of sitagliptin was not assessed.

Warfarin: Multiple daily doses of sitagliptin did not meaningfully alter the pharmacokinetics, as assessed by measurement of S(-) or R(+) warfarin enantiomers, or pharmacodynamics (as assessed by measurement of prothrombin INR) of a single dose of warfarin. Since S(-) warfarin is primarily metabolized by CYP2C9, these data also support the conclusion that sitagliptin is not a CYP2C9 inhibitor.

Oral Contraceptives: Co-administration with sitagliptin did not meaningfully alter the steady-state pharmacokinetics of norethindrone or ethinyl estradiol.

Digoxin: Sitagliptin had a minimal effect on the pharmacokinetics of digoxin. Following administration of 0.25 mg digoxin concomitantly with 100 mg of JANUVIA[®] daily for 10 days, the plasma AUC of digoxin was increased by 11%, and the plasma C_{max} by 18%. These increases are not considered likely to be clinically meaningful. No dosage adjustment of digoxin or JANUVIA[®] is recommended.

Cyclosporine: A study was conducted to assess the effect of cyclosporine, a potent inhibitor of p-glycoprotein, on the pharmacokinetics of sitagliptin. Coadministration of a single 100-mg oral dose of JANUVIA[®] and a single 600-mg oral dose of cyclosporine increased the AUC and C_{max} of sitagliptin by approximately 29% and 68%, respectively. These modest changes in sitagliptin pharmacokinetics were not considered to be clinically meaningful. The renal clearance of sitagliptin was also not meaningfully altered. Therefore, meaningful interactions would not be expected with other p-glycoprotein inhibitors. No dosage adjustment for JANUVIA[®] is recommended when co-administered with cyclosporine or other p-glycoprotein inhibitors (e.g., ketoconazole).

Drug-Food Interactions

There are no known interactions with food.

Drug-Herb Interactions

Interactions with herbal products have not been established.

Drug-Laboratory Interactions

Interactions with laboratory tests have not been established.

Drug-Lifestyle Interactions

No studies of the effects of JANUVIA[®] on the ability to drive and use machines have been performed. However, JANUVIA[®] is not expected to affect the ability to drive and use machines. When JANUVIA[®] is used in combination with metformin a sulfonylurea or in combination with insulin (with or without metformin) patients should be advised to take precautions to avoid hypoglycemia while driving or using machinery.

DOSAGE AND ADMINISTRATION

Dosing Considerations

JANUVIA[®] can be taken with or without food.

Recommended Dose and Dosage Adjustment

The recommended dose of JANUVIA[®] is 100 mg once daily as monotherapy or as combination therapy with metformin, with metformin and a sulfonylurea, with insulin (with or without metformin), or with pioglitazone (with or without metformin).

When JANUVIA[®] is used in combination with metformin and a sulfonylurea or with insulin (with or without metformin); a lower dose of sulfonylurea or insulin may be considered to reduce the risk of hypoglycemia (see WARNINGS AND PRECAUTIONS, Hypoglycemia).

Patients with Renal Insufficiency: JANUVIA[®] is renally excreted. To achieve plasma concentrations of JANUVIA[®] similar to those in patients with normal renal function, dosage recommendations are as follows:

For patients with mild renal insufficiency (creatinine clearance [CrCl] ≥ 50 mL/min), no dosage adjustment for JANUVIA[®] is required.

For patients with moderate renal insufficiency (CrCl ≥ 30 to < 50 mL/min), the dose of JANUVIA[®] is 50 mg once daily.

For patients with severe renal insufficiency (CrCl < 30 mL/min) or with end-stage renal disease (ESRD) requiring hemodialysis or peritoneal dialysis, the dose of JANUVIA[®] is 25 mg once daily. JANUVIA[®] may be administered without regard to the timing of dialysis.

Because there is a dosage adjustment based upon renal function, assessment of renal function is recommended prior to initiation of JANUVIA[®] and periodically thereafter.

Creatinine clearance can be estimated from serum creatinine using the Cockcroft-Gault formula (see WARNINGS AND PRECAUTIONS, ACTION AND CLINICAL PHARMACOLOGY and CLINICAL TRIALS, Patients with Renal Insufficiency).

When considering the use of sitagliptin in combination with another anti-diabetic product, its conditions for use in patients with renal impairment should be followed.

Patients with Hepatic Insufficiency: Use of sitagliptin in patients with severe hepatic insufficiency is not recommended.

Geriatrics: No dosage adjustment is necessary for geriatric patients.

Pediatrics: There are no data available on the use of JANUVIA[®] in patients younger than 18 years of age. Therefore, use of JANUVIA[®] in pediatric patients is not recommended.

Missed Dose

If a dose of JANUVIA[®] is missed, it should be taken as soon as the patient remembers. A double dose of JANUVIA[®] should not be taken on the same day.

OVERDOSAGE

For management of a suspected drug overdose, contact your regional Poison Control Centre.

During controlled clinical trials in healthy subjects, single doses of up to 800 mg JANUVIA[®] were generally well tolerated. Minimal increases in QTc, not considered to be clinically relevant, were observed in one study at a dose of 800 mg JANUVIA[®] (see ACTION AND CLINICAL PHARMACOLOGY). There is no experience with doses above 800 mg in clinical trials.

In the event of an overdose, it is reasonable to employ the usual supportive measures, e.g., remove unabsorbed material from the gastrointestinal tract, employ clinical monitoring (including obtaining an electrocardiogram), and institute supportive therapy if required.

Sitagliptin is modestly dialyzable. In clinical studies, approximately 13.5% of the dose was removed over a 3- to 4-hour hemodialysis session. Prolonged hemodialysis may be considered if clinically appropriate. It is not known if sitagliptin is dialyzable by peritoneal dialysis.

ACTION AND CLINICAL PHARMACOLOGY

Mechanism of Action

JANUVIA[®] is an orally-active, potent, and highly selective inhibitor of the dipeptidyl peptidase 4 (DPP-4) enzyme for the treatment of type 2 diabetes. The DPP-4 inhibitors are a novel class of agents that act as incretin enhancers.

Incretin hormones, including glucagon-like peptide-1 (GLP-1) and glucose-dependent insulinotropic polypeptide (GIP), are released by the intestine throughout the day, and levels are increased in response to a meal. The incretins are part of an endogenous system involved in the physiologic regulation of glucose homeostasis. When blood glucose concentrations are normal or elevated, GLP-1 and GIP increase insulin synthesis and release from pancreatic beta cells by intracellular signaling pathways involving cyclic AMP. Progressive beta-cell failure is a feature characterizing the pathogenesis of type 2 diabetes. Treatment with GLP-1 or with DPP-4 inhibitors in animal models of type 2 diabetes has been demonstrated to improve beta cell responsiveness to glucose and stimulate insulin biosynthesis and release. With higher insulin levels, tissue glucose uptake is enhanced.

In addition, GLP-1 lowers glucagon secretion from pancreatic alpha cells. Decreased glucagon concentrations, along with higher insulin levels, lead to reduced hepatic glucose production, resulting in a decrease in blood glucose levels. When blood glucose concentrations are low, stimulation of insulin release and suppression of glucagon secretion by GLP-1 are not observed. GLP-1 does not impair the normal glucagon response to hypoglycemia.

The activity of GLP-1 and GIP is limited by the DPP-4 enzyme, which rapidly hydrolyzes the incretin hormones to produce inactive products. Sitagliptin prevents the hydrolysis of incretin hormones by DPP-4, thereby increasing plasma concentrations of the active forms of GLP-1 and GIP. By enhancing active incretin levels, sitagliptin increases insulin release and decreases glucagon levels in a glucose-dependent manner.


In patients with type 2 diabetes with hyperglycemia, these changes in insulin and glucagon levels lead to lower hemoglobin A_{1c} (HbA_{1c}) and lower fasting and postprandial glucose concentrations. Sitagliptin demonstrates selectivity for DPP-4, and does not inhibit the DPP-8 or DPP-9 activity *in vitro* at concentrations approximating those from therapeutic doses. Inhibition of DPP-8 or DPP-9, but not DPP-4, is associated with toxicity in preclinical animal models and alteration of immune function *in vitro*.

Pharmacodynamics

In patients with type 2 diabetes, administration of single oral doses of JANUVIA[®] leads to inhibition of DPP-4 enzyme activity for a 24-hour period, resulting in a 2- to 3-fold increase in circulating levels of active GLP-1 and GIP, increased plasma levels of insulin and C-peptide, decreased glucagon concentrations, reduced fasting glucose, and reduced glucose excursion following an oral glucose load or a meal.

In a study of patients with type 2 diabetes inadequately controlled on metformin monotherapy (N=26), glucose levels monitored throughout the day were significantly lower ($p < 0.001$) in patients who received sitagliptin 100 mg per day (50 mg twice daily) in combination with metformin compared with patients who received placebo with metformin (see Figure 1).

Figure 1 – 24-hour plasma glucose profile after 4-week treatment with Sitagliptin 50 mg BID with metformin or placebo with metformin



In studies with healthy subjects, JANUVIA[®] did not lower blood glucose or cause hypoglycemia, suggesting that the insulinotropic and glucagon suppressive actions of the drug are glucose dependent.

Cardiac Electrophysiology: In a randomized, placebo-controlled crossover study, 79 healthy subjects were administered a single oral dose of JANUVIA[®] 100 mg, JANUVIA[®] 800 mg (8 times the recommended dose), and placebo. At the recommended dose of 100 mg, there was no effect on the QTc interval obtained at the peak plasma concentration, or at any other time during the study.

Following the 800-mg dose, the maximum increase in the placebo-corrected mean change in QTc from baseline at 3 hours postdose was 8.0 msec (90% CI 5.5, 10.6). At the 800-mg dose, peak sitagliptin plasma concentrations were approximately 11 times higher than the peak concentrations following a 100-mg dose.

In patients with type 2 diabetes administered JANUVIA[®] 100 mg (N=81) or JANUVIA[®] 200 mg (N=63) daily, there were no meaningful changes in QTc interval based on ECG data obtained at the time of expected peak plasma concentration.

Pharmacokinetics

Table 9 – Summary of sitagliptin’s pharmacokinetic parameters in healthy volunteers

	C_{max} nM	t_½ (h)	AUC_{0-∞} μM•hr	Renal Clearance mL/min	Volume of distribution (L)*
Single oral dose (100 mg) mean	950	12.4	8.52	350	198

* Volume of distribution at steady state following an I.V. dose.

The pharmacokinetics of sitagliptin have been extensively characterized in healthy subjects and patients with type 2 diabetes. After oral administration of a 100-mg dose to healthy subjects, sitagliptin was rapidly absorbed, with peak plasma concentrations (median T_{max}) occurring 1 to 4 hours post-dose. Plasma AUC of sitagliptin increased in a dose-proportional manner. Following a single oral 100-mg dose to healthy volunteers, mean plasma AUC of sitagliptin was 8.52 μM•hr, C_{max} was 950 nM, and apparent terminal half-life (t_{1/2}) was 12.4 hours. Plasma AUC of sitagliptin increased approximately 14% following 100-mg doses at steady-state compared to the first dose. The intra-subject and inter-subject coefficients of variation for sitagliptin AUC were small (5.8% and 15.1%). The pharmacokinetics of sitagliptin were generally similar in healthy subjects and in patients with type 2 diabetes.

Absorption: The absolute bioavailability of sitagliptin is approximately 87%. Since coadministration of a high-fat meal with JANUVIA[®] had no effect on the pharmacokinetics, JANUVIA[®] may be administered with or without food.

Distribution: The mean volume of distribution at steady state following a single 100-mg intravenous dose of sitagliptin to healthy subjects is approximately 198 liters. The fraction of sitagliptin reversibly bound to plasma proteins is low (38%).

Metabolism: Sitagliptin is primarily eliminated unchanged in urine, and metabolism is a minor pathway. Approximately 79% of sitagliptin is excreted unchanged in the urine.

Following a [¹⁴C] sitagliptin oral dose, approximately 16% of the radioactivity was excreted as metabolites of sitagliptin. Six metabolites were detected at trace levels and are not expected to contribute to the plasma DPP-4 inhibitory activity of sitagliptin. *In vitro* studies indicated that the primary enzyme responsible for the limited metabolism of sitagliptin was CYP3A4, with contribution from CYP2C8.

Excretion: Following administration of an oral [¹⁴C] sitagliptin dose to healthy subjects, approximately 100% of the administered radioactivity was eliminated in feces (13%) or urine (87%) within one week of dosing. The apparent terminal t_{1/2} following a 100-mg oral dose of sitagliptin was approximately 12.4 hours and renal clearance was approximately 350 mL/min.

Elimination of sitagliptin occurs primarily via renal excretion and involves active tubular secretion. Sitagliptin is a substrate for human organic anion transporter-3 (hOAT-3), which may be involved in the renal elimination of sitagliptin. The clinical relevance of hOAT-3 in sitagliptin transport has not been established. Sitagliptin is also a substrate of p-glycoprotein, which may also be involved in mediating the renal elimination of sitagliptin. However, cyclosporine, a p-glycoprotein inhibitor, did not reduce the renal clearance of sitagliptin.

Special Populations and Conditions

Pediatrics: No studies with JANUVIA[®] have been performed in pediatric patients.

Geriatrics: No dosage adjustment is required based on age. Age did not have a clinically meaningful impact on the pharmacokinetics of sitagliptin based on a population pharmacokinetic analysis of Phase I and Phase II data. Elderly subjects (65 to 80 years) had approximately 19% higher plasma concentrations of sitagliptin compared to younger subjects.

Gender: No dosage adjustment is necessary based on gender. Gender had no clinically meaningful effect on the pharmacokinetics of sitagliptin based on a composite analysis of Phase I pharmacokinetic data and on a population pharmacokinetic analysis of Phase I and Phase II data.

Race: No dosage adjustment is necessary based on race. Race had no clinically meaningful effect on the pharmacokinetics of sitagliptin based on a composite analysis of Phase I pharmacokinetic data and on a population pharmacokinetic analysis of Phase I and Phase II data, including subjects of white, Hispanic, black and Asian racial groups.

Hepatic Insufficiency: In patients with moderate hepatic insufficiency (Child-Pugh score 7 to 9), mean AUC and C_{max} of sitagliptin increased approximately 21% (90% CI: 1%, 46%) and 13% (90% CI: -9%, 42%), respectively, compared to healthy matched controls following administration of a single 100-mg dose of JANUVIA[®].

Renal Insufficiency: A single-dose, open-label study was conducted to evaluate the pharmacokinetics of a reduced dose of JANUVIA[®] (50 mg) in patients with varying degrees of chronic renal insufficiency compared to normal healthy control subjects. The study included patients with renal insufficiency classified on the basis of creatinine clearance as mild (50 to <80 mL/min), moderate (30 to <50 mL/min), and severe (<30 mL/min), as well as patients with end-stage renal disease (ESRD) on hemodialysis. Compared to normal healthy control subjects, an approximate 1.1 to 1.6-fold increase in plasma AUC of sitagliptin was observed in patients with mild renal insufficiency, which is not a clinically meaningful increase to require dosage adjustment. An approximately 2-fold increase in the plasma AUC of sitagliptin was observed in patients with renal insufficiency classified on the basis of creatinine clearance as moderate (30 to <50 mL/min) and an approximately 4-fold increase was observed in patients with severe (<30 mL/min) renal insufficiency and in patients with ESRD on hemodialysis, as compared to normal healthy control subjects.

Sitagliptin was modestly removed by hemodialysis (13.5% over a 3- to 4-hour hemodialysis session starting 4 hours postdose). To achieve plasma concentrations of sitagliptin similar to those in patients with normal renal function, lower dosages are recommended in patients with moderate and severe renal insufficiency, as well as in ESRD patients requiring dialysis. The dose of JANUVIA[®] should be reduced to 50 mg once daily in patients with moderate renal insufficiency or 25 mg in patients with severe renal insufficiency and in patients with ESRD on hemodialysis (see WARNINGS AND PRECAUTIONS, DOSAGE AND ADMINISTRATION and CLINICAL TRIALS, Patients with Renal Insufficiency).

STORAGE AND STABILITY

Store at room temperature (15°C to 30°C).

DOSAGE FORMS, COMPOSITION AND PACKAGING

Tablets JANUVIA[®], 25 mg, are pink, round, film-coated tablets with “221” on one side. They are supplied in bottles of 30.

Tablets JANUVIA[®], 50 mg, are light beige, round, film-coated tablets with “112” on one side. They are supplied in bottles of 30.

Tablets JANUVIA[®], 100 mg, are beige, round, film-coated tablets with “277” on one side. They are supplied in bottles of 30 and 100.

Each film-coated tablet of JANUVIA[®] contains 32.13, 64.25, or 128.5 mg of sitagliptin phosphate monohydrate, which is equivalent to 25, 50 or 100 mg, respectively, of free base.

Each film-coated tablet of JANUVIA[®] contains the following inactive ingredients: microcrystalline cellulose, anhydrous dibasic calcium phosphate (calcium hydrogen phosphate, anhydrous), croscarmellose sodium, magnesium stearate, and sodium stearyl fumarate. In addition, the film coating contains the following inactive ingredients: polyvinyl alcohol, polyethylene glycol (macrogol), talc, titanium dioxide, red iron oxide, and yellow iron oxide.

PART II: SCIENTIFIC INFORMATION

PHARMACEUTICAL INFORMATION

Drug Substance


Common name: sitagliptin phosphate monohydrate

Chemical name: 7-[(3*R*)-3-amino-1-oxo-4-(2,4,5-trifluorophenyl)butyl]-5,6,7,8-tetrahydro-3-(trifluoromethyl)-1,2,4-triazolo[4,3-*a*]pyrazine phosphate (1:1) monohydrate.

Molecular formula: $C_{16}H_{15}F_6N_5O \cdot H_3PO_4 \cdot H_2O$

Molecular mass: 523.32

Structural formula:



Physicochemical properties:

Sitagliptin phosphate monohydrate is a white to off-white, crystalline, non-hygroscopic powder. It is soluble in water and N, N-dimethyl formamide; slightly soluble in methanol; very slightly soluble in ethanol, acetone, and acetonitrile; and insoluble in isopropanol and isopropyl acetate.

CLINICAL TRIALS

Study demographics and trial design

Table 10 – Summary of patient demographics for clinical trials in specific indication

Study #	Trial design	Dosage, route of administration and duration	Study subjects (n=number)	Mean age (Range)	Gender
Study 1 (P020)	Multicentre, randomized, double-blind, placebo-controlled	JANUVIA® 100 mg once daily + ≥1500 mg/day Metformin or Placebo + ≥1500 mg/day Metformin Oral 24-week	701	54.5 years (19–78)	Male: 400 Female: 301
Study 2 (P024)	Multicentre, randomized, double-blind, with an active comparator	JANUVIA® 100 mg/day + ≥1500 mg/day Metformin or Glipizide 5–20 mg/day + ≥1500 mg/day Metformin Oral 52-week	1172	Male 23–79 Female 22–78	Male: 694 Female: 478
Study 3 (P023)	Multicentre, randomized, double-blind, placebo-controlled	Placebo or 100 mg or 200 mg JANUVIA® once daily Oral 18-week	521	55.1 years (27–76)	Male: 283 Female: 238
Study 4 (P021)	Multicentre, randomized, double-blind, placebo-controlled	Placebo or 100 mg or 200 mg JANUVIA® once daily Oral 24-week	741	54.2 years (18–75)	Male: 383 Female: 358
Study 5 (P049)	Multicentre, randomized double-blind active-controlled	JANUVIA® 100 mg/day or Metformin 500 mg/day and titrated to 1500 to 2000 mg/day Oral 24 Weeks	1050	56.0 years (20–78)	Male: 484 Female: 566

Table 10 – Summary of patient demographics for clinical trials in specific indication

Study #	Trial design	Dosage, route of administration and duration	Study subjects (n=number)	Mean age (Range)	Gender
Study 6 (P035)	Multicentre, randomized, double-blind, placebo-controlled	JANUVIA® 100 mg/day + Glimepiride ≥4 mg/day in combination with Metformin ≥1500 mg/day or Placebo + Glimepiride ≥4 mg/day in combination with Metformin ≥1500 mg/day Oral 24-week	229	58.0 years (33–75)	Male: 120 Female: 109
Study 7 (P051)	Multicentre, randomized, double-blind placebo-controlled	JANUVIA® 100 mg/day + Stable dose insulin (alone or in combination with metformin ≥1500 mg/day) or Placebo + stable dose insulin (alone or in combination with metformin ≥1500 mg/day) 24-week	641	57.8 years (25–82)	Male: 326 Female: 315
Study 8 (P028)	Multicentre, randomized, double-blind, placebo- and active- controlled	JANUVIA® 25 or 50 mg/day or Placebo for 12 weeks followed by glipizide 2.5 to 20 mg/day for 42 weeks Oral 54 weeks	91	67.9 years (41–92)	Male: 47 Female: 44

Table 10 – Summary of patient demographics for clinical trials in specific indication

Study #	Trial design	Dosage, route of administration and duration	Study subjects (n=number)	Mean age (Range)	Gender
Study 9 (P063)	Multicentre, randomized, double-blind, active-controlled	JANUVIA® 25 or 50 mg/day or Glipizide 2.5 to 20 mg/day Oral 54 weeks	423	64.2 years (33–89)	Male: 253 Female: 170
Study 10 (P073)	Multicentre, randomized, double-blind, active-controlled	JANUVIA® 25 mg/day or Glipizide 2.5 to 20 mg/day Oral 54 weeks	129	59.5 years (38–83)	Male: 77 Female: 52
Study 11 (P019)	Multicentre, randomized, double-blind, placebo-controlled	JANUVIA® 100 mg/day + 30 or 45 mg/day Pioglitazone or Placebo + 30 or 45 mg/day Pioglitazone Oral 24-week	353	56.2 years (24–87)	Male: 196 Female: 157
Study 12 (P128)	Multicentre, randomized, double-blind, placebo-controlled	JANUVIA® 100 mg/day + Pioglitazone ≥30 mg/day + Metformin ≥1500 mg/day or Placebo + Pioglitazone ≥30 mg/day + Metformin ≥1500 mg/day Oral 26-week	313	56.1 (22–78)	Male: 195 Female: 118

Study results

Monotherapy – Placebo-Controlled Study:

A total of 1262 patients with type 2 diabetes participated in two double-blind, placebo-controlled studies, one of 18-week and another of 24-week duration, to evaluate the efficacy and safety of JANUVIA[®] monotherapy. Patients with inadequate glycemic control (HbA_{1c} 7% to 10%) were randomized to receive a 100-mg or 200-mg dose of JANUVIA[®] or placebo once daily.

Treatment with JANUVIA[®] at 100 mg daily provided significant improvements in HbA_{1c}, FPG, and 2-hour PPG compared to placebo (Table 11). The improvement in HbA_{1c} compared to placebo was not affected by gender, age, race, prior antihyperglycemic therapy or baseline BMI. Patients with a shorter length of time since diagnosis of diabetes (<3 years) or with higher baseline HbA_{1c} had greater reductions in HbA_{1c}. Overall, the 200-mg daily dose did not provide greater glycemic efficacy than the 100-mg daily dose. The effect of JANUVIA[®] on lipid endpoints was similar to placebo. Body weight did not increase from the baseline with JANUVIA[®] (mean weight loss of 0.6 kg in the 18-week study and 0.2 kg in the 24-week study). Patients on placebo lost more weight (mean weight loss 0.7 kg in the 18-week study and 1.1 kg in the 24-week study) than patients on JANUVIA[®].

Table 11 – Glycemic parameters in 18- and 24-week placebo-controlled studies of JANUVIA[®] in patients with type 2 diabetes[†]

	18-Week Study		24-Week Study	
	JANUVIA [®] 100 mg	Placebo	JANUVIA [®] 100 mg	Placebo
HbA_{1c} (%)	N=193	N=103	N=229	N=244
Baseline (mean)	8.0	8.1	8.0	8.0
Change from Baseline (adjusted mean [‡])	-0.5	0.1	-0.6	0.2
Difference from Placebo (adjusted mean [‡])	-0.6 [§]		-0.8 [§]	
Patients (%) achieving HbA _{1c} <7%	69 [§] (35.8%)	16 (15.5%)	93 [§] (40.6%)	41 (16.8%)
FPG (mmol/L)	N=201	N=107	N=234	N=247
Baseline (mean)	10.0	10.2	9.5	9.8
Change from baseline (adjusted mean [‡])	-0.7	0.4	-0.7	0.3
Difference from Placebo (adjusted mean [‡])	-1.1 [§]		-1.0 [§]	
2-hour PPG (mmol/L)	%	%	N=201	N=204
Baseline (mean)			14.3	15.0
Change from baseline (adjusted mean [‡])			-2.7	-0.1
Difference from Placebo (adjusted mean [‡])			-2.6 [§]	

[†] All Patients Treated Population (an intention-to-treat analysis).

[‡] Least squares means adjusted for prior antihyperglycemic therapy status and baseline value.

[§] p<0.001 compared to placebo.

[%] Data not available.

Monotherapy – Active-Controlled Study with Metformin:

The efficacy of JANUVIA[®] compared to that of metformin was evaluated in a 24-week, double-blind, metformin-controlled trial in patients with type 2 diabetes and inadequate glycemic control on diet and exercise and who were not on antihyperglycemic therapy (off therapy for at least 4 months). In this study, patients with an HbA_{1c} of 6.5% to 9.0% were randomized to receive either JANUVIA[®] 100 mg daily (N=528) or metformin (N=522) for 24 weeks. Patients receiving metformin were given an initial dosage of 500 mg/day and then titrated to a dose of 1500 to 2000 mg/day over a period of up to 5 weeks based on tolerability. The mean dose of metformin after the titration period was approximately 1900 mg/day. Glycemic endpoints measured included HbA_{1c} and fasting glucose.

Both treatments resulted in a statistically significant improvement in glycemic control from baseline. At 24 weeks, the reduction from baseline in HbA_{1c} was -0.43% for JANUVIA[®] 100 mg daily and -0.57% for metformin in the per protocol population analysis.

The reduction in FPG was -0.64 mmol/L for JANUVIA[®] and -1.08 mmol/L for metformin. Body weight decreased from baseline in both treatment groups (JANUVIA[®], -0.6 kg; metformin -1.9 kg).

Sitagliptin in Combination with Metformin – Placebo-Controlled Study:

A total of 701 patients with type 2 diabetes participated in a 24-week, randomized, double-blind, placebo-controlled study designed to assess the efficacy of JANUVIA[®] in combination with metformin. All patients were started on metformin monotherapy and the dose increased to at least 1500 mg per day. Patients were randomized to the addition of either 100 mg of JANUVIA[®] or placebo, administered once daily. Patients with congestive heart failure requiring pharmacological treatment were excluded from this study.

Glycemic parameters and body weight at final visit (24-week study) for JANUVIA[®] in combination with metformin are shown in Table 12.

Table 12 – Glycemic parameters and body weight at final visit (24-week study) for JANUVIA® in combination with metformin†

	JANUVIA® 100 mg + Metformin	Placebo + Metformin
HbA_{1c} (%)	N=453	N=224
Baseline (mean)	8.0	8.0
Change from baseline (adjusted mean‡)	-0.7	-0.0
Difference from placebo + metformin (adjusted mean‡)	-0.7§	
Patients (%) achieving HbA _{1c} <7%	213 (47.0%)	41 (18.3%)
FPG (mmol/L)	N=454	N=226
Baseline (mean)	9.4	9.6
Change from baseline (adjusted mean‡)	-0.9	0.5
Difference from placebo + metformin (adjusted mean‡)	-1.4§	
2-hour PPG (mmol/L)	N=387	N=182
Baseline (mean)	15.3	15.1
Change from baseline (adjusted mean‡)	-3.4	-0.6
Difference from placebo + metformin (adjusted mean‡)	-2.8§	
Body Weight (kg) %	N=399	N=169
Baseline (mean)	86.9	87.6
Change from baseline (adjusted mean‡)	-0.7	-0.6
Difference from placebo + metformin (adjusted mean‡)	-0.1¶	

† All Patients Treated Population (an intention-to-treat analysis).

‡ Least squares means adjusted for prior antihyperglycemic therapy and baseline value.

§ p<0.001 compared to placebo + metformin.

% All Patients as Treated (APaT) population, excluding patients given glycemic rescue therapy.

¶ Not statistically significant (p≥0.05) compared to placebo + metformin.

Sitagliptin in Combination with Metformin – Active-Controlled (Sulfonylurea Agent) Study:

Long-term maintenance of effect was evaluated in a 52-week, double-blind, glipizide-controlled trial in patients with type 2 diabetes and inadequate glycemic control on metformin monotherapy at ≥ 1500 mg/day. In this study, patients were randomized to the addition of either JANUVIA[®] 100 mg daily (N=588) or glipizide (N=584) for 52 weeks. Patients receiving glipizide were given an initial dosage of 5 mg/day and then electively titrated by the investigator to a target FPG of 6.1 mmol/L, without significant hypoglycemia, over the next 18 weeks. A maximum dosage of 20 mg/day was allowed to optimize glycemic control. Thereafter, the glipizide dose was to have been kept constant. The mean daily dose of glipizide after the titration period was 10.3 mg.

Both treatments resulted in a statistically significant improvement in glycemic control from baseline. After 52 weeks, the reduction from baseline in HbA_{1c} was 0.67% for JANUVIA[®] 100 mg daily and 0.67% for glipizide, confirming the non-inferiority of JANUVIA[®] compared to glipizide. The reduction in FPG was 0.6 mmol/L for JANUVIA[®] and 0.4 mmol/L for glipizide. In this study, the proinsulin to insulin ratio, a marker of efficiency of insulin synthesis and release, improved with JANUVIA[®] relative to glipizide. The incidence of hypoglycemia in the JANUVIA[®] group (4.9%) was significantly lower than that in the glipizide group (32.0%). Patients treated with JANUVIA[®] exhibited a significant mean decrease from baseline in body weight compared to a significant weight gain in patients administered glipizide (-1.5 kg vs. +1.1 kg).

Add-on Combination Therapy with Metformin plus Glimepiride:

In a 24-week, randomized, double-blind, placebo-controlled study designed to assess the efficacy of sitagliptin 100 mg once daily (N=116) compared to placebo (N=113), 229 patients were on the combination of glimepiride (≥ 4 mg per day) and metformin (≥ 1500 mg per day); the results of the glycemic endpoints, including HbA_{1c} and FPG, are described below.

The combination of sitagliptin, glimepiride, and metformin provided significant reduction from baseline in HbA_{1c} and FPG compared to placebo (see Table 13). Mean reductions from baseline in HbA_{1c} compared with placebo were generally greater for patients with higher baseline HbA_{1c} values. Patients treated with sitagliptin, had a modest increase in body weight (0.4 kg) compared to those given placebo who had a significant decrease in body weight (-0.7 kg).

Table 13 – Glycemic Parameters and Body Weight at Final Visit (24-Week Study) for JANUVIA® in Add-on Combination Therapy with Metformin plus Glimepiride[†]

	JANUVIA® 100 mg + Metformin + Glimepiride	Placebo + Metformin + Glimepiride
HbA_{1c} (%)	N=115	N=105
Baseline (mean)	8.27	8.28
Change from baseline (adjusted mean [‡])	-0.59	0.30
Difference from placebo (adjusted mean [‡])	-0.89 [§]	
Patients (%) achieving HbA _{1c} <7%	26 (22.6)	1 (1.0)
FPG (mmol/L)	N=115	N=109
Baseline (mean)	9.95	9.93
Change from baseline (adjusted mean [‡])	-0.43	0.72
Difference from placebo (adjusted mean [‡])	-1.15 [§]	
Body Weight (kg)[¶]	N=102	N=74
Baseline (mean)	86.5	84.6
Change from baseline (adjusted mean [‡])	0.4	-0.7
Difference from placebo (adjusted mean [‡])	1.1 ^{††}	

[†] All Patients Treated Population (an intention-to-treat analysis).

[‡] Least squares means adjusted for prior antihyperglycemic therapy status and baseline value.

[§] p<0.001 compared to placebo.

[¶] All Patients as Treated (APaT) population, excluding patients given glycemic rescue therapy.

^{††} p=0.007 compared to placebo.

Add-on Combination Therapy with Insulin (with or without Metformin):

A total of 641 patients with type 2 diabetes participated in a 24-week, randomized, double-blind, placebo-controlled study designed to assess the efficacy of JANUVIA® as add-on combination therapy with a stable dose of insulin (with or without metformin). Patients with an HbA_{1c} of 7.5% to 11.0% while on a stable regimen of pre-mixed, long-acting, or intermediate-acting insulin with or without metformin (≥1500 mg per day) were randomized to the addition of either 100 mg of JANUVIA® or placebo, administered once daily. Patients using pre-meal short-acting or rapid-acting insulins that were not components of a pre-mixed insulin formulation, or that were administered via insulin pumps, were not included in this study. Glycemic endpoints measured included HbA_{1c}, fasting glucose, and 2-hour post-prandial glucose.

In combination with insulin (including patients taking and not taking metformin), JANUVIA® provided significant improvements in HbA_{1c}, FPG, and 2-hour PPG compared to placebo (Table 14). There was no significant difference between JANUVIA® and placebo in body weight change.

Table 14 – Glycemic Parameters and Body Weight at Final Visit (24-Week Study) for JANUVIA® as Add-on Combination Therapy with a Stable dose of Insulin (with or without Metformin[†])

	JANUVIA® 100 mg + Insulin (+/- Metformin)	Placebo + Insulin (+/- Metformin)
HbA_{1c} (%)	N=305	N=312
Baseline (mean)	8.7	8.6
Change from baseline (adjusted mean [‡])	-0.6	-0.0
Difference from placebo (adjusted mean ^{‡,§})	-0.6 [%]	
Patients (%) achieving HbA _{1c} <7%	39 (12.8)	16 (5.1)
FPG (mmol/L)	N=310	N=313
Baseline (mean)	9.7	9.8
Change from baseline (adjusted mean [‡])	-1.0	-0.2
Difference from placebo (adjusted mean [‡])	-0.8	
2-hour PPG (mmol/L)	N=240	N=257
Baseline (mean)	16.0	16.1
Change from baseline (adjusted mean [‡])	-1.7	0.3
Difference from placebo (adjusted mean [‡])	-2.0 [%]	
Body Weight (kg)[¶]	N=266	N=266
Baseline (mean)	86.6	87.4
Change from baseline (adjusted mean [‡])	0.1	0.1
Difference from placebo (adjusted mean [‡])	0.0 [#]	

[†] All Patients Treated Population (an intention-to-treat analysis).

[‡] Least squares means adjusted for metformin use at Visit 1 (yes/no), insulin use at Visit 1 (pre-mixed vs. non-pre-mixed [intermediate- or long-acting]), and baseline value.

[§] Treatment by stratum interaction was not significant (p>0.10) for metformin stratum and for insulin stratum.

[%] p<0.001 compared to placebo.

[¶] All Patients as Treated (APaT) population, excluding data following glycemic rescue therapy.

[#] Not statistically significant (p≥0.05) compared to placebo.

Table 15 – Glycemic Parameters at Final Visit (24-Week Study) for JANUVIA® as Add-on Combination Therapy with a Stable Dose of Insulin (with or without Metformin[†])

	JANUVIA® 100 mg + Insulin	Placebo + Insulin	JANUVIA® 100 mg + Insulin + Metformin	Placebo + Insulin + Metformin
HbA_{1c} (%)	N=82	N=83	N=223	N=229
Baseline (mean)	8.7	8.8	8.7	8.6
Change from baseline (adjusted mean [‡])	-0.6	0.1	-0.7	-0.1
Difference from placebo (adjusted mean [‡])	-0.7 [%]		-0.5 [%]	
Patients (%) achieving HbA _{1c} <7%	7 (8.5)	4 (4.8)	32 (14.3)	12 (5.2)
FPG (mmol/L)	N=85	N=84	N=225	N=229
Baseline (mean)	10.1	10.5	9.6	9.8
Change from baseline (adjusted mean [‡])	-0.7	-0.3	-1.2	-0.2
Difference from placebo (adjusted mean [‡])	-0.3 [§]		-1.0 [%]	
2-hour PPG (mmol/L)	N=58	N=68	N=182	N=189
Baseline (mean)	17.9	18.0	15.6	15.6
Change from baseline (adjusted mean [‡])	-1.0	0.3	-2.2	0.1
Difference from placebo (adjusted mean [‡])	-1.3 [#]		-2.2 [%]	

[†] All Patients Treated Population (an intention-to-treat analysis).

[‡] Least squares means adjusted for insulin use at Visit 1 (pre-mixed vs. non-pre-mixed [intermediate- or long-acting]), and baseline value.

[%] p<0.001 compared to placebo.

[§] Not statistically significant (p≥0.05) compared to placebo.

[#] p=0.037 compared to placebo.

Add-on Combination Therapy with Pioglitazone:

Combination Therapy with Pioglitazone: A total of 353 patients with type 2 diabetes participated in a 24-week, randomized, double-blind, placebo-controlled study designed to assess the efficacy of JANUVIA® in combination with pioglitazone. All patients were on pioglitazone monotherapy at a dose of 30–45 mg per day. Patients were randomized to the addition of either 100 mg of JANUVIA® or placebo, administered once daily. Glycemic endpoints measured included HbA_{1c} and fasting glucose.

In combination with pioglitazone, JANUVIA® provided significant improvements in HbA_{1c} and FPG compared to placebo with pioglitazone (Table 16). The improvement in HbA_{1c} was not affected by baseline HbA_{1c}, prior anti-hyperglycemic therapy, gender, age, race, baseline BMI, length of time since diagnosis of diabetes, presence of metabolic syndrome (according to NCEP criteria), or standard indices of insulin resistance (HOMA-IR) or insulin secretion (HOMA-β). Treatment with JANUVIA® did not significantly increase body weight from baseline compared to placebo.

Table 16 – Glycemic parameters and body weight at final visit (24-week study) for JANUVIA® in combination with pioglitazone†

	JANUVIA® 100 mg + Pioglitazone	Placebo + Pioglitazone
HbA_{1c} (%)	N=163	N=174
Baseline (mean)	8.1	8.0
Change from baseline (adjusted mean‡)	-0.9	-0.2
Difference from pioglitazone alone (adjusted mean‡)	-0.7§	
Patients (%) achieving HbA _{1c} <7%	74 (45.4%)	40 (23.0%)
FPG (mmol/L)	N=163	N=174
Baseline (mean)	9.3	9.2
Change from baseline (adjusted mean‡)	-0.9	0.1
Difference from pioglitazone alone (adjusted mean‡)	-1.0§	
Body Weight (kg)¶	N=133	N=136
Baseline (mean)	90.0	85.6
Change from baseline (adjusted mean‡)	1.8	1.5
Difference from pioglitazone alone (adjusted mean‡)	0.2¶	

† All Patients Treated Population (an intention-to-treat analysis).

‡ Least squares means adjusted for prior antihyperglycemic therapy status and baseline value.

§ p<0.001 compared to placebo + pioglitazone.

¶ All Patients as Treated (APaT) population, excluding patients given glycemic rescue therapy.

¶ Not statistically significant (p≥0.05) compared to placebo + pioglitazone.

Combination Therapy with Pioglitazone and Metformin:

Sitagliptin Add-on Therapy in Patients Inadequately Controlled on Pioglitazone and Metformin

A total of 313 patients with type 2 diabetes participated in a 26-week, randomized, double-blind, placebo-controlled study designed to assess the efficacy of sitagliptin in combination with pioglitazone and metformin. Patients with inadequate glycemic control on a stable regimen of pioglitazone (30 or 45 mg per day) and metformin (≥1500 mg per day) were randomized to the addition of either 100 mg of sitagliptin or placebo, administered once daily.

In combination with pioglitazone and metformin, sitagliptin provided significant improvements in HbA_{1c}, FPG, and 2-hour PPG compared to placebo with pioglitazone and metformin (Table 17). Lipid effects were generally neutral. The difference between sitagliptin and placebo in body weight change was not significant.

Table 17 – Glycemic Parameters and Body Weight at Final Visit (26-Week Study) for JANUVIA® as Add-on Combination Therapy with Pioglitazone and Metformin†

	Sitagliptin 100 mg + Pioglitazone 30 or 45 mg + Metformin	Placebo + Pioglitazone 30 or 45 mg + Metformin
HbA_{1c} (%)	N=152	N=153
Baseline (mean)	8.8	8.6
Change from baseline (adjusted mean‡)	-1.2	-0.4
Difference from placebo (adjusted mean‡)	-0.7§	
Patients (%) achieving HbA _{1c} <7%	38 (25.0)	15 (9.8)
FPG (mmol/L)	N=155	N=153
Baseline (mean)	10.0	9.6
Change from baseline (adjusted mean‡)	-1.1	-0.2
Difference from placebo (adjusted mean‡)	-1.0§	
2-hour PPG (mmol/L)	N=141	N=135
Baseline (mean)	15.3	14.7
Change from baseline (adjusted mean‡)	-3.0	-0.8
Difference from placebo (adjusted mean‡)	-2.2§	
Body Weight (kg)%	N=146	N=128
Baseline (mean)	81.4	82.0
Change from baseline (adjusted mean‡)	1.3	1.1
Difference from placebo (adjusted mean‡)	0.1¶	

† Full Analysis Set population (an intention-to-treat analysis).

‡ Least squares mean adjusted for baseline value.

§ p<0.001 compared to placebo.

% All Patients as Treated (APaT) population, excluding data following glycemic rescue therapy.

¶ Not statistically significant (p≥0.05) compared to placebo.

Patients with Renal Insufficiency:

A study comparing sitagliptin at 25 or 50 mg once daily to glipizide at 2.5 to 20 mg/day was conducted in patients with moderate to severe renal impairment. In this study, 277 patients with chronic renal impairment were included in the Per-protocol population (135 patients on JANUVIA®: moderate [n=98], severe [n=37]; and 142 patients on glipizide: moderate [n=106], severe [n=36]). After 54 weeks, the mean reduction from baseline in HbA_{1c} was -0.76% with sitagliptin and -0.64% with glipizide (Per-Protocol Analysis). In this study, the efficacy and safety profile of sitagliptin at 25 or 50 mg once daily was generally similar to that observed in other monotherapy studies in patients with normal renal function. The incidence of hypoglycemia in the sitagliptin group (6.2%) was significantly lower than that in the glipizide group (17.0%).

Another study comparing sitagliptin at 25 mg once daily to glipizide at 2.5 to 20 mg/day was conducted in 129 patients with ESRD who were on dialysis (64 patients on JANUVIA®; and 65 patients on glipizide). After 54 weeks, the mean reduction from baseline in HbA_{1c} was -0.72% with sitagliptin and -0.87% with glipizide. In this study, the efficacy and safety profile of sitagliptin at 25 mg once daily was generally similar to that observed in other monotherapy studies in patients with normal renal function. The incidence of hypoglycemia was not significantly different between the treatment groups (sitagliptin, 6.3%; glipizide, 10.8%).

In a study involving 91 patients with type 2 diabetes and chronic renal impairment (creatinine clearance <50 mL/min), the safety and tolerability of treatment with sitagliptin at 25 or 50 mg once daily were generally similar to placebo.

TECOS Cardiovascular Safety Study

The Trial Evaluating Cardiovascular Outcomes with Sitagliptin (TECOS) was a randomized double-blind, placebo-controlled, parallel-group, event-driven, multicentre study in patients with type 2 diabetes mellitus (HbA1c ≥ 6.5 to 8.0%) and established vascular disease (coronary artery disease, ischemic cerebrovascular disease, atherosclerotic peripheral artery disease). The study included 14,671 (70.7% male, 29.3% female) patients in the intention-to-treat population who received JANUVIA[®] (N=7,332) 100 mg daily (or 50 mg daily if the baseline eGFR was ≥ 30 and <50 mL/min/1.73 m²) or placebo (N=7,339) added to usual care targeting regional standards for HbA1c and CV risk factors. The median duration of treatment was 31 months and the median duration of follow-up was 36 months. Patients with an eGFR <30 mL/min/1.73 m² were not to be enrolled in the study. The study population included 10,863 patients with coronary artery disease, 3,588 patients with cerebrovascular disease, 2,433 patients with peripheral artery disease, 2,643 patients with prior congestive heart failure (including 373 with New York Heart Association [NYHA] class 3 or higher), 2,004 patients ≥ 75 years of age, and 3,324 patients with renal impairment (eGFR <60 mL/min/1.73 m²).

The primary cardiovascular endpoint was a composite of the first occurrence of cardiovascular death, non-fatal myocardial infarction, non-fatal stroke, or hospitalization for unstable angina. Secondary cardiovascular endpoints included a composite of the first occurrence of cardiovascular death, non-fatal myocardial infarction, or non-fatal stroke; as well as first occurrence of the following independent CV endpoints: cardiovascular death, myocardial infarction (fatal + non-fatal), stroke (fatal + non-fatal), hospitalization for unstable angina, hospitalization for heart failure, and all-cause mortality. A composite endpoint of first occurrence of death due to heart failure or hospitalization for congestive heart failure was also assessed

JANUVIA[®], when added to usual care, did not increase the risk of major adverse cardiovascular events or the risk of death or hospitalization for heart failure compared to usual care without JANUVIA[®] in patients with type 2 diabetes. Superiority to placebo was not demonstrated for any endpoint (Table 18).

Table 18 -Rates of Composite Cardiovascular Outcomes and Key Secondary Outcomes Censored at the End of Follow-up (Intention-to-Treat Population)

	JANUVIA [®] (N=7,332)		Placebo (N=7,339)		Hazard Ratio (95% CI)	p-value [†]
	Subjects with Events N (%)	Incidence Rate per 100 Patient-Years*	Subjects with Events N (%)	Incidence Rate per 100 Patient-Years*		
Primary Composite Endpoint (Cardiovascular death, non-fatal myocardial infarction, non-fatal stroke, or hospitalization for unstable angina)	839 (11.4)	4.1	851 (11.6)	4.2	0.98 (0.89, 1.08)	<0.001
Secondary Composite Endpoint (Cardiovascular death, non-fatal myocardial infarction, or non-fatal stroke)	745 (10.2)	3.6	746 (10.2)	3.6	0.99 (0.89–1.10)	<0.001
Secondary Outcome						
Cardiovascular death	380 (5.2)	1.7	366 (5.0)	1.7	1.03 (0.89-1.19)	0.711
All myocardial infarction (fatal and non-fatal)	300 (4.1)	1.4	316 (4.3)	1.5	0.95 (0.81–1.11)	0.487
All stroke (fatal and non-fatal)	178 (2.4)	0.8	183 (2.5)	0.9	0.97 (0.79–1.19)	0.760
Hospitalization for unstable angina	116 (1.6)	0.5	129 (1.8)	0.6	0.90 (0.70–1.16)	0.419
Death from any cause	547 (7.5)	2.5	537 (7.3)	2.5	1.01 (0.90–1.14)	0.875
Hospitalization for heart failure [‡]	228 (3.1)	1.1	229 (3.1)	1.1	1.00 (0.83–1.20)	0.983
Death due to heart failure or hospitalization for heart failure [‡]	237 (3.2)	1.1	240 (3.3)	1.1	0.99 (0.83, 1.18)	0.909

* Incidence rate per 100 patient-years is calculated as $100 \times (\text{total number of patients with } \geq 1 \text{ event during eligible exposure period per total patient-years of follow-up})$.

[†] Based on a Cox model stratified by region. For composite endpoints, the p-values correspond to a test of non-inferiority seeking to show that the hazard ratio is less than 1.3. For all other endpoints, the p-values correspond to a test of differences in hazard rates.

[‡] The analysis of hospitalization for heart failure was adjusted for a history of heart failure at baseline.

DETAILED PHARMACOLOGY

Sitagliptin was assessed for its ability to improve glucose tolerance in lean and diet-induced obese (DIO) mice following dextrose challenge and in diabetic (db/db) mice. In lean and DIO mice, single oral doses of sitagliptin reduced blood glucose levels in a dosage-dependent manner. Acute lowering of blood glucose was also demonstrated in diabetic db/db mice. A 2- to 3-fold increase in

active GLP-1 was seen at the maximally effective dose of 1 mg/kg sitagliptin in lean mice. These results are consistent with the action of sitagliptin as an anti-hyperglycemic agent.

Treatment with GLP-1 or with DPP-4 inhibitors in animal models of type 2 diabetes has been demonstrated to improve beta cell responsiveness to glucose, stimulate insulin biosynthesis and release, increase beta cell neogenesis, and decrease beta cell death. The effects on beta cell neogenesis and beta cell death have not been studied in humans.

TOXICOLOGY

Acute Toxicity

The approximate LD50 of sitagliptin given orally to rats is >3000 mg/kg (maximum dose tested). This dose is equivalent to ≥ 200 times the human exposure based on the recommended daily adult human dose of 100 mg/day. In mice the approximate oral LD50 of sitagliptin is 4000 mg/kg. This dose is equivalent to >385 times the human exposure based on recommended daily adult human dose of 100 mg/day.

Chronic Toxicity

The toxicity potential of sitagliptin was evaluated in a series of repeated dose toxicity studies of up to 53 weeks in dogs and up to 27 weeks in rats. In dogs administered sitagliptin orally at dosages of 2, 10 and 50 mg/kg/day, the no-observed effect level was 10 mg/kg/day (up to 6 times the human exposure based on the recommended daily adult human dose of 100 mg/day). Treatment-related physical signs observed in the 50-mg/kg/day group included open-mouth breathing, salivation, white foamy emesis, ataxia, trembling, decreased activity, and/or hunched posture. These signs were transient, slight in degree, and occurred with decreased incidence during the course of the study. In addition, very slight to slight skeletal muscle degeneration was observed histologically in the 14- and 27-week toxicity studies at the 50-mg/kg/day dose. However, no skeletal muscle degeneration was found in the 53-week toxicity study, indicating the lack of reproducibility or progression of this change with increased duration of treatment. The 50-mg/kg/day dose in dogs resulted in systemic exposure values 26 times the human exposure at the recommended daily adult human dose of 100 mg/day. In rats, sitagliptin administered orally at dosages of up to 180 mg/kg/day (up to 23 times the human exposure based on the recommended daily adult human dose of 100 mg/day), no significant toxicity was observed. The only drug-related effect observed was postdose salivation, likely related to poor palatability of the drug, at doses of 60 mg/kg/day and 180 mg/kg/day.

The treatment-related changes noted in animals do not suggest any clinical concerns at the recommended therapeutic dosages in humans.

Carcinogenicity

A two-year carcinogenicity study was conducted in male and female rats given oral doses of sitagliptin of 50, 150, and 500 mg/kg/day. There was an increased incidence of hepatic adenomas and carcinomas in the high-dose males and hepatic carcinomas in the high-dose females. This dose in rats results in approximately 58 times the human exposure based on the recommended daily adult human dose of 100 mg/day. This dose level was associated with hepatotoxicity in rats. The no-observed effect level for induction of hepatic neoplasia was 150 mg/kg/day, approximately 19-fold the human exposure at the 100-mg recommended dose. Since hepatotoxicity has been shown to correlate with

induction of hepatic neoplasia in rats, this increased incidence of hepatic tumors in rats was likely secondary to chronic hepatic toxicity at this high dose. The clinical significance of these findings for humans is unknown.

A two-year carcinogenicity study was conducted in male and female mice at oral doses of 50, 125, 250, and 500 mg/kg/day. Sitagliptin did not increase tumor incidence in mice in any organ at doses up to 500 mg/kg/day (approximately 68 times the human exposure based on the recommended daily adult human dose of 100 mg/day).

Mutagenesis

Sitagliptin was not mutagenic or clastogenic in a battery of genetic toxicology studies, including the Ames bacterial assay (microbial mutagenesis test), Chinese hamster ovary cells (CHO cells) chromosome aberration assay, an *in vitro* cytogenetics assay using CHO cells, an *in vitro* rat hepatocyte DNA alkaline elution assay (an assay which measures the compound's ability to induce single strand breaks in DNA), and an *in vivo* micronucleus assay.

Reproduction

No adverse effects upon fertility were observed in male and female rats given sitagliptin orally at doses up to 1000 mg/kg daily (up to approximately 100 times the human exposure based on the recommended daily adult human dose of 100 mg/day) prior to and throughout mating.

Development

Sitagliptin was not teratogenic in rats at oral doses up to 250 mg/kg or in rabbits given up to 125 mg/kg during organogenesis (up to 32 and 22 times the human exposure based on the recommended daily adult human dose of 100 mg/day). A slight, treatment-related increased incidence of fetal rib malformations (absent, hypoplastic and wavy ribs) was observed in the offspring of rats at oral doses of 1000 mg/kg/day (approximately 100 times the human exposure based on the recommended daily adult human dose of 100 mg/day). The no-observed effect level for developmental effects was 250 mg/kg/day (32 times the human exposure based on the recommended daily adult human dose of 100 mg/day). Treatment-related decreases in the mean preweaning body weight of both sexes and postweaning body weight gain of male animals was observed in offspring of rats at oral doses of 1000 mg/kg.

REFERENCES

1. Canadian Diabetes Association Clinical Practice Guidelines Expert Committee. Canadian Diabetes Association 2013 Clinical Practice Guidelines for the Prevention and Management of Diabetes in Canada. *Can J Diabetes* 2013;37 (Suppl 1):S1-S212.
2. Charbonnel B, Karasik A, Liu J, Wu M, Meininger G. Efficacy and safety of the dipeptidyl peptidase-4 inhibitor sitagliptin added to ongoing metformin therapy in patients with type 2 diabetes inadequately controlled with metformin alone. *Diabetes Care* 2006;29(12):2638-43.
3. Drucker DJ. Development of glucagon-like peptide-1-based pharmaceuticals as therapeutic agents for the treatment of diabetes. *Curr Pharm Des* 2001;7(14):1399-412.
4. Nauck MA, Meininger G, Sheng D, Terranella L, Stein PP. Efficacy and safety of the dipeptidyl peptidase-4 inhibitor, sitagliptin, compared with the sulfonylurea, glipizide, in patients with type 2 diabetes inadequately controlled on metformin alone: a randomized, double-blind, non-inferiority trial. *Diabetes Obes Metab* 2007;9(2):194-205.
5. Williams-Herman D, Engel SS, Round E, Johnson J, Golm GT et al. Safety and tolerability of sitagliptin in clinical studies: a pooled analysis of data from 10,246 patients with type 2 diabetes. *BMC Endocr Disord* 2010;10(7).
6. Vilsbøll T, Rosenstock J, Yki-Järvinen H, Cefalu WT, Chen Y, Luo E et al. Efficacy and safety of sitagliptin when added to insulin therapy in patients with type 2 diabetes. *Diabetes Obes Metab* 2010;12:167-77.
7. Rosenstock J, Brazg R, Andryuk J. P, Lu K, Stein P. Efficacy and Safety of the Dipeptidyl Peptidase-4 inhibitor Sitagliptin Added to Ongoing Pioglitazone Therapy in Patients with Type 2 Diabetes: A 24-Week, Multicenter, Randomized, Double-Blind, Placebo-Controlled, Parallel-Group Study. *Clinical Therapeutics* 2006;28:1556-1568.
8. Green JB, Bethel MA, Armstrong PW, Buse JB et al. Effect of sitagliptin on cardiovascular outcomes in type 2 diabetes. *N Engl J Med* 2015;373: 232-42.

PART III: CONSUMER INFORMATION

JANUVIA[®]
sitagliptin tablets
(as sitagliptin phosphate monohydrate)

This leaflet is part III of a three-part “Product Monograph” published when JANUVIA[®] was approved for sale in Canada and is designed specifically for Consumers. This leaflet is a summary and will not tell you everything about JANUVIA[®]. Contact your physician or pharmacist if you have any questions about the drug.

Please read this information carefully before you start to take your medicine, even if you have just refilled your prescription. Some of the information may have changed.

Remember that your physician has prescribed this medicine only for you. Never give it to anyone else.

ABOUT THIS MEDICATION

What the medication is used for:

- JANUVIA[®] can be used to improve blood sugar levels in adult patients with type 2 diabetes mellitus in addition to diet and exercise:
 - alone in patients who cannot take metformin
 - in combination with metformin
 - in combination with metformin and a sulfonylurea (e.g. glyburide, gliclazide or glimepiride)
 - in combination with premixed or long/intermediate acting insulin (with or without metformin)
 - in combination with pioglitazone (with or without metformin)

What it does:

JANUVIA[®] is a member of a class of medicines called DPP-4 inhibitors (dipeptidyl peptidase-4 inhibitors). JANUVIA[®] helps to improve the levels of insulin when blood sugar level is high, especially after a meal. JANUVIA[®] also helps to decrease the amount of sugar made by the body. JANUVIA[®] is unlikely to cause low blood sugar (hypoglycemia).

What is type 2 diabetes?

Type 2 diabetes is a condition in which your body does not make enough insulin, and/or does not use the insulin that your body produces as well as it should. When this happens, sugar (glucose) builds up in the blood. This can lead to serious problems.

When it should not be used:

Do not take JANUVIA[®] if you are allergic to any of the ingredients in JANUVIA[®].

What the medicinal ingredient is:
 sitagliptin phosphate monohydrate

What the important non-medicinal ingredients are:
 Each film-coated tablet of JANUVIA[®] contains the following inactive ingredients: microcrystalline cellulose, anhydrous dibasic calcium phosphate (calcium hydrogen phosphate, anhydrous), croscarmellose sodium, magnesium stearate, and sodium stearyl fumarate. In addition, the film coating contains the following inactive ingredients: polyvinyl alcohol, polyethylene glycol, talc, titanium dioxide, red iron oxide, and yellow iron oxide.

What dosage forms it comes in:
 Tablets: Each tablet contains 25, 50 or 100 mg sitagliptin.

WARNINGS AND PRECAUTIONS

Cases of inflammation of the pancreas (pancreatitis) which may be severe and lead to death have been reported in patients taking JANUVIA[®].

Cases of a skin reaction called bullous pemphigoid that can require treatment in a hospital have been reported in patients receiving JANUVIA[®]. Tell your doctor if you develop blisters or the breakdown of your skin (erosion). Your doctor may tell you to stop taking JANUVIA[®].

BEFORE you use JANUVIA[®] talk to your physician or pharmacist if:

- you have or have had pancreatitis (inflammation of the pancreas) or any risk factors for pancreatitis such as gallstones (solid particles that form in the gall bladder), a history of alcoholism, high triglyceride levels.
- you have type 1 diabetes.
- you have or have had diabetic ketoacidosis (increased ketones in the blood or urine).
- you are taking a sulfonylurea. When JANUVIA[®] is used in combination with a sulfonylurea and metformin, or in combination with insulin (with or without metformin), low blood sugar can occur. Your physician may consider lowering the dose of the sulfonylurea or insulin. Take precautions to avoid low blood sugar while driving or using machinery.
- you have or have had an allergic reaction to JANUVIA[®].
- you have or have had kidney problems.
- you have liver problems.
- you are pregnant or plan to become pregnant. JANUVIA[®] is not recommended for use during pregnancy.
- you are breast-feeding or plan to breast-feed. It is not known if JANUVIA[®] passes into breast milk.

JANUVIA[®] may cause abnormal kidney function. Your doctor will do blood tests to monitor how well your kidneys are working while you are taking JANUVIA[®].

JANUVIA[®] is not recommended for use in children under 18 years of age.

INTERACTIONS WITH THIS MEDICATION

Tell your physician or pharmacist about all the medicines you take. This includes prescription and non-prescription medicines, and herbal supplements.

PROPER USE OF THIS MEDICATION

Take JANUVIA[®] exactly as your physician has prescribed.

Usual dose:

100 mg once daily by mouth with or without food. The daily dose may be adjusted in patients with kidney problems.

Overdose:

In case of drug overdose, contact a health care practitioner, hospital emergency department or regional Poison Control Centre immediately, even if there are no symptoms.

Missed Dose:

If you miss a dose, take it as soon as you remember. If you do not remember until it is time for your next dose, skip the missed dose and go back to your regular schedule. Do not take a double dose of JANUVIA[®] on the same day.

SIDE EFFECTS AND WHAT TO DO ABOUT THEM

Like all medicines, JANUVIA[®] may cause side effects.

The most common side effects of JANUVIA[®] include: stuffy or runny nose and sore throat.

Tell your physician or pharmacist if you develop any unusual side effects, or any of the above side effects that do not go away or gets worse.

When JANUVIA[®] is used with metformin and a sulfonylurea medicine, or with insulin (with or without metformin), low blood sugar (hypoglycemia) can occur. Lower doses of the sulfonylurea medicine or insulin may be required while you use JANUVIA[®]. The signs and symptoms of low blood sugar may include headache, drowsiness, weakness, dizziness, confusion, irritability, hunger, fast heartbeat, sweating, and feeling jittery. Discuss with your physician or pharmacist how to treat low blood sugar.

Additional side effects have been reported in general use with JANUVIA[®]:

- Allergic reactions, which may be serious, including rash, hives, and swelling of the face, lips, tongue, and throat that may cause difficulty in breathing or swallowing. If you have an allergic reaction, stop taking JANUVIA[®] and call your physician right

away. Your physician may prescribe a medication to treat your allergic reaction and a different medication for your diabetes.

- Inflammation of the pancreas
- Kidney problems (sometimes requiring dialysis)
- Vomiting
- Constipation
- Headache
- Joint pain
- Muscle aches
- Arm or leg pain
- Back pain
- Itching
- Blisters

SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM			
Symptoms / Effects	Talk with your physician or pharmacist		Stop taking drug and call your physician or pharmacist
	Only if severe	In all cases	
Very common			
Hypoglycemia (when used with metformin and a sulfonylurea, or when used with insulin with or without metformin)		√	
Rare			
Pancreatitis: symptoms of pancreatitis (prolonged severe abdominal pain which may be accompanied by vomiting)		√	√
Allergic reactions including rash, hives, and swelling of the face, lips, tongue, and throat that may cause difficulty in breathing or swallowing.			√
Bullous pemphigoid (blisters or breakdown of your skin)		√	
Acute kidney failure: symptoms may include nausea, loss of appetite and weakness, pass little or no urine, breathlessness.		√	

This is not a complete list of side effects. For any unexpected effects while taking JANUVIA[®], contact your physician or pharmacist.

HOW TO STORE IT

Store at room temperature (15°C to 30°C).

Keep JANUVIA[®] and all medicines safely away from children.

REPORTING SUSPECTED SIDE EFFECTS

Reporting Side Effects

You can help improve the safe use of health products for Canadians by reporting serious and unexpected side effects to Health Canada. Your report may help to identify new side effects and change the product safety information.

3 ways to report:

- Online at [MedEffect](http://hc-sc.gc.ca/dhp-mps/medeff/index-eng.php) (<http://hc-sc.gc.ca/dhp-mps/medeff/index-eng.php>);
- By calling 1-866-234-2345 (toll-free);
- By completing a Consumer Side Effect Reporting Form and sending it by:

- Fax to 1-866-678-6789 (toll-free), or
- Mail to: Canada Vigilance Program
Health Canada, Postal Locator
1908C
Ottawa, ON
K1A 0K9

Postage paid labels and the Consumer Side Effect Reporting Form are available at [MedEffect](http://hc-sc.gc.ca/dhp-mps/medeff/index-eng.php) (<http://hc-sc.gc.ca/dhp-mps/medeff/index-eng.php>)

NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice. »

To report an adverse event related to JANUVIA[®], please contact 1-800-567-2594.

MORE INFORMATION

If you want more information about JANUVIA[®]:

- Talk to your healthcare professional
- Find the full product monograph that is prepared for healthcare professionals and includes this Consumer Information by visiting the [Health Canada website](http://www.healthcanada.gc.ca) or Merck Canada website www.merck.ca or by calling Merck Canada at 1-800-567-2594

This leaflet was prepared by Merck Canada Inc.

Last revised: March 13, 2017

[®] Merck Sharp & Dohme Corp. Used under license.

© 2011, 2017 Merck Canada Inc. All rights reserved.