

PRODUCT MONOGRAPH

Maximum Strength Acid Reducer Without Prescription
Ranitidine Tablets, USP
(as Ranitidine Hydrochloride)

150 mg

Histamine H₂-receptor antagonist

DOMINION PHARMACAL
6111 Royalmount Ave., Suite #100
Montréal, Québec
H4P 2T4

Date of Preparation
July 10, 2017

Submission Control No.: 206134

PRODUCT MONOGRAPH

Maximum Strength Acid Reducer Without Prescription Ranitidine Tablets, USP (as Ranitidine Hydrochloride) 150 mg

Histamine H₂-receptor antagonist

CLINICAL PHARMACOLOGY

Ranitidine is an antagonist of histamine at gastric H₂-receptor sites. Thus, ranitidine inhibits both basal gastric secretion and gastric acid secretion induced by histamine, pentagastrin and other secretagogues. Inhibition of gastric acid secretion has been observed following intravenous, intraduodenal and oral administration of ranitidine. This response is dose-related, a maximum response being achieved at an oral dose of 300 mg/day.

Pepsin secretion is also inhibited but secretion of gastric mucus is not affected. Ranitidine does not alter the secretion of bicarbonate or enzymes from the pancreas in response to secretin and pancreozymin.

Ranitidine is rapidly absorbed after oral administration, peak plasma concentrations being achieved within 2 to 3 hours. These plasma concentrations are not significantly influenced by the presence of food in the stomach at the time of the oral administration nor by regular doses of antacids.

Bioavailability of oral ranitidine is approximately 50%. Serum protein binding of ranitidine in man is in the range 10 to 19%. The elimination half-life is approximately 3 hours. The principal route of excretion is the urine (40% recovery of free and metabolized drug in 24 hours).

There is a significant linear correlation between the dose administered and the inhibitory effect upon gastric acid secretion for single oral doses up to 300 mg. In healthy subjects a single 75 mg dose of ranitidine significantly reduced meal-stimulated intragastric acidity ($[H^+]$ AUC) compared with placebo. The effect of ranitidine on intragastric acidity and pH is also dose-related.

The important pharmacologic activity of ranitidine is inhibition of gastric acid and fluid secretion in basal and stimulated states, which increases the pH and decreases the volume of secretions. A single 75 mg dose, compared to placebo, has an early onset of action; significantly elevating gastric pH within one hour, and lasting for up to 13 hours post dosing. After correcting for onset of action (within one hour), the duration of acid suppression for ranitidine 75 mg is up to 12 hours (i.e. all day or all night). In the same multicentre, randomized, cross-over study, the onset of acid suppression effect for ranitidine 75 mg was statistically superior to famotidine 10 mg at only one and two hours post-dosing. The duration and degree of acid suppression of ranitidine 75 mg (63.1%, n=75) were superior to cimetidine 200 mg (37.8%, n=52) over the 10-hour daytime evaluation period.

Table 1. Summary of weighted mean hydrogen ion activity (H⁺ AUC mmol/L.h.)

| Time Period | Placebo N=75 subjects | Ranitidine 75 mg N=75 subjects | Cimetidine 200 mg N=52 subjects | Famotidine 10 mg N=22 subjects |
|-----------------------|--------------------------|--------------------------------------|---------------------------------------|--------------------------------------|
| Total (20-h) | 30.89 | 18.21*† | 25.08* | 19.32* |
| Day (11.00h-22.30h) | 32.76 | 18.19*† | 22.06* | 13.23* |
| Night (22.30h-08.30h) | 28.83 | 23.23*† | 28.09 | 25.41* |

* p<0.05 compared with paired placebo group

† p<0.05 compared with paired cimetidine group

Table 2. Changes in intragastric acidity and pH

| | Median percentage decrease in acidity | | | Median intragastric pH | |
|--------------------------------|---------------------------------------|------------------|--------------------|------------------------|--------------------|
| | total | day ^a | night ^a | day ^a | night ^a |
| Ranitidine 75 mg (N=75) | 44.1% † | 63.1% † | 21.2% † | 2.10* | 1.80* |
| Famotidine 10 mg (N=22) | 38.9% | 58.9% | 20.1% | 2.06* | 1.90 |
| Cimetidine 200 mg (N=52) | 23.0% | 37.8% | 1.8% | 1.69* | 1.77 |
| Placebo (N=75) | | | | 1.48 | 1.70 |

*p<0.05 compared with paired placebo group

†p<0.05 compared with paired cimetidine group

^aday=12.30h to 22.30h; night=22.30h to 8.30h

In a large, multicentre, dose-ranging, placebo-controlled trial in patients with episodic heartburn, a single 75 mg dose relieved symptoms within 30 minutes and provided relief for the duration of the 4-hour evaluation period.

Volunteers treated with an oral dose of ranitidine have reported no significant gastrointestinal or central nervous system side effects; moreover pulse rate, blood pressure, electrocardiogram and electroencephalogram are not significantly affected in man following ranitidine administration. In healthy human volunteers and patients, ranitidine, when administered orally did not influence plasma levels of the following hormones: cortisol, testosterone, oestrogens, growth hormone, follicle-stimulating hormone, luteinizing hormone, thyroid-stimulating hormone, aldosterone or gastrin; although like cimetidine, ranitidine reduced vasopressin output. Treatment for up to 6 weeks with ranitidine 150 mg twice daily by mouth did not affect the human hypothalamic-pituitary-testicular-ovarian or - adrenal axes.

The safety and efficacy of 75 mg ranitidine for treatment of episodic heartburn were established in two large replicate Phase III studies involving 2,985 patients. These two pivotal studies demonstrated that one ranitidine 75 mg tablet was statistically and clinically superior to placebo in providing relief of episodic heartburn beginning at 30 minutes.

In two subsequent heartburn treatment trials evaluating total pain relief, both the 75 mg and 150 mg strengths of ranitidine were demonstrated to be statistically significantly superior to placebo

as shown in the table below.

Table 3 presents pooled data for the primary efficacy variable of total pain relief for the first treated episode (TOTPAR) along with its associated p values.

Table 3. Pooled Treatment Data Primary Variable (Intent-to-treat subjects)

| Efficacy Variable | Ranitidine 150 mg (N=637) | Ranitidine 75 mg (N=637) | Placebo (N=635) | Ranitidine 75 mg vs placebo (p value) | Ranitidine 150 mg vs placebo (p value) | Ranitidine 150 mg vs Ranitidine 75 mg (p value) |
|--|---------------------------|--------------------------|-----------------|---------------------------------------|--|---|
| TOTPAR for 1 st drug treated episode (mean) | 20.8 | 20.4 | 18.3 | <0.001 | 0.004 | 0.673 |

The safety and efficacy of 150 mg ranitidine for the prevention or reduction of severity of meal-induced heartburn when taken immediately before consuming food and beverages anticipated to provoke heartburn was studied in two pivotal and one supporting large replicate Phase III well-controlled, multi-center, double-blind, randomized, placebo-controlled studies involving 2,484 patients. The results of the 2 pivotal and supporting studies are summarized below.

Table 4. Pooled Prevention Data for Key Efficacy Variables (Intent-to-Treat Subjects) Pivotal Studies

| Efficacy Variable | Ranitidine 150 mg (N=518) | Ranitidine 75 mg (N=524) | Placebo (N=521) |
|---|---------------------------|---------------------------|-----------------|
| Treatment Meal AUC (mm hr) | 87.0 ± 3.96 *† | 91.1 ± 3.84 ^{*a} | 107.5 ± 4.17 |
| Reduction in Treatment Meal AUC (%) | 49.1 ± 2.44* | 44.0 ± 2.29 ^{*b} | 32.7 ± 2.93 |
| Treatment Meal peak heartburn severity (mm) | 40.2 ± 1.29* | 41.6 ± 1.23 ^{*c} | 47.0 ± 1.28 |
| Peak heartburn reduction (%) | 42.6 ± 1.78* | 38.7 ± 1.73 ^{*d} | 31.2 ± 1.79 |
| Largest consecutive time points with no heartburn | 5.3 ± 0.24* | 4.7 ± 0.23* | 4.2 ± 0.22 |

*p<0.05 versus placebo, †p<0.05 versus ranitidine 75 mg, ^aN=520, ^bN=518, ^cN=521, ^dN=519

Table 5. Prevention Data for Key Efficacy Variables (Intent-to-Treat Subjects) Supporting Study

| Efficacy Variable | Ranitidine 150 mg (N=306) | Ranitidine 75 mg (N=309) | Placebo (N=306) |
|-------------------------------------|---------------------------|--------------------------|-----------------|
| Treatment Meal AUC (mm hr) | 94.6 ± 5.91 | 88.6 ± 4.97 | 98.8 ± 4.83 |
| Reduction in Treatment Meal AUC (%) | 45.4 ± 3.08 | 43.4 ± 3.20 | 39.2 ± 3.05 |

| Efficacy Variable | Ranitidine 150 mg (N=306) | Ranitidine 75 mg (N=309) | Placebo (N=306) |
|---|------------------------------|-----------------------------|--------------------|
| Treatment Meal peak heartburn severity (mm) | 42.4 ± 1.80 | 43.4 ± 1.60 | 46.8 ± 1.61 |
| Peak heartburn reduction (%) | 39.7 ± 2.47 | 37.4 ± 2.30 | 32.8 ± 2.29 |
| Largest consecutive time points with no heartburn | 5.5 ± 0.32* | 5.2 ± 0.30 | 4.4 ± 0.29 |

*p≤0.05 versus placebo

In pooled data of the 2 studies reflected in Table 4, ranitidine 150 mg was effective in preventing meal-related heartburn. Ranitidine 75 mg was statistically significantly better than placebo for the primary endpoint in one of the 2 pivotal clinical trials and in the pooled analysis, but was less consistently effective than ranitidine 150 mg in the prevention pivotal trials.

Comparative Bioavailability Studies

A single center, randomized, single-dose, blinded, 2-period, 2-sequence crossover comparative bioavailability study was performed under fasting conditions on Ranitidine Hydrochloride tablets using DOMINION PHARMACAL ranitidine 300 mg tablets versus the reference product, ZANTAC[®] 300 mg Tablets, by GlaxoSmithKline Inc. Canada. Twenty-two healthy male volunteers took part in the study. The pharmacokinetic data calculated for the ranitidine 300 mg tablets and ZANTAC[®] 300 mg tablets formulation are tabulated below:

SUMMARY TABLE OF THE COMPARATIVE BIOAVAILABILITY DATA

| Ranitidine (1 x 300 mg tablet) From measured data uncorrected for potency Geometric Mean Arithmetic Mean (CV %) | | | | |
|---|-------------------------|-------------------------|-------------------------------|------------------------------|
| Parameter | Ranitidine | ZANTAC [†] | % Ratio of Geometric Means | Confidence Interval (90%) |
| AUC _T (ng·h/mL) | 4656.9 4801.8 (25.2) | 4636.2 4775.4 (23.1) | 100.45 | 94.03 – 107.31 |
| AUC _I (ng·h/mL) | 4781.1 4929.1 (25.0) | 4759.3 4901.2 (23.0) | 100.46 | 94.15 – 107.19 |
| C _{max} (ng/mL) | 987.2 1032.4 (30.5) | 915.9 961.8 (32.4) | 107.79 | 98.86 – 117.52 |
| T _{max} [§] (h) | 2.84 (1.33 – 4.00) | 3.00 (1.67 – 4.00) | | |
| T _½ ^² (h) | 2.79 (11.7) | 2.85 (16.4) | | |

[†] ZANTAC was manufactured by GlaxoSmithKline Inc., Canada, and was purchased in Canada. McNeil Consumer Healthcare is the current manufacturer for ZANTAC.

[§] Expressed as the median (range)

^² Expressed as the arithmetic mean (CV%)

INDICATIONS AND CLINICAL USE

MAXIMUM STRENGTH ACID REDUCER WITHOUT PRESCRIPTION Tablets are indicated for fast and effective relief, treatment, and prevention, day or night, of the burning and discomfort of acid indigestion (dyspepsia), heartburn, hyperacidity, sour stomach, and upset stomach associated with excess stomach acid. These symptoms may be brought on by consuming food and beverages.

CONTRAINDICATIONS

MAXIMUM STRENGTH ACID REDUCER WITHOUT PRESCRIPTION (ranitidine hydrochloride) is contraindicated for patients known to have hypersensitivity to any component of the preparation.

WARNINGS

Gastric Carcinoma

Treatment with a histamine H₂-antagonist may mask symptoms associated with carcinoma of the stomach and, therefore, may delay diagnosis of that condition. Accordingly, patients should be advised to consult a physician if they have difficulty swallowing or persistent abdominal discomfort or if symptoms get worse or persist for more than 2 weeks.

Use in Patients with a History of Acute Porphyria

Rare clinical reports suggest that ranitidine may precipitate acute porphyric attacks. Therefore, ranitidine should be avoided in patients with a history of acute porphyria.

Use in Pregnancy and Nursing Mothers

The safety of ranitidine hydrochloride in the treatment of conditions where a controlled reduction of gastric secretion is required during pregnancy has not been established. Reproduction studies performed in rats and rabbits at higher doses have revealed no evidence of ranitidine induced impaired fertility or harm to the fetus. Nevertheless, if the administration of ranitidine is considered to be necessary, its use requires that the potential benefits be weighed against possible hazards to the patient and to the fetus.

Ranitidine is secreted in breast milk in lactating mothers but the clinical significance of this has not been fully evaluated. Women who are breast-feeding are advised to speak with their doctor before taking MAXIMUM STRENGTH ACID REDUCER WITHOUT PRESCRIPTION Tablets.

PRECAUTIONS

Use in Impaired Renal Function

Ranitidine is excreted via the kidneys and, in the presence of severe renal impairment, plasma levels of ranitidine are increased and elimination prolonged. Accordingly, MAXIMUM STRENGTH ACID REDUCER WITHOUT PRESCRIPTION should be used under physician supervision for these patients.

Interaction with Other Drugs

Although ranitidine has been reported to bind weakly to cytochrome P₄₅₀ *in vitro*, recommended doses of the drug do not inhibit the action of the hepatic cytochrome P₄₅₀-linked oxygenase enzymes. A review of selected publications of controlled clinical drug interaction studies, at the level of hepatic elimination has indicated ranitidine is unlikely to cause clinically significant potentiation of actions of drugs which are inactivated by the hepatic cytochrome P₄₅₀ enzyme system; these drugs may include: diazepam, phenytoin, propranolol, theophylline, and warfarin. Sporadic cases (approximately 1 case per 4 million patient treatments) of drug interactions have been reported in elderly patients involving both hypoglycaemic drugs and theophylline. The significance of these reports cannot be determined at present, as controlled clinical trials have not shown interactions. These reports are based on use for prescription indications and dosage. Patients consuming NSAIDs may have dyspepsia as a side effect of these medicines and should consult a physician or a pharmacist before taking MAXIMUM STRENGTH ACID REDUCER WITHOUT PRESCRIPTION.

Antacids: concurrent administration of high-dose antacids (75 mEq) with ranitidine is not recommended. The absorption of ranitidine may be decreased. Patients should be cautioned not to take antacids within ½ to 1 hour of ranitidine ingestion.

Ketoconazole: Simultaneous administration of ketoconazole and ranitidine may result in reduction of the absorption of ketoconazole by some mechanism as yet unidentified (e.g., a pH dependent effect on absorption or a change of volume of distribution). Patients should be cautioned not to take ranitidine for at least 2 hours after ketoconazole. These reports are based on use of prescription indications and dosage.

Sucralfate: If high doses of sucralfate (two grams) are co-administered with MAXIMUM STRENGTH ACID REDUCER WITHOUT PRESCRIPTION, the absorption of MAXIMUM STRENGTH ACID REDUCER WITHOUT PRESCRIPTION may be reduced. This effect is not seen if sucralfate is taken at least two hours after ranitidine hydrochloride administration. These reports are based on use of prescription indications and dosage.

Procinamide: Ranitidine is believed to compete for active renal tubular secretion and/or inhibit the absorption of procainamide. Data indicates adjustment of procainamide doses in the elderly, patients with renal insufficiency, and patients receiving high doses of ranitidine are warranted. These reports are based on use for prescription indications and dosage.

Interactions with the following drugs have been reported with prescription strength ranitidine: Alendronate, amoxicillin, clavulanic acid, cefditoren, cefpodoxime, ceftibuten, delavirdine,

dicumarol, diltiazem, enoxacin, fluvastatin, fosphenytoin, gefitinib, glyburide, itraconazole, metformin, midazolam, miglitol, nisoldipine, pancreatin, pancrelipase, pancuronium, pentoxifylline, sucralfate, tolazoline, triamterene, triazolam, tubocurarine.

In a ranitidine-triazolam drug-drug interaction study, triazolam plasma concentrations were higher during b.i.d. dosing of ranitidine than triazolam given alone. The mean area under the triazolam concentration-time curve (AUC) values in 18- to 60-year-old subjects were 10% and 28% higher following administration of 75 mg and 150 mg ranitidine tablets, respectively, than triazolam given alone. In subjects older than 60 years of age, the mean AUC values were approximately 30% higher following administration of 75 mg and 150 mg ranitidine tablets. It appears that there were no changes in pharmacokinetics of triazolam and (alpha)-hydroxytriazolam, a major metabolite, and in their elimination. Reduced gastric acidity due to ranitidine may have resulted in an increase in the availability of triazolam. The clinical significance of this triazolam and ranitidine pharmacokinetic interaction is unknown.

Ethanol: Consumption of moderate amounts of alcohol is unlikely to result in clinically important alterations in blood alcohol concentrations and/or alcohol metabolism.

The co-administration of a single oral dose of ranitidine 75 mg and ethanol 0.15 g/Kg has no clinically relevant effect on ethanol pharmacokinetics as shown in a double-blind placebo-controlled, crossover study in 25 healthy subjects.

Use in the Elderly

Since malignancy is more common in the elderly, particular consideration must be given to this before therapy with MAXIMUM STRENGTH ACID REDUCER WITHOUT PRESCRIPTION is instituted. Elderly patients receiving NSAIDs concomitantly with MAXIMUM STRENGTH ACID REDUCER WITHOUT PRESCRIPTION should be closely supervised. As with all medication, in the elderly, consideration should be given to concurrent drug therapy.

ADVERSE REACTIONS

Ranitidine has been shown to be generally very well tolerated. In various clinical trials involving either 75 mg or 150 mg of ranitidine the adverse reaction rates were comparable with the most frequently reported adverse events being: headache, nausea, vomiting, and diarrhea: common [frequent] >1% and <10%. Overall adverse event incidence among ranitidine-treated subjects was comparable to that seen in placebo-treated subjects, (no statistical difference) independent of demographic characteristics.

The following adverse reactions have been reported as events in clinical trials, in post-marketing surveillance, or in the routine management of patients treated with prescription doses of ranitidine hydrochloride. The majority of these events have been observed following oral administration of higher prescription doses of ranitidine, and a causal relationship to ranitidine hydrochloride has not always been established.

Central Nervous System: headache, sometimes severe; malaise; dizziness; somnolence; insomnia; vertigo; dystonia; and reversible blurred vision suggestive of a change in

accommodation. Isolated cases of reversible mental confusion, delusion, delirium, mania, agitation, depression, hallucinations have been reported, predominantly in severely ill elderly patients.

Cardiovascular: As with other H₂ receptor antagonists, there have been rare reports of arrhythmias such as tachycardia, premature ventricular beats, bradycardia, and atrioventricular block.

Gastrointestinal: Constipation, diarrhea, nausea/vomiting and abdominal discomfort/pain. Rebound hypersecretion has been reported upon withdrawal of ranitidine therapy.

Hepatic: Transient and reversible changes in liver function tests can occur (increase in ALT and AST values). With oral administration, there have been occasional reports of hepatitis, hepatocellular or hepatocanalicular or mixed with or without jaundice and cholestatic jaundice. In such circumstances, ranitidine should be discontinued immediately. These are usually reversible, but in exceedingly rare circumstances, death has occurred.

Musculoskeletal: Rare reports of arthralgia and myalgia.

Haematologic: Blood count changes (anemia, leukopenia, thrombocytopenia) have occurred in a few patients. These are usually reversible. Rare cases of agranulocytosis, pancytopenia, or neutropenia, sometimes with marrow hypoplasia or aplasia have been reported.

Endocrine: No clinically significant interference with endocrine or gonadal function has been reported. There have been rare reports of breast symptoms in men taking ranitidine.

Dermatologic: Rash, including cases suggestive of mild erythema multiforme. Alopecia, contact dermatitis, photosensitivity.

Other: Rare cases of hypersensitivity reactions (including angina, chest pain, bronchospasm, fever, rash, eosinophilia, anaphylaxis, urticaria, angioneurotic edema, hypotension) and small increases in serum creatinine have occasionally occurred after a single dose. Acute pancreatitis, meningitis, nephrotoxicity has been reported rarely. Rare cases of reversible involuntary motor disturbances and intraocular pressure changes have been reported.

Symptoms and Treatment of Overdosage

There is no experience to date with deliberate overdosage. The usual measures to remove unabsorbed drug from the gastrointestinal tract (including activated charcoal or syrup of ipecac), clinical monitoring and supportive therapy should be employed. Also, if need be, the drug can be removed from the plasma by haemodialysis. Up to 6 g per day has been administered without untoward effect.

For management of a suspected drug overdose, contact your regional Poison Control Centre immediately

DOSAGE AND ADMINISTRATION

ADULTS AND CHILDREN 16 YEARS OF AGE AND OLDER

One tablet should be taken when symptoms appear, day or night. If symptoms persist for more than 1 hour or return after 1 hour, a second tablet may be taken. The maximum dosage is 2 tablets (300 mg ranitidine) in a 24-hour period.

For prevention of symptoms brought on by consuming food or beverages, 1 tablet should be taken 30 to 60 minutes before eating a meal or consuming a beverage that is expected to cause symptoms.

Patients are advised to consult their physician if symptoms get worse or continue after 14 days of treatment.

CHILDREN UNDER 16 YEARS

Children under 16 years of age should be supervised by a physician.


PHARMACEUTICAL INFORMATION

Drug Substance

Proper Name: Ranitidine Hydrochloride

Chemical Name: (n- {2- [(5-[(dimethylamino)-methyl]-2-furanyl)-methyl] thio]ethyl}-n-methyl-2-nitro-1,1 -ethenediamine, hydrochloride.

Structural Formula:



Molecular Formula: $C_{13}H_{22}N_4O_3S$ HCl

Molecular Mass: 350.87 g/mol (as hydrochloride salt)

Description: Ranitidine hydrochloride is a practically odourless white to pale yellow granular substance. At room temperature, ranitidine hydrochloride is soluble in water, methanol, ethanol, and chloroform.

Composition

MAXIMUM STRENGTH ACID REDUCER WITHOUT PRESCRIPTION (ranitidine hydrochloride) Tablets

Medicinal Ingredients: 150 mg of ranitidine as ranitidine hydrochloride.

Non-medicinal Ingredients: Tablets also contain Colloidal Silicon Dioxide, Croscarmellose Sodium, Iron Oxide Red, Iron Oxide Yellow, Magnesium Stearate, Microcrystalline Cellulose, Polyvinyl Alcohol, Soy Lecithin, Talc, Titanium Dioxide, and Xanthan Gum.

Stability and Storage Recommendations

Store between 15°C and 30°C. Protect from light and moisture.

Availability of Dosage Forms

MAXIMUM STRENGTH ACID REDUCER WITHOUT PRESCRIPTION Tablets are salmon color, shield-shaped, five-sided biconvex, coated tablet debossed with “150” on one side and nothing on the other side. Available in bottles of 50 tablets, and in blister packs of 8, and 24.

PHARMACOLOGY

Animal Pharmacology

Ranitidine is a potent competitive reversible, selective antagonist of histamine at H₂-receptors *in vitro* and *in vivo*. Thus, ranitidine antagonised the actions of histamine at H₂-receptors in the rat isolated uterus and in the guinea-pig isolated atrium. Ranitidine is not an anticholinergic agent. On a molar basis, ranitidine is 4 to 5 times more active than cimetidine with a pA₂ value of 7.2. In concentrations 1,000 times greater than those required to block H₂- receptors, it failed to block either H₁-receptors or muscarinic receptors in the guinea-pig isolated ileum. The beta-adrenoceptor responses of the rat uterus and guinea pig atrium to isoprenaline were also unaffected by ranitidine.

Blockade of histamine H₂- receptors in the stomach *in vivo* is the pharmacological action of ranitidine with greatest immediate clinical relevance. Ranitidine inhibits gastric secretion induced by various secretagogues in both the rat and dog.

In the conscious dog with a Heidenhain pouch, ranitidine given orally or intravenously antagonised gastric acid secretion induced by histamine, pentagastrin and bethanechol. Ranitidine was 5 to 10 times more active than cimetidine. However, both ranitidine and cimetidine had similar time curves of action. Ranitidine also inhibited the gastric secretory response to food in the conscious fistulated dog.

Ranitidine inhibited acid secretion in the perfused stomach of the anaesthetised rat, and acetylsalicylic acid-induced gastric lesion formation in the conscious rat, both in the presence and absence of excess hydrochloric acid. Measurements of the ratio of mucosal blood flow to acid secretion show that the inhibitory action of ranitidine upon gastric acid secretion cannot be attributed to changes in blood flow.

There were no behavioural effects in the mouse and rat after oral administration of 800 mg/kg ranitidine. Cats and dogs dosed with ranitidine 80 mg/kg orally, exhibited no behavioural effects indicative of an action on the central nervous system, although at this high dose level in the dog there was an indication of peripheral vasodilation and skin irritation due to released histamine.

Ranitidine, when co-administered with the following CNS modulating preparations; codeine, hexobarbitone, ethyl alcohol, chlordiazepoxide, chlorpromazine, imipramine, methyl dopa, reserpine, apomorphine or pentylenetetrazol, did not alter the pharmacological effects of either preparation.

At a dose level 45 times the antisecretory ED₅₀, intravenous infusion of ranitidine had no effect on the heart rate, blood pressure or electrocardiogram of the anaesthetised dog. The respiratory system was unaffected by ranitidine after oral doses in the mouse, rat, rabbit, cat and dog and after intravenous doses in the dog.

In the conscious dog, ranitidine had no appreciable effect on blood pressure or heart rate when administered orally at 10 mg/kg. There were short-lived falls in diastolic blood pressure after an intravenous dose of 10 mg/kg, 370 times the antisecretory dose level. There was no evidence of

arrhythmia nor of any electrocardiographic abnormality.

Long- term toxicity studies have shown that ranitidine does not possess antiandrogenic activity nor does it displace dihydrotestosterone from the androgen binding sites.

Metoclopramide, atropine and acetylsalicylic acid in the rat produced no change in the antisecretory activity of ranitidine.

The effect of ranitidine on anti-inflammatory drugs was varied. There was no effect on the anti-inflammatory action of prednisolone, but the anti-inflammatory action of indomethacin was enhanced. Administration of ranitidine reduced the frequency of acetylsalicylic acid- and indomethacin-induced gastric erosions. The antinociceptive action of acetylsalicylic acid was reduced after ranitidine treatment.

Ranitidine, unlike cimetidine, does not inhibit the hepatic mixed function oxygenase system. Spectral interaction studies have shown that whilst cimetidine binds strongly to cytochrome P₄₅₀, ranitidine has only weak affinity for this enzyme. Cimetidine is known to impair the metabolism of pentobarbitone and warfarin. In doses of up to 166 mg/kg in the rat, ranitidine had no effect on the pentobarbitone sleeping time or the pharmacokinetics and pharmacodynamics of warfarin.

Metabolism, Distribution and Excretion

The metabolism of ranitidine hydrochloride has been studied in four species of laboratory animal (mouse, rat, rabbit and dog) using radiolabelled drug. The drug was rapidly absorbed after oral administration. In the mouse, rat and rabbit between 30% and 60% of the administered radioactivity was excreted in the urine, the remainder being recovered in the faeces.

In the mouse 47% was excreted in the urine within 24 hours. In the rat, N-demethylation of ranitidine was the major route of metabolism. 30% of the administered dose was excreted in the urine as unchanged drug, up to 14% as desmethylranitidine, 3-6% as the N-oxide and 4% as the S-oxide. In rat bile the major radioactive components were ranitidine and an unidentified metabolite known as "Fast-Running Metabolite" (FRM) which is thought to be a charge transfer complex of ranitidine with bile pigments.

In the rabbit, sulphoxidation of ranitidine was the major route of metabolism, 18% of the administered dose being excreted in the urine as unmetabolised ranitidine, 8% as S-oxide, 2-4% as the N-oxide, and 2-4% as desmethylranitidine.

In the dog up to 70% of the administered dose was excreted in the first 24 hours. About 40% of the drug was excreted in the urine as unchanged ranitidine and up to 30% as the N-oxide, N-oxidation being the main route of metabolism of ranitidine in the dog. The N-oxide was also the major radioactive component present in dog bile together with small amounts of unchanged ranitidine and FRM.

In the rat, rabbit and dog, less than 10.1% of ranitidine in plasma is protein bound. Within one to seven days of administration of radio-labelled drug in the rat and dog over 99% of the radioactivity was cleared from the body. In common with many drugs, radioactivity persisted in the uveal tract of these two species, the half-life in the dog uveal tract being of the order of 6 months. Ranitidine and its S-oxide have greater affinity for melanin than the desmethyl metabolite; the N-oxide is bound only to a small extent.

The placental transfer of radioactive ranitidine and its metabolites has been studied in the pregnant rat and rabbit. Whole body autoradiography of rat and rabbit foetuses showed that small amounts of radioactivity were present in the uveal tract of the foetal eye in both species, in the gall bladder and intestine of the rabbit foetus and in the bladder of the rat foetus. Radioactivity was also detected in the salivary and mammary glands of the maternal rat and at very low concentration, in the milk.

Human Pharmacokinetics

Serum concentrations necessary to inhibit 50% of stimulated gastric acid secretion are estimated to be 36 to 94 ng/mL. Following a single oral dose of 150 mg, serum concentrations of ranitidine are in this range for up to 12 hours. There is a relationship between plasma concentrations of ranitidine and suppression of gastric acid production but wide interindividual variability exists.

Lower than prescription doses of ranitidine significantly reduce meal-stimulated intragastric acidity in human subjects. The effect is dose-related and acidity declines linearly with increasing doses of ranitidine.

Ranitidine is 50% absorbed after oral administration compared to an IV injection with mean peak levels of 440 to 545 ng/mL occurring two to three hours after a 150 mg dose. The elimination half-life is 1.5 to 3 hours.

The pharmacokinetics of lower doses of ranitidine has also been studied. Peak plasma concentrations and AUC showed a linear relationship over the dose range 20-80 mg; T_{max} , $t_{1/2}$ and clearance were independent of dose.

Ranitidine is absorbed very rapidly after an intramuscular injection. Mean peak levels of 576 ng/mL occur within 15 minutes or less following a 50 mg intramuscular dose. Absorption from intramuscular sites is virtually complete, with a bioavailability of 90% to 100% compared with intravenous administration.

The principal route of excretion is the urine, with approximately 30% of the orally administered dose collected in the urine as unchanged drug in 24 hours. Renal clearance is about 530 mL/min, indicating active tubular excretion, with a total clearance of 760 mL/min. The volume of distribution is 1.4 L/kg. Serum protein binding averages 15%.

Studies in patients with hepatic dysfunction (compensated cirrhosis) indicate that there are minor, but clinically insignificant alterations in ranitidine half life, distribution, clearance and bioavailability.

The gastric antisecretory activity of ranitidine metabolites has been examined. In man, both the principal metabolite in the urine, the N-oxide (4% of the dose) and the S-oxide (1%) possess weak H_2 -receptor blocking activity but desmethylranitidine (1 %) is only 4 times less potent than ranitidine in the rat and half as potent as ranitidine in the dog.

TOXICOLOGY

Animals

Toxicology, Impairment of Fertility, Carcinogenesis and Mutagenesis

Ranitidine hydrochloride has been subjected to exhaustive toxicological testing which has demonstrated the lack of any specific target organ or any special risk associated with its clinical use.

Acute Toxicity Studies

In mice and rats, the intravenous LD₅₀ is of the order of 75 mg/kg, whereas orally, even doses of 1000 mg/kg are not lethal. In dogs, the oral minimum lethal dose is 450 mg/kg/day. High single doses of ranitidine (up to 80 mg/kg orally) show only minimal and reversible signs of toxicity, some of which are related to transitory histamine releases.

Long-Term Toxicity Studies

In the long-term toxicity and carcinogenicity studies, very high doses of ranitidine were given daily to mice (up to 2000 mg/kg/day) throughout their normal life-span, and to dogs (up to 450 mg/kg/day) for periods of up to one year.

These doses produced massive plasma ranitidine concentrations far in excess of those found in human patients receiving ranitidine at the recommended therapeutic dose. For example, in the dogs, peak plasma concentrations were in excess of 115 mcg/mL and in mice basal plasma levels were in the range of 4-9 mcg/mL. In man, after oral administration of 150 mg ranitidine, the mean peak plasma concentration (C_{max}) was between 360 and 650 ng/mL.

In the rat, doses as high as 2000 mg/kg/day were well tolerated, the only morphological change seen was the increased incidence of accumulations of foamy alveolar macrophages in the lungs. The accumulations of these cells are natural phenomena in aging rats and chronic administration of a wide variety of drugs has been known to contribute to this process. Therefore, it is unlikely that the pharmacologic concentrations of ranitidine administered to these rats contributed to this natural process.

In the six-week and six-month oral studies in the dog (100 mg/kg/day) loose faeces were occasionally detected, while in the six-month study loose stools were accompanied on eight occasions by mucus-like material and sometimes by blood, mostly from one dog. Loose faeces, salivation and vomiting were observed in the 54-week dog study.

In isolated cases, dogs passed red-stained faeces which occasionally tested positive for occult blood. When the dose level was increased from 100 mg/kg/day to 225-450 mg/kg/day, no further red-stained faeces were seen, suggesting that any relationship to ranitidine is unlikely. Post-mortem examination of the dogs revealed no ranitidine-induced changes in the alimentary tract.

One dog had marginally raised levels of plasma alanine aminotransferase and alkaline phosphatase during the six-week study. This same dog also showed some necrotic foci in the liver. Small lesions of focal necrosis and fibrosis were also seen in one piece of liver from one female dog treated with 100 mg/kg for six months. No other differences were detected by light and electron microscopic examination of the treated and control livers. Since the focal lesions were seen in only one dog and were restricted to one piece of liver, it suggests that they were not caused by ranitidine.

Muscular tremors, an inability to stand, and rapid respiration were seen on occasion in dogs treated with 225 mg/kg/day in the 54-week study. The prevalence of these observations was increased when the dose was increased to a toxic level of 450 mg/kg/day. One dog died: no specific pathological changes or reason for the death was discovered.

Changes in the colour or granularity of the tapetum lucidum of the eye were detected in three dogs receiving the highest dose of ranitidine (450 mg/kg/day) during the 54-week study. In one dog this change was considered to be related to treatment. The change, a pallor of the tapetum, was reversible. No changes were seen with light or electron microscopic examination of the eye. The changes in the tapetum are of no clinical significance in humans since (i) humans do not have a tapetum lucidum and (ii) the changes were only seen at toxic pharmacological concentrations of ranitidine.

The mean serum glutamic pyruvic transaminase values for dogs treated at 450 mg/kg/day were significantly greater, albeit marginally, than the control values. These enzyme increases were not accompanied by any histological changes.

Studies in which ranitidine was administered parenterally were performed. No sign of specific local irritation attributable to ranitidine was detected. In the rat, no biochemical or histopathological changes were observed at intravenous dose levels as high as 20 mg/kg. Specifically, no significant changes were found in the veins or subcutis. Mild lesions in some muscle samples were observed: usually, the cells were basophilic and smaller than normal; and the nuclei were swollen, more numerous, and sometimes had migrated to the centre of the cell.

In the rabbit, slight infiltration of the pannicular muscle by mononuclear cells were noted. This minor subcutaneous reaction was uncommon and showed no group related distribution. There was no apparent difference in irritancy between ranitidine injection and placebo injection. In the rat, intravenous ranitidine at dose levels of 5.0 and 10.0 mg/kg daily for 15 days and 28 days produced no treatment related changes of biological importance in the haematopoietic system.

In Beagle dogs, intravenous ranitidine injection in doses up to 10 mg/kg/day for 28 and 42 days, produced no drug-related change in circulating erythrocytes or leukocytes and had no adverse effects on the haematopoietic system. No dose related changes were seen in electrocardiograms of Beagle dogs receiving up to 10 mg/kg ranitidine by intravenous injection. At dosage levels of up to 30 mg/kg, administered twice daily to Beagle dogs for 14 or 15 days, intravenous ranitidine injection produced no changes of biological significance in haematology, clinical chemistry or urinalysis.

No changes were observed in the eyes of dogs (specifically the tapetum lucidum) receiving ranitidine in doses up to 30 mg/kg twice daily for 15 days. At intravenous doses above 1.25 mg/kg, ranitidine injection produced immediate and transient reactions in the Beagle dog. The following reactions were typically produced by the administration of 1.25 mg/kg: bloodshot eyes, closing and watering of eyes, defaecation, diarrhoea, erythema, flatus, licking of lips, running nose, salivation, subdued behaviour, swallowing, tachycardia, and trembling. The range and severity of the effects was aggravated by increased dosage.

Reproduction Studies (Impairment of Fertility)

Reproduction studies were carried out in the rat and rabbit. Rats were exposed to ranitidine before and during mating, throughout pregnancy, lactation and during the weaning period. No effects on the reproductive process were seen and there was no evidence of an anti-androgenic effect.

A total of 2,297 foetuses from rats treated with ranitidine were examined. There was no evidence that ranitidine is a rat teratogen. Cleft palates occurred in foetuses from both treatment groups, however, there were significantly more in the control rat population.

A total of 944 foetuses from rabbits treated with ranitidine were examined; no drug-related adverse events or abnormalities in the foetuses were observed.

Rabbits receiving a bolus intravenous injection of ranitidine (10 mg/kg) once daily on gestation days 7-16 exhibited a reduction in weight gain. Their foetuses weighed significantly less than foetuses of untreated controls. In addition, 12.4% of ranitidine exposed foetuses had cleft palates. Reanalysis of this and a companion study performed to assess reproducibility demonstrated a lack of data reproducibility. Therefore, the effects observed in the first trial are aberrant, and should not form the basis for maternal or foetal toxicity.

In the subsequent study, no evidence of maternal or foetal toxicity was observed in rabbits dosed with 100 mg/kg ranitidine orally during days 2-29 of pregnancy. The peak plasma levels of ranitidine after a 100 mg/kg oral dose are similar to those obtained one minute after a 10 mg/kg dose administered intravenously (20-25 mcg/mL). Therefore, no teratogenic effects of ranitidine have been demonstrated at doses of 10 mg/kg (IV) and 100 mg/kg (Tablets) in rabbits.

Carcinogenicity Studies

There is no evidence that ranitidine is a carcinogen. Long term toxicity and carcinogenicity studies have involved the treatment of 600 mice and 636 rats at doses up to 2,000 mg/kg for two years and 129 weeks respectively and 42 dogs at doses up to 450 mg/kg/day for periods up to one year. These dose levels are far in excess of those to be used therapeutically in man. None of these animals had any intestinal metaplasia. There was no evidence of a tumourigenic effect of ranitidine in any other tissue.

Mutagenesis

Ranitidine is not mutagenic at doses as great as 30 mg/plate in the Ames Assay utilizing *Salmonella typhimurium* (TA 1538, TA 98, TA 100 and TA 1537) or in doses of 9 mg/plate utilizing *Escherichia Coli* (WP2 and WP2 uvrA) with or without activation.

Ranitidine at concentrations of 20-30 mg/plate had a weak direct mutagenic action in *S. Typhimurium* TA 1535 and at 9 mg/plate in *E. Coli* WP67. Ranitidine hydrochloride was not mutagenic at a concentration of 2 mg/mL in *E. Coli* or *S. Typhimurium* in the more sensitive Oral Solution microtitre fluctuation assay method. This weak direct mutagenic effect is of no clinical significance; the magnitudes of ranitidine concentration used in these assays are thousands of times greater than that attained therapeutically in human plasma.

The principal metabolites of ranitidine in man were not significantly mutagenic. This conclusion is supported by the following experiment. A test solution obtained by interacting ranitidine (10mM) and sodium nitrite (40mM) was mutagenic in *S. Typhimurium* (TA 1535) but not in *S. Typhimurium* (TA 1537) or in *E. Coli* (WP67 or WP2 uvrA). This positive result is attributable to the presence of a nitrosonitrolic acid derivative AH 23729, which was mutagenic. When the sodium nitrite concentration was reduced to 15mM or less, the solution was not mutagenic in any of the test microorganisms. The formation of AH 23729 requires concentrations of nitrous acid far in excess of those encountered in any probable physiological conditions. The other nitrosation products were not mutagenic in any of the microorganisms tested. There is no reason, therefore, for supposing that ranitidine is likely to be mutagenic in animals or man as a consequence of nitrosation in the stomach.

There is no evidence for long term toxicology, carcinogenicity and mutagenicity studies in animals to suggest that ranitidine is likely to have any deleterious effects in man when administered at therapeutic dose levels.

REFERENCES

1. Andersen M, Schou JS. Adverse reactions to H₂-receptor antagonists in Denmark before and after transfer of cimetidine and ranitidine to over-the-counter status. *Pharmacol Toxicol* 1991; 69:253-8.
2. Boehning W. Effect of cimetidine and ranitidine on plasma theophylline in patients with chronic obstructive airways disease treated with theophylline and corticosteroids. *Eur J Clin Pharmacol* 1990; 38:43-5.
3. Bye A, Lacey LF, Lettis S, Dixon JS, Felgate LA. Effect of ranitidine (150mg bd) on the pharmacokinetics of increasing doses of alcohol (0.15, 0.3, 0.6g/kg). *Am J Gastroenterol* 1993; 88:1590 A437.
4. Callaghan JT, Nyhart EH. Drug interactions between H₂-blockers and theophylline or warfarin. *Pharmacologist* 1988; 30:A14.
5. Das AF, Freston JW, Jacobs J, Fox NA, Morton RE. An evaluation of safety in 37, 252 patients treated with cimetidine or ranitidine. *Internal Medicine* 1990; 11:127-49.
6. Dent J, Dodds WJ, Friedman RH et al. Mechanism of gastroesophageal reflux in recumbent asymptomatic human subjects. *J Clin Invest* 1980; 65:256-67.
7. Desmond PV, Mashford ML, Harman PJ et al. Decreased oral warfarin clearance after ranitidine and cimetidine. *Clin Pharmacol Ther* 1984; 35:338-41.
8. Eandi M, De Carli GF, Recchia G, Colonna CV. Ranitidine: Eight years of post-marketing drug surveillance. *Post Marketing Surveillance* 1990; 4:1-8.
9. Furman D, Mensh R, Winan G et al. A double-blind trial comparing high dose liquid antacid to placebo and cimetidine in improving symptoms and objective parameters in gastroesophageal reflux. *Gastroenterol* 1982; 82:1062.
10. Garcia-Rodriguez LA, Jick H. Risk of gynaecomastia associated with cimetidine, omeprazole, and other anti-ulcer drugs. *Brit Med J* 1994; 308:503-6.
11. Graham DY, Patterson DJ. Double-blind comparison of liquid antacid and placebo in the treatment of symptomatic reflux oesophagitis. *Dig Dis Sci* 1983; 28:559-63.
12. Graham DY, Smith JL, Patterson DJ. Why do apparently healthy people use antacid tablets? *Am J Gastroenterol* 1983; 78:257-60.
13. Hansten PD. 1994 Drug Interaction of H₂-receptor antagonists.
14. Inman WAW. Drug Surveillance Research Unit. University of Southampton PEM. Prescription Event Monitoring News 1983; 1:9-13.

15. Grove O, Bekker C, Jeppe-Hansen MG et al. Ranitidine and high-dose antacid in reflux oesophagitis. *Scand J Gastroenterol* 1985; 20:457-61.
16. Johnsen R, Bernersen B, Straume B, Førde OH, Bostad L, Burhol PG. Prevalences of endoscopic and histological findings in subjects with and without dyspepsia. *Brit Med J* 1991; 302:749-52.
17. Johnson CD, Milward-Sadler GH, Jones R. Endoscopic findings in patients with heartburn who have not sought medical advice. *Hellenic J Gastroenterol* 1992; 5 (Suppl):141(A563).
18. Jones RH, Lydeard SE, Hobbs FDR et al. Dyspepsia in England and Scotland. *Gut* 1990; 31: 401-5.
19. Jones R, Lydeard S. Prevalence of symptoms of dyspepsia in the community. *Brit Med J* 1989; 298:30-2.
20. Kelly HW, Powell JR, Donohue JF. Ranitidine at very large doses does not inhibit theophylline elimination. *Clin Pharmacol Ther* 1986; 39:577-81.
21. Klauser AG, Schindlbeck NE, Müller-Lissner SA. Symptoms in gastro-oesophageal reflux disease. *Lancet* 1990; 335:205-8.
22. Koss MA, Hogan DL, Lane J, Steinbach JH, Isenberg JI. Anti-secretory effects and pharmacokinetics of low dose ranitidine. *Aliment Pharmacol Ther* 1993; 7:417-22.
23. Levitt MD. Lack of clinical significance of the interaction between H₂-receptor antagonists and ethanol. *Aliment Pharmacol Ther* 1993; 7:131-8.
24. Mills JG, Clancy A, Bond B et al. A comparison of the effects of cimetidine, ranitidine, omeprazole and placebo on the metabolism and clearance of racemic warfarin. *Br J Clin Pharmacol* 1986; 21: 566P-567P.
25. Mitchard M, Harris A, Mullinger BM. Ranitidine drug interactions: a literature review. *Pharmacol Ther* 1987; 32:293-325.
26. Müller-Lissner SA, Koch EMW, Geerke H. Uncomplicated gastro-oesophageal reflux disease. Symptomatic treatment with ranitidine. *Munch Med Wschr* 1992; 134:212-5.
27. Murdoch RH, Pappa KA, Giefer EE, Payne JE, Sanders M, Sirgo M. Endoscopic findings in a target population for over-the-counter treatment of heartburn. *Gastroenterology* 1994; 106(4, pt2):A146.
28. Nebel OT, Fornes MF, Castell DO. Symptomatic gastroesophageal reflux: incidence and precipitating factors. *Dig Dis* 1976; 21:953-6.

29. O'Reilly RA. Comparative interaction of cimetidine and ranitidine with racemic warfarin in man. *Arch Intern Med* 1984; 144:989-91.
30. Perrin VL. Safety evaluation of ranitidine in *The Clinical Use of Ranitidine - Proceedings of an International Symposium* Ed Wesdorp ICE Theracom 1982; pp48-54.
31. Ruth M, Månsson I, Sandberg N. The prevalence of symptoms suggestive of oesophageal disorders. *Scand J Gastroenterol* 1991; 26:73-81.
32. Serlin MJ, Sibeon RG, Breckenridge AM. Lack of effect of ranitidine on warfarin action. *Br J Clin Pharmacol* 1981; 12:791-4.
33. Smith SR, Kendall MJ. Ranitidine versus cimetidine. A comparison of their potential to cause clinically important drug interactions. *Clin Pharmacokinet* 1988; 15:44-56.
34. Sontag S, Robinson M, McCallum RW et al. Ranitidine therapy for gastro-oesophageal reflux disease. Results of a large double-blind trial. *Arch Int Med* 1987; 147:1485-92.
35. Toon S, Hopkins KJ, Garstang FM, Rowland M. Comparative effects of ranitidine and cimetidine on the pharmacokinetics and pharmacodynamics of warfarin in man. *Eur J Clin Pharmacol* 1987; 32:165-72.
36. Watts RW, Hetzel DJ, Bochner F et al. Lack of interaction between ranitidine and phenytoin. *Brit J Clin Pharmacol* 1983; 15:499-500.
37. Weberg R, Berstad A, Osnes M. Comparison of low-dose antacids, cimetidine and placebo on 24-hour intragastric acidity in healthy volunteers. *Dig Dis Sci* 1992; 37:1810-14.
38. Weberg R, Berstad A. Symptomatic effect of a low-dose antacid regimen in reflux esophagitis, *Scand J Gastroenterol* 1989; 24:401-6.
39. Wienbeck M, Berges W. Oesophageal disorders in the aetiology and pathophysiology of dyspepsia. *Scand J Gastroenterol* 1985; 20(suppl 109):133-43.
40. ZANTAC[®] Maximum Strength Non-Prescription Product Monograph, McNeil Consumer Healthcare, Date of revision: November 15, 2007; Control No 116984

Last revised July 10, 2017