



THE
REGULATORY & COMPLIANCE
LEADERS™

CANADIAN DRUG LABEL TEMPLATE

PRODUCT NAME	PRO-SENSE® ROUNDWORM LIQUID DEWORMER DOG
PRIMARY PACKAGE TYPE (i.e. bottle, tube)	Bottle

PRIMARY (INNER) LABEL (FINAL LABEL TO BE IN FRENCH & ENGLISH)

PRO-SENSE®

Roundworm Liquid Dewormer Dog

Piperazine (as citrate) Solution

FOR VETERINARY USE ONLY

Warnings: Keep out of reach of children.

Net Contents:

118 mL (4 fl. oz.)

DIN : XXXXXXXXX

Active Ingredients: 250 mg Piperazine base (as Piperazine Citrate) per teaspoon (5 mL)

Inactive Ingredients: Caramel Colour, Artificial Chocolate Flavour, Glycerin, Methylparaben, Potassium Sorbate, Sucrose, Vanillin, Water

Indications: This product is an anthelmintic to be given orally for use as an aid in the treatment and/or prevention of roundworms (*Toxocara canis* and *Toxascaris leonina*) in dogs and puppies 6 weeks of age or older.

Dosage and Administration: Consult your veterinarian in the diagnosis, treatment and control of parasitism.

Dogs and puppies (6 weeks or older): ½ teaspoon (tsp.) per 2.3 kg (5 lbs) of body weight. After initial



THE
REGULATORY & COMPLIANCE
LEADERS™

dose on day one, do not administer again until day 14. On day 14, repeat initial dose. Then administer every 30 days to help prevent re-infection. For puppies, administer dose every 14 days until 8 to 12 weeks of age, then monthly to 6 months of age, then every 30 days to help prevent re-infection. Treatment should be given shortly after pet has eaten or with food. Dosage may be given directly or added to food. For multiple pet households, give directly to individual dog or puppy to ensure proper dosage.

NOTE: A ½ tsp. is included for ease of administration.

Contra-Indications: Do not deworm a puppy under 6 weeks of age. Never deworm a sick dog or puppy. Not recommended for dogs under 2.3 kg (5 lbs) of body weight. Not recommended for pregnant or nursing females.

Cautions: Dogs and puppies 7 months of age or older should have a recent negative heartworm blood test before deworming. **Keep this product out of reach of animals to avoid unintended consumption.**

Adverse Reactions: Although uncommon at recommended doses, adverse reactions such as nausea, vomiting, diarrhea, ataxia or muscular tremors may occur. The recommended dosage should be followed carefully. If clinical signs occur, consult a veterinarian immediately.

Storage: Store at Room Temperature (15 – 30 °C). Keep bottle closed when not in use.

Manufactured by:
United Pet Group, Inc., a division of Spectrum Brands, Inc.
3001 Commerce Street, Blacksburg,
VA 24060 United States

Imported by:
Quality & Compliance Services Inc., Mississauga, Ontario L5N 5R1

Lot:

Exp:



THE
REGULATORY & COMPLIANCE
LEADERS™

CANADIAN DRUG LABEL TEMPLATE

PRODUCT NAME	PRO-SENSE® ROUNDWORM LIQUID DEWORMER DOG
SECONDARY PACKAGE TYPE (i.e. carton, shrink wrap)	Carton

SECONDARY (OUTER) LABEL (FINAL LABEL TO BE IN FRENCH & ENGLISH)

MAIN (FRONT) PANEL

Pro-Sense®
Roundworm Liquid Dewormer Dog

Piperazine (as citrate) Solution

FOR VETERINARY USE ONLY

Logos:

Pro-Sense®

Warnings: Keep out of reach of children.

Optional Marketing Language:
For the Treatment and Control of Roundworms
For Dogs and Puppies 6 Weeks and Older

DIN:XXXXXXXXX

Net Contents : 118 mL (4 fl. oz.)

ALTERNATE BOX PANEL

Optional Marketing Language:

- Easy to Administer Liquid Dewormer
- Provide ½ tsp. (included) per 2.3 kg (5 lbs) of Body Weight to Dog

ACTIVE INGREDIENTS: 250 mg Piperazine base (as Piperazine Citrate), per teaspoon (5 mL)

INACTIVE INGREDIENTS: Caramel Colour, Artificial Chocolate Flavour, Glycerin, Methylparaben, Potassium Sorbate, Sucrose, Vanillin, Water

INDICATIONS: This product is an anthelmintic to be given orally for use as an aid in the treatment and/or prevention of roundworms (*Toxocara canis* and *Toxascaris leonina*) in dogs and puppies 6 weeks of age or older.

DOSAGE AND ADMINISTRATION: Consult your veterinarian in the diagnosis, treatment and control of parasitism.

Dogs and puppies (6 weeks or older): ½ teaspoon (tsp.) per 2.3 kg (5 lbs) of body weight. After initial dose on day one, do not administer again until day 14. On day 14, repeat initial dose. Then administer every 30 days to help prevent re-infection. For puppies, administer dose every 14 days until 8 to 12 weeks of age, then monthly to 6 months of age, then every 30 days to help prevent re-infection. Treatment should be given shortly after pet has eaten or with food. Dosage may be given directly or added to food. For multiple pet households, give directly to individual dog or puppy to ensure proper dosage.

NOTE: A ½ tsp. is included for ease of administration.

Pro-Sense Roundworm Liquid Dewormer Dog solution is an easy to administer liquid dewormer. Simply follow the steps below to treat your dog or puppy (6 weeks or older).

[Infographic]

STEP-BY-STEP GUIDE

STEP 1

- Weigh yourself
- Weigh you and your pet
- Subtract your weight to get pet's weight

STEP 2

- If weight is in kilograms (kg), divide pet weight by 2.3 kg.
 - Example: if pet's weight is 5 kg: $5 \text{ kg} \div 2.3 \text{ kg} = 2 =$ number of ½ teaspoons to give
- If weight is in pounds (lbs), divide pet weight by 5 lbs.
 - Example: if pet's weight is 10 lbs: $10 \text{ lbs} \div 5 \text{ lbs} = 2 =$ number of ½ teaspoons to give
- The table below is provided as a guide:

Pet's weight	Number of ½ teaspoons (included) to give
2.3 kg (5 lbs)	1
4.6 kg (10 lbs)	2
6.9 kg (15 lbs)	3
9.2 kg (20 lbs)	4
11.5kg (25 lbs)	5
13.8 kg (30 lbs)	6
18.4 kg (40 lbs)	8
23 kg (50 lbs)	10
27.6 kg (60 lbs)	12

STEP 3

- Administer the appropriate amount of ½ teaspoon (tsp.) per 2.3 kg (5 lbs) of body weight to dog utilizing provided ½ teaspoon (tsp.)

STEP 4

- Mix dose into food or give directly

STEP 5

- Repeat 14 days later

STEP 6

- Repeat every 30 days to help prevent re-infestation

CONTRA-INDICATIONS: Do not deworm a puppy under 6 weeks of age. Never deworm a sick dog or puppy. Not recommended for dogs under 2.3 kg (5 lbs) of body weight. Not recommended for pregnant or nursing females.

CAUTIONS: Dogs and puppies 7 months of age or older should have a recent negative heartworm blood test before deworming. **KEEP THIS PRODUCT OUT OF THE REACH OF ANIMALS TO AVOID UNINTENDED CONSUMPTION.**

ADVERSE REACTIONS: Although uncommon at recommended doses, adverse reactions such as nausea, vomiting, diarrhea, ataxia or muscular tremors may occur. The recommended dosage should be followed carefully. If clinical signs occur, consult a veterinarian immediately.

STORAGE: Store between Room Temperature (15 – 30 °C). Keep bottle closed when not in use.

ALTERNATE BOX PANEL



THE
REGULATORY & COMPLIANCE
LEADERS™

QUESTIONS OR COMMENTS?

Call 1-800-645-5154 or visit our website at www.prosensepet.com

Manufactured by:

United Pet Group, Inc., a division of Spectrum Brands, Inc.

3001 Commerce Street, Blacksburg, VA 24060 United States

Item # [insert item number here]

® Registered Trademark of Spectrum Brands, Inc.

© [insert year] Spectrum Brands, Inc.

Imported by: Quality & Compliance Services Inc., Mississauga, Ontario L5N 5R1

SKU: J1715FL

LOT:

EXP.: