


C26500

DIN 02228386

VETERINARY USE ONLY

CYDECTIN®
**Moxidectin
(Pour-On for Beef and Dairy Cattle)**

(Each mL contains 5 mg of moxidectin)

For the treatment of infections and infestations due to internal and external parasites in cattle, including lactating dairy cows.

FOR EXTERNAL USE ONLY

INTRODUCTION

CYDECTIN® Pour-On contains moxidectin, a second generation endectocide of the milbemycin class of compounds. It is highly effective when administered at a dosage of 0.5 mg/kg body weight against a broad range of economically important endoparasites and ectoparasites that infect beef and dairy cattle, including lactating dairy cows.

Mode of action studies indicate that CYDECTIN® eliminates parasites by causing paralysis through interfering with neurotransmission of the gamma-aminobutyric acid (GABA)-gated or glutamate-gated chloride channels.

COMPOSITION

CYDECTIN® Pour-On comes ready to use as a violet solution containing 5 mg/mL (0.5% w/v) moxidectin.

INDICATIONS

CYDECTIN® Pour-On is indicated for the treatment of healthy cattle of all ages and breeds including pregnant cows, breeding bulls and lactating dairy cows, which are parasitized by the following gastrointestinal roundworms (including inhibited larvae of *Ostertagia ostertagi*), lungworms, grubs, mange mites and biting and sucking lice.

GASTROINTESTINAL ROUNDWORMS

<i>Ostertagia ostertagi</i>	Adults, Developing L ₄ , Inhibited L ₄
<i>Haemonchus placei</i>	Adults, Developing L ₄
<i>Trichostrongylus axei</i>	Adults, Developing L ₄
<i>Trichostrongylus colubriformis</i>	Adults, Developing L ₄
<i>Cooperia oncophora</i>	Adults, Developing L ₄
<i>Cooperia pectinata</i>	Adults
<i>Cooperia punctata</i>	Adults, Developing L ₄
<i>Cooperia surrabadana</i>	Adults, Developing L ₄
<i>Bunostomum phlebotomum</i>	Adults
<i>Oesophagostomum radiatum</i>	Adults, Developing L ₄
<i>Nematodirus helvetianus</i>	Adults, Developing L ₄
<i>Trichuris discolor</i>	Adults, Developing L ₄

LUNGWORMS

<i>Dictyocaulus viviparus</i>	Adults, Developing L ₄
-------------------------------	-----------------------------------

MANGE MITES

<i>Chorioptes bovis</i>
<i>Psoroptes ovis</i> (<i>Psoroptes communis</i> var. <i>bovis</i>)
<i>Sarcoptes bovis</i>

BITING AND SUCKING LICE

<i>Linognathus vituli</i>
<i>Haematopinus eurysternus</i>
<i>Solenopotes capillatus</i>
<i>Damalinea (Bovicola) bovis</i>

GRUBS (Migrating Larvae)

<i>Hypoderma lineatum</i>
<i>Hypoderma bovis</i>

Persistent Activity: CYDECTIN® Pour-On prevents reinfection with *Haemonchus placei* for 14 days, *Oesophagostomum radiatum*, and *Ostertagia ostertagi* for at least 28 days and *Dictyocaulus viviparus* for at least 42 days following a single application at the recommended dose.

Research data, based on faecal nematode egg output (EPG counts), lice counts and controlled rainfall simulation, indicate that the application of the pour-on formulation to wet skin or application prior to a typical rain event does not affect the efficacy of the product.

DOSAGE AND ADMINISTRATION

Dosage and administration: EXTERNAL USE ONLY. CYDECTIN® Pour-On should be applied to healthy skin along the top of the back from the withers to the base of the tail. Use 1 mL per 10 kg body weight to administer a dose of 0.5 mg/kg.

CYDECTIN® Pour-On comes ready to use. A number of different applicators have been tested with CYDECTIN® Pour-On and several have proved suitable. Use of applicators is recommended when using the 2.5, 5 and 10 liter sizes and when any large group of animals is to be treated.

DOSAGE TABLE

Body weight (kg)	Dose (mL)	Doses per pack				Body weight (kg)	Dose (mL)	Doses per pack			
		1 L	2.5 L	5 L	10 L			1 L	2.5 L	5 L	10 L
40	4	250	625	1250	2500	250	25	40	100	200	400
50	5	200	500	1000	2000	300	30	33	83	166	332
60	6	166	416	833	1666	350	35	28	71	142	284
70	7	142	357	714	1428	400	40	25	62	125	250
80	8	125	312	625	1250	450	45	22	55	111	222
90	9	111	277	555	1110	500	50	20	50	100	200
100	10	100	250	500	1000	550	55	16	45	90	180
110	11	90	227	454	908	600	60	16	41	83	166
120	12	83	208	416	832	650	65	15	38	76	152
130	13	76	192	384	768	700	70	14	35	70	140
140	14	71	178	357	714	750	75	13	33	66	132
150	15	66	166	333	666	800	80	12	31	62	124
200	20	50	125	250	500	850	85	11	29	58	116

If the animals are likely to be reinfected, a strategic parasite control program should be established.

SAFETY

CYDECTIN® Pour-On has a very broad spectrum of activity and wide safety margin of 10X the recommended dosage of 500 µg moxidectin per kg of body weight. This safety margin is expected to be valid for cattle of most breeds, types and ages. It can be safely administered to breeding animals or replacement animals kept for breeding purposes.

CYDECTIN® Pour-On does not cause irritation to the skin or hide damage when used as directed and is not harmful to the eyes of cattle.

SUPPLIED

CYDECTIN® Pour-On is supplied in 1 L, 2.5 L, 5 L and 10 L plastic bottles.

STORAGE CONDITIONS

CYDECTIN® Pour-On should be stored between 4-25°C. AVOID FREEZING. Protect from light. Store bottle in box.

CAUTIONS

- Do not apply at the same time as any other pour-on product.
- Apply to healthy skin. Avoid applying to mange scabs, lesions or foreign matter.
- Dispose of any containers and any residual content safely (e.g., bury or incinerate). Bury away from water courses.
- For external application only. Do not administer orally or parenterally to animals.
- To avoid secondary reactions when treating *Hypoderma* spp. infections, consult your veterinarian on the correct timing of treatment.

WARNINGS

- Treated cattle must not be slaughtered for use in food for at least 15 days after the latest treatment with this drug.
- Do not use in milk-fed calves. Do not use in calves less than 14 weeks of age to be processed for veal.
- Keep this and all drugs out of reach of children.
- Avoid direct contact with skin and eyes.

NOTE TO USER – When used according to label directions, no milk withholding time is required following the application of CYDECTIN® Pour-On on dairy cattle.

12170
C26500

Boehringer Ingelheim (Canada) Ltd.
5180 South Service Road
Burlington, Ontario L7L 5H4

Boehringer Ingelheim

Revision 2
Operator Ish
Date 05Aug10
Product ID#... C26500
Resource #... I26500
Product Cydectin
Type/Size Insert
Template # ... 594
Dimensions .. 8-1/2" x 11"
File Path V1/PF/BI Labeling/C26500
PDF 06Aug10 hal
Rev Level 0

BLACK

Boehringer Ingelheim	Revision 1	Template # ... 181
	Operator Ish	Dimensions .. 2-1/2" x 3-1/2" <i>(special shape)</i>
	Date 04Aug10	File Path V1/PF/BI Labeling/ C2653L
	Product ID# . C2653L	PDF 05Aug10 jaf
	Resource # .. L2653C	Rev Level L
	Product Cydectin	
Type/Size ... Label (front) 1L		

- BLACK
- PMS 1795
- UNCOL GREY
- PMS 322
- VARNISH

Click here with mover tool to bring varnish plate to front

SEPARATIONS
CHECKED by
hal/26Jan10

DIN 02228386

VETERINARY USE ONLY

CYDECTIN®

Moxidectin
(Pour-On for Beef and Dairy Cattle)

(Each mL contains 5 mg of moxidectin)

For the treatment of infections and infestations due to internal and external parasites in cattle, including lactating dairy cows.

WARNINGS

1. Treated cattle must not be slaughtered for use in food for at least 10 days after the latest treatment with this drug.
2. Do not use in milk-fed calves. Do not use in calves less than 14 weeks of age to be processed for veal.
3. Keep this and all drugs out of reach of children.
4. Avoid direct contact with skin and eyes.

NOTE TO USER - When used according to label directions, no milk withholding time is required following the application of CYDECTIN Pour-On on dairy cattle.

See package insert for complete information on dosage and administration.



EXTERNAL USE ONLY

Store between 4-25°C. **AVOID FREEZING.**
Protect from light. Store below 10 mm.

Dosage and administration: EXTERNAL USE ONLY: CYDECTIN® Pour-On should be applied to healthy skin along the top of the back from the withers to the base of the tail. Use 1 mL per 10 kg bodyweight to administer a dose of 0.5 mg/kg.

Net Contents
1 L

Boehringer Ingelheim (Canada) Ltd.
5180 South Service Road
Burlington, Ontario L7L 5H4



**Boehringer
Ingelheim**

Lot: Exp:

785922

Boehringer Ingelheim	Revision 1	Type/Size Label (front) 2.5L
	Operator lsh	Template # ... 182
	Date 04Aug10	Dimensions .. 6" x 5.592"
	Product ID# . C2655L	File Path V1/PF/BI Labeling/ C2655L
	Resource #.. L2655C	PDF..... 05Aug10 jaf
	Product Cydectin	Rev Level L

- BLACK**
- PMS 1795**
- COOL GREY 3**
- PMS 322**
- VARNISH**

SEPARATIONS
CHECKED
by hal / 26Jan10

DIN 02228386

**VETERINARY
USE ONLY**

CYDECTIN®

Moxidectin (Pour-On for Beef and Dairy Cattle)

(Each mL contains 5 mg of moxidectin)

For the treatment of infections and infestations due to internal and external parasites in cattle, including lactating dairy cows.

EXTERNAL USE ONLY

WARNINGS:

1. Treated cattle must not be slaughtered for use in food for at least 15 days after the latest treatment with this drug.
2. Do not use in milk-fed calves. Do not use in calves less than 14 weeks of age to be processed for veal.
3. Keep this and all drugs out of reach of children.
4. Avoid direct contact with skin and eyes.

NOTE TO USER – When used according to label directions, no milk withholding time is required following the application of CYDECTIN Pour-on on dairy cattle.

Dosage and administration: EXTERNAL USE ONLY. CYDECTIN® Pour-On should be applied to healthy skin along the top of the back from the withers to the base of the tail. Use 1 mL per 10 kg bodyweight to administer a dose of 0.5 mg/kg.

See package insert for complete information on dosage and administration.

Store between 4-25°C. AVOID FREEZING.
Protect from light. Store bottle in box.

Boehringer Ingelheim (Canada) Ltd.
5180 South Service Road
Burlington, Ontario L7L 5H4



Net Contents 2.5 L



**Boehringer
Ingelheim**

Lot: C2655L
 Exp.: 12/160

Boehringer
Ingelheim

Revision 1
Operator Ish
Date 04Aug10
Product ID# . C2657K
Resource # . . L2657C
Product Cydectin
Type/Size ... Label (front) 5L

Template # ... 155
Dimensions .. 7.059" x 6.579"
File Path..... V1/PF/BI Labeling/
C2657K
PDF..... 05Aug10 jaf
Rev Level..... K

BLACK

PMS 1795

COOL GREY 3

PMS 322

VARNISH

SEPARATIONS
CHECKED
by hal / 26Jan10

DIN 02228386

**VETERINARY
USE ONLY**

CYDECTIN[®]

**Moxidectin (Pour-On
for Beef and Dairy Cattle)**

(Each mL contains 5 mg of moxidectin)

For the treatment of infections and infestations due to internal and
external parasites in cattle, including lactating dairy cows.

EXTERNAL USE ONLY

WARNINGS:

1. Treated cattle must not be slaughtered for use in food for at least 15 days after the latest treatment with this drug.
2. Do not use in milk-fed calves. Do not use in calves less than 14 weeks of age to be processed for veal.
3. Keep this and all drugs out of reach of children.
4. Avoid direct contact with skin and eyes.

NOTE TO USER – When used according to label directions, no milk withholding time is required following the application of
CYDECTIN Pour-On on dairy cattle.

Dosage and administration: EXTERNAL USE ONLY. CYDECTIN[®]
Pour-On should be applied to healthy skin along the top of the
back from the withers to the base of the tail. Use 1 mL per 10 kg
bodyweight to administer a dose of 0.5 mg/kg.

See package insert for complete information on dosage and
administration.

Store between 4-25°C. AVOID FREEZING.
Protect from light. Store bottle in box.

Boehringer Ingelheim (Canada) Ltd.
5180 South Service Road
Burlington, Ontario L7L 5H4



Net Contents 5 L



Boehringer
Ingelheim

12160

C2657K

Lot:
Exp.:

Boehringer
Ingelheim

Revision 1
Operator Ish
Date 04Aug10
Product ID# . C2659M
Resource # . L2659C
Product Cydectin
Type/Size ... Label / 10L

Template # ... 131
Dimensions .. 5-3/4" x 7-1/4"
File Path V1/PF/BI Labeling/
C2659M
PDF 06Aug10 hal
Rev Level M

BLACK

PMS 1795

COOL GREY 3

PMS 322

VARNISH

SEPARATIONS
CHECKED
by hal / 26jan10

Click here with mover tool to bring varnish plate to front

DIN 02228386

VETERINARY USE ONLY / USAGE VÉTÉRINAIRE SEULEMENT

CYDECTIN[®]

Moxidectin (Pour-On for Beef and Dairy Cattle) Moxidectin (Solution à verser pour bovins de boucherie et bovins laitiers)

(Each mL contains 5 mg of moxidectin) / (Chaque mL renferme 5 mg de moxidectin)

For the treatment of infections and infestations due to internal and external parasites in cattle, including lactating dairy cows.

Pour le traitement des infections et des infestations attribuables aux parasites internes et externes chez les bovins, y inclus les vaches laitières en lactation.

EXTERNAL USE ONLY / USAGE EXTERNE SEULEMENT

WARNINGS:

1. Treated cattle must not be slaughtered for use in food for at least 15 days after the latest treatment with this drug.
2. Do not use in milk-fed calves. Do not use in calves less than 14 weeks of age to be processed for veal.
3. Keep this and all drugs out of reach of children.
4. Avoid direct contact with skin and eyes.

NOTE TO USER – When used according to label directions, no milk withholding time is required following the application of CYDECTIN Pour-On on dairy cattle.

Dosage and administration: EXTERNAL USE ONLY. CYDECTIN[®] Pour-On should be applied to healthy skin along the top of the back from the withers to the base of the tail. Use 1 mL per 10 kg body weight to administer a dose of 0.5 mg/kg.

See package insert for complete information on dosage and administration.
Store between 4-25°C. AVOID FREEZING.
Protect from light. Store bottle in box.

Boehringer Ingelheim (Canada) Ltd.
5180 South Service Road
Burlington, Ontario L7L 5H4



MISES EN GARDE :

1. Les animaux traités ne doivent pas être abattus à des fins alimentaires dans un délai d'au moins 15 jours après le dernier traitement à l'aide de ce médicament.
2. Ne pas utiliser chez les veaux de lait. Ne pas utiliser chez les veaux de moins de 14 semaines d'âge élevés pour la production de viande de veau.
3. Garder ce produit, ainsi que tout autre médicament de la portée des enfants.
4. Éviter le contact direct avec la peau ou les yeux.

AVIS À L'UTILISATEUR – Lorsque le produit est utilisé conformément aux directives de l'étiquette, aucune période d'attente pour l'utilisation du lait n'est requise à la suite de l'administration de la solution à verser CYDECTIN à 0.5% chez les bovins laitiers.

Posologie et administration : La solution à verser CYDECTIN[®], à usage externe seulement, devrait être appliquée sur une peau saine, le long du dos, du garrot jusqu'à l'attache de la queue. Utiliser 1 mL par 10 kg de poids vif pour administrer une dose de 0,5 mg/kg.

Voir le dépliant pour obtenir toutes les informations sur la posologie et l'administration.

Garder entre 4 et 25 °C. ÉVITER LE GEL.
Garder à l'abri de la lumière. Entreposer la bouteille dans la boîte.

Boehringer Ingelheim (Canada) Ltée
5180 South Service Road
Burlington (Ontario) L7L 5H4

Net Contents/Contenu Net
10 L



Boehringer
Ingelheim

Lot:
Exp. Date:

C2659M

12/60

Boehringer
Ingelheim

Revision 1
Operator lsh
Date 05Aug10
Product ID# .. CC2653L
Resource # ... 8548
Product Cydectin
Type/Size..... Carton / 1 L

Template # ... 758
Dimensions ... 5-3/8" x 2-3/4" x 9-19/32"
File Path V1/PF/BI Labeling/
CC2653L
PDF 06Aug10 hal
Rev Level L

BLACK
PMS 1795
CORN PRNT 2
PMS 322
PMS 488888001

SEPARATIONS
CHECKED BY
hal/26Jan10

50265320

Moxidectin
(Solution à verser
pour bovins de
boucherie et
bovins laitiers)



Moxidectin
(Pour-On for
Beef and
Dairy Cattle)

CYDECTIN®

DIN 02228386

USAGE VÉTÉRINAIRE SEULEMENT

CYDECTIN®

**Moxidectin
(Solution à verser pour
bovins de boucherie et
bovins laitiers)**

(Chaque mL renferme 5 mg de moxidectin)

Pour le traitement des infections et des infestations attribuables
aux parasites internes et externes chez les bovins,
y inclus les vaches laitières en lactation.

USAGE EXTERNE SEULEMENT

MISES EN GARDE :

1. Les animaux traités ne doivent pas être abattus à des fins alimentaires dans un délai d'au moins 15 jours après le dernier traitement à l'aide de ce médicament.
2. Ne pas utiliser chez les veaux de lait. Ne pas utiliser chez les veaux de moins de 14 semaines d'âge élevés pour la production de viande de veau.
3. Garder ce produit, ainsi que tout autre médicament de la portée des enfants.
4. Éviter le contact direct avec la peau ou les yeux.

AVIS À L'UTILISATEUR - Lorsque le produit est utilisé conformément aux directives de l'étiquette, aucune période d'attente pour l'utilisation du lait n'est requise à la suite de l'administration de la solution à verser CYDECTIN à 0,5% chez les bovins laitiers.

Garder entre 4 et 25 °C. ÉVITER LE GEL.
Garder à l'abri de la lumière.
Entreposer la bouteille dans la boîte.

Boehringer Ingelheim (Canada) Ltée
5180 South Service Road
Burlington (Ontario) L7L 5H4



Boehringer
Ingelheim

Contenu Net
1 L

Posologie et administration : La solution à verser CYDECTIN® à usage externe seulement, devrait être appliquée sur une peau saine, le long du dos, du grand pectoral à l'arrière de la queue. Utiliser 1 mL par 10 kg de poids vif pour administrer une dose de 0,5 mg/kg.

TABLÉAU POSOLOGIQUE

Poids vif (kg)	Dose par race		Poids vif (kg)	Dose par race	
	(mL)	(1.1)		(mL)	(1.1)
40	4	250	250	25	40
50	5	250	300	30	33
60	6	156	250	25	29
70	7	142	400	40	25
80	8	125	400	40	22
90	9	111	500	50	20
100	10	100	550	55	18
110	11	90	600	60	16
120	12	83	650	65	15
130	13	76	700	70	14
140	14	71	750	75	13
150	15	66	800	80	12
200	20	50	850	85	11

Voir le dépliant à l'intérieur pour obtenir toutes les informations sur le posologie et l'administration.

Voir le dessous ou le côté pour le numéro de lot et la date de péremption.



Net Contents/Contenu Net
1 L

VETERINARY USE ONLY

DIN 02228386

CYDECTIN®

**Moxidectin
(Pour-On for Beef
and Dairy Cattle)**

(Each mL contains 5 mg of moxidectin)

For the treatment of infections and infestations due
to internal and external parasites in cattle,
including lactating dairy cows.

EXTERNAL USE ONLY

WARNINGS:

1. Treated cattle must not be slaughtered for use in food for at least 15 days after the latest treatment with this drug.
2. Do not use in milk-fed calves. Do not use in calves less than 14 weeks of age to be processed for veal.
3. Keep this and all drugs out of reach of children.
4. Avoid direct contact with skin and eyes.

NOTE TO USER - When used according to label directions, no milk withholding time is required following the application of CYDECTIN Pour-On on dairy cattle.

Store between 4-25°C. AVOID FREEZING.
Protect from light. Store bottle in box.

Boehringer Ingelheim (Canada) Ltd.
5180 South Service Road
Burlington, Ontario L7L 5H4



Net Contents
1 L

Boehringer
Ingelheim

Dosage and administration: EXTERNAL USE ONLY. CYDECTIN® Pour-On should be applied to healthy skin along the top of the back, from the withers to the base of the tail. Use 1 mL per 10 kg body weight to administer a dose of 0.5 mg/kg.

DOSAGE TABLE

Body weight (kg)	Dose per head		Body weight (kg)	Dose per head	
	(mL)	(1.1)		(mL)	(1.1)
40	4	250	250	25	40
50	5	200	300	30	33
60	6	166	350	35	28
70	7	142	400	40	25
80	8	125	450	45	22
90	9	111	500	50	20
100	10	100	550	55	18
110	11	90	600	60	16
120	12	83	650	65	15
130	13	76	700	70	14
140	14	71	750	75	13
150	15	66	800	80	12
200	20	50	850	85	11

See package insert for complete information on dosage and administration.

See bottom or side for lot number and expiration date.



8548
12170
CC2653L
8548
CC2653L
8548
CC2653L

Boehringer Ingelheim
 Part No. 1
 Description 04kg/10
 Product ID# CC2855L
 Revision # 04/0
 Product Cydectin
 Type/Use Cattle / 2.5 L

Template # 782
 Dimensions - 8.18" x 3.91"
 x 11.98"
 File Path Y:\OFFICE Labeling
 CC2855L
 Product Code 08kg/10 ml
 Rev Level L

BLACK
 8549 12160
 PMS 222
 8549 12160

SEPARATIONS
 CHECKED BY
 HAVZJaa10

Moxidectin
 (Solution à verser pour bovins de
 boucherie et bovins laitiers)



Moxidectin
 (Pour-On
 for
 Beef and
 Dairy Cattle)

CYDECTIN®

USAGE VÉTÉRINAIRE SEULEMENT

DIN 02228386

CYDECTIN®

Moxidectin
(Solution à verser pour bovins
de boucherie et bovins laitiers)

(Chaque mL renferme 5 mg de moxidectin)

Pour le traitement des infections et des infestations attribuables
 aux parasites internes et externes chez les bovins, y inclus les
 vaches laitières en lactation.

USAGE EXTERNE SEULEMENT

MISES EN GARDE :

1. Les animaux traités ne doivent pas être abattus à des fins alimentaires dans un délai d'au moins 15 jours après le dernier traitement à l'aide de ce médicament.
2. Ne pas utiliser chez les veaux de lait. Ne pas utiliser chez les veaux de moins de 14 semaines d'âge élevés pour la production de viande de veau.
3. Garder ce produit, ainsi que tout autre médicament de la portée des enfants.
4. Éviter le contact direct avec la peau ou les yeux.

AVIS À L'UTILISATEUR - Lorsque le produit est utilisé conformément aux directives de
 l'étiquette, aucune période d'attente pour l'utilisation du lait n'est requise à la suite de
 l'administration de la solution à verser CYDECTIN à 0,5% chez les bovins laitiers.

Garder entre 4 et 25 °C. ÉVITER LE GEL.
 Garder à l'abri de la lumière. Entreposer la bouteille dans la boîte.

Boehringer Ingelheim (Canada) Ltée
 5180 South Service Road
 Burlington (Ontario) L7L 5H4



Boehringer
 Ingelheim

Contenu Net
 2,5 L

CYDECTIN®

Moxidectin
(Pour-On for
Beef and Dairy Cattle)

Moxidectin
(Pour-On for
Beef and Dairy Cattle)

(Each mL contains 5 mg of moxidectin)

For the treatment of infections and infestations due to
 internal and external parasites in cattle,
 including lactating dairy cows.

EXTERNAL USE ONLY

WARNINGS:

1. Treated cattle must not be slaughtered for use in food for at least 15 days after the latest treatment with this drug.
2. Do not use in milk-fed calves. Do not use in calves less than 14 weeks of age to be processed for veal.
3. Keep this and all drugs out of reach of children.
4. Avoid direct contact with skin and eyes.

NOTE TO USER - When used according to label directions, no milk withholding time is required following the
 application of CYDECTIN Pour-On on dairy cattle.

Store between 4-25°C. AVOID FREEZING.
 Protect from light. Store bottle in box.

Boehringer Ingelheim (Canada) Ltd.
 5180 South Service Road
 Burlington, Ontario L7L 5H4



Boehringer
 Ingelheim

Net Contents
 2.5 L

Net Contents/Contenu Net
 2,5 L

Posologie et administration : La solution à verser CYDECTIN®
 à usage externe seulement, devrait être appliquée sur une peau
 saine, le long du dos, du garrot jusqu'à l'attache de la queue.
 Utiliser 1 mL par 10 kg de poids et pour administrer une dose
 de 0,5 mg/kg.

DOSAGE TABLE

Poids vil (kg)	Dose (mL)	Dose façon (2,5 L)	Poids vil (kg)	Dose (mL)	Dose façon (2,5 L)
45	4	625	25	25	100
55	5	500	30	30	83
65	6	416	35	35	71
75	7	357	40	40	62
85	8	312	45	45	55
95	9	277	50	50	50
105	10	250	55	55	45
115	11	227	60	60	41
125	12	208	65	65	38
135	13	192	70	70	35
145	14	178	75	75	33
155	15	166	80	80	31
200	20	125	85	85	28

Voir le séquent à l'extérieur pour obtenir toutes les informations sur la
 posologie et l'administration.
 Voir le dessous ou le côté pour le numéro de lot et la date de péremption.

Dosage and administration: EXTERNAL USE ONLY
 CYDECTIN® Pour-On should be applied to healthy skin along
 the top of the back from the withers to the base of the tail.
 Use 1 mL per 10 kg body weight to administer a dose of
 0.5 mg/kg.

DOSAGE TABLE

Body weight (kg)	Dose (mL)	Dose per pack (2.5 L)	Body weight (kg)	Dose (mL)	Dose per pack (2.5 L)
45	4	625	25	25	100
55	5	500	30	30	83
65	6	416	35	35	71
75	7	357	40	40	62
85	8	312	45	45	55
95	9	277	50	50	50
105	10	250	55	55	45
115	11	227	60	60	41
125	12	208	65	65	38
135	13	192	70	70	35
145	14	178	75	75	33
155	15	166	80	80	31
200	20	125	85	85	28

See package insert for complete information on dosage and administration.

See bottom or side for lot number and expiration date.

8549
 12160
 CC2855L
 8549
 CC2855L
 8549
 CC2855L

Boehringer Ingelheim
 Revlon 2
 Couleur Ind
 Date 05Aug10
 Product ID# CC2657L
 Revision # 0504
 Product Cydectin
 Type/Size Carton / 5 L
 Template # 773
 Dimensions 9.811" x 5.813" x 13.157"
 File Path V:\PDS Labeling
 CC2657L
 PDF 11Aug10 Ind
 Rev Level L
 BLACK
 PMS 1795
 PMS 322
 SEPARATIONS CHECKED by
 10/27/10

Moxidectin (Solution à verser pour bovins de boucherie et bovins laitiers)



Moxidectin (Pour-On for Beef and Dairy Cattle)

CYDECTIN

DIN 02228386

USAGE VÉTÉRINAIRE SEULEMENT

CYDECTIN

Moxidectin
(Solution à verser pour bovins de boucherie et bovins laitiers)

(Chaque mL renferme 5 mg de moxidectin)

Pour le traitement des infections et des infestations attribuables aux parasites internes et externes chez les bovins, y inclus les vaches laitières en lactation.

USAGE EXTERNE SEULEMENT

MISES EN GARDE :

1. Les animaux traités ne doivent pas être abattus à des fins alimentaires dans un délai d'au moins 15 jours après le dernier traitement à l'aide de ce médicament.
2. Ne pas utiliser chez les veaux de lait. Ne pas utiliser chez les veaux de moins de 14 semaines d'âge élevés pour la production de viande de veau.
3. Garder le produit, ainsi que tout autre médicament de la portée des enfants.
4. Éviter le contact direct avec le nez ou les yeux.


AVIS À L'UTILISATEUR - Lorsque le produit est utilisé conformément aux directives de l'étiquette, aucune période d'attente pour l'utilisation du lait n'est requise à la suite de l'administration de la solution à verser CYDECTIN à 0,5% chez les bovins laitiers.

Garder entre 4 et 25 °C. ÉVITER LE GEL.
Garder à l'abri de la lumière. Entreposer la bouteille dans la boîte.

Boehringer Ingelheim (Canada) Ltée
5180 South Service Road
Burlington (Ontario) L7L 5H4



Contenu Net
5 L



Net Contents/Contenu Net
5 L

CYDECTIN

Moxidectin
(Pour-On for Beef and Dairy Cattle)

VETERINARY USE ONLY

CYDECTIN

Moxidectin
(Pour-On for Beef and Dairy Cattle)

(Each mL contains 5 mg of moxidectin)

For the treatment of infections and infestations due to internal and external parasites in cattle including lactating dairy cows.

EXTERNAL USE ONLY

WARNINGS:

1. Treated cattle must not be slaughtered for use in food for at least 15 days after the latest treatment with this drug.
2. Do not use in milk-fed calves. Do not use in calves less than 14 weeks of age to be processed for veal.
3. Keep this and all drugs out of reach of children.
4. Avoid direct contact with skin and eyes.


NOTE TO USER - When used according to label directions, no milk withholding time is required following the application of CYDECTIN Pour-On on dairy cattle.

Store between 4-25 °C. AVOID FREEZING.
Protect from light. Store bottle in box.

Boehringer Ingelheim (Canada) Ltd.
5180 South Service Road
Burlington, Ontario L7L 5H4



Net Contents
5 L



CYDECTIN

Moxidectin
(Pour-On for Beef and Dairy Cattle)

Dosage and administration: EXTERNAL USE ONLY. CYDECTIN® Pour-On should be applied to healthy pigs along the top of the back from the withers to the base of the tail. Use 1 mL per 10 kg body weight to administer a dose of 0.5 mg/kg.

DOSAGE TABLE

Body weight (kg)	Dose (mL)	Dose per pig (mL)	Body weight (kg)	Dose (mL)	Dose per pig (mL)
40	4	1000	200	40	1000
50	5	1250	250	50	1250
60	6	1500	300	60	1500
70	7	1750	350	70	1750
80	8	2000	400	80	2000
90	9	2250	450	90	2250
100	10	2500	500	100	2500
110	11	2750	550	110	2750
120	12	3000	600	120	3000
130	13	3250	650	130	3250
140	14	3500	700	140	3500
150	15	3750	750	150	3750
200	20	5000	800	200	5000

See package insert for complete information on dosage and administration. See bottom of side for lot number and expiration date.

8844
2180
CC2657L
8844
CC2657L



Boehringer Ingelheim
 Moxidectin
 Solution à verser
 pour bovins de boucherie et bovins laitiers
 (Pour-On for Beef and Dairy Cattle)
 Moxidectin
 Solution à verser pour bovins de boucherie et bovins laitiers
 (Pour-On for Beef and Dairy Cattle)
 Moxidectin
 Solution à verser pour bovins de boucherie et bovins laitiers
 (Pour-On for Beef and Dairy Cattle)



Moxidectin
 Solution à verser pour
 bovins de boucherie et
 bovins laitiers
 (Pour-On for
 Beef and
 Dairy Cattle)
 Moxidectin
 Solution à verser pour
 bovins de boucherie et
 bovins laitiers
 (Pour-On for
 Beef and
 Dairy Cattle)
 Moxidectin
 Solution à verser pour
 bovins de boucherie et
 bovins laitiers
 (Pour-On for
 Beef and
 Dairy Cattle)

CYDECTIN®

DIN 02228386

USAGE VÉTÉRINAIRE SEULEMENT

CYDECTIN®

**Moxidectin
(Solution à verser pour bovins de
boucherie et bovins laitiers)**

(Chaque mL renferme 5 mg de moxidectin)
 Pour le traitement des infections et des infestations
 attribuables aux parasites internes et externes chez les
 bovins, y inclus les vaches laitières en lactation.

USAGE EXTERNE SEULEMENT

MISES EN GARDE :

1. Les animaux traités ne doivent pas être abattus à des fins alimentaires dans un délai d'au moins 15 jours après le dernier traitement à l'aide de ce médicament.
2. Ne pas utiliser chez les vœux de lait. Ne pas utiliser chez les vœux de moins de 14 semaines d'âge élevés pour la production de viande de vœux.
3. Garder ce produit, ainsi que tout autre médicament de la portée des enfants.
4. Éviter le contact direct avec la peau ou les yeux.

AVIS À L'UTILISATEUR - Lorsque le produit est utilisé conformément aux directives de l'étiquette, aucune période d'attente pour l'utilisation du lait n'est requise à la suite de l'administration de la solution à verser CYDECTIN à 0,5% chez les bovins laitiers.

Garder entre 4 et 25 °C. ÉVITER LE GEL.
 Garder à l'abri de la lumière. Entourer la bouteille dans la boîte.

Boehringer Ingelheim (Canada) Ltd.
 5180 South Service Road
 Burlington (Ontario) L7L 5H4



Contenu Net
10 L



Net Contents/Contenu Net
10 L

Posologie et administration : Solution à verser CYDECTIN® à usage externe seulement, devrait être appliquée sur une peau saine, le long du dos, du garrot jusqu'à l'attache de la queue. Utiliser 1 mL par 10 kg de poids vif pour administrer une dose de 0.5 mg/kg.


TABLEAU POSOLOGIQUE

Poids vif (kg)	Dose (mL)	Doses per flacon (10 L)	Poids vif (kg)	Dose (mL)	Doses per flacon (10 L)
40	4	2500	250	25	400
50	5	2000	300	30	332
60	6	1666	350	35	284
70	7	1428	400	40	250
80	8	1250	450	45	222
90	9	1110	500	50	200
100	10	1000	550	55	180
110	11	908	600	60	166
120	12	832	650	65	152
130	13	768	700	70	140
140	14	714	750	75	122
150	15	666	800	80	124
200	20	500	850	85	116

Voir le dépliant à l'intérieur pour obtenir toutes les informations sur la posologie et l'administration. Voir le dessous ou le côté pour le numéro de lot et date de péremption.



Net Contents
10 L



**Moxidectin
(Pour-On for Beef and
Dairy Cattle)**

(Each mL contains 5 mg of moxidectin)
 For the treatment of infections and infestations due to
 internal and external parasites in cattle,
 including lactating dairy cows.

EXTERNAL USE ONLY

WARNINGS:

1. Treated cattle must not be slaughtered for use in food for at least 15 days after the latest treatment with this drug.
2. Do not use in milk-fed calves. Do not use in calves less than 14 weeks of age to be processed for veal.
3. Keep this and all drugs out of reach of children.
4. Avoid direct contact with skin and eyes.

NOTE TO USER - When used according to label directions, no milk withholding time is required following the application of CYDECTIN Pour-On on dairy cattle.

Store between 4-25°C. AVOID FREEZING.
 Protect from light. Store bottle in box.

Boehringer Ingelheim (Canada) Ltd.
 5180 South Service Road
 Burlington, Ontario L7L 5H4



Dosage and administration: EXTERNAL USE ONLY. CYDECTIN® Pour-On should be applied to healthy skin along the top of the back from the withers to the base of the tail. Use 1 mL per 10 kg body weight to administer a dose of 0.5 mg/kg.

DOSAGE TABLE

Body weight (kg)	Dose (mL)	Doses per pack (10 L)	Body weight (kg)	Dose (mL)	Doses per pack (10 L)
40	4	2500	250	25	400
50	5	2000	300	30	332
60	6	1666	350	35	284
70	7	1428	400	40	250
80	8	1250	450	45	222
90	9	1110	500	50	200
100	10	1000	550	55	180
110	11	908	600	60	166
120	12	832	650	65	152
130	13	768	700	70	140
140	14	714	750	75	122
150	15	666	800	80	124
200	20	500	850	85	116

See package insert for complete information on dosage and administration. See bottom or side for lot number and expiration date.

