

## **PRODUCT MONOGRAPH**

### **Pr SANDOZ BICALUTAMIDE**

Bicalutamide tablets

50 mg tablets

Non-steroidal Anti-androgen

Sandoz Canada Inc.  
145 Jules-Léger  
Boucherville, QC, Canada  
J4B 7K8

Date of Revision: September 8, 2017

Control Number: 208772

## Table of Contents

<b>PART I: HEALTH PROFESSIONAL INFORMATION .....</b>	<b>3</b>
SUMMARY PRODUCT INFORMATION.....	3
INDICATIONS AND CLINICAL USE .....	3
CONTRAINDICATIONS.....	3
WARNINGS AND PRECAUTIONS .....	4
ADVERSE REACTIONS .....	7
DRUG INTERACTIONS .....	11
DOSAGE AND ADMINISTRATION .....	12
OVERDOSAGE.....	12
ACTION AND CLINICAL PHARMACOLOGY .....	13
STORAGE AND STABILITY .....	13
DOSAGE FORMS, COMPOSITION AND PACKAGING.....	13
<b>PART II: SCIENTIFIC INFORMATION .....</b>	<b>15</b>
PHARMACEUTICAL INFORMATION .....	15
CLINICAL TRIALS .....	16
DETAILED PHARMACOLOGY .....	18
TOXICOLOGY.....	19
REFERENCES.....	25
<b>PART III: CONSUMER INFORMATION .....</b>	<b>28</b>

**Pr Sandoz Bicalutamide**  
(Bicalutamide tablets 50 mg)

**PART I: HEALTH PROFESSIONAL INFORMATION**

**SUMMARY PRODUCT INFORMATION**

<b>Route of Administration</b>	<b>Dosage Form / Strength</b>	<b>Clinically Relevant Nonmedicinal Ingredients</b>
Oral	Tablet / 50 mg	lactose monohydrate  <i>For a complete listing see DOSAGE FORMS, COMPOSITION AND PACKAGING section.</i>

**INDICATIONS AND CLINICAL USE**

Sandoz Bicalutamide (bicalutamide) 50 mg is indicated for use in combination therapy with either a luteinizing hormone releasing hormone (LHRH) analogue or surgical castration in the treatment of metastatic (Stage D2) prostate cancer.

**Pediatrics:**

The safety and effectiveness of bicalutamide in children has not been established.

**CONTRAINDICATIONS**

Sandoz Bicalutamide is contraindicated in the following:

- Patients who are hypersensitive to the drug or any of its components. For a complete listing, see the DOSAGE FORMS, COMPOSITION AND PACKAGING section of the Product Monograph.
- Patients with localized prostate cancer otherwise undergoing watchful waiting (see WARNINGS AND PRECAUTIONS).
- Women: The safety and effectiveness of bicalutamide in women has not been studied.
- Children: The safety and effectiveness of bicalutamide in children has not been studied.

## WARNINGS AND PRECAUTIONS

Sandoz Bicalutamide should only be prescribed by a qualified healthcare professional who is experienced with the treatment of prostate cancer and the use of anti-androgens.

- Sandoz Bicalutamide 150mg/day dose should not be used (see WARNINGS & PRECAUTIONS, General).
- Rare hepatic failure, including fatal outcomes (see WARNINGS & PRECAUTIONS, Hepatic).
- Uncommon interstitial lung disease, including fatal outcomes (see WARNINGS & PRECAUTIONS, Respiratory).

### **General**

During treatment with bicalutamide, somnolence has been reported and those patients who experience this symptom should observe caution when driving or using machines.

### **Localized Prostate Cancer Patients**

Bicalutamide 150 mg is NOT to be administered.

Evidence from a large on-going clinical study demonstrates that at 5.4 year median follow-up, the use of bicalutamide 150 mg as immediate therapy for the treatment of localized prostate cancer in patients otherwise undergoing watchful waiting is associated with increased mortality. Health Canada previously assessed bicalutamide 150 mg *versus* castration in the locally advanced patient population and found level 1 scientific evidence (one of the 2 randomized clinical trials) of increased mortality in bicalutamide 150 mg treated patients.

Patients taking bicalutamide 50 mg per day for the treatment of metastatic prostate cancer are not affected by this new information.

### **Anti-androgen Withdrawal Syndrome**

In some patients with metastatic prostate cancer, anti-androgens (steroidal and non-steroidal), may promote, rather than inhibit, the growth of prostate cancer. A decrease in PSA and/or clinical improvement following discontinuation of antiandrogens has been reported. It is recommended that patients prescribed an antiandrogen, who have PSA progression, should have the antiandrogen discontinued immediately and be monitored for 6 - 8 weeks for a withdrawal response prior to any decision to proceed with other prostate cancer therapy.

### **Cardiovascular**

Bicalutamide is indicated for use in combination with either an LHRH analogue or surgical castration. Combined androgen blockade with an anti-androgen plus LHRH analogue or surgical castration increases risk of cardiovascular disease (heart attack, cardiac failure, sudden cardiac death) and adversely affects independent cardiovascular risk factors (serum lipoproteins, insulin sensitivity and obesity). Physicians should carefully consider whether the benefits of combined androgen blockade outweigh the potential cardiovascular risk. Assessment of cardiovascular risk factors, monitoring for signs and symptoms suggestive of development of cardiovascular disease, and management according to local clinical practice and guidelines should be considered.

### **Effect on QT/QTc interval**

Bicalutamide is indicated for use in combination with either an LHRH analogue or surgical castration. Combined androgen blockade with an anti-androgen plus LHRH analogue or surgical castration has the potential to prolong QT/QTc interval on ECG. In patients with a history of or who have risk factors for QT prolongation including congenital long QT syndrome, electrolyte abnormalities, or congestive heart failure and in patients taking concomitant medicinal products that may prolong the QT interval including Class IA (e.g. quinidine, procainamide), Class III (e.g. amiodarone, sotalol, dofetilide, ibutilide), or Class IC (e.g. flecainide, propafenone) antiarrhythmic medications (see DRUG INTERACTIONS), physicians should assess the benefit risk ratio, including the potential for Torsade de Pointes prior to initiating bicalutamide.

### **Endocrine and Metabolism**

A reduction in glucose tolerance and/or glycated hemoglobin (HbA1c) has been observed in males receiving bicalutamide in combination with LHRH analogues. This may manifest as diabetes or loss of glycemic control in those with pre-existing diabetes. Consideration should therefore be given to monitoring blood glucose and/or glycated hemoglobin (HbA1c) in patients receiving bicalutamide in combination with LHRH analogues.

### **Gynaecomastia, Breast Pain**

Gynaecomastia has been reported in patients receiving bicalutamide. For metastatic (M1) patients receiving bicalutamide 50 mg, concomitant surgical or medical castration may reduce the effects of gynaecomastia.

### **Hematologic**

Anemia is a known physiologic consequence of testosterone suppression. Assessment of anemia risk and management according to local clinical practice and guidelines should be considered. There may be an increased risk of hemorrhage in patients taking coumarin anticoagulants that are started on bicalutamide. (see ADVERSE REACTIONS and DRUG INTERACTIONS).

### **Hepatic**

Bicalutamide is extensively metabolized in the liver. Data suggests that the elimination of bicalutamide may be slower in subjects with severe hepatic impairment and this could lead to increased accumulation of bicalutamide. Therefore, Sandoz Bicalutamide should be used with caution in patients with moderate to severe hepatic impairment.

Hepatotoxicity including rare hepatic failure has been observed with bicalutamide, and fatal outcomes have been reported. Sandoz Bicalutamide therapy should be discontinued if changes are severe (also see Post-Market Adverse Drug Reactions).

### **Musculoskeletal**

#### **Changes in Bone Density**

Bicalutamide is indicated for use in combination with either an LHRH analogue or surgical castration. Decreased bone mineral density can be anticipated with long term combined androgen blockade with an anti-androgen plus LHRH analogue or surgical castration. Combined androgen blockade is associated with increased risks of osteoporosis and skeletal bone fractures. The risk of skeletal fracture increases with the duration of combined androgen blockade. Assessment of

osteoporosis risk and management according to clinical practice and guidelines should be considered.

In patients with significant risk factors for decreased bone mineral content and/or bone mass such as chronic alcohol and/or tobacco use, presumed or strong family history of osteoporosis or chronic use of drugs that can reduce bone mass such as anticonvulsants or corticosteroids, combined androgen blockade may pose an additional risk. In these patients, risk versus benefit must be weighed carefully before combined androgen blockade is instituted.

### **Respiratory**

Uncommon cases of interstitial lung disease (some cases have been fatal) have been reported with bicalutamide (also see Post-Market Adverse Drug Reactions). Interstitial lung disease has been reported most often at doses greater than 50 mg. Bicalutamide 150 mg is NOT to be administered.

If patients present with worsening of respiratory symptoms such as dyspnoea, cough and fever, bicalutamide should be interrupted and prompt investigation initiated. If Interstitial Lung Disease is confirmed, Sandoz Bicalutamide should be discontinued and the patient treated appropriately.

### **Sexual Function/Reproduction**

Sandoz Bicalutamide may cause reversible impairment of male fertility. Animal studies (see TOXICOLOGY) indicate that bicalutamide can affect spermatogenesis (spermatozoa count and motility) and increase the interval to successful mating. A period of subfertility or infertility should be assumed in men taking Sandoz Bicalutamide. Although the long-term effects of Sandoz Bicalutamide on male fertility have not been evaluated in humans, patients and their partners should follow adequate contraception during and for at least 130 days after Sandoz Bicalutamide therapy.

### **Skin**

In rare cases, photosensitivity reactions have been reported for patients taking bicalutamide. Patients should be advised to avoid direct exposure to excessive sunlight or UV-light while on Sandoz Bicalutamide and the use of sunscreens may be considered. In cases where the photosensitivity reaction is more persistent and/or severe, an appropriate symptomatic treatment should be initiated.

### **Special Populations**

**Pregnant and Nursing Women:** Sandoz Bicalutamide is contraindicated in females and must not be given to pregnant women. Bicalutamide may cause fetal harm when administered to pregnant women. Sandoz Bicalutamide is contraindicated during breast-feeding.

The male offspring of rats (but not rabbits) receiving doses of 10 mg/kg/day and above, were observed to have reduced anogenital distance and hypospadias in reproductive toxicology studies. These pharmacological effects have been observed with other antiandrogens. No other teratogenic effects were observed in rabbits (receiving doses up to 200 mg/kg/day) or rats (receiving doses up to 250 mg/kg/day).

**Pediatrics:** The safety and effectiveness of bicalutamide (non-steroidal antiandrogen) in children has not been established.

### **Monitoring and Laboratory Tests**

Regular assessments of serum Prostate Specific Antigen (PSA) may be helpful in monitoring patients' response.

Anemia has been observed in patients treated with bicalutamide. Hemoglobin levels should be monitored.

Baseline risk factors of cardiovascular diseases should be assessed. Patients receiving bicalutamide should be monitored periodically for risk factors, signs and symptoms of cardiovascular diseases. In addition, baseline ECG recording and serum potassium, calcium, and magnesium levels are recommended. Monitoring of ECG and serum electrolyte levels during treatment should also be considered for those at risk for electrolyte abnormality and QTc prolongation.

Since transaminase abnormalities and jaundice, rarely severe, have been reported with the use of bicalutamide, periodic liver function tests should be considered. If clinically indicated, discontinuation of therapy should be considered. Abnormalities are usually reversible upon discontinuation.

Since bicalutamide may elevate plasma testosterone and estradiol levels, fluid retention could occur. Accordingly, Sandoz Bicalutamide should be used with caution in those patients with cardiac disease.

A reduction in glucose tolerance has been observed in males receiving bicalutamide in combination with LHRH analogues. This may manifest as diabetes or loss of glycaemic control in those with pre-existing diabetes. Consideration should therefore be given to monitoring blood glucose and/or glycated hemoglobin (HbA1c) in patients receiving bicalutamide in combination with LHRH analogues, especially diabetic patients (See WARNINGS AND PRECAUTIONS, Endocrine and Metabolism).

Potential of coumarin anticoagulant effects have been reported in patients receiving concomitant Sandoz Bicalutamide therapy, which may result in increased Prothrombin Time (PT) and International Normalized Ratio (INR). Close monitoring of PT/INR is advised and anticoagulant dose adjustment should be considered (see ADVERSE REACTIONS and DRUG INTERACTIONS).

## **ADVERSE REACTIONS**

### **Adverse Drug Reaction Overview**

#### **Bicalutamide in Metastatic Patients**

In patients with advanced prostate cancer, treated in the multicentre, double-blind controlled clinical trial comparing bicalutamide 50 mg once daily with flutamide 250 mg three times a day, each in combination with an LHRH analogue, the most frequent adverse experiences included: hot flushes (53%), asthenia (22%), constipation (22%), nausea (14%), peripheral edema (13%), anemia (13%), haematuria (12%), abdominal pain (11%), dizziness (10%), gynecomastia (9%), rash (9%), chest pain (8%), erectile dysfunction (7%), flatulence (7%), dyspepsia (7%), decreased appetite (6%), breast tenderness (6%), weight increase (5%), cardiac failure (4%), depression (4%), dry skin (4%), alopecia (4%), pruritus (3%), somnolence (3%), myocardial infarction (3%), decreased libido (2%), hirsutism (2%), and hypersensitivity reactions (1%) including angioedema and urticaria.

Adverse event reports of abnormal liver function test results occurred in 7% of patients. These changes were frequently transient and rarely severe, resolving or improving with continued therapy or following cessation of therapy.

Hepatic failure and interstitial lung disease (see WARNINGS AND PRECAUTIONS) have been observed in post-marketed data and fatal outcomes have been reported for both.

Bicalutamide, in general has been well tolerated with few withdrawals due to adverse events. The most common adverse events leading to withdrawal of study medication were abnormal liver function tests (1.5%), hot flushes (1.0%), and nausea and vomiting (0.7%).

After a 160 week follow-up, there were 213/401 deaths in the bicalutamide-LHRH arm and 235/407 deaths in the flutamide-LHRH arm of the trial. There were 30 vs. 18 deaths due to adverse events in the two arms respectively and in both arms, the most common causes of death due to adverse events were attributed to the cardiovascular system (see ‘Cardiovascular’ under the ‘Clinical Trial Adverse Drug Reactions’ sub-section below).

Myocardial infarction and cardiac failure were observed in a pharmaco-epidemiology study of LHRH agonists and anti-androgens used in the treatment of prostate cancer. The risk appeared to be increased when bicalutamide was used in combination with LHRH agonists. Fatal outcomes of myocardial infarction have been reported.

### **Clinical Trial Adverse Drug Reactions**

The following adverse experiences were reported within the same clinical trial with an incidence of  $\geq 5\%$ , regardless of causality.

**Table 1 Incidence of Adverse Events ( $\geq 5\%$  in Either Treatment Group) Regardless of Causality**

Adverse Event	Treatment Group Number of Patients (%)	
	Bicalutamide Plus LHRH Analogue (n=401)	Flutamide Plus LHRH Analogue (n=407)
Hot Flushes	211 (53)	217 (53)
Pain (General)	142 (35)	127 (31)
Back Pain	102 (25)	105 (26)
Asthenia	89 (22)	87 (21)



Adverse Event	Treatment Group Number of Patients (%)	
	Bicalutamide Plus LHRH Analogue (n=401)	Flutamide Plus LHRH Analogue (n=407)
Constipation	87 (22)	69 (17)
Pelvic Pain	85 (21)	70 (17)
Infection	71 (18)	57 (14)
Nausea	56 (14)	54 (13)
Peripheral Edema	53 (13)	42 (10)
Anemia <sup>a</sup>	51 (13)	60 (15)
Dyspnea	51 (13)	32 (8)
Diarrhea	49 (12)	107 (26)
Nocturia	49 (12)	55 (14)
Hematuria	48 (12)	26 (6)
Abdominal Pain	46 (11)	46 (11)
Dizziness	41 (10)	35 (9)
Bone Pain	37 (9)	43 (11)
Gynecomastia	36 (9)	30 (8)
Rash	35 (9)	30 (7)
Urinary Tract Infection	35 (9)	36 (9)
Chest Pain	34 (8)	34 (8)
Hypertension	34 (8)	29 (7)
Cough Increased	33 (8)	24 (6)
Pharyngitis	32 (8)	23 (6)
Paresthesia	31 (8)	40 (10)
Increased Liver Enzyme Test <sup>b</sup>	30 (7)	46 (11)
Weight Loss	30 (7)	39 (10)
Headache	29 (7)	27 (7)
Flu Syndrome	28 (7)	20 (5)
Myasthenia	27 (7)	19 (5)
Insomnia	27 (7)	39 (10)
Erectile Dysfunction	27 (7)	35 (9)
Flatulence	26 (7)	22 (5)
Hyperglycemia	26 (7)	27 (7)
Dyspepsia	26 (7)	23 (6)
Decreased Appetite	25 (6)	29 (7)
Sweating	25 (6)	20 (5)
Bronchitis	24 (6)	11 (3)
Breast Pain (tenderness)	23 (6)	15 (4)
Urinary Frequency	23 (6)	29 (7)
Alkaline Phosphatase Increased	22 (5)	24 (6)
Weight-Increased	22 (5)	18 (4)
Arthritis	21 (5)	29 (7)
Anxiety	20 (5)	9 (2)
Urinary Retention	20 (5)	14 (3)
Urinary Impaired	19 (5)	15 (4)
Pneumonia	18 (4)	19 (5)
Pathological Fracture	17 (4)	32 (8)
Depression	16 (4)	33 (8)
Vomiting	16 (4)	28 (7)
Rhinitis	15 (4)	22 (5)
Urinary Incontinence	15 (4)	32 (8)

<sup>a</sup> Anemia includes hypochromic anemia and iron deficiency anemia.

<sup>b</sup> Abnormal liver function tests reported as adverse events.

In addition, the following adverse experiences were reported by investigators within the same clinical trial (as possible adverse drug reactions in the opinion of investigating clinicians) with a

frequency of less than 5%, during treatment with bicalutamide 50 mg plus an LHRH analogue. These experiences are not necessarily considered as causally related to drug treatment.

**Cardiovascular:** In the pivotal trial of 813 patients comparing bicalutamide 50 mg once daily with Flutamide 250 mg three times a day, each in combination with an LHRH analogue, an imbalance of deaths related to cardiovascular adverse events was noted (bicalutamide-LHRH therapy: 18 deaths; Flutamide-LHRH therapy: 9 deaths) however, there is difficulty in interpreting this imbalance as the exposure was longer on the bicalutamide-LHRH arm by a mean of 13 weeks. Other cardiovascular-related experiences reported include angina pectoris, congestive heart failure, myocardial infarction, heart arrest, coronary artery disorder, syncope, atrial fibrillation, cerebrovascular accident, deep thrombophlebitis, arrhythmia, bradycardia, cerebral ischemia, hemorrhage.

**Central Nervous System:** hypertonia, confusion, somnolence, decreased libido, neuropathy, nervousness

**Endocrine System:** diabetes mellitus

**Gastrointestinal:** melena, rectal hemorrhage, dry mouth, dysphagia, gastrointestinal disorder, periodontal abscess, gastrointestinal carcinoma, rectal disorder, intestinal obstruction, gastritis

**Hematological:** ecchymosis, thrombocytopenia

**Immune System Disorders:** hypersensitivity, angioedema and urticaria

**Metabolic & Nutritional:** edema, BUN increased, creatinine increased, dehydration, gout, hypercholesteremia, hypoglycemia, hypercalcemia

**Musculoskeletal:** leg cramps, bone disorders, myalgia

**Respiratory System:** lung disorder, asthma, epistaxis, sinusitis, pleural effusion, voice alteration

**Skin & Appendages:** dry skin, alopecia, pruritus, herpes zoster, skin carcinoma, skin disorder, skin hypertrophy, hirsutism, skin ulcer

**Special Senses:** cataract, abnormal vision, conjunctivitis

**Urogenital:** dysuria, urinary urgency, hydronephrosis, urinary tract disorder, bladder stenosis, kidney calculus, prostatic disorder, balanitis

**Whole Body:** neoplasm, neck pain, fever, chills, sepsis, hernia, cyst, injection site reaction, allergic reaction, neck rigidity, face edema

#### **Abnormal Hematologic and Clinical Chemistry Findings**

Laboratory abnormalities including elevated AST, ALT, bilirubin, BUN, creatinine and decreased hemoglobin and white cell count have been reported in both bicalutamide-LHRH

analogue treated and flutamide-LHRH analogue treated patients. Increased liver enzyme tests and decreases in hemoglobin were reported less frequently with bicalutamide-LHRH analogue therapy. Other changes were reported with similar incidence in both treatment groups.

### **Post-Market Adverse Drug Reactions**

The following adverse reactions have been identified during post-approval use of bicalutamide:

**Cardiovascular:** Myocardial infarction (fatal outcomes have been reported, cardiac failure, sudden cardiac death)

**Hepato-biliary disorders:** Hepatic failure (fatal outcomes have been reported)

**Hematologic:** Anaemia, increased PT/INR have been reported due to drug interaction between bicalutamide and coumarin anticoagulants.

**Respiratory:** Interstitial lung disease (fatal outcomes have been reported)

**Skin and subcutaneous tissue disorders:** Photosensitivity reaction

## **DRUG INTERACTIONS**

### **Drug-Drug Interactions**

Clinical studies with bicalutamide have not demonstrated any drug/drug interactions with LHRH analogues.

*In vitro* studies have shown that the (R)-enantiomer is an inhibitor of CYP3A4, with lesser inhibitory effects on CYP 2C9, 2C19 and 2D6 activity. Although *in vitro* studies have suggested the potential for bicalutamide to inhibit cytochrome 3A4, a number of clinical studies show the magnitude of any inhibition is unlikely to be of clinical significance for the majority of substances which are metabolised by cytochrome P450. Nevertheless, such an increase in AUC could be of clinical relevance for drugs with a narrow therapeutic index (e.g. cyclosporin).

*In vitro* studies have shown that bicalutamide, which is highly protein bound can displace coumarin anticoagulants such as warfarin, from protein binding sites. In patients receiving bicalutamide and warfarin concomitantly, there have been reports of increased effects of anticoagulation such as increased INR above therapeutic levels and risk of hemorrhagic events. It is therefore recommended that if Sandoz Bicalutamide is administered in patients who are receiving coumarin anticoagulation, PT/INR should be closely monitored and adjustments of anticoagulant dose considered (see WARNINGS AND PRECAUTIONS and ADVERSE REACTIONS)..

Sandoz Bicalutamide is indicated for use in combination with either an LHRH analogue or surgical castration. Since combined androgen blockade prolongs the QTc interval, the combination use of bicalutamide with an LHRH analogue, and medicinal products known to prolong the QTc interval or able to induce Torsades de Pointes should be carefully evaluated. Such medicinal products include but are not limited to the examples that follow: Class IA (e.g.

quinidine, disopyramide), Class III (e.g. amiodarone, sotalol, dofetilide, ibutilide, dronedarone), or Class IC (e.g. flecainide, propafenone) antiarrhythmic medicinal products, antipsychotics (e.g. chlorpromazine), antidepressants (e.g. amitriptyline, nortriptyline), opioids (e.g. methadone), macrolide antibiotics and analogues (e.g. erythromycin, clarithromycin, azithromycin), quinolone antibiotics (e.g. moxifloxacin), antimalarials (e.g. quinine), azole antifungals, 5-hydroxytryptamine (5-HT<sub>3</sub>) receptor antagonists (e.g. ondansetron), and beta-2 adrenoceptor analogues (e.g. salbutamol).

### **Drug-Food Interactions**

Interactions with food have not been established.

### **Drug-Herb Interactions**

Interactions with herbal products have not been established.

### **Drug-Laboratory Interactions**

Interactions with laboratory tests have not been established.

## **DOSAGE AND ADMINISTRATION**

### **Recommended Dose and Dosage Adjustment**

**Sandoz Bicalutamide 50 mg in metastatic disease:** The recommended dose for Sandoz Bicalutamide therapy in combination with an LHRH analogue or surgical castration is one 50 mg tablet once daily with or without food. Bicalutamide treatment should be started at the same time as treatment with an LHRH analogue or after surgical castration.

### **Dosing Considerations in Special Populations**

**Renal or Hepatic Impairment:** No dosage adjustment is necessary for patients with renal or mild hepatic impairment. Increased accumulation may occur in patients with moderate to severe hepatic impairment (see WARNINGS AND PRECAUTIONS).

## **OVERDOSAGE**

A single dose of bicalutamide that results in symptoms of an overdose considered to be life-threatening has not been established. In animal studies, bicalutamide demonstrated a low potential acute toxicity. The LD<sub>50</sub> in mice and rats was greater than 2000 mg/kg. Long-term clinical trials have been conducted with doses up to 200 mg of bicalutamide daily and these doses have been well tolerated.

There is no specific antidote; treatment of an overdose should be symptomatic. In the management of an overdose with bicalutamide, vomiting may be induced if the patient is alert. It should be remembered that in this patient population multiple drugs may have been taken. Dialysis is not likely to be helpful since bicalutamide is highly protein bound and is extensively metabolized. General supportive care, including frequent monitoring of vital signs and close observation of the patient, is indicated.

For management of a suspected drug overdose, contact your regional Poison Control Centre.

## **ACTION AND CLINICAL PHARMACOLOGY**

### **Pharmacodynamics**

Bicalutamide is a non-steroidal antiandrogen, devoid of other endocrine activity. Bicalutamide competitively inhibits the action of androgens by binding to cytosol androgen receptors in target tissue. This inhibition results in regression of prostatic tumours. Bicalutamide is a racemate and the (R)-enantiomer is primarily responsible for the antiandrogenic activity of bicalutamide.

### **Pharmacokinetics**

The absorption, distribution, metabolism and excretion of bicalutamide have been investigated after administration of a single 50 mg oral dose to volunteers. The results indicated that the dose was extensively absorbed and was excreted almost equally in urine (36%) and faeces (43%) over a 9 day collection period. There is no evidence of any clinically significant effect of food on bioavailability. Steady state plasma concentrations of the (R)-enantiomer of approximately 9 mcg/mL are observed during daily administration of 50 mg doses of bicalutamide. At steady state, the active (R)-enantiomer accounts for 99% of the circulating plasma bicalutamide concentration. Bicalutamide is highly protein bound (racemate 96%, (R)-enantiomer 99.6%). On daily administration, the (R)-enantiomer accumulates about 10-fold in plasma, consistent with an elimination half-life of approximately one week. The (S)-enantiomer is very rapidly cleared relative to the (R)-enantiomer. Bicalutamide is extensively metabolized via both oxidation and glucuronidation with approximately equal renal and biliary elimination of the metabolites.

### **Special Populations and Conditions**

**Pediatrics:** The pharmacokinetics of the (R)-enantiomer are unaffected by age.

**Geriatrics:** The pharmacokinetics of the (R)-enantiomer are unaffected by age.

**Hepatic Insufficiency:** The pharmacokinetics of the (R)-enantiomer are unaffected by mild to moderate hepatic impairment. Patients with severe hepatic impairment eliminate the (R)-enantiomer from plasma more slowly.

**Renal Insufficiency:** The pharmacokinetics of the (R)-enantiomer are unaffected by renal impairment.

## **STORAGE AND STABILITY**

Store between 15 and 30°C.

## **DOSAGE FORMS, COMPOSITION AND PACKAGING**

Sandoz Bicalutamide 50 mg tablets are white, or almost white, round, biconvex film-coated tablets, with a sign “B” on one side and “50” on the other side. Available in blister packs of 30 tablets, and in 30 and 1000 count bottles.


In addition to the active ingredient bicalutamide, each tablet contains the following inactive ingredients: lactose monohydrate, sodium starch glycolate, povidone, colloidal anhydrous silica, magnesium stearate, hydroxypropyl methylcellulose, polyethylene glycol, titanium dioxide, triacetin.

## PART II: SCIENTIFIC INFORMATION

### PHARMACEUTICAL INFORMATION

#### Drug Substance

- Proper name: Bicalutamide
- Chemical Name: (±)-N-[4-cyano-3-(trifluoromethyl)phenyl]-3-[(4-fluorophenyl)sulfonyl]-2-hydroxy-2-methyl-propanamide
- Molecular Formula: C<sub>18</sub>H<sub>14</sub>F<sub>4</sub>N<sub>2</sub>O<sub>4</sub>S
- Molecular Mass: 430.37
- Structural Formula:



#### Physicochemical Properties

- Description: Bicalutamide is a fine white or creamy white crystalline powder.
- Solubility: Practically insoluble in water, slightly soluble in isopropanol, dichloromethane, and ethanol. Sparingly soluble in methanol. Soluble in acetonitrile.
- Freely soluble in acetone and dimethylformamide.

#### Octanol/Water Partition

- pKa: Approximately 12.
- Chirality: Sandoz Bicalutamide is a racemate with its' antiandrogen activity being predominately exhibited by (R)-enantiomer of bicalutamide

## CLINICAL TRIALS

### Comparative Bioavailability Studies

A comparative, randomized, single-dose, double-blinded, two-treatment, single period, parallel design bioavailability study of Sandoz Bicalutamide 50 mg tablets was performed vs the Canadian reference, CASODEX<sup>®</sup> 50 mg, in healthy adult males under fasting conditions. The summary of the comparative bioavailability study carried out by Sandoz Bicalutamide is presented in the Table 2.

**Table 2 Comparative Bioavailability Study**

Bicalutamide (1 x 50 mg film-coated tablet) From measured data Geometric Mean Arithmetic Mean (CV %)				
Parameter	Sandoz Bicalutamide 50 mg film-coated tablet	Casodex <sup>®</sup> † 50 mg film-coated tablet	% Ratio of Geometric Means	90% Confidence Interval
AUC <sub>0-72</sub> (ng·h/mL)	54653.30 55163.51 (13.81)	53000.26 53534.57 (14.06)	103.12%	97.31%-109.27%
AUC <sub>∞</sub> (ng·h/mL)	181496.45 193824.8 (44.03)	223466.24 243288.82 (42.69)	81.22%	60.13%-109.70%
C <sub>max</sub> (ng/mL)	890.51 898.46 (13.64)	853.73 864.23 (15.79)	104.31%	98.26%-110.72%
T <sub>max</sub> <sup>*</sup> (h)	25.80 (53.41)	27.91 (54.42)		
T <sub>1/2</sub> <sup>*</sup> (h)	136.18 (49.35)	206.29 (76.27)		

† Casodex<sup>®</sup> is manufactured by AstraZeneca Canada Inc. Casodex<sup>®</sup> was purchased in Canada.

\*Expressed as the arithmetic mean (CV%) only.

### Study demographics and trial design

Time to treatment failure was the primary endpoint of a large, double-blinded, multicentre, non-inferiority clinical trial. Eight-hundred thirteen (813) patients with previously untreated advanced prostate cancer were randomized to receive bicalutamide 50 mg once daily (404 patients) or flutamide 250 mg (409 patients) three times a day, each in combination with luteinizing hormone-releasing hormone (LHRH) analogues (either goserelin acetate implant or leuprolide acetate depot).


### Study results

Approval was based on a median follow-up of 49 weeks which showed bicalutamide-LHRH analogue therapy was associated with a statistically significant ( $p = 0.005$ ) improvement in time to treatment failure. With a longer follow-up (median 95 weeks), improvement in time to treatment failure was no longer statistically significant ( $p = 0.10$ ).

In a survival analysis conducted after a median follow-up of 160 weeks was reached, 213 (52.7%) patients treated with bicalutamide-LHRH analogue therapy and 235 (57.5%) patients treated with Flutamide-LHRH analogue therapy had died. There was no significant difference in survival between treatment groups (see Figure 1). The hazard ratio for survival was 0.87 (95% confidence interval 0.72 to 1.05,  $p = 0.15$ ).




**Figure 1** The Kaplan-Meier probability of survival for both antiandrogen treatment groups.



————— Bicalutamide plus luteinizing hormone releasing hormone-A  
----- Flutamide plus luteinizing hormone releasing hormone-A

There was no significant difference in time to objective tumour progression between treatment groups (see Figure 2). Objective tumour progression was defined as the appearance of any bone metastases or the worsening of any existing bone metastases on bone scan attributable to metastatic disease, or an increase by 25% or more of any existing measurable extraskelatal metastases. The hazard ratio for time to progression of bicalutamide plus LHRH analogue to that of flutamide plus LHRH analogue was 0.93 (95% confidence interval, 0.79 to 1.10,  $p = 0.41$ ).

**Figure 2 Kaplan-Meier curve for time to progression for both antiandrogen treatment groups.**



Quality of life was assessed using a self-administered patient questionnaire on pain, bed disability, activity limitation, physical capacity, social functioning, emotional well-being, vitality, overall health, general symptoms, and treatment-related symptoms. At a median follow-up of 95 weeks, no significant differences were noted between the two treatment groups.

## DETAILED PHARMACOLOGY

### Animal Pharmacology

#### Pharmacodynamics

##### *In vitro*

Bicalutamide binds to rat, dog and human prostate and rat pituitary androgen receptors. In radioligand displacement assays, graded doses of bicalutamide inhibit the binding of the synthetic androgen [<sup>3</sup>H]-R-1881. Using the rat prostate androgen receptor, the displacement curves for bicalutamide, the antiandrogen hydroxyflutamide, R-1881 and the natural ligand, 5 $\alpha$ -dihydrotestosterone are parallel.

Bicalutamide binds around fifty times less effectively than 5 $\alpha$ -dihydrotestosterone and around 100 times less effectively than R-1881 to the rat androgen receptor but has an affinity around 4-fold higher for the prostate and 10 times higher for the pituitary androgen receptor than hydroxyflutamide. The relative affinities of bicalutamide for dog and human prostate androgen receptors are similar to those for the rat and are again higher than for hydroxyflutamide. Bicalutamide has no effect on prostate steroid 5 $\alpha$ -reductase and has negligible affinity for the sex hormone-binding globulin and no affinity for corticosteroid-binding globulin.

### ***In vivo***

**Rat:** In the rat, bicalutamide and the (R)-enantiomer are at least 1000 times more potent as antiandrogens than the (S)-enantiomer which had very low potency. In immature castrated rats, 0.5 mg/kg oral bicalutamide prevents stimulation of the growth of the seminal vesicles and ventral prostate gland in response to daily subcutaneous injections of testosterone propionate (200 mcg/kg). In intact mature rats, several studies show that bicalutamide causes a dose-related reduction in accessory sex organ weights. In these studies bicalutamide had only a minimal effect on serum luteinizing hormone and testosterone.

**Dog:** Studies show that bicalutamide is an effective antiandrogen at the dog prostate but does not elevate serum testosterone concentrations. The ED<sub>50</sub> value for inducing prostate atrophy in the dog following daily oral treatment for 6 weeks is about 0.1 mg/kg. At all doses tested up to 100 mg/kg, bicalutamide has no effect on serum testosterone concentrations.

**Monkey:** Longitudinal studies in monkeys, where prostate and seminal vesicle sizes were followed by magnetic resonance imaging, show bicalutamide to be a highly potent (1-5 mg/kg) antiandrogen with negligible effect on serum testosterone, although there was wide intra- and inter-animal variability.

### **Pharmacokinetics**

Bicalutamide displays enantioselective pharmacokinetics in rats, dogs and man with the (R)-enantiomer being slowly eliminated, particularly in the dog and man, and consequently accumulating on daily administration. Steady state ratios (R)-enantiomer to (S)-enantiomer are highest in man (~ 100:1), lower in the rat (~ 14:1) and even lower in the dog (~ 3:1).

## **TOXICOLOGY**

### **Acute Toxicity**

In animal studies, bicalutamide demonstrated a low potential acute toxicity. The LD<sub>50</sub> in mice, rats and dogs was greater than 2000 mg/kg. The LD<sub>50</sub> in rabbits was greater than 200 mg/kg.

### **Long-Term Toxicity**

Multiple dose studies include one, six and twelve month studies in the rat and dog (see following table).

**Table 3 Long-Term Toxicity**

Species	Duration	No. of Animals/Group	Route	Dose mg/kg/day	Effects
Rat Wistar	1 month	28-40 M* + 28-40 F*	Oral	0, 25, 100, 500	Minor, reversible drug-related increases (<10%) in plasma total protein & albumin in Groups III and IV. Small prostate and seminal vesicles at all doses and reversible, drug-related increase in liver weight (21% and 35% for Groups III and IV males, 36%, 55% and 90% for Groups II-IV females) and adrenal weights (24% and 50% for Group III and IV males, 16% and 27% for Group III and IV females). Microscopic changes were consistent with anti-androgen activity (e.g. atrophy of ventral prostate & seminal vesicles, Leydig cell hyperplasia). There were changes consistent with enzyme induction in the liver in bicalutamide-dosed groups and a minimal to mild increases in cortical single cell necrosis in adrenal glands in bicalutamide-dosed groups and a minimal to mild hypertrophy of follicular epithelium and reduced colloid, in the thyroid gland from dosed groups. There was a dose dependant increase in basophilia and RNA content of hepatocyte cytoplasm in all bicalutamide-dosed groups and an increase in smooth ER in some Group IV animals.
Rat Wistar	6 month	30-57 M* + 30-57 F*	Oral	0, 10, 50, 250	There were small reductions in body weight and a reduction in alkaline phosphatase in dosed males. A small, reversible increase in plasma protein and albumin, a decrease in packed cell volume and hæmoglobin was seen in all bicalutamide-dosed groups. Expected reversible size reduction in prostate and seminal vesicles (all dosed) and testes (Groups III & IV); some Group IV males had enlarged testes. Increased adrenal gland weight in all groups - increased weight of liver, kidneys, heart (females only) and brain, not accompanied by important histological change. Histopathological changes were seen in the prostate and seminal vesicles (atrophy), testes (atrophy of seminiferous tubules and Leydig cell hyperplasia), ovaries (granulosa-thecal cell hyperplasia), adrenals (cortical hypertrophy to cortical vacuolation), pituitary glands in males (castration cells) and thyroid gland (epithelial cell hypertrophy). Many of these changes were reduced or reversed in the drug withdrawal period - the adrenal cortical vacuolation and castration cells in pituitary were largely unchanged.

\* Reflects group related extra animals (e.g. For pharmacokinetic, coagulation, hæmatology and drug withdrawal).

**Table 3 Long-Term Toxicity (Continued)**

Species	Duration	No. of Animals/Group	Route	Dose mg/kg/day	Effects
Rat Wistar	12 month	33-45 M* + 33-45 F*	Oral in diet	0, 5, 15, 75	Increased incidence of small/flaccid testes in Groups III & IV, small reduction in male body weight, alkaline phosphatase, alanine aminotransferase and aspartate aminotransferase and a small reversible reduction in haemoglobin & related indices in Group IV females. There was a small increase in plasma total protein. There was an increase in liver weight in Groups III & IV accompanied by hepatocyte hypertrophy and basophilia, related to MFO induction. Other histological changes were limited to the reproductive and some endocrine organs-increased adrenal weight, hypertrophy of the thyroid follicular epithelium, follicular epithelium hyperplasia and colloid basophilia, testicular tubular atrophy (Group III & IV), atrophy of prostate and seminal vesicles - except for testicular atrophy, changes reversed or showed signs of recovery following withdrawal. There was an increase in tumours in three hormone-sensitive organs, benign testicular Leydig cell tumours (all dosed groups), thyroid follicular adenomas (Group IV) and uterine carcinomas (Group IV) at the end of the withdrawal period.
Dog Beagle	6 weeks	2 M + 2 F	Oral	0, 25, 75, 150	There was a reduction in the weight of the testes, epididymides & prostate gland in dosed groups and atrophy of the seminiferous tubules and diffuse Leydig cell hyperplasia; the epididymides showed minimal/mild microcystic degeneration and spermatozoa were absent. Adrenal glands of dogs given bicalutamide were increased in weight; there was cytoplasmic vacuolation of the cortex (changes related to bicalutamide administration); there were no bicalutamide-related changes in the female reproductive tract. Significant increases in heart rate (28-39 BPM) were seen in all groups by week 5. The P-R interval was reduced in all groups (21-26 msec, week 5); there were no important differences in blood pressure and no changes were seen on the electrocardiogram for any dog. There was an increase in plasma cholesterol (1.5 times control) at all time points for Groups III & IV; there was a mild phenobarbital-like induction of cytochrome P450.

\* Reflects group related extra animals (e.g. For pharmacokinetic, coagulation, hematology and drug withdrawal).

**Table 3 Long-Term Toxicity (Continued)**

Species	Duration	No. of Animals/Group	Route	Dose mg/kg/day	Effects
Dog Beagle	6 months	5-8 M* + 5-8 F*	Oral	0, 2.5, 10, 100	Two males (Group III & IV) were killed because of infection, 1 Group IV female with an infection during week 20 recovered. Body weight (8%) and food intake were reduced in the first 6 weeks in Group IV; this group gained weight in the withdrawal period. A dose-related reduction in P-R interval was seen; the changes (Group IV) reversed 4 weeks after drug withdrawal; there were no histological findings in the heart associated with these changes. There was a reduction in weight & diffuse atrophy of the prostate gland (all doses), Leydig cell hyperplasia, seminiferous tubule atrophy, arrested spermatogenesis of the testes, ductal atrophy of the epididymides, endometrial gland reduction of the uterus, increased keratinization of the cervix and vagina, atrophy of the mammary gland and increased weight, cortical vacuolation and cortical hypertrophy of the adrenal glands; effects associated with anti-androgenic activity. Following 16 weeks drug withdrawal Group IV animals showed no evidence of atrophy of the prostate; other changes in the male and female reproductive tract were absent or less marked. Cortical vacuolation of the adrenal gland was still present.
Dog Beagle	12 months	5-8 M* + 5-8 F*	Oral	0, 1, 2.5, 50	There was a decrease in mean P-R interval in Group III & IV (7-16% & 16-22% respectively). There was a higher liver weight and small increases in alkaline phosphatase in Group IV because of enzyme-induction. Increases in plasma glucose, urea, cholesterol and in alanine aminotransferase in Group IV, were a result of antiandrogenic activity. There was decrease in weight and atrophy of the prostate gland, changes in the testes (Leydig cell hyperplasia, exfoliated seminiferous epithelial cells, maturation arrest) and epididymides (stromal hyperplasia, ductal atrophy, sperm reduction) at all doses and mammary glands (reduced acinar development) in Groups III & IV; these were anti-androgen related effects. Following 6 months withdrawal there was no compound-related changes in the male reproductive tract; reduced acinar development was present in all high-dose females. There was increased weight and cortical vacuolation of the adrenal glands (all dosed groups) and cortical hypertrophy and hyperplasia (Groups III & IV); on withdrawal both adrenal weight and vacuolation showed evidence of reversibility but cortical hyperplasia was still evident.

\* Reflects group related extra animals (e.g. for pharmacokinetic, coagulation, hematology and drug withdrawal).

### **Carcinogenicity**

Two-year oral oncogenicity studies in both male and female rats and mice at doses of 5, 15 or 75 mg/kg/day of bicalutamide have been completed. A variety of tumour target organ effects were identified and were attributed to the antiandrogenicity of bicalutamide, namely testicular benign interstitial (Leydig) cell tumours in rats at all dose levels (the steady state plasma concentration with the 5 mg/kg/day dose is comparable to a human oral 50 mg/day dose) and uterine adenocarcinoma in rats at 75 mg/kg/day (3 times greater than the human plasma concentration, based on a maximum dose of 50 mg/day of bicalutamide for an average 70 kg patient). There is no evidence of Leydig cell hyperplasia in patients treated in combination with LHRH analogues. Uterine tumours are not relevant to the indicated patient population.

A small increase in incidence of hepatocellular carcinoma in male mice given 75 mg/kg/day of bicalutamide (plasma concentration 4 times greater than the human concentration) and an increased incidence of benign thyroid follicular cell adenomas in rats given 5 mg/kg and above were recorded. These neoplastic changes were progressions of non-neoplastic changes related to hepatic enzyme induction observed in animal toxicity studies. Enzyme induction has not been observed following bicalutamide administration in man. There were no tumourigenic effects suggestive of genotoxic carcinogenesis.

### **Mutagenicity**

A comprehensive battery of both *in vitro* and *in vivo* genotoxicity tests has demonstrated that bicalutamide does not have genotoxic activity.

### **Reproduction & Teratology**

Reproduction and teratology studies have been conducted in the rat and rabbit (see following table).

**Table 4      Reproduction and Teratology**

Species	Duration	No. of Animals/Dose	Route	Dose mg/kg/day	Type of Study	Effects
Rat Wistar	11 weeks	25 M + 150 F	Oral	0, 0.25, 5, 250	Male Fertility	In male rats dosed at 250 mg/kg/day, the precoital interval and time to successful mating were increased in the first pairing but no effects on fertility following successful mating were seen. These effects were reversed by 7 weeks after the end of an eleven week period of dosing.
Rat Wistar	2 wks before mating through pregnancy & lactation*	6 M + 6 F	Oral	0, 10, 250	Female Fertility	No effects on dosed females (10 and 250 mg/kg/day) or their female offspring were observed. As an antiandrogen, there was feminization of the male offspring of all dosed females leading to hypospadias. Affected male offspring were also impotent.
Rat Wistar	days 6-15	20 pregnant per group**	Oral	0, 1, 10, 50, 250	Teratology	The offspring of rats dosed at 0, 1, 10, 50 and 250 mg/kg/day and rabbits dosed at 0, 10, 50 and 200 mg/kg/day did not show evidence of any developmental or teratogenic effect. The only developmental abnormality seen was a predictable reduction of anogenital distance due to the androgenic properties of the drug in only male fetuses at doses of 10, 50 and 250 mg/kg/day; no effect was seen at 1 mg/kg/day. Feminization of the male offspring of all females dosed at 10 and 50 mg/kg/day was reported in a fertility and reproductive study in rats.
Rabbit Dutch Belled	days 6-18	20 pregnant per group***	Oral	0, 10, 50, 200	Teratology	

\* Up to twelve weeks

\*\* An extra 4 females were added for pharmacokinetic sample

\*\*\* An extra 6 females were added for pharmacokinetic sample



## REFERENCES

1. Bauer J, Connely R, et al. Biostatistical modeling using traditional preoperative and pathological prognostic variables in the selection of men at high risk for disease recurrence after radical prostatectomy for prostate cancer. *The Journal of Urology* 1998;159: 929-933.
2. Blackledge G. Casodex<sup>®</sup> - Mechanisms of Action and Opportunities for Usage. *Cancer* 1993; 72 (12). Dec 15 Supplement: 3830-3833.
3. Blute M, Bergstralh E, et al. Validation of partin tables for predicting pathological stage of clinically localized prostate cancer. *The Journal of Urology* 2000;164:1591-1595.
4. Boyle GW, McKillop D, Phillips PJ, Harding JR, Pickford R, McCormick AD. Metabolism of Casodex<sup>®</sup> in laboratory animals. *Xenobiotica* 1993; 23(7): 781-798.
5. Cockshott ID, Cooper KJ, Sweetmore DS, Blacklock NJ, Denis L, Study Group. The Pharmacokinetics of Casodex<sup>®</sup> in Prostate Cancer Patients after Single and During Multiple Dosing. *European Urology* 1990; 18 Suppl 1 3:10-17.
6. Cockshott ID, Sotaniemi EI, Cooper KJ, Jones DC. The pharmacokinetics of Casodex<sup>®</sup> enantiomers in subjects with impaired liver function. *British Journal of Clinical Pharmacology* 1993; 36(4): 339-343.
7. D'Amico A, Whittington R, et al. Pretreatment nomogram for prostate-specific antigen recurrence after radical prostatectomy or external-beam radiation therapy for clinically localized prostate cancer. *Journal of Clinical Oncology*, Vol 17, No.1 (Jan), 1999:168-172.
8. D'Amico A, Whittington R, et al. The combination of preoperative prostate specific antigen and postoperative pathological findings to predict prostate specific antigen outcome in clinically localized prostate cancer. *The Journal of Urology* 1998;160:2096-2101.
9. Freeman SN, Mainwaring WIP, Furr BJA. A possible explanation for the peripheral selectivity of a novel non-steroidal pure antiandrogen, Casodex<sup>®</sup> (ICI 176, 334). *British Journal of Cancer* 1989; 60:664-668.
10. Furr BJA. Casodex<sup>®</sup> (ICI 176, 334) - a new, pure, peripherally-selective antiandrogen: preclinical studies. *Hormone Research* 1989; 32 Suppl 1:69-76. Proceedings of the International Symposium on Endocrine Therapy, Monaco, 19-21 Nov 1988.
11. Furr BJA. Casodex<sup>®</sup>: Preclinical Studies. *European Urology* 1990; 18 Suppl 3: 2-9.

12. Furr BJA. Pharmacological properties and potential clinical utility of ICI 176, 334: a novel, non-steroidal, peripheral selective antiandrogen. *Hormonal Therapy of Prostatic diseases: Basic and Clinical Aspects* 1987; 148-161.
13. Kattan M, Eastham J, et al. A preoperative nomogram for disease recurrence following radical prostatectomy for prostate cancer. *Journal of the National Cancer Institute*, May 1998, vol 90 (10):766-771.
14. Kennealey GT, Furr BJA. Use of Non-steroidal Antiandrogen Casodex<sup>®</sup> in Advanced Prostatic Carcinoma. (Review, 43 Refs). *Urologic Clinics of North America* 1991;18(1) Feb: 99-110.
15. Levine et al. Androgen-Deprivation Therapy in Prostate Cancer and Cardiovascular Risk. *Circulation* 2010; 121; 833-840.
16. Lunglmayr G. Casodex<sup>®</sup> (ICI 176, 334) A New, Non-steroidal Antiandrogen. *Early Clinical Results. Hormone Research* 1989; 32 Suppl 1: 77-81.
17. Mahler C, Denis L. Clinical profile of a new non-steroidal antiandrogen. *Journal of Steroid Biochemistry and Molecular Biology* 1990; 37 (6): 921-924.
18. McConnell JD. Physiologic Basis of Endocrine Therapy for Prostatic Cancer (Review 96 refs.). *Urologic Clinics of North America* 1991; 18(1): 1-13.
19. McKillop D, Boyle GW, Cockshott ID, Jones DC, Phillips PJ, Yates RA. Metabolism and enantioselective pharmacokinetics of Casodex<sup>®</sup> in man. *Xenobiotica* 1993; 23(11): 1241-1253.
20. McLaren D, McKenzie M, Duncan G, Pickles T. Watchful waiting or watchful progression? Prostate Specific Antigen Doubling times and clinical behavior in patients with Early Untreated Prostate Carcinoma. *American Cancer Society* 1998;82(2) 342-348.
21. Narayan P, Gajendran V, et al. The role of transrectal ultrasound-guided biopsy-based staging, preoperative serum prostate-specific antigen, and biopsy gleason score in prediction of final pathologic diagnosis in prostate cancer. *Urology* 1995, 46(2):205-212.
22. Newling DWW. The response of advanced prostatic cancer to a new non-steroidal antiandrogen. Results of a multicentre open phase II study of Casodex<sup>®</sup> European *Urology* 1990; 18 Suppl 3:18-21.
23. Partin A, Kattan M, et al. Combination of prostate-specific antigen, clinical stage, and Gleason score to predict pathological stage of localized prostate cancer. *JAMA* 1997;277(18):1445-1451.

24. Partin A, Yoo J, et al. The use of prostate specific antigen, clinical stage and gleason score to predict pathological stage in men with localized prostate cancer. *The Journal of Urology* 1993, 150:110-114.
25. Pisansky T, Kahn M, Bostwick D. An enhanced prognostic system for clinically localized carcinoma of the prostate. *Cancer* 1997;79:2154-2161.
26. Pruthi R, Johnstone I, Tu I, Stamey T. Prostate-specific antigen doubling times in patients who have failed radical prostatectomy: Correlation with histologic characteristics of the Primary Cancer. *Adult Urology* 1997;49(5):737-742.
27. Schellhammer P, Sharifi R, Block N, et al. A controlled trial of bicalutamide versus flutamide, each in combination with luteinizing hormone-releasing hormone analogue therapy, in patients with advanced prostate cancer. *Urology* May 1995; 45(5): 745-752.
28. Schellhammer P, Sharafi, R, Block N, et al. Clinical benefits of bicalutamide compared with flutamide in combined androgen blockade for patients with advanced prostatic carcinoma: final report of a double-blind, randomized, multicenter trial. *Urology* 1997; 50(3): 330-336.
29. See, William A., Wirth Manfred. P. Bicalutamide as immediate therapy either alone or as adjuvant to standard care of patients with localized and locally advanced prostate cancer: First analysis of the Early Prostate Cancer Program. *J. of Urology*, August 2002, 168:429-435.
30. Zagars G, Pollack A, Eschenbach. A prognostic factors for clinically localized prostate carcinoma. *Cancer* 1997;79:1370-1380.
31. AstraZeneca Canada Inc., Product Monograph, Casodex<sup>®</sup> (bicalutamide 50 mg tablets), Control number: 205811, Date of revision: July 13, 2017.

**PART III: CONSUMER INFORMATION**

**PrSANDOZ® BICALUTAMIDE,  
50 mg  
(Bicalutamide Tablets)**

This leaflet is Part III of a three-part "Product Monograph" published when Sandoz Bicalutamide was approved for sale in Canada and is designed specifically for Consumers. This leaflet is a summary and will not tell you everything about Sandoz Bicalutamide. Contact your doctor or pharmacist if you have any questions about the drug.

**ABOUT THIS MEDICATION**

**What the medication is used for:**

Sandoz Bicalutamide is used in the treatment of advanced prostate cancer in combination with other drugs (LHRH analogues) which reduce the levels of androgens in the body or surgery.

**What it does:**

Androgens are male sex hormones within the body which can cause tumour growth within the prostate. Sandoz Bicalutamide belongs to a group of medicines called non-steroidal anti-androgens. This means that Sandoz Bicalutamide interferes with some of the actions of androgens to prevent the tumour from growing.

**What are the Stages of Prostate Cancer:**

- **Localized disease** – the early stages of disease when prostate cancer is confined to the prostate gland.
- **Locally advanced disease** – the disease progresses and the cancer spreads to other tissues within the pelvis.
- **Advanced or metastatic disease** – the disease progresses to other parts of the body.

The PSA (Prostate Specific Antigen) test is a simple blood test for a protein produced by the prostate (PSA). This test has helped in the detection of prostate cancer resulting in an increase in the number of men whose prostate cancer is detected at an early stage.

**What are the Treatment Options for Localized Prostate Cancer:** The optimal treatment for a given individual will depend on the specific circumstances of his case. For localized disease, patients are usually offered one of the following:

- Surgery to remove the prostate
- Targeted radiotherapy to kill the cancer cells in the prostate

- No treatment initially (watchful waiting) whereby the patient is monitored until there are signs of progression before treatment is started.

**When it should not be used:**

- Do not take Sandoz Bicalutamide if you have early phase (localized) prostate cancer requiring watchful waiting.
- Do not take Sandoz Bicalutamide if you are allergic to bicalutamide or any of the nonmedicinal ingredients in Sandoz Bicalutamide.
- Sandoz Bicalutamide must not be taken by women, including pregnant women or mothers who are breast feeding their babies.
- Sandoz Bicalutamide must not be given to children.

**What the medicinal ingredient is:**

The active ingredient in Sandoz Bicalutamide is bicalutamide.

**What the important nonmedicinal ingredients are:**

Each tablet contains the following inactive ingredients:  
Lactose monohydrate, Sodium starch glycolate, Povidone, Colloidal anhydrous silica, Magnesium stearate, Hydroxypropyl methylcellulose, Polyethylene glycol, Titanium dioxide, Triacetin.

**What dosage forms it comes in:**

Sandoz Bicalutamide is available in 50 mg tablets.

**WARNINGS AND PRECAUTIONS**

- **Sandoz Bicalutamide should only be prescribed by a doctor experienced with the treatment of prostate cancer.**
- **Sandoz Bicalutamide 150mg/day dose should not be used.**
- **Sandoz Bicalutamide may rarely be associated with liver failure; some cases have been fatal.**
- **Sandoz Bicalutamide may be associated with uncommon cases of interstitial lung disease; some cases have been fatal.**

BEFORE you use Sandoz Bicalutamide talk to your doctor or pharmacist if:

- You have liver disease.
- You have lung disease.
- You have low bone mineral density (BMD).
- You have low red blood cell count (anemia).
- You are taking blood thinners or medicines to prevent blood clots.

- You have heart or blood vessel conditions, including heart disease, a heart condition called ‘long QT syndrome’ or family history of this heart condition, heart rhythm problems (arrhythmia), or are being treated with medicines for these conditions. The risk of heart rhythm problems may be increased in such patients when using Sandoz Bicalutamide.

If you go into hospital let the medical staff know you are taking Sandoz Bicalutamide.

If you are taking Sandoz Bicalutamide, you and/or your partner should use birth control while you are taking Sandoz Bicalutamide and for at least 130 days after stopping Sandoz Bicalutamide. Talk to your doctor if you have any questions about birth control.

Sandoz Bicalutamide may have an effect on male fertility which could be reversible.

Avoid direct exposure to excessive sunlight or UV-light while you are taking Sandoz Bicalutamide.

Sandoz Bicalutamide may make you feel sleepy. Do not drive or use machines until you know how the drug affects you.

### **INTERACTIONS WITH THIS MEDICATION**

Please inform your doctor if you are taking or have recently taken any other medicines, even those not prescribed.

- In particular please inform your doctor if you are taking blood thinners like warfarin or medicines to prevent blood clots.
- If you are taking any medicines that may increase the risk of having an abnormal heart rhythm.
- If you are taking medicines used to treat heart rhythm problems.

### **PROPER USE OF THIS MEDICATION**

Follow your doctor’s instructions about when and how to take your tablets. Ask your doctor or pharmacist if you are not sure.

- The usual adult dose is 50 mg daily.
- Swallow the tablet(s) whole with a drink of water.
- Try to take your dose at the same time each day.

During the first few months of use, you may be monitored by your physician for signs of changes in your liver function. In approximately 2.0% of patients, such changes may lead to withdrawal of therapy.

If you experience a rise in PSA while taking Sandoz Bicalutamide, your physician may discontinue Sandoz Bicalutamide for several weeks in order to monitor your condition off treatment.

### **Overdose**

In case of drug overdose, contact a health care practitioner, hospital emergency department or regional Poison Control Centre immediately, even if there are no symptoms.

### **Missed Dose**

You should take Sandoz Bicalutamide as prescribed. However, if you miss a dose do not take an extra dose, just resume your usual schedule.

## **SIDE EFFECTS AND WHAT TO DO ABOUT THEM**

Like all medicines, Sandoz Bicalutamide 50 mg can have side effects.

Side effects that are very common (more than 10 in every 100 patients are likely to have them):

- dizziness
- nausea
- hot flushes
- feeling weak
- decreased red blood cell count (anemia)
- puffiness/swelling
- constipation

Side effects that are common (1 to 10 in every 100 patients are likely to have them):

- loss of appetite
- reduced sex drive
- depression
- sleepiness
- indigestion
- flatulence
- loss of hair or hair re-growth
- rash
- itching
- dry skin
- impotence
- chest pain
- tender or enlarged breast tissue
- weight gain
- heart failure
- heart attack

Occasionally Sandoz Bicalutamide may be associated with changes in your blood which may require your doctor to do certain blood tests.

**SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM**

Symptom/Effect	Talk with your doctor or pharmacist		Stop taking drug and call your doctor or pharmacist
	Only if severe	In all cases	
<b>Very common (more than 10 in every 100 patients are likely to have them)</b>			
Blood in urine		√	
Abdominal pain		√	
<b>Common (1 to 10 in every 100 patients are likely to have them)</b>			
Yellow skin and eyes (jaundice). These may be symptoms of liver damage.		√	
Heart failure (reduced heart function)		√	
Heart attack		√	
<b>Uncommon (1 to 10 in every 1000 patients are likely to have them)</b>			
Serious breathlessness, or sudden worsening of breathlessness, possibly with a cough or fever. Some patients taking Sandoz Bicalutamide 50 mg get an inflammation of the lungs called interstitial lung disease.		√	
Severe itching of the skin (with raised lumps) or swelling of the face, lips, tongue and/or throat, which may cause difficulty in swallowing		√	
<b>Rare (1 to 10 in every 10,000 patients are likely to have them)</b>			
increased skin sensitivity to sunlight		√	

Tell your doctor or pharmacist if you think you have any of these or any other problems with your tablets.

*This is not a complete list of side effects. For any unexpected effects while taking Sandoz Bicalutamide, contact your doctor or pharmacist.*

**HOW TO STORE IT**

- Keep your tablets in the container they came in.
- Do not take your tablets after the expiry date on the container. Dispose of them in an appropriate way.
- Keep your tablets in a safe place where children cannot see or reach them. Your tablets could harm them.
- Keep your tablets between 15 and 30°C.

**REPORTING SIDE EFFECTS**

You can report any suspected side effects associated with the use of health products to the Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<http://www.hc-sc.gc.ca/dhp-mps/medeff/report-declaration/index-eng.php>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345

*NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.*

**MORE INFORMATION**

This document, plus the full Product Monograph prepared for health professionals, can be obtained by contacting the sponsor, Sandoz Canada Inc., at:

1-800-361-3062

or by written request at:

145, Jules-Léger  
Boucherville, (QC), Canada  
J4B 7K8

or by e-mail at :

[medinfo@sandoz.com](mailto:medinfo@sandoz.com)

This leaflet was prepared by Sandoz Canada Inc.

Last revised: September 8, 2017