

PRODUCT MONOGRAPH

^{Pr}METHYLPREDNISOLONE SODIUM SUCCINATE FOR INJECTION USP

Sterile Powder
500 mg, 1 g Vials

Glucocorticoid

Hospira Healthcare Corporation
17300 Trans-Canada Highway
Kirkland, Québec
H9J 2M5

Date of Revision: October 18, 2017

Submission Control No: 209610

Table of Contents

PART I: HEALTH PROFESSIONAL INFORMATION	3
SUMMARY PRODUCT INFORMATION.....	3
INDICATIONS AND CLINICAL USE	3
CONTRAINDICATIONS.....	5
WARNINGS AND PRECAUTIONS	5
ADVERSE REACTIONS	15
DRUG INTERACTIONS.....	18
DOSAGE AND ADMINISTRATION	25
OVERDOSAGE.....	28
ACTION AND CLINICAL PHARMACOLOGY.....	29
STORAGE AND STABILITY	31
DOSAGE FORMS, COMPOSITION AND PACKAGING.....	31
PART II: SCIENTIFIC INFORMATION.....	34
PHARMACEUTICAL INFORMATION	34
CLINICAL TRIALS	34
DETAILED PHARMACOLOGY	36
TOXICOLOGY.....	36
REFERENCES.....	37
PART III: CONSUMER INFORMATION	40

PART I: HEALTH PROFESSIONAL INFORMATION

SUMMARY PRODUCT INFORMATION

Route of Administration	Dosage Form / Strength	All Nonmedicinal Ingredients
Intravenous or intramuscular injection or by intravenous infusion	sterile powder 500 mg and 1 g and diluent with preservative	Powder : Sodium dihydrogen phosphate and Disodium hydrogen phosphate Diluent : Bacteriostatic Water for Injection with Benzyl alcohol 0.9%v/v as preservative <i>see Dosage Forms, Composition and Packaging section.</i>

INDICATIONS AND CLINICAL USE

Intravenous administration of Methylprednisolone Sodium Succinate for Injection USP is indicated in situations in which a rapid and intense hormonal effect is required. These include the following:

Hypersensitivity and dermatologic conditions

- Status asthmaticus
- Anaphylactic reactions (see text)
- Drug reactions
- Contact dermatitis
- Urticaria
- Generalized neurodermatitis
- Reactions to insect bites
- Pemphigus foliaceus and vulgaris
- Exfoliative dermatitis
- Erythema multiforme

In anaphylactic reactions, epinephrine or norepinephrine should be administered first for an immediate hemodynamic effect followed by intravenous injection of Methylprednisolone Sodium Succinate for Injection USP and other accepted procedures. There is evidence that the corticoids through their prolonged hemodynamic effect are of value in preventing recurrent attacks of acute anaphylactic reactions.

In sensitivity reactions, such as in serum sickness, allergic dermatosis (urticaria) and reactions to insect bites, Methylprednisolone Sodium Succinate for Injection USP is capable of providing relief within 1/2 to 2 hours. In some asthmatic patients it may be advantageous to administer Methylprednisolone Sodium Succinate for Injection USP by slow intravenous drip over a period of hours.

As adjunctive therapy in

- Acute systemic lupus erythematosus
- Acute rheumatic fever
- Acute gout

In these conditions, Methylprednisolone Sodium Succinate for Injection USP may be given by slow intravenous administration over a period of several minutes. Thereafter, the patient should be placed on intramuscular or oral therapy as required for continued relief of symptoms. In these conditions, other accepted measures of therapy should also be instituted.

Ulcerative colitis

In addition to the above conditions, colonic instillation of Methylprednisolone Sodium Succinate in retention enemas or by continuous drip, have been shown to be a useful adjunct in the treatment of patients with ulcerative colitis.

Shock

In severe hemorrhagic or traumatic shock, adjunctive use of intravenous methylprednisolone sodium succinate may aid in achieving hemodynamic restoration. Corticoid therapy should not replace standard methods of combating shock, but present evidence indicates that concurrent use of large doses of corticoids with other measures may improve survival rates.

Organ transplants

Corticosteroids, both parenterally and orally, in high doses have been used following organ transplantation as part of multi-faceted attempts to reduce the rejection phenomenon. Methylprednisolone Sodium Succinate for Injection USP is suitable for such indications.

Cerebral oedema of non traumatic origin

Administration of Methylprednisolone Sodium Succinate for Injection USP immediately prior to intracranial surgery and in the immediate post-operative period has reduced the duration of post-operative complications related to cerebral oedema.

CONTRAINDICATIONS

Methylprednisolone Sodium Succinate for Injection USP is contraindicated:

- for systemic fungal infections
- in patients administered with live or live attenuated vaccines while receiving immunosuppressive doses of corticosteroids.
- for intrathecal or epidural administration. Reports of serious medical events have been associated with these routes of administration.
- for use in premature infants because the diluent for reconstitution contains benzyl alcohol. See WARNINGS AND PRECAUTIONS, General; Pediatrics.
- Intramuscular corticosteroid preparations are contraindicated for idiopathic thrombocytopenic purpura.

Except when used for short-term or emergency therapy, Methylprednisolone Sodium Succinate for Injection USP is contraindicated in patients with:

- arrested tuberculosis
- herpes simplex keratitis
- acute psychoses
- Cushing's syndrome

- peptic ulcer
- markedly elevated serum creatinine
- vaccinia and varicella.

WARNINGS AND PRECAUTIONS

Methylprednisolone Sodium Succinate for Injection USP should not be used in premature infants. The diluent for reconstitution of the Vials contains benzyl alcohol which has been reported to be associated with fatal “Gasping Syndrome” in premature infants. Exposure to excessive amounts of benzyl alcohol has been associated with toxicity (hypotension, metabolic acidosis), particularly in neonates, and an increased incidence of kernicterus, particularly in small preterm infants. There have been rare reports of deaths, primarily in preterm infants, associated with exposure to excessive amounts of benzyl alcohol. The amount of benzyl alcohol at which toxicity may occur is not known. See also CONTRAINDICATIONS; WARNINGS AND PRECAUTIONS, Pediatrics.

Benzyl alcohol is potentially toxic when administered locally to neural tissue.

General

Methylprednisolone Sodium Succinate for Injection USP should not be administered by any route other than those listed under SUMMARY PRODUCT INFORMATION. It is critical that, during administration of Methylprednisolone Sodium Succinate for Injection USP, appropriate technique be used and care taken to assure proper route of administration.

Dosage must be decreased or discontinued gradually when the drug has been administered for more than a few days.

Patients should be advised to inform subsequent physicians of the prior use of Methylprednisolone Sodium Succinate for Injection USP.

Serious medical events have been reported in association with the intrathecal/epidural routes of administration (see CONTRAINDICATIONS and ADVERSE REACTIONS).

In patients on corticosteroid therapy subjected to unusual stress, increased dosage of rapidly acting corticosteroids before, during and after the stressful situation is indicated.

The slower rate of absorption by intramuscular administration should be recognized.

The lowest possible dose of corticosteroid should be used to control the condition under treatment. Since complications of treatment with glucocorticoids are dependent on the size of the dose and the duration of treatment, a risk/benefit decision must be made in each individual case as to dose and duration of treatment and as to whether daily or intermittent therapy should be used.

The existence of diabetes, osteoporosis, renal insufficiency, chronic psychosis, hypertension, myasthenia gravis or predisposition to thrombophlebitis requires that Methylprednisolone Sodium Succinate for Injection USP be administered with extreme caution.

Carcinogenesis and Mutagenesis

Kaposi's sarcoma has been reported to occur in patients receiving corticosteroid therapy. Discontinuation

of corticosteroids may result in clinical remission.

Animal studies found corticosteroids to have a possible tumorigenic and mutagenic potential (See TOXICOLOGY, Carcinogenesis and also TOXICOLOGY, Mutagenesis).

Cardiovascular

There are reports of cardiac arrhythmias and/or circulatory collapse and/or cardiac arrest following the rapid administration of large intravenous doses of methylprednisolone sodium succinate (greater than 0.5 g administered over a period of less than 10 minutes). Bradycardia has been reported during or after the administration of large doses of methylprednisolone sodium succinate, and may be unrelated to the speed or duration of infusion.

Literature reports suggest an apparent association between the use of corticosteroids and left ventricular free wall rupture after a recent myocardial infarction; therefore, therapy with corticosteroids should be used with great caution in these patients.

Thrombosis including venous thromboembolism has been reported to occur with corticosteroids. As a result, corticosteroids should be used with caution in patients who have or may be predisposed to thromboembolic disorders.

As sodium retention with resultant oedema and potassium loss may occur in patients receiving corticosteroids, these agents should be used with caution, and only if strictly necessary, in patients with congestive heart failure. Corticosteroids should be used with caution in hypertension, or renal insufficiency. See also WARNINGS AND PRECAUTIONS, Endocrine and Metabolism.

Adverse effects of glucocorticoids on the cardiovascular system, such as dyslipidemia and hypertension, may predispose treated patients with existing cardiovascular risk factors to additional cardiovascular effects, if high doses and prolonged courses are used. Accordingly, corticosteroids should be employed judiciously in such patients and attention should be paid to risk modification and additional cardiac monitoring if needed.

Endocrine and Metabolism

Corticosteroid administration may result in hypothalamic-pituitary-adrenal (HPA) suppression (secondary adrenocortical insufficiency). The degree and duration of adrenocortical insufficiency produced is variable among patients and depends on the dose, frequency, time of administration and duration of glucocorticoid therapy. Drug induced secondary adrenocortical insufficiency may be minimized by gradual reduction of dosage. This type of relative insufficiency may persist for months after discontinuation of therapy; therefore, in any situation of stress occurring during that period, hormone therapy may need to be reinstated. In addition, acute adrenal insufficiency leading to a fatal outcome may occur if glucocorticoids are withdrawn abruptly.

Because glucocorticoids can produce or aggravate Cushing's syndrome, glucocorticoids should be avoided in patients with Cushing's disease.

Average and large doses of corticosteroids can cause elevation of blood pressure, salt and water retention, and increased excretion of potassium. These effects are less likely to occur with the synthetic derivatives except when used in large doses. Dietary salt restriction and potassium supplementation may be necessary. All corticosteroids increase calcium excretion. See also WARNINGS AND PRECAUTIONS,

Cardiovascular.

Corticosteroids, including methylprednisolone, can increase blood glucose, worsen pre-existing diabetes, and predispose those on long-term corticosteroid therapy to diabetes mellitus.

There is an enhanced effect of corticosteroids in patients with hypothyroidism. Metabolic clearance of corticosteroids is decreased in hypothyroid patients and increased in hyperthyroid patients. Changes in thyroid status of the patient may necessitate adjustment in dosage.

Pheochromocytoma crisis, which can be fatal, has been reported after administration of systemic corticosteroids, including methylprednisolone. Corticosteroids should only be administered to patients with suspected or identified pheochromocytoma after an appropriate risk/benefit evaluation.

Gastrointestinal

Corticosteroids should be used with caution in non-specific ulcerative colitis if there is a probability of impending perforation, abscess or other pyogenic infection and in diverticulitis, fresh intestinal anastomoses, and active or latent peptic ulcer, since they may increase the risk of a perforation. Signs of peritoneal irritation following gastrointestinal perforation in patients receiving corticosteroids may be minimal or absent.

Glucocorticoid therapy may mask the symptoms of peptic ulcer so that perforation or haemorrhage may occur without significant pain. In combination with NSAIDs, such as Aspirin (acetylsalicylic acid), the risk of developing gastrointestinal ulcers is increased.

Hematologic

Aspirin (acetylsalicylic acid) should be used cautiously in conjunction with corticosteroids in hypoprothrombinemia.

Hepatic/Biliary/Pancreatic

Drug-induced liver injury such as acute hepatitis can result from cyclical pulsed intravenous methylprednisolone (usually at initial dose \geq 1 gm/day). The time to onset of acute hepatitis can be several weeks or longer. Resolution of the adverse event has been observed after treatment was discontinued.

There is an enhanced effect of corticosteroids in patients with cirrhosis.

High doses of corticosteroids may produce acute pancreatitis.

Immune

Corticosteroids may suppress the immune system and may mask some signs of infection, and new infections may appear during their use. There may be decreased resistance and inability to localize infection when corticosteroids are used. Infections with any pathogen including viral, bacterial, fungal, protozoan or helminthic organisms, in any location in the body, may be associated with the use of corticosteroids alone or in combination with other immunosuppressive agents that affect cellular immunity, humoral immunity, or neutrophil function. These infections may be mild, but can be severe and at times fatal. With increasing doses of corticosteroids, the rate of occurrence of infectious complications increases.

Recent studies suggest that corticosteroids should not be used in septic shock (an unapproved indication), and suggest that increased mortality may occur in some subgroups at higher risk (e.g., elevated serum

creatinine greater than 2.0 mg/dL or secondary infections).

Do not use intra-articularly, intrabursally or for intratendinous administration for local effect in the presence of acute local infection.

Fungal infections

Corticosteroids may exacerbate systemic fungal infections and therefore should not be used in the presence of such infections unless they are needed to control drug reactions. There have been cases reported in which concomitant use of amphotericin B and hydrocortisone was followed by cardiac enlargement and congestive heart failure. See also CONTRAINDICATIONS; DRUG INTERACTIONS.

Special pathogens

Latent disease may be activated or there may be an exacerbation of intercurrent infections due to pathogens, including those caused by *Amoeba*, *Candida*, *Cryptococcus*, *Mycobacterium*, *Nocardia*, *Pneumocystis*, *Toxoplasma*.

It is recommended that latent amebiasis or active amebiasis be ruled out before initiating corticosteroid therapy in any patient who has spent time in the tropics or in any patient with unexplained diarrhea.

Similarly, corticosteroids should be used with great care in patients with known or suspected *Strongyloides* (threadworm) infestation. In such patients, corticosteroid-induced immunosuppression may lead to *Strongyloides* hyperinfection and dissemination with widespread larval migration, often accompanied by severe enterocolitis and potentially fatal gram-negative septicemia.

Corticosteroids should not be used in cerebral malaria. There is currently no evidence of benefit from steroids in this condition.

If corticosteroids are indicated in patients with latent tuberculosis or tuberculin reactivity, close observation is necessary as reactivation of the disease may occur. During prolonged corticosteroid therapy, these patients should receive chemoprophylaxis.

Vaccination

Administration of live or live, attenuated vaccines is contraindicated in patients receiving immunosuppressive doses of corticosteroids (see CONTRAINDICATIONS). Killed or inactivated vaccines may be administered; however, the response to such vaccines may be diminished. Indicated immunization procedures may be undertaken in patients receiving non immunosuppressive doses of corticosteroids.

While on corticosteroid therapy patients should not be vaccinated against smallpox. Other immunization procedures should not be undertaken in patients who are on corticosteroids, especially in high doses, because of possible hazards of neurological complications and lack of antibody response.

Viral infections

Chicken pox and measles can have a more serious or even fatal course in pediatric and adult patients on corticosteroids. In pediatric and adult patients who have not had these diseases, particular care should be taken to avoid exposure. The contribution of the underlying disease and/or prior corticosteroid treatment to the risk is also not known. If exposed to chicken pox, prophylaxis with varicella zoster immune globulin (VZIG) may be indicated. If exposed to measles, prophylaxis with immune globulin (IG) may be indicated. (See the respective package inserts for complete VZIG and IG prescribing information.) If

chicken pox develops, treatment with antiviral agents should be considered.

Recent studies do not support Methylprednisolone Sodium Succinate for Injection USP use during septic shock, and suggest that increased mortality may occur in some subgroups at higher risk (e.g., elevated serum creatinine greater than 2.0 mg/dL or secondary infections).

Musculoskeletal

An acute myopathy has been observed with the use of high doses of corticosteroids, most often occurring in patients with disorders of neuromuscular transmission (e.g. myasthenia gravis, see WARNINGS AND PRECAUTIONS - Neurologic), or in patients receiving concomitant therapy with anticholinergics such as neuromuscular blocking drugs (e.g. pancuronium). This acute myopathy is generalized, may involve ocular and respiratory muscles, and may result in quadriparesis. Elevations of creatine kinase may occur. Clinical improvement or recovery after stopping corticosteroids may require weeks to years.

Corticosteroids decrease bone formation and increase bone resorption both through their effect on calcium regulation (e.g., decreasing absorption and increasing excretion) and inhibition of osteoblast function. This, together with a decrease in the protein matrix of the bone secondary to an increase in protein catabolism, and reduced sex hormone production, may lead to inhibition of bone growth in pediatric patients and the development of osteoporosis at any age. Special consideration should be given to patients at increased risk of osteoporosis (e.g., postmenopausal women) before initiating corticosteroid therapy.

Osteoporosis is a common but infrequently recognized adverse effect associated with a long-term use of large doses of glucocorticoid.

Neurologic

Convulsions have been reported with concurrent use of methylprednisolone and cyclosporine. Since concurrent administration of these agents results in a mutual inhibition of metabolism, it is possible that convulsions and other adverse events associated with the individual use of either drug may be more apt to occur.

Systemic corticosteroids are not indicated for, and therefore should not be used to treat traumatic brain injury. A multicenter study revealed an increased mortality at 2 weeks and 6 months after injury in patients administered methylprednisolone sodium succinate compared to placebo.

Corticosteroids should be used with caution in patients with seizure disorders.

Corticosteroids should be used with caution in patients with myasthenia gravis.

There have been reports of epidural lipomatosis in patients taking corticosteroids (including reports in children).

Ophthalmologic

Use of corticosteroids may produce posterior subcapsular cataracts and nuclear cataracts (particularly in children), exophthalmos, or increased intraocular pressure, which may result in glaucoma with possible damage to the optic nerves, and may enhance the establishment of secondary ocular infections due to bacteria, fungi, or viruses. Intraocular pressure may become elevated in some individuals. If steroid therapy is continued for more than 6 weeks, intraocular pressure should be monitored.

The use of oral corticosteroids is not recommended in the treatment of optic neuritis and may lead to an increase in the risk of new episodes. Corticosteroids should be used cautiously in patients with ocular herpes simplex because of possible corneal perforation.

Corticosteroid therapy has been associated with central serous chorioretinopathy, which may lead to retinal detachment.

Psychiatric

Psychic derangements may appear when corticosteroids are used, ranging from euphoria, insomnia, mood swings, personality changes, and severe depression to frank psychotic manifestations. Also, existing emotional instability or psychotic tendencies may be aggravated by corticosteroids, and therefore these patients should be treated with caution.

Potentially severe psychiatric adverse reactions may occur with systemic steroids. Symptoms typically emerge within a few days or weeks of starting treatment. Most reactions recover after either dose reduction or withdrawal, although specific treatment may be necessary.

Psychological effects have been reported upon withdrawal of corticosteroids; the frequency is unknown. Patients/caregivers should be encouraged to seek medical attention if psychological symptoms develop in the patient, especially if depressed mood or suicidal ideation is suspected. Patients/caregivers should be alert to possible psychiatric disturbances that may occur either during or immediately after dose tapering/withdrawal of systemic steroids.

Sensitivity/Resistance

Allergic reactions may occur. Because rare instances of skin reactions and anaphylactic/anaphylactoid reactions have occurred in patients receiving corticosteroid therapy, appropriate precautionary measures should be taken prior to administration, especially when the patient has a history of allergy to any drug (see ADVERSE REACTIONS).

Sexual Function/Reproduction

Steroids may increase or decrease motility and number of spermatozoa in some patients.

Skin

Injection of Methylprednisolone Sodium Succinate for Injection USP may result in dermal and/or subdermal changes forming depressions in the skin at the injection site. In order to minimize the incidence of dermal and subdermal atrophy, care must be exercised not to exceed recommended doses in injections. Injection into the deltoid muscle should be avoided because of a high incidence of subcutaneous atrophy.

Special Populations

Pregnant Women: Corticosteroids readily cross the placenta. Corticosteroids have been shown to be teratogenic in many species when given in doses equivalent to the human dose. Administration of corticosteroids to pregnant animals can cause fetal malformations (cleft palate, skeletal malformations) and intra-uterine growth retardation (see TOXICOLOGY, Reproductive toxicity).

One retrospective study found an increased incidence of low birth weights in infants born of mothers receiving corticosteroids. Cataracts have been observed in infants born to mothers undergoing long-term treatment corticosteroids during pregnancy.

Since adequate human reproductive studies have not been done with methylprednisolone sodium succinate, Methylprednisolone Sodium Succinate for Injection USP should be used during pregnancy only at the lowest possible dose, only if clearly needed, where the potential benefit to the mother justifies the potential risk to the embryo or fetus.

Infants born to mothers who received substantial doses of corticosteroids during pregnancy should be observed for signs of hypoadrenalism and appropriate measures instituted if such signs are present. There are no known effects of corticosteroids on labour and delivery.

Benzyl alcohol can be introduced into the solution when Methylprednisolone Sodium Succinate for Injection USP Plain Vials are reconstituted with Bacteriostatic Water for Injection USP (benzyl alcohol as preservative) as directed.

Benzyl alcohol can cross the placenta.

Nursing Women: Corticosteroids are excreted in breast milk. Corticosteroids distributed into breast milk may suppress growth, interfere with endogenous corticosteroid production, or cause other untoward effects.

Because of the potential for serious adverse reactions in nursing infants from corticosteroids, a careful benefit-risk assessment should be conducted and a decision should be made whether to discontinue nursing, or discontinue the drug, taking into account the importance of the drug to the mother.

Pediatrics: Methylprednisolone Sodium Succinate for Injection USP is contraindicated for use in premature infants (see CONTRAINDICATIONS). Benzyl alcohol can be introduced into the solution; when Methylprednisolone Sodium Succinate for Injection USP Plain Vials are reconstituted with Bacteriostatic Water for Injection USP (benzyl alcohol as preservative). Benzyl alcohol has been associated with serious adverse events and death, particularly in pediatric patients, including the “gasping syndrome” in neonate and low-birth weight infants. The “gasping syndrome” is characterized by central nervous system depression, metabolic acidosis, gasping respirations, and high levels of benzyl alcohol and its metabolites found in the blood and urine. Additional symptoms may include gradual neurological deterioration, seizures, intracranial hemorrhage, hematologic abnormalities, skin breakdown, hepatic and renal failure, hypotension, bradycardia, and cardiovascular collapse. The minimum amount of benzyl alcohol at which toxicity may occur is not known. Premature and low-birth weight infants, as well as patients receiving high doses (>90 mg/kg/day), may be more likely to develop toxicity. Practitioners administering this and other medications containing benzyl alcohol should consider the combined daily metabolic load of benzyl alcohol from all sources.

Pediatric patients may experience a decrease in their growth velocity at low systemic doses and in the absence of laboratory evidence of HPA axis suppression. Growth velocity may therefore be a more sensitive indicator of systemic corticosteroid exposure in pediatric patients than some commonly used tests of HPA axis function. In order to minimize the potential growth effects of corticosteroids, pediatric patients should be titrated to the lowest effective dose.

Growth and development of infants and children on prolonged corticosteroid therapy should be carefully observed. Like adults, pediatric patients should be carefully observed with frequent measurements of blood pressure, weight, height, intraocular pressure, and clinical evaluation for the presence of infection, psychosocial disturbances, thromboembolism, peptic ulcers, cataracts, and osteoporosis.

Infants and children on prolonged corticosteroid therapy are at special risk from raised intracranial pressure.

High doses of corticosteroids may produce pancreatitis in children.

Geriatrics: In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy.

Caution is recommended with prolonged corticosteroid treatment in the elderly due to a potential increased risk for osteoporosis, as well as increased risk for fluid retention with possible resultant hypertension.

Monitoring and Laboratory testing

Corticosteroids may suppress reactions to skin tests.

Since methylprednisolone suppresses endogenous adrenocortical activity, it is highly important that the patient receiving Methylprednisolone Sodium Succinate for Injection USP be under careful observation, not only during the course of treatment but for some time after treatment is terminated. Monitoring for signs and symptoms of drug-induced secondary adrenocortical insufficiency may be necessary for up to one year following cessation of long-term or high-dose corticosteroid therapy.

ADVERSE REACTIONS

The following Adverse Reactions have been reported with the systemic use of Methylprednisolone Sodium Succinate for Injection USP and other corticosteroid preparations.

MedDRA (v15) System Organ Class		Undesirable Effect
<i>Blood and lymphatic system disorders</i>		Leukocytosis
<i>Infections and infestations</i>		Infection; opportunistic infection; injection site infections following non-sterile administration; decreased resistance to infection
<i>Immune system disorders</i>		Drug hypersensitivity (anaphylactoid reaction; anaphylactic reaction(with or without circulatory collapse))
<i>Endocrine disorders</i>		Cushingoid; hypopituitarism; steroid withdrawal syndrome; moon face; abnormal fat deposits; glycosuria; hypertrichosis; secondary adrenocortical and pituitary unresponsiveness (particularly in times of stress, as in trauma, surgery or illness) A steroid “withdrawal syndrome,” seemingly unrelated to adrenocortical insufficiency, may also occur following abrupt discontinuance of glucocorticoids. This syndrome includes symptoms such as: anorexia, nausea, vomiting, lethargy,

MedDRA (v15) System Organ Class		Undesirable Effect
		headache, fever, joint pain, desquamation, myalgia, weight loss, and/or hypotension. These effects are thought to be due to the sudden change in glucocorticoid concentration rather than to low corticosteroid levels.
<i>Metabolism and nutrition disorders</i>		Lipomatosis; sodium retention; sodium excretion; fluid retention; alkalosis hypokalemic; dyslipidaemia; metabolic acidosis; glucose tolerance impaired; increased requirement for insulin (or oral hypoglycemic agents in diabetics); nitrogen balance negative (due to protein catabolism); blood urea increased; increased appetite (which may result in weight increased); diuresis
<i>Psychiatric disorders</i>		Affective disorder (including affect lability, depressed mood, euphoric mood, drug dependence, suicidal ideation); psychotic disorder (including mania, delusion, hallucination, schizophrenia [aggravation of]); mental disorder; insomnia; mood swings; personality change; confusional state; abnormal behavior; anxiety; irritability; emotional instability
<i>Nervous system disorders</i>		Epidural lipomatosis; intracranial pressure increased (with papilloedema [idiopathic intracranial hypertension] usually following discontinuation of treatment); convulsion; amnesia; cognitive disorder; dizziness; headache; seizures; neuritis; neuropathy; paresthesia
<i>Eye disorders</i>		Central serous chorioretinopathy; cataract; glaucoma; exophthalmos; rare instances of blindness associated with periocular injections.
<i>Ear and labyrinth disorders</i>		Vertigo
<i>Cardiac disorders</i>		Cardiac failure congestive (in susceptible patients); arrhythmia; cardiac arrest; bradycardia; tachycardia; cardiac enlargement; circulatory collapse; hypertrophic cardiomyopathy in premature infants; myocardial rupture following recent myocardial infarction; pulmonary oedema; syncope
<i>Vascular disorders</i>		Hypertension; hypotension; thromboembolism; thrombophlebitis, thrombosis, vasculitis
<i>Respiratory, thoracic and mediastinal disorders</i>		Hiccups; bronchospasm, pulmonary embolism
<i>Gastrointestinal disorders</i>		Peptic ulcer (with possible peptic ulcer perforation and peptic ulcer haemorrhage); intestinal perforation; gastric haemorrhage; pancreatitis; esophagitis ulcerative; oesophagitis; abdominal distension; abdominal pain; diarrhoea; dyspepsia; nausea;

MedDRA (v15) System Organ Class		Undesirable Effect
		vomiting; dysgeusia; peritonitis (peritonitis may be the primary presenting sign or symptom of a gastrointestinal disorder such as perforation, obstruction or pancreatitis)
<i>Hepatic disorders</i>		Hepatomegaly, hepatitis, drug-induced liver injury, liver failure
<i>Skin and subcutaneous disorders</i>		Angioedema; hirsutism; petechiae; ecchymoses; skin atrophy; erythema; hyperhidrosis; skin striae; rash; pruritus; urticaria; acne; skin hypopigmentation; skin hyperpigmentation; allergic dermatitis; burning or tingling (especially in the perineal area after intravenous injection); cutaneous and subcutaneous atrophy; dry scaly skin; sterile abscess; thinning scalp hair; Kaposi's sarcoma
<i>Musculoskeletal and connective tissue disorders</i>		Muscular weakness; myalgia; myopathy; muscle atrophy; osteoporosis; osteonecrosis; pathological; fracture; neuropathic arthropathy; arthralgia; growth retardation
<i>Reproductive system and breast disorders</i>		Menstruation irregular; increased or decreased motility and number of spermatozoa.
<i>General disorders and administration site conditions</i>		Impaired healing; fatigue; malaise; injection site reaction; oedema peripheral
<i>Investigations</i>		Urine calcium increased; blood potassium decreased; Carbohydrate tolerance decreased; intraocular pressure increased; aminotransferase increased; aspartate aminotransferase increased; blood alkaline phosphatase increased; suppression of reactions to skin tests; post-injection flare (following intra-articular use); blood urea increased
<i>Injury, poisoning and procedural complications</i>		Spinal compression fracture; tendon rupture (particularly of the Achilles tendon)

The following adverse reactions have been reported with the following routes of administration:
Intrathecal/Epidual: Arachnoiditis, functional gastrointestinal disorder, bladder dysfunction, headache, meningitis, paraparesis/paraplegia, seizures, sensory disturbances.

DRUG INTERACTIONS

Overview

Methylprednisolone is a cytochrome P450 enzyme (CYP) substrate and is mainly metabolized by the CYP3A enzyme. CYP3A4 is the dominant enzyme of the most abundant CYP subfamily in the liver of adult humans. It catalyzes 6 β -hydroxylation of steroids, the essential Phase I metabolic step for both endogenous and synthetic corticosteroids. Many other compounds are also substrates of CYP3A4, some of which (as well as other drugs) have been shown to alter glucocorticoid metabolism by induction (upregulation) or inhibition of the CYP3A4 enzyme.

CYP3A4 INHIBITORS – Drugs that inhibit CYP3A4 activity generally decrease hepatic clearance and

increase the plasma concentration of CYP3A4 substrate medications, such as methylprednisolone. In the presence of a CYP3A4 inhibitor, the dose of methylprednisolone may need to be titrated to avoid steroid toxicity.

CYP3A4 INDUCERS – Drugs that induce CYP3A4 activity generally increase hepatic clearance, resulting in decreased plasma concentration of medications that are substrates for CYP3A4. Coadministration may require an increase in methylprednisolone dosage to achieve the desired result.

CYP3A4 SUBSTRATES – In the presence of another CYP3A4 substrate, the hepatic clearance of methylprednisolone may be affected, with corresponding dosage adjustments required. It is possible that adverse events associated with the use of either drug alone may be more likely to occur with coadministration.

NON-CYP3A4-MEDIATED EFFECTS – Other interactions and effects that occur with methylprednisolone are described in the Table below.

Drug-Drug Interactions

The following table provides a list and description of the most common and/or clinically important drug interactions or effects with methylprednisolone.

COMMON INTERACTIONS SEEN WITH METHYLPREDNISOLONE SODIUM SUCCINATE AND OTHER DRUG PRODUCTS				
CLASS OF DRUG	DRUG(S) INVOLVED		CLINICAL IMPLICATION	MECHANISM
Antibiotics/ Antifungals	Troleandomycin Erythromycin Ketoconazole Itraconazole		Enhanced clinical effects and side effects of methylprednisolone.	Enzyme inhibition: Reduced MP elimination.
	Isoniazid		There is a potential effect of methylprednisolone to increase the acetylation rate and clearance of isoniazid	CYP3A4 Inhibitor
	Rifampin		May reduce efficacy; dosage adjustment may be required.	Enzyme induction, increased clearance.
	Clarithromycin Erythromycin			CYP3A4 Inhibitor (and Substrate)
Anticholinergics	Pancuronium		Partial reversal of neuromuscular block.	
	Neuromuscular blockers		1) An Acute myopathy has been reported with concomitant high doses of corticosteroids and anticholinergics, such as neuromuscular blocking drugs. 2) Antagonism of the	

			neuromuscular blocking effects of pancurinium and vecuronium has been reported in patients taking corticosteroids. This interaction may be expected with all competitive neuromuscular blockers.	
Anticholinesterases	Neostigmine, Pyridostigmine		Precipitation of myasthenic crisis. Anticholinesterase agents should be withdrawn at least 24 hours before initiating corticosteroid therapy.	
	Steroids		Steroids may reduce the effects of anticholinesterases in myasthenia gravis.	
Anticoagulants	Oral Anticoagulants or Heparin		The effect of methylprednisolone on oral anticoagulants is variable. There are reports of enhanced as well as diminished effects of anticoagulants when given concurrently with corticosteroids. Therefore, coagulation indices should be monitored to maintain the desired anticoagulant effects.	
	Warfarin		Inhibition of response to warfarin. Coagulation indices should be monitored frequently to maintain desired anticoagulant effect.	
Anticonvulsants	e.g. Phenobarbital Phenytoin Carbamazepine		May reduce methylprednisolone efficacy. Monitor clinical response. Adjust dose if necessary.	Enzyme induction: increased clearance of methylprednisolone
Antidiabetics	e.g. Insulin, Glibenclamide, Metformin		Because corticosteroids may increase corticosteroid concentrations, dosage adjustments of antidiabetic agents may be required.	Diabetogenic effects of corticosteroid.
Antiemetics	Aprepitant Fosaprepitant			CYP3A4 Inhibitor
Antihyper-Cholesterolemics and Antidiarrheals	Cholestyramine		May increase the clearance of corticosteroids.	

Antihypertensive Agents	All Antihypertensives		May result in partial loss of hypertensive control.	Mineralocorticoid effect of corticosteroid leading to raised blood pressure.
Antitubercular Drugs	Isoniazid		Serum concentrations of isoniazid may be decreased.	
Antivirals	HIV-Protease inhibitors		Protease inhibitors, such as indinavir and ritonavir, may increase plasma levels of corticosteroids. Corticosteroids may induce the metabolism of HIV-protease inhibitors resulting in reduced plasma concentrations.	CYP3A4 Inhibitor (and Substrate)
Aromatase Inhibitors	Aminoglutethimide		May lead to a loss of corticosteroid-induced adrenal suppression. Aminoglutethimide-induced adrenal suppression may exacerbate endocrine changes caused by prolonged glucocorticoid treatment.	
Cardioactive Drugs	Digoxin and related Glycosides Calcium Channel Blockers e.g. Diltiazem		Potential of digoxin toxicity. Increased risk of arrhythmias due to hypokalemia.	Corticosteroid induced potassium loss (mineralocorticoid effect) CYP3A4 Inhibitor and Substrate)
Potassium- Depleting Agents	All potassium losing diuretics e.g. Furosemide Amphotericin B Xanthenes Beta2 agonists		Enhanced toxicity. Monitor K ⁺ levels and supplement if necessary. Development of hypokalemia. When corticosteroids are administered concomitantly with potassium-depleting agents (i.e. diuretics), patients should be observed closely for development of hypokalemia. There is also an increased risk for hypokalemia with concurrent use of corticosteroids with amphotericin B, xanthenes, or beta2 agonists.	Potassium loss.

Estrogens, Including Oral Contraceptives	Ethinylestradiol/ Norethindrone		May decrease hepatic metabolism of certain corticosteroids, thereby increasing their effect.	CYP3A4 Inhibitor
Immunizing Agents	Live Vaccine: Poliomyelitis, BCG, Mumps, Measles, Rubella, Smallpox		May see increased toxicity from vaccine. Disseminated viral disease may occur.	Corticosteroid induced immunosuppression
	Killed Virulent Vaccines		Reduced response to vaccine.	Impaired immune response
Immuno- suppressants	Methotrexate Azathioprine		May allow reduced dose of corticosteroid.	Synergistic effect on disease state.
	Cyclosporine (CYA)		1) Mutual inhibition of metabolism occurs with concurrent use of cyclosporine and methylprednisolone, which may increase the plasma concentrations of either or both drugs. Therefore, it is possible that adverse events associated with the use of either drug alone may be more likely to occur upon coadministration. 2) Convulsions have been reported with concurrent use of methylprednisolone and cyclosporine.	CYP3A4 Inhibitor (and Substrate)
	Cyclosporine Tacrolimus			CYP3A4 Substrate
	Cyclophosphamide Tacrolimus			
Psychotherapeutic	Anxiolytics Antipsychotics		Recurrence or poor control of CNS symptoms. May require dose adjustment.	CNS effects of corticosteroid.

Nonsteroidal Anti-inflammatory Agents (NSAIDs)	e.g. Aspirin		<p>1) Concomitant use of nonsteroidal anti-inflammatory agents and corticosteroids increases risk of gastrointestinal side effects.</p> <p>2) Aspirin should be used cautiously in conjunction with corticosteroids in hypoprothrombinemia.</p> <p>3) There may be increased incidence of gastrointestinal bleeding and ulceration when corticosteroids are given with NSAIDs.</p> <p>4) Methylprednisolone may increase the clearance of high-dose aspirin, which can lead to decreased salicylate serum levels. Discontinuation of methylprednisolone treatment can lead to raised salicylate serum levels, which could lead to an increased risk of salicylate toxicity.</p>	<p>Increased clearance and decreased plasma level.</p> <p>CYP3A4 Inhibitor</p>
Sympathomimetic Agents	e.g. Salbutamol		Increased efficacy and potentially increased toxicity.	Increased response to sympathetic agents.

Incompatibilities

To avoid compatibility and stability problems, it is recommended that methylprednisolone sodium succinate be administered separately from other compounds that are administered via the IV route of administration. Drugs that are physically incompatible in solution with methylprednisolone sodium succinate include, allopurinol sodium, doxapram hydrochloride, tigecycline, diltiazem hydrochloride, calcium gluconate, vecuronium bromide, rocuronium bromide, cisatracurium besylate, glycopyrrolate, propofol (see also DOSAGE AND ADMINISTRATION, Compatibility).

Drug-Food Interactions

Interactions with food have not been established.

Grapefruit juice is a CYP3A4 inhibitor. See DRUG INTERACTIONS, CYP3A4 INHIBITORS above.

Drug-Herb Interactions

Interactions with herbal products have not been established.

Drug-Laboratory Interactions

Skin tests: Corticosteroids may suppress reactions to skin tests.

Drug-Lifestyle Interactions

Dizziness, vertigo, visual disturbances and fatigue are possible side effects associated with corticosteroid

use. If affected, patients should not drive or operate machinery.

DOSAGE AND ADMINISTRATION

Recommended Dose and Dosage Adjustment

Corticosteroid therapy is an adjunct to, and not replacement for, conventional therapy.

As adjunctive therapy in life threatening conditions (e.g., shock states), the recommended dose of Methylprednisolone Sodium Succinate for Injection USP is 30 mg per kg, given intravenously over a period of at least 30 minutes. This large dose may be repeated every 4 - 6 hours for up to 48 hours.

In other indications, initial dosage will vary from 10 to 500 mg depending on the clinical problem being treated. Larger doses may be required for short-term management of severe, acute conditions. Therapy may be initiated by administering Methylprednisolone Sodium Succinate for Injection USP intravenously over a period of at least 5 minutes (e.g., doses up to 250 mg) to at least 30 minutes (e.g., doses greater than 250 mg). Subsequent doses may be given intravenously or intramuscularly at intervals dictated by the patient's response and clinical condition.

Methylprednisolone Sodium Succinate for Injection USP in doses of 40 to 120 mg administered as retention enemas or by continuous drip three to seven times weekly for periods of two or more weeks have been shown to be a useful adjunct in the treatment of some patients with ulcerative colitis. Many patients can be controlled with 40 mg of Methylprednisolone Sodium Succinate for Injection USP administered in from 1 to 10 fluid ounces of water depending on the degree of involvement of the inflamed colonic mucosa. Other accepted therapeutic measures should, of course, be instituted.

Administration

Methylprednisolone Sodium Succinate for Injection USP may be administered by intravenous or intramuscular injection or by intravenous infusion, the preferred method for initial emergency use being intravenous injection.

To administer intravenous (or intramuscular) injection, prepare solution as per instructions.

Reconstitution:

1. Remove the protective plastic flip-top seal.
2. Swab the rubber stopper with an antiseptic solution and introduce the required quantity of the diluent by means of a syringe into the vial.
3. Shake the vial thoroughly to dissolve the powder content.
4. Withdraw the dose in the usual manner with the help of a syringe, unused portion should be discarded.

Reconstitute with Sterile Water for Injection, or, if required, Bacteriostatic Water for Injection as follows:

RECONSTITUTION TABLE

Size (mg/vial)	Quantity of Diluent (mL)	Approx. withdrawable volume (mL)	Nominal Concentration (mg/mL)
500	7.8	8	62.5
1000	15.6	16	62.5

The supplied Bacteriostatic Water for Injection in each pack of Methylprednisolone Sodium Succinate for Injection USP contains benzyl alcohol as preservative. **SEE WARNINGS AND PRECAUTIONS, Pediatrics.**

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit.

Store powder or reconstituted solution at 15 to 25 C. When reconstituted with Bacteriostatic Water for Injection, the resulting solution may be stored for up to 48 hours at 15 -25 C. When reconstituted with Sterile Water for Injection, the resulting solution may be stored for up to 24 hours at 15 -25 C due to risk of microbial contamination. The solution with a slight haze should be discarded. The vials are single dose vials. Discard unused portion.

To prepare solutions for IV infusion:

First prepare the solution for injection as directed. Further dilution should be done immediately after reconstitution.

If desired, the medication may be administered in diluted solutions by admixing the reconstituted product with 0.9% Sodium Chloride Injection or with Dextrose 5% Injection. Concentrations of 1mg/mL are physically and chemically stable for 24 hours at 15 to 25 C.

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit.

Compatibility

The compatibility and stability of Methylprednisolone Sodium Succinate for Injection USP, in solutions and with other drugs in intravenous admixtures is dependent on admixture pH, concentration, time, temperature, and the ability of methylprednisolone to solubilize itself. Thus, to avoid compatibility and stability problems, whenever possible it is recommended that Methylprednisolone Sodium Succinate for Injection USP be administered separate from other drugs and as either I.V. push, through an I.V. medication chamber, or as an I.V. "piggy-back" solution.

OVERDOSAGE

Treatment of acute overdosage is by supportive and symptomatic therapy. For chronic overdosage in the face of severe disease requiring continuous steroid therapy, the dosage of the corticosteroid may be reduced only temporarily, or alternate day treatment may be introduced. Continuous overdosage would require careful gradual reduction of dosage in order to prevent the occurrence of acute adrenal insufficiency.

Methylprednisolone is dialyzable.

For management of a suspected drug overdose, contact your regional Poison Control Centre.

ACTION AND CLINICAL PHARMACOLOGY

Pharmacodynamics

Methylprednisolone is a potent anti-inflammatory steroid.

Methylprednisolone sodium succinate has the same metabolic and anti-inflammatory actions as methylprednisolone. When given parenterally and in equimolar quantities, the two compounds are equivalent in biologic activity. The relative potency of methylprednisolone sodium succinate and hydrocortisone sodium succinate, following intravenous administration, is at least four to one. This is in good agreement with the relative oral potency of methylprednisolone and hydrocortisone.

The metabolism and excretion of methylprednisolone sodium succinate is similar to that of other corticosteroids.

It influences carbohydrate, protein, fat and purine metabolism, electrolyte and water balance, and the functional capacities of the cardiovascular system, the kidney, skeletal muscle, the nervous system and other organs and tissues.

Exceeding prednisolone in anti-inflammatory potency and having even less tendency than prednisolone to induce retention of sodium and water, methylprednisolone sodium succinate offers the use of lower doses with an enhanced split between anti-inflammatory and mineralocorticoid activities. Thus methylprednisolone sodium succinate may be indicated for emergency use in patients in whom increased sodium retention would be hazardous.

The relative potency of methylprednisolone sodium succinate and hydrocortisone sodium succinate, as indicated by depression of eosinophil count, following intravenous administration, is at least four to one. This is in good agreement with the relative oral potency of methylprednisolone and hydrocortisone. Studies indicate that the administration of methylprednisolone results in an appreciable prolongation of plasma steroid levels over those obtained following equivalent doses of hydrocortisone or prednisolone. The following table illustrates this prolongation of blood levels expressed as the half-life in minutes of the 17-hydroxy-corticosteroid levels obtained following intravenous administration of methylprednisolone, prednisolone and hydrocortisone.

COMPOUND	DOSE	HALF-LIFE (minutes)
Methylprednisolone	25 mg	188
Prednisolone	25 mg	69
Hydrocortisone	25 mg	57

Pharmacokinetics

Methylprednisolone pharmacokinetics are linear, independent of the route of administration.

Absorption: After a 40 mg intramuscular dose of methylprednisolone sodium succinate to fourteen

healthy adult male volunteers, the average peak concentration of 454 mg/mL was achieved at 1 hour. At 12 hours, the methylprednisolone plasma concentration declined to 31.9 mg/mL. No methylprednisolone was detected 18 hours after dosing. Based on the area under the time-concentration curve, an indication of total drug absorbed, intramuscular methylprednisolone sodium succinate was found to be bioequivalent to the same dose administered intravenously.

The sodium succinate ester of methylprednisolone is rapidly and extensively converted to the active methylprednisolone moiety after all routes of administration. Extent of absorption of free methylprednisolone following IV and IM administrations were found to be bioequivalent and significantly greater than those following administration of the oral solution and oral methylprednisolone tablets. Since the extent of methylprednisolone absorbed following the IV and IM treatment was equivalent, in spite of the greater amount of the hemisuccinate ester reaching the general circulation after IV administration, it appears that the ester is converted in the tissue after IM injection, with subsequent absorption as free methylprednisolone.

Distribution: Methylprednisolone is widely distributed into the tissues, crosses the blood-brain barrier and is secreted in breast milk. Its apparent volume of distribution is approximately 1.4 L/kg. The plasma protein binding of methylprednisolone in humans is approximately 77%.

Metabolism: In humans, methylprednisolone is metabolized in the liver to inactive metabolites, primarily 20 α -hydroxymethylprednisolone and 20 β -hydroxymethylprednisolone. Metabolism in the liver occurs primarily via the CYP3A4 (see DRUG INTERACTIONS).

Methylprednisolone, like many CYP3A4 substrates, may also be a substrate for the ATP-binding cassette (ABC) transport protein p-glycoprotein (P-gp), influencing tissue distribution and interactions with other medicines modulated by P-gp.

Excretion: The mean elimination half-life for total methylprednisolone is in the range of 1.8 to 5.2 hours. Total clearance is approximately 5 to 6 mL/min/kg.

STORAGE AND STABILITY:

Unreconstituted vials:

Store at 15 -25°C. Protect from light.

Reconstituted Solution:

When reconstituted with Bacteriostatic Water for Injection, the resulting solution may be stored for up to 48 hours at 15 -25 °C.

When reconstituted with Sterile Water for Injection, the resulting solution may be stored for up to 24 hours at 15 -25 °C due to risk of microbial contamination. The solution with a slight haze should be discarded. Discard unused portion.

DOSAGE FORMS, COMPOSITION AND PACKAGING

Methylprednisolone Sodium Succinate for Injection USP is a sterile, lyophilized powder available in clear glass vial, equivalent to methylprednisolone 500 mg/vial and 1000 mg/vial. Each pack of Methylprednisolone Sodium Succinate for Injection USP 500 mg per vial or 1000 mg per

vial includes respectively a vial of 10 mL and 20 mL of Bacteriostatic Water for Injection (supplied as diluent) with Benzyl alcohol 0.9%v/v as preservative.

Composition

The respective vials of Methylprednisolone Sodium Succinate for Injection USP contain:

Ingredients	500 mg vial	1000 mg vial
POWDER		
Methylprednisolone as the sodium succinate (mg/vial)	500 mg	1 000 mg
Sodium Dihydrogen Phosphate	6.4 mg	12.8 mg
Disodium Hydrogen Phosphate	69.8 mg	139.6 mg
DILUENT		
Bacteriostatic Water for Injection (with Benzyl Alcohol 0.9% v/v as preservative)	10 mL	20 mL

PART II: SCIENTIFIC INFORMATION

PHARMACEUTICAL INFORMATION


Drug Substance

Proper name: methylprednisolone sodium succinate for injection USP
(methylprednisolone sodium succinate is prepared in situ from methylprednisolone hemisuccinate with the aid of sodium hydroxide)

Chemical name: pregna-1,4-diene-3,20-dione,21-(3-carboxy-1-oxopropoxy)-11,17-dihydroxy-6-methyl-monosodium salt, (6 α ,11 β)

Molecular mass: 496.53

Structural formula:



Physicochemical properties: white, or nearly white, odourless hygroscopic, amorphous solid, very soluble in water and in alcohol, insoluble in chloroform, very slightly soluble in acetone, melting point of 228° to 237°C, pka of 4.6, partition coefficient (butyronitrile-water) of 0.03 at pH 8.5

CLINICAL TRIALS

Hypersensitivity and Dermatologic Conditions

Status Asthmaticus

In a double-blind, placebo-controlled, randomized trial, the use of intravenous methylprednisolone (125 mg), given on presentation in the emergency room in addition to standard emergency treatments for asthma, reduced the need for hospital admission in acutely ill patients with bronchial asthma. Nine of 48 patients (19 percent) treated with methylprednisolone required hospital admission compared with 23 of 49 patients (47 percent) in the control group ($p < 0.003$).

Pemphigus Vulgaris

A small ($n=15$) retrospective study compared high-dose pulsed methylprednisolone sodium succinate to oral prednisone in patients with pemphigus vulgaris. Methylprednisolone sodium succinate was administered intravenously ($n=9$); the dose varied from 250 to 1000 mg/day for 2 to 5 days. Four of 6 responders to methylprednisolone sodium succinate maintained a remission without prednisone for almost 2 years. Patients in the control group ($n=6$) treated with prednisone required long-term treatment with higher doses of prednisone, and none of the patients maintained a long-term remission.

Acute Systemic Lupus Erythematosus

High-dose, intravenous methylprednisolone pulse therapy in 34 patients (30 adults and 4 adolescents) with lupus nephritis was evaluated. The 30 adult patients received 1 g of methylprednisolone intravenously over 30 minutes on 3 successive days, while the 4 adolescents received a 15 mg/kg/day dose for 3 days. Twelve of the 34 patients responded to treatment, as indicated by at least a 20% improvement in renal function and corresponding improvement in creatinine clearance levels. These improvements were maintained for at least 6 months in 60% of patients who responded to treatment.

Ulcerative Colitis

In a prospective, single-blind study of 60 patients with active ulcerative colitis, patients were randomized to receive either sucralfate enemas (20 g/100 ml) or methylprednisolone enemas (20 mg/100 ml). The enemas were administered twice daily for the first week and then once daily for three weeks. Results showed similar reductions in diarrhea and rectal bleeding at two weeks and at four weeks in the two groups. Sigmoidoscopic examination of the rectal mucosa demonstrated similar significant improvement in the macroscopic appearance of the rectal mucosa in both groups (8.28 to 6.20 in sucralfate group, $p < 0.02$; and 8.72 to 6.36 in the methylprednisolone treated group, $p < 0.04$). Histological assessment of the rectal biopsies taken at entry into the study and following four weeks of therapy also revealed similar improvements in the two groups.

Organ Transplants

A prospective, controlled study was conducted among 100 renal transplant patients to compare two different regimens of immunosuppressive therapy. In the study, 86 patients received kidneys from cadavers and 14 patients received kidneys from living, related donors. Patients were assigned to receive either double therapy (methylprednisolone plus cyclosporine) or triple therapy (methylprednisolone plus cyclosporine and azathioprine). In both groups, patients were given intravenous pulse doses of 0.5 g methylprednisolone at the moment of transplantation. Oral methylprednisolone was subsequently administered in a single morning dose of 16 mg until the end of the third month. Patients then received 12 mg/day oral methylprednisolone until the end of month 6, and a maintenance dosage of 8 mg/day thereafter. The results were similar with both regimens. No significant differences between groups were reported in the 2-year patient and kidney survival rates.

DETAILED PHARMACOLOGY

See **ACTION AND CLINICAL PHARMACOLOGY, Pharmacodynamics and Pharmacokinetics** sections.

TOXICOLOGY

Acute Toxicology

The acute LD₅₀ of methylprednisolone sodium succinate intraperitoneally in the mouse is 850 mg/kg. The oral LD₅₀ of this drug in the rat is 5150 mg/kg. Dogs receiving single intravenous injections of methylprednisolone sodium succinate in doses of 4.4 to 6.4 mg/kg were free from clinical signs of drug intoxication during the 24 hour post-injection observation period.

Carcinogenesis:

Methylprednisolone has not been evaluated in rodent carcinogenicity studies.

Variable results have been obtained with other glucocorticoids tested for carcinogenicity in mice and rats. However, published data indicate that several related glucocorticoids including budesonide, prednisolone, and triamcinolone acetonide can increase the incidence of hepatocellular adenomas and carcinomas after oral administration in drinking water to male rats. These tumorigenic effects occurred at doses which were less than the typical clinical doses on a mg/m² basis.

Mutagenesis:

Methylprednisolone has not been evaluated for genotoxicity.

However, methylprednisolone sulfonate, which is structurally similar to methylprednisolone sodium succinate, was not mutagenic with or without metabolic activation in *Salmonella typhimurium*, or in a mammalian cell gene mutation assay using Chinese hamster ovary cells. Methylprednisolone sulteptanate did not induce unscheduled DNA synthesis in primary rat hepatocytes. Moreover, a review of published data indicates that prednisolone farnesylate (PNF), which is structurally similar to methylprednisolone, was not mutagenic with or without metabolic activation in *Salmonella typhimurium* and *Escherichia coli* strains. In a Chinese hamster fibroblast cell line, PNF produced a slight increase in the incidence of structural chromosomal aberrations with metabolic activation at the highest concentration tested.

Reproductive toxicity:

Corticosteroids have been shown to reduce fertility when administered to rats. Male rats were administered corticosterone at doses of 0, 10, and 25 mg/kg/day by subcutaneous injection once daily for 6 weeks and mated with untreated females. The high dose was reduced to 20 mg/kg/day after Day 15. Decreased copulatory plugs were observed, which may have been secondary to decreased accessory organ weight. The numbers of implantations and live fetuses were reduced in untreated females mated with males treated at the administered doses of 10 and 25 mg/kg/day.

Corticosteroids have been shown to be teratogenic in many species when given in doses equivalent to the human dose. In animal reproduction studies, glucocorticoids such as methylprednisolone have been shown to increase the incidence of malformations (cleft palate, skeletal malformations), embryo-fetal lethality (e.g., increase in resorptions), and intra-uterine growth retardation.

REFERENCES

1. Antman EM, Braunwald E. Acute myocardial infarction. In: Fauci AS, *et al*, eds. Harrison's Principles of Internal medicine. 1998:1356-7.
2. Baethge BA, Lidsky MD. Intractable hiccups associated with high-dose intravenous methylprednisolone therapy. *Ann Intern Med.* 1986;104:58-9.
3. Bagheri H, Cismondo S, Montastruc JL. Drug-induced hiccup: A review of the France pharmacologic vigilance database. *Therapie.* 1999;54(1):35-9. (French)
4. Burkle WS. Cyclosporin pharmacokinetics and blood level monitoring. *Drug Intell Clin Pharm* 1985;19:101-5.
5. Cowley RA, Trump BF. Pathophysiology of shock, anoxia and ischemia. Baltimore, MD. Williams and Wilkins 1982;692-3.
6. CRASH trial collaborators. Effect of intravenous corticosteroids on death within 14 days in 10,008 adults with clinically significant head injury (MRC CRASH trial): randomized placebo-controlled trial. *The Lancet* 2004; 364: 1321-8.
7. Dietzman RH, Bloch JH, Lillehei RC. Treatment of shock. *GP* 1967;36(3):135-41.
8. Dietzman RH, Manax WG, Lillehei RC. Shock: Mechanism and Therapy. *Can Anaes Soc J* 1967;14(4):276-86.
9. Dugger JA. Parenteral methylprednisolone acetate in management of allergy and contact dermatitis in children. *Clinical Medicine* 1962;69(10):2253-7.
10. Fekety R. Infections associated with corticosteroids and immunosuppressive therapy. In: Gorbach SL, Bartlett JG, Blacklow NR, eds. *Infectious Diseases.* Philadelphia: WB Saunders Company 1992:1050-1.
11. Fox JL. Use of methylprednisolone in intracranial surgery. *Medical Annals of District of Columbia* 1965;34(6):261-6.
12. Goldberg LC. Routine use of intravenous 6-methylprednisolone. *Journal of Michigan State Medical Society* 1960;59:1049-51.
13. Kimberly RP, Lockshin MD, Sherman RL *et al*. High-dose intravenous methylprednisolone pulse therapy in systemic lupus erythematosus. *Am J Med.* 1981;70:817-24.
14. Lillehei RC, Longerbeam JK, Bloch JH, Manax WG. The nature of irreversible shock: Experimental and clinical observations. *Ann Surg* 1964;160(4):682-710.
15. Lillehei RC, Longerbeam JK, Bloch JH, Manax WG. The modern treatment of shock based on physiologic principals. *Clin Pharmacol Ther* 1964;5(1):63-101.

16. Littenberg B, Gluck EH. A controlled trial of methylprednisolone in the emergency treatment of acute asthma. *N Engl J Med*. 1986;314(3):150-2.
17. Marcus PS, Sheehan JC, Comunale FL. Corticosteroids in hypovolemic shock. *Anesthesiology*, Dept. of Anesthesiology, Boston City Hospital 1966;27(2):221.
18. Melby JC. Corticosteroids in shock in man and in animals. *The New Physician* 1964;13:425-8.
19. Panes J, Esteve M, Cabre E et al. Comparison of heparin and steroids in the treatment of moderate and severe ulcerative colitis. *Gastroenterology* 2000; 119(4):903-8.
20. Ponticelli C, Tarantino A, Montagnino G et al. A randomized trial comparing triple-drug and double-drug therapy in renal transplantation. *Transplantation*. 1988;45:913-18.
21. Robins J. Modern Concepts in the management of shock. *Obstet Gynecol (View and Reviews Section)* 1966;28(1):130-8.
22. Sauerland S, Maegele M. A CRASH landing in severe head injury. *The Lancet* 2004; 364:1291-92.
23. Sholter DE, Armstrong PW. Adverse effects of corticosteroids on the cardiovascular system. *Can J Cardiol*. 2000;16:505-11.
24. Speed Rogers J, Hudson FM. Parenteral corticosteroids in the management of acute brain stem injuries. *W V Med J* 1963;59(8):18-20
25. Stinson EB, Dong E Jr, Bieber CP, Schroeder JS, Shumway NE. Cardiac Transplantation in Man. I. Early Rejection. *JAMA* 1969;207(12):2233-47.
26. Stubbs SS, Morrell RM. Intravenous methylprednisolone sodium succinate: Adverse reactions reported in association with immunosuppressive therapy. *Transplantation Proceedings* 1973;5(2):1145-6.
27. Stuck AE, Minder CE, Frey FJ. Risk of infectious complications in patients taking glucocorticoids. *Rev Infect Dis* 1989;11(6):954-63.
28. Turcotte JG, et al. Rejection crises in human renal transplant recipients. *Arch Surg* 1972;105:230-6.
29. Turlapaty PDMV, Altura BM. Extracellular magnesium ions control calcium exchange and content of vascular smooth muscle. *Eur J Pharmacol* 1978;52(3-4):421-3.
30. Weissman G, Thomas L. The effects of corticosteroids upon connective tissue and lysosomes. *Recent Prog Horm Res* 1964;20:215-45.
31. Werth VP. Treatment of pemphigus vulgaris with brief, high-dose intravenous glucocorticoids. *Arch Dermatol* 1996;132:1435-9.

32. Wilson RD, Steroids stem “Burn Lung Syndrome”. Medical News Section, JAMA 1969;208 (1):26.
33. Woods JE, et al. High-dosage intravenously administered methylprednisolone in renal transplantation. JAMA 1973;223(8):896-9.
34. Zahony I, Winnie AP, Collins VJ. Delayed postoperative effects of antihypertensive therapy. Illinois Medical Journal 1962;Aug:155-8.
35. Solu-Medrol Product Monograph, Pfizer Canada, Control No.205330, 13 July 2017.

PART III: CONSUMER INFORMATION**^{Pr}Methylprednisolone Sodium Succinate for Injection USP**

This leaflet is part III of a three-part "Product Monograph" published when Methylprednisolone Sodium Succinate for Injection USP (methylprednisolone sodium succinate) was approved for sale in Canada and is designed specifically for Consumers. This leaflet is a summary and will not tell you everything about Methylprednisolone Sodium Succinate for Injection USP. Contact your doctor or pharmacist if you have any questions about the drug.

ABOUT THIS MEDICATION**What the medication is used for:**

Methylprednisolone Sodium Succinate for Injection USP is used in the treatment of various conditions such as allergic reactions or inflammation.

What it does:

Methylprednisolone Sodium Succinate for Injection USP contains a synthetic corticosteroid and is usually used for short periods in severe conditions to decrease inflammation.

When it should not be used:

- you have any allergies to this medicine or to any of the ingredients of this medication.
- for fungal infections
- for patients taking live or live, attenuated vaccines
- for injection into the brain or the spine
- for a blood condition called idiopathic thrombocytopenic purpura (low platelet count), if Methylprednisolone Sodium Succinate for Injection USP is administered intramuscularly

Except for short-term or emergency use such as severe allergic reactions, Methylprednisolone Sodium Succinate for Injection USP should not be given to patients with:

- viral diseases including vaccinia (cowpox), varicella (chickenpox), and herpes simplex of the eye
- tuberculosis
- serious mental disorder (psychoses)
- Cushing's syndrome (abnormal bodily condition caused by excess corticosteroids)
- a stomach ulcer
- altered kidney function

Methylprednisolone Sodium Succinate for Injection USP should not be given to premature infants because diluent for reconstitution contains benzyl alcohol.

Patients taking Methylprednisolone Sodium Succinate for Injection USP should not receive live vaccines.

What the medicinal ingredient is:

methylprednisolone sodium succinate

What the important nonmedicinal ingredients are:

Methylprednisolone Sodium Succinate for Injection USP contains the following nonmedicinal ingredients: sodium phosphate, sodium acid phosphate, sodium hydroxide and water for injection-

The diluent for reconstitution of the vials is Bacteriostatic Water for Injection, which contains benzyl alcohol. Benzyl alcohol has been reported to be associated with fatal "Gaspings Syndrome" in premature infants.

What dosage forms it comes in:

Methylprednisolone Sodium Succinate for Injection USP comes in vials containing sterile powder for intravenous or intramuscular injection or for intravenous infusion. The available formulations are:

- 500 mg Vial
- 1 g Vial

WARNINGS AND PRECAUTIONS**BEFORE you use Methylprednisolone Sodium Succinate for Injection USP talk to your doctor or pharmacist if:**

- you have an infection (such as herpes simplex, chicken pox, tuberculosis, threadworm)
- you recently had myocardial infarction (heart attack)
- you have thromboembolic disorders (bleeding or blood clotting problems)
- you have brittle bone (osteoporosis)
- you have high blood pressure (hypertension)
- you have water retention (oedema)
- you have seizures (fits)
- you have thyroid problems (hypothyroidism or hyperthyroidism)
- you have muscle pain or muscle weakness (such as myasthenia gravis)
- you have skin cancer (Kaposi's sarcoma)
- you have certain eye diseases such as glaucoma, cataracts, herpes infection
- you have kidney disease
- you have liver disease
- you have heart disease
- you have diabetes (high blood sugar)
- you have certain mental or mood conditions (such as depression)
- you have stomach or gut problems (ulcer, ulcerative colitis)
- you have low potassium or calcium
- you have Cushing's disease (caused by an excess of cortisol hormone)
- you have weak immune response
- you have thrombophlebitis (vein inflammation)
- you are pregnant, planning to become pregnant or are breast-feeding (nursing).
- you had any prior use of Methylprednisolone Sodium Succinate for Injection USP.

INTERACTIONS WITH THIS MEDICATION

Tell your doctor or pharmacist about all prescription and non-prescription medications you are using. The following may interact with Methylprednisolone Sodium Succinate for Injection USP:

- Antibiotics/Antifungals (e.g. rifampin, ketoconazole, troleandomycin, erythromycin and amphotericin B)
- Anticholinesterase (drugs that prevent the elimination of a neurotransmitter, acetylcholine. e.g. neostigmine and pyridostigmine)
- Drugs that prevent blood clotting (e.g. warfarin or heparin)
- Epilepsy medication (e.g. phenytoin)
- Diabetes medication (e.g. insulin or metformin)
- High blood pressure treatment (e.g. amlodipine or quinapril)
- Diuretics (e.g. furosemide)
- Heart medication (e.g. digoxin, calcium channel blockers)
- Vaccines
- Drugs that suppress the immune system (methotrexate or cyclosporine)
- Neuromuscular Blocking Agents (agents that block signals between nerves and muscles. e.g. pancuronium)
- Drugs that act on the nervous system (e.g. diazepam or clozapine)
- Aspirin and non-steroidal anti-inflammatory medicines (also called NSAIDs) such as ibuprofen
- Sympathomimetic Agents (agents that mimic the effects of adrenaline. e.g. salbutamol)
- Drugs to treat high cholesterol (e.g., cholestyramine)
- Drugs to treat diarrhea
- Drugs to treat tuberculosis
- Hormone replacement therapy or hormonal oral contraceptives
- Aromatase inhibitors (drugs to treat breast or ovarian cancer)
- Immunosuppressants (drugs that suppress or reduce the strength of the body's immune system)

Do not drink grapefruit juice while taking Methylprednisolone Sodium Succinate for Injection USP.

Driving and Using Machines

Methylprednisolone Sodium Succinate for Injection USP may cause dizziness, vertigo, vision problems and fatigue. If you experience these side effects, you should not drive or operate machinery.

PROPER USE OF THIS MEDICATION

Methylprednisolone Sodium Succinate for Injection

USP may be administered by intravenous or intramuscular injection or by intravenous infusion, the preferred method for initial emergency use being intravenous injection.

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit.

Usual dose:

Initial dosage will vary from 10 to 500 mg depending on the clinical problem being treated. Larger doses may be required for short-term management of severe, acute conditions. Therapy may be initiated by administering Methylprednisolone Sodium Succinate for Injection USP intravenously over a period of at least 5 minutes (e.g., doses up to 250 mg) to at least 30 minutes (e.g., doses greater than 250 mg). Subsequent doses may be given intravenously or intramuscularly at intervals dictated by the patient's response and clinical condition. Corticosteroid therapy is used in combination with, and not replacement for, conventional therapy. The dose needs to be gradually decreased when the medication needs to be discontinued after several days of treatment.

Overdose:

If an overdose occurs, Methylprednisolone Sodium Succinate for Injection USP can be eliminated through dialysis. Continuous overdosing would require careful gradual reduction of the dose of the medication.

In case of overdose, contact a health care practitioner, hospital emergency department or regional poison control centre immediately, even if there are no symptoms.

SIDE EFFECTS AND WHAT TO DO ABOUT THEM

Like all medicines, Methylprednisolone Sodium Succinate for Injection USP can have side effects although not everybody gets them.

Methylprednisolone Sodium Succinate for Injection USP may hide symptoms of infections, may cause latent infections becoming active, and may induce infections by normally inoffensive organisms due to lowered body resistance.

Potential side effects with Methylprednisolone Sodium Succinate for Injection USP include:

Allergic Reactions:

- Anaphylaxis (a severe, life-threatening allergic reaction)
- Cardiac arrest
- Bronchospasm (narrowing of the airway)
- Rapid swelling of the skin

Cardiovascular:

- Heart failure
- Heart attack
- Arrhythmia (irregular heartbeat)
- Slow heart beat
- High and low blood pressure
- Fainting

- Blood clots
- Thrombophlebitis (vein inflammation)

Dermatologic:

- Thin fragile skin
- Impaired wound healing
- Ecchymoses (spots caused by ruptured blood vessels)
- Petechiae (reddish spot containing blood that appears in skin)
- Skin changes (depressions) at the injection
- Acne
- Rash
- Itchiness
- Dry scaly skin
- Swelling
- Redness
- Increased sweating
- Lightening or darkening of an area of the skin
- Abscess
- Thinning scalp hair
- Injection site infections

Endocrine and Metabolism:

- Development of Cushingoid state (abnormal bodily condition caused by excess corticosteroids)
- Moon face (enlargement of face and forehead)
- Weight gain
- Abnormal fat deposits
- Suppression of pituitary-adrenal axis (a condition that could lead to disabling the body's responses to physiological stress such as severe infections or trauma)
- Suppression of growth in children
- Abnormal hair growth
- Sodium retention and excretion
- Fluid retention
- Increased urination
- Decreased carbohydrate (sugar) tolerance
- New symptoms of diabetes
- Need for higher doses of insulin or sugar lowering pills in diabetics

Gastrointestinal:

- Stomach ulcer
- Stomach bleeding
- Inflammation of the pancreas and esophagus
- Perforation of the bowel
- Nausea
- Vomiting or altered sense of taste (with rapid administration of large doses)
- Abdominal pain
- Bloating
- Bowel/bladder dysfunction
- Increased or decreased appetite

- Peritonitis

Hepatic:

- Enlarged liver
- Liver injury
- Hepatitis

Musculoskeletal:

- Muscle disease
- Muscle weakness
- Muscle pain
- Loss of muscle mass
- Malaise (feeling of general discomfort or uneasiness)
- Osteoporosis
- Aseptic necrosis (tissue death) Pathologic fractures
- Vertebral compression fractures
- Tendon rupture, particularly of the Achilles tendon Charcot joint disease
- Pain and inflammation of the tissues surrounding the injection site

Neurologic:

- Increased pressure within the skull with oedema and inflammation of the optic nerve
- Seizures
- Headache
- Pain and tenderness
- Impaired sensation, strength and reflexes
- Sensation of tingling, tickling, prickling or burning of a person's skin
- Vertigo
- Meningitis
- Amnesia
- Dizziness

Ophthalmologic:

- Cataracts
- Protrusion of the eyeball
- Increased intraocular pressure
- Glaucoma
- Blindness.

Psychiatric:

- Mental illness
- Depression
- Emotional instability
- Euphoria (intense feelings of well-being, elation, happiness, excitement, and joy)
- Insomnia (difficulty sleeping)
- Mood swings
- Personality changes
- Thoughts of suicide
- Delusion
- Hallucination
- Confusion

- Schizophrenia
- Anxiety
- Irritability

Sexual Function/Reproduction:

- Menstrual irregularities
- Increased or decreased motility and number of sperm

Hematology:

- Above normal white blood cell count
- Above normal cholesterol or triglycerides
- Abnormal blood tests (ex. Liver enzymes and urea)

Other: Methylprednisolone Sodium Succinate for Injection USP may cause abnormal liver tests and may suppress reactions to skin tests. Methylprednisolone Sodium Succinate for Injection USP may also cause hiccups and fatigue.

may be headaches or generally feeling unwell)			
Fast/pounding/irregular heartbeat		√	
Swelling		√	
Cramps and spasms		√	
Vision changes		√	
Increased thirst/urination		√	
Mental/mood changes (such as mood swings, depression, suicidal thinking, agitation, anxiety)		√	
Tendon pain			√
Bone/joint pain			√
Easy bruising/bleeding			√
Pain/redness/swelling at the injection site	√		
Thinning skin	√		
Poor wound healing	√		
Unusual hair growth	√		
Unusual skin growth (nodules or blotches that may be red, purple, brown or black and may be raised)		√	

SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM

Symptom/effect	Talk with your doctor or pharmacist		Seek IMMEDIATE medical attention
	Only if severe	In all cases	
Burst or bleeding ulcers: symptoms of which are stomach pain, bleeding from the rectum, black or bloodstained stools and/or vomiting blood			√
Flare up of a previous Tuberculosis: symptoms of which could be coughing blood or pain in the chest.			√
Serious allergic reaction: symptoms of which include rash, itching/swelling (especially of the face/tongue/throat), severe dizziness and trouble breathing			√
Signs of infection (such as persistent fever/cough/sore throat, painful urination, eye pain/discharge)		√	
High blood pressure (symptoms of which		√	

HOW TO STORE IT

Before Reconstitution: Store Methylprednisolone Sodium Succinate for Injection USP at 15° - 25°C. Protect from light. Keep out of the reach of children.

After Reconstitution: When reconstituted with Bacteriostatic Water for Injection, the resulting solution may be stored for up to 48 hours at 15 -25 °C. When reconstituted with Sterile Water for Injection, the resulting solution may be stored for up to 24 hours at 15 -25 °C due to risk of microbial contamination. Protect from light. Keep out of the reach of children.

REPORTING SIDE EFFECTS

You can help improve the safe use of health products for Canadians by reporting serious and unexpected side effects to Health Canada. Your report may help to identify new side effects and change the product safety information.

3 ways to report:

- *Online at <https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada/adverse-reaction-reporting.html>;*
- *By calling 1-866-234-2345 (toll-free);*
- *By completing a Consumer Side Effect Reporting Form and sending it by:*
 - *Fax to 1-866-678-6789 (toll-free), or*
 - *Mail to: Canada Vigilance Program*
Health Canada, Postal Locator 0701E
Ottawa, ON K1A 0K9

Postage paid labels and the Consumer Side Effect Reporting Form are available at <https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada/adverse-reaction-reporting.html>.

NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

MORE INFORMATION

This document plus the full product monograph, prepared for health professionals may be obtained by contacting the sponsor, Hospira Healthcare Corporation at:
1-800-463-6001.

This leaflet was prepared by
Hospira Healthcare Corporation.
Kirkland, Québec
H9J 2M5

Last revised: October 18, 2017