

PRODUCT MONOGRAPH

Pr CYTARABINE INJECTION BP

100 mg / mL

(100 mg / 1 mL, 500 mg / 5 mL, 1 g / 10 mL, 2 g / 20 mL)

Antileukemic Agent

Sandoz Canada Inc.
145, Jules-Léger
Boucherville, Quebec
J4B 7K8

Date of Revision:
December 06, 2017

Submission Control No: 199911

Table of contents

PART I: HEALTH PROFESSIONAL INFORMATION.....	3
SUMMARY PRODUCT INFORMATION	3
INDICATIONS AND CLINICAL USE.....	3
CONTRAINDICATIONS	3
WARNINGS AND PRECAUTIONS.....	4
ADVERSE REACTIONS.....	9
DRUG INTERACTIONS	13
DOSAGE AND ADMINISTRATION	14
OVERDOSAGE	21
ACTION AND CLINICAL PHARMACOLOGY	21
STORAGE AND STABILITY.....	23
SPECIAL HANDLING INSTRUCTIONS	23
DOSAGE FORMS, COMPOSITION AND PACKAGING	24
PART II: SCIENTIFIC INFORMATION	25
PHARMACEUTICAL INFORMATION.....	25
DETAILED PHARMACOLOGY	26
TOXICOLOGY	26
REFERENCES	28
APPENDIX A.....	36
PART III: CONSUMER INFORMATION.....	38

Pr CYTARABINE INJECTION BP

Cytarabine injection

PART I: HEALTH PROFESSIONAL INFORMATION

SUMMARY PRODUCT INFORMATION

Route of Administration	Dosage Form/Strength	Clinically Relevant Nonmedicinal Ingredients
Intravenous infusion Subcutaneous injection Intrathecal injection	Solution for Injection 100mg / mL (100 mg / 1 mL, 500 mg / 5 mL, 1000 mg / 10 mL and 2000 mg / 20 mL)	Not applicable <i>For a complete listing see DOSAGE FORMS, COMPOSITION and PACKAGING section.</i>

INDICATIONS AND CLINICAL USE

Cytarabine Injection BP (cytarabine) is indicated primarily for induction and maintenance of remission in acute leukemia in both adults and children.

It has been found useful in the treatment of acute myelocytic leukemia, chronic myelocytic leukemia (blast phase), acute lymphocytic leukemia and erythroleukemia. Cytarabine Injection BP may be used alone or in combination with other antineoplastic agents; the best results are obtained with combination therapy.

Children with non-Hodgkin's lymphoma have benefited from a combination drug program (LSA₂L₂) that included cytarabine.

Cytarabine has been used intrathecally in newly diagnosed children with acute lymphocytic leukemia as well as in the treatment of meningeal leukemia.

Cytarabine, in high dose 2-3 g/m² as an IV infusion over 1-3 hours given every 12 hours for 2-6 days with or without additional cancer chemotherapeutic agents, has been shown to be effective in the treatment of poor-risk leukemia, refractory leukemia, and relapsed acute leukemia.

Remissions induced by cytarabine not followed by maintenance treatment have been brief.

CONTRAINDICATIONS

Cytarabine Injection BP is contraindicated in those patients who are hypersensitive to the drug.

Anaphylactic reactions have occurred with cytarabine treatment (see WARNINGS AND PRECAUTIONS, Sensitivity/Resistance).

WARNINGS AND PRECAUTIONS

Serious Warnings and Precautions

Cytarabine Injection BP should be prescribed only by physicians experienced with cancer therapy drugs. Patients should be monitored and blood counts as well as renal and hepatic function tests should be performed regularly (see WARNINGS AND PRECAUTIONS, Hematologic, Hepatic/Biliary/Pancreatic, Renal, Monitoring and Laboratory Tests and OVERDOSAGE).

Do not use a diluent that contains benzyl alcohol when giving to premature or low birth weight infants as benzyl alcohol has been associated with the “gasping syndrome” (see WARNINGS AND PRECAUTIONS, General and Special Populations, Pediatrics). Do not use a diluent that contains benzyl alcohol for high dose therapy or when using intrathecally (see ADVERSE REACTIONS, High Dose Therapy).

The following are clinically significant adverse events:

- Cardiomyopathy with subsequent death (see WARNINGS AND PRECAUTIONS, Cardiovascular and ADVERSE REACTIONS, High Dose Therapy).
- GI toxicity, at times fatal (see WARNINGS AND PRECAUTIONS, Gastrointestinal and ADVERSE REACTIONS, High Dose Therapy).
- Acute pancreatitis (see WARNINGS AND PRECAUTIONS, Hepatic/Biliary/Pancreatic).
- CNS toxicity, severe neurological adverse reactions, paraplegia, necrotizing leukoencephalopathy and spinal cord toxicity. Patients with impaired hepatic or renal function may be at increased risk after high dose Cytarabine Injection BP (see WARNINGS AND PRECAUTIONS, Hepatic/Biliary/Pancreatic, Neurologic and Renal; ADVERSE REACTIONS, High Dose Therapy and Intrathecal Therapy; DRUG INTERACTIONS, Serious Interactions; DOSAGE AND ADMINISTRATION, Meningeal Leukemia – Intrathecal Use, OVERDOSAGE, and ACTION AND CLINICAL PHARMACOLOGY).
- Infection (see WARNINGS AND PRECAUTIONS, Immune and ADVERSE REACTIONS, Infections and Infestations).
- Pulmonary toxicity, adult respiratory distress syndrome and pulmonary edema (see WARNINGS AND PRECAUTIONS, Respiratory and ADVERSE REACTIONS, High Dose Therapy).
- Myelosuppression (see WARNINGS AND PRECAUTIONS, Hematologic; ADVERSE REACTIONS, Blood and Lymphatic System Disorders and OVERDOSAGE).

General

Before instituting a program of combined therapy, the physician should be familiar with the literature, adverse reactions, warnings and precautions, and contraindications applicable to all the

drugs in the program (see DOSAGE AND ADMINISTRATION, Combined Chemotherapy).

For induction therapy, patients should be treated in a facility with laboratory and supportive resources sufficient to monitor drug tolerance and protect and maintain a patient compromised by drug toxicity. The main toxic effect of Cytarabine Injection BP is bone marrow suppression with leukopenia, thrombocytopenia and anemia. Less serious toxicity includes nausea, vomiting, diarrhea and abdominal pain, oral ulceration, and hepatic dysfunction (see ADVERSE REACTIONS).

The physician must judge possible benefit to the patient against known toxic effects of this drug in considering the advisability of therapy with Cytarabine Injection BP. Before making this judgment or beginning treatment, the physician should be familiar with the following text.

When large intravenous doses are given quickly, patients are frequently nauseated and may vomit for several hours post injection. This problem tends to be less severe when the drug is infused.

High dose schedules: Do not further dilute Cytarabine Injection BP with diluents containing benzyl alcohol. Benzyl alcohol has been reported to be associated with a fatal "Gasping Syndrome" in pediatric patients. As premature and low birth weight infants may be at increased risk of developing this toxicity, they should not be given cytarabine when using a diluent containing benzyl alcohol (see WARNINGS AND PRECAUTIONS, Special Populations, Pediatrics).

Carcinogenesis and Mutagenesis

Extensive chromosomal damage, including chromatoid breaks have been produced by cytarabine and malignant transformation of rodent cells in culture has been reported (see DETAILED PHARMACOLOGY).

Cardiovascular

High dose schedules: An increase in cardiomyopathy with subsequent death has been reported following experimental high dose cytarabine and cyclophosphamide therapy when used for bone marrow transplant preparation. This may be schedule dependent (see also DRUG INTERACTIONS).

Gastrointestinal

Abdominal tenderness (peritonitis) and typhlitis with concurrent neutropenia and thrombocytopenia have been reported in patients treated with conventional doses of cytarabine in combination with other drugs. Patients have responded to nonoperative medical management.

High dose schedule: Severe and at times fatal, GI toxicity (different from that seen with conventional therapy regimens of cytarabine) has been reported following high dose (2-3 g / m²) schedules of cytarabine). These reactions include severe gastrointestinal ulceration, including pneumatosis cystoides intestinalis, leading to peritonitis, bowel necrosis; and necrotizing colitis.

Genitourinary

Tumor Lysis Syndrome: Like other cytotoxic drugs, Cytarabine Injection BP may induce hyperuricemia secondary to rapid lysis of neoplastic cells. The clinician should monitor the patient's blood uric acid level and be prepared to use such supportive and pharmacologic measurements as might be necessary to control this problem.

Hematologic

Cytarabine Injection BP is a potent bone marrow suppressant; the severity depends on the dose of the drug and schedule of administration. Therapy should be started cautiously in patients with pre-existing drug-induced bone marrow suppression. Patients receiving this drug must be under close medical supervision and during induction therapy, should have leukocyte and platelet counts performed daily. Bone marrow examinations should be performed frequently after blasts have disappeared from the peripheral blood. Facilities should be available for management of complications (possibly fatal) of bone marrow suppression (infection resulting from granulocytopenia and other impaired body defenses, and hemorrhage secondary to thrombocytopenia). Periodic checks of bone marrow should be performed in patients receiving Cytarabine Injection BP.

Hepatic/Biliary/Pancreatic

The human liver may detoxify a substantial fraction of an administered cytarabine dose. In particular, patients with hepatic function impairment may have a higher likelihood of CNS toxicity after high dose treatment with Cytarabine Injection BP. Use the drug with caution and at reduced dose in patients whose liver function is poor.

Periodic checks of liver function should be performed in patients receiving Cytarabine Injection BP.

Pancreatitis: Acute pancreatitis has been reported to occur in patients being treated with cytarabine in combination with other drugs.

High dose schedules: Other reactions have been reported following high dose (2-3 g / m²) schedules of cytarabine and include sepsis and liver abscess, and liver damage with increased hyperbilirubinemia.

Immune

Immunosuppressant Effects/Increased Susceptibility to Infections: Administration of live or live-attenuated vaccines in patients immunocompromised by chemotherapeutic agents, including cytarabine, may result in serious or fatal infections. Vaccination with a live vaccine should be avoided in patients receiving cytarabine. Killed or inactivated vaccines may be administered; however, the response to such vaccines may be diminished.

Neurologic

High dose schedules: Severe and at times fatal, CNS toxicity (different from that seen with conventional therapy regimens of cytarabine) has been reported following high dose (2-3 g / m²) schedules of cytarabine. These reactions include cerebral and cerebellar dysfunction including personality changes, somnolence, convulsion and coma, usually reversible.

Delayed progressive ascending paralysis resulting in death has been reported in children with AML following intrathecal and intravenous cytarabine at conventional doses in combination with other drugs.

Cases of severe neurological adverse reactions that ranged from headache to paralysis, coma and stroke-like episodes have been reported mostly in pediatric patients given intravenous cytarabine in combination with intrathecal methotrexate.

Ophthalmologic

High dose schedules: The following reactions have been reported following high dose (2-3 g / m²) schedules of cytarabine: reversible corneal toxicity and hemorrhagic conjunctivitis, which may be prevented or diminished by prophylaxis with a local corticosteroid eye drop.

Renal

Patients with renal function impairment may have a higher likelihood of CNS toxicity after high dose treatment with Cytarabine Injection BP. Periodic checks of kidney function should be performed in patients receiving Cytarabine Injection BP.

Respiratory

High dose schedules: Severe and sometimes fatal pulmonary toxicity, adult respiratory distress syndrome and pulmonary edema have occurred following high dose schedules with cytarabine therapy. A syndrome of sudden respiratory distress, rapidly progressing to pulmonary edema and radiographically pronounced cardiomegaly has been reported following experimental high dose cytarabine therapy used for the treatment of relapsed leukemia.

Sensitivity/Resistance

Anaphylactic reactions have occurred with cytarabine treatment. Anaphylaxis that resulted in acute cardiopulmonary arrest and required resuscitation has been reported. This occurred immediately after the intravenous administration of cytarabine.

Sexual Function/Reproduction

Male Fertility: Cytarabine may present in the semen. Male patients who are not surgically sterile must agree to use effective contraception during treatment with Cytarabine Injection BP to prevent pregnancy in female partners (see WARNINGS AND PRECAUTIONS, Special Populations, Pregnant Women and TOXICOLOGY).

Skin

Palmar plantar erythrodysesthesia: Palmar plantar erythrodysesthesia (PPE) has occurred with cytarabine treatment in adults and children. Severe cytarabine associated PPE that resulted in treatment discontinuation has been reported.

High dose schedules: Rarely, severe skin rash, leading to desquamation has been reported. Complete alopecia is more commonly seen with high dose therapy than with standard cytarabine treatment programs.

Special Populations

Pregnant Women: Cytarabine is embryotoxic and teratogenic and produced peri- and postnatal toxicity in various species. Sperm head abnormalities were observed following cytarabine treatment in mice (see TOXICOLOGY).

There are no studies on the use of cytarabine in pregnant women. Use of this drug in women who are or who may become pregnant should be undertaken only after due consideration of potential benefit and potential hazard to both mother and child. Women of childbearing potential should be advised to avoid becoming pregnant (see also WARNINGS AND PRECAUTIONS, Sexual Function/Reproduction).

Normal infants have been born to mothers exposed to cytarabine during pregnancy (alone or in combination with other drugs); some of these infants were premature or of low birthweight. Some of the normal infants were followed up at ages ranging from six weeks to seven years following exposure, and showed no abnormalities. One apparently normal infant died at 80 days of gastroenteritis.

Congenital abnormalities have been reported, particularly when the fetus has been exposed to systemic therapy with cytarabine during the first trimester. These include upper and lower distal limb defects, and extremity and ear deformities.

Reports of pancytopenia, leucopenia, anemia, thrombocytopenia, electrolyte abnormalities, transient eosinophilia, increased IgM levels and hyperpyrexia, sepsis and death have occurred during the neonatal period to infants exposed to cytarabine in utero. Some of these infants were also premature.

Therapeutic abortions have been done in pregnant women on cytarabine. Normal fetuses have been reported while other reported fetal effects included enlarged spleen and Trisomy C chromosome abnormality in the chorionic tissue.

Because of the potential for abnormalities with cytotoxic therapy, particularly during the first trimester, a patient who is or who becomes pregnant while on Cytarabine Injection BP should be apprised of the potential risk to the fetus and the advisability of pregnancy continuation. There is a definite, but considerably reduced risk if therapy is initiated during the second or third trimester. Although normal infants have been delivered to patients treated in all three trimesters of pregnancy, follow-up of such infants would be advisable.

Do not further dilute Cytarabine Injection BP with diluents containing benzyl alcohol. Benzyl alcohol can cross the placenta (see WARNINGS AND PRECAUTIONS, Special Populations, Pediatrics).

Nursing Women: It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk and because of the potential for serious adverse reactions in nursing infants from cytarabine, a decision should be made whether to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother.

Pediatrics: The safety of this drug for use in infants (under 1 year of age) is not established.

Gasping Syndrome: Cytarabine Injection BP should not be given to premature and low birth weight infants when using a diluent that contains benzyl alcohol. The preservative benzyl alcohol has been associated with serious adverse events, including the “gasping syndrome”, and death in pediatric patients. Symptoms of gasping syndrome may include metabolic acidosis, seizure, bradycardia, gasping respiration and cardiovascular collapse. Although normal therapeutic doses of this product ordinarily deliver amounts of benzyl alcohol that are substantially lower than those reported in association with the “gasping syndrome”, the minimum amount of benzyl alcohol at which toxicity may occur is not known. The risk of benzyl alcohol toxicity depends on the quantity administered and the hepatic capacity to detoxify the chemical. Premature and low-birth weight infants may be more likely to develop toxicity. If Cytarabine Injection BP is used in high dose or intrathecal therapy, do not further dilute with benzyl alcohol. The preservative-free 0.9% sodium chloride can be used for dilution (see also SERIOUS WARNINGS AND PRECAUTIONS).

See also WARNINGS AND PRECAUTIONS, Neurologic and Skin.

Monitoring and Laboratory Tests

Patients receiving Cytarabine Injection BP must be monitored closely. Frequent platelet and leukocyte counts and bone marrow examinations are mandatory. Consider suspending or modifying therapy when drug-induced marrow depression has resulted in a platelet count under 50,000 or a polymorphonuclear granulocyte count under 1000 / mm³. Counts of formed elements in the peripheral blood may continue to fall after the drug is stopped and reach lowest values after drug-free intervals of 12 of 24 days. When indicated, restart therapy when definite signs of marrow recovery appear (on successive bone marrow studies). Patients whose drug is withheld until "normal" peripheral blood values are attained, may escape from control.

ADVERSE REACTIONS

Adverse Drug Reaction Overview

The following listing is based on adverse events reported in clinical trials and/or spontaneous adverse event reports from post-marketing experience. When a frequency cannot be estimated from the available data it is classified as “not known”.

Blood and Lymphatic System Disorders

Because Cytarabine Injection BP is a bone marrow suppressant, anemia, leukopenia, thrombocytopenia, megaloblastosis, and reduced reticulocytes can be expected as a result of its administration. The severity of these reactions is dose and schedule dependent. Cellular changes in the morphology of bone marrows and peripheral smears can be expected.

Following 5-day constant infusions or acute injections of 50 mg / m² to 600 mg / m², white cell depression follows a biphasic course. Regardless of initial count, dosage level, or schedule, there is an initial fall starting the first 24 hours with a nadir at days 7 to 9. This is followed by a brief rise which peaks around the twelfth day. A second and deeper fall reaches nadir at days 15 to

24. Then there is a rapid rise to above baseline in the next 10 days. Platelet depression is noticeable at 5 days with a peak depression occurring between days 12 to 15. Thereupon, a rapid rise to above baseline occurs in the next 10 days.

Infections and Infestations

Viral, bacterial, fungal, parasitic, or saprophytic infections, in any location on the body, may be associated with the use of Cytarabine Injection BP alone or in combination with other immunosuppressive agents following immunosuppressive doses that affect cellular or humoral immunity. These infections may be mild, but can be severe and at times fatal.

Musculoskeletal and Connective Tissue Disorders

The Cytarabine Syndrome

A cytarabine syndrome has been described by Castleberry et al. 1981. It is characterized by fever, myalgia, bone pain, occasionally chest pain, maculopapular rash, conjunctivitis and malaise. It usually occurs 6 to 12 hours following drug administration. Corticosteroids have been shown to be beneficial in treating or preventing this syndrome. If the symptoms of the syndrome are deemed treatable, corticosteroids should be contemplated as well as continuation of therapy with Cytarabine Injection BP.

Other Adverse Reactions

Conventional Dose Therapy

Nausea and vomiting are most frequent following rapid intravenous injection.

Table 1 – Frequencies of Adverse Reactions with Cytarabine Conventional Dose Therapy

The reported adverse reactions are listed below by MedDRA System Organ Class and by frequency.

ADR frequencies are based on CIOMS convention: Very common (>10%), Common (>1%, ≤10%), Uncommon (>0.1%, ≤1%), Rare (>0.01%, ≤0.1%), and Frequency not known (cannot be estimated from available data).

Blood and Lymphatic System Disorders:	
Very common	Bone marrow failure, thrombocytopenia, anaemia, anaemia megaloblastic, leukopenia, reticulocyte count decreased
Frequency not known	Bleeding (all sites)
Cardiac Disorders:	
Frequency not known	Pericarditis
Eye Disorders:	
Frequency not known	Conjunctivitis ^a
Gastrointestinal Disorders:	
Very common	Stomatitis, mouth ulceration, anal ulcer, anal inflammation, diarrhoea, vomiting, nausea, abdominal pain
Frequency not known	Bowel necrosis, Pancreatitis, oesophageal ulcer, oesophagitis
General Disorders and Administration Site Conditions:	
Very common	Pyrexia
Frequency not known	Chest pain, injection site reaction ^b

Hepatobiliary Disorders:	
Very common	Hepatic function abnormal
Frequency not known	Jaundice
Immune System Disorders:	
Frequency not known	Anaphylactic reaction, allergic oedema
Infections and Infestations:	
Very common	Sepsis, pneumonia, infection ^c
Frequency not known	Injection site cellulitis
Investigations:	
Very common	Biopsy bone marrow abnormal, blood smear test abnormal
Metabolism and Nutrition Disorders:	
Frequency not known	Decreased appetite
Musculoskeletal, Connective Tissue and Bone Disorders:	
Very common	Cytarabine syndrome
Nervous System Disorders:	
Frequency not known	Neurotoxicity, neuritis, dizziness, headache
Renal and Urinary Disorders:	
Frequency not known	Renal impairment, urinary retention
Respiratory, Thoracic and Mediastinal Disorders:	
Frequency not known	Dyspnea, oropharyngeal pain
Skin and Subcutaneous Tissue Disorders:	
Very common	Alopecia, rash
Common	Skin ulcer
Frequency not known	Palmar-plantar erythrodysesthesia syndrome, urticaria, pruritus, freckling
Vascular Disorders:	
Frequency not known	Thrombophlebitis

^a May occur with rash and may be hemorrhagic with high dose therapy.

^b Pain and inflammation at subcutaneous injection site.

^c May be mild, but can be severe and at times fatal.

High Dose Therapy

Severe and at times fatal CNS, GI and pulmonary toxicity (different from that seen with conventional therapy regimens of cytarabine) has been reported following high dose schedules (2.0 g to 3.0 g / m² given every 12 hours for 12 doses).

Table 2 – Frequencies of Adverse Reactions with Cytarabine High Dose Therapy

The reported adverse reactions are listed below by MedDRA System Organ Class and by frequency.

ADR frequencies are based on CIOMS convention: Very common (>10%), Common (>1%, ≤10%), Uncommon (>0.1%, ≤1%), Rare (>0.01%, ≤0.1%), and Frequency not known (cannot be estimated from available data).

Cardiac Disorders:	
Frequency not known	Cardiomyopathy ^a
Eye Disorders:	
Very common	Corneal disorder
Frequency not known	hemorrhagic conjunctivitis ^b
Gastrointestinal Disorders:	
Common	Necrotising colitis
Frequency not known	Gastrointestinal necrosis, gastrointestinal ulcer, pneumatosis intestinalis, peritonitis
Hepatobiliary Disorders:	
Frequency not known	Liver injury, hyperbilirubinaemia
Infections and Infestations:	
Very common	Sepsis
Frequency not known	Liver abscess
Nervous System Disorders:	
Very common	Cerebral disorder, cerebellar disorder, somnolence
Frequency not known	Coma, convulsion, peripheral motor neuropathy, peripheral sensory neuropathy
Psychiatric Disorders:	
Frequency not known	Personality change ^c
Respiratory, Thoracic and Mediastinal Disorders:	
Very common	Acute respiratory distress syndrome, pulmonary oedema
Skin and Subcutaneous Tissue Disorders:	
Common	Skin exfoliation

^a With subsequent death.

^b May be prevented or diminished by prophylaxis with a local corticosteroid eyedrop.

^c Personality change was reported in association with cerebral and cerebellar dysfunction.

Peripheral motor and sensory neuropathies after consolidation with high-dose cytarabine, daunorubicin, and asparaginase have occurred in adult patients with acute non lymphocytic leukemia. Patients treated with high-dose Cytarabine Injection BP should be observed for neuropathy since dose schedule alterations may be needed to avoid irreversible neurologic disorders.

Corneal toxicity consisting of ocular pain, tearing, foreign-body sensation, photophobia and blurred vision has been reported.

Rarely, severe skin rash, leading to desquamation has been reported. Complete alopecia is more commonly seen with high dose therapy than with standard cytarabine treatment programs.

Do not further dilute Cytarabine Injection BP with diluents containing benzyl alcohol.

Intermediate Dose Therapy

A diffuse interstitial pneumonitis without clear cause that may have been related to cytarabine was reported in patients treated with experimental intermediate doses of cytarabine (1 gm / m²)

with and without other chemotherapeutic agents (meta-AMSA, daunorubicin, VP-16).

Intrathecal Therapy

Cytarabine Injection BP given intrathecally may cause systemic toxicity and careful monitoring of the hemopoietic system is indicated. Modification of other anti-leukemia therapy may be necessary. Major toxicity is rare. The most frequently reported reactions after intrathecal administration were nausea, vomiting and fever; these reactions are mild and self-limiting. Paraplegia has been reported. Necrotizing leukoencephalopathy with or without convulsion has been reported; in some cases, patients had also been treated with intrathecal methotrexate and/or hydrocortisone, as well as by central nervous system radiation. Isolated neurotoxicity has been reported. Blindness occurred in two patients in remission whose treatment had consisted of combination systemic chemotherapy, prophylactic central nervous system radiation and intrathecal cytarabine. When Cytarabine Injection BP is administered both intrathecally and intravenously within a few days, there is an increased risk of spinal cord toxicity, however, in serious life-threatening disease, concurrent use of intravenous and intrathecal Cytarabine Injection BP is left to the discretion of the treating physician.

DRUG INTERACTIONS

Serious Drug Interactions

Methotrexate: Intravenous cytarabine given concomitantly with intrathecal methotrexate may increase the risk of severe neurological adverse reactions such as headache, paralysis, coma and stroke-like episodes.

Drug-Drug Interactions

Digoxin: Reversible decreases in steady-state plasma digoxin concentrations and renal glycoside excretion were observed in patients receiving beta-acetyldigoxin and chemotherapy regimens containing cyclophosphamide, vincristine and prednisone with or without cytarabine or procarbazine. Steady-state plasma digitoxin concentrations did not appear to change. Therefore, monitoring of plasma digoxin levels may be indicated in patients receiving similar combination chemotherapy regimens. The utilization of digitoxin for such patients may be considered as an alternative.

Gentamicin: An *in vitro* interaction study between gentamicin and cytarabine showed a cytarabine related antagonism for the susceptibility of *K. pneumoniae* strains. This study suggests that in patients on cytarabine being treated with gentamicin for a *K. pneumoniae* infection, the lack of a prompt therapeutic response may indicate the need for re-evaluation of antibacterial therapy.

Fluorocytosine: Clinical evidence showed possible inhibition of fluorocytosine efficacy therapy with cytarabine. This may be due to potential competitive inhibition of its uptake.

Experimental high dose cytarabine and cyclophosphamide therapy: An increase in cardiomyopathy with subsequent death has been reported when used for bone marrow transplant

preparation. This may be schedule dependent (see WARNINGS AND PRECAUTIONS, Cardiovascular).

Drug-Food Interaction

Interactions with food have not been established.

Drug-Herb Interactions

Interactions with herbal product have not been established.

Drug-Laboratory Test Interactions

Interactions with laboratory tests have not been established.

Drug-Lifestyle Interactions

Interactions associated with lifestyle have not been established.

DOSAGE AND ADMINISTRATION

Dosing Considerations

Clinical experience accumulated to date suggests that success with cytarabine is dependent more on adeptness in modifying day-to-day dosage to obtain maximum leukemic cell kill with tolerable toxicity than on the basic treatment schedule chosen at the outset of therapy. Toxicity necessitating dosage alteration almost always occurs.

In many chemotherapeutic programs, Cytarabine Injection BP is used in combination with other cytotoxic drugs. The addition of these cytotoxic drugs has necessitated changes and dose alterations. The dosage schedules for combination therapy outlined below have been reported in the literature (see REFERENCES).

Recommended Dose and Dosage Adjustment

Acute myelocytic leukemia - induction remission: adults

Cytarabine Injection BP 200 mg / m² daily by continuous infusion for 5 days (120 hours) - total dose 1000 mg / m². This course is repeated approximately every 2 weeks. Modifications must be made based on hematologic response.

Acute myelocytic leukemia - maintenance: adults

Maintenance programs are modifications of induction programs and, in general, use similar schedules of drug therapy as were used during induction. Most programs have a greater time spacing between courses of therapy during remission maintenance.

Acute myelocytic leukemia - induction and maintenance in children

Numerous studies have shown that childhood AML responds better than adult AML given similar regimens. Where the adult dosage is stated in terms of body weight or surface area, the children's dosage may be calculated on the same basis. When specified amounts of a drug are indicated for the adult dosage, these should be adjusted for children on the basis of such factors as age, body weight or body surface area.

Acute myelocytic leukemia – adults and children

The following tables outline the results of treatment with cytarabine alone and in combination with other chemotherapeutic agents, in the treatment of acute myelocytic leukemia in adults and children.

The treatment regimens outlined in the tables should not be compared for efficacy. These were independent studies with a number of variables involved, such as patient population, duration of disease, and previous treatment.

The responsiveness and course of childhood acute myelocytic leukemia (AML) appear to be different from that in adults. Numerous studies show response rates to be higher in children than in adults with similar treatment schedules. Experience indicates that at least with induction and initial drug responsiveness, childhood AML appears to be more similar to childhood acute lymphocytic leukemia (ALL) than to its adult variant.

Patients with hepatic impairment: Cytarabine and dose adjustment has not been studied in individuals with hepatic impairment (see also WARNINGS AND PRECAUTIONS, Hepatic/Biliary/Pancreatic).

Patients with renal impairment: Cytarabine and dose adjustment has not been studied in individuals with renal impairment (see also WARNINGS AND PRECAUTIONS, Renal).

Table I: Acute Myelocytic Leukemia - Remission Induction: Adults

Drug Dosage Schedule*		No. of Patients Evaluated	Complete Remissions	Investigators
Cytarabine Single-Dose Therapy	(Infusion) 10 mg / m ² 12 hrs/day	12	2 (17%)	Ellison (1968)
	30 mg / m ² 12 hrs/day	41	10 (24%)	
	10 mg / m ² 24 hrs/day	9	2 (22%)	
	30 mg / m ² 24 hrs/day	36	2 (6%)	
	(Infusion) 200 mg / m ² 24 hrs/5 days	36	9 (25%)	Bodey (1969)
	10 mg / m ² IV injection initially, then infusions of 30 mg/m ² per 12 hrs or 60 mg/m ² /day for 4 days	49	21 (43%)	Goodell (1970)
	(Infusion Therapy) 800 mg / m ² / 2 days	53	12 (23%)	Southwest Oncology Group.(1974)
1000 mg / m ² / 5 days	60	24 (40%)		
100 mg / m ² /day 1 hr infusion	49	7 (14%)	Carey (1975)	

Drug Dosage Schedule*		No. of Patients Evaluated	Complete Remissions	Investigators
	5-12.5 mg /kg / 12 hr infusion following IV synchronizing dose**	5	5 (100%)	Lampkin (1976)
Combined Therapy	Cytarabine – doxorubicin	41	30 (73%)	Preisler (1979)
	Cytarabine - thioguanine daunorubicin	28	22 (79%)	Gale (1977)
	Cytarabine - doxorubicin vincristine – prednisolone	35	23 (66%)	Weinstein (1980)
	Cytarabine - daunorubicin thioguanine – prednisone vincristine	139	84 (60%)	Glucksberg (1981)
	Cytarabine – daunorubicin	21	14 (67%)	Cassileth (1977)
High Dose Therapy	Cytarabine	7	6 (86%)	Lister (1983)
	Cytarabine	21	12 (57%)	Herzig (1983)
	Cytarabine	11	8 (73%)	Preisler (1983)
	Cytarabine - doxorubicin	14	7 (50%)	Willemze (1982)
	Cytarabine - asparaginase	13	9 (69%)	Capizzi (1983)

* Unless otherwise stated, all doses given until drug effect-modifications then based on hematologic reasons. See references.

** Highly experimental - requires ability to study mitotic indices.

Table II: Acute Myelocytic Leukemia - Remission Induction: Children (21 and under)

Drug Therapy	No. of Patients Evaluated	Complete Remissions	Investigator
Cytarabine (5-12.5 mg / kg following IV synchronizing dose**)	16	12 (75%)	Lampkin (1976)
Cytarabine, vincristine, doxorubicin, prednisolone	48	35 (73%)	Weinstein (1980)
Cytarabine, thioguanine, doxorubicin	11	8 (72%)	Hagbin (1975)
Cytarabine, thioguanine	47	20 (43%)	Pizzo (1976)
Cytarabine, cyclophosphamide	12	7 (58%)	Pizzo (1976)

** Highly experimental - requires ability to study mitotic indices.

Acute lymphocytic leukemia

In general, dosage schedules are similar to those used in acute myelocytic leukemia with some modification. Cytarabine has been used in the treatment of acute lymphocytic leukemia in both adults and children. When cytarabine was used with other antineoplastic agents as part of a total therapy program, results were equal to or better than reported with such programs which did not include cytarabine. Used singly, or in combination with other agents, cytarabine has also been effective in treating patients who had relapsed on other therapy. Table III and IV summarize the results obtained in previously treated patients. Since these are independent studies with such variables as patient population, duration of disease and previous treatment, results shown should not be used for comparing the efficacy of the outlined treatment programs.

Table III: Acute Lymphocytic Leukemia - Remission Induction Previously Treatment Patients Adults and Children

Drug Therapy	No. of Patients Evaluated	Complete Remissions	Response	Investigator
Cytarabine 3-5 mg / kg / day (IV injection)	43	2 (5%)	15 (35%)	Howard (1968)
Cytarabine - asparaginase	9	8 (89%)	8 (89%)	McElwain (1969)
Cytarabine - cyclophosphamide	11	7 (64%)	9 (82%)	Bodey 1970
Cytarabine - prednisone	83	-	(49%)	Nesbitt (1970)
Cytarabine 150-200 mg / m ² / 5 days (infusion)	34	1 (3%)	4 (12%)	Wang (1970)
Cytarabine - L - asparaginase - prednisone - vincristine - doxorubicin	91	72 (79%)	-	Klemperer (1978)
Cytarabine - L - asparaginase - prednisone - vincristine - doxorubicin	55	42 (76%)	-	Klemperer (1978)
Cytarabine - asparaginase	22	13 (59%)	15 (68%)	Ortega (1972)
Cytarabine - thioguanine	19	9 (47%)	9 (47%)	Bryan (1974)

Table IV

Drug Therapy		No. of Patients Evaluated	Complete Remissions	Investigator
High Dose Therapy	Cytarabine	8	3 (38%)	Rohatinar (1983)
	Cytarabine - doxorubicin	3	2 (67%)	Willemze (1982)
	Cytarabine - asparaginase	10	3 (30%)	Capizzi (1983)

Non-Hodgkin's lymphoma in children

Cytarabine has been used as part of multi-drug program (LSA₂L₂) to treat non-Hodgkin's lymphoma in children. See Appendix A for complete dosage schedule.

High Dose Chemotherapy

Before instituting a program of high dose chemotherapy, the physician should be familiar with the literature, adverse reactions, precautions, contraindications, and warnings applicable to all the drugs involved in the program.

Cytarabine Injection BP

Cytarabine Injection BP: 2 g / m² infused over 3 hours every 12 hours x 12 doses (Days 1-6).

Cytarabine Injection BP

Cytarabine Injection BP: 3 g / m² infused over 1 hour every 12 hours x 12 doses (Days 1-6).

Cytarabine Injection BP

Cytarabine Injection BP: 3 g / m² infused over 75 minutes every 12 hours x 12 doses (Days 1-6).

Cytarabine Injection BP - doxorubicin

Cytarabine Injection BP: 3 g / m² infused over 2 hours every 12 hours x 12 doses (Days 1-6).

Doxorubicin: 30 mg / m² IV on Days 6-7.

Cytarabine Injection BP - asparaginase

Cytarabine Injection BP: 3 g / m² infused over 3 hours at 0 hours, 12 hours, 24 hours, and 36 hours. At 42 hours, 6000 units / m² of asparaginase IM (Days 1-2); repeat same schedule Days 8-9.

Combined Chemotherapy

Before instituting a program of combined chemotherapy, the physician should be familiar with the literature, adverse reactions, precautions, contraindications, and warnings applicable to all the drugs involved in the program.

Cytarabine Injection BP, doxorubicin

Cytarabine Injection BP: 100 mg / m² / day, continuous IV infusion (Days 1-10)
Doxorubicin: 30 mg / m² / day, IV infusion of 30 minutes (Days 1-3)

Additional (complete or modified) courses as necessary at 2-4 week intervals if leukemia is persistent.

Cytarabine Injection BP, thioguanine, daunorubicin

Cytarabine Injection BP: 100 mg / m², IV infusion over 30 minutes every 12 hours (Days 1-7).
Thioguanine: 100 mg / m², orally every 12 hours (Days 1-7).
Daunorubicin: 60 mg / m²/day, IV infusion (Days 5-7).

Additional (complete or modified) courses as necessary at 2-4 week intervals if leukemia is persistent.

Cytarabine Injection BP, doxorubicin, vincristine, prednisone

Cytarabine Injection BP: 100 mg / m² / day, continuous IV infusion (Days 1-7).
Doxorubicin: 30 mg / m² / day, IV infusion (Days 1-3).
Vincristine: 1.5 mg / m²/day, IV infusion (Days 1, 5)
Prednisolone: 40 mg / m² / day, IV infusion every 12 hours (Days 1-5).

Additional (complete or modified) courses as necessary at 2-4 week intervals if leukemia is persistent.

Cytarabine Injection BP, daunorubicin, thioguanine, prednisone, vincristine

Cytarabine Injection BP: 100 mg / m² / day, IV infusion (Days 1-10).
Daunorubicin: 70 mg / m² / day, IV infusion (Days 1-3).
Thioguanine: 100 mg / m², orally every 12 hours (Days 1-7).
Prednisone: 40 mg / m² / day, orally (Days 1-7).
Vincristine: 1 mg / m² / day, IV infusion (Days 1, 7)

Additional (complete or modified) courses as necessary at 2-4 week intervals if leukemia is persistent.

Cytarabine Injection BP, daunorubicin

Cytarabine Injection BP: 100 mg / m² / day, continuous IV infusion (Days 1-7).

Daunorubicin: 45 mg / m² / day, IV. push (Days 1-3).

Additional (complete or modified) courses as necessary at 2-4 week intervals if leukemia is persistent.

Meningeal Leukemia - Intrathecal Use

Cytarabine has been used intrathecally in acute leukemia in doses ranging from 5 mg / m² to 75 mg / m² of body surface area. The frequency of administration varied from once a day for 4 days to once every 4 days. The most frequently used dose was 30 mg / m² every 4 days until cerebrospinal fluid findings were normal, followed by one additional treatment. The dosage schedule is usually governed by the type and severity of central nervous system manifestations and the response to previous therapy.

Cytarabine has been used intrathecally with hydrocortisone sodium succinate and methotrexate, both as prophylaxis in newly diagnosed children with acute lymphocytic leukemia, as well as in the treatment of meningeal leukemia. Sullivan et al. has reported that prophylactic triple therapy has prevented late CNS disease and given overall cure and survival rates similar to those seen in patients in whom CNS radiation and intrathecal methotrexate was used as initial CNS prophylaxis. The dose of cytarabine was 30 mg / m², hydrocortisone sodium succinate 15 mg / m², and methotrexate 15 mg / m² (an absolute maximum single dose of 15 mg of methotrexate). The physician should be aware of this regimen and note that methotrexate dosage in pediatric patients is otherwise based on age rather than body surface area. Prescribers should consult related Product Monographs for more information.

Prophylactic triple therapy following the successful treatment of the acute meningeal episode may be useful. The physician should familiarize himself with the current literature before instituting such a program.

Cytarabine Injection BP given intrathecally may cause systemic toxicity and careful monitoring of the hemopoietic system is indicated. Modification of the anti-leukemia therapy may be necessary. Major toxicity is rare. The most frequently reported reactions after intrathecal administration were nausea, vomiting and fever; these reactions are mild and self-limiting. Paraplegia has been reported. Necrotizing leukoencephalopathy occurred in 5 children; these patients had also been treated with intrathecal methotrexate and hydrocortisone, as well as by central nervous system radiation. Isolated neurotoxicity has been reported.

Blindness occurred in two patients in remission whose treatment had consisted of combination systemic chemotherapy, prophylactic central nervous system radiation and intrathecal cytarabine.

Focal leukemic involvement of the central nervous system may not respond to intrathecal Cytarabine Injection BP and may better be treated with radiotherapy.

If used intrathecally, dilute with preservative free saline and use immediately. Do not further dilute Cytarabine Injection BP with diluents containing benzyl alcohol.

Dosage Modification

The dosage of Cytarabine Injection BP must be modified or suspended when signs of serious hematologic depression appear. In general, consider discontinuing the drug if the patient has less than 50000 platelets or 1000 polymorphonuclear granulocytes/mm³ in his peripheral blood. These guidelines may be modified depending on signs of toxicity in other systems and on the rapidity of fall in formed blood elements. Restart the drug when there are signs of marrow recovery and the above platelet and granulocyte levels have been attained. Withholding therapy until the patient's blood values are normal may result in escape of the patient's disease from control by the drug.

Hepatic Insufficiency: Use cytarabine with caution or possibly at reduced doses in patients whose liver function is poor (see also WARNINGS AND PRECAUTIONS, Hepatic/Biliary/Pancreatic).

Renal Insufficiency: Use cytarabine with caution or possibly at reduced doses in patients whose kidney function is poor (see also WARNINGS AND PRECAUTIONS, Renal).

Administration

Cytarabine is not active orally. The schedule and method of administration varies with the program of therapy to be used. Cytarabine Injection BP may be given by intravenous infusion, injection/subcutaneously or intrathecally. When preparing cytarabine for intravenous high dose therapy or intrathecal use, do not further dilute with benzyl alcohol (see SERIOUS WARNINGS AND PRECAUTIONS). It is recommended that Cytarabine Injection BP be diluted with preservative-free 0.9% sodium chloride for injection and used immediately.

Thrombophlebitis has occurred at the site of drug injection or infusion in some patients, and rarely patients have noted pain and inflammation at subcutaneous injection sites. In most instances, however, the drug has been well tolerated.

Patients can tolerate higher total doses when they receive the drug by rapid intravenous injection as compared with slow infusion. This phenomenon is related to the drug's rapid inactivation and brief exposure of susceptible normal and neoplastic cells to significant levels after rapid injection. Normal and neoplastic cells seem to respond to somewhat parallel fashion to these different modes of administration and no clear-cut clinical advantage has been demonstrated for either.

Relatively constant plasma levels can be achieved by continuous intravenous infusion.

FOR INTRATHECAL USE: DO NOT USE DILUENT CONTAINING BENZYL ALCOHOL. DILUTE WITH PRESERVATIVE-FREE 0.9% SODIUM CHLORIDE FOR INJECTION. USE IMMEDIATELY.

FOR HIGH DOSE USE: DO NOT FURTHER DILUTE CYTARABINE INJECTION BP WITH DILUENTS CONTAINING BENZYL ALCOHOL.

OVERDOSAGE

There is no antidote for Cytarabine Injection BP overdose.

Discontinuation of the drug and supportive therapy are of course indicated. Transfusions of platelets should be given if there is any sign of hemorrhage. Patients should be carefully observed for intercurrent infection and if such appears they should be rapidly and rigorously treated with appropriate antibiotic therapy.

Chronic overdose may cause serious bone marrow suppression. Daily hematological evaluation should be performed to prevent overdose. Nausea and vomiting, although a general side effect of the drug, may be an additional warning of overdose. Severe hemorrhage into the gastrointestinal tract may indicate overdose as may severe generalized infections.

Doses exceeding recommended dosage schedules have been used clinically and have been tolerated. The major toxicity with the use of $3 \text{ g} / \text{m}^2$ intravenous infusion over 1 hour every 12 hours for 12 doses and $3 \text{ g} / \text{m}^2$ continuous infusion for 4 days, other than reversible bone marrow suppression has been reversible corneal, cerebral and cerebellar dysfunction. Doses of $4.5 \text{ g} / \text{m}^2$ intravenous infusion over 1 hour every 12 hours for 12 doses has caused an unacceptable increase in irreversible CNS toxicity and death.

For management of a suspected drug overdose, contact your regional Poison Control Centre immediately.

ACTION AND CLINICAL PHARMACOLOGY

Pharmacodynamics

Cytarabine is capable of obliterating immune responses in man during administration. Suppression of antibody responses to *E-coli*-VI antigen and tetanus toxoid has been demonstrated. This suppression was obtained during both primary and secondary antibody responses.

Cytarabine also suppressed the development of cell-mediated immune responses such as delayed hypersensitivity skin reaction to dinitrochlorobenzene. However, it has no effect on already established delayed hypersensitivity reactions.

Following 5-day courses of intensive therapy with cytarabine the immune response was suppressed, as indicated by the following parameters: macrophage ingress into skin windows; circulating antibody response following primary antigenic stimulation; lymphocyte blastogenesis with phytohemagglutinin. A few days after termination of therapy there was a rapid return to normal.

Pharmacokinetics

Absorption:

Cytarabine is rapidly metabolized and is not effective orally; less than 20% of the orally administered dose is absorbed from the gastrointestinal tract.

After subcutaneous or intramuscular administration of cytarabine, peak plasma levels of radioactivity are achieved about 20 to 60 minutes after injection and are considerably lower than those after intravenous administration.

Distribution:

Cerebrospinal fluid levels of cytarabine are low in comparison to plasma levels after single intravenous injection. However, in one patient in whom cerebrospinal levels were examined after 2 hours of constant intravenous infusion, levels approached 40% of the steady state plasma level. With intrathecal administration, levels of cytarabine in the cerebrospinal fluid declined with a first order half-life of about 2 hours. Because cerebrospinal fluid levels of deaminase are low, little conversion to ara-U was observed.

Metabolism:

Cytarabine is metabolized by deoxycytidine kinase and other nucleotide kinases to the nucleotide triphosphate, an effective inhibitor of DNA polymerase; it is inactivated by pyrimidine nucleoside deaminase which converts it to the non-toxic uracil derivative. It appears that the balance of kinase and deaminase levels may be an important factor in determining sensitivity or resistance of the cell to cytarabine.

Excretion:

Following rapid intravenous injection of cytarabine, the disappearance from plasma is biphasic. There is an initial distributive phase with a half-life of about 10 minutes, followed by a second elimination phase with a half-life of about 1 to 3 hours. After the distributive phase, over 80% of plasma radioactivity can be accounted for by the inactive metabolite 1-β-D-arabinofuranosyluracil (ara-U). Within 24 hours about 80% of the administered radioactivity can be recovered in the urine, approximately 90% of which is excreted as ara-U.

Special Populations and Conditions

Hepatic Insufficiency: Use cytarabine with caution or possibly at reduced doses in patients whose liver function is poor (see WARNINGS AND PRECAUTIONS, Hepatic/Biliary/Pancreatic and DOSAGE AND ADMINISTRATION).

Renal Insufficiency: Use cytarabine with caution or possibly at reduced doses in patients whose kidney function is poor (see WARNINGS AND PRECAUTIONS, Renal and DOSAGE AND ADMINISTRATION).

STORAGE AND STABILITY

Stability and Storage Recommendations

Store between 15°C and 25°C. Protect from light.

Chemical Stability and Compatibility

In-use Stability

Cytarabine Injection BP solution is stable after the vial has been punctured for 24 hours when stored at ambient temperature (15°C – 25°C) and for 48 hours when stored at refrigerated conditions (2°C - 8°C).

Stability in Infusion Solutions

Chemical and physical stability studies of Cytarabine Injection BP have demonstrated that cytarabine is stable for seven days at room temperature or at 2°C - 8°C when admixed at 0.1 mg / mL in plastic IV bags with: 5% Dextrose injection; 0.9% Sodium Chloride injection and Lactated Ringer's Injection solutions.

Also when similarly admixed at 32.5 mg / mL in plastic IV bags, cytarabine is stable for seven days at room temperature or at 2°C - 8°C in 0.9% Sodium Chloride Injection Solution.

Drug Incompatibilities

Cytarabine has been known to be physically incompatible with heparin, insulin, 5-fluorouracil, penicillin G, and methylprednisolone sodium succinate.

AS WITH ALL INTRAVENOUS ADMIXTURES, DILUTION SHOULD BE MADE JUST PRIOR TO ADMINISTRATION AND THE RESULTING UNPRESERVED SOLUTION USED WITHIN 24 HOURS.

As with all parenteral drug products, intravenous admixtures should be inspected visually for clarity, particulate matter, precipitate, discoloration and leakage prior to administration whenever solution and container permit. Solutions showing haziness, particulate matter, precipitate, discoloration or leakage should not be used.

SPECIAL HANDLING INSTRUCTIONS

CAUTION

The following precautionary measures are recommended in proceeding with the preparation and handling of cytotoxic agents such as Cytarabine Injection BP.

1. The procedure should be carried out in a vertical laminar flow hood (Biological Safety Cabinet - Class II).
2. Personnel should wear: PVC gloves, safety glasses, disposable gowns and masks.
3. All needles, syringes, vials, and other materials which have come in contact with Cytarabine Injection BP should be segregated and destroyed by incineration (sealed

containers may explode). If incineration is not available, neutralization should be carried out using 5% sodium hypochlorite, or 5% sodium thiosulfate.

4. Personnel regularly involved in the preparation and handling of Cytarabine Injection BP should have bi-annual haematologic examinations.

DOSAGE FORMS, COMPOSITION AND PACKAGING

Each mL of sterile unpreserved solution of Cytarabine Injection BP contains 100 mg of cytarabine in water for injection. Hydrochloric acid solution and/or sodium hydroxide solution is added to adjust the pH. pH of the solution is approximately 8.

Cytarabine Injection BP is supplied in single use vials of 100 mg / 1 mL (5 vials per box), 500 mg / 5 mL (5 vials per box), 1000 mg / 10 mL (1 vial per box) and 2000 mg / 20 mL (1 vial per box).

PART II: SCIENTIFIC INFORMATION

PHARMACEUTICAL INFORMATION


Drug Substance

Proper name: cytarabine

Chemical name: 4-amino-1- β - D-arabinofuranosyl-1*H*-pyrimidin-2-one

Molecular formula and molecular mass: $C_9H_{13}N_3O_5$ 243.2 g / mol

Structural formula:



Description:

- white to almost white, crystalline powder
- melting point 215°C

Solubility:

- freely soluble in water
- slightly soluble in alcohol and methylene chloride

DETAILED PHARMACOLOGY

Cell Culture Studies

Cytarabine is cytotoxic to a wide variety of proliferating mammalian cells in culture. It exhibits cell phase specificity, primarily killing cells undergoing DNA synthesis (S-phase) and under certain conditions blocking the progression of cells from the G₁ phase to S-phase. Although the mechanism of action is not completely understood, it appears that cytarabine acts through the inhibition of DNA polymerase. A limited, but significant, incorporation of cytarabine into both DNA and RNA has also been reported. Extensive chromosomal damage, including chromatoid breaks has been produced by cytarabine and malignant transformation of rodent cells in culture has been reported. Deoxycytidine prevents or delays (but does not reverse) the cytotoxic activity.

Animal Studies

In experimental studies with mouse tumors, cytarabine was most effective in those tumors with a high growth fraction. The effect was dependent on the treatment schedule; optimal effects were achieved when the schedule (multiple closely spaced doses or constant infusion) ensured contact of the drug with the tumor cells when the maximum number of cells was in the susceptible S-phase. The best results were obtained when courses of therapy were separated by intervals sufficient to permit adequate host recovery.

TOXICOLOGY

Animal Studies

Toxicity of cytarabine in experimental animals, as well as activity, is markedly influenced by the schedule of administration. For example, in mice, the LD₁₀ for single intraperitoneal administration is greater than 6000 mg / m². However, when administered in 8 doses, each separated by 3 hours, the LD₁₀ is less than 750 mg / m² total dose. Similarly, although a total dose of 1920 mg / m² administered as 12 injections at 6-hour intervals was lethal to beagle dogs (severe bone marrow hypoplasia with evidence of liver and kidney damage), dogs receiving the same total dose administered as 8 injections (again at 6-hour intervals) over a 48-hour period survived with minimal signs of toxicity.

The most consistent observation in surviving dogs was elevated transaminase levels. In all experimental species the primary limiting toxic effect is marrow suppression with leukopenia. In addition, cytarabine causes abnormal cerebellar development in the neonatal hamster and is teratogenic to the rat fetus.

The major dose-limiting toxicity of cytarabine observed in all tested species is myelosuppression, manifested by megaloblastosis, reticulocytopenia, leukopenia, and thrombocytopenia. Other target organs include liver, kidney, and brain. Extensive chromosomal damage, including chromatoid breaks have been produced by cytarabine and malignant transformation of rodent cells in culture has been reported. Cytarabine is embryotoxic and teratogenic and produced peri- and postnatal toxicity in various species. No formal fertility

studies have been reported however sperm head abnormalities were observed following cytarabine treatment in mice.

REFERENCES

1. Zaky DA, Betts RF, Douglas RG, et al. Varicella-Zoster virus and subcutaneous cytarabine: Correlation of *in vitro* sensitivities to blood levels. *Antimicrob Agents Chemother* 1975;7:229-32.
2. Davis CM, VanDersarl JV, Coltman CA Jr. Failure of cytarabine in varicella-zoster infections. *JAMA* 1973;224:122-3.
3. Betts RF, Zaky DA, Douglas RG, et al. Ineffectiveness of subcutaneous cytosine arabinoside in localized herpes zoster. *Ann Intern Med* 1975;82:778-83.
4. Dennis DT, Doberstyn EB, Awoke S, et al. Failure of cytosine arabinoside in treating smallpox, a double-blind study. *Lancet* 1974;2:377-9.
5. Gray GD. ARA-C and derivatives as examples of immunosuppressive nucleoside analogs. *Ann NY Acad Sci* 1975;255:372-9.
6. Mitchell MS, Wade ME, DaConti RC, et al. Immunosuppressive effects of cytosine arabinoside and methotrexate in man. *Ann Intern Med* 1969;70:525-47.
7. Frei E, Ho DHW, Body GP, et al. Pharmacologic and cytotoxic studies of arabinosyl cytosine. In unifying concepts of leukemia. *Bibl Hematol* No. 39 Karger, Base 1, 1973, p 1085-7.
8. Woolner N, Burchenal JH, Lieberman PH, et al. Non-Hodgkin's lymphoma in children - A comparative study of two modalities of therapy. *Cancer* 1976;37:123-34.
9. Woolner N, Exelby PR, Lieberman PH. Non-Hodgkin's lymphoma in children - A progress report on the original patients treated with the LSA₂-L₂ protocol. *Cancer* 1979;44:1990-99.
10. Sullivan MP, Pullen J, Moore T, et al: Pediatric oncology group trial of LSA₂-L₂ therapy in Non-Hodgkin's lymphoma. Abstracted, *Proc AACR and ADCO* 1981;22:C-180.
11. Ellison RR, Holland JF, Weil M, et al. Arabinosyl cytosine: A useful agent in the treatment of acute leukemia in adults. *Blood* 1968;32:507-23.
12. Bodey GP, Freireich EJ, Monto RW, et al. Cytosine arabinoside (NSC-63878) therapy for acute leukemia in adults. *Cancer Chemother Rep* 1969;53:59-66.
13. Goodell B, Leventhall B, Henderson E. Cytosine arabinoside in acute granulocytic leukemia. *Clin Pharmacol Ther* 1970;12:599-606.
14. Southwest Oncology Group. Cytarabine for acute leukemia in adults. *Arch Intern Med* 1974;133:251-9.

15. Carey RW, Ribas-Mundo M, Ellison RR, et al. Comparative study of cytosine arabinoside therapy alone and combined with thioguanine, mercaptopurine or daunorubicin in acute myelocytic leukemia. *Cancer* 1975;36:1560-66.
16. Lampkin BC, McWilliams NB, Mauer AM, et al. Manipulation of the mitotic cycle in the treatment of acute myelogenous leukemia. *Br J Haematol* 1976;32:29-40.
17. Preisler H, Bjornsson S, Henderson ES, et al. Remission induction in acute non-lymphocytic leukemia - Comparison of a seven-day and ten-day infusion of cytosine arabinoside in combination with adriamycin. *Med Pediatr Oncol* 1979;7:269-75.
18. Gale RP, Cline MJ. High remission - induction rate in acute myeloid leukemia. *Lancet* 1977;1:497-9.
19. Weinstein JH, Mayer RJ, Rosenthal DS, et al. Treatment of acute myelogenous leukemia in children and adults. *N Engl J Med* 1980;303:473-8.
20. Glucksberg H, Cheever MA, Farewell UT, et al. High-dose combination chemotherapy for acute non-lymphoblastic leukemia in adults. *Cancer* 1981;48:1073-81.
21. Cassileth PA, Katz ME. Chemotherapy for adult acute non-lymphocytic leukemia with daunorubicin and cytosine arabinoside. *Cancer Treat Rep* 1977;61:1441-5.
22. Hagbin M. Acute non-lymphoblastic leukemia: Clinical and morphological characterization. *Mod Prob Pediatr* 1975;16:39-58.
23. Pizzo PA, Henderson ES, Leventhal BG. Acute myelogenous leukemia in children. A preliminary report of combination chemotherapy. *J Pediatr* 1976;88:125-30.
24. Report of the Medical Research Council's Working Party on Leukemia in Adults: Treatment of acute myeloid leukemia with daunorubicin, cytosine arabinoside, mercaptopurine, L-asparaginase, prednisone and thioguanine: Results of treatment with five multiple-drug schedules. *Br J Haematol* 1974;27:373-89.
25. Ansari BM, Thompson EN, Whittaker JA. A comparative study of acute myeloblastic leukemia in children and adults. *Br J Haematol* 1975;31:269-77.
26. Gee TS, Haghbin M, Dowling MD Jr, et al. Acute lymphoblastic leukemia. In adults and children: Differences in response with similar therapeutic regimens. *Cancer* 1976;37:1256-64.
27. Spiers ASD, Roberts PD, Marsh GW, et al. Acute lymphoblastic leukemia: Cyclical chemotherapy with three combinations of four drugs (COAP-POMP-CART Regimen). *Brit Med J* 1975;4:614-7.

28. Howard JP, Albo V, Newton WA Jr. Cytosine arabinoside: Results of a cooperative study in acute childhood leukemia. *Cancer* 1968;21:341-5.
29. McElwain TJ, Hardisty RM. Remission induction with cytosine arabinoside and L-asparaginase in acute lymphoblastic leukemia. *Brit Med J* 1969;4:596-8.
30. Bodey GP, Rodriguez V, Hart J, et al. Therapy of acute leukemia with the combination of cytosine arabinoside (NSC-63878) and cyclophosphamide (NSC-26271). *Cancer Chemother Rep* 1970;54:255-62.
31. Nesbitt ME Jr, Hammond D. Cytosine arabinoside (ARAC) and prednisone therapy of previously treated acute lymphoblastic and undifferentiated leukemia (ALL/AUL) of childhood. *Proc Am Assoc Cancer Res* 1970;11:59.
32. Wang JJ, Selawry OS, Vietti TJ, et al. Prolonged infusion of arabinosyl cytosine in childhood leukemia. *Cancer* 1970;25:1-6.
33. Klemperer M, Coccia P, Albo V, et al. Reinduction of remission after first bone marrow relapse in childhood acute lymphoblastic leukemia.. *Proc Am Assoc Cancer Res* 1978;19:414.
34. Ortega JA, Finklestein JZ, Ertel I, et al. Effective combination treatment of advanced acute lymphocytic leukemia with cytosine arabinoside (NSC-63878) and L-asparaginase (NSC-109229). *Cancer Chemother Rep* 1972;56:363-8.
35. Bryan JH, Henderson ES, Leventhal BG. Cytosine arabinoside and 6-thioguanine in refractory acute lymphocytic leukemia. *Cancer* 1974;33:539-44.
36. Proceedings of the chemotherapy conference on ARA-C: Development and application (cytosine arabinoside hydrochloride - NSC 63878), Oct 10, 1969.
37. Lay HN, Colebatch JH, Ekert H. Experiences with cytosine arabinoside in childhood leukemia and lymphoma. *Med J Aust* 1971;2:187-92.
38. Halikowsli B, Cyklis R, Armata J, et al. Cytosine arabinoside administered intrathecally in cerebromeningeal leukemia. *Acta Paediatr Scand* 1970;59:164-8.
39. Wang JJ, Pratt CG. Intrathecal arabinosyl cytosine in meningeal leukemia. *Cancer* 1970;25:531-4.
40. Band PR, Holland JF, Bernard J, et al. Treatment of central nervous system leukemia with intrathecal cytosine arabinoside. *Cancer* 1973;32:744-8.
41. Sullivan MP, Dymont P, Hvizdala E, et al. Favourable Comparison of all out #2 with "total" therapy in the treatment of childhood leukemia - The equivalence of intrathecal chemotherapy and radiotherapy as CNS prophylaxis. Abstracted, Proc of AACR and

- ASCO 1981;22:675.
42. Saiki JH, Thompson S, Smith F, et al. Paraplegia following intrathecal chemotherapy. *Cancer* 1972;29:370-74.
 43. Rubinstein LJ, Herman MM, Long TF, et al. Disseminated necrotizing leukoencephalopathy: A complication of treated central nervous system leukemia and lymphoma. *Cancer* 1975;35:291-305.
 44. Marmont AM, Damasio EE. Neurotoxicity of intrathecal chemotherapy for leukemia. *Brit Med J* 1973;4:47.
 45. Margileth DA, Poplack DG, Pizzo PA, et al. Blindness during remission in two patients with acute lymphoblastic leukemia. *Cancer* 1977;39:58-61.
 46. Hopen G, Mondino BJ, Johnson BL, et al. Corneal toxicity with systemic cytarabine. *Am J Ophthalmol* 1981;91:500-504.
 47. Lazarus HM, Herzig RH, Herzig GP, et al. Central nervous system toxicity of high-dose systemic cytosine arabinoside. *Cancer* 1981;48(12):2577-82.
 48. Slavin RE, Dias MA, Soral R. Cytosine arabinoside induced gastrointestinal toxic alterations in sequential chemotherapeutic protocols - A clinical pathologic study of 33 patients. *Cancer* 1978;42:1747-59.
 49. Haupt HM, Hutchins GM, Moore GW. Ara-C lung: Noncardiogenic pulmonary edema complicating cytosine arabinoside therapy of leukemia. *Am J Med* 1981;70:256-61.
 50. Shafer AI. Teratogenic effects of antileukemic chemotherapy. *Arch Intern Med* 1981;141:514-5.
 51. Wagner VM, et al. Congenital abnormalities in baby born to cytarabine treated mother. *Lancet* 1980;2:98-9.
 52. Frei E III, Bickets JN, Hewlet JS, et al. Dose schedule and antitumor studies of arabinosyl cytosine (NSC 63878). *Cancer Res* 1969;29:1325-32.
 53. Bell WR, Whang JJ, Carbone PP, et al. Cytogenetic and morphologic abnormalities in human bone marrow cells during cytosine arabinoside therapy. *J Hematol* 1966;27:771-81.
 54. Burke PJ, Serpick AA, Carbone PP, et al. A clinical evaluation of dose and schedule of administration of cytosine arabinoside (NSC 63878). *Cancer Res* 1968;28:274-9.
 55. Castleberry RP, Crist WM, Holbrook T, et al. The cytosine arabinoside (Ara-C) syndrome. *Med Pediatr Oncol* 1981;9:257-64.

56. Slevin ML, Pfall EM, et al. The pharmacokinetics of subcutaneous cytosine arabinoside in patients with acute myelogenous leukemia. *Br J Clin Pharmacol* 1981;12:507-10.
57. Munson WJ, Kubiak EJ, Cohon MS. Cytosine arabinoside stability in intravenous admixtures with sodium bicarbonate and in plastic syringes. *Drug Intell Clin Pharm* 1982;16:765-7.
58. Athanikar N, Boyer B, Deamer R, et al. Visual Compatibility of 30 additives with a parenteral nutrient solution. *Am J Hosp Pharm* 1979;36:511-3.
59. Cradack JC, Kleinman LM, Rahman A. Evaluation of some pharmaceutical aspects of intrathecal methotrexate sodium, cytarabine and hydrocortisone sodium succinate. *Am J Hosp Pharm* 1978;35:402-406.
60. Keller JH, Ensminger WD. Stability of cancer chemotherapeutic agents in a totally implanted drug delivery system. *Am J Hosp Pharm* 1982;39:1321-3.
61. Benvenuto JA, Anderson RW, Kerkof K, et al. Stability and compatibility of antitumor agents in glass and plastic containers. *Am J Hosp Pharm* 1981;38:1914-8.
62. McRae MP, King JC. Compatibility of antineoplastic, antibiotic and corticosteroid drugs in intravenous admixtures. *Am J Hosp Pharm* 1976;33:1010-13.
63. Ho D. Potential advances in the clinical use of arabinosylcytosine. *Cancer Treat Rep* 1977;61:717-22.
64. Pfall E, et al. Cytosine arabinoside: Pharmacokinetics following different routes of administration. *Biochem Soc Trans* 1982;10:512-3.
65. Fulton DS, et al. Intrathecal cytosine arabinoside for the treatment of meningeal metastases from malignant brain tumors and systemic tumors. *Cancer Chemother Pharmacol* 1982;8:285-91.
66. Dahl S, et al. Therapeutic efficacy of preventive intrathecal (IT) chemotherapy for children with acute lymphocytic leukemia (ALL) who relapse after cessation of therapy. Abstracted, *Proc of AACR and ASCO* 1979;20:628.
67. Altman AJ, et al. Remission induction in acute non-lymphocytic leukemia (ANLL) with low-dose cytosine arabinoside (ARA-C). *Abstract Pediatr Res* 1982;16(4,Part 2):197A(714).
68. Lister TA, Robatiner AZS. The treatment of acute myelogenous leukemia in adults. *Semin Haematol* 1982;19:3,172-92.
69. Mitrou PS for the AIO. Sequential combination therapy (COP-Bleo+AVP) in non-

- Hodgkin's lymphomas (NHL) of high-grade malignancy stage III and IV. A phase II study. *J Cancer Res Clin Oncol* 1982;103 Suppl:A23.
70. Pichler E, et al. Results of LSA₂-L₂ therapy in 26 children with non-Hodgkin's lymphoma. *Cancer* 1982;50:2740-46.
 71. Preisler HD. High dose cytosine arabinoside therapy in acute non-lymphocytic leukemia. *Eur J Cancer Clin Oncol* 1984;20(2):297-300.
 72. Rohatiner AZS, Slevin ML, Dhaliwal HS, et al. High dose cytosine arabinoside: Response to therapy in acute leukemia and non-Hodgkin's lymphoma. *Cancer Chemother Pharmacol* 1983;12:90-93.
 73. Herzig RH, Wolff SN, Larzaus HM, et al. High-dose cytosine arabinoside therapy for refractory leukemia. *Blood* 1983;62(2):361-9.
 74. Preisler HD, Early AP, Raza A, et al. Therapy of secondary acute non-lymphocytic leukemia with cytarabine. *N Eng J Med* 1983;308(1):21-3.
 75. Willemze R, Zwaan FE, Colpin G, et al. High dose cytosine arabinoside in the management of refractory acute leukemia. *Scand J Haematol* 1982;29:141-6.
 76. Capizzi RL, Poole M, Cooper MR, et al. Treatment of poor risk acute leukemia with sequential high dose ARA-C and asparaginase. *Blood* 1984;63(3):694-700.
 77. Johnson H, Smith TJ, Desforges J. Cytosine arabinoside induced colitis and peritonitis: Non-operative management. *J Clin Oncol* 1985;3(5):607-12.
 78. Dunton SF, Ruprecht N, Spruce W, et al. Progressive ascending paralysis following administration of intrathecal and intravenous cytosine arabinoside. *Cancer* 1986;57:1083-8.
 79. Takvorian T, Anderson K, Ritz J. A fetal cardiomyopathy associated with high dosage ARA-C (HIDAC) and cyclophosphamide (CTX) in bone marrow transplantation (BMTx). Abstract submitted for 1985 AACR meetings in Houston, Texas.
 80. Anderson BS, Cogan B, Keating MJ, Estey EH, et al. Subacute pulmonary failure complicating therapy with high dose ARA-C in acute leukemia. *Cancer* 1985;56(9):2181-4.
 81. Altman AJ, Dindorf P, Quinn JJ. Acute pancreatitis in association with cytosine arabinoside therapy. *Cancer* 1982;49:1384-6.
 82. Powell BL, Capizzi RL, Lysterly EW, et al. Peripheral neuropathy after high-dose cytosine arabinoside, daunorubicin, and asparaginase consolidation for acute non-lymphocytic leukemia. *J Clin Oncol* 1986;4(1):95-7.

83. Peters WG, Willenze R, Coely LP. Results of induction and consolidation treatment with intermediate and high dose ARA-C and m-AMSA containing regimens in patients with primarily failed or relapsed acute leukemia and non-Hodgkin's lymphoma. *Scand J Haematol* 1986;36 Suppl 44:7-16.
84. Marmont AM, Dimasio EE. Neurotoxicity of intrathecal chemotherapy for leukemia. *Brit Med J* 1973;4:47.
85. Margileth DA, Peplack DG, Pizzo PA, et al. Blindness during remission in two patients with acute lymphoblastic leukemia. *Cancer* 1977;39:58-61.
86. Trissel LA. Handbook on injectable drugs, 7th Ed. American Society of Hospital Pharmacists 1992;267-71.
87. Crampton JD, Cohon MS, Lummis WL, et al: Cytosine Arabinoside Stability in Three Intravenous Infusion Solutions at Three Temperatures, Upjohn Technical Report, Code No. 7262/82/7262/037, December 10, 1982.
88. Hassing, DH: 8-Day Stability of CYTOSAR in a Dextrose-NaCl-KCl Infusion Solution, Upjohn Interoffice Memo to J.R. Kline, April 7, 1978.
89. Kuhlman J: Inhibition of Digoxin Absorption but not of Digitoxin During Cytostatic Drug Therapy, *Arzneim Forsch* 32:698-704, 1982.
90. Moody MR, Morris MJ, Yang VM, et al: Effect of Two Cancer Chemotherapeutic Agents on the Antibacterial Activity of Three Antimicrobial Agents, *Antimicrob. Agents Chemother.*, 14:737-742, 1978.
91. Holt RJ: Clinical Problems with 5-fluorocytosine, *Mykosen*, 21(11):363-369, August, 1978.
92. Polak A, Grenson M: Interference Between the Uptake of Pyrimidines and Purines in Yeasts, *Path. Microbiol.*, 39:37-38, 1973.
93. Nand et al, Neurotoxicity Associated With Systemic High-Dose Cytosine Arabinoside, *J Clin Oncol* 1986;4:571-5.
94. Damon et al, The Association Between High-Dose Cytarabine Neurotoxicity and Renal Insufficiency, *J Clin Oncol* 1989;7:1563-8.
95. Reykdal S, Sham R, Kouides P: Cytarabine-Induced Pericarditis: A Case Report and Review of the Literature of the Cardio-Pulmonary Complications of Cytarabine Therapy. *Leukemia Research* 1995; 19:141-144.
96. Watterson J, Toogood I, Nieder M, et al: Excessive Spinal Cord Toxicity From Intensive

Central Nervous System-Directed Therapies. *Cancer* 1994; 74:3034-3041.

97. Pharmascience Inc., Product Monograph ^{Pr}CYTOSAR, Control No. 204769, Date of Revision: May 17, 2017.

APPENDIX A

LSA₂-L₂ Protocol

Woolner N, Burchenal JH, Lieberman PH, et al: Non-Hodgkin's Lymphoma in Children - A Comparative Study of Two Modalities of Therapy. *Cancer* 1976;37:123-134.

Induction Phase

Day 1. Cyclophosphamide 1200 mg / m² single push injection.

Day 3 to 31. Prednisone 60 mg / m² PO divided into three daily doses.

Day 3, 10, 17, 24. Vincristine 1.5 to 2.25 mg / m² intravenously.

Day 5, 27, 30. Spinal tap and intrathecal injection of Methotrexate 6.25 mg/m²

Day 12, 13. Daunomycin 60 mg / m² intravenously.

At the end of induction (last dose of intrathecal methotrexate) patient rests for 3-5 days before consolidation.

Consolidation Phase

Day 34 or 36, daily intravenous injections of cytosine arabinoside (Ara-C) 150 mg / m² for a total of 15 injections are given. (Injections are given from Monday through Friday.)

Thioguanine 75 mg / m² is given orally, 8-12 hours after the injection of Ara-C. If the white blood count is 1500 or more and the platelet count 150000 or more on the 5th day of Ara-C, the patient continues to receive the same dosage of thioguanine over the weekend. However, both are discontinued temporarily when there is evidence of marrow depression; this usually occurs after the initial seventh to tenth doses of the combination and ordinarily recovers within 7-10 days. Hence, the patients may receive more than 15 doses of thioguanine orally, but receive only 15 doses of IV cytosine arabinoside (Ara-C). This first phase of the consolidation takes an average of 30-35 days. The second phase of the consolidation should be started immediately after completion of the 15 doses of Ara-C; it entails daily IV administration of L-asparaginase, 60000 U/m² for a total of 12 injections, excluding weekends.

Two days after the last injection of the L-asparaginase, two more intrathecal (IT) injections of methotrexate are given 2 days apart. Three days after the last IT methotrexate, BCNU [1, 3-Bis (2 chloroethyl 1-1-nitrosourea)] 60 mg / m² is given IV, which completes the consolidation. The average duration of the induction and consolidation is 85-100 days.

Maintenance Phase

The maintenance period consists of five cycles of 5 days each and is started 3-4 days after completion of consolidation.

Cycle I: Oral thioguanine 300 mg / m² for 4 consecutive days: i.v. cyclophosphamide 600 mg / m² on the 5th day.

Rest 7-10 days.

Cycle II: Oral hydroxyurea 2400 mg / m² for 4 consecutive days: i.v. daunomycin 45 mg / m² on the 5th day.

Rest 7-10 days.

Cycle III: Oral methotrexate 10 mg/m² for 4 consecutive days: IV BCNU 60 mg / m² on the 5th day.

Rest 7-10 days.

Cycle IV: I.V. Ara-C 150 mg / m² for 4 consecutive days: i.v. vincristine 1.5 mg / m² on day 5.

Rest 7-10 days.

Cycle V: Two doses of IT methotrexate 6.25 mg / m² 2-3 days apart.

Rest 7-10 days and restart with Cycle I.

PART III: CONSUMER INFORMATION

Pr Cytarabine Injection BP
(cytarabine injection)

This leaflet is part III of a three-part "Product Monograph" published when Cytarabine Injection BP was approved for sale in Canada and is designed specifically for Consumers. This leaflet is a summary and will not tell you everything about Cytarabine Injection BP. Contact a member of your healthcare team if you have any questions about the drug.

ABOUT THIS MEDICATION

What the medication is used for:

Cytarabine Injection BP is used to treat patients with cancer of the blood (leukemia) or cancer of the lymph nodes (lymphoma). It is used alone or in combination with other medicines.

What it does:

Cytarabine Injection BP slows or stops the growth of cancer cells.

When it should not be used:

Do not take Cytarabine Injection BP:

- If you/the child in your care are allergic (hypersensitive) to cytarabine or any of the other ingredients in Cytarabine Injection BP (see "What the nonmedicinal ingredients are" section of this leaflet).

What the medicinal ingredient is:

Cytarabine.

What the nonmedicinal ingredients are:

Hydrochloric acid solution and/or sodium hydroxide solution to adjust the pH, water for injection.

What dosage forms it comes in:

Cytarabine Injection BP is supplied in single use vials of 100 mg / 1 mL (5 vials per box), 500 mg/5 mL (5 vials per box), 1000 mg / 10 mL (1 vial per box) and 2000 mg / 20 mL (1 vial per box).

WARNINGS AND PRECAUTIONS

Serious Warnings and Precautions

Cytarabine Injection BP should be prescribed and monitored only by doctors with experience with cancer medicines.

The following are serious side effects of Cytarabine Injection BP :

- **Serious Allergic Reaction:** Symptoms include sudden wheeziness, difficulty in breathing, swelling of eyelids, face or lips, rash or itching (especially affecting the whole body), hives.
- **Cytarabine Injection BP can cause damage to the heart.** Tell your doctor right away if you/the child in your care have chest pain, shortness of breath, swelling of the legs or irregular heartbeat.
- **Cytarabine Injection BP can cause changes to the lungs.** Tell

- your doctor right away if you/the child in your care develop wheezing, cough, fever or feeling of breathlessness, or if existing breathing problems get worse.
- **Cytarabine Injection BP can have harmful effects on the nervous system.** Tell your doctor right away if you/the child in your care feel drowsy or confused, dizzy or unsteady, get headaches or personality changes.
- **Cytarabine Injection BP can have harmful effects on the stomach and gut that can sometimes be fatal.** Tell your doctor right away if you/the child in your care feel sick or vomit, have diarrhea, a loss of appetite or abdominal pain.
- **Cytarabine Injection BP can cause a decrease in the number of white blood cells, red blood cells, and platelets** (low blood cell counts). This means that you/the child in your care may bruise or bleed more easily. Tell your doctor right away if you/the child in your care get infection, bleeding, fever, or chills with shivering, bruising or rash.

While you/the child in your care are being given Cytarabine Injection BP your doctor will monitor your blood counts (white blood cells, red blood cells, platelets) as well as your liver and kidney function by doing regular blood tests.

A preservative called **benzyl alcohol** is sometimes added to Cytarabine Injection BP. This preservative should not be given to low birth weight or premature babies.

Cytarabine Injection BP may cause Tumour Lysis Syndrome (TLS). This happens when cytarabine makes the cancer cells break down very quickly. This releases uric acid (a waste product) into the blood. The kidneys usually get rid of uric acid but may not be able to cope with large amounts. This can cause serious imbalances in the blood that affect the kidneys and the heart. Tell your doctor immediately if you/the child in your care have palpitations/irregular heartbeats; vomiting; fatigue/weakness; difficulty concentrating/trouble thinking; swelling, numbness or tingling in hands, face or feet; back pain; muscle cramps; fainting or trouble breathing.

Vaccination with a live vaccine should be avoided while being treated with Cytarabine Injection BP. Tell your doctor that you/the child in your care are on Cytarabine Injection BP before getting any vaccine.

Cases of sudden inflammation of the pancreas and cases of paralysis, at times fatal in children, have been reported with the use of cytarabine in combination with other drugs.

Serious nervous system side effects that ranged from headache to paralysis, coma and stroke-like episodes have been reported mostly in children (under 18 years of age) given intravenous (injected into the vein) cytarabine in combination with intrathecal (injected into the spinal cord) methotrexate.

The safety of Cytarabine Injection BP in infants (under 1 year of age) is not known.

Before you start treatment with Cytarabine Injection BP, tell your doctor if you, or the child in your care, have any of the following:

- Liver or kidney problems;
- Heart problems;
- Lung problems;
- Stomach or gut problems
- Low blood cell counts;
- Skin problems.

Pregnancy, Breastfeeding and Fertility:

Cytarabine Injection BP may harm your baby/unborn baby.

Do not become pregnant while being treated with Cytarabine Injection BP. Women who may become pregnant must use effective birth control during treatment and for 3 months after treatment has finished.

If you are pregnant, think you might be pregnant or are planning to have a baby, ask your doctor for advice before starting treatment with Cytarabine Injection BP.

Tell your doctor immediately if you become pregnant.

Do not breastfeed while you are being treated with Cytarabine Injection BP.

Male Fertility:

Do not father a child while being treated with Cytarabine Injection BP and for 3 months after stopping treatment. Use condoms and do not donate sperm during treatment and for 3 months after your treatment has finished. If you plan to father a child, talk to your doctor before starting treatment with Cytarabine Injection BP.

Driving and using machines:

If you feel drowsy or dizzy, do not drive or use machinery.

INTERACTIONS WITH THIS MEDICATION

Serious Drug Interactions

If Cytarabine Injection BP is given to you/the child in your care with methotrexate (another drug used to treat cancer), you have more chances of having serious side effects on your nervous system such as headache, paralysis, coma and stroke-like episodes.

Tell your doctor or pharmacist about any medicines you or the child in your care are on or have taken (including the ones you don't need a prescription for), especially the following:

- 5-Fluorocytosine (a medicine used to treat fungal infections)
- Digoxin (a heart medicine)
- Gentamicin (an antibiotic)
- Cyclophosphamide, vincristine and prednisone

PROPER USE OF THIS MEDICATION

Cytarabine Injection BP will be given to you or the child in your care as an injection or an infusion. It can be given:

- Into the spinal cord
- Into a vein (through a "drip")
- Under the skin

Chemotherapy is usually given during several cycles of treatment over a few months. The length of your/the child in your care's treatment and the number of cycles you or the child in your care need will depend on the type of cancer you/they have. Your doctor will discuss your treatment plan with you.

Usual dose:

The dose of Cytarabine Injection BP you or the child in your care will be given will be calculated by your doctor based on your/the child's weight and height.

Overdose:

In case of drug overdose, contact a healthcare practitioner, hospital emergency department or regional poison control centre, even if there are no symptoms.

Missed dose:

Call your doctor for instructions if you/the child in your care miss an appointment for your cytarabine injection.

SIDE EFFECTS AND WHAT TO DO ABOUT THEM

Side effects of Cytarabine Injection BP may include:

- Viral, bacterial, or fungal infections: Infections can be serious and may lead to death. Contact your doctor if you or the child in your care have fever, chills, or any other signs or symptoms of a possible infection.
- Cytarabine Syndrome: Cytarabine Injection BP may cause a reaction called Cytarabine Syndrome 6 to 12 hours after it has been given. Contact your doctor if you or the child in your care develop fever, muscle pain, bone pain, chest pain, rash, eye problems (pain, itching, redness, discharge, blurred vision), or generally feel unwell.
- Feeling tired or weak.
- Headaches or feeling dizzy, fainting.
- Feeling of pins and needles.
- Nausea, vomiting, diarrhea, loss of appetite, abdominal pain.
- Eye infection, irritation, pain and blurred vision.
- Hair loss, skin rash or open sores, peeling of the skin, itching or increased freckles.
- Swelling of the throat, heartburn, sores and bleeding in the mouth, lips, or on the anus.
- Feeling hot and feverish.
- Sore throat.
- Muscle pain, bone pain.
- Fast heartbeat.
- Rash or blisters on the palms of the hands and soles of the feet.

If any of these affects you severely, tell your doctor, nurse or pharmacist.

During treatment, you or the child in your care will need to have regular blood tests. Your doctor will tell you how often this should be done. It is important that you do not miss any of these tests.

SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM

Symptom / effect	Talk with your doctor or pharmacist		Stop treatment and seek immediate medical help
	Only if severe	In all cases	
Very Common	Anemia: symptoms include fatigue, loss of energy, weakness, shortness of breath.		✓
	Decreased platelets: symptoms include bruising, bleeding, fatigue and weakness.		✓
	Decreased white blood cells: symptoms include infections, fever, chills with shivering, fatigue, aches pains and flu-like symptoms.		✓
	Infection of the blood: symptoms include feeling dizzy or faint, confusion or disorientation, diarrhea, nausea, vomiting, slurred speech, severe muscle pain.		✓
	Megaloblastic anemia: symptoms include fatigue, weakness, loss of appetite, nausea, diarrhea, fast heartbeat, smooth or tender tongue, tingling or numbness in hands and feet.		✓
	Pneumonia: symptoms include cough with or without mucus, fever, chills, shortness of breath.		✓
	Serious stomach or gut problems: symptoms include severe vomiting, severe diarrhea (increased number of bowel movements, watery or bloody stool, stomach pain and/or cramps).		✓
Frequency not known	Serious allergic reaction: symptoms include rash, hives, swelling of the face, lips tongue or throat, difficulty swallowing or breathing. It may lead to a heart attack.		✓
	Oedema: symptoms include swelling of the stomach, legs, ankles or feet.		✓
	Inflammation of the pancreas: symptoms include abdominal pain that lasts and gets worse when you lie down, fever, nausea, vomiting.		✓
	Injection site reaction: symptoms include pain, redness, warmth, swelling at the injection site or along the vein.		✓

SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM			
Symptom / effect	Talk with your doctor or pharmacist		Stop treatment and seek immediate medical help
	Only if severe	In all cases	
Kidney disorder: symptoms include decreased urination, nausea, vomiting, swelling of extremities, fatigue, difficulty or pain when urinating, blood in the urine.		✓	
Liver disorder: symptoms include yellowing of the skin or eyes, dark urine, abdominal pain, nausea, vomiting, loss of appetite.		✓	
Serious bleeding problems: symptoms include blood in your stool or urine, bleeding that lasts for a long time or that you cannot control, coughing up blood or blood clots, increased bruising, feel dizzy or weak, confusion, change in your speech, or a headache that lasts a long time.			✓
Serious eye problems: symptoms include sensitivity to light, blurry vision, eye pain, tearing, feeling like there is something stuck in your eye.		✓	
Serious heart problems: symptoms include shortness of breath, swelling of the legs, irregular heartbeat, chest pain.			✓
Serious nervous system problems: symptoms include headache, paralysis, coma, stroke-like episodes, drowsiness or confusion, dizziness or unsteadiness, personality changes, shaking and fits, speech problems, involuntary movements.			✓
Tumor Lysis Syndrome: symptoms include nausea, vomiting, decreased urination, irregular heartbeat, confusion, delirium, seizures.			✓

This is not a complete list of side effects. For any unexpected effects while taking Cytarabine Injection BP, contact your doctor or pharmacist.

HOW TO STORE IT

Keep out of reach and sight of children.

This drug will be given to you in a hospital or doctor's office. You will not store it at home.

Medicines should not be thrown down the drain or in the garbage. Ask your pharmacist how to dispose of medicines you no longer need. This will help to protect the environment.

REPORTING SIDE EFFECTS

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<http://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada/adverse-reaction-reporting.html>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

MORE INFORMATION

This document plus the full product monograph, prepared for health professionals can be found by contacting the sponsor, Sandoz Canada Inc., at:

1-800-361-3062

or by written request at:
145 Jules-Léger
Boucherville QC
J4B 7K8

Or by e-mail at:
medinfo@sandoz.com

This leaflet was prepared by Sandoz Canada Inc.

Last revised: December 06, 2017