

Main panel

DIN 01983253



VETERINARY USE ONLY  
USAGE VÉTÉRINAIRE SEULEMENT  
ANTIBIOTIC / ANTIBIOTIQUE

 **DEPOCILLIN®**

**Benzylpenicillin procaine suspension Mfr. Std.**  
**Suspension de benzylpénicilline procaine Norme fabr.**

sterile / stérile

Intervet Canada Corp.,  
subsidiary of/filiale de Merck & Co., Inc.  
16750, route Transcanadienne  
Kirkland, QC H9H 4M7 1-866-683-7838

Side panel

XXX mL

Lot:  
Exp:

**Active Ingredients:** 300,000 I.U. (300 mg) Benzylpenicillin procaine (Ph.Eur.) per mL.





**Preservatives:** 1.1 mg Methylparahydroxybenzoate per mL.

**Indications:** For the treatment of infections caused by bacteria sensitive to penicillin in cattle, swine, horses and sheep. Consult package insert for complete product information.

**Warnings:** Withdrawal period for Cattle and Sheep–meat: 10 days; –milk: 96 hours. Swine–meat: 8 days. Horses–meat: Do not eat. Keep out of reach of children. See package insert for complete warnings.

**Dosage and administration:** Deep intramuscular injection only. Read entire insert before use.

**Dosing chart per species/ Tableau de dosage par espèce**

 horse/ cheval 500 kg 35 mL	 cow/ vache 400 kg 28 mL	 pig/ porc 50 kg 2.5 mL	 sheep/ mouton 50 kg 3.5 mL
---	--	--	---

Note to reviewer: this table is inverted

**DEPOCILLIN® Inner label (100, 250 or 500 mL) - MIR #1**  
**Version 11 SEP 2017**

**Ingrédients actifs:** 300 000 U.I. (300 mg) benzylpénicilline procaine (Ph.Eur.) par mL.

**Agents de conservation:** 1,1 mg méthylparahydroxybenzoate par mL.

**Indications:** Pour le traitement des infections causées par des bactéries sensibles à la pénicilline chez les bovins, les porcs, les chevaux et les moutons. Voir la notice pour de l'information complète sur le produit.

**Mises en garde:** Délais d'attente pour Bovins et Moutons–viande: 10 jours; –lait: 96 heures. Porcs–viande: 8 jours. Chevaux–viande: Ne pas consommer. Garder hors de la portée des enfants. Voir la notice pour les mises en garde complètes.

**Posologie et administration:** Administrer seulement par voie intramusculaire profonde. Lire la notice au complet avant l'utilisation du produit.

Main panel English

DIN 01983253

XXX mL

VETERINARY USE ONLY  
ANTIBIOTIC

 **DEPOCILLIN®**  
**Benzylpenicillin Procaine suspension Mfr. Std.**

sterile

**Warnings:** Cattle and Sheep: Treated animals must not be slaughtered for use in food for at least 10 days after the latest treatment with this drug. Milk taken from treated animals during treatment and within 96 hours after the latest treatment with this drug must not be used as food. To avoid illegal residues in milk: Do not use in dairy animals with clinical or subclinical mastitis and do not exceed the recommended dosage. Swine: Treated animals must not be slaughtered for use in food for at least 8 days after the latest treatment with this drug. Horses: This drug is not to be administered to horses that are to be slaughtered for use in food. To avoid illegal residues, do not inject subcutaneously. Keep out of reach of children. See package insert for complete warnings.



Side panel English

**Each mL contains:**

Active Ingredients:

Benzylpenicillin procaine (Ph.Eur.)..... 300,000 I.U. (300 mg)

Preservatives:

Methylparahydroxybenzoate.....1.1 mg

Non-medicinal Ingredients:

Sodium citrate dihydrate.....17.60 mg

Potassium Di-Hydrogen Phosphate.....0.54 mg

Edetate Disodium dihydrate.....0.11 mg

Povidone K30.....1.50 mg

Lecithin.....3.30 mg

Sodium hydroxide.....q.s.

Phosphoric acid.....q.s.

Water for Injection.....ad 1 mL





**Indications:**

For the treatment of infections caused by bacteria susceptible to penicillin in cattle, swine, horses and sheep. Consult package insert for complete product information.

**DEPOCILLIN® Outer label (100, 250 or 500 mL) - MIR #1**  
**Version 11 SEP 2017**

**Dosage and administration:** DEPOCILLIN should be given by deep intramuscular injection only. Read entire insert before use. Shake well before using.

**Dosing chart per species**

 horse 500 kg 35 mL	 cow 400 kg 28 mL	 pig 50 kg 2.5 mL	 sheep 50 kg 3.5 mL
---	---	--	---

**Storage:** Store below 15°C. Do not freeze. Protect from light.

Intervet Canada Corp.  
subsidiary of Merck & Co., Inc.  
16750, route Transcanadienne  
Kirkland, QC H9H 4M7  
1-866-683-7838

---

*Top panel:*

 **DEPOCILLIN®**  
**VETERINARY USE ONLY**  
**USAGE VÉTÉRINAIRE SEULEMENT**

XXX mL



---

*Bottom panel:*

Lot:  
Exp:

**DEPOCILLIN® English insert - MIR #1**  
**Version 11 SEP 2017**

DIN 01983253

VETERINARY USE ONLY  
ANTIBIOTIC

 **DEPOCILLIN®**

**Benzylpenicillin procaine suspension Mfr. Std.**

**DESCRIPTION:**

DEPOCILLIN is a sterile aqueous antibiotic suspension for injection, containing per mL:

**ACTIVE INGREDIENTS:**

Benzylpenicillin Procaine (Ph.Eur.).....300,000 I.U. (300 mg)

**PRESERVATIVES:**

Methylparahydroxybenzoate..... 1.1 mg

**NON-MEDICINAL INGREDIENTS:**

Sodium citrate dihydrate.....17.60 mg

Potassium Di-Hydrogen Phosphate.....0.54 mg

Edetate Disodium dihydrate.....0.11 mg

Povidone K30.....1.50 mg

Lecithin.....3.30 mg

Sodium hydroxide.....q.s.

Phosphoric acid.....q.s.

Water for Injection.....ad 1 mL

**INDICATIONS:**

DEPOCILLIN is indicated for the treatment of infections caused by bacteria susceptible to penicillin in cattle, swine, horses and sheep:

Cattle: Bacterial pneumonia; calf diphtheria; footrot; metritis; wound infections.

Swine: Bacterial pneumonia; erysipelas; wound infections.

Horses: Bacterial pneumonia; metritis; strangles; wound infections.

Sheep: Bacterial pneumonia; metritis; wound infections.

**DOSAGE AND ADMINISTRATION:**

DEPOCILLIN should be given by deep intramuscular injection only, using a sterile hypodermic needle and a syringe. Shake well before using.

**Cattle and Sheep:**

21,000 I.U./kg of body weight (7 mL/100 kg or 0.7 mL/10 kg) intramuscularly once daily until 2 days after clinical signs disappear. Do not exceed 5 days of treatment. Do not exceed 15 mL per injection site in cattle and 5 mL per injection site in sheep.

**Swine:**

15,000 I.U./kg of body weight (1 mL/20 kg) intramuscularly once daily until 2 days after clinical signs disappear. Do not exceed 5 days of treatment. Do not exceed 10 mL per injection site.

**Horses:**

21,000 I.U./kg of body weight (7 mL/100 kg) intramuscularly twice daily (at 12 hour intervals) until 2 days after clinical signs disappear or, in the case of respiratory

**DEPOCILLIN® English insert - MIR #1**  
**Version 11 SEP 2017**

infections, until 5 days after clinical signs disappear. Do not exceed 15 mL per injection site. Rotate injection sites for succeeding doses.

If animals do not respond within 48 hours of administration of the first dose, consult a veterinarian. An alternative antibacterial therapy should be considered.

**CAUTIONS:**

In case of anaphylactic reaction (acute respiratory distress, mouth breathing, recumbency), administer epinephrine immediately to prevent death.

In pigs, the administration of this product may occasionally cause a fever, vomiting, shivering, listlessness, incoordination and possibly, death. In pregnant sows and gilts, it may result in abortions.

In horses, Benzylpenicillin procaine is occasionally associated with excitement and may cause pain at the site of injection. Repeated use of the same injection site may intensify these side effects.

**WARNINGS:**

**Cattle and Sheep:** Treated animals must not be slaughtered for use in food for at least 10 days after the latest treatment with this drug. Milk taken from treated animals during treatment and within 96 hours after the latest treatment with this drug must not be used as food.

To avoid illegal residues in milk: Do not use in dairy animals with clinical or subclinical mastitis and do not exceed the recommended dosage.

The use of on-farm tests for detecting subclinical mastitis prior to treatment (C.M.T. or other tests) and/or for detecting residues prior to marketing milk from treated animals is recommended.

**Swine:** Treated animals must not be slaughtered for use in food for at least 8 days after the latest treatment with this drug.

**Horses:** This drug is not to be administered to horses that are to be slaughtered for use in food.

To avoid illegal residues, do not inject subcutaneously.  
Keep out of reach of children.

**STORAGE:**

Store below 15°C. Do not freeze. Protect from light.

**HOW SUPPLIED:**

100, 250 and 500 mL plastic bottles.

Intervet Canada Corp.,  
subsidiary of Merck & Co., Inc.  
16750, route Transcanadienne,  
Kirkland, QC H9H 4M7  
1-866-683-7838



® Intervet International B.V. Used under license.

Version 11 SEP 2017