

Pr OXYTETRACYCLINE 100

GRANULAR PREMIX

VETERINARY USE ONLY

DIN 00654787

ANTIBIOTIC

ACTIVE INGREDIENT

Oxytetracycline hydrochloride.....220 g/kg

INDICATIONS, DOSAGE and DIRECTIONS FOR USE

To reduce the development of antimicrobial resistance and maintain effectiveness, use this antibiotic prudently and for the shortest duration required to achieve the desired clinical outcome.

SWINE

INDICATIONS	Required activity of oxytetracycline HCl (g/tonne)	kg OXY 100 per tonne of complete feed	FEEDING DIRECTIONS
As an aid in the prevention of bacterial enteritis (scours).	220	1	Feed as the sole ration up to 6 weeks of age.
	55	0.25	Feed as the sole ration from six weeks of age up to 27-32 kg body weight
As an aid in the treatment of bacterial enteritis (scours).	220	1	Feed as the sole ration following the appearance of symptoms and until three days after symptoms disappear — for pigs up to six weeks of age.
	110	0.5	Feed as the sole ration following the appearance of symptoms and until three days after symptoms disappear — for pigs greater than six weeks of age. <u>Note:</u> Pigs off feed should be given individual treatment.
As an aid in reducing the incidence of abortion caused by leptospirosis. <u>Note:</u> Good sanitation procedures must be practiced to prevent spread of infections.	550	2.5	When abortion in swine has been diagnosed due to leptospirosis, feed at once as the sole ration to pregnant and affected sows for a two week period.

BEEF CATTLE

INDICATION	DOSAGE & DIRECTIONS FOR USE
As an aid in reducing the incidence of bloat in young cattle on pasture and in feedlots.	Feed continuously. Level of oxytetracycline hydrochloride in the complete feed is to be such that each animal will receive 75 mg per day. 55 g of oxytetracycline hydrochloride (0.25 kg of OXY 100) per tonne (1000 kg) of complete feed when fed at the rate of 1.36 kg per head per day meets the 75 mg level stated above.

CALVES

INDICATIONS	Required activity of oxytetracycline HCl (g/tonne)	kg OXY 100 per tonne of complete feed	FEEDING DIRECTIONS
As an aid in the prevention of	55	0.25	Feed continuously during the period of

bacterial enteritis (scours).			early growth (up to 136 kg body weight).
-------------------------------	--	--	--

TURKEYS

INDICATIONS	Required activity of oxytetracycline HCl (g/tonne)	kg OXY 100 per tonne of complete feed	FEEDING DIRECTIONS
As an aid in the prevention of infectious sinusitis of turkeys.	110	0.5	Feed as the sole ration for as long as the birds are expected to be exposed to the infection.
As an aid in the prevention of infectious synovitis of turkeys.	110	0.5	Feed as the sole ration for as long as the birds are expected to be exposed to the infection.
As an aid in the treatment of infectious synovitis of turkeys	220	1	Feed as the sole ration when symptoms appear continue until at least two weeks after symptoms disappear.

LAMBS

INDICATIONS	Required activity of oxytetracycline HCl (g/tonne)	kg OXY 100 per tonne of complete feed	FEEDING DIRECTIONS
As an aid in the reduction of bacterial enteritis (scours) in creep-fed suckling lambs.	110	0.5	Feed as the sole creep feed during the suckling period.
As an aid in the reduction of losses due to Enterotoxemia (overeating disease) in feedlot lambs.	22	0.1	Feed as the sole ration while lambs are in the feedlot.

CAUTIONS

This drug premix must be thoroughly mixed in complete feeds.
Do not feed undiluted.

WARNINGS

Treated animals must not be slaughtered for use in food for at least 7 days for swine and turkeys, 5 days for cattle and calves, and 4 days for lambs after the latest treatment with this drug.
Do not administer to lactating dairy cattle.
When handling the product, avoid inhalation, oral exposure and direct contact with skin or eyes.
Keep out of reach of children.

STORAGE

Store in a cool dry place below 30°C.

NET CONTENTS 25 kg

MANUFACTURED BY

Bio Agri Mix LP

PO Box 399
Mitchell, ON
N0K 1N0



**CONTAINS
ANTIMICROBIAL
USE RESPONSIBLY**
**CONTIENT UN
ANTIMICROBIEN
UTILISEZ DE FAÇON RESPONSABLE**

LOT NO / NUMERO DE LOT:
EXP.
MADE IN CANADA