

English Panel

DIN 01939807

Sodium Sulfamethazine 25% Solution

Veterinary Use Only
Antibiotic

Net 4 L

ACTIVE INGREDIENT:

A non-sterile solution containing 25 grams of sodium sulfamethazine in each 100 mL.

INDICATIONS:

Cattle & Sheep: As an aid in the treatment of the following conditions caused by bacteria susceptible to sulfamethazine: foot rot, shipping fever, bacterial enteritis/scours, metritis, mastitis and bacterial respiratory diseases. As an aid in the treatment of coccidiosis in sheep.

Calves: As an aid in the treatment of the following conditions caused by bacteria susceptible to sulfamethazine: bacterial respiratory infections and secondary infections in bacterial enteritis/scours. As an aid in the prevention of coccidiosis.

Swine: As an aid in treatment of bacterial enteritis, mastitis, metritis, and bacterial respiratory diseases caused by bacteria susceptible to sulfamethazine.

Horses: As an aid in the treatment of strangles, navel ill, joint ill (septic arthritis), bacterial enteritis and many secondary bacterial infections, associated with respiratory virus infections, caused by bacteria susceptible to sulfamethazine.

Poultry: As an aid in the treatment of caecal coccidiosis in chickens.

DOSAGE AND ADMINISTRATION:

To reduce the development of antimicrobial resistance and maintain effectiveness, use this antibiotic prudently.

For oral use.

Dairy Cattle, Calves, Sheep, Swine and Horses: First day: 45 mL of 25% sodium sulfamethazine for each 50 kg of body weight. Second and following days: one half of the above dosage. Do not treat for more than 5 days.

Beef Cattle: First day: for each 100 kg of body weight, add 100 mL of 25% sodium sulfamethazine into the amount of water that will be consumed daily by the animals. Second and following days: one half of the above dosage. Do not treat for more than 5 days.

Poultry: Mix 35 mL of product per 9 litres of drinking water for 2 days - then reduce dosage to 17.5 mL of product per 9 litres of drinking water for 5 additional days. Allow access to medicated water only.

WARNINGS:

Milk taken from treated animals during treatment and within 96 hours after the latest treatment must not be used as food. Treated animals must not be slaughtered for use in food for at least 12 days after the latest treatment with this drug. This product must not be added to swine feeds. Do not use in laying


birds. Sulfonamides can cause allergic reactions in sensitized individuals. When handling the product, avoid inhalation, oral exposure and direct contact with skin or eyes. Keep out of reach of children.

STORAGE:

Store at room temperature (15-30°C).

Manufactured by/Fabriqué par:
Citadel Animal Health/Santé Animale
Cambridge, ON N3C 2W4

Ditributed by/Distribué par:
Vetoquinol N.-A. Inc.
2000, ch. Georges, Lavaltrie, QC
CANADA J5T 3S5



8JUG014A Rev. 09/17
Code 1SUL001