

<p><b>Indications:</b> For use in retained placenta shreds, metritis and post partum infections in cattle.</p> <p><b>Active Ingredients:</b> Each bolus contains: Urea ..... 13.44 g Sulfanilamide ..... 1.92 g Sulfathiazole ..... 320 mg</p> <p><b>Directions for use:</b> <i>To reduce the development of antimicrobial resistance and maintain effectiveness, use this antibiotic prudently.</i> Insert one (1) or two (2) boluses deeply into the uterus.</p> <p><b>Warnings:</b> Milk taken from treated cattle during treatment and within 96 hours after the latest treatment must not be used in food. Treated animals must not be slaughtered for use in food for at least 10 days after the latest treatment with this drug. Sulfonamides can cause allergic reactions in sensitized individuals. When handling the product, avoid inhalation, oral exposure and direct contact with skin or eyes. Keep out of reach of children.</p> <p><b>Storage:</b> Store at room temperature (15-30°C).</p> <p>Professional Veterinary Laboratories. 1199 Sanford Street Winnipeg, Manitoba R3E 3A1 1-800-465-7122 - <a href="http://www.domvet.com">www.domvet.com</a></p>	<p>DIN 00308226</p> <p><b>Pr AFTER-CALF BOLUS</b></p> <p>Veterinary use only Usage vétérinaire seulement</p> <p>Antibiotic/Antibiotique</p> <p><b>Warnings/Mises en garde :</b> Refer to the side panel for complete warnings. Référez-vous au panneau latéral pour les mises en garde complètes.</p> <p><b>Net Contents/Contenu Net:</b> X</p>	<p><b>Indications :</b> A utiliser dans les lambeaux de placenta retenus, en cas de métrite ou d'infection puerpérales chez les bovins.</p> <p><b>Ingrédients actifs :</b> Chaque bol contient : Urea ..... 13,44 g Sulfanilamide ..... 1,92 g Sulfathiazole ..... 320 mg</p> <p><b>Mode d'emploi :</b> <i>Pour réduire le développement de résistance aux antimicrobiens et maintenir l'efficacité de cet antibiotique, l'utiliser avec prudence.</i> Insérez un ou deux bols profondément dans l'utérus.</p> <p><b>Mises en garde :</b> Le lait provenant des bovins traités dans un délai de 96 heures après la dernière dose ne doit pas être utilisé comme aliment. Les animaux traités ne doivent pas être abattus à des fins alimentaires dans un délai d'au moins 10 jours après le dernier traitement avec ce médicament. Les sulfamidés peuvent causer des réactions allergiques chez les individus sensibilisés. Lors de la manipulation du produit, éviter l'inhalation, l'exposition orale et le contact direct avec la peau ou les yeux. Garder hors de la portée des enfants.</p> <p><b>Entreposage :</b> Conserver entre 15 et 30°C.</p>
--	---	--

LABEL

X: 12 Boluses, 20 Boluses – VER 9 NOV 2017

(L) EXP		
------------	--	--

LABEL

X: 12 Boluses, 20 Boluses – VER 9 NOV 2017