

<p>Active ingredients: Each gram contains: Sulfanilamide 38.8 mg Sulfathiazole 38.8 mg Urea 98.8 mg</p> <p>Indications: A topical antibacterial agent for use in cattle, dogs, goats, horses, sheep and swine for the treatment of infections caused by bacteria susceptible to sulfonamides. Also for use in the treatment of skin conditions.</p> <p>Directions for use: <i>To reduce the development of antimicrobial resistance and maintain effectiveness, use this antibiotic prudently and for the shortest duration required to achieve the desired clinical outcome.</i> Apply an amount sufficient to cover the affected tissue twice daily.</p> <p>Warnings: Sulfonamides can cause allergic reactions in sensitized individuals. When handling the product, avoid inhalation, oral exposure and direct contact with skin or eyes. Keep out of reach of children.</p> <p>Storage: Store at room temperature (15-30°C).</p> <p>Professional Veterinary Laboratories 1199 Sanford Street Winnipeg, Manitoba R3E 3A1 1-800-465-7122 - www.domvet.com</p> <p>(L) EXP</p>	<p>DIN 00502650</p> <p>Pr SULFA UREA CREAM</p> <p>Veterinary use only Usage vétérinaire seulement</p> <p>Antibiotic/Antibiotique</p> <p>Warnings/Mises en garde : See the side panel for complete warnings. Voir le panneau latéral pour les mises en garde complètes</p> <p>Net Contents/Contenu Net : 400 g</p>	<p>Ingrédients actifs : Chaque gramme contiens : Sulfanilamide 38,8 mg Sulfathiazole 38,8 mg Urea 98,8 mg</p> <p>Indications : Un agent bactériostatique topique pour utiliser dans les bovins, les chiens, les chèvres, les chevaux, les moutons et les porcs pour le traitement des infections causées par les bactéries sensibles aux sulfamidés. Utilisé pour le traitement de la pyoderme.</p> <p>Mode d'emploi : <i>Pour réduire le développement de résistance aux antimicrobiens et maintenir l'efficacité de cet antibiotique, l'utiliser avec prudence et pendant la plus courte durée de temps requise pour obtenir le résultat clinique attendu.</i> Appliquez un montant suffisant pour couvrir le tissu affecté deux fois par jour.</p> <p>Mises en garde : Les sulfamidés peuvent causer des réactions allergiques chez les individus sensibilisés. Lors de la manipulation du produit, éviter l'inhalation, l'exposition orale et le contact direct avec la peau ou les yeux. Garder hors de la portée des enfants.</p> <p>Entreposage : Conserver entre 15 et 30°C.</p>
--	--	---

LABEL

REV#1 – 9 NOV 2017

LABEL

REV#1 – 9 NOV 2017