

<p>Active Ingredient: Each gram contains 55 mg (175 mg / teaspoonful) of tetracycline hydrochloride in a water soluble base.</p> <p>Indications: For the treatment of infections caused by microorganisms susceptible to tetracycline.</p> <p>Directions for Use: <i>To reduce the development of antimicrobial resistance and maintain effectiveness, use this antibiotic prudently and for the shortest duration required to achieve the desired clinical outcome.</i></p> <p>Swine: For bacterial enteritis and respiratory diseases (bacterial) give 2 level teaspoons per 4.5 L of water or milk.</p> <p>Calves: For bacterial scours and bacterial diarrhea, give 1 level teaspoonful once daily (for calves under 45 kg) or 2 level teaspoons once daily (for calves over 45 kg).</p> <p>Lambs: For bacterial enteritis and bacterial diarrhea, give ½ level teaspoon in 60 mL of milk daily as a drench.</p> <p>Poultry: For bluecomb, hexamitiasis, and C.R.D., give 2 level teaspoons per 4.5 L of water.</p> <p>Warnings: Treated animals must not be slaughtered for use in food for at least 5 days after the latest treatment with this drug. Do not feed to laying hens. When handling the product, avoid inhalation, oral exposure and direct contact with skin or eyes. Keep out of reach of children.</p> <p>Storage: Store at room temperature (15-30°C).</p>	<p>DIN 00283851</p> <p style="text-align: center;">Pr KANADOM TETRACYCLINE HYDROCHLORIDE</p> <p style="text-align: center;">Veterinary use only Usage vétérinaire seulement</p> <p style="text-align: center;">Antibiotic/Antibiotique</p> <p style="text-align: center;">Soluble Powder Poudre Soluble</p> <p style="text-align: center;">Warnings/Mises en garde: Refer to the side panel for complete warnings. Référez-vous au panneau latéral pour les mises en garde complètes.</p> <p style="text-align: center;">Net Contents/Contenu Net: 400 g</p>	<p>Ingrédient actif : Chaque gramme contient 55 mg (175 mg / cuillerée à thé) de chlorhydrate de tétracycline sous base hydrosoluble.</p> <p>Indications : Pour le traitement des infections causées par les organismes sensible à tétracycline.</p> <p>Mode d'emploi : <i>Pour réduire le développement de résistance aux antimicrobiens et maintenir l'efficacité de cet antibiotique, l'utiliser avec prudence et pendant la plus courte durée de temps requise pour obtenir le résultat clinique attendu.</i></p> <p>Porcs : Pour maladies respiratoires et entérite d'origine bactérienne donné deux cuillerée à thé rase par 4,5 L d'eau ou de lait.</p> <p>Veaux : Pour diarrhée d'origine bactérienne donné une cuillerée à thé rase par jour (veaux en dessous 45 kg) ou deux cuillerée à thé rase par jour (veaux au dessus 45 kg).</p> <p>Agneaux : Pour entérite et diarrhée d'origine bactérienne donné une demie cuillerée à thé dans 60 mL de lait par jour en potion.</p> <p>Volailles : Pour Crête-Bleu, Hexamitiasis, et C.R.D., donné deux cuillerée à thé rase par 4,5 L d'eau.</p> <p>Mises en garde : Les animaux traités ne doivent pas être abattus à des fins alimentaires dans un délai d'au moins 5 jours après le dernier traitement avec ce médicament. Lors de la manipulation du produit, éviter l'inhalation, l'exposition orale et le contact direct avec la peau ou les yeux. Garder hors de la portée des enfants.</p> <p>Entreposage :</p>
---	---	---

LABEL

VER 9 NOV 2017

<p>Dominion Veterinary Laboratories Ltd. 1199 Sanford Street Winnipeg, Manitoba R3E 3A1 1-800-465-7122 - www.domvet.com</p> <p>(L) EXP</p>		<p>Conserver entre 15 et 30°C.</p>
--	--	------------------------------------

LABEL

VER 9 NOV 2017