

Pr CHLORTETRACYCLINE VITAMIN PREMIX CRUMBLES

Chlortetracycline vitamin premix crumbles

FOR VETERINARY USE ONLY
ANTIBIOTIC

DIN 02370263

ACTIVE INGREDIENTS:

Chlortetracycline Hydrochloride (1)	4.4 g/kg
Vitamin A	550 000 IU/kg
Vitamin D3	55 000 IU/kg
Vitamin E	550 IU/kg

(1). Source: Chlor 100 Premix (DIN 00698210), Bio Agri Mix LP.

Note: For cattle, the amounts of vitamins A and D contained in this premix will supply approximately 50% of the daily requirements. The remaining 50% of vitamins must be supplied by other feed ingredients.

WARNINGS:

1. Treated animals must not be slaughtered for use in food for at least 5 days for cattle and calves, and 7 days for swine after the last treatment with this drug.
2. Do not administer to lactating dairy cattle.
3. When handling the product, avoid inhalation, oral exposure and direct contact with skin or eyes.
4. Keep out of the reach of children.

INDICATIONS:

Medicate with chlortetracycline hydrochloride at 4.4 g/kg as an aid in the following conditions of cattle and swine:

Beef cattle and non-lactating dairy cattle: Prevention of foot rot.

Calves: Prevention of bacterial diarrhea.

Swine:

1. Prevention of bacterial enteritis (scours).
2. Treatment of bacterial enteritis (scours).

Note: Pigs off feed should be given individual treatment.

DIRECTIONS FOR USE:

To reduce the development of antimicrobial resistance and maintain effectiveness, use this antibiotic prudently and for the shortest duration required to achieve the desired clinical outcome.

Species and Indication	Dose Level of Chlortetracycline Hydrochloride	Amount of Chlortetracycline Vitamin Premix Crumbles to mix with other feed ingredients to make a complete ration		Feeding Instructions
		Animal Weight (kg)	Amount per Animal (g)	
Beef and non-lactating dairy cattle Foot rot - Prevention	70 mg/head/day or 0.22 mg/kg of bodyweight per day	-	16	Feed this medicated feed continuously during the period the animals are exposed to this disease.
Beef cattle on full feed may occasionally show digestive disturbances following the sudden introduction of the full dosage of the premix. In such cattle, it is recommended that one-quarter of the dose be fed for 7 days, then gradually increase the dose during the next 7 days up to 16 g of Chlortetracycline Vitamin Premix Crumbles per head per day.				
Calves – Bacterial Diarrhea - Prevention	55 mg/kg of complete feed	45-65 65-90 90-115 115-136	23 33 43 50	Feed the medicated feed continuously up to 136 kg of body weight.
Swine 1. Prevention of bacterial enteritis (scours)	220 mg/kg of complete feed	up to 13	23	Feed this medicated feed as the sole ration up to and including 6 weeks of age.
	55 mg/kg of complete feed	14-23	11	Feed this medicated feed as the sole ration up to 32 kg of body weight.
23-32		16		
2. Treatment of bacterial enteritis (scours)	220 mg/kg of complete feed	up to 13	23	Feed this medicated feed as the sole ration up to and including 6 weeks of age.
	110 mg/kg of complete feed	14-23	23	Feed this medicated feed as the sole ration following the appearance of symptoms and until 3 days after symptoms disappear. Pigs off feed should be given individual treatment.
		23-32	32	

Some common measures and their approximate metric equivalents are as follows: one level teaspoonful = 3 g, one level tablespoonful = 9 g, and one cup = 140 g.

STORAGE: Store in a dry place between 5°C and 30°C.

Net Contents: 25 kg

Manufactured by:
Bio Agri Mix LP
PO Box 399
Mitchell, ON
N0K 1N0



CONTAINS
ANTIMICROBIAL
USE RESPONSIBLY

CONTIENT UN
ANTIMICROBIEN
UTILISEZ DE FAÇON RESPONSABLE

Lot No.:
EXP.: