

P^rLINCOMYCIN SPECTINOMYCIN 100 SOLUBLE POWDER

lincomycin hydrochloride and spectinomycin hydrochloride soluble powder

FOR VETERINARY USE ONLY
ANTIBIOTIC

ACTIVE INGREDIENTS:

Each 150 g jar contains: Lincomycin base (as lincomycin hydrochloride)	33.3 g
Spectinomycin base (as spectinomycin hydrochloride)	66.6 g
Total antibiotic amount	100 g

INDICATIONS:

Swine: For the treatment of swine dysentery (bloody scours) associated with *Serpulina (Treponema) hyodysenteriae*.

Poultry: As an aid in the prevention and treatment of Chronic Respiratory Disease (CRD, Air sacculitis) associated with mycoplasmal and coliform infections of growing chickens.

DOSAGE AND ADMINISTRATION:

To reduce the development of antimicrobial resistance and maintain effectiveness, use this antibiotic prudently.

Swine: Add the contents of this jar to 1500 liters of drinking water (provides 67 mg antibiotic activity per liter). For automatic medicators set to deliver 7.8 mL of stock solution per liter (1 oz per U.S. gallon) prepare stock solution by dissolving the contents of this jar in 11.7 liters of water.

Treatment: At the first sign of disease, administer medication at the above dosage as the sole source of drinking water for 4 to 7 days.

Poultry: Add the contents of this jar to 120 liters of drinking water (provides 833 mg antibiotic activity per liter). For automatic medicators set to deliver 7.8 mL of stock solution per liter of drinking water (1 oz per U.S. gallon), prepare stock solution by dissolving contents of this jar in 0.9 liters of water.

Prevention: To prevent losses due to CRD, administer at the above dosage for the first 3 to 5 days of life and again for one or two days at about 4 weeks of age. The longer period (5 days) of initial administration is recommended if disease has been a problem in the past on the premises.

Treatment: Begin treatment at the above dosage at the first sign of disease. Administer as the sole source of drinking water for 5 to 7 days.

Important: Fresh medicated drinking water should be provided daily. Store stock solutions in closed glass containers in a cool place and discard after 7 days.

WARNINGS:

1. Treated animals must not be slaughtered for use in food for at least 72 hours for poultry and 5 days for swine after the latest treatment with this drug.
2. This product must not be used in laying birds.
3. When handling the product, avoid inhalation, oral exposure and direct contact with skin or eyes.
4. Keep out of reach of children.

CAUTIONS:

LINCOMYCIN SPECTINOMYCIN 100 Soluble Powder

English package insert

1. Treated pigs may develop some stool softening or mild diarrhea within 18 to 36 hours. Signs of mild irritation and swelling of the anus and/or vulva may be associated with the stool changes. These adverse reactions are transient and are self-correcting by the fifth to seventh day of treatment. Should these reactions be more severe than described above, discontinue the use of the drug.

If no improvement in disease condition is noted after 3 to 5 days of treatment, the diagnosis should be re-evaluated.

2. Do not allow rabbits, hamsters, guinea pigs, horses, dairy cattle, or other ruminants access to feeds and/or drinking water containing lincomycin. Ingestion by these species may result in severe gastrointestinal effects or metabolic disorders (i.e. Ketosis in dairy cattle).

STORAGE:

Store in a dry place between 2 and 30 °C.



Bio Agri Mix LP

PO Box 399
Mitchell, ON
N0K 1N0



CONTAINS
ANTIMICROBIAL
USE RESPONSIBLY
CONTIENT UN
ANTIMICROBIEN
UTILISEZ DE FAÇON RESPONSABLE

Rev: BAM Oct-30-17

Rev: BAM Oct 30/17

LINCOMYCIN SPECTINOMYCIN 100 Soluble Powder
Label bilingual main panel

DIN 02230947

Pr **LINCOMYCIN SPECTINOMYCIN 100 SOLUBLE POWDER**
Lincomycin Hydrochloride and Spectinomycin Hydrochloride Soluble Powder
Poudre soluble de chlorhydrate de lincomycine et de chlorhydrate de spectinomycine
VETERINARY USE ONLY / POUR USAGE VÉTÉRINAIRE SEULEMENT
ANTIBIOTIC / ANTIBIOTIQUE
Net contents / contenu net: 150 g

Manufactured by / Fabriqué par



Bio Agri Mix LP
PO Box 399
Mitchell, ON
N0K 1N0



CONTAINS
ANTIMICROBIAL
USE RESPONSIBLY
CONTIENT UN
ANTIMICROBIEN
UTILISEZ DE FAÇON RESPONSABLE

LINCOMYCIN SPECTINOMYCIN 100 Soluble Powder

Label English side panel

ACTIVE INGREDIENTS:

Each 150 g jar contains: Lincomycin base (as lincomycin hydrochloride)	33.3 g
Spectinomycin base (as spectinomycin hydrochloride)	66.6 g
Total antibiotic amount	100 g

INDICATIONS: Swine – For the treatment of swine dysentery (bloody scours) associated with *Serpulina (Treponema) hyodysenteriae*.

Poultry – As an aid in the prevention and treatment of Chronic Respiratory Disease (CRD, Air sacculitis) associated with mycoplasmal and coliform infections of growing chickens.

DOSAGE:

To reduce the development of antimicrobial resistance and maintain effectiveness, use this antibiotic prudently.

Swine – 1 jar (150 g) per 1500 litres of drinking water.

Poultry – 1 jar (150 g) per 120 litres of drinking water.

WARNINGS: Treated animals must not be slaughtered for use in food for at least 72 hours for poultry and 5 days for swine, after the latest treatment with this drug. This product must not be used in laying birds. Keep out of reach of children.

STORAGE: Store in a dry place between 2 and 30°C.

See package insert for complete product information.