

<p>Active Ingredient: Sodium Sulfathiazole 1000 mg/g</p> <p>Indication: For the treatment of infectious Coryza in poultry.</p> <p>Directions for Use: <i>To reduce the development of antimicrobial resistance and maintain effectiveness, use this antibiotic prudently.</i> Mix 3.5 grams of Sulfazole per 4.5 Litres of water and give as the sole source of drinking water for 3-5 days.</p> <p>Cautions: Prepare fresh medicated water daily. If the birds do not respond to treatment within 5 days, discontinue use and re-evaluated the diagnosis.</p> <p>Warnings: This product must not be added to swine feed. Eggs from laying birds must not be used for 5 days following the latest treatment with this drug. Sulfonamides can cause allergic reactions in sensitized individuals. When handling the product, avoid inhalation, oral exposure and direct contact with skin or eyes. Keep out of reach of children.</p> <p>Storage: Store at room temperature (15-30°C).</p> <p>Dominion Veterinary Laboratories Ltd. 1199 Sanford Street Winnipeg, Manitoba R3E 3A1 1-800-465-7122 - www.domvet.com</p>	<p>DIN 00467340</p> <p>Pr SULFAZOLE POWDER Sodium Sulfathiazole Sulfathiazole sodique</p> <p>Veterinary Use Only Usage vétérinaire seulement Antibiotic/Antibiotique</p> <p>Warnings/Mises en garde : See the side panel for complete warnings. Voir le panneau latéral pour les mises en garde complètes.</p> <p>Net Contents/Contenu net : 400 g</p>	<p>Ingrédient actif : Sulfathiazole sodique 1000 mg / g</p> <p>Indication : Pour le traitement d'infectieuse Coryza dans les volailles.</p> <p>Mode d'emploi : <i>Pour réduire le développement de résistance aux antimicrobiens et maintenir l'efficacité de cet antibiotique, l'utiliser avec prudence.</i> Melange 3,5 grammes de Sulfazole par 4,5 Litres d'eau et administré comme la seule source d'eau d'abreuvement pour 3-5 jours.</p> <p>Précautions : Preparez une nouvelle solution chaque jour. Si la condition des volailles n'améliore pas dans un délai de moins 5 jours, arrêtez l'usage et reconsidérer le diagnostic.</p> <p>Mises en garde : Ce produit ne doit pas être ajoutée aux aliments pour porcs. Les oeufs provenant des volailles traitées dans un délai de 5 jours après la dernière dose ne doit pas être utilisé comme aliment. Les sulfamidés peuvent causer des réactions allergiques chez les individus sensibilisés. Lors de la manipulation du produit, éviter l'inhalation, l'exposition orale et le contact direct avec la peau ou les yeux. Garder hors de la portée des enfants.</p> <p>Entreposage : Conserver entre 15 et 30°C.</p> <p>(L) EXP</p>
---	---	--

--	--	--

LABEL

Version 30 NOV 2017