

**Pr BENZAPRO**

Sterile suspension of benzylpenicillin benzathine and benzylpenicillin procaine

Veterinary Use Only

Antibiotic

**ACTIVE INGREDIENTS:**

Benzylpenicillin benzathine BP	150,000 I.U./mL
Benzylpenicillin procaine BP	150,000 I.U./mL

**PRESERVATIVES:**

Methylparaben	0.9 mg/mL
Propylparaben	0.1 mg/mL

**NON-MEDICINAL INGREDIENTS:**

Sodium citrate dihydrate	10.87 mg/mL
Potassium dihydrogen phosphate	0.3 mg/mL
Dimethicone	0.16 mg/mL
Sorbitan monopalmitate	0.5 mg/mL
Polysorbate 40	0.5 mg/mL
Sodium chloride	3.0 mg/mL
Lecithin	5.0 mg/mL
Water for injection	Ad 1 mL

**INDICATIONS:**

BENZAPRO is indicated as an aid in the treatment of local and systemic infections caused by bacteria susceptible to penicillin including metritis, strangles and tetanus in **horses**; blackleg, foot rot, metritis and shipping fever complex in **cattle**; clostridial diseases and foot rot in **sheep**; erysipelas, mastitis and metritis in **swine** and metritis, respiratory infections and wound infections in **dogs** and **cats**, where a prolongation of the therapeutic period from a single injection is required.

**DOSAGE AND ADMINISTRATION:**

Shake well before using. Administer by deep intramuscular injection only. The usual aseptic precautions should be followed. Accurate determination of body weight is important to ensure proper dosage. Please read package insert before use.

**WARNINGS:**

Treated animals must not be slaughtered for use in food for at least 14 days after the latest treatment with this drug. Do not use in lactating dairy cattle. Do not use in horses intended for food. Penicillins can cause allergic reactions in sensitized individuals. When handling the product, avoid inhalation, oral exposure and direct contact with skin or eyes. Keep out of reach of children.

Benzapro label – English - **Front label**

Net contents: 100 mL

Lot no.:

Exp.:



**Bio Agri Mix LP**  
PO Box 399  
Mitchell, ON  
N0K 1N0



CONTAINS  
**ANTIMICROBIAL**  
USE RESPONSIBLY  
CONTIENT UN  
**ANTIMICROBIEN**  
UTILISEZ DE FAÇON RESPONSABLE

Pr **BENZAPRO**

**Sterile suspension of benzylpenicillin benzathine and benzylpenicillin procaine**

**Veterinary Use Only**

**Antibiotic**

**ACTIVE INGREDIENTS:**

Benzylpenicillin benzathine BP	150,000 I.U./mL
Benzylpenicillin procaine BP	150,000 I.U./mL

**PRESERVATIVES:**

Methylparaben	0.9 mg/mL
Propylparaben	0.1 mg/mL

**NON-MEDICINAL INGREDIENTS:**

Sodium citrate dihydrate	10.87 mg/mL
Potassium dihydrogen phosphate	0.3 mg/mL
Dimethicone	0.16 mg/mL
Sorbitan monopalmitate	0.5 mg/mL
Polysorbate 40	0.5 mg/mL
Sodium chloride	3.0 mg/mL
Lecithin	5.0 mg/mL
Water for injection	Ad 1 mL

**INDICATIONS:**

**BENZAPRO** is indicated as an aid in the treatment of local and systemic infections caused by bacteria susceptible to penicillin including metritis, strangles and tetanus in horses; blackleg, foot rot, metritis and shipping fever complex in cattle; clostridial diseases and foot rot in sheep; erysipelas, mastitis and metritis in swine and metritis, respiratory infections and wound infections in dogs and cats, where a prolongation of the therapeutic period from a single injection is required.

**DOSAGE AND ADMINISTRATION:**

Shake well before using. Administer by deep intramuscular injection only. The usual aseptic precautions should be followed. Accurate determination of body weight is important to ensure proper dosage.

The following dosage rates are recommended:

*Horses and Cattle:*

6000 I.U. each of Benzylpenicillin procaine and Benzylpenicillin benzathine/kg of body weight (corresponds to 1 mL/25 kg, 10 mL/250 kg, 16 mL/400 kg, or 20 mL/500 kg of body weight)

*Sheep and Swine:*

7500 I.U. each of Benzylpenicillin procaine and Benzylpenicillin benzathine /kg of body weight (corresponds to 1 mL/20 kg, 2 mL/40 kg, or 3 mL/60 kg of body weight)

*Dogs and Cats:*

Benzapro label – English – **Package Insert**

15,000 I.U. each of Benzylpenicillin procaine and Benzylpenicillin benzathine /kg of body weight  
(corresponds to 0.5 mL/5 kg, 1 mL/10 kg, or 2 mL/20 kg of body weight)

**Note:**

Repeat the dose in 3 to 5 days if required. If no improvement of the condition is noted within 36-48 hours, the treatment and/or diagnosis should be re-evaluated.

**CAUTIONS:**

BENZAPRO should not be administered to animals known to be allergic to penicillin. In case of an anaphylactic reaction administer epinephrine immediately. In suckling and fattening pigs, administration of benzylpenicillin procaine will occasionally cause a transient pyrexia, vomiting, shivering and incoordination. Abortions have also been reported in pregnant sows and gilts.

**WARNINGS:**

Treated animals must not be slaughtered for use in food for at least 14 days after the latest treatment with this drug. Do not use in lactating dairy cattle. Do not use in horses intended for food. Penicillins can cause allergic reactions in sensitized individuals. When handling the product, avoid inhalation, oral exposure and direct contact with skin or eyes. Keep out of reach of children.

**STORAGE:**

Store below 15° C and keep from freezing.

Net contents: 100 mL



**Bio Agri Mix LP**  
PO Box 399  
Mitchell, ON  
N0K 1N0



**CONTAINS  
ANTIMICROBIAL  
USE RESPONSIBLY  
CONTIENT UN  
ANTIMICROBIEN  
UTILISEZ DE FAÇON RESPONSABLE**

**Pr BENZAPRO**

Sterile suspension of benzylpenicillin benzathine and benzylpenicillin procaine  
Veterinary Use Only  
Antibiotic

**ACTIVE INGREDIENTS:**

Benzylpenicillin benzathine BP	150,000 I.U./mL
Benzylpenicillin procaine BP	150,000 I.U./mL

**PRESERVATIVES:**

Methylparaben	0.9 mg/mL
Propylparaben	0.1 mg/mL

**NON-MEDICINAL INGREDIENTS:**

Sodium citrate dihydrate	10.87 mg/mL
Potassium dihydrogen phosphate	0.3 mg/mL
Dimethicone	0.16 mg/mL
Sorbitan monopalmitate	0.5 mg/mL
Polysorbate 40	0.5 mg/mL
Sodium chloride	3.0 mg/mL
Lecithin	5.0 mg/mL
Water for injection	Ad 1 mL

**INDICATIONS:**

BENZAPRO is indicated as an aid in the treatment of local and systemic infections caused by bacteria susceptible to penicillin including metritis, strangles and tetanus in **horses**; blackleg, foot rot, metritis and shipping fever complex in **cattle**; clostridial diseases and foot rot in **sheep**; erysipelas, mastitis and metritis in **swine** and metritis, respiratory infections and wound infections in **dogs** and **cats**, where a prolongation of the therapeutic period from a single injection is required.

**DOSAGE AND ADMINISTRATION:**

Shake well before using. Administer by deep intramuscular injection only. The usual aseptic precautions should be followed. Accurate determination of body weight is important to ensure proper dosage. Please read package insert before use.

**WARNINGS:**

Treated animals must not be slaughtered for use in food for at least 14 days after the latest treatment with this drug. Do not use in lactating dairy cattle. Do not use in horses intended for food. Penicillins can

Benzapro 250 label – English - **Front label**

cause allergic reactions in sensitized individuals. When handling the product, avoid inhalation, oral exposure and direct contact with skin or eyes. Keep out of reach of children.

Net contents: 250 mL

Lot no.:

Exp.:



**Bio Agri Mix LP**  
PO Box 399  
Mitchell, ON  
N0K 1N0



**CONTAINS  
ANTIMICROBIAL  
USE RESPONSIBLY**  
**CONTIENT UN  
ANTIMICROBIEN  
UTILISEZ DE FAÇON RESPONSABLE**

**Pr BENZAPRO**

**Sterile suspension of benzylpenicillin benzathine and benzylpenicillin procaine  
Veterinary Use Only  
Antibiotic**

**ACTIVE INGREDIENTS:**

Benzylpenicillin benzathine BP	150,000 I.U./mL
Benzylpenicillin procaine BP	150,000 I.U./mL

**PRESERVATIVES:**

Methylparaben	0.9 mg/mL
Propylparaben	0.1 mg/mL

**NON-MEDICINAL INGREDIENTS:**

Sodium citrate dihydrate	10.87 mg/mL
Potassium dihydrogen phosphate	0.3 mg/mL
Dimethicone	0.16 mg/mL
Sorbitan monopalmitate	0.5 mg/mL
Polysorbate 40	0.5 mg/mL
Sodium chloride	3.0 mg/mL
Lecithin	5.0 mg/mL
Water for injection	Ad 1 mL

**INDICATIONS:**

**BENZAPRO** is indicated as an aid in the treatment of local and systemic infections caused by bacteria susceptible to penicillin including metritis, strangles and tetanus in horses; blackleg, foot rot, metritis and shipping fever complex in cattle; clostridial diseases and foot rot in sheep; erysipelas, mastitis and metritis in swine and metritis, respiratory infections and wound infections in dogs and cats, where a prolongation of the therapeutic period from a single injection is required.

**DOSAGE AND ADMINISTRATION:**

Shake well before using. Administer by deep intramuscular injection only. The usual aseptic precautions should be followed. Accurate determination of body weight is important to ensure proper dosage. The following dosage rates are recommended:

*Horses and Cattle:*

6000 I.U. each of Benzylpenicillin procaine and Benzylpenicillin benzathine /kg of body weight (corresponds to 1 mL/25 kg, 10 mL/250 kg, 16 mL/400 kg, or 20 mL/500 kg of body weight)

*Sheep and Swine:*

7500 I.U. each of Benzylpenicillin procaine and Benzylpenicillin benzathine /kg of body weight (corresponds to 1 mL/20 kg, 2 mL/40 kg, or 3 mL/60 kg of body weight)

Benzapro label – English – **Package Insert**

*Dogs and Cats:*

15,000 I.U. each of Benzylpenicillin procaine and Benzylpenicillin benzathine /kg of body weight (corresponds to 0.5 mL/5 kg, 1 mL/10 kg, or 2 mL/20 kg of body weight)

Note:

Repeat the dose in 3 to 5 days if required. If no improvement of the condition is noted within 36-48 hours, the treatment and/or diagnosis should be re-evaluated.

**CAUTIONS:**

BENZAPRO should not be administered to animals known to be allergic to penicillin. In case of an anaphylactic reaction administer epinephrine immediately. In suckling and fattening pigs, administration of benzylpenicillin procaine will occasionally cause a transient pyrexia, vomiting, shivering and incoordination. Abortions have also been reported in pregnant sows and gilts.

**WARNINGS:**

Treated animals must not be slaughtered for use in food for at least 14 days after the latest treatment with this drug. Do not use in lactating dairy cattle. Do not use in horses intended for food. Penicillins can cause allergic reactions in sensitized individuals. When handling the product, avoid inhalation, oral exposure and direct contact with skin or eyes. Keep out of reach of children.

**STORAGE:**

Store below 15° C and keep from freezing.

Net contents: 250 mL



**Bio Agri Mix LP**  
PO Box 399  
Mitchell, ON  
N0K 1N0



CONTAINS  
**ANTIMICROBIAL**  
USE RESPONSIBLY  
CONTIENT UN  
**ANTIMICROBIEN**  
UTILISEZ DE FAÇON RESPONSABLE



**Pr BENZAPRO**

Sterile suspension of benzylpenicillin benzathine and benzylpenicillin procaine  
Veterinary Use Only  
Antibiotic

**ACTIVE INGREDIENTS:**

Benzylpenicillin benzathine BP	150,000 I.U./mL
Benzylpenicillin procaine BP	150,000 I.U./mL

**PRESERVATIVES:**

Methylparaben	0.9 mg/mL
Propylparaben	0.1 mg/mL

**NON-MEDICINAL INGREDIENTS:**

Sodium citrate dihydrate	10.87 mg/mL
Potassium dihydrogen phosphate	0.3 mg/mL
Dimethicone	0.16 mg/mL
Sorbitan monopalmitate	0.5 mg/mL
Polysorbate 40	0.5 mg/mL
Sodium chloride	3.0 mg/mL
Lecithin	5.0 mg/mL
Water for injection	Ad 1 mL

**INDICATIONS:**

BENZAPRO is indicated as an aid in the treatment of local and systemic infections caused by bacteria susceptible to penicillin including metritis, strangles and tetanus in **horses**; blackleg, foot rot, metritis and shipping fever complex in **cattle**; clostridial diseases and foot rot in **sheep**; erysipelas, mastitis and metritis in **swine** and metritis, respiratory infections and wound infections in **dogs** and **cats**, where a prolongation of the therapeutic period from a single injection is required.

**DOSAGE AND ADMINISTRATION:**

Shake well before using. Administer by deep intramuscular injection only. The usual aseptic precautions should be followed. Accurate determination of body weight is important to ensure proper dosage. Please read package insert before use.

**WARNINGS:**

Treated animals must not be slaughtered for use in food for at least 14 days after the latest treatment with this drug. Do not use in lactating dairy cattle. Do not use in horses intended for food. Penicillins can

Benzapro 500 mL label – English - **Front label**

cause allergic reactions in sensitized individuals. When handling the product, avoid inhalation, oral exposure and direct contact with skin or eyes. Keep out of reach of children.

Net contents: 500 mL

Lot no.:

Exp.:



**Bio Agri Mix LP**  
PO Box 399  
Mitchell, ON  
N0K 1N0



CONTAINS  
**ANTIMICROBIAL**  
USE RESPONSIBLY  
CONTIENT UN  
**ANTIMICROBIEN**  
UTILISEZ DE FAÇON RESPONSABLE

**Pr**  
**BENZAPRO**

**Sterile suspension of benzylpenicillin benzathine and benzylpenicillin procaine**  
**Veterinary Use Only**  
**Antibiotic**

**ACTIVE INGREDIENTS:**

Benzylpenicillin benzathine BP	150,000 I.U./mL
Benzylpenicillin procaine BP	150,000 I.U./mL

**PRESERVATIVES:**

Methylparaben	0.9 mg/mL
Propylparaben	0.1 mg/mL

**NON-MEDICINAL INGREDIENTS:**

Sodium citrate dihydrate	10.87 mg/mL
Potassium dihydrogen phosphate	0.3 mg/mL
Dimethicone	0.16 mg/mL
Sorbitan monopalmitate	0.5 mg/mL
Polysorbate 40	0.5 mg/mL
Sodium chloride	3.0 mg/mL
Lecithin	5.0 mg/mL
Water for injection	Ad 1 mL

**INDICATIONS:**

**BENZAPRO** is indicated as an aid in the treatment of local and systemic infections caused by bacteria susceptible to penicillin including metritis, strangles and tetanus in horses; blackleg, foot rot, metritis and shipping fever complex in cattle; clostridial diseases and foot rot in sheep; erysipelas, mastitis and metritis in swine and metritis, respiratory infections and wound infections in dogs and cats, where a prolongation of the therapeutic period from a single injection is required.

**DOSAGE AND ADMINISTRATION:**

Shake well before using. Administer by deep intramuscular injection only. The usual aseptic precautions should be followed. Accurate determination of body weight is important to ensure proper dosage.

The following dosage rates are recommended:

*Horses and Cattle:*

6000 I.U. each of Benzylpenicillin procaine and Benzylpenicillin benzathine /kg of body weight (corresponds to 1 mL/25 kg, 10 mL/250 kg, 16 mL/400 kg, or 20 mL/500 kg of body weight)

*Sheep and Swine:*

7500 I.U. each of Benzylpenicillin procaine and Benzylpenicillin benzathine /kg of body weight (corresponds to 1 mL/20 kg, 2 mL/40 kg, or 3 mL/60 kg of body weight)

*Dogs and Cats:*

## Benzapro 500 mL label – English – **Package Insert**

15,000 I.U. each of Benzylpenicillin procaine and Benzylpenicillin benzathine /kg of body weight (corresponds to 0.5 mL/5 kg, 1 mL/10 kg, or 2 mL/20 kg of body weight)

### Note:

Repeat the dose in 3 to 5 days if required. If no improvement of the condition is noted within 36-48 hours, the treatment and/or diagnosis should be re-evaluated.

### **CAUTIONS:**

BENZAPRO should not be administered to animals known to be allergic to penicillin. In case of an anaphylactic reaction administer epinephrine immediately. In suckling and fattening pigs, administration of benzylpenicillin procaine will occasionally cause a transient pyrexia, vomiting, shivering and incoordination. Abortions have also been reported in pregnant sows and gilts.

### **WARNINGS:**

Treated animals must not be slaughtered for use in food for at least 14 days after the latest treatment with this drug. Do not use in lactating dairy cattle. Do not use in horses intended for food. Penicillins can cause allergic reactions in sensitized individuals. When handling the product, avoid inhalation, oral exposure and direct contact with skin or eyes. Keep out of reach of children.

### **STORAGE:**

Store below 15° C and keep from freezing.

Net contents: 500 mL



**Bio Agri Mix LP**  
PO Box 399  
Mitchell, ON  
N0K 1N0



CONTAINS  
**ANTIMICROBIAL**  
USE RESPONSIBLY  
CONTIENT UN  
**ANTIMICROBIEN**  
UTILISEZ DE FAÇON RESPONSABLE