

DIN 00780774

Pr OXYSOL-110

Oxytetracycline hydrochloride medicated premix

FOR VETERINARY USE ONLY
ANTIBIOTIC

ACTIVE INGREDIENT: Oxytetracycline hydrochloride 110 g/kg

INDICATIONS

BEEF CATTLE: As an aid in reducing the incidence of bloat in young cattle on pasture or in feedlots.

CALVES: As an aid in the prevention of bacterial enteritis.

LAMBS: As an aid in the reduction of bacterial enteritis in creep-fed suckling lambs. As an aid in the reduction of losses due to enterotoxaemia in feedlot lambs.

SWINE: As an aid in the treatment or prevention of bacterial enteritis. As an aid in reducing the incidence of abortion in swine caused by leptospirosis.

TURKEYS: As an aid in the prevention of infectious sinusitis of turkeys. As an aid in the prevention or treatment of infectious synovitis of turkeys.

DIRECTIONS FOR USE

To reduce the development of antimicrobial resistance and maintain effectiveness, use this antibiotic prudently and for the shortest duration required to achieve the desired clinical outcome.

Must be thoroughly mixed in feeds. Do not feed undiluted.

SPECIES	DISEASE OR CONDITION	INDICATIONS	kg of OXYSOL-110 PER TONNE COMPLETE FEED	Concentration in % of Oxytetracycline Hydrochloride in complete feed	FEEDING DIRECTIONS
SWINE	BACTERIAL ENTERITIS (scours)	As an aid in the prevention of bacterial enteritis.	2.0	0.022%	Feed as the sole ration up to six weeks of age.
			0.5	0.0055%	Feed as the sole ration from six weeks of age up to 27-32 kg bodyweight.
	As an aid in the treatment of bacterial enteritis.	2.0	0.022%	Feed as the sole ration following the appearance of clinical signs and until three days after clinical signs disappear.-- for pigs up to six weeks of age.	
		1.0	0.011%	As above -for pigs greater than six weeks of age. NOTE: Pigs off feed should be given individual treatment.	
LEPTOSPIROSIS (abortion in swine)	As an aid in reducing the incidence of abortion caused by leptospirosis. NOTE: Good sanitation procedures must be practiced to prevent spread of infections.	5.0	0.055%	When abortion in swine has been diagnosed as due to leptospirosis, feed at once as the sole ration to pregnant and affected sows for a two-week period.	
BEEF CATTLE	BLOAT	As an aid in reducing the incidence of bloat in young cattle on pasture and in feed lots.	Feed continuously. Level of oxytetracycline hydrochloride in the complete feed is to be such that each animal will receive 75 mg per day. 55 g of oxytetracycline hydrochloride (0.5 kg of OXYSOL-110) per tonne complete feed when fed at the rate of 1.36 kg per head per day meets the 75 mg level stated above.		
CALVES	BACTERIAL ENTERITIS (scours)	As an aid in the prevention of bacterial enteritis.	0.5	0.0055%	Feed continuously during the period of early growth (up to 136 kg bodyweight).
TURKEYS	INFECTIOUS SINUSITIS	As an aid in the prevention of infectious sinusitis of turkeys.	1.0	0.011%	Feed as the sole ration for as long as the birds are expected to be exposed to the infection.
	INFECTIOUS SYNOVITIS	As an aid in the prevention of infectious synovitis of turkeys.	1.0	0.011%	Feed as the sole ration for as long as the birds are expected to be exposed to the infection.
		As an aid in the treatment of infectious synovitis of turkeys.	2.0	0.022%	Feed as the sole ration when clinical signs appear and continue until at least two weeks after the clinical signs disappear.
LAMBS	BACTERIAL ENTERITIS (scours)	As an aid in the reduction of bacterial enteritis in creep-fed suckling lambs.	1.0	0.011%	Feed as the sole creep feed during the suckling period.
	ENTEROTOXEMIA (overeating disease)	As an aid in the reduction of losses due to enterotoxemia in feedlot lambs.	0.20	0.0022%	Feed as the sole ration while lambs are in the feedlot

WARNINGS: Treated animals must not be slaughtered for use in food for at least 7 days for swine and turkeys, 5 days for cattle and calves, and 4 days for lambs after the latest treatment with this drug. Do not use in lactating dairy cattle. When handling the product, avoid inhalation, oral exposure and direct contact with skin or eyes. Keep out of reach of children.

STORAGE: Store in a cool dry place below 23°C.

NET CONTENTS: 25 kg

Lot:

Exp.:

Manufactured by:

Bio Agri Mix LP

PO Box 399
Mitchell, ON
N0K 1N0



**CONTAINS
ANTIMICROBIAL
USE RESPONSIBLY**

**CONTIENT UN
ANTIMICROBIEN
UTILISEZ DE FAÇON RESPONSABLE**