

Pr **TIAMULIN SOLUBLE POWDER**
(Tiamulin fumarate USP)

VETERINARY USE ONLY
ANTIBIOTIC

ACTIVE INGREDIENT

Each 152.5 g jar contains:

Tiamulin (as tiamulin fumarate USP).....55.7 g

NET CONTENTS 152.5 g

INDICATION

For the treatment of swine dysentery caused by *Serpulina hyodysenteriae*.

DIRECTIONS FOR USE

To reduce the development of antimicrobial resistance and maintain effectiveness, use this antibiotic prudently.

Add the contents of this jar to 1136 litres (250 Imp. Gal.) of drinking water (provides 49 mg antibiotic activity per litre). Administer at this dosage as the sole source of drinking water for 5 days.

For automatic medicators set to deliver 7.8 mL of stock solution per litre (1 oz. per US gallon), prepare stock solution by dissolving the contents of this jar in 8.87 litres of water.

Preparation of medicated solutions -

Pour the contents a Tiamulin Soluble Powder jar into a suitable clean container. Do not use old galvanized or corroded containers as they may compromise the efficacy of this product.

Add approximately 4 L of drinking water.

Stir thoroughly until the powder is dissolved.

Add this solution to the required quantity of drinking water or make up to the correct volume for the stock solution for automatic medicators.

Notes -

Prepare fresh medicated water every day.

Water medicated with Tiamulin Soluble Powder should be the only source of drinking water during the treatment period.

During treatment, swine should be housed under conditions of adequate space and sanitation.

CONTRAINDICATION

Swine being medicated with Tiamulin Soluble Powder should not have access to or be treated with incompatible polyether ionophores (e.g. monensin, salinomycin, narasin).

Tiamulin Soluble Powder
English Product Insert

CAUTIONS

For use in the drinking water of swine only. This drug should not be administered to sows during 4 weeks after service.

PRECAUTIONS

If no response is obtained within 5 days, re-establish the diagnosis. Failure of response may be related to the presence of non-susceptible organisms or other complicating disease conditions.

WARNINGS

Treated animals must not be slaughtered for use in food for at least 3 days after the latest with this drug. Direct contact with skin or mucous membranes during mixing may cause irritation. When handling the product, avoid inhalation, oral exposure and direct contact with skin or eyes. Keep out of reach of children

ADVERSE REACTIONS

Overdoses of tiamulin have sometimes produced transitory salivation, vomiting and an apparent calming effect on pigs.

In rare cases, redness of the skin, primarily over the ham and underline, has been observed during medication.

If signs of toxicity or redness of the skin occurs, promptly discontinue use of medicated water and provide fresh clean water for drinking.

STORAGE

Store in a dry place between 15 and 30°C.



Bio Agri Mix LP
PO Box 399
Mitchell, ON
N0K 1N0



**CONTAINS
ANTIMICROBIAL
USE RESPONSIBLY**
**CONTIENT UN
ANTIMICROBIEN
UTILISEZ DE FAÇON RESPONSABLE**

Rev: Oct 30/17

DIN 02324407

PrTIAMULIN SOLUBLE POWDER
(Tiamulin fumarate USP)

VETERINARY USE ONLY
ANTIBIOTIC

ACTIVE INGREDIENT

Each 152.5 g jar contains:

Tiamulin (as tiamulin fumarate USP).....55.7 g

NET CONTENTS 152.5 g

INDICATION

Swine - For the treatment of swine dysentery caused by *Serpulina hyodysenteriae*.

DIRECTIONS FOR USE

To reduce the development of antimicrobial resistance and maintain effectiveness, use this antibiotic prudently.

Swine - 1 jar (152.5 g) per 1136 litres (250 Imp. Gal.) of drinking water continuously for 5 days.

See package insert for detailed instructions.

CONTRAINDICATION

Swine being medicated with Tiamulin Soluble Powder should not have access to or be treated with tiamulin incompatible polyether ionophores (e.g. monensin, salinomycin, narasin).

CAUTIONS

For use in the drinking water of swine only. This drug should not be administered to sows during 4 weeks after service.

WARNINGS

Treated animals must not be slaughtered for use in food for at least 3 days after the latest treatment with this drug.

Direct contact with skin or mucous membranes during mixing may cause irritation. When handling the product, avoid inhalation, oral exposure and direct contact with skin or eyes.

Keep out of reach of children.

Tiamulin Soluble Powder
English Product Label Panel

STORAGE

Store in a dry place, between 15 and 30°C.



Bio Agri Mix LP
PO Box 399
Mitchell, ON
N0K 1N0



**CONTAINS
ANTIMICROBIAL
USE RESPONSIBLY**
**CONTIENT UN
ANTIMICROBIEN
UTILISEZ DE FAÇON RESPONSABLE**

Rev: Oct 30/17

Tiamulin Soluble Powder
English Outer Label Panel

Lot No.

Exp.