DIN 02209098

PrLINCOMYCIN SPECTINOMYCIN 4.4% G PREMIX

lincomycin hydrochloride and spectinomycin hydrochloride granular premix

FOR VETERINARY USE ONLY ANTIBIOTIC

ACTIVE INGREDIENTS:

Lincomycin base (as lincomycin hydrochloride)

22 g/kg

Spectinomycin base (as spectinomycin hydrochloride)

22 g/kg

INDICATIONS:

For the prevention of swine dysentery (vibrionic dysentery, bloody scours) in growing swine up to 57 kg of bodyweight.

DIRECTIONS FOR USE:

To reduce the development of antimicrobial resistance and maintain effectiveness, use this antibiotic prudently and for the shortest duration required to achieve the desired clinical outcome.

For the prevention of swine dysentery, complete feeds containing 44 grams of antibiotic activity per tonne should be fed to growing swine up to a weight of 57 kg.

LINCOMYCIN SPECTINOMYCIN 4.4% G PREMIX should be mixed with feed before use.

Mixing Directions:

For making medicated complete feed. Add 1 kg of LINCOMYCIN SPECTINOMYCIN 4.4% G PREMIX per tonne of complete feed. The resulting medicated complete feed will contain 44 grams of antibiotic activity per metric tonne. This should be fed as the sole ration to growing swine up to a weight of 57 kg.

For supplements designed to be incorporated into complete feed. Add the amount of LINCOMYCIN SPECTINOMYCIN 4.4% G PREMIX listed in the following table to the supplement to prepare a medicated supplement.

Supplement rate per tonne* of complete feed	Amount of LINCOMYCIN SPECTINOMYCIN 4.4% G PREMIX to be added to supplement to make medicated supplement
25 kg supplement and 975 kg other ingredients	40 kg LINCOMYCIN SPECTINOMYCIN 4.4% G PREMIX per tonne of supplement (1760 grams antibiotic activity/tonne)
50 kg supplement and 950 kg other ingredients	20 kg LINCOMYCIN SPECTINOMYCIN 4.4% G PREMIX per tonne of supplement (880 grams antibiotic activity/tonne)
100 kg supplement and 900 kg other ingredients	10 kg LINCOMYCIN SPECTINOMYCIN 4.4% G PREMIX per tonne of supplement

Rev: BAM Sep-20-17

Lincomycin Spectinomycin 4.4% Premix English Label

	(440 grams antibiotic activity/tonne)
150 kg supplement and 850 kg other	7.5 kg LINCOMYCIN SPECTINOMYCIN 4.4% G
ingredients	PREMIX per tonne of supplement
	(330 grams antibiotic activity/tonne)
200 kg supplement and 800 kg other	5 kg LINCOMYCIN SPECTINOMYCIN 4.4% G PREMIX
ingredients.	per tonne of supplement
A	(220 grams antibiotic activity/tonne)
225 kg supplement and 775 kg other	4.5 kg LINCOMYCIN SPECTINOMYCIN 4.4% G
ingredients.	PREMIX per tonne of supplement
1980 - 19	(200 grams antibiotic activity/tonne)
250 kg supplement and 750 kg other	4 kg LINCOMYCIN SPECTINOMYCIN 4.4% G PREMIX
ingredients	per tonne of supplement
	(175 grams antibiotic activity/tonne)

^{*} The resultant medicated complete feed containing 44 grams of antibiotic activity per metric tonne should be fed as the sole ration to growing swine up to a weight of 57 kg.

WARNINGS:

- 1. Treated swine must not be slaughtered for use in food for at least 24 hours after the latest treatment with this drug.
- 2. When handling the product, avoid inhalation, oral exposure and direct contact with skin or eyes.
- 3. Keep out of reach of children.

CAUTIONS:

- 1. Do not allow rabbits, hamsters, guinea pigs, horses, dairy cattle or other ruminants access to feeds containing lincomycin. Ingestion by these species may result in severe gastrointestinal effects or metabolic disorders (e.g. Ketosis in dairy cattle).
- 2. Good manufacturing practices should be observed in preparing feeds containing this premix. This includes appropriate clean-out procedures to avoid cross contamination.
- 3. Only for use in growing swine up to 57 kg bodyweight.
- 4. Treated pigs may develop some stool softening or mild diarrhea within 36 hours of treatment initiation. Signs of mild irritation and swelling of the anus and/or vulva may be associated with the stool changes. These adverse reactions are transient and self-correcting by the fifth to eighth day of treatment. Should these reactions be more severe than described above discontinue the use of the drug.
- 5. Feed only to swine intended for slaughter.

STORAGE:

Store in a cool dry place below 30°C.

Net contents 25 kg.

Manufactured by

Bio Agri Mix LP

PO Box 399 Mitchell, ON N0K 1N0



Lot: Exp:

Rev: BAM Sep-20-17