

PRODUCT MONOGRAPH

^{Pr}**SANDOZ MOXIFLOXACIN**

Moxifloxacin Ophthalmic Solution

0.5% w/v (as moxifloxacin hydrochloride)

Sterile

Antibacterial (ophthalmic)

Sandoz Canada Inc.
145 Jules-Léger
Boucherville, Quebec
J4B 7K8

Date of Preparation:
February 2, 2018

Control number: 213238

Table of Contents

PART I: HEALTH PROFESSIONAL INFORMATION.....	3
SUMMARY PRODUCT INFORMATION	3
INDICATIONS AND CLINICAL USE.....	3
CONTRAINDICATIONS	4
WARNINGS AND PRECAUTIONS	4
ADVERSE REACTIONS	6
DRUG INTERACTIONS.....	7
DOSAGE AND ADMINISTRATION.....	7
OVERDOSAGE	8
ACTION AND CLINICAL PHARMACOLOGY	8
STORAGE AND STABILITY	9
DOSAGE FORMS, COMPOSITION AND PACKAGING	9
PART II: SCIENTIFIC INFORMATION.....	10
PHARMACEUTICAL INFORMATION	10
CLINICAL TRIALS	11
DETAILED PHARMACOLOGY.....	11
MICROBIOLOGY	14
TOXICOLOGY.....	16
REFERENCES.....	20
PATIENT MEDICATION INFORMATION.....	22

Pr SANDOZ MOXIFLOXACIN

moxifloxacin ophthalmic solution

PART I: HEALTH PROFESSIONAL INFORMATION

SUMMARY PRODUCT INFORMATION

Route of Administration	Dosage Form / Strength	All Nonmedicinal Ingredients
Ophthalmic (topical)	Solution/ 0.5% w/v moxifloxacin (as moxifloxacin hydrochloride)	Boric acid, purified water and sodium chloride. May also contain hydrochloric acid/sodium hydroxide to adjust pH.

INDICATIONS AND CLINICAL USE

Sandoz Moxifloxacin (moxifloxacin ophthalmic solution) is indicated for the treatment of patients 1 year of age and older with bacterial conjunctivitis caused by susceptible strains of the following organisms:

Aerobic, Gram-Positive

Staphylococcus aureus
Staphylococcus epidermidis
Staphylococcus haemolyticus
Staphylococcus hominis
Streptococcus pneumoniae
Streptococcus viridans group

Aerobic, Gram-Negative

Acinetobacter species
Haemophilus influenza

To reduce the development of the drug-resistant bacteria and maintain the effectiveness of Sandoz Moxifloxacin and other antibacterial drugs, Sandoz Moxifloxacin should be used only to treat infections that are proven or strongly suspected to be caused by susceptible bacteria. When culture and susceptibility information are available, they should be considered in selecting or modifying antibacterial therapy. In the absence of such data, local epidemiology and susceptibility patterns may contribute to the empiric selection of therapy.

Geriatrics (> 65 years of age):

No overall differences in safety and effectiveness have been observed between elderly and other adult patients.

Pediatrics (< 1 year of age):

The safety and efficacy of Sandoz Moxifloxacin in patients less than one year of age have not been established.

CONTRAINDICATIONS

Sandoz Moxifloxacin is contraindicated in patients with:

- Hypersensitivity to moxifloxacin or to any ingredient in the formulation or component of the container (for a complete listing, see the Dosage Forms, Composition and Packaging section of the Product Monograph).
- Hypersensitivity to other quinolones.

WARNINGS AND PRECAUTIONS**General****For ocular use only.**

Sandoz Moxifloxacin is not for injection into the eye.

Sandoz Moxifloxacin should not be injected subconjunctivally, nor should it be introduced directly into the anterior chamber of the eye.

Prescribing Sandoz Moxifloxacin in the absence of a proven or strongly suspected bacterial infection is unlikely to provide benefit to the patient and risks the development of drug-resistant bacteria.

In patients receiving systemically administered quinolones, serious and occasionally fatal hypersensitivity (anaphylactic) reactions have been reported, some following the first dose. Some reactions were accompanied by cardiovascular collapse, loss of consciousness, angioedema (including laryngeal, pharyngeal or facial edema), airway obstruction, dyspnea, urticaria, and itching. If an allergic reaction to moxifloxacin occurs, discontinue use of Sandoz Moxifloxacin. Serious acute hypersensitivity reactions may require immediate emergency treatment. Oxygen and airway management should be administered as clinically indicated.

Serious and sometimes fatal events, some due to hypersensitivity and some due to uncertain etiology, have been reported in patients receiving therapy with all oral antibiotics. These events may be severe and generally occur following the administration of multiple doses. Clinical manifestations may include one or more of the following: fever, rash or severe dermatologic reactions (e.g. toxic epidermal necrolysis, Stevens-Johnson Syndrome), vasculitis, arthralgia, myalgia, serum sickness, allergic pneumonitis, interstitial nephritis, acute renal insufficiency or failure, hepatitis, jaundice, acute hepatic necrosis or failure, anemia including hemolytic and aplastic, thrombocytopenia including thrombotic thrombocytopenic purpura, leukopenia, agranulocytosis, pancytopenia, and/or other hematologic abnormalities.

As with other anti-infectives, prolonged use may result in overgrowth of non-susceptible organisms, including fungi. If superinfection occurs, discontinue use and institute alternative therapy. Whenever clinical judgment dictates, the patient should be examined with the aid of magnification, such as slit-lamp biomicroscopy, and, where appropriate, fluorescein staining.

Tendon inflammation and rupture may occur with systemic fluoroquinolone therapy including moxifloxacin, particularly in elderly patients and in those treated concurrently with corticosteroids. Treatment with moxifloxacin hydrochloride ophthalmic solution should be discontinued at the first sign of tendon inflammation.

Sandoz Moxifloxacin may cause temporary blurred vision or other visual disturbances, which may affect the ability to drive or use machines. If blurred vision occurs at application, the patient must wait until the vision clears before driving or using machinery.

Ophthalmologic

Patients with signs and symptoms of bacterial conjunctivitis should be advised not to wear contact lenses.

Sexual Function/Reproduction

There are no studies on the effect of ocular administration of moxifloxacin hydrochloride ophthalmic solution on fertility.

Special Populations

Pregnant Women: There are no adequate and well-controlled studies in pregnant women. Sandoz Moxifloxacin should only be used during pregnancy if the potential benefit justifies the potential risk to the fetus.

Moxifloxacin hydrochloride ophthalmic solution has not been studied in pregnant animals. Oral and IV studies in pregnant animals indicated that moxifloxacin is not teratogenic. Decreased fetal birth weights and slightly delayed fetal skeletal development was observed only at doses >4000 times the highest recommended total daily human ophthalmic dose (see TOXICOLOGY).

Nursing Women: Moxifloxacin is excreted in the breast milk of rats following oral and intravenous administration. Because of the potential for unknown effects from moxifloxacin in infants being nursed by mothers taking moxifloxacin hydrochloride, a decision should be made to either discontinue nursing or discontinue the administration of Sandoz Moxifloxacin, taking into account the importance of Sandoz Moxifloxacin therapy to the mother and the possible risk to the infant (see TOXICOLOGY).

Geriatrics (>65 years of age): No overall differences in safety and effectiveness have been observed between elderly and other adult patients.

Pediatrics (<1 years of age): The safety and efficacy of moxifloxacin hydrochloride ophthalmic solution in patients less than one year of age have not been established.

Pediatrics (<18 years of age): The effect of moxifloxacin hydrochloride ophthalmic solution on

weight bearing joints has not been assessed. Oral administration of some quinolones, including moxifloxacin, has been shown to cause arthropathy in immature Beagle dogs (see TOXICOLOGY). The significance of these findings to humans is unknown.

ADVERSE REACTIONS

Clinical Trial Adverse Drug Reactions

Because clinical trials are conducted under very specific conditions the adverse reaction rates observed in the clinical trials may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse drug reaction information from clinical trials is useful for identifying drug-related adverse events and for approximating rates.

In clinical trials involving 1068 subjects/patients, moxifloxacin hydrochloride was administered twice-daily for three days, three-times-daily for four to fourteen days and eight-times-daily for fourteen days. During treatment with moxifloxacin hydrochloride, 6.6% (71 out of 1068) subjects/patients experienced treatment-related adverse drug reactions and of these only two (0.2%) discontinued study participation. No serious ophthalmic or systemic adverse reactions related to moxifloxacin hydrochloride were reported.

The most frequently reported treatment-related adverse drug reactions were transient eye irritation (3.9%) (burning and/or stinging) and eye pruritus (1.1%).

Less Common Clinical Trial Adverse Drug Reactions (<1%)

Eye disorders: abnormal sensation in eye, conjunctival haemorrhage, conjunctivitis, corneal epithelium defect, eyelid oedema, eye pain, keratoconjunctivitis sicca, ocular discomfort, ocular hyperaemia, visual acuity reduced;

General disorders and administration site conditions: sensation of foreign body.

Investigations: alanine aminotransferase increased, corneal staining.

Nervous system disorders: dysgeusia, headache.

Respiratory, thoracic, and mediastinal disorders: pharyngolaryngeal pain.

Post-Market Adverse Drug Reactions

Adverse reactions identified from spontaneous reporting and subsequent clinical trials are listed below.

Blood and lymphatic system disorders: hemoglobin decreased.

Cardiac disorders: palpitations.

Eye disorders: anterior chamber cells, asthenopia, blepharitis, conjunctival edema, corneal deposits, corneal disorder, corneal infiltrates, dry eye, endophthalmitis, erythema of eyelid, eye discharge, eye irritation, eye swelling, keratitis, lacrimation increased, photophobia, punctate keratitis, ulcerative keratitis, vision blurred, visual acuity reduced.

Gastrointestinal disorders: nausea, vomiting.

Hepatobiliary disorders: gamma-glutamyltransferase increased.

Immune system disorders: hypersensitivity NOS.

Nervous system disorders: dizziness, paresthesia.

Respiratory, thoracic and mediastinal disorders: dyspnea, nasal discomfort

Skin and subcutaneous tissue disorders: erythema, pruritis, rash, urticaria.

DRUG INTERACTIONS

Overview

Specific drug interaction studies have not been conducted with moxifloxacin hydrochloride ophthalmic solution. There is limited information available on the concurrent use of moxifloxacin hydrochloride ophthalmic solution and other ophthalmic products.

Drug-Drug Interactions

Following oral administration, no clinically significant drug-drug interactions between theophylline, warfarin, digoxin, oral contraceptives or glyburide have been observed with moxifloxacin.

Theophylline, digoxin, probenecid, and ranitidine have been shown not to alter the pharmacokinetics of moxifloxacin. *In vitro* studies indicate that moxifloxacin does not inhibit CYP3A4, CYP2D6, CYP2C9, CYP2C19 or CYP1A2 indicating that moxifloxacin is unlikely to alter the pharmacokinetics of drugs metabolized by these cytochrome P450 isozymes.

Moxifloxacin can be chelated by polyvalent ions such as Mg^{++} , Al^{+++} , Fe^{++} and Zn^{++} .

Tendon inflammation and rupture may occur with systemic fluoroquinolone therapy including moxifloxacin, particularly in those treated concurrently with corticosteroids (see WARNINGS AND PRECAUTIONS, General).

Drug-food, drug-herb and drug-laboratory interactions have not been studied.

DOSAGE AND ADMINISTRATION

Recommended Dose

The recommended dosage regimen for patients one year of age and older is one drop in the affected eye(s) 3 times a day for 7 days.

Missed Dose

If a dose is missed, the missed dose should be administered as soon as possible. Treatment should then be continued with the next dose as planned.

Administration

To prevent contamination of the dropper tip and solution, care must be taken not to touch the eyelids, surrounding areas or other surfaces with the dropper tip of the bottle.

OVERDOSAGE

No information is available on overdose of moxifloxacin hydrochloride in humans. A topical overdose of Sandoz Moxifloxacin may be flushed from the eye(s) with warm tap water.

In an oral (gavage) monkey study, doses of moxifloxacin hydrochloride up to 15 mg/kg/day did not produce any toxicity. This dose is at least 10 times higher than the accidental ingestion of the contents of a 3 mL bottle of Sandoz Moxifloxacin by a 10 kg child.

No toxic effects are expected with an ocular overdose of the product, or in the event of accidental ingestion of the contents of one bottle.

For management of a suspected drug overdose, contact your regional Poison Control Centre.

ACTION AND CLINICAL PHARMACOLOGY

Mechanism of Action

Moxifloxacin is a synthetic fluoroquinolone antibacterial agent active *in vitro* against a broad spectrum of Gram-positive and Gram-negative ocular pathogens, atypical microorganisms and anaerobes.

The antibacterial action of moxifloxacin results from inhibition of topoisomerase II (DNA gyrase) and topoisomerase IV. DNA gyrase is an essential enzyme that is involved in the replication, transcription and repair of bacterial DNA. Topoisomerase IV is an enzyme known to play a key role in the partitioning of the chromosomal DNA during bacterial cell division (see MICROBIOLOGY).

Pharmacodynamics/Pharmacokinetics

Following topical ocular administration of moxifloxacin hydrochloride ophthalmic solution, moxifloxacin was absorbed into the systemic circulation. Plasma concentrations of moxifloxacin were measured in 21 male and female adult subjects who received bilateral topical ocular doses of moxifloxacin hydrochloride every 8 hours for a total of 13 doses. The mean steady-state C_{max} and AUC were 2.7 ng/mL and 41.9 ng.hr/mL, respectively. These systemic exposure values were at least 1600 and 1000 times lower than the mean C_{max} and AUC reported after therapeutic 400 mg oral doses of moxifloxacin. The plasma half-life of moxifloxacin was estimated to be 13 hours. Moxifloxacin is widely distributed in the body and is excreted in feces or urine either unchanged or as glucuronide or sulfate conjugates.

Tear film concentrations were studied in 31 healthy male and female adult volunteers who were administered 1 drop of moxifloxacin hydrochloride to both eyes every 8 hours for a total of 10 doses. Mean tear concentrations at 5 minutes following the first and last topical dose were 46.0 and 55.2 mcg/mL, respectively. Thereafter, they decline rapidly in a biphasic manner with the means ranging approximately 1 to 4 mcg/mL over the 1 to 8 hour sampling period. Pre-dose morning tear concentrations on Days 2 to 4 averaged over 4 mcg/mL. Studies conducted in animals indicate penetration into the conjunctiva and ocular tissues with prolonged binding to melanin.

Special Populations and Conditions

Geriatrics: The effects of age on the pharmacokinetic parameters of oral moxifloxacin have been studied. Plasma levels were 24 to 29% higher in the elderly than in young subjects. But, when normalized for body weight, the differences were minimized.

Gender: Gender differences in the steady-state C_{max} and AUC were seen. However, when adjusted for body weight, the differences were minimized and not clinically relevant (see DETAILED PHARMACOLOGY, Human Pharmacokinetics).

Race: Subgroup analysis by race (Caucasian, Asian) showed no meaningful differences in the mean steady-state pharmacokinetic parameters of moxifloxacin (see DETAILED PHARMACOLOGY, Human Pharmacokinetics).

Hepatic Insufficiency: The pharmacokinetic parameters of oral moxifloxacin were not significantly altered in patients with mild to moderate hepatic insufficiency (see DETAILED PHARMACOLOGY, Special Populations).

Renal Insufficiency: The pharmacokinetic parameters of oral moxifloxacin were not significantly altered by mild, moderate or severe renal impairment (see DETAILED PHARMACOLOGY, Special Populations).

STORAGE AND STABILITY

Store at 4° C - 25°C. Discard 28 days after opening. Keep out of the reach and sight of children.

DOSAGE FORMS, COMPOSITION AND PACKAGING

Each mL of Sandoz Moxifloxacin contains

Medicinal ingredient: Moxifloxacin 0.5% (5.45 mg moxifloxacin hydrochloride equivalent to 5 mg moxifloxacin base)

Preservative: None. Product is self-preserved.

Non-medicinal ingredients: boric acid, purified water and sodium chloride. May also contain hydrochloric acid/sodium hydroxide to adjust pH.

Sandoz Moxifloxacin is isotonic and formulated at pH 6.8 with an osmolality of approximately 290 mOsm/kg.

Sandoz Moxifloxacin is supplied as a 3 mL sterile ophthalmic solution in the DROP-TAINER[®] dispensing system consisting of a natural low density polyethylene bottle and dispensing plug and white polypropylene closure.

Tamper evidence is provided by a closure with an extended skirt that locks to the bottle finish on application and breaks away from the closure on opening. After cap is removed, if tamper evident snap collar is loose, remove before using product.

PART II: SCIENTIFIC INFORMATION

PHARMACEUTICAL INFORMATION


Drug Substance

Proper Name: Moxifloxacin Hydrochloride

Chemical Name: 1-Cyclopropyl-6-fluoro-1,4-dihydro-8-methoxy-7-[(4*aS*,7*aS*)-octahydro-6*H*-pyrrolol [3,4-*b*]pyridin-6-yl]-4-oxo-3-quinolinecarboxylic acid, monohydrochloride. Moxifloxacin differs from other quinolones in that it has a methoxy function at the 8 position, and an *S,S*- configured diazabicyclononyl ring moiety at the 7-position.

Molecular formula and molecular mass: $C_{21}H_{24}FN_3O_4 \cdot HCl$; 437.9 g/mol

Structural Formula:



Physicochemical properties: Slightly yellow to yellow crystalline powder.

CLINICAL TRIALS

Study demographics and trial design

A summary of the patient demographics for the two studies relevant to the evaluation of the efficacy of moxifloxacin hydrochloride ophthalmic solution is provided in Table 1. Overall, these demographics are representative of the population that would be expected to receive this medicinal product.

Table 1: Summary of Patient Demographics for Clinical Trials

Study #	Trial design	Dosage and route of administration and duration	Treatment duration	No. Patients (Intent to Treat)
C-00-55	Double-masked, randomized, vehicle controlled	0.5% moxifloxacin hydrochloride ophthalmic solution: 1 drop, TID Vehicle: 1 drop, TID	4 days	ITT = 544 270 moxifloxacin hydrochloride ophthalmic solution TID 274 Vehicle TID
C-00-46	Double-masked, randomized, active-controlled	0.5% moxifloxacin hydrochloride ophthalmic solution: 1 drop, TID Ocuflow: 1 drop, QID	4 days	ITT = 554 277 moxifloxacin hydrochloride ophthalmic solution TID 277 Ocuflow QID

Study results

In two, randomized, double-masked, multicenter, controlled trials in which 547 patients dosed with moxifloxacin hydrochloride 3 times a day for 4 days, moxifloxacin hydrochloride produced clinical cures on day 5 to 6 in 66% to 69% of patients treated for bacterial conjunctivitis. Microbiological success rates for the eradication of the baseline pathogens ranged from 84% to 94% at the test-of-cure visit (day 9). Please note that microbiologic eradication does not always correlate with clinical outcome in anti-infective trials.

DETAILED PHARMACOLOGY

Animal Pharmacokinetics

Ocular tissue concentrations of moxifloxacin were determined in pigmented rabbits following a single bilateral 30 mcL topical administration of 0.3% ophthalmic solution of moxifloxacin (n=3 rabbits sampled at each time point). Mean maximum concentrations (C_{max}) in cornea and aqueous humor were 12.5 ± 3.8 mcg/g and 1.78 ± 0.39 mcg/mL, respectively, and were achieved within 30 minutes after dosing. In iris-ciliary body, a moxifloxacin C_{max} of 10.4 ± 5.6 mcg/g was observed at 1 hour and declined slowly relative to other tissues, presumably due to binding to melanin pigment, which is characteristic of fluoroquinolones. The accumulation in ocular tissues of moxifloxacin after multiple dosing has not been studied. Maximum plasma concentrations were low (approximately 0.01 mcg/mL) and declined rapidly.

The distribution of radiolabeled moxifloxacin was also studied in pigmented rabbits after a single

unilateral 30 mcL dose of a 0.3% ¹⁴C-moxifloxacin solution (n=4 rabbits sampled at each time point). Mean C_{max} values in cornea, conjunctiva, aqueous humor and iris-ciliary body were 10.6 ± 2.8 mcg/g, 2.54 ± 0.40 mcg/g, 1.36 ± 0.33 mcg/mL and 7.54 ± 3.34 mcg/g, respectively. Maximum concentrations and half-lives in ocular tissues are summarized in Table 2.

Table 2: Maximum Concentrations and Half-Lives of Radiolabeled Moxifloxacin in Ocular Tissues from Pigmented Rabbits

Tissue	C _{max} (mcg equivalents/g) ± SD	t _{1/2} (hours)
Cornea	10.6 ± 2.8	92
Conjunctiva	2.54 ± 0.40	43
Aqueous Humor	1.36 ± 0.33	5.6
Iris-Ciliary Body	7.54 ± 3.34	649
Lens	0.08 ± 0.06	37
Anterior Sclera	2.86 ± 1.01	1080
Posterior Sclera	0.09 ± 0.03	92
Choroid	0.441 ± 0.178	872
Retina	0.066 ± 0.016	48

Tear film concentrations of moxifloxacin were measured in pigmented rabbits (n=3) after single unilateral administration of 30 mcL of a 0.3% moxifloxacin ophthalmic solution. The mean concentration of moxifloxacin was 366 ± 214 mcg/mL at the first sampling point of 1 minute after dosing. The levels then declined rapidly such that by 5 minutes after dosing the concentrations were approximately 20 mcg/mL. The concentrations in the tear film were 1.73 ± 1.50 mcg/mL at 6 hours post-dosing. Tear concentration data are summarised in Table 3.

Table 3: Tear Concentrations of Moxifloxacin Following Administration of a 0.3% Moxifloxacin Solution to Pigmented Rabbits

Time After Dose (minutes)	Mean Concentration ± SD (mcg/mL)	Sample Size
1	366 ± 214	3
2	74.2 ± 70.6	3
3	60.9 ± 11.9	3
5	23.7 ± 17.2	3
10	19.4 ± 4.03	3
20	23.4 ± 11.6	3
30	10.3 ± 3.6	3
45	1.21 ± 0.65	3
60	7.14 ± 6.12	3
90	2.69 ± 1.32	3
120	7.27 ± 9.96	2*
180	1.67 ± 1.06	2*
360	1.73 ± 1.50	2*

* 1 of 3 samples below quantitation limit of the assay. These samples were assigned a value of one half the limit of quantitation for calculation of the mean (1 mcg/mL/2 = 0.5 mcg/mL).

Human Pharmacokinetics


Plasma concentrations were studied in 21 healthy male and female subjects who were administered moxifloxacin hydrochloride to both eyes every 8 hours for a total of 13 doses. The results showed measurable plasma concentrations of moxifloxacin (≥ 0.75 ng/mL) in 16 of 21 subjects at 4-hours

following the first dose, and in all subjects following the last dose. Figure 1 shows the mean moxifloxacin plasma concentrations following the last dose.

The mean steady-state estimates for C_{\max} and AUC were 2.7 ng/mL and 41.9 ng.hr/mL, respectively. The steady-state parameter estimates for C_{\max} and AUC were at least 1600 and 1000 fold lower than mean C_{\max} and AUC values reported after therapeutic 400 mg oral doses of moxifloxacin. The steady-state plasma half-life of moxifloxacin was estimated to be 13 hours.

Subgroup analysis by race (Caucasian, Asian) showed no meaningful differences in the mean steady-state pharmacokinetic parameters of moxifloxacin. Gender differences in the steady-state C_{\max} and AUC were seen; however, when adjusted for body weight, the differences were minimized and not clinically relevant.

Figure 1: Mean (\pm SD) Moxifloxacin Plasma profile Following the Last Topical Ocular Dose of Moxifloxacin Hydrochloride Ophthalmic Solution in Healthy Subjects



Tear film concentrations of moxifloxacin were studied in 31 healthy male and female adult volunteers who were administered 1 drop of moxifloxacin hydrochloride to both eyes every 8 hours for a total of 10 doses.

Mean tear concentrations at 5 minutes following the first and last topical dose were 46.0 and 55.2 mcg/mL, respectively. Thereafter, mean tear concentrations rapidly declined in a biphasic manner with means ranging from approximately 1 to 4 mcg/mL over the 1 to 8 hour sampling period. Pre-dose morning tear concentrations on Days 2 to 4 averaged over 4 mcg/mL, demonstrating that concentrations are above the MICs for most of the common organisms in conjunctivitis over the 24-hour period.

Elimination and Metabolism: Moxifloxacin is widely distributed in the body tissues and approximately 50% is bound to serum proteins. Animal studies indicate some penetration into conjunctiva and ocular tissues with prolonged binding to melanin. Approximately 45% of an oral dose is excreted as unchanged drug, and most of the rest as glucuronide and sulfate conjugates in

feces and urine. The cytochrome P450 enzyme system is not involved in metabolizing the drug.

Drug-Drug Interactions: Specific drug-drug pharmacokinetic interaction studies were not conducted with moxifloxacin hydrochloride. Given the low systemic exposure observed for moxifloxacin after topical ocular administration of Sandoz Moxifloxacin solution, clinically relevant drug-drug interactions through protein binding, renal elimination or hepatic metabolism are unlikely following topical ocular administration. Moxifloxacin can be chelated by polyvalent ions such as Mg^{++} , Al^{+++} , Fe^{++} and Zn^{++} .

In vitro studies with cytochrome P450 isozymes have shown that moxifloxacin does not inhibit CYP3A4, CYP2D6, CYP2C9, CYP2C19 or CYP1A2 indicating that moxifloxacin is unlikely to alter the pharmacokinetics of drugs metabolized by these enzymes.

Special Populations: The pharmacokinetics of moxifloxacin hydrochloride has not been studied in patients with hepatic or renal impairment. However, the pharmacokinetics of orally administered moxifloxacin has been studied in these special populations.

The pharmacokinetic parameters of oral moxifloxacin are not significantly altered by mild, moderate or severe renal impairment. No dosage adjustment of moxifloxacin hydrochloride is necessary in patients with renal impairment.

Pharmacokinetic parameters of oral moxifloxacin were not significantly altered in patients with mild to moderate hepatic insufficiency (Child Pugh Classes A and B). Studies were not performed in patients with severe hepatic impairment (Child Pugh Class C). Because of the low systemic exposure by the topical route of administration, no dosage adjustment of moxifloxacin hydrochloride is needed in patients with hepatic impairment.

MICROBIOLOGY

Moxifloxacin has *in vitro* activity against a wide range of Gram-positive and Gram-negative microorganisms.

The antibacterial action of moxifloxacin results from inhibition of topoisomerase II (DNA gyrase) and topoisomerase IV. DNA gyrase is an essential enzyme that is involved in the replication, transcription and repair of bacterial DNA. Topoisomerase IV is an enzyme known to play a key role in the partitioning of the chromosomal DNA during bacterial cell division. The presence of the bulky bicycloamine substituent at the C-7 position prevents active efflux, a proposed mechanism of fluoroquinolone resistance.

Moxifloxacin concentrations at twice the MIC are sufficient to be bactericidal for most strains of *Staphylococcus aureus*, *Streptococcus pneumoniae*, and *Haemophilus influenzae*. Concentrations of moxifloxacin somewhat greater than twice the MIC were bactericidal for strains of *Escherichia coli* while those greater than ten times the MIC were bactericidal for *Streptococcus pyogenes*.

Resistance: The mechanism of resistance of quinolones, including moxifloxacin, is different from

that of macrolides, aminoglycosides, tetracyclines or β -lactams. Therefore, moxifloxacin may be active against pathogens that are resistant to these antibiotics and these antibiotics may be active against pathogens that are resistant to moxifloxacin. There is no cross-resistance between moxifloxacin and the aforementioned classes of antibiotics. Cross resistance has been observed between systemic moxifloxacin and some other quinolones.

In vitro resistance to moxifloxacin develops slowly via multiple-step mutations and occurs *in vitro* at a general frequency of between 1.8×10^{-9} to less than 1×10^{-11} in one strain of *Staphylococcus aureus* and one strain of *Streptococcus pneumoniae*.

Moxifloxacin has been shown to be active against most strains of the following microorganisms (see Table 4), both *in vitro* and in clinical infections from the US and India (see INDICATIONS AND USE)

Table 4: Moxifloxacin *In Vitro* Activity Against Clinical Isolates

Pathogen	N	MIC Range mcg/mL	MIC ₅₀ mcg/mL	MIC ₉₀ mcg/mL
Aerobic, Gram-Positive				
<i>Staphylococcus aureus</i>	49	≤ 0.016 - 2.0	0.06	1.0
<i>Staphylococcus epidermidis</i>	119	≤ 0.016 - 2.0	0.06	0.25
<i>Staphylococcus haemolyticus</i>	22	0.03 - 2.0	0.13	1.0
<i>Staphylococcus hominis</i>	11	0.06 - 1.0	0.06	0.13
<i>Streptococcus pneumoniae</i>	42	0.03 - 0.25	0.13	0.25
<i>Streptococcus viridans</i> group	22	0.06 - 2.0	0.25	0.25
Aerobic, Gram-Negative				
<i>Acinetobacter</i> species	15	≤ 0.016 - 0.25	0.03	0.06
<i>Haemophilus influenzae</i>	68	≤ 0.016 - 0.25	0.06	0.13

The following *in vitro* data (Table 5) are also available, but their clinical significance in ophthalmic infections is unknown. The safety and effectiveness of moxifloxacin hydrochloride in treating ophthalmic infections due to these organisms have not been established in adequate and well-controlled trials. The following organisms are considered susceptible when evaluated using systemic breakpoints. However, a correlation between the *in vitro* systemic breakpoint and ophthalmic efficacy has not been established. This list of organisms (Table 5) is provided as guidance only in assessing the potential treatment of conjunctival infections. Moxifloxacin exhibits *in vitro* minimal inhibitory concentrations (MICs) of 2 mcg/mL or less (systemic breakpoint susceptibility) against most (greater than or equal to 90%) strains of the following ocular isolates:

Table 5: Susceptibility of Bacterial Conjunctivitis Isolates to Moxifloxacin

Bacterial Species	N	MIC Range mcg/mL	MIC ₅₀ mcg/mL	MIC ₉₀ mcg/mL
Aerobic Gram-positive Microorganisms				
<i>Bacillus cereus</i>	15	0.032 - 0.25	0.13	0.13
<i>Corynebacterium</i> species	35	0.016 - 16	0.25	2.0
<i>Kocuria</i> species	11	0.25 - 0.50	0.25	0.50
<i>Micrococcus luteus</i>	35	0.03 - 1.0	0.5	1.0
<i>Staphylococcus capitis</i>	68	0.03 - 1.0	0.13	0.25
<i>Staphylococcus caprae</i>	13	0.06 - 0.13	0.06	0.13

Bacterial Species	N	MIC Range mcg/mL	MIC ₅₀ mcg/mL	MIC ₉₀ mcg/mL
<i>Staphylococcus lugdunensis</i>	36	0.06 - 1.0	0.13	0.25
<i>Staphylococcus pasteurii</i>	15	0.06 - 1.0	0.13	0.25
<i>Staphylococcus saprophyticus</i>	18	0.13 - 0.25	0.13	0.25
<i>Staphylococcus warneri</i>	10	0.06 - 0.13	0.13	0.13
<i>Streptococcus mitis</i>	76	0.06 - 0.25	0.13	0.25
<i>Streptococcus oralis</i>	10	0.13 - 0.25	0.13	0.25
<i>Streptococcus parasanguinis</i>	18	0.06 - 1.0	0.13	0.25
Aerobic, Gram-negative Microorganisms				
<i>Acinetobacter baumannii</i>	23	0.03 - 0.50	0.13	0.25
<i>Acinetobacter junii</i>	27	0.03 - 8.0	0.06	0.13
<i>Acinetobacter schindleri</i>	10	0.03 - 0.06	0.03	0.06
<i>Acinetobacter ursingii</i>	10	0.06 - 1.0	0.25	0.50
<i>Citrobacter koseri</i>	12	0.016 - 0.25	0.03	0.13
<i>Enterobacter hormaechei</i>	13	0.06 - 8.0	0.13	0.5
<i>Escherichia coli</i>	21	0.03 - 32	0.06	1.0
<i>Klebsiella pneumoniae</i>	17	0.06 - 2.0	0.13	0.5
<i>Moraxella osloensis</i>	13	0.03 - 0.25	0.06	0.25
<i>Moraxella catarrhalis</i>	25	0.06 - 0.13	0.06	0.13
<i>Pseudomonas stutzeri</i>	67	0.03 - 2.0	0.25	0.50
<i>Serratia marcescens</i>	20	0.25 - 2.0	0.5	1.0
<i>Stenotrophomonas maltophilia</i>	18	0.25 - 2.0	0.5	2.0

Susceptibility Tests: There are currently no NCCLS approved standards for assessing *in vitro* susceptibility of conjunctival isolates to topical antibiotics, including moxifloxacin. Standardized systemic susceptibility tests may not be appropriate to predict clinical effectiveness in treating conjunctivitis.

TOXICOLOGY

Topical Ocular Studies: Ophthalmic solutions of moxifloxacin were evaluated in repeat-dose topical ocular studies in rabbits (pigmented) and Cynomolgus monkeys (see Table 6).

Table 6: Results of Topical Ocular Studies

Species/No. per Group	Dose/Route	Duration of Treatment	Findings
Rabbits (pigmented)/ 4 male, 4 female	0.5%, 1%, 3% (80 mcL unilateral, 4 times daily) / topical ocular	1 month	Low ocular irritation potential; no significant ocular or systemic effects
Cynomolgus monkeys/ 4 male, 4 female	0.5%, 1%, 3% (80 mcL unilateral, 6 times daily Days 1-16, 3 times daily thereafter) / topical ocular	3 months	Low ocular irritation potential; no significant ocular or systemic effects

Ocular Toxicity Study: A special ocular toxicity study was conducted in dogs following systemic (oral) administration of moxifloxacin (see Table 7). The daily dosages of moxifloxacin evaluated in this study are significantly higher than the recommended daily dose of Moxifloxacin hydrochloride.

Table 7: Results of Ocular Toxicity Study

Species/No. per Group	Dose/Route	Duration of Treatment	Findings
Dog (Beagle)/ 4 males	30, 60, 90 mg/kg moxifloxacin / orally 100 mg/kg nalidixic acid (positive control) / orally	2 weeks (with 8 week recovery period)	↓ in group mean amplitude of a- and b- waves at 60 and 90 mg/kg moxifloxacin and with nalidixic acid; Histopath: slight to marked atrophy in outer nuclear and plexiform layers and rod and cone layers of two high dose animal; NOEL = 30 mg/kg orally (over 1300 times > the human dose of moxifloxacin hydrochloride solution)

Single and Repeat-Dose Oral and IV Studies: Oral and intravenous single-dose studies conducted with moxifloxacin are summarized in Table 8, and repeat-dose systemic studies that included ocular evaluations are summarized in Table 9. The daily dose levels of moxifloxacin evaluated in these studies are significantly higher than the recommended daily dose of moxifloxacin hydrochloride.

Table 8: Single-Dose Systemic Studies

Species	Strain/Sex	No./Group	Route of Administration	LD ₅₀ mg/kg BW (Conf. Int. for 95%)
Mouse	NMRI/male	5	PO	Approx. 435
	NMRI/female		PO	Approx. 758 (440-1305)
	NMRI/male		IV	Approx. 105 (84-132)
	NMRI/female		IV	Approx. 130 (116-145)
	WU/male		PO	Approx. 1320
	WU/female		PO	Approx. 1320
	WU/male		IV	Approx. 112
	WU/female		IV	Approx. 146
Monkey	Cynomolgus/ Male	2	PO	Approx. 1500

Table 9: Repeat-Dose Systemic Studies

Species/No. per Group	Dose/Route	Duration of Treatment	Findings
Wistar rats/ 15 male, 15 female	0, 20, 100, 500, 750 mg/kg / orally by gavage	13 weeks for all groups; 1 group examined after a 4 week recovery period	↓ body wt. gain at 100, 500, 750 mg/kg males; ocular evaluations (indirect ophthalmoscope and slit-lamp) unremarkable; ↑ASAT, ALAT, LDH at 500, 750 mg/kg males and females at 750 mg/kg; histopath unremarkable; NOAEL for females 100 mg/kg, 20 mg/kg for males
Wistar rats/ 20 male, 20 female	0, 20, 100, 500 mg/kg / orally by gavage	28 weeks	↓ body wt. gain at 500 mg/kg both sexes; ↑ASAT, ALAT, LDH, bilirubin 500 mg/kg males; ocular evaluations (indirect ophthalmoscope and slit- lamp) unremarkable; histopath 500 mg/kg both sexes, thyroid 500 mg/kg males NOAEL females 100 mg/kg, males 20 mg/kg

Species/No. per Group	Dose/Route	Duration of Treatment	Findings
Young Beagle pups/ 4 male, 4 female	0, 10, 30, 90 mg/kg/PO	4 weeks	Vacuolization of subcapsular lens cortex (indirect ophthalmoscope and slit-lamp) at 90 mg/kg; no evidence of co-cataractogenesis; prolongation of QT interval at 90 mg/kg; histopath chondropathy at 30 and 90 mg/kg
Young Beagle pups/ 2 male, 2 female	0, 10, 30, 90 mg/kg/PO	4 weeks	Vomiting, salivation, ↓body wt. gain at 90 mg/kg; ocular evaluations (indirect ophthalmoscope) unremarkable; histopath blistering of articular cartilage at 30 and 90 mg/kg
Rhesus monkeys/ 3 male, 3 female	0, 100, 150 mg/kg/orally by gavage	4 weeks	↓body wt. gain at 150 mg/kg; ocular evaluations (indirect ophthalmoscope) unremarkable; histopath liver and bone marrow at 100 and 150 mg/kg
Rhesus monkeys/ 4 male, 4 female	0, 15, 45, 135 mg/kg/orally by gavage	13 weeks	Salivation at 15 mg/kg; salivation, vomiting, ↓body wt. gain at 135 mg/kg; ocular evaluations (indirect ophthalmoscope) unremarkable; NOEL 15 mg/kg
Rhesus monkeys/ 4 male, 4 female	0, 15, 45, 135 mg/kg/orally by gavage	26 weeks	1 mortality at 135 mg/kg; ocular evaluations (indirect ophthalmoscope) unremarkable; ↑ALAT and GLDH at 45 mg/kg; histopath liver and bone marrow at 135 mg/kg; NOEL 15 mg/kg

Mutagenicity: Moxifloxacin was not mutagenic in four bacterial strains used in the Ames *Salmonella* reversion assay. As with other quinolones, the positive response observed with moxifloxacin in strain TA 102 using the same assay may be due to the inhibition of DNA gyrase. Moxifloxacin was not mutagenic in the CHO/HGPRT mammalian cell gene mutation assay. An equivocal result was obtained in the same assay when v79 cells were used. Moxifloxacin was clastogenic in the v79 chromosome aberration assay, but it did not induce unscheduled DNA synthesis in cultured rat hepatocytes. There was no evidence of genotoxicity *in vivo* in a micronucleus test or dominant lethal test in mice.

Carcinogenicity: Long term studies in animals to determine the carcinogenic potential of moxifloxacin have not been performed. However, in an accelerated study with initiators and promoters, moxifloxacin was not carcinogenic following up to 38 weeks of oral dosing at 500 mg/kg/day.

Reproduction and Teratology: Moxifloxacin had no effect on fertility in male and female rats at oral doses as high as 500 mg/kg/day, approximately 21,700 times the highest recommended total daily human ophthalmic dose.

Moxifloxacin was not teratogenic when administered to pregnant rats during organogenesis at oral doses as high as 500 mg/kg/day (approximately 21,700 times the highest recommended total human ophthalmic dose); however, decreased fetal body weights and slightly delayed fetal skeletal development were observed. When ¹⁴C-moxifloxacin was administered orally to pregnant rats, radioactivity penetrated the placenta and was absorbed to a moderate extent by the fetus. The ratio for AUC (0-24 h) for fetal plasma to maternal plasma was 0.656.

There was no evidence of teratogenicity when pregnant *Cynomolgus* monkeys were given oral doses as high as 100 mg/kg/day (approximately 4300 times the highest recommended total daily human ophthalmic dose). An increased incidence of smaller fetuses was observed at 100 mg/kg/day. In an oral peri/postnatal development study conducted in rats, marginal effects observed at 500 mg/kg/day included extended duration of pregnancy, increased prenatal loss, reduced birth weight and decreased survival index. Maternal mortality occurred at 500 mg/kg/day.

In an intravenous rabbit study, moxifloxacin at 20 mg/kg (approximately 860 times the highest recommended total daily human ophthalmic dose) was found to decrease the gestation rate, decrease fetal weights and delay ossification.

REFERENCES

1. Blondeau JM. A review of the comparative *in-vitro* activities of 12 antimicrobial agents, with a focus on five new 'respiratory quinolones'. J Antimicrob Chemother 1999; 43 Suppl B:1-11.
2. Blondeau JM, Felmingham D. *In vitro* and *in vivo* activity of moxifloxacin against community respiratory tract pathogens. Clin Drug Invest 1999;18:57-78.
3. Boswell FJ, Andrews JM, Wise R, Dalhoff A. Bactericidal properties of moxifloxacin and post-antibiotic effect. J Antimicrob Chemother 1999;43(Suppl B):43-9.
4. Brueggemann AB, Coffman SL, Rhomberg PR, Huynh HK, Almer L, Nilius A, Flamm R, Doern GV. Fluoroquinolone resistance in *Streptococcus pneumoniae* in United States since 1994-1995. Antimicrob Agents Chemother 2002;46:680-8.
5. Dalhoff A, Petersen U, Endermann R. *In vitro* activity of Bay 12-8039, a new 8-methoxyquinolone. Chemotherapy 1996; 42:410-25.
6. Doern GV, Heilmann KP, Huynh HK, Rhomberg PR, Coffman SL, Brueggemann AB. Antimicrobial resistance among clinical isolates of *Streptococcus pneumoniae* in the United States during 1999-2000, including a comparison of resistance rates since 1994-1995. Antimicrob Agents Chemother 2001;45:1721-9.
7. Esposito S, Noviello S, Ianniello F. Comparative *in vitro* activity of older and newer fluoroquinolones against respiratory tract pathogens. Chemotherapy 2000;46:309-314.
8. Jones ME, Sahm DF, Martin N, Scheuring S, Heisig P, Thornsberry C, Kohrer K, Schmitz F-J. Prevalence of *gyrA*, *gyrB*, *parC*, and *parE* mutations in clinical isolates of *Streptococcus pneumoniae* with decreased susceptibilities to different fluoroquinolones and originating from worldwide surveillance studies during 1997-1998 respiratory season. Antimicrob Agents Chemother 2000;44(2):462-6.
9. Klepser ME, Ernst EJ, Petzold R, Rhomberg P, Doern GV. Comparative bactericidal activities of ciprofloxacin, clinafloxacin, grepafloxacin, levofloxacin, moxifloxacin, and trovafloxacin against *Streptococcus pneumoniae* in a dynamic *in vitro* model. Antimicrob Agents Chemother 2001;45:673-8.
10. Mather R, Karenchak LM, Romanowski EG, Kowalski RP. Fourth generation fluoroquinolones: new weapons in the arsenal of ophthalmic antibiotics. Am J Ophthalmol 2002;133:463-6.
11. Pong A, Thomson KS, Moland, ES, Chartrand SA, Sanders CC. Activity of moxifloxacin against pathogens with decreased susceptibility to ciprofloxacin. J Antimicrob Chemother 1999;44(5):621-7.

12. Schedletzky H, Wiedemann B, Heisig P. The effect of moxifloxacin on its target topoisomerases from *Escherichia coli* and *Staphylococcus aureus*. J Antimicrob Chemother 1999; 43 Suppl B:31-7.
13. Schmitz FJ, Hofmann B, Hansen B, Scheuring S, Luckefahr M, Klootwijk M, Verhoef J, Fluit A, Heinz HP, Kohrer K, Jones ME. Relationship between ciprofloxacin, ofloxacin, levofloxacin, sparfloxacin and moxifloxacin (BAY 12-8039) MICs and mutations in *grlA*, *grlB*, *gyrA* and *gyrB* in 116 unrelated clinical isolates of *Staphylococcus aureus*. J Antimicrob Chemother 1998;41: 41-484.
14. Siefert HM, Domdey-Bette A, Henninger K, Hucke F, Kohlsdorfer C, Stass HH. Pharmacokinetics of the 8-methoxyquinolone, moxifloxacin: a comparison in humans and other mammalian species. J Antimicrob Chemother 1999;43(Suppl B):69-76.
15. Stass H, Kubitz D. Pharmacokinetics and elimination of moxifloxacin after oral and intravenous administration. J Antimicrob Chemother 1999;43(Suppl B):83-90.
16. Tankovic J, Bachoual R, Ouabdesselam S, Boudjadja A, Soussy C-J. *In vitro* activity of moxifloxacin against fluoroquinolone-resistant strains of aerobic gram-negative bacilli and *Enterococcus faecalis*. J Antimicrob Chemother 1999; 43:19-23.
17. Novartis Pharmaceuticals Canada Inc. Product Monograph: VIGAMOX. Control #: 204458. Date of Revision: December 1, 2017.

READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE

PATIENT MEDICATION INFORMATION

SANDOZ MOXIFLOXACIN **Moxifloxacin ophthalmic solution** **0.5% w/v (as moxifloxacin hydrochloride)**

Read this carefully before you start taking Sandoz Moxifloxacin and each time you get a refill. This leaflet is a summary and will not tell you everything about this drug. Talk to your healthcare professional about your medical condition and treatment and ask if there is any new information about Sandoz Moxifloxacin.

What is Sandoz Moxifloxacin used for?

Sandoz Moxifloxacin is an antibiotic used to treat bacterial infection of the eye (pink eye).

How does Sandoz Moxifloxacin work?

Sandoz Moxifloxacin is an antibiotic that stops the growth of bacteria. This kills the bacteria and reduces eye infections.

What are the ingredients in Sandoz Moxifloxacin?

Medicinal ingredients: Moxifloxacin 0.5% w/v (as moxifloxacin hydrochloride)

Non-medicinal ingredients: Boric acid, purified water and sodium chloride. May also contain hydrochloric acid and/or sodium hydroxide to adjust pH.

Sandoz Moxifocacin comes in the following dosage forms:

Sandoz Moxifloxacin is a clear greenish-yellow solution and comes in a 3 mL oval plastic bottle.

Do not use Sandoz Moxifloxacin if you are:

- Allergic to moxifloxacin or any of the ingredients in Sandoz Moxifloxacin (see What are the ingredients in Sandoz Moxifloxacin?)
- Allergic to other quinolones.

To help avoid side effects and ensure proper use, talk to your healthcare professional before you take Sandoz Moxifloxacin. Talk about any health conditions or problems you may have, including if you:

- Develop an allergic reaction (see Serious side effects and what to do about them). Stop taking Sandoz Moxifloxacin and get immediate medical help.
- Notice your infection gets worse. Stop taking Sandoz Moxifloxacin and get immediate medical help. As with any antibiotic, use of Sandoz Moxifloxacin for a long time may lead to other infections. Do NOT use Sandoz Moxifloxacin longer than your doctor tells you to.
- Develop pain or swelling in your tendons. Stop taking Sandoz Moxifloxacin and get immediate medical help. This is more likely to happen if you are elderly or taking corticosteroids at the same time as Sandoz Moxifloxacin.

Other warnings you should know about:

- Sandoz Moxifloxacin can be used in children as young as 1 year of age. Sandoz Moxifloxacin should not be used in children younger than 1 year of age.
- Do NOT wear contact lenses if you have an eye infection.
- Your vision may be temporarily blurry after using Sandoz Moxifloxacin. Wait until your vision clears before driving or using machinery.
- If you are pregnant or planning to become pregnant or are breast-feeding or planning to breastfeed, talk to your doctor or pharmacist before using Sandoz Moxifloxacin.
- Antibacterial drugs like Sandoz Moxifloxacin treat only bacterial infections. They do not treat viral infections, such as the common cold. Although you may feel better early in treatment, Sandoz Moxifloxacin should be taken exactly as directed. Misuse or overuse of Sandoz Moxifloxacin could lead to the growth of bacteria that will not be killed by Sandoz Moxifloxacin (resistance). This means that Sandoz Moxifloxacin may not work for you in the future. Do not share your medicine.

Tell your healthcare professional about all the medicines you take, including any drugs, vitamins, minerals, natural supplements or alternative medicines.

The following may interact with Sandoz Moxifloxacin:

- Tell your doctor or pharmacist if you are:
 - taking or have recently taken any prescription or non-prescription medicines,
 - using any other eye products,
 - taking corticosteroids as this may increase your chance of developing pain or swelling of you tendons.

How to take Sandoz Moxifloxacin:

1. Get the bottle of Sandoz Moxifloxacin eye drops and a mirror.
2. Wash your hands.
3. Open the bottle being careful not to touch the dropper tip.
4. Hold the bottle, pointing down, between your thumb and fingers.
5. Tilt your head back.
6. Pull down your eyelid with a clean finger, until there is a 'pocket' between the eyelid and eye. The drop will go in there.
7. Bring the bottle close to the eye. Use the mirror if it helps.
8. **Do not touch your eye or eyelid, surrounding areas or other surfaces with the dropper.** It could contaminate the drops.
9. Gently squeeze the bottle to release one drop of Sandoz Moxifloxacin.
10. If the drop misses your eye, wipe it up and try again.
11. If you need drops in both eyes, repeat the steps for your other eye.

12. Keep the bottle tightly closed when not in use.

Usual dose:

Apply one drop in the affected eye(s) three times a day (morning, afternoon, and at night) unless your doctor tells you otherwise.

Use Sandoz Moxifloxacin for seven days or as long as your doctor tells you to.

Overdose:

If you use too much Sandoz Moxifloxacin, rinse it out of your eyes with warm water. Do not put any more drops of Sandoz Moxifloxacin in until it's time for your next dose.

If you accidentally swallow Sandoz Moxifloxacin, talk to your doctor or pharmacist.

If you think you have taken too much Sandoz Moxifloxacin, contact your healthcare professional, hospital emergency department or regional Poison Control Centre immediately, even if there are no symptoms.

Missed Dose:

If you miss a dose, apply the missed dose as soon as possible and then go back to your regular dosing schedule. If the drop misses your eye, try again.

What are possible side effects from using Sandoz Moxifloxacin?

These are not all the possible side effects you may feel when taking Sandoz Moxifloxacin. If you experience any side effects not listed here, contact your healthcare professional.

Common eye side effects while using Sandoz Moxifloxacin include:

- mild temporary burning or stinging,
- itching or redness.

Other less common eye side effects include:

- dryness,
- sensation of pressure, discomfort, corneal inflammation, broken blood vessels in the white part of the eye, swelling of the eye or eyelid,
- blurry vision,
- temporary reduction of vision,
- pain, inflammation of the eye surface or eyelid, tired eyes, redness of the eyelid,
- watery eyes, sensitivity to light, eye discharge.

You may also experience reactions in the other areas of your body, including:

- bad taste,
- headache,
- throat pain,
- abnormal liver blood tests,
- abnormal skin sensation,

- vomiting,
- nose discomfort,
- dizziness,
- irregular heart rhythm,
- shortness of breath,
- nausea,
- allergic reaction,
- skin redness or itching,
- rash or hives.

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
UNKNOWN Severe allergic reaction: <ul style="list-style-type: none"> • swelling of hands • feet • ankles • face • lips • mouth or throat • difficulty breathing • fever • rash of hives • large fluid-filled blisters • sores 			√

If you have a troublesome symptom or side effect that is not listed here or becomes bad enough to interfere with your daily activities, talk to your healthcare professional.

Reporting Side Effects

You can report any suspected side effects associated with the use of health products to the Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<http://www.hc-sc.gc.ca/dhp-mpps/medeff/report-declaration/index-eng.php>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345

NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

Storage

Store between 4°C and 25°C. Keep out of the reach and sight of children.

Do not use Sandoz Moxifloxacin after the expiry date (shown as "EXP" on the package). Discard 28 days after opening.

If you want more information about Sandoz Moxifloxacin:

- Talk to your healthcare professional.
- This document, plus the full Product Monograph prepared for health professionals, can be obtained by contacting the sponsor, Sandoz Canada Inc., at:
 - 1-800-361-3062
 - or by written request at:
145, Jules-Léger
Boucherville, Quebec
J4B 7K8
 - or by e-mail at: medinfo@sandoz.com

This leaflet was prepared by Sandoz Canada Inc.

Last revised: February 2, 2018