

**PRODUCT MONOGRAPH**

**Pr APO-TEMOZOLOMIDE**

**Temozolomide Capsules**

**5 mg, 20 mg, 100 mg, 140 mg, 180 mg and 250 mg**

**Antineoplastic Agent**

**APOTEX INC.  
150 Signet Drive  
Toronto, Ontario  
M9L 1T9**

**DATE OF REVISION:  
February 1, 2018**

**Control No: 207074**

**Table of Contents**

**PART I: HEALTH PROFESSIONAL INFORMATION..... 3**

SUMMARY PRODUCT INFORMATION ..... 3

INDICATIONS AND CLINICAL USE ..... 3

CONTRAINDICATIONS ..... 3

WARNINGS AND PRECAUTIONS ..... 4

ADVERSE REACTIONS ..... 8

DRUG INTERACTIONS ..... 18

DOSAGE AND ADMINISTRATION ..... 19

OVERDOSAGE ..... 21

ACTION AND CLINICAL PHARMACOLOGY ..... 22

STORAGE AND STABILITY ..... 22

SPECIAL HANDLING INSTRUCTIONS ..... 22

DOSAGE FORMS, COMPOSITION AND PACKAGING ..... 23

**PART II: SCIENTIFIC INFORMATION ..... 24**

PHARMACEUTICAL INFORMATION ..... 24

CLINICAL TRIALS ..... 26

DETAILED PHARMACOLOGY ..... 32

TOXICOLOGY ..... 35

REFERENCES ..... 50

**PART III: CONSUMER INFORMATION..... 51**

<sup>Pr</sup> **APO-TEMOZOLOMIDE**

**Temozolomide Capsules**

**5 mg, 20 mg, 100 mg, 140 mg, 180 mg and 250 mg**

**PART I: HEALTH PROFESSIONAL INFORMATION**

**SUMMARY PRODUCT INFORMATION**

| <b>Route of Administration</b> | <b>Dosage Form / Strength</b>                          | <b>All Nonmedicinal Ingredients</b>  |
|--------------------------------|--|--|
| Oral                           | Capsule/5 mg, 20 mg, 100 mg, 140 mg, 180 mg and 250 mg | <p>colloidal silicon dioxide, lactose anhydrous, sodium starch glycolate, stearic acid and tartaric acid.</p> <p>The empty capsule shells contain the non-medicinal ingredients FD&amp;C Blue No. 2 (5 mg, 140 mg), gelatin, iron oxide red (100 mg, 180 mg), iron oxide yellow (5 mg, 20 mg, 180 mg), talc and titanium dioxide.</p> <p>The capsule shells are imprinted with black edible ink containing ammonium hydroxide, iron oxide black, isopropyl alcohol, N-Butyl alcohol, propylene glycol and shellac glaze.</p> |

**INDICATIONS AND CLINICAL USE**

APO-TEMOZOLOMIDE (temozolomide) is indicated for:

- treatment of adult patients with newly diagnosed glioblastoma multiforme concomitantly with radiotherapy and then as maintenance treatment.
- treatment of adult patients with glioblastoma multiforme or anaplastic astrocytoma and documented evidence of recurrence or progression after standard therapy.

**CONTRAINDICATIONS**

- APO-TEMOZOLOMIDE is contraindicated in patients who have a history of hypersensitivity reaction to its components or to dacarbazine (DTIC).
- The use of APO-TEMOZOLOMIDE is not recommended in patients with severe myelosuppression.

## WARNINGS AND PRECAUTIONS

### Serious Warnings and Precautions

APO-TEMOZOLOMIDE should be prescribed by a qualified healthcare professional who is experienced in the use of antineoplastic therapy.

The following are clinically significant adverse events:

- Myelosuppression including Neutropenia and Thrombocytopenia and prolonged pancytopenia, which may result in aplastic anemia, which in some cases has resulted in a fatal outcome (see **WARNINGS AND PRECAUTIONS / Hematologic/Myelosuppression**).
- Hepatic injury, including fatal hepatic failure, has been reported in patients treated with temozolomide (see **WARNINGS AND PRECAUTIONS / Hepatic /Biliary/Pancreatic**).

APO-TEMOZOLOMIDE may have to be discontinued or the dose may have to be adjusted (see **DOSAGE AND ADMINISTRATION**).

### **General**

The treating physician should use his discretion with respect to the use of APO-TEMOZOLOMIDE in patients with poor performance status, severe debilitating diseases or infection when the risk of treatment outweighs the potential benefit to the patient.

### **Drug Interactions:**

Co-administration with valproic acid was associated with a small but statistically significant decrease in clearance of temozolomide.

The combination of temozolomide with other chemotherapeutic agents has not been fully evaluated. Combination with other alkylating agents is likely to result in increased myelosuppression.

### **Gastrointestinal**

#### Antiemetic therapy:

Nausea and vomiting are very commonly associated with temozolomide, and guidelines are provided:

Patients with newly diagnosed glioblastoma multiforme:

- anti-emetic prophylaxis is recommended prior to the initial dose of concomitant temozolomide,
- anti-emetic prophylaxis is strongly recommended during the maintenance phase.

Patients with recurrent or progressive glioma:

Patients who have experienced severe (Grade 3 or 4) vomiting in previous treatment cycles may require anti-emetic therapy.

## **Hematologic**

### **Myelosuppression:**

Temozolomide is an alkylating antitumor drug. Severe myelosuppression can occur, and is a dose limiting side effect. Temozolomide is associated with Grade 3 and Grade 4 neutropenia and Grade 3 and Grade 4 thrombocytopenia. Prior to dosing and during treatment, proper hematologic monitoring must be performed. APO-TEMOZOLOMIDE may have to be discontinued or the dose may have to be adjusted (see **WARNINGS AND PRECAUTIONS/ Monitoring and Laboratory Tests, ADVERSE REACTIONS** and **DOSAGE AND ADMINISTRATION/Administration**).

Patients treated with temozolomide who experience myelosuppression, may experience prolonged pancytopenia, which may result in aplastic anemia, which in some cases has resulted in a fatal outcome. In some cases, exposure to concomitant medications associated with aplastic anemia, including carbamazepine, phenytoin, and sulfamethoxazole/trimethoprim, complicates assessment.

## **Hepatic/Biliary/Pancreatic**

Hepatotoxicity, including liver enzyme elevation, hyperbilirubinemia, cholestasis and hepatitis, has been observed with temozolomide use in the post-market setting (see **ADVERSE REACTIONS/Post-Market Adverse Drug Reactions**). Hepatic injury, including fatal hepatic failure, has been reported in patients treated with temozolomide. Baseline liver function tests should be performed prior to treatment initiation. If abnormal, physicians should assess the benefit/risk prior to initiating temozolomide including the potential for fatal hepatic failure. For patients on a 42 day treatment cycle liver function tests should be repeated midway during this cycle. For all patients, liver function tests should be checked after each treatment cycle. For patients with significant liver function abnormalities, physicians should assess the benefit/risk of continuing treatment. Liver toxicity may occur several weeks or more after the last treatment with temozolomide. In the absence of formal studies in patients suffering from severe hepatic dysfunction the treating physician should use his discretion in weighing the benefits of using APO-TEMOZOLOMIDE in this patient population against the potential risks.

Additionally, hepatitis due to hepatitis B virus (HBV) reactivation, in some cases resulting in death, has been reported. Patients should be screened for HBV infection before treatment initiation. Patients with evidence of current or prior HBV infection should be monitored for clinical and laboratory signs of hepatitis or HBV reactivation during and for several months following treatment with APO-TEMOZOLOMIDE. Therapy should be discontinued for patients with evidence of active hepatitis B infection.

## **Infection**

Cases of herpes simplex encephalitis (HSE), including cases with fatal outcomes, were reported mostly in association with concomitant radiotherapy. All patients, particularly those with previous herpes simplex infection need to be monitored for signs and symptoms of HSE during the treatment.

### **Renal**

In the absence of formal studies in patients suffering from severe renal failure the treating physician should use his discretion in weighing the benefits of using APO-TEMOZOLOMIDE in this patient population against the potential risks.

### **Respiratory**

Patients who received concomitant temozolomide and radiotherapy in a pilot trial for the prolonged 42 day schedule were shown to be at particular risk for developing *Pneumocystis carinii* pneumonia. Thus prophylaxis against *Pneumocystis carinii* pneumonia (PCP) is required for all patients receiving concomitant temozolomide and radiotherapy for the 42 day regimen (with a maximum of 49 days). There may be a higher occurrence of PCP when temozolomide is administered during a longer dosing regimen. However, all patients receiving APO-TEMOZOLOMIDE, particularly patients receiving steroids should be observed closely for the development of PCP regardless of the regimen.

Cases of interstitial pneumonitis/pneumonitis have been reported in post-marketing experience. These events have the potential to be fatal.

### **Sexual Function/Reproduction**

**Male patients:** Temozolomide can have genotoxic effects. Effective contraception should also be used by male patients taking APO-TEMOZOLOMIDE. Men being treated with APO-TEMOZOLOMIDE are advised not to father a child during or up to 6 months after treatment and to seek advice on cryoconservation of sperm prior to treatment because of the possibility of irreversible infertility due to therapy with APO-TEMOZOLOMIDE

### **Skin**

Serious dermatologic reactions including Stevens-Johnson syndrome (SJS) and toxic epidermal necrolysis (TEN) have been reported in post-marketing experience. These events have the potential to be fatal. When SJS/TEN is suspected, appropriate action should be taken, including close monitoring of the patient. Discontinuation of all concomitant medications suspected to contribute to SJS/TEN and temozolomide should be evaluated.

### **Special Populations**

**Pregnant Women:** There are no studies in pregnant women. In preclinical studies in rats and rabbits administered 150 mg/m<sup>2</sup>, teratogenicity and/or fetal toxicity were demonstrated. Therefore, APO-TEMOZOLOMIDE should not be administered to pregnant women. If use during pregnancy must be considered, the patient should be apprised of the potential risks to the fetus. Women of childbearing potential should be advised to avoid pregnancy while they are receiving APO-TEMOZOLOMIDE therapy and in the six months after discontinuation of treatment.

**Nursing Women:** It is not known whether temozolomide is excreted in human milk. Lactating mothers should be advised to stop lactation while under treatment.

**Pediatrics (<18 years and >3 years):** The safety and effectiveness of temozolomide in pediatric patients has not yet been fully established.

**Geriatrics (>70 years of age):** Elderly patients appear to be at increased risk of neutropenia and thrombocytopenia, compared with younger patients.

### **Monitoring and Laboratory Tests**

Baseline liver function tests should be performed prior to treatment initiation. If abnormal, physicians should assess the benefit/risk prior to initiating temozolomide including the potential for fatal hepatic failure. For patients on a 42 day treatment cycle liver function tests should be repeated midway during this cycle. For all patients, liver function tests should be checked after each treatment cycle.

Liver toxicity may occur several weeks or more after the last treatment with temozolomide.

Patients should also be screened for HBV infection before treatment initiation. Patients with evidence of current or prior HBV infection should be monitored for clinical and laboratory signs of hepatitis or HBV reactivation during and for several months following treatment with APO-TEMOZOLOMIDE. Therapy should be discontinued for patients with evidence of active hepatitis B infection.

### **Concomitant phase for adult patients with newly diagnosed glioblastoma multiforme:**

Temozolomide is administered at 75 mg/m<sup>2</sup> daily for 42 days concomitant with radiotherapy (60 Gy administered in 30 fractions). A complete blood count should be obtained prior to initiation of treatment and weekly during treatment. APO-TEMOZOLOMIDE dosing should be interrupted or discontinued during concomitant phase according to the hematological and non-hematological toxicity criteria (see **DOSAGE AND ADMINISTRATION**).

### **Maintenance phase for adults with newly diagnosed glioblastoma multiforme or treatment for patients with malignant gliomas showing recurrence or progression after standard therapy:**

Temozolomide is administered at a dose of 150 or 200 mg/m<sup>2</sup> once daily for 5 days per 28-day cycle. Prior to dosing, on Day 1 of each cycle, the following values must be met: absolute neutrophil count (ANC) >1.5 x 10<sup>9</sup>/L and platelets >100 x 10<sup>9</sup>/L. A complete blood count must also be obtained on Day 22 (21 days after the first dose) or within 48 hours of that day, and weekly until ANC is above 1.5 x 10<sup>9</sup>/L and platelet count exceeds 100 x 10<sup>9</sup>/L. If the ANC falls to <1.0 x 10<sup>9</sup>/L or the platelet count is <50 x 10<sup>9</sup>/L during any cycle, the next cycle should be reduced by one dose level, based upon the nadir blood count (see **DOSAGE AND ADMINISTRATION**). Dose levels include 100 mg/m<sup>2</sup>, 150 mg/m<sup>2</sup> and 200 mg/m<sup>2</sup>. The lowest recommended dose is 100 mg/m<sup>2</sup>.

## ADVERSE REACTIONS

### Clinical trial experience in patients treated with temozolomide capsules

#### Newly Diagnosed Patients with Glioblastoma Multiforme

Table 1 provides treatment emergent adverse events, in (causality not determined during clinical trials) patients with newly diagnosed glioblastoma multiforme during the concomitant and maintenance phases of treatment.

| <i>Table 1. Temozolomide and radiotherapy: Treatment-emergent events during concomitant and maintenance treatment</i> |   |   |                                  |
|---|---|---|----------------------------------|
| <b>Body System</b>  | <b>Temozolomide + concomitant radiotherapy<br/>n=288*<br/>n (%)</b> | <b>Temozolomide maintenance therapy<br/>n=224<br/>n (%)</b> | <b>Total<br/>n=288<br/>n (%)</b> |
| <u>Infections and Infestations</u>  |   |   |                                  |
| Candidiasis oral  | 4 (1%)  | 5 (2%)  | 7 (2%)                           |
| Herpes simplex  | 4 (1%)  | 2 (1%)  | 6 (2%)                           |
| Herpes zoster   | 0 (0%)  | 3 (1%)  | 3 (1%)                           |
| Infection   | 4 (1%)  | 8 (4%)  | 12 (4%)                          |
| Influenza-like symptoms   | 0 (0%)  | 3 (1%)  | 3 (1%)                           |
| Pharyngitis   | 2 (1%)  | 1 (<1%)   | 3 (1%)                           |
| Wound infection   | 2 (1%)  | 0 (0%)  | 2 (1%)                           |
| <u>Blood and the lymphatic system disorders</u>   |   |   |                                  |
| Anemia  | 3 (1%)  | 4 (2%)  | 6 (2%)                           |
| Febrile neutropenia   | 2 (1%)  | 4 (2%)  | 6 (2%)                           |
| Leukopenia  | 6 (2%)  | 5 (2%)  | 10 (3%)                          |
| Lymphopenia   | 7 (2%)  | 2 (1%)  | 7 (2%)                           |
| Neutropenia   | 6 (2%)  | 7 (3%)  | 10 (3%)                          |
| Thrombocytopenia  | 11 (4%)   | 19 (8%)   | 29 (10%)                         |
| Petechiae   | 1 (<1%)   | 2 (1%)  | 3 (1%)                           |
| <u>Endocrine disorders</u>  |   |   |                                  |
| Cushingoid  | 4 (1%)  | 2 (1%)  | 6 (2%)                           |
| <u>Metabolism and nutrition disorders</u>   |   |   |                                  |
| Anorexia  | 56 (19%)  | 61 (27%)  | 91 (32%)                         |
| Alkaline phosphatase increased  | 3 (1%)  | 1 (<1%)   | 4 (1%)                           |
| Hyperglycemia   | 7 (2%)  | 3 (1%)  | 9 (3%)                           |
| Hypokalemia   | 2 (1%)  | 1 (<1%)   | 3 (1%)                           |
| Weight decreased  | 5 (2%)  | 7 (3%)  | 11 (4%)                          |
| Weight increased  | 4 (1%)  | 3 (1%)  | 6 (2%)                           |



**Table 1. Temozolomide and radiotherapy: Treatment-emergent events during concomitant and maintenance treatment**

| <b>Body System</b>              | <b>Temozolomide + concomitant radiotherapy<br/>n=288*<br/>n (%)</b> | <b>Temozolomide maintenance therapy<br/>n=224<br/>n (%)</b> | <b>Total<br/>n=288<br/>n (%)</b> |
|---------------------------------|---|---|----------------------------------|
| <u>Psychiatric disorders</u>    |   |   |                                  |
| Agitation                       | 2 (1%)  | 1 (<1%)   | 3 (1%)                           |
| Amnesia                         | 0 (0%)  | 2 (1%)  | 2 (1%)                           |
| Anxiety                         | 5 (2%)  | 8 (4%)  | 10 (3%)                          |
| Apathy                          | 2 (1%)  | 1 (<1%)   | 3 (1%)                           |
| Behavior disorder               | 2 (1%)  | 1 (<1%)   | 2 (1%)                           |
| Depression                      | 3 (1%)  | 6 (3%)  | 8 (3%)                           |
| Emotional lability              | 5 (2%)  | 7 (3%)  | 10 (3%)                          |
| Hallucination                   | 2 (1%)  | 2 (1%)  | 4 (1%)                           |
| Insomnia                        | 14 (5%)   | 9 (4%)  | 18 (6%)                          |
| <u>Nervous system disorders</u> |   |   |                                  |
| Aphasia                         | 9 (3%)  | 5 (2%)  | 11 (4%)                          |
| Ataxia                          | 3 (1%)  | 3 (1%)  | 5 (2%)                           |
| Cerebral hemorrhage             | 2 (1%)  | 0 (0%)  | 2 (1%)                           |
| Balance impaired                | 5 (2%)  | 4 (2%)  | 9 (3%)                           |
| Cognition impaired              | 2 (1%)  | 0 (0%)  | 2 (1%)                           |
| Concentration impaired          | 6 (2%)  | 6 (3%)  | 10 (3%)                          |
| Confusion                       | 11 (4%)   | 12 (5%)   | 22 (8%)                          |
| Consciousness decreased         | 5 (2%)  | 1 (<1%)   | 6 (2%)                           |
| Convulsions                     | 17 (6%)   | 25 (11%)  | 36 (13%)                         |
| Coordination abnormal           | 0 (0%)  | 2 (1%)  | 2 (1%)                           |
| Dizziness                       | 12 (4%)   | 12 (5%)   | 22 (8%)                          |
| Dysphasia                       | 4 (1%)  | 9 (4%)  | 10 (3%)                          |
| Extrapyramidal disorder         | 2 (1%)  | 0 (0%)  | 2 (1%)                           |
| Gait abnormal                   | 4 (1%)  | 3 (1%)  | 7 (2%)                           |
| Headache                        | 56 (19%)  | 51 (23%)  | 87 (30%)                         |
| Hemiparesis                     | 4 (1%)  | 8 (4%)  | 10 (3%)                          |
| Hemiplegia                      | 0 (0%)  | 2 (1%)  | 2 (1%)                           |
| Hyperesthesia                   | 2 (1%)  | 2 (1%)  | 3 (1%)                           |
| Hypoesthesia                    | 2 (1%)  | 1 (<1%)   | 3 (1%)                           |
| Memory impairment               | 8 (3%)  | 16 (7%)   | 21 (7%)                          |
| Neurological disorder (NOS)     | 3 (1%)  | 6 (3%)  | 7 (2%)                           |
| Neuropathy                      | 8 (3%)  | 6 (3%)  | 12 (4%)                          |
| Paresthesia                     | 6 (2%)  | 4 (2%)  | 7 (2%)                           |
| Peripheral neuropathy           | 2 (1%)  | 4 (2%)  | 5 (2%)                           |
| Sensory disturbance             | 0 (0%)  | 2 (1%)  | 2 (1%)                           |
| Somnolence                      | 5 (2%)  | 5 (2%)  | 10 (3%)                          |
| Speech disorder                 | 6 (2%)  | 9 (4%)  | 14 (5%)                          |
| Status epilepticus              | 2 (1%)  | 0 (0%)  | 2 (1%)                           |
| Tremor                          | 7 (2%)  | 9 (4%)  | 14 (5%)                          |

**Table 1. Temozolomide and radiotherapy: Treatment-emergent events during concomitant and maintenance treatment**

| <b>Body System</b>                                     | <b>Temozolomide + concomitant radiotherapy<br/>n=288*<br/>n (%)</b> | <b>Temozolomide maintenance therapy<br/>n=224<br/>n (%)</b> | <b>Total<br/>n=288<br/>n (%)</b> |
|--|---|---|----------------------------------|
| <u>Eye disorders</u>                                   |   |   |                                  |
| Diplopia   | 1 (<1%)   | 5 (2%)  | 6 (2%)                           |
| Eye pain   | 3 (1%)  | 2 (1%)  | 4 (1%)                           |
| Eyes dry   | 1 (<1%)   | 2 (1%)  | 2 (1%)                           |
| Hemianopia   | 2 (1%)  | 1 (<1%)   | 2 (1%)                           |
| Vision blurred   | 26 (9%)   | 17 (8%)   | 33 (11%)                         |
| Vision disorder  | 2 (1%)  | 2 (1%)  | 4 (1%)                           |
| Visual acuity reduced                                  | 2 (1%)  | 3 (1%)  | 4 (1%)                           |
| Visual field defect                                    | 4 (1%)  | 5 (2%)  | 7 (2%)                           |
| <u>Ear and labyrinth disorders</u>                     |   |   |                                  |
| Deafness   | 1 (<1%)   | 2 (1%)  | 2 (1%)                           |
| Earache  | 3 (1%)  | 3 (1%)  | 5 (2%)                           |
| Hearing impairment                                     | 8 (3%)  | 10 (4%)   | 13 (5%)                          |
| Hyperacusis  | 2 (1%)  | 1 (<1%)   | 2 (1%)                           |
| Otitis media   | 2 (1%)  | 0 (0%)  | 2 (1%)                           |
| Tinnitus   | 4 (1%)  | 4 (2%)  | 6 (2%)                           |
| Vertigo  | 1 (<1%)   | 3 (1%)  | 3 (1%)                           |
| <u>Cardiac disorders</u>                               |   |   |                                  |
| Palpitation  | 2 (1%)  | 0 (0%)  | 2 (1%)                           |
| <u>Vascular disorders</u>                              |   |   |                                  |
| Deep venous thrombosis                                 | 5 (2%)  | 4 (2%)  | 8 (3%)                           |
| Edema  | 6 (2%)  | 2 (1%)  | 8 (3%)                           |
| Edema leg  | 6 (2%)  | 4 (2%)  | 9 (3%)                           |
| Edema peripheral                                       | 0 (0%)  | 3 (1%)  | 3 (1%)                           |
| Embolism pulmonary                                     | 0 (0%)  | 2 (1%)  | 2 (1%)                           |
| Hemorrhage   | 7 (2%)  | 7 (3%)  | 13 (5%)                          |
| Hypertension   | 2 (1%)  | 1 (<1%)   | 3 (1%)                           |
| <u>Respiratory, thoracic and mediastinal disorders</u> |   |   |                                  |
| Bronchitis   | 0 (0%)  | 2 (1%)  | 2 (1%)                           |
| Coughing   | 15 (5%)   | 19 (8%)   | 26 (9%)                          |
| Dyspnea  | 11 (4%)   | 12 (5%)   | 19 (7%)                          |
| Nasal congestion                                       | 2 (1%)  | 1 (<1%)   | 3 (1%)                           |
| Pneumonia  | 4 (1%)  | 2 (1%)  | 6 (2%)                           |
| Upper respiratory infection                            | 4 (1%)  | 2 (1%)  | 6 (2%)                           |
| Sinusitis  | 1 (<1%)   | 2 (1%)  | 3 (1%)                           |

**Table 1. Temozolomide and radiotherapy: Treatment-emergent events during concomitant and maintenance treatment**

| <b>Body System</b>                                     | <b>Temozolomide + concomitant radiotherapy<br/>n=288*<br/>n (%)</b> | <b>Temozolomide maintenance therapy<br/>n=224<br/>n (%)</b> | <b>Total<br/>n=288<br/>n (%)</b> |
|--|---|---|----------------------------------|
| <u>Gastrointestinal disorders</u>                      |   |   |                                  |
| Abdominal distension                                   | 1 (<1%)   | 2 (1%)  | 3 (1%)                           |
| Abdominal pain   | 7 (2%)  | 11 (5%)   | 15 (5%)                          |
| Constipation   | 53 (18%)  | 49 (22%)  | 87 (30%)                         |
| Diarrhea   | 18 (6%)   | 23 (10%)  | 36 (13%)                         |
| Dyspepsia  | 9 (3%)  | 4 (2%)  | 10 (3%)                          |
| Dysphagia  | 6 (2%)  | 6 (3%)  | 9 (3%)                           |
| Fecal incontinence                                     | 0 (0%)  | 2 (1%)  | 2 (1%)                           |
| Gastrointestinal disorder                              | 1 (<1%)   | 2 (1%)  | 3 (1%)                           |
| Gastroenteritis  | 0 (0%)  | 2 (1%)  | 2 (1%)                           |
| Hemorrhoids  | 1 (<1%)   | 2 (1%)  | 3 (1%)                           |
| Mouth dry  | 1 (<1%)   | 5 (2%)  | 6 (2%)                           |
| Nausea   | 105 (36%)   | 110 (49%)   | 165 (57%)                        |
| Stomatitis   | 19 (7%)   | 20 (9%)   | 36 (13%)                         |
| Vomiting   | 57 (20%)  | 66 (29%)  | 106 (37%)                        |
| <u>Skin and subcutaneous tissue disorders</u>          |   |   |                                  |
| Alopecia   | 199 (69%)   | 124 (55%)   | 208 (72%)                        |
| Dermatitis   | 8 (3%)  | 1 (<1%)   | 9 (3%)                           |
| Dry skin   | 7 (2%)  | 11 (5%)   | 17 (6%)                          |
| Erythema   | 14 (5%)   | 2 (1%)  | 16 (6%)                          |
| Exfoliation dermatitis                                 | 4 (1%)  | 0 (0%)  | 4 (1%)                           |
| Photosensitivity reaction                              | 2 (1%)  | 0 (0%)  | 2 (1%)                           |
| Pigmentation abnormal                                  | 4 (1%)  | 2 (1%)  | 5 (2%)                           |
| Pruritus   | 11 (4%)   | 11 (5%)   | 20 (7%)                          |
| Rash   | 56 (19%)  | 29 (13%)  | 74 (26%)                         |
| Sweating increased                                     | 1 (<1%)   | 2 (1%)  | 3 (1%)                           |
| <u>Musculoskeletal and connective tissue disorders</u> |   |   |                                  |
| Arthralgia   | 7 (2%)  | 14 (6%)   | 17 (6%)                          |
| Back pain  | 2 (1%)  | 3 (1%)  | 5 (2%)                           |
| Musculoskeletal pain                                   | 2 (1%)  | 4 (2%)  | 6 (2%)                           |
| Muscle weakness  | 8 (3%)  | 6 (3%)  | 11 (4%)                          |
| Myalgia  | 3 (1%)  | 7 (3%)  | 9 (3%)                           |
| Myopathy   | 3 (1%)  | 3 (1%)  | 5 (2%)                           |
| <u>Renal and urinary disorders</u>                     |   |   |                                  |
| Dysuria  | 1 (<1%)   | 2 (1%)  | 2 (1%)                           |
| Micturition frequency                                  | 5 (2%)  | 1 (<1%)   | 6 (2%)                           |
| Urinary incontinence                                   | 6 (2%)  | 4 (2%)  | 10 (3%)                          |

**Table 1. Temozolomide and radiotherapy: Treatment-emergent events during concomitant and maintenance treatment**

| <b>Body System</b>  | <b>Temozolomide + concomitant radiotherapy<br/>n=288*<br/>n (%)</b> | <b>Temozolomide maintenance therapy<br/>n=224<br/>n (%)</b> | <b>Total<br/>n=288<br/>n (%)</b> |
|---|---|---|----------------------------------|
| <u>Reproductive system and breast disorders</u>             |   |   |                                  |
| Amenorrhea  | 0 (0%)  | 1 (1%)  | 1 (1%)                           |
| Breast pain   | 0 (0%)  | 1 (1%)  | 1 (1%)                           |
| Impotence   | 1 (1%)  | 0 (0%)  | 1 (1%)                           |
| Menorrhagia   | 0 (0%)  | 1 (1%)  | 1 (1%)                           |
| Vaginal haemorrhage   | 0 (0%)  | 1 (1%)  | 1 (1%)                           |
| Vaginitis   | 0 (0%)  | 1 (1%)  | 1 (1%)                           |
| <u>General disorders and administration site conditions</u> |   |   |                                  |
| Allergic reaction   | 13 (5%)   | 6 (3%)  | 17 (6%)                          |
| Asthenia  | 3 (1%)  | 2 (1%)  | 5 (2%)                           |
| Condition aggravated  | 2 (1%)  | 2 (1%)  | 4 (1%)                           |
| Face edema  | 8 (3%)  | 3 (1%)  | 9 (3%)                           |
| Fatigue   | 156 (54%)   | 137 (61%)   | 205 (71%)                        |
| Fever   | 12 (4%)   | 8 (4%)  | 18 (6%)                          |
| Flushing  | 2 (1%)  | 1 (<1%)   | 3 (1%)                           |
| Hot flushes   | 2 (1%)  | 1 (<1%)   | 2 (1%)                           |
| Pain  | 5 (2%)  | 5 (2%)  | 9 (3%)                           |
| Parosmia  | 2 (1%)  | 0 (0%)  | 2 (1%)                           |
| Radiation injury  | 20 (7%)   | 5 (2%)  | 22 (8%)                          |
| Rigors  | 2 (1%)  | 3 (1%)  | 4 (1%)                           |
| Taste perversion  | 18 (6%)   | 11 (5%)   | 22 (8%)                          |
| Thirst  | 3 (1%)  | 0 (0%)  | 3 (1%)                           |
| Tooth disorder  | 0 (0%)  | 2 (1%)  | 2 (1%)                           |
| Tongue discolouration                                       | 2 (1%)  | 0 (0%)  | 2 (1%)                           |
| <u>Investigation</u>  |   |   |                                  |
| Gamma GT increased  | 4 (1%)  | 0 (0%)  | 4 (1%)                           |
| Hepatic enzymes increased                                   | 3 (1%)  | 1 (<1%)   | 3 (1%)                           |
| SGOT increased  | 3 (1%)  | 0 (0%)  | 3 (1%)                           |
| SGPT increased  | 12 (4%)   | 5 (2%)  | 13 (5%)                          |

\*A patient who was randomised to the RT arm only, received temozolomide + RT

Laboratory results: Myelosuppression, (neutropenia and thrombocytopenia), which are known dose limiting toxicities for most cytotoxic agents, including temozolomide, were observed. When laboratory abnormalities and adverse events were combined across concomitant and maintenance treatment phases, Grade 3 or Grade 4 neutrophil abnormalities including neutropenic events were observed in 8% of the patients. Grade 3 or Grade 4 platelets abnormalities, including thrombocytopenic events were observed in 14% of the patients who received temozolomide.

| <b>Table 2. Grade 3 or Grade 4 Abnormalities Related to Neutrophils and Platelets</b><br><i>Protocol No. P00458</i>   |                     |
|---|---------------------|
|   | <b>Temozolomide</b> |
| <b>Neutrophils</b>  | 8% (24/288)         |
| <b>Platelets</b>  | 14% (39/288)        |
| Includes patients with Grade 3 or 4 abnormalities based on either the lowest observed post-baseline laboratory values (Common Toxicity Criteria) for hematology assessments and/or adverse events related to hematological abnormalities. |                     |

| <b>Table 3. Temozolomide + Radiotherapy: Grade 3/4 Abnormalities During Concomitant and Maintenance Phases Related to Neutrophils and Platelets</b> |                                |                             |
|---|--------------------------------|-----------------------------|
|   | <b>Concomitant Phase n=288</b> | <b>Maintenance n=224</b>    |
| <b>Neutrophil Abnormalities</b>   | <b>13 (5%)<sup>1</sup></b>     | <b>14 (6%)<sup>1</sup></b>  |
| Febrile Neutropenia   | 2 (1%)                         | 3 (1%)                      |
| Neutropenia   | 2 (1%)                         | 5 (2%)                      |
| Lab Only  | 9 (3%) <sup>2</sup>            | 6 (3%)                      |
| <b>Platelet Abnormalities</b>   | <b>12 (4%)<sup>3</sup></b>     | <b>28 (13%)<sup>3</sup></b> |
| Cerebral hemorrhage   | 2 (1%)                         | 0                           |
| Hemorrhage*   | 4 (1%)                         | 3 (1%)                      |
| Thrombocytopenia  | 8 (3%) <sup>--</sup>           | 8 (4%)                      |
| Lab Only  | 2 (1%)                         | 18 (8%)                     |

<sup>1</sup> Three patients reported neutrophil abnormalities in both phases. A total of 24 patients (8%) reported Grade 3/4 neutropenia.

<sup>2</sup> Two of the 9 patients (182 & 194) reported event of neutropenia in Maintenance phase and Lab Only neutropenia in Concomitant Phase and are included in both categories.

<sup>3</sup> One patient reported platelet abnormality in both phases. A total of 39 patients (14%) reported Grade 3/4 platelet abnormalities.

\* All reports of hemorrhage were associated with Grade 3/4 thrombocytopenia

-- One of 8 events of thrombocytopenia was Grade 5 = fatal

### **Malignant Gliomas Showing Recurrence or Progression after Standard Therapy:**

A total of 1030 patients with advanced malignancies, among which 400 recurrent glioma patients, were treated with temozolomide in clinical trials. The most common treatment-related adverse events in the total population analyzed for safety were gastrointestinal disturbances, specifically nausea (43%) and vomiting (36%). These effects were usually Grade 1 or 2 mild to moderate in severity (0 to 5 episodes of vomiting in 24 hours), and were either self-limiting or readily controlled with standard anti-emetic therapy. The incidence of severe nausea and vomiting was 4% each.

The grade 3 or 4 treatment-related hematologic adverse events (defined as those laboratory hematologic events leading to discontinuation, hospitalization, or transfusion) of thrombocytopenia, neutropenia, and anemia, occurred in 9%, 3%, and 3% of the total population analyzed for safety (1030 patients), respectively. In the recurrent glioma population (400 patients), these events occurred in 9%, 4%, and 1% of patients, respectively.

Myelosuppression was predictable (typically within the first 2 to 4 cycles with platelet and neutrophil nadirs between Days 21 to 28) and recovery was rapid, usually within 2 weeks. Myelosuppression was not cumulative. Pancytopenia and leukopenia have been reported. Lymphopenia has been commonly reported.

| <b>Table 4. Treatment-related Grade 3 and 4 Adverse Events for All Cycles – Recurrent Glioma Population</b> |   |  |
|---|---|--|
| <b>Body System/Adverse Event</b>  | <b>Number (%) of Patients; N=400</b>                          |  |
|   | <b>Grade 3 Adverse Events Reported in At Least 2 Patients</b> | <b>Grade 4 Adverse Events Reported in All Patients</b> |
| <b>No. of Subjects with any AE</b>  | <b>87 (22%)</b>   | <b>26 (7%)</b>   |
| <b>Body as a Whole, General</b>   | <b>25 (6%)</b>  | <b>2 (&lt;1%)</b>                                      |
| Asthenia  | 6 (2%)  | 2 (<1%)  |
| Fatigue   | 9 (2%)  | 0  |
| Fever   | 2 (<1%)   | 0  |
| Headache  | 6 (2%)  | 0  |
| <b>Central and Peripheral Nervous System</b>  | <b>11 (3%)</b>  | <b>1 (&lt;1%)</b>                                      |
| Confusion   | 2 (<1%)   | 0  |
| Consciousness decreased   | 0   | 1 (<1%)  |
| Convulsions   | 2 (<1%)   | 0  |
| Hemiparesis   | 2 (<1%)   | 0  |
| Paresis   | 2 (<1%)   | 0  |
| Transient ischemic attack   | 0   | 1 (<1%)  |
| <b>Gastrointestinal System</b>  | <b>33 (8%)</b>  | <b>1 (&lt;1%)</b>                                      |
| Abdominal pain  | 2 (<1%)   | 0  |
| Constipation  | 2 (<1%)   | 0  |
| Dehydration   | 2 (<1%)   | 0  |
| Diarrhea  | 2 (<1%)   | 0  |
| Nausea  | 18 (5%)   | 0  |
| Vomiting  | 14 (4%)   | 1 (<1%)  |
| <b>Metabolic and Nutritional</b>  | <b>2 (&lt;1%)</b>   | <b>0</b>   |
| Hyperglycemia   | 2 (<1%)   | 0  |
| <b>Platelet, Bleeding &amp; Clotting</b>  | <b>17 (4%)</b>  | <b>19 (5%)</b>   |
| Thrombocytopenia  | 17 (4%)   | 19 (5%)  |
| <b>Psychiatric Disorders</b>  | <b>3 (1%)</b>   | <b>0</b>   |
| Somnolence  | 3 (1%)  | 0  |

| <b>Table 4. Treatment-related Grade 3 and 4 Adverse Events for All Cycles – Recurrent Glioma Population</b>  |   |  |
|--|---|--|
| <b>Body System/Adverse Event</b>   | <b>Number (%) of Patients; N=400</b>                          |  |
|  | <b>Grade 3 Adverse Events Reported in At Least 2 Patients</b> | <b>Grade 4 Adverse Events Reported in All Patients</b> |
| <b>No. of Subjects with any AE</b>   | <b>87 (22%)</b>   | <b>26 (7%)</b>   |
| <b>Red Blood Cells</b>   | <b>3 (1%)</b>   | <b>3 (1%)</b>  |
| Anemia   | 2 (<1%)   | 2 (<1%)  |
| Pancytopenia   | 1 (<1%)   | 1 (<1%)  |
| <b>Respiratory System</b>  | <b>3 (1%)</b>   | <b>1 (&lt;1%)</b>                                      |
| Pneumonia  | 2 (<1%)   | 0  |
| Pulmonary Infection  | 1 (<1%)   | 1 (<1%)  |
| <b>Vascular (extracardiac)</b>   | <b>1 (&lt;1%)</b>   | <b>5 (1%)</b>  |
| Embolism pulmonary   | 0   | 1 (<1%)  |
| Hemorrhage intracranial  | 0   | 1 (<1%)  |
| Hemorrhage, NOS  | 0   | 2 (<1%)  |
| Purpura  | 1 (<1%)   | 0  |
| Thrombophlebitis, deep   | 0   | 2 (<1%)  |
| <b>White Cell and RES</b>  | <b>14 (4%)</b>  | <b>10 (3%)</b>   |
| Leukopenia   | 10 (3%)   | 6 (2%)   |
| Neutropenia  | 7 (2%)  | 7 (2%)   |
| Only lab abnormalities that led to discontinuation, hospitalization or transfusion were reported as AEs and are included in this table. A patient is counted only once if >1 occurrence of a specific AE. Body system total numbers and percentages reflect all patients reporting any AE within that body system. |   |  |

Among all patients treated with temozolomide, changes in hematologic laboratory data from Grade 0 to 2 at Baseline to Grade 3 to 4 during treatment (thrombocytopenia, neutropenia, and anemia) occurred in 19%, 17% and 7% of the total population analyzed for safety, respectively and in 20%, 14%, and 5% of recurrent glioma patients respectively.

| <b>Table 5. Changes in Hematologic laboratory Data from Grade 0 to 2 at Baseline to Grade 3 to 4 During Treatment (Overall and Recurrent Glioma Population)</b> |  |  |
|---|--|--|
|   | <b>Overall Population (N=1030)<sup>a</sup></b> | <b>Recurrent Glioma Population (N=400)<sup>a</sup></b> |
| Platelets   | 19% (180/950)                                  | 20% (79/394)   |
| Neutrophils   | 17% (154/907)                                  | 14% (52/366)   |
| Hemoglobin  | 7% (63/969)                                    | 5% (20/397)  |
| a: Percents were based on the number of patients with data available at baseline and at least one subsequent visit for each parameter.                          |  |  |

In a population pharmacokinetics analysis of clinical trial experience there were 101 female and 169 male subjects for whom nadir neutrophil counts were available and 110 female and 174 male subjects for whom nadir platelet counts were available. There were higher rates of Grade 4 neutropenia (ANC <500 cells/mcL), 12% versus 5%, and thrombocytopenia (<20,000 cells/mcL), 9% versus 3%, in women vs. men in the first cycle of therapy. In a 400-subject recurrent glioma data set, Grade 4 neutropenia occurred in 8% of female versus 4% of male subjects and Grade 4 thrombocytopenia in 8% of female vs. 3% of male subjects in the first cycle of therapy. In a study of 288 subjects with newly diagnosed glioblastoma multiforme, Grade 4 neutropenia occurred in 3% of female vs 0% of male subjects and Grade 4 thrombocytopenia in 1% of female vs 0% of male subjects in the first cycle of therapy.

Other adverse events reported frequently in the total population analysed for safety included fatigue (22%), constipation (17%), and headache (14%). Anorexia (11%), diarrhea (8%), rash, fever, asthenia, and somnolence (6% each) were also reported. Less common adverse events (2% to 5%) and in descending order of frequency, were abdominal pain, pain, dizziness, weight decrease, malaise, dyspnea, alopecia, rigors, pruritus, dyspepsia, taste perversion, paresthesia and petechiae.

The table below shows the treatment-related adverse events reported in  $\geq 2\%$  of patients in clinical trials involving a total of 400 glioma patients treated with temozolomide.

| <b>Table 6. Treatment-Related Adverse Events Reported in <math>\geq 2\%</math> of recurrent Glioma Patients</b> |                               |
|---|-------------------------------|
| <b>Body System/Adverse Event</b>  | <b>Number (%) of Patients</b> |
| <b>No. of Subjects with any AE</b>  | <b>304 (76%)</b>              |
| <u>Body as a Whole, General</u>   | <u>154 (39%)</u>              |
| Fatigue   | 90 (23%)                      |
| Headache  | 42 (11%)                      |
| Fever   | 15 (4%)                       |
| Asthenia  | 19 (5%)                       |
| Pain  | 10 (3%)                       |
| Malaise   | 7 (2%)                        |
| Rigors  | 2 (<1%)                       |
| Weight decrease   | 4 (1%)                        |
| <u>Central and Peripheral Nervous System</u>  | <u>52 (13%)</u>               |
| Convulsions   | 10 (3%)                       |
| Dizziness   | 9 (2%)                        |
| Paresthesia   | 6 (2%)                        |
| <u>Gastrointestinal System</u>  | <u>230 (58%)</u>              |
| Nausea  | 162 (41%)                     |
| Vomiting  | 137 (34%)                     |
| Constipation  | 60 (15%)                      |
| Anorexia  | 35 (9%)                       |
| Diarrhea  | 28 (7%)                       |
| Abdominal pain  | 13 (3%)                       |



| <b>Table 6. Treatment-Related Adverse Events Reported in <math>\geq 2\%</math> of recurrent Glioma Patients</b> |                               |
|---|-------------------------------|
| <b>Body System/Adverse Event</b>  | <b>Number (%) of Patients</b> |
| <b>No. of Subjects with any AE</b>  | <b>304 (76%)</b>              |
| Dyspepsia   | 9 (2%)                        |
| <u>Musculo-skeletal System</u>  | 8 (2%)                        |
| Myalgia   | 3 (1%)                        |
| <u>Platelet, Bleeding &amp; Clotting</u>  | 35 (9%)                       |
| Thrombocytopenia  | 35 (9%)                       |
| <u>Psychiatric Disorders</u>  | 37 (9%)                       |
| Somnolence  | 18 (4%)                       |
| Depression  | 4 (1%)                        |
| Insomnia  | 6 (2%)                        |
| <u>Red Blood Cells</u>  | 10 (2%)                       |
| Anemia  | 8 (2%)                        |
| Pancytopenia  | 2 (<1%)                       |
| <u>Resistance Mechanism</u>   | 31 (8%)                       |
| Candidiasis Oral  | 9 (2%)                        |
| <u>Respiratory System</u>   | 27 (7%)                       |
| Dyspnea   | 6 (2%)                        |
| <u>Special Senses</u>   | 4 (1%)                        |
| Taste Perversion  | 4 (1%)                        |
| <u>Skin and Appendages</u>  | 73 (18%)                      |
| Rash  | 21 (5%)                       |
| Alopecia  | 15 (4%)                       |
| Pruritus  | 12 (3%)                       |
| Petechiae   | 14 (4%)                       |
| <u>White Cell and RES</u>   | 21 (5%)                       |
| Neutropenia   | 14 (4%)                       |
| Leukopenia  | 15 (4%)                       |

Only lab abnormalities that led to discontinuation, hospitalization or transfusion were reported as AEs and are included in this table. A patient is counted only once if >1 occurrence of a specific AE. Body system total numbers and percentages reflect all patients reporting any AE within that body system.

In the phase II malignant recurrent glioma trials, serious adverse events were reported in 278 (70%) patients treated with temozolomide. The majority of serious adverse events were hospitalizations due to disease progression or disease-related complications, and were unrelated to temozolomide. Hematologic toxicity, usually grade 3 or 4 thrombocytopenia or neutropenia, was the most common serious adverse event. The majority of these reports were at the 200 mg/m<sup>2</sup>/day dose level, and most cases resolved with one dose level reduction. Non-hematologic serious adverse events were uncommon.

Within 30 days of the last dose of temozolomide, forty recurrent glioma patients died, the majority due to disease progression or disease-related complications. Two deaths were judged as possibly related to the administration of temozolomide (grade 4 intratumoral hemorrhage with grade 3 cerebral edema in one patient and grade 4 cerebral ischemia in one patient).

### **Post-Market Adverse Drug Reactions**

The following adverse events have been reported from post-marketing experience:

- Allergic reactions, including anaphylaxis
- Erythema multiforme, toxic epidermal necrolysis (TEN), Stevens-Johnson syndrome (SJS)
- Opportunistic infections including *Pneumocystis carinii* pneumonia (PCP) and primary and reactivated cytomegalovirus (CMV) infection, and reactivation of hepatitis B infection, including some cases with fatal outcomes (see **WARNINGS AND PRECAUTIONS**)
- Cases of herpes simplex encephalitis, including cases with fatal outcomes
- Myelodysplastic syndrome (MDS) and secondary malignancies including myeloid leukemia
- Pancytopenia, which may result in aplastic anemia has been reported, and in some cases has resulted in a fatal outcome
- Interstitial pneumonitis/pneumonitis and pulmonary fibrosis
- Hepatotoxicity including elevations of liver enzymes, hyperbilirubinemia, cholestasis and hepatitis. Hepatic injury, including fatal hepatic failure, has been reported (see **WARNINGS AND PRECAUTIONS**).
- Diabetes insipidus.

### **DRUG INTERACTIONS**

#### **Drug interactions with oral temozolomide:**

Antiemetic therapy may be administered prior to or following administration of APO-TEMOZOLOMIDE

Administration of temozolomide with ranitidine or with food did not result in clinically significant alterations in the extent of absorption of temozolomide. Analyses of data obtained from population pharmacokinetics in the phase II studies demonstrated that co-administration of dexamethasone, prochlorperazine, phenytoin, carbamazepine, ondansetron, H<sub>2</sub>-receptor antagonists, or phenobarbital did not alter the clearance of temozolomide. Co-administration with valproic acid was associated with a small but statistically significant decrease in clearance of temozolomide.

No studies have been conducted to determine the effect of temozolomide on the metabolism or elimination of other medicinal products. However, since temozolomide does not require hepatic metabolism, has a short half-life, and exhibits low protein binding, it is unlikely that it would affect the pharmacokinetics of other medicinal products.

The combination of temozolomide with other chemotherapeutic agents has not been fully evaluated. Combination with other alkylating agents is likely to result in increased myelosuppression.

#### **Drug-Food Interactions**

Temozolomide interactions with food have not been established.

#### **Drug-Herb Interactions**

Temozolomide interactions with herbal products have not been established.

**Drug-Laboratory Interactions**

Temozolomide interactions with laboratory tests have not been established.

**DOSAGE AND ADMINISTRATION**

Do NOT open, divide, crush or chew the capsule. Swallow the capsule whole with a glass of water.

**Recommended Dose and Dosage Adjustment**

**Adults Patients with Newly Diagnosed Glioblastoma Multiforme:**

**Concomitant Phase**

APO-TEMOZOLOMIDE is administered at a dose of 75 mg/m<sup>2</sup> daily for 42 days concomitant with radiotherapy (60 Gy administered in 30 fractions) followed by maintenance APO-TEMOZOLOMIDE for 6 cycles. No dose reductions are recommended; however, dose interruptions may occur based on patient tolerance. The APO-TEMOZOLOMIDE dose can be continued throughout the 42 day concomitant period up to 49 days if all of the following conditions are met: absolute neutrophil count  $\geq 1.5 \times 10^9/L$ ; platelet count  $\geq 100 \times 10^9/L$ ; common toxicity criteria (CTC) non-hematological toxicity Grade  $\leq 1$  (except for alopecia, nausea and vomiting). During treatment a complete blood count should be obtained weekly. APO-TEMOZOLOMIDE dosing should be interrupted or discontinued during concomitant phase according to the hematological and non-hematological toxicity criteria as noted in Table 7.

***Table 7. APO-TEMOZOLOMIDE Dosing Interruption or Discontinuation During Concomitant Radiotherapy and APO-TEMOZOLOMIDE***

| <b>Toxicity</b>  | <b>APO-TEMOZOLOMIDE Interruption<sup>a</sup></b> | <b>APO-TEMOZOLOMIDE Discontinuation</b> |
|--|--|---|
| Absolute Neutrophil Count  | $\geq 0.5$ and $< 1.5 \times 10^9/L$             | $< 0.5 \times 10^9/L$                   |
| Platelet Count   | $\geq 10$ and $< 100 \times 10^9/L$              | $< 10 \times 10^9/L$                    |
| CTC Non-hematological Toxicity (except for alopecia, nausea, vomiting)   | CTC Grade 2                                      | CTC Grade 3 or 4                        |
| a: Treatment with concomitant APO-TEMOZOLOMIDE could be continued when all of the following conditions were met: absolute neutrophil count $\geq 1.5 \times 10^9/L$ ; platelet count $\geq 100 \times 10^9/L$ ; CTC non-hematological toxicity Grade $\leq 1$ (except for alopecia, nausea, vomiting). CTC = Common Toxicity Criteria. |  |   |

**Maintenance Phase**

Four weeks after completing the APO-TEMOZOLOMIDE + RT (Radiotherapy) phase, APO-TEMOZOLOMIDE is administered for an additional 6 cycles of maintenance treatment. Dosage in Cycle 1 (maintenance) is 150 mg/m<sup>2</sup> once daily for 5 days followed by 23 days without treatment.

At the start of Cycle 2, the dose is escalated to 200 mg/m<sup>2</sup>, if the CTC non-hematologic toxicity for Cycle 1 is Grade ≤2 (except for alopecia, nausea and vomiting), absolute neutrophil count (ANC) is ≥1.5 x 10<sup>9</sup>/L, and the platelet count is ≥100 x 10<sup>9</sup>/L. If the dose was not escalated at Cycle 2, escalation should not be done in subsequent cycles. The dose remains at 200 mg/m<sup>2</sup> per day for the first 5 days of each subsequent cycle except if toxicity occurs. Dose reductions during the maintenance phase should be applied according to Tables 8 and 9.

During treatment a complete blood count should be obtained on day 22 (21 days after the first dose of APO-TEMOZOLOMIDE). The APO-TEMOZOLOMIDE dose should be reduced or discontinued according to Table 9.

| <b>Table 8. APO-TEMOZOLOMIDE Dose Levels for Maintenance Treatment</b> |                                    |  |
|--|------------------------------------|--|
| <b>Dose Level</b>  | <b>Dose (mg/m<sup>2</sup>/day)</b> | <b>Remarks</b>                                   |
| -1   | 100                                | Reduction for prior toxicity                     |
| 0  | 150                                | Dose during Cycle 1                              |
| 1  | 200                                | Dose during Cycles 2 to 6 in absence of toxicity |

| <b>Table 9. APO-TEMOZOLOMIDE Dose Reduction or Discontinuation During Maintenance Treatment</b> |  |                                     |
|---|--|-------------------------------------|
| <b>Toxicity</b>   | <b>Reduce APO-TEMOZOLOMIDE by 1 Dose Level<sup>a</sup></b> | <b>Discontinue APO-TEMOZOLOMIDE</b> |
| Absolute Neutrophil Count   | <1.0 x 10 <sup>9</sup> /L                                  | See footnote b                      |
| Platelet Count  | <50 x 10 <sup>9</sup> /L                                   | See footnote b                      |
| CTC Non-hematological Toxicity (except for alopecia, nausea, vomiting)                          | CTC Grade 3  | CTC Grade 4 <sup>b</sup>            |

a: APO-TEMOZOLOMIDE dose levels are listed in Table 8.  
b: APO-TEMOZOLOMIDE is to be discontinued if dose reduction to <100 mg/m<sup>2</sup> is required or if the same Grade 3 non-hematological toxicity (except for alopecia, nausea, vomiting) recurs after dose reduction.  
CTC = Common Toxicity Criteria.

**Malignant Gliomas Showing Recurrence or Progression after Standard Therapy:**

Adult patients: In patients previously untreated with chemotherapy, APO-TEMOZOLOMIDE is administered at a dose of 200 mg/m<sup>2</sup> once daily for 5 days per 28-day cycle. For patients previously treated with chemotherapy, the initial dose is 150 mg/m<sup>2</sup> once daily for 5 days, to be increased in the second cycle to 200 mg/m<sup>2</sup> once daily for 5 days, providing there is no hematologic toxicity (see **WARNINGS AND PRECAUTIONS**).

In the reference controlled trial of GBM, the majority of patients treated with Temozolomide (90%) received more than one cycle and 22% of patients received 6 or more cycles. These patients

received a total of 484 cycles of Temozolomide in total; 60% of cycles at 200 mg/m<sup>2</sup>/day and 36% at 150 mg/m<sup>2</sup>/day. In the single arm AA trial, 93% of patients received more than one cycle and 25% of patients continued on study for 12 months or greater. Eighty-eight percent of patients were receiving either their initial dose or a higher dose at the last cycle. However, limited experience is available on the prolonged use of Temozolomide in this patient population.

APO-TEMOZOLOMIDE therapy can be continued until disease progression.

### Administration

Prior to dosing and during treatment, proper hematologic monitoring must be performed (see **WARNINGS AND PRECAUTIONS**) to ensure that the following laboratory parameters are met: absolute neutrophil count (ANC)  $\geq 1.5 \times 10^9/L$  and platelets  $\geq 100 \times 10^9/L$ . If the ANC falls to  $< 1.0 \times 10^9/L$  or the platelet count is  $< 50 \times 10^9/L$  during any cycle, the next cycle should be reduced one dose level. Dose levels include 100 mg/m<sup>2</sup>, 150 mg/m<sup>2</sup>, and 200 mg/m<sup>2</sup>. The lowest recommended dose is 100 mg/m<sup>2</sup>. Dose modification for APO-TEMOZOLOMIDE should be based on toxicities according to nadir ANC or platelet counts.

Since women taking Temozolomide were reported to have a higher incidence of grade 4 neutropenia and thrombocytopenia than men in the first cycle of therapy, they must be closely monitored for abnormal neutrophil and platelet counts.

APO-TEMOZOLOMIDE should be administered in the fasting state, at least one hour before a meal. Antiemetic therapy may be administered prior to or following administration of APO-TEMOZOLOMIDE. If vomiting occurs after the dose is administered, a second dose should not be administered.

Store APO-TEMOZOLOMIDE capsules at room temperature 15°C to 30°C. Protect from moisture.

### **OVERDOSAGE**

Doses of 500, 750, 1,000, and 1,250 mg/m<sup>2</sup> (total dose per cycle over 5 days) have been evaluated clinically in patients. Dose-limiting toxicity was hematological and was reported at any dose but is expected to be more severe at higher doses. An overdose of 2,000 mg per day for 5 days was taken by one patient and the adverse events reported were pancytopenia, pyrexia, multi-organ failure and death. There are reports of patients who have taken more than 5 consecutive days of treatment (up to 64 consecutive days) with adverse events reported including bone marrow suppression, with or without infection, in some cases severe and prolonged and resulting in death. In the event of an overdose, hematologic evaluation is needed. Supportive measures should be provided as necessary.

|   |
|---|
| For management of a suspected drug overdose, contact your regional Poison Control Centre immediately. |
|---|

## **ACTION AND CLINICAL PHARMACOLOGY**

### **Mechanism of Action**

Temozolomide is an imidazotetrazine alkylating agent with antitumor activity that can be used orally. It undergoes rapid chemical conversion in the systemic circulation at physiologic pH to the active compound, MTIC. The cytotoxicity of MTIC is thought to be due primarily to alkylation at the O<sup>6</sup> position of guanine with additional alkylation also occurring at the N<sup>7</sup> position. Cytotoxic lesions that develop subsequently are thought to involve aberrant repair of the methyl adduct.

After oral administration to adult patients, temozolomide is absorbed rapidly with peak plasma concentrations reached as early as 20 minutes post-dose (mean T<sub>max</sub> range between 0.5 and 1.5 hours).

Plasma concentrations are dose-dependent, while plasma clearance, volume of distribution and half-life are independent of dose. Temozolomide demonstrates low protein binding (10% to 20%), and thus is not expected to interact with highly protein bound agents. After oral administration of <sup>14</sup>C labelled temozolomide, mean fecal elimination of <sup>14</sup>C over 7 days post-dose was 0.8% indicating complete absorption. Following oral administration, approximately 5% to 10% of the dose is recovered unchanged in the urine over 24 hours, and the remainder excreted as AIC (4-amino-5-imidazole-carboxamide hydrochloride) or unidentified polar metabolites.

Analysis of population based pharmacokinetics of temozolomide revealed that plasma temozolomide clearance was independent of age, renal function, hepatic function, or tobacco use.

Pediatric patients (<18 years old and >3 years old) had a higher area under the curve (AUC) than adult patients; however, the maximum tolerated dose (MTD) was 1000 mg/m<sup>2</sup> per cycle both in children and in adults.

## **STORAGE AND STABILITY**

Store APO-TEMOZOLOMIDE capsules at room temperature 15°C to 30°C. Protect from moisture.

## **SPECIAL HANDLING INSTRUCTIONS**

APO-TEMOZOLOMIDE Capsules must not be opened or chewed, but are to be swallowed whole with a glass of water. If a capsule becomes damaged, avoid contact of the powder contents with skin or mucous membrane. In the case of accidental contact with skin or mucous membrane, flush with water.

KEEP OUT OF REACH AND SIGHT OF CHILDREN.

## DOSAGE FORMS, COMPOSITION AND PACKAGING

**APO-TEMOZOLOMIDE 5 mg Capsules:** Each white opaque body, green opaque cap, hard gelatin capsule, imprinted “APO TEM 5” in black ink, white to light pink powder fill, contains 5 mg of temozolomide. Available in bottles of 5 and 20, and unit dose sachets of 1 capsule (5 or 20 sachets per box).

**APO-TEMOZOLOMIDE 20 mg Capsules:** Each white opaque body, gold opaque cap, hard gelatin capsule, imprinted “APO TEM 20” in black ink, white to light pink powder fill, contains 20 mg of temozolomide. Available in bottles of 5 and 20, and unit dose sachets of 1 capsule (5 or 20 sachets per box).

**APO-TEMOZOLOMIDE 100 mg Capsules:** Each white opaque body, flesh opaque cap, hard gelatin capsule, imprinted “APO TEM 100” in black ink, white to light pink powder fill, contains 100 mg of temozolomide. Available in bottles of 5 and 20, and unit dose sachets of 1 capsule (5 or 20 sachets per box).

**APO-TEMOZOLOMIDE 140 mg Capsules:** Each white opaque body, blue transparent cap, hard gelatin capsule, imprinted “APO TEM 140” in black ink, white to light pink powder fill, contains 140 mg of temozolomide. Available in bottles of 5 and 20, and unit dose sachets of 1 capsule (5 or 20 sachets per box).

**APO-TEMOZOLOMIDE 180 mg Capsules:** Each white opaque body, swedish orange opaque cap, hard gelatin capsule, imprinted “APO TEM 180” in black ink, white to light pink powder fill, contains 180 mg of temozolomide. Available in bottles of 5 and 20, and unit dose sachets of 1 capsule (5 or 20 sachets per box).

**APO-TEMOZOLOMIDE 250 mg Capsules:** Each white opaque body, white opaque cap, hard gelatin capsule, imprinted “APO TEM 250” in black ink, white to light pink powder fill, contains 250 mg of temozolomide. Available in bottles of 5 and unit dose sachets of 1 capsule (5 or 20 sachets per box).

In addition to the active ingredient, temozolomide, each capsule also contains the non-medicinal ingredients: colloidal silicon dioxide, lactose anhydrous, sodium starch glycolate, stearic acid and tartaric acid.

The empty capsule shells contain the non-medicinal ingredients FD&C Blue No. 2 (5 mg, 140 mg), gelatin, iron oxide red (100 mg, 180 mg), iron oxide yellow (5 mg, 20 mg, 180 mg), talc and titanium dioxide.

The capsule shells are imprinted with black edible ink containing ammonium hydroxide, iron oxide black, isopropyl alcohol, N-Butyl alcohol, propylene glycol and shellac glaze.

## PART II: SCIENTIFIC INFORMATION

### PHARMACEUTICAL INFORMATION

#### Drug Substance


Proper name: Temozolomide

Chemical name: 3,4-Dihydro-3-methyl-4-oxoimidazo[5,1-*d*]-1,2,3,5-tetrazine-8-carboxamide; Imidazo[5,1-*d*]-1,2,3,5-tetrazine-8-carboxamide, 3,4-dihydro-3-methyl-4-oxo

Molecular formula: C<sub>6</sub>H<sub>6</sub>N<sub>6</sub>O<sub>2</sub>

Molecular weight: 194.15 g/mol

Structural formula:



Physicochemical properties:

*Physical form:*

White to light pink/light tan powder

*Solubility:*

Temozolomide is soluble in dimethylsulfoxide; sparingly soluble in water; practically insoluble in toluene.

*pKa/pH :*

Temozolomide contains no functional groups that can be protonated or deprotonated between pH 1 and pH 13, and therefore, does not have a dissociation constant (pKa) in this pH range. The pH of a 10 mg/mL aqueous dispersion of temozolomide is about 5.8.

*Partition coefficient:*

Temozolomide partitions primarily into the organic phase and the pH of the aqueous phase has little, if any effect, on the partition coefficient:



| <i>Solvent</i>                  | <i>Partition Coefficient<br/>(octanol/aqueous)</i> |
|---------------------------------|--|
| water                           | 22.4   |
| phosphate buffer pH 7.0 (0.1 M) | 22.0   |
| 0.1N HCl                        | 20.8   |

*Melting point:*

Temozolomide does not show a true melting point but undergoes decomposition from about 182°C to 200°C.

*Aqueous pH solubility:*

The aqueous solubility of Temozolomide as a function of pH determined at 37°C is displayed in the table below:

| <i>Solvent Media*</i>            | <i>Final<br/>pH</i> | <i>Solubility<br/>(mg/mL)</i> | <i>Highest Dose<br/>(250 mg)/solubility<br/>volume</i> |
|----------------------------------|---------------------|-------------------------------|--|
| 0.1N HCl                         | 1.4                 | 7.0                           | 36 mL  |
| pH 2.5 Phosphate Buffer (0.05 M) | 2.6                 | 6.7                           | 38 mL  |
| pH 3.5 Phosphate Buffer (0.05 M) | 3.8                 | 6.5                           | 39 mL  |
| pH 4.5 Phosphate Buffer (0.05 M) | 5.0                 | 5.1                           | 50 mL  |
| Water                            | 5.9                 | 5.3                           | 46 mL  |

\*At pH >6.0, temozolomide degrades to a large extent, preventing the accurate determination of solubility.

## CLINICAL TRIALS

### Comparative Bioavailability Studies

A randomized, single-dose, multiple centre, double blinded, two period crossover bioavailability study, conducted under fasting conditions, was performed on an adult patient population with diagnosed glioblastoma multiforme or refractory anaplastic astrocytoma. The following table summarizes T/R ratios and confidence intervals obtained from 22 patients. The rate and extent of absorption of temozolomide was measured and compared following a single oral dose (1 x 250 mg capsule) of Apo-Temozolomide (temozolomide) 250 mg capsule (Apotex Inc.) and Temodal<sup>®</sup> (temozolomide) 250 mg capsule (Merck Canada Inc.).


| Summary Table of the Comparative Bioavailability Data<br>Temozolomide<br>(A single 250 mg dose: 1 x 250 mg capsule)<br>From Measured Data/Fasting Conditions<br>Geometric Mean <sup>#</sup><br>Arithmetic Mean (CV%) |              |                        |                              |                             |
|--|--------------|------------------------|------------------------------|-----------------------------|
| Parameter  | Test*        | Reference <sup>†</sup> | Ratio of Geometric Means (%) | 90% Confidence Interval (%) |
| AUC <sub>t</sub> (ng•h/mL)   | 21367.2      | 21247.2                | 100.6                        | 96.7 – 104.6                |
|  | 22014.6 (25) | 21392.7 (25)           |                              |                             |
| AUC <sub>inf</sub> (ng•h/mL)   | 22020.6      | 21969.8                | 100.2                        | 96.6 – 104.0                |
|  | 22676.9 (25) | 22079.0 (24)           |                              |                             |
| C <sub>max</sub> (ng/mL)   | 6714.2       | 6795.1                 | 98.8                         | 85.6 – 114.1                |
|  | 7189.6 (39)  | 7052.1 (35)            |                              |                             |
| T <sub>max</sub> <sup>§</sup> (h)  | 1.36 (48)    | 1.29 (62)              |                              |                             |
| T <sub>half</sub> <sup>§</sup> (h)   | 1.78 (7)     | 1.85 (14)              |                              |                             |
| * Apo-Temozolomide (temozolomide) 250 mg capsules (Apotex Inc.)  |              |                        |                              |                             |
| † Temodal <sup>®</sup> (temozolomide) 250 mg capsules (Merck Canada Inc.) was purchased in Canada.   |              |                        |                              |                             |
| # For unbalanced treatment sequence, results are based on Least Squares Means (LSM).   |              |                        |                              |                             |
| § Expressed as arithmetic means (CV%) only.  |              |                        |                              |                             |

### Newly Diagnosed Glioblastoma Multiforme

Five hundred seventy three subjects were randomized to receive either temozolomide + Radiotherapy (RT) (n=287) or RT alone (n=286). Patients in the temozolomide + RT arm received concomitant temozolomide (75 mg/m<sup>2</sup>) once daily, starting the first day of RT until the last day of RT, for 42 days (with a maximum of 49 days). This was followed by maintenance temozolomide (150 or 200 mg/m<sup>2</sup>) on day 1 to 5 of every 28-day cycle for 6 cycles, starting 4 weeks after the end of RT. Patients in the control arm received RT only. *Pneumocystis carinii* pneumonia (PCP) prophylaxis was required during RT and combined temozolomide therapy, and was to continue until recovery of lymphopenia to grade <1.

Temozolomide was administered as salvage therapy in the follow-up phase in 161 patients of the 282 (57%) in the RT alone arm, and 62 patients of the 277 (22%) in the temozolomide + RT arm.

The hazard ratio (HR) for overall survival was 1.59 (95% CI for HR=1.33-1.91) with a log-rank  $P < 0.0001$  in favour of the temozolomide arm. The estimated probability of surviving 2 years or more (26% vs 10%) is higher for the RT + temozolomide arm. The addition of concomitant and maintenance temozolomide to radiotherapy in the treatment of patients with newly diagnosed glioblastoma multiforme demonstrated a statistically significant improved overall survival compared with radiotherapy alone (Figure 1).



**Figure 1** Kaplan-Meier Curves for Overall Survival (Intent To Treat) ITT Population

### Malignant Gliomas Showing Recurrence or Progression after Standard Therapy:

Consistent patient selection criteria were used in the 3 phase II studies. In all trials, adult patients  $\geq 18$  years of age with histologically confirmed supratentorial GBM or AA at first relapse, a baseline Karnofsky performance status (KPS) of at least 70, and a life expectancy  $> 12$  weeks were eligible. Patients had unequivocal evidence of tumor recurrence or progression (first relapse) and evaluable enhancing residual disease. They failed a conventional course of radiation therapy for initial disease and no more than one prior regimen of adjuvant chemotherapy (with either a single

agent or a regimen containing a nitrosourea).

In the phase II studies, consistent criteria based on neuroimaging and clinical neurologic examination were used to define overall response and to determine disease progression for the progression-free survival analysis. Objective assessments of overall response were based upon tumor assessments interpreted in light of steroid use and, to a lesser extent, neurologic status. Overall response was based on the following:

- Complete response (CR): Disappearance of all enhancing tumor (measurable or non-measurable) on consecutive magnetic resonance imaging (MRI) scans at least one month apart, off steroids except for physiologic doses which may have been required following prolonged therapy and neurologically stable or improved.
- Partial response (PR): For patients with lesions which were either all measurable or all non-measurable, greater than or equal to a 50% reduction (<100%) in the sum of the products of the largest perpendicular diameters of contrast enhancement for all measurable lesions or +2 rating (definitely better) for all non-measurable lesions on consecutive MRI scans at least one month apart, steroids stable for 7 days prior to each scan at the same dose administered at the time of the previous scan or at a reduced dose, and neurologically stable or improved. No new lesions could arise.
- Progressive disease (PD): Greater than or equal to a 25% increase in size of the product of the largest perpendicular diameters of contrast enhancement for any measurable lesions or -2 rating (definitely worse) for any non-measurable lesions or any new tumor on MRI scans, steroids stable for 7 days prior to each scan at the same dose administered at the time of the previous scan or at an increased dose, with or without neurologic progression. The investigator had to carefully exclude non-tumor-related causes of clinical or radiological worsening (i.e. pseudoprogression).
- Stable disease (SD): All other situations.

Temozolomide has been shown to be effective in prolonging progression-free survival and maintaining or improving health-related quality of life (HQL) in adult patients with recurrent high grade glioma. Both patients with anaplastic astrocytoma (AA) and glioblastoma multiforme (GBM) experienced clinically meaningful efficacy and HQL benefits.

In an open-label, active-reference study in which patients received either temozolomide or procarbazine, temozolomide demonstrated efficacy in GBM patients at first relapse based on improvements in progression-free survival, event-free survival and overall survival relative to the reference agent, procarbazine. This study was not designed nor powered to make statistically valid comparisons between the two drugs.

Two hundred ten patients were determined by central review as having histologically confirmed GBM or gliosarcoma and comprise the eligible histology population. In the temozolomide group, the median age was 52 years and 69% were male. Karnofsky performance status was  $\geq 80$  in 70% of patients. At the time of initial diagnosis, 86% of patients in the temozolomide group had undergone surgical resection, with all patients subsequently receiving radiation therapy. Chemotherapy was administered in 65% of patients in the temozolomide group. The median time from initial diagnosis

to first relapse was 7.0 months for temozolomide patients. At first relapse, 20% of patients had surgical resection.

Results from this controlled trial are summarized in the table below:

**Efficacy Results: Controlled Study**

| Study   | Histology | No. Pts. | Drug Study | PFS at 6 mos (95% CI) | Median PFS (Months) | Median OS (Months) | 6-month Survival Rate |
|---------|-----------|----------|------------|-----------------------|---------------------|--------------------|-----------------------|
| C94-091 | GBM       | 112      | TMZ        | 21%<br>(13%–29%)      | 2.99                | 7.34               | 60%                   |
| C94-091 | GBM       | 113      | PROC       | 8%<br>(3%–15%)        | 1.97                | 5.82               | 44%                   |

PFS: Progression-free survival    TMZ: Temozolomide  
 CI: Confidence Intervals        PROC: Procarbazine  
 OS: Overall Survival

Objective response (partial response; PR) as determined by Gd-MRI scan after independent central review was achieved in 5% (6/112) of temozolomide patients and 6% (6/113) of procarbazine patients. Including stable disease (SD), the objective response (PR and SD) rate was 46% for temozolomide and 33% for procarbazine.

In patients with prior exposure to chemotherapy, the benefit of temozolomide was limited to those with KPS  $\geq$ 80. In patients who were progression-free at 6 months, quality of life was maintained or improved.

Results from a large, non-comparative trial provide further evidence of the efficacy of temozolomide in patients with relapsing GBM. Of the 128 patients with eligible histologies, all but two had GBM, the remaining two had gliosarcoma. The median age was 54 years and 62% were male. Karnofsky performance status was  $\geq$ 80 in 57%. At the time of initial diagnosis, 89% of patients underwent surgical resection, with all patients subsequently receiving radiation therapy. Eighty-six percent of patients were treated with standard dose fractionation. Nitrosourea-based chemotherapy was administered in 29% of patients. The median time from initial diagnosis to first relapse was 8.1 months. At first relapse, 13% of patients had surgical resection. The primary endpoint, progression-free survival at 6 months, was 19% (95% CI: 12% to 26%) for the intent-to-treat (ITT) population. The median progression-free survival was 2.1 months. Median overall survival was 5.4 months. The objective response (CR/PR) as determined by Gd-MRI scan after independent central review was 8% (11/138) for the ITT population. Including stable disease, the objective response (CR, PR and SD) was 51% (71/138). Both overall response as objectively assessed and maintenance in progression-free status were associated with HQL benefits.

In a large phase II study, temozolomide demonstrated clinically meaningful efficacy in AA patients in relapse. A total of 162 patients were enrolled and comprise the ITT population. A total of

111 patients was determined by central review as having histologically confirmed AA or AOA (anaplastic oligoastrocytoma) and comprises the eligible histology population who received temozolomide. Fifty one patients were excluded from the eligible histology population. The median age was 42 years and 57% were male. Karnofsky performance status was  $\geq 80$  in 67%. At the time of initial diagnosis, 68% of patients underwent surgical resection, with all patients subsequently receiving radiation therapy. Ninety-one percent of patients were treated with standard dose fractionation. Nitrosourea-based chemotherapy was administered in 60% of patients. The median time from initial diagnosis to first relapse was 14.9 months. At first relapse, 18% of patients had surgical resection.

Progression-free survival at 6 months was 46% (95% CI: 39%-54%). The median progression-free survival was 5.4 months. Twenty four percent of patients remained progression-free after 12 months. The median overall survival was 14.6 months. Fifty-eight percent of patients remained alive after 12 months.

The objective response rate (CR/PR) as determined by Gd-MRI scan after independent central review was 35% (13 CR and 43 PR) for the ITT population. Including stable disease, the objective response rate (CR, PR and SD) was 61% (99/162). For the 13 complete responders, the progression-free survival range was 11 to 26 months, with 7 patients remaining in complete response beyond 16 months; the overall survival for these patients ranged from 15 to 30 months, with 8 patients alive beyond 20 months. For the 43 partial responders, the median progression-free survival was 11 months and the median overall survival was 21 months.

**Comparative Bioavailability Study** (temozolomide capsules for oral administration vs. temozolomide powder for solution for IV injection)

In a randomized, multi-center, open-label two-way crossover bioequivalence study, twenty-two patients (12 male and 10 female) with primary CNS malignancies received temozolomide once daily for 5 days in the fasting state during a 4-week treatment cycle. Nineteen of the patients were included in the pharmacokinetic analyses. On Days 1, 2, and 5, patients received temozolomide (200 mg/m<sup>2</sup>/day) as an oral capsule dose; on Days 3 and 4, patients received temozolomide (150 mg/m<sup>2</sup>/day) as an oral dose on one day and as a 90 minute intravenous infusion using temozolomide powder for solution for injection on the other day. Temozolomide and MTIC were found to be bioequivalent for C<sub>max</sub> and AUC. Following administration of 150 mg/m<sup>2</sup> by 90-minute IV infusion, mean C<sub>max</sub> values for temozolomide and MTIC were 7.4 mcg/mL and 320 ng/mL, respectively. Mean AUC<sub>(t)</sub> values for temozolomide and MTIC were 25 mcg·hr/mL and 1004 ng·hr/mL, respectively.

**Summary Table of the Comparative Bioavailability Data for Temozolomide (TMZ)**

| <p align="center"><b>TMZ</b><br/> <b>(150 mg/m<sup>2</sup>)</b><br/> <b>From measured data</b></p> <p align="center"><b>Geometric Mean</b><br/> <b>Arithmetic Mean (CV %)</b></p> |  |   |                                   |                                |
|---|--|---|-----------------------------------|--------------------------------|
| <b>Parameter</b>  | <b>IV Formulation (Test)<sup>†</sup></b> | <b>Capsules (Reference)<sup>‡</sup></b> | <b>% Ratio of Geometric Means</b> | <b>90% Confidence Interval</b> |
| <b>AUC<sub>T</sub> (mcg·hr/mL)</b>  | 23.1<br>23.4 (18)                        | 21.8<br>22.0 (14)                       | 106                               | 103–109                        |
| <b>AUC<sub>I</sub> (mcg·hr/mL)</b>  | 24.6<br>25.0 (18)                        | 23.4<br>23.6 (15)                       | 105                               | 102–108                        |
| <b>C<sub>max</sub> (mcg/mL)</b>   | 7.29<br>7.44 (21)                        | 7.54<br>7.68 (19)                       | 97                                | 91–102                         |
| <b>T<sub>max</sub><sup>§</sup> (hr)</b>   | 1.50<br>(0.92–2.00)                      | 1.00<br>(0.25–2.00)                     |                                   |                                |
| <b>T<sub>1/2</sub><sup>*</sup> (hr)</b>   | 1.81 (12)                                | 1.91 (13)                               |                                   |                                |

<sup>†</sup>temozolomide powder for solution for injection

<sup>‡</sup>temozolomide capsules

<sup>§</sup> Expressed as the median (range)

\* Expressed as the arithmetic mean (CV %) only

N=19 patients

## Summary Table of the Comparative Bioavailability Data for MTIC

| <b>MTIC</b><br><b>(150 mg/m<sup>2</sup> TMZ)</b><br><b>From measured data</b><br><br><b>Geometric Mean</b><br><b>Arithmetic Mean (CV %)</b> |                                    |                                   |                            |                         |
|---|------------------------------------|-----------------------------------|----------------------------|-------------------------|
| Parameter   | IV Formulation (Test) <sup>†</sup> | Capsules (Reference) <sup>‡</sup> | % Ratio of Geometric Means | 90% Confidence Interval |
| <b>AUC<sub>T</sub></b><br>(ng•hr/mL)  | 837<br>941 (53)                    | 815<br>944 (60)                   | 103                        | 98–108                  |
| <b>AUC<sub>I</sub></b><br>(ng•hr/mL)  | 891<br>1004 (54)                   | 864<br>1003 (60)                  | 103                        | 98–108                  |
| <b>C<sub>max</sub></b><br>(ng/mL)   | 276<br>320 (61)                    | 282<br>333 (62)                   | 98                         | 91–105                  |
| <b>T<sub>max</sub></b> <sup>§</sup><br>(hr)   | 1.50<br>(1.25–1.75)                | 1.00<br>(0.25–2.00)               |                            |                         |
| <b>T<sub>½</sub></b> <sup>*</sup><br>(hr)   | 1.80 (16)                          | 1.77 (11)                         |                            |                         |

<sup>†</sup>temozolomide powder for solution for injection

<sup>‡</sup>temozolomide capsules

<sup>§</sup> Expressed as the median (range)

<sup>\*</sup> Expressed as the arithmetic mean (CV %) only

MTIC (monomethyl triazeno imidazole carboxamide)

TMZ: temozolomide

N=19 patients

## DETAILED PHARMACOLOGY

### Animal Pharmacology

#### Pharmacodynamics

The anti-tumor properties of temozolomide have been demonstrated *in vitro* and *in vivo*, with tumor cell lines and xenograft models. The cytotoxicity of temozolomide results from DNA methylation and correlates specifically with the O<sup>6</sup>-methylation of guanine residues.

Temozolomide showed marked *in vivo* anti-tumor activity in murine xenograft models. Murines with subcutaneous or intracranial implanted human CNS tumor were either long term, tumor-free survivors or their tumors had substantial growth delays.



Among a panel of human tumor cell lines, U373MG astrocytoma and U87MG glioblastoma were revealed as the most sensitive to temozolomide. In another *in vitro* study, with a broader profile of human glioma and medulloblastoma, CNS cell lines were as sensitive as U373MG astrocytoma to temozolomide.

In another study, temozolomide given orally to mice in early stage subcutaneous implanted astrocytoma xenograft model revealed dose-dependent anti-tumor activity: 60 to 100% of mice were tumor-free on Day 54. Of 60 U251 glioblastoma xenografts treated with temozolomide, all 57 surviving animals showed complete tumor regression.

Temozolomide showed greater tumor growth delay than BCNU or procarbazine with all four CNS tumor xenografts models studied.

Some studies showed that temozolomide would have potential synergistic effects with other cytotoxic drugs such as O<sup>6</sup>-Benzylguanine, cisplatin, topotecan, 3-aminobenzamine or chloroethylnitrosoureas.

Temozolomide safety pharmacology was assessed in cell lines, mice, rats and dogs. It was shown that it affected hematological parameters, increased total bilirubin and  $\gamma$ -glutamyl-transferase. Temozolomide also decreased food consumption, body weight and body weight gain; it even produced weight loss. Temozolomide did not affect the blood pressure and electrocardiogram in dogs. Temozolomide did not cause gastric mucosal lesions nor affect intestinal transit after a single oral dose. Temozolomide caused a moderate inhibition of gastric emptying. It increased urine volume and BUN values and decreased urine osmolality in rats. Finally, temozolomide had CNS effects when given at lethal doses: hypoactivity, hunched posture, partial closure of the eyes, tremors, prostration, emesis and salivation.

### Pharmacokinetics

Temozolomide is hydrolysed at physiological pH to MTIC, the metabolite responsible for DNA alkylation. The latter then breaks down into a reactive methyl-diazonium cation and AIC. AIC is an intermediate on the biosynthetic pathway to purines and ultimately to nucleic acids. Temozolomide is stable in acidic pH (<5) and labile at pH >7, and MTIC is unstable at pH <7 and more stable at alkaline pH.

Temozolomide was given to mice, rats and dogs under various forms of administration: orally (PO), intraperitoneally (IP), intraarterially (IA) and intravenously (IV) to determine its pharmacokinetics properties. It also has been studied *in vitro* in an aqueous buffer to assess its rate of chemical degradation.

C<sub>max</sub> was attained in mice 10 minutes after temozolomide PO and IP administration. Following oral administration in rats, temozolomide was rapidly absorbed and was completely bioavailable 0.25 hours later. Its mean half-life was found to be 1.2 hours and it was independent of the route of administration. This value was lower than the value reported for the degradation in aqueous buffer due to the renal clearance contribution.

Terminal phase half-life of temozolomide was similar in sick rats, compared to the value found in healthy rats. The volume of distribution at steady state was larger than in healthy rats and is probably due to the hyperpermeable state and neovascularization of the tumor.

Following PO dosing in healthy dogs, temozolomide was rapidly and completely absorbed. Its absolute bioavailability ranged from 95 to 110%. Bioavailability of the toxicology capsule was compared to the clinical capsule in dogs. There was no significant formulation effects seen in  $C_{\max}$  or  $AUC_{(t)}$  but there was a decrease in  $T_{\max}$  value indicating a more rapid absorption following administration of the clinical capsule.

Temozolomide was mainly excreted in urine and in small amounts in feces. 1.39% (IV) and 1.45% (PO) of the radiocarbon administered to rats was excreted in bile collected 48 hours postdose.

After repeated administration,  $AUC_{(t)}$  values for Day 1 and Day 5 of each cycle were the same for all dose levels in both rat and dog except for the 800 mg/m<sup>2</sup> given to male rats where the mean  $AUC_{(t)}$  value was higher for Day 5. Since temozolomide was shown to have a short elimination half-life, no accumulation with multiple dosing was expected.

Tissue distribution was assessed in rats in two studies. <sup>14</sup>C-temozolomide extensively distributed to all tissues. In both studies, high concentrations of radiocarbon were noted in tissues at the late sampling times due to the incorporation of <sup>14</sup>C-AIC into the purine biosynthetic pool. Results suggest that temozolomide crosses the blood-brain barrier rapidly and is present in the cerebrospinal fluid. Concentrations in brain and testes appeared highest at 1 hour postdose then decreased slowly; higher levels of radioactivity remained in the kidneys, liver, large and small intestinal wall, salivary gland and testes. No difference was found in tissue concentration related to gender.

No metabolites were identified in mouse during an *in vitro* study. In an *in vivo* study, it was found that 39% of temozolomide was excreted unchanged and that a small amount of TMA (temozolomide acid metabolite) was also excreted. No other metabolites were seen.

In rat, no metabolites were detected through 6 hours. Females excreted the same percentage of parent drug as males did. For dogs, temozolomide represented about 30% of the radiocarbon in plasma by 8 hours postdose.

## **Human Pharmacology**

### **Clinical Pharmacology**

Temozolomide was rapidly and completely absorbed when administered orally at therapeutic doses to humans.  $C_{\max}$  and AUC increased in a dose-proportional manner. No accumulation occurred on multiple dosing. The volume of distribution, clearance, and half-life were dose-independent, had very low coefficient of variation, and were predictable and reproducible. The major pathways for elimination of temozolomide from plasma were non-enzymatic hydrolysis to MTIC and renal excretion of parent drug. TMA was the only metabolite of significance and accounted for <3% of the dose excreted in urine.

Cytochrome P450 (CYP450)-mediated metabolism as assessed by measuring TMA levels did not contribute significantly to the plasma clearance of temozolomide. Consequently, clearance of temozolomide should not be affected to a clinically meaningful degree by interaction of concurrent medications with specific isozymes of CYP450 nor would administration of temozolomide alter by competitive inhibition the metabolism of other drugs. Analysis of data from phase II studies confirmed that clearance of temozolomide was unaffected by 7 medications commonly used by this

patient population (i.e., phenytoin, phenobarbital, carbamazepine, dexamethasone, H<sub>2</sub>-receptor antagonists, prochlorperazine, and ondansetron). Valproic acid was associated with a statistically significant (p=0.019) but clinically insignificant 4.7% decrease in the clearance of temozolomide. Renal disease should not affect temozolomide clearance. This is in agreement with experimental data which demonstrated that age, renal function, hepatic function and use of tobacco did not alter clearance of temozolomide. Female patients had a clinically insignificantly lower clearance of temozolomide than did male patients. Administration of temozolomide with food delayed absorption of temozolomide and resulted in a clinically insignificant 9% decrease in exposure. Compared to adults, pediatric patients over three years of age had higher plasma temozolomide concentrations. This is probably due to their higher body surface area to weight ratio.

MTIC degrades to AIC at a much faster rate than its rate of formation from temozolomide. Following oral dosing with temozolomide, the plasma t<sub>1/2</sub> for MTIC was the same as that for temozolomide (1.8 hours). Since the volume of distribution for temozolomide and MTIC are approximately the same, the AUC for MTIC could be predicted. The AUC for MTIC was approximately 2 to 4% of that of temozolomide.

Pharmacodynamic evaluations indicated that the primary hematologic toxicities of temozolomide (severe thrombocytopenia and neutropenia) were uncommon during the first cycle. Increasing dose and AUC of temozolomide were associated with an increased incidence of neutropenia and thrombocytopenia. Patients >70 years of age appeared to be at increased risk of neutropenia, although the number of patients in this age subgroup was small (8 patients). The incidence of thrombocytopenia and neutropenia was approximately three times higher in females. Pediatric patients appeared to tolerate higher plasma concentrations of temozolomide before reaching dose limiting toxicity. This is likely due to increased bone marrow reserves in pediatric patients.

## **TOXICOLOGY**

### **Acute Toxicity**

Acute toxicity studies were conducted in both mice and rats. In single dose studies conducted in mice, calculated LD<sub>50</sub> values were 891 (males) and 1072 (females) mg/m<sup>2</sup> for oral administration and 1297 (males) and 891 (females) mg/m<sup>2</sup> for intraperitoneal administration of temozolomide. In rats, LD<sub>50</sub> values were 1937 mg/m<sup>2</sup> when temozolomide was given orally and 1414 mg/m<sup>2</sup> for intraperitoneal administration. Antemortem observations for both mice and rats included hypoactivity, hunched posture and partial closure of the eyes (dose ≥1000 mg/m<sup>2</sup> generally). Tremors (≥1000 mg/m<sup>2</sup> PO, ≥2000 mg/m<sup>2</sup> IP), prostration (≥2000 mg/m<sup>2</sup>) and ataxia (≥4000 mg/m<sup>2</sup> IP) were also observed in mice. At necropsy, dark-red areas were observed in the stomachs of male mice at doses ≥3000 mg/m<sup>2</sup> (PO) or ≥2000 mg/m<sup>2</sup> (IP) and in female mice at doses ≥1000 mg/m<sup>2</sup> of temozolomide.

Observations for rats included abnormal or few feces (≥1500 mg/m<sup>2</sup> PO) and dyspnea (≥2500 mg/m<sup>2</sup> PO). When doses reached 5000 mg/m<sup>2</sup> orally or more, poor appetite, thin appearance, few or abnormal feces, anorexia and dyspnea were noted. Anorexia and swollen heads were also noted in rats at intraperitoneal doses of ≥2000 mg/m<sup>2</sup> of temozolomide. At necropsy, dark-red areas were observed in the stomach of rats at oral doses ≥1500 mg/m<sup>2</sup> and intraperitoneal doses ≥2000 mg/m<sup>2</sup>. Dark areas were also noted in the brain, reproductive organs, lymph nodes, lung, pancreas, cecum and subcutaneous tissue at oral doses ≥1500 mg/m<sup>2</sup>. At intraperitoneal doses ≥2000 mg/m<sup>2</sup>, dark areas were observed in the small intestine

(males, 4000 mg/m<sup>2</sup>), lymph nodes, lung and subcutaneous tissue.

Clinical observations in dogs which received a total dose of 3500 mg/m<sup>2</sup> of temozolomide over 6 days included emesis, hypoactivity, ataxia, polypnea, mydriasis and discolored mucoid feces. At necropsy, dark-red areas were observed in the stomach and dark-red to brown material in the gastrointestinal tract.

Emesis, salivation and abnormal or few feces were noted in dogs administered single oral doses  $\geq 200$  mg/m<sup>2</sup> of temozolomide. All dogs which received 200 or 400 mg/m<sup>2</sup> survived the 14-day observation period; dogs administered 600, 1000 or 1500 mg/m<sup>2</sup> of temozolomide died or were sacrificed in poor condition before the 14-day period was completed. Necropsy observations at doses 1000 mg/m<sup>2</sup> included dark areas in the stomach, lymph nodes, cecum, small intestine, heart, urinary bladder and subcutaneous tissue. There was no gross lesion observed at doses  $< 1000$  mg/m<sup>2</sup>.

### **Multiple-Dose Toxicity**

The toxicity of temozolomide was evaluated in single-cycle, three-cycle and six-cycle studies, in rats and dogs. Results are reported in the following tables.

|  | RATS                  |  | DOGS                  |   |
|--|-----------------------|--|-----------------------|---|
|  | DOSES                 | TOXIC EFFECTS  | DOSES                 | TOXIC EFFECTS   |
| SINGLE-CYCLE STUDIES<br><br><b>Rats:</b><br><br><b>Dogs:</b> | 200 mg/m <sup>2</sup> | <ul style="list-style-type: none"> <li>• 1 male died</li> <li>• ↓ mean food consumption, body weight and body weight gain</li> <li>• ↓ mean erythrocytic and leukocytic values</li> <li>• ↓ mean platelet, lymphocyte and segmented neutrophil counts</li> <li>• ↑ total bilirubin, GGT and BUN</li> <li>• ↓ total protein and albumin</li> <li>• ↓ organ weights:               <ul style="list-style-type: none"> <li>• thymus</li> <li>• prostate</li> <li>• spleen/testes</li> </ul> </li> <li>• <b>necropsy findings:</b> <ul style="list-style-type: none"> <li>• dark areas on stomach, lung, testes, lymph nodes</li> <li>• pale areas on liver and kidneys</li> <li>• enlarged seminal vesicles</li> <li>• degeneration of testes</li> </ul> </li> <li>• <b>histopathologic findings:</b> <ul style="list-style-type: none"> <li>• lymphoid depletion of thymus</li> <li>• hypertrophy/reduced colloid in thyroid gland</li> <li>• syncytial cells in the testes</li> <li>• lymphoid depletion of spleen</li> <li>• crypt necrosis</li> <li>• hypocellularity in bone marrow</li> <li>• degeneration of testes</li> <li>• hyperplasia/mucosal epithelium disruption of small intestine</li> </ul> </li> </ul> | 200 mg/m <sup>2</sup> | <ul style="list-style-type: none"> <li>• all dogs died or were sacrificed</li> <li>• emesis</li> <li>• hypoactivity</li> <li>• dehydration</li> <li>• anorexia</li> <li>• abnormal feces</li> <li>• ↓ food consumption</li> <li>• ↓ body weight/weight gain</li> <li>• ↓ mean erythrocytic and leukocytic values</li> <li>• <b>necropsy findings:</b> <ul style="list-style-type: none"> <li>• enlarged, dark lymph nodes</li> <li>• dark areas in the intestine, urinary bladder, esophagus, heart, thymus, subcutaneous tissue</li> <li>• pale/raised areas of the spleen</li> <li>• small thymus glands</li> </ul> </li> <li>• <b>histopathologic findings:</b> <ul style="list-style-type: none"> <li>• lymphoid depletion of the thymus</li> <li>• syncytial cells in the testes</li> <li>• atrophy of bone marrow</li> <li>• lymphoid depletion of the spleen, lymph nodes and small intestine</li> <li>• hemorrhage, crypt necrosis and congestion of small intestine</li> </ul> </li> </ul> |
|  | 400 mg/m <sup>2</sup> | <ul style="list-style-type: none"> <li>• 9 males/9 females died</li> <li>• hypoactivity</li> <li>• hunched posture</li> </ul>  | 500 mg/m <sup>2</sup> | <ul style="list-style-type: none"> <li>• all dogs died or were sacrificed</li> <li>• emesis</li> <li>• hypoactivity</li> </ul>  |

|  | RATS  |   | DOGS  |   |
|--|-------|---|-------|---|
|  | DOSES | TOXIC EFFECTS   | DOSES | TOXIC EFFECTS   |
|  |       | <ul style="list-style-type: none"> <li>• thin appearance</li> <li>• few feces</li> <li>• ↓ mean food consumption, body weight and body weight gain</li> <li>• bilateral pallor of fundus of the eyes (10 rats)</li> <li>• ↓ mean erythrocytic and leukocytic values</li> <li>• ↓ mean platelet, lymphocyte and segmented neutrophil counts</li> <li>• ↑ urine volume, ↓ urine osmolality</li> <li>• ↓ organ weights: <ul style="list-style-type: none"> <li>• thymus</li> <li>• prostate</li> <li>• spleen/testes</li> <li>• pituitary gland</li> <li>• salivary gland</li> <li>• heart</li> <li>• ovary, epididymis</li> </ul> </li> <li>• <b>necropsy findings:</b> <ul style="list-style-type: none"> <li>• dark areas on stomach, lung, testes, lymph nodes</li> <li>• pale areas on liver and kidneys</li> <li>• enlarged seminal vesicles</li> <li>• degeneration of testes</li> </ul> </li> <li>• <b>histopathologic findings:</b> <ul style="list-style-type: none"> <li>• lymphoid depletion of thymus</li> <li>• hypertrophy/reduced colloid in thyroid gland</li> <li>• syncytial cells in the testes</li> <li>• lymphoid depletion of spleen</li> <li>• crypt necrosis</li> </ul> </li> </ul> |       | <ul style="list-style-type: none"> <li>• dehydration</li> <li>• anorexia</li> <li>• abnormal feces</li> <li>• ↓ food consumption</li> <li>• ↓ body weight/weight gain</li> <li>• ↓ mean erythrocytic and leukocytic values</li> <li>• <b>necropsy findings:</b> <ul style="list-style-type: none"> <li>• enlarged, dark lymph nodes</li> <li>• dark areas in the intestine, urinary bladder, esophagus, heart, thymus, subcutaneous tissue</li> <li>• pale/raised areas of the spleen</li> <li>• small thymus glands</li> </ul> </li> <li>• <b>histopathologic findings:</b> <ul style="list-style-type: none"> <li>• lymphoid depletion of the thymus</li> <li>• syncytial cells in the testes</li> <li>• atrophy of bone marrow</li> <li>• lymphoid depletion of the spleen, lymph nodes and small intestine</li> <li>• hemorrhage, crypt necrosis and congestion of small intestine</li> </ul> </li> </ul> |

|  | <b>RATS</b> |  | <b>DOGS</b> |               |
|--|-------------|--|-------------|---------------|
|  | DOSES       | TOXIC EFFECTS  | DOSES       | TOXIC EFFECTS |
|  |             | <ul style="list-style-type: none"> <li>• hypocellularity in bone marrow</li> <li>• degeneration of testes</li> <li>• hyperplasia/mucosal epithelium disruption of small intestine</li> </ul> |             |               |

|  | RATS  |   | DOGS                      |   |
|--|---|---|---------------------------|---|
|  | DOSES   | TOXIC EFFECTS   | DOSES                     | TOXIC EFFECTS   |
|  | 800/male<br>or<br>600/female<br>mg/m <sup>2</sup> | <ul style="list-style-type: none"> <li>• all rats died or sacrificed by Day 21</li> <li>• hypoactivity</li> <li>• hunched posture</li> <li>• thin appearance</li> <li>• few feces</li> <li>• ↓ mean food consumption, body weight and body weight gain</li> <li>• ↓ mean erythrocytic and leukocytic values</li> <li>• ↓ mean platelet, lymphocyte and segmented neutrophil counts</li> <li>• ↑ urine volume, ↓ urine osmolality</li> <li>• ↓ organ weights: <ul style="list-style-type: none"> <li>• thymus</li> <li>• prostate</li> <li>• spleen/testes</li> </ul> </li> <li>• <b>necropsy findings:</b> <ul style="list-style-type: none"> <li>• dark areas on stomach, lung, testes, lymph nodes</li> <li>• pale areas on liver and kidneys</li> <li>• enlarged seminal vesicles</li> <li>• degeneration of testes</li> </ul> </li> <li>• <b>histopathologic findings:</b> <ul style="list-style-type: none"> <li>• lymphoid depletion of thymus</li> <li>• hypertrophy/reduced colloid in thyroid gland</li> <li>• syncytial cells in the testes</li> <li>• retinal degeneration/necrosis</li> <li>• lymphoid depletion of spleen</li> <li>• crypt necrosis</li> <li>• hypocellularity in bone marrow</li> <li>• degeneration of testes</li> </ul> </li> </ul> | 1000<br>mg/m <sup>2</sup> | <ul style="list-style-type: none"> <li>• all dogs died or were sacrificed</li> <li>• emesis</li> <li>• hypoactivity</li> <li>• dehydration</li> <li>• anorexia</li> <li>• abnormal feces</li> <li>• ↓ food consumption, ↓ body weight and weight gain</li> <li>• ↓ mean erythrocytic and leukocytic values</li> <li>• <b>necropsy findings:</b> <ul style="list-style-type: none"> <li>• enlarged, dark lymph nodes</li> <li>• dark areas in the intestine, urinary bladder, esophagus, heart, thymus, subcutaneous tissue</li> <li>• pale/raised areas of the spleen</li> <li>• small thymus glands</li> <li>• prominent lymphoid tissue in the intestine</li> </ul> </li> <li>• <b>histopathologic findings:</b> <ul style="list-style-type: none"> <li>• lymphoid depletion of the thymus</li> <li>• syncytial cells in the testes</li> <li>• atrophy of bone marrow</li> <li>• lymphoid depletion of the spleen, lymph nodes and small intestine</li> <li>• hemorrhage, crypt necrosis and congestion of small intestine</li> <li>• degeneration/necrosis of the outer layer of the retina</li> </ul> </li> </ul> |



|  | RATS                  |  | DOGS                  |  |
|--|-----------------------|--|-----------------------|--|
|  | DOSES                 | TOXIC EFFECTS  | DOSES                 | TOXIC EFFECTS  |
|  |                       | <ul style="list-style-type: none"> <li>• hyperplasia/mucosal epithelium disruption of small intestine</li> </ul>   |                       |  |
|  | 25 mg/m <sup>2</sup>  | <ul style="list-style-type: none"> <li>• ↓ organ weights: <ul style="list-style-type: none"> <li>• thymus</li> </ul> </li> <li>• <b>necropsy findings:</b> <ul style="list-style-type: none"> <li>• dark lung (1 female)</li> </ul> </li> <li>• <b>histopathologic findings:</b> <ul style="list-style-type: none"> <li>• lymphoid depletion of thymus</li> <li>• hypertrophy/reduced colloid in thyroid gland</li> <li>• syncytial cells in testes</li> </ul> </li> </ul>                                       | 25 mg/m <sup>2</sup>  |  |
|  | 50 mg/m <sup>2</sup>  | <ul style="list-style-type: none"> <li>• ↓ mean platelet, lymphocyte and segmented neutrophil counts</li> <li>• ↓ organ weights: <ul style="list-style-type: none"> <li>• thymus</li> </ul> </li> <li>• <b>histopathologic findings:</b> <ul style="list-style-type: none"> <li>• lymphoid depletion of thymus</li> <li>• hypertrophy/reduced colloid in thyroid gland</li> <li>• syncytial cells in testes</li> </ul> </li> </ul>   | 50 mg/m <sup>2</sup>  | <ul style="list-style-type: none"> <li>• emesis</li> </ul>   |
|  | 100 mg/m <sup>2</sup> | <ul style="list-style-type: none"> <li>• ↓ mean erythrocytic and leukocytic values</li> <li>• ↓ mean platelet, lymphocyte and segmented neutrophil counts</li> <li>• ↓ organ weights: <ul style="list-style-type: none"> <li>• thymus</li> <li>• spleen/testes</li> </ul> </li> <li>• <b>histopathologic findings:</b> <ul style="list-style-type: none"> <li>• lymphoid depletion of thymus</li> <li>• hypertrophy/reduced colloid in thyroid gland</li> <li>• syncytial cells in testes</li> </ul> </li> </ul> | 125 mg/m <sup>2</sup> | <ul style="list-style-type: none"> <li>• 1 male died</li> <li>• hypoactivity</li> <li>• <b>histopathologic findings:</b> <ul style="list-style-type: none"> <li>• lymphoid depletion of the thymus</li> <li>• syncytial cells in the testes</li> </ul> </li> </ul> |

|  | RATS                  |  | DOGS  |               |
|--|-----------------------|--|-------|---------------|
|  | DOSES                 | TOXIC EFFECTS  | DOSES | TOXIC EFFECTS |
|  |                       | <ul style="list-style-type: none"> <li>• lymphoid depletion of spleen</li> <li>• crypt necrosis</li> </ul>   |       |               |
|  | 150 mg/m <sup>2</sup> | <ul style="list-style-type: none"> <li>• ↓ mean erythrocytic and leukocytic values</li> <li>• ↓ mean platelet, lymphocyte and segmented neutrophil counts</li> <li>• ↓ organ weights: <ul style="list-style-type: none"> <li>• thymus</li> <li>• spleen/testes</li> </ul> </li> <li>• <b>histopathologic findings:</b> <ul style="list-style-type: none"> <li>• lymphoid depletion of thymus</li> <li>• hypertrophy/reduced colloid in thyroid gland</li> <li>• syncytial cells in testes</li> <li>• lymphoid depletion/spleen</li> <li>• crypt necrosis</li> <li>• hypocellularity/bone marrow</li> <li>• degeneration of testes</li> <li>• hyperplasia/mucosal epithelium disruption of small intestine</li> </ul> </li> </ul> |       |               |
|  | 200 mg/m <sup>2</sup> | <ul style="list-style-type: none"> <li>• ↓ mean food consumption, body weight and body weight gain</li> <li>• ↓ mean erythrocytic and leukocytic values</li> <li>• ↓ mean platelet, lymphocyte and segmented neutrophil counts</li> <li>• ↑ total bilirubin, GGT and BUN</li> <li>• ↓ total protein and albumin</li> <li>• ↓ organ weights: <ul style="list-style-type: none"> <li>• thymus</li> <li>• spleen/testes</li> </ul> </li> </ul>  |       |               |

|  | RATS  |   | DOGS  |               |
|--|-------|---|-------|---------------|
|  | DOSES | TOXIC EFFECTS   | DOSES | TOXIC EFFECTS |
|  |       | <ul style="list-style-type: none"> <li>• <b>histopathologic findings:</b></li> <li>• lymphoid depletion of thymus</li> <li>• hypertrophy/reduced colloid in thyroid gland</li> <li>• syncytial cells in testes</li> <li>• lymphoid depletion/spleen</li> <li>• crypt necrosis</li> <li>• hypocellularity/bone marrow</li> <li>• degeneration of testes</li> <li>• hyperplasia/mucosal epithelium disruption of small intestine</li> </ul> |       |               |

|   | RATS                  |  | DOGS                  |  |
|---|-----------------------|--|-----------------------|--|
|   | DOSES                 | TOXIC EFFECTS  | DOSES                 | TOXIC EFFECTS  |
| THREE-CYCLE STUDIES<br><br><b>Rats:</b><br><br><b>Dogs:</b> | 25 mg/m <sup>2</sup>  | <ul style="list-style-type: none"> <li>• ↓ food consumption (during 1st week of cycle one)</li> <li>• <b>necropsy findings:</b> <ul style="list-style-type: none"> <li>• ↓ mean thymus weight (interim)</li> </ul> </li> <li>• <b>histopathologic changes:</b> <ul style="list-style-type: none"> <li>• lymphoid depletion/thymus</li> </ul> </li> </ul>   | 25 mg/m <sup>2</sup>  | <ul style="list-style-type: none"> <li>• emesis in several dogs</li> <li>• ↓ lactate dehydrogenase in males</li> </ul>   |
|   | 50 mg/m <sup>2</sup>  | <ul style="list-style-type: none"> <li>• ↓ food consumption (during 1st week of cycle one)</li> <li>• <b>necropsy findings:</b> <ul style="list-style-type: none"> <li>• ↓ mean thymus weight (interim)</li> <li>• small thymus</li> <li>• alopecia</li> </ul> </li> <li>• <b>histopathologic changes:</b> <ul style="list-style-type: none"> <li>• lymphoid depletion of thymus</li> </ul> </li> </ul>  | 50 mg/m <sup>2</sup>  | <ul style="list-style-type: none"> <li>• emesis in several dogs</li> <li>• hypoactivity in a few dogs</li> <li>• ↓ lactate dehydrogenase in males and females</li> </ul> <p><b>NO-OBSERVABLE-EFFECT LEVEL</b><br/>(with minor exceptions)</p>  |
|   | 200 mg/m <sup>2</sup> | <ul style="list-style-type: none"> <li>• hair loss</li> <li>• alopecia (dose-related)</li> <li>• palpable subcutaneous masses along the thorax and abdomen (2 males and 19 females)</li> <li>• ↓ mean food consumption, body weights and body weight gains</li> <li>• ↓ erythrocyte, reticulocyte and platelet counts</li> <li>• ↓ hemoglobin and hematocrit</li> <li>• ↓ total and corrected leukocyte, segmented neutrophils and lymphocyte counts</li> <li>• <b>necropsy findings:</b> <ul style="list-style-type: none"> <li>• ↓ mean thymus weight (interim)</li> <li>• ↓ testes and epididymides weights (terminal)</li> <li>• masses (in 2/10 females)/interim</li> <li>• masses in 2/20 males and 17/20 females /terminal</li> <li>• small thymuses</li> </ul> </li> </ul> | 125 mg/m <sup>2</sup> | <ul style="list-style-type: none"> <li>• emesis in all dogs</li> <li>• pale gums in some dogs</li> <li>• hypoactivity in a few dogs</li> <li>• ↓ platelet, leukocyte, neutrophil and/or lymphocyte (during and after dosing period)</li> <li>• ↓ lactate dehydrogenase in males and females</li> <li>• <b>postmortem findings:</b> <ul style="list-style-type: none"> <li>• ↓ thymus weight in females</li> </ul> </li> <li>• <b>histopathologic findings:</b> <ul style="list-style-type: none"> <li>• lymphoid depletion in the thymus and spleen</li> <li>• ↑ syncytial cells in the testes</li> <li>• ↑ immature/abnormal sperm forms in the epididymal ducts</li> </ul> </li> </ul> |

|  | RATS  |   | DOGS  |               |
|--|-------|---|-------|---------------|
|  | DOSES | TOXIC EFFECTS   | DOSES | TOXIC EFFECTS |
|  |       | <ul style="list-style-type: none"> <li>• alopecia</li> <li>• <b>histopathologic changes:</b></li> <li>• bone marrow hypocellularity and hemorrhage</li> <li>• necrosis of crypt epithelium of small and large intestine</li> <li>• lymphoid depletion of the thymus</li> <li>• lymphoid depletion of the spleen</li> <li>• reduced colloid and hypertrophy of follicular epithelium in some thyroid glands</li> </ul> |       |               |

|   | RATS                  |  | DOGS                  |   |
|---|-----------------------|--|-----------------------|---|
|   | DOSES                 | TOXIC EFFECTS  | DOSES                 | TOXIC EFFECTS   |
| SIX-CYCLE STUDIES<br><br><b>Rats:</b><br><br><b>Dogs:</b> | 25 mg/m <sup>2</sup>  | <ul style="list-style-type: none"> <li>• 1 death (male)</li> <li>• lymphoid depletion of thymus (interim)</li> <li>• mammary gland carcinoma and carcinoma <i>in situ</i> (few females)</li> </ul>   | 25 mg/m <sup>2</sup>  | <ul style="list-style-type: none"> <li>• emesis</li> </ul>  |
|   | 50 mg/m <sup>2</sup>  | <ul style="list-style-type: none"> <li>• 1 death (male)</li> <li>• ↓ mean body weight for females (terminal sacrifice)</li> <li>• ↓ weekly food consumption and body weight gain</li> <li>• ↓ mean thymus weight (females)</li> <li>• ↓ testes weights (terminal sacrifice)</li> <li>• lymphoid depletion of thymus (interim)</li> <li>• mammary gland carcinoma and carcinoma <i>in situ</i> (few females)</li> </ul>   | 50 mg/m <sup>2</sup>  | <ul style="list-style-type: none"> <li>• emesis</li> </ul> <p><b>NO-OBSERVABLE-EFFECT LEVEL</b><br/>(with minor exceptions)</p>   |
|   | 125 mg/m <sup>2</sup> | <ul style="list-style-type: none"> <li>• 18 deaths (8 males and 10 females) <ul style="list-style-type: none"> <li>• most female deaths: carcinomas</li> </ul> </li> <li>• hair loss (moderate)</li> <li>• swollen areas of the body</li> <li>• palpable masses in males (5/35) and females (31/35)</li> <li>• hunched posture, hypoactivity (females)</li> <li>• pale coloring (females)</li> <li>• ↓ mean absolute body weight, weekly food consumption and body weight gain</li> <li>• ↓ erythrocyte count, hemoglobin and hematocrit</li> <li>• ↓ leukocyte and lymphocyte counts</li> </ul> | 125 mg/m <sup>2</sup> | <ul style="list-style-type: none"> <li>• emesis</li> <li>• pale gums</li> <li>• discolored feces</li> <li>• body weight loss</li> <li>• mean platelet, total leukocyte, segmented neutrophil and lymphocyte values vary in a cyclic manner</li> <li>• mild cyclic changes in erythrocyte parameters for females</li> <li>• <b>postmortem findings:</b> <ul style="list-style-type: none"> <li>• histomorphologic alterations of the spleen, kidneys, testes and epididymides</li> </ul> </li> </ul> |

|  | RATS  |   | DOGS  |  |
|--|-------|---|-------|--|
|  | DOSES | TOXIC EFFECTS   | DOSES | TOXIC EFFECTS  |
|  |       | <ul style="list-style-type: none"> <li>• ↓ total protein, albumin and globulin (cycles 5 and 6)</li> <li>• ↓ mean thymus weight</li> <li>• ↑ mean absolute organ weights, organ-to-body weight ratio, organ-to-brain weight ratio ( for liver, kidneys and adrenal glands)/females at interim sacrifice</li> <li>• ↑ liver and spleen weights (terminal sacrifice) for females</li> <li>• ↑ adrenal weights (terminal sacrifice) for males</li> <li>• ↓ testes weights</li> <li>• histopathologic changes in hematopoietic system, testes and epididymides, mammary gland, adrenal cortex and skin</li> <li>• ↑ incidence of miscellaneous neoplasms</li> <li>• lymphoid depletion of thymus (interim and terminal)</li> <li>• mammary gland carcinoma and carcinoma <i>in situ</i> (most females)</li> <li>• keratoacanthomas of the skin (54%) and basal cell adenoma (infrequently) in males</li> <li>• various mesenchymal neoplasms</li> </ul> |       | <ul style="list-style-type: none"> <li>• ↑ extramedullary hematopoiesis</li> <li>• pigmented spleen</li> <li>• syncytial cells in the testes</li> <li>• ↑ in immature/abnormal sperm form</li> </ul> |

These studies demonstrated that temozolomide was absorbed in a dose-related manner, without sex differences and no evidence of accumulation. The overall carcinogenic potential of temozolomide in rats does not appear significantly different from other chemotherapeutic drugs. Hematologic changes seem to be cyclic: they happened after dosing and were followed by a recovery period.

### **Carcinogenicity**

Carcinogenicity studies of temozolomide have not been conducted. However, the results of the six-cycle study in rats can be used to evaluate the carcinogenic potential of temozolomide.

Many types of neoplasms were observed in the six-cycle rat study. They included mammary carcinoma, carcinoma *in situ*, keratoacanthoma of the skin and basal cell adenoma. Mesenchymal neoplasms included fibrosarcoma, malignant schwannoma, endometrial stromal sarcoma, sarcoma, hemangiosarcoma and fibroma. No tumors or indication of preneoplastic changes were observed in the dog studies. Considering that temozolomide is a prodrug of an alkylating agent, MTIC, its carcinogenic potential is not unexpected.

### **Mutagenicity**

Temozolomide was found to be mutagenic in two studies: an Ames Assay for bacterial mutagenicity and a human peripheral blood lymphocyte assay. Additional *in vitro* toxicity studies are not being conducted as both assays were positive for mutagenic potential, and neoplasia has been observed *in vivo*. Since these findings are consistent with other drugs in this class, it is unlikely that *in vivo* assays would provide additional information that could impact the clinical use of temozolomide or aid in the assessment of human risk. Therefore, no *in vivo* mutagenic potential studies were conducted.

### **Reproductive Toxicity**

Segment I studies were not conducted with temozolomide. In pregnant rats and rabbits, temozolomide did not affect pregnancy maintenance.

The results of the multiple-cycle studies indicate testicular toxicity: reduced absolute testes weights occurred in rats at doses of 50 mg/m<sup>2</sup> and syncytial cells were observed in the testes of both rats and dogs at doses of 125 mg/m<sup>2</sup>. These results suggest additional potential reproductive effects including infertility and possibly genetic damage to germ cells.

Testing for reproductive toxicity was limited to dose range finding studies in rats and rabbits. No significant maternal toxicity was observed and pregnancy rates were not affected in either species. Dosing did not influence implantation rates or lengths of gestation. Resorptions and post implantation loss were increased at the 150 mg/m<sup>2</sup>/day dose level, compared to 5, 25 and 50 mg/m<sup>2</sup>/day dose levels. Fetal weights were reduced at 50 (slight) and 150 mg/m<sup>2</sup>/day. No external variations or malformations were observed in the rat study. In the rabbit study, 18 different types of malformations were observed in the fetuses of rabbits dosed with 125 mg/m<sup>2</sup>/day. Based on these results, the developmental NOEL is approximately 50 mg/m<sup>2</sup>/day. These data indicate that temozolomide, like other alkylating agents, has potential to produce embryoletality and malformations in rats and rabbits.



Segment III studies of temozolomide were not conducted. Considering that temozolomide's therapeutic intent is to interfere with mitosis, postnatal growth and development of offspring may be adversely affected by exposure to temozolomide if present in mothers' milk.

The preclinical toxicology profile of temozolomide for IV administration is comparable to that of the oral (capsule) formulation and consistent with that of other marketed alkylating anticancer agents. While the IV formulation produced local irritation at the site of injection in both rabbits and rats, the irritation was transient and not associated with lasting tissue damage.

## REFERENCES

1. Bower M, Newlands ES, Bleehen NM, Brada M, et al. Multicentre CRC phase II trial of temozolomide in recurrent or progressive high-grade glioma. *Canc Chemother Pharmacol* 1997;40:484-488.
2. Brock CS, Matthews JC, Brown G, Newlands ES, et al. *In vivo* demonstration of <sup>11</sup>C temozolomide uptake by human recurrent high grade astrocytomas. *Br J Cancer* 1997; 75(8):1241.
3. Devineni D, Klein-Szanto A, Gallo JM. Uptake of temozolomide in a rat glioma model in the presence and absence of the angiogenesis inhibitor TNP-470. *Cancer Research* 1996;56:1983-1987.
4. Newlands ES, Stevens MFG, Wedge SR, Wheelhouse RT, Brock C. Temozolomide: a review of its discovery, chemical properties, pre-clinical development and clinical trials. *Canc Treat Rev* 1997;23:35-61.
5. Osoba D, Aaronson NK, Muller M, Sneeuw K, et al. Effect of neurological dysfunction on health-related quality of life in patients with high-grade glioma. *J Neuro-Oncol* 1997;34:263-278.
6. Tsang LLH, Quarterman CP, Gescher A, Slack JA. Comparison of the cytotoxicity in vitro of temozolomide and dacarbazine, prodrugs of 3-methyl-(triazene-1-yl)imidazole-4-carboxamide. *Cancer Chemother Pharmacol* 1991;27:342-346.
7. Stupp R, Mason WP, van den Bent MJ, Weller M, et al., Radiotherapy plus Concomitant and Adjuvant Temozolomide for Glioblastoma. *N Engl J Med* 2005;352:987-96.
8. Product Monograph – Temodal® (temozolomide 5 mg, 20 mg, 100 mg, 140 mg and 250 mg Capsules). Merck Canada Inc., Submission Control No: 204940. Date of Revision: June 23, 2017.

**PART III: CONSUMER INFORMATION**

<sup>Pr</sup>  
**APO-TEMOZOLOMIDE**  
**Temozolomide Capsules**  
**5 mg, 20 mg, 100 mg, 140 mg, 180 mg and 250 mg**

**This leaflet is part III of a three-part “Product Monograph” published when APO-TEMOZOLOMIDE was approved for sale in Canada and is designed specifically for Consumers. This leaflet is a summary and will not tell you everything about APO-TEMOZOLOMIDE. Contact your doctor or pharmacist if you have any questions about the drug.**

**Please read this leaflet carefully before you start to take your medicine. Keep this leaflet. You may want to read it again. Remember, this medicine is for you and must be used as prescribed by your doctor. Never give it to anyone else.**

**ABOUT THIS MEDICATION**

**What the medication is used for:**

- APO-TEMOZOLOMIDE in combination with radiotherapy is used in the treatment of adult patients with newly diagnosed glioblastoma multiforme (GBM) (a form of brain tumor) and then as maintenance therapy.
- APO-TEMOZOLOMIDE alone is used in the treatment of adult patients with recurrent or progressive GBM or anaplastic astrocytoma (AA) after standard therapy.

**What it does:**

APO-TEMOZOLOMIDE is an antitumor agent. APO-TEMOZOLOMIDE acts on cancer cells. Normal cells may also be affected which may lead to side effects (see Warnings and Precautions section).

**When it should not be used:**

This medicine should not be used:

- If you are allergic to temozolomide or to any of its ingredients.
- If you have had an allergic reaction to dacarbazine (DTIC), another drug used to treat cancer.
- If you have low blood cell counts (severe myelosuppression).

**What the medicinal ingredient is:**

APO-TEMOZOLOMIDE medicinal ingredient is temozolomide.

**What the non-medicinal ingredients are:**

APO-TEMOZOLOMIDE Capsule’s non-medicinal ingredients: colloidal silicon dioxide, lactose anhydrous, sodium starch glycolate, stearic acid and tartaric acid.

The empty capsule shells contain the non-medicinal ingredients FD&C Blue No. 2 (5 mg, 140 mg), gelatin, iron oxide red (100 mg, 180 mg), iron oxide yellow (5 mg, 20 mg, 180 mg), talc and titanium dioxide.

The capsule shells are imprinted with black edible ink containing ammonium hydroxide, iron oxide black, isopropyl alcohol, N-Butyl alcohol, propylene glycol and shellac glaze.

**What dosage forms it comes in:**

APO-TEMOZOLOMIDE 5 mg Capsules: Each white opaque body, green opaque cap, hard gelatin capsule, imprinted “APO TEM 5” in black ink, white to light pink powder fill, contains 5 mg of temozolomide. Available in bottles of 5 and 20, and unit dose sachets of 1 capsule (5 or 20 sachets per box).

APO-TEMOZOLOMIDE 20 mg Capsules: Each white opaque body, gold opaque cap, hard gelatin capsule, imprinted “APO TEM 20” in black ink, white to light pink powder fill, contains 20 mg of temozolomide. Available in bottles of 5 and 20, and unit dose sachets of 1 capsule (5 or 20 sachets per box).

APO-TEMOZOLOMIDE 100 mg Capsules: Each white opaque body, flesh opaque cap, hard gelatin capsule, imprinted “APO TEM 100” in black ink, white to light pink powder fill, contains 100 mg of temozolomide. Available in bottles of 5 and 20, and unit dose sachets of 1 capsule (5 or 20 sachets per box).

APO-TEMOZOLOMIDE 140 mg Capsules: Each white opaque body, blue transparent cap, hard gelatin capsule, imprinted “APO TEM 140” in black ink, white to light pink powder fill, contains 140 mg of temozolomide. Available in bottles of 5 and 20, and unit dose sachets of 1 capsule (5 or 20 sachets per box).

APO-TEMOZOLOMIDE 180 mg Capsules: Each white opaque body, swedish orange opaque cap, hard gelatin capsule, imprinted “APO TEM 180” in black ink, white to light pink powder fill, contains 180 mg of temozolomide. Available in bottles of 5 and 20, and unit dose sachets of 1 capsule (5 or 20 sachets per box).

APO-TEMOZOLOMIDE 250 mg Capsules: Each white opaque body, white opaque cap, hard gelatin capsule, imprinted “APO TEM 250” in black ink, white to light

pink powder fill, contains 250 mg of temozolomide. Available in bottles of 5 and unit dose sachets of 1 capsule (5 or 20 sachets per box).

**WARNINGS AND PRECAUTIONS**

**Serious Warnings and Precautions**

APO-TEMOZOLOMIDE should be prescribed by doctor experienced with the use of cancer drugs.

APO-TEMOZOLOMIDE may cause a severe decrease in the production of blood cells which may be life threatening.

APO-TEMOZOLOMIDE may cause liver problems which may be life threatening.

Nausea and vomiting are very common with the use of APO-TEMOZOLOMIDE.

APO-TEMOZOLOMIDE combination with radiotherapy may cause severe pneumonia (*Pneumocystis carinii*).

BEFORE you use APO-TEMOZOLOMIDE talk to your doctor or pharmacist if you:

- have liver problems,
- have kidney problems,
- have a history of hepatitis B or current hepatitis B infection,
- are pregnant or planning to become pregnant,
- are breast feeding, or
- plan to father a child (or seek advice on cryoconservation, a laboratory technique which involves freezing of sperm).

In some cases, patients who have had hepatitis B might have a repeat attack of hepatitis. Tell the doctor if you think you have had hepatitis B in the past.

Infection with hepatitis B virus causes inflammation of the liver which may show as mild fever, feeling of sickness, fatigue, loss of appetite, joint and/or abdominal pain and yellowing of whites of the eyes, skin and tongue. If you experience any of these symptoms immediately contact your doctor.

APO-TEMOZOLOMIDE may cause harm to your unborn child, both male and female patients should use effective method of birth control while taking APO-TEMOZOLOMIDE and for 6 months after the last dose of

APO-TEMOZOLOMIDE.

Male patients should also be advised that APO-TEMOZOLOMIDE may cause irreversible infertility.

Do not drive or use machines until you know how you react to APO-TEMOZOLOMIDE.

**INTERACTIONS WITH THIS MEDICATION**

To avoid the possibility of one drug affecting another drug, be sure to advise your doctor or pharmacist of any other medications you are taking. Valproic Acid is an example of such drug interaction.

**PROPER USE OF THIS MEDICATION**

Your doctor will determine the dose of APO-TEMOZOLOMIDE based on your height and weight (m<sup>2</sup>). Take APO-TEMOZOLOMIDE as instructed by your doctor.

**Usual dose:**

Adult Dose:

Newly diagnosed Glioblastoma Multiforme (GBM):

Concomitant with Radiotherapy: 75 mg/m<sup>2</sup> daily for 42 days (up to 49 days).

Maintenance phases: 150 mg/m<sup>2</sup> daily for 5 days for the first cycle, then 200 mg/m<sup>2</sup> daily for cycle 2 to 6 if tolerated (a cycle equals 28 days).

**GBM or anaplastic astrocytoma (AA) Recurrence or Progression after Standard Therapy:**

Previously untreated with chemotherapy: 200 mg/m<sup>2</sup> daily for 5 days every 4 weeks per 28 day cycle.

Previously treated with chemotherapy: 150 mg/m<sup>2</sup> daily for 5 days for the first cycle to be increased in the second cycle to 200 mg/m<sup>2</sup> once daily for 5 days, if no hematologic toxicity.

**How APO-TEMOZOLOMIDE is taken:**

APO-TEMOZOLOMIDE Capsules are taken by mouth, on an empty stomach, at least one hour before a meal.

Swallow the capsule whole with a glass of water. DO NOT open, divide, crush or chew the capsule.

Avoid contact with your skin, eyes, and nose.

You may be given other medicines to prevent nausea and vomiting.

**Overdose:**

If you think you have taken too much APO-TEMOZOLOMIDE contact your health care professional, hospital emergency department or regional Poison Control Centre immediately, even if there are no symptoms.

**Missed Dose:**

APO-TEMOZOLOMIDE Capsules: If you miss a dose, or vomit after taking a dose, contact your doctor for instructions.

**SIDE EFFECTS AND WHAT TO DO ABOUT THEM**

Like all medicines, APO-TEMOZOLOMIDE can have unintended or undesirable, so called side effects.

Contact your doctor immediately if you have a severe allergic reaction (which may include hives, wheezing or other breathing difficulty).

Common side effects of APO-TEMOZOLOMIDE during concomitant and maintenance treatment, in descending order of frequency, include: hair loss, fatigue, nausea (feeling sick), vomiting, loss of appetite or weight, constipation, headache, rash, diarrhoea, blurred vision, anemia (reduction in blood cells), fever, muscle weakness, and sleepiness.

In case of vomiting, ask your doctor about controlling the vomiting, and the best time to take APO-TEMOZOLOMIDE until the vomiting is under control.

APO-TEMOZOLOMIDE treatment can cause a reduction in certain kinds of blood cells. This may cause you to have increased bruising or bleeding, anemia, fever, and/or a reduced resistance to infections. The reduction of blood cells is usually transient, but in some cases may be prolonged, and may lead to a very severe form of anemia (aplastic anemia) which may be life threatening. Your doctor will monitor your blood regularly for any changes, and will decide if any specific treatment is needed. In some cases, your APO-TEMOZOLOMIDE dose will be reduced or discontinued.

If you are receiving APO-TEMOZOLOMIDE for the 42 day regimen, in combination with radiation treatment, your doctor will also prescribe medicine to help prevent a serious form of pneumonia called *Pneumocystis carinii* pneumonia (PCP).

Less common adverse events, in descending order of frequency, include: convulsions, inflammation of the mouth, cough, radiation injury, dizziness, change in taste, abnormal blood values, shortness of breath, confusion/memory impairment, itching, allergic reaction, insomnia, pain, joint pain, skin dryness, skin redness, abdominal pain, bleeding, chills, hearing impairment, speech disorder, tremor, infection, blood sugar elevation, anxiety, depression, emotional lability, and tingling sensation.

Cases of rash with skin swelling, including on the palms of the hands and soles of the feet, have been observed. Tell your doctor if this occurs.

Cases of lung side effects have been observed with APO-TEMOZOLOMIDE. Patients usually present with shortness of breath and cough. Tell your doctor if you notice any of these symptoms.

Cases of painful reddening of the skin and/or blister on the body or the mouth, have been observed. Tell your doctor if you notice any of these symptoms.

ADVISE YOUR DOCTOR OR PHARMACIST OF ANY UNDESIRABLE OR TROUBLESOME EFFECT NOT LISTED.

| <b>SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM</b> |                  |                                     |              |   |
|--|------------------|-------------------------------------|--------------|---|
| Symptom / effect   |                  | Talk with your doctor or pharmacist |              | Stop taking drug and get immediate medical help |
|  |                  | Only if severe                      | In all cases |   |
| <b>Very Common</b>   | Blurred vision   |                                     | √            |   |
|  | Loss of appetite |                                     | √            |   |
|  | Rash             |                                     | √            |   |
|  | Vomiting         |                                     | √            |   |

**IMPORTANT: PLEASE READ**

|                  |  |  |   |   |
|------------------|--|--|---|---|
| <b>Common</b>    | Confusion  |  | √ |   |
|                  | Convulsion   |  | √ |   |
|                  | Diarrhea   |  | √ |   |
|                  | Fever, other signs of infection (such as fever, chills, cough)   |  | √ |   |
|                  | Increased bruising or bleeding   |  | √ |   |
|                  | Loss of weight   |  | √ |   |
|                  | Memory impairment  |  | √ |   |
|                  | <i>Pneumocystis carinii</i> pneumonia (symptoms such as cough that does not go away, trouble breathing, and fever) |  | √ |   |
| <b>Un-common</b> | Severe allergic reactions including: hives, wheezing or other breathing difficulty                                 |  |   | √ |

|                |  |  |   |   |
|----------------|--|--|---|---|
| <b>Unknown</b> | Rash with skin swelling, including on the palms of the hands and soles of the feet (erythema multiforme)                                     |  | √ |   |
|                | Fatigue, pale skin, shortness of breath, rapid heart beat, fever, and bleeding (aplastic anemia)   |  | √ |   |
|                | Painful reddening of the skin and/or blister on the body or the mouth [toxic epidermal necrolysis (TEN), Stevens-Johnson syndrome (SJS)]     |  |   | √ |
|                | Shortness of breath and cough (interstitial pneumonitis)   |  | √ |   |
|                | Jaundice and hepatitis. Liver injury, including liver failure which may be life threatening.   |  | √ |   |
|                | Herpes simplex encephalitis (symptoms such as fever, headache, personality change, seizures, and/or vomiting) which may be life threatening. |  |   | √ |

*This is not a complete list of side effects. For any unexpected effects while taking APO-TEMOZOLOMIDE, contact your doctor or pharmacist.*

## HOW TO STORE IT

### **APO-TEMOZOLOMIDE Capsules:**

Do not use this product after the expiration date on the package.

Store at room temperature 15°C to 30°C. Protect from moisture.

Keep out of reach and sight of children.

Tell your pharmacist if you notice any change in the appearance of the capsules.

### **Reporting Side Effects**

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting <https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada/adverse-reaction-reporting.html>. for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

*NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.*

## MORE INFORMATION

For more information, please contact your doctor, pharmacist or other healthcare professional.

This leaflet plus the full product monograph, prepared for health professionals, can be obtained by contacting DISpedia, Apotex's Drug Information Service at:

1-800-667-4708

This leaflet can also be found at:  
<http://www.apotex.ca/products>.

This leaflet was prepared by Apotex Inc., Toronto, Ontario, M9L 1T9.

Last revised: February 1, 2018