

Product Monograph

Pr Fomepizole for Injection

1.5 g / 1.5 mL (1 g/mL)

Synthetic Alcohol Dehydrogenase Inhibitor

SteriMax Inc.
2770 Portland Drive
Oakville, ON, L6H 6R4

Date of Revision: June 12, 2018

Control No.: 216459

PRODUCT NAME

Pr Fomepizole for Injection

1.5 g/1.5 mL (1 g/mL)

THERAPEUTIC CLASSIFICATION

Synthetic Alcohol Dehydrogenase Inhibitor

ACTION AND CLINICAL PHARMACOLOGY

Mechanism of Action:

Fomepizole for Injection is a competitive inhibitor of alcohol dehydrogenase (ADH). Alcohol dehydrogenase catalyzes the oxidation of ethanol to acetaldehyde. Alcohol dehydrogenase also catalyzes the initial steps in the metabolism of ethylene glycol and methanol to their toxic metabolites.

Ethylene glycol, the main component of most antifreezes and coolants, is metabolized to glycoaldehyde, which undergoes subsequent sequential oxidations to yield glycolate, glyoxylate, and oxalate. Glycolate and oxalate are the metabolic by-products primarily responsible for the metabolic acidosis and renal damage seen in ethylene glycol toxicosis which presents with the following morbidities: nausea/vomiting, seizures, cardiac arrhythmias, stupor, coma, calcium oxaluria, acute tubular necrosis and death, depending on the amount of ethylene glycol ingested and the time elapsed since ingestion. The lethal dose of ethylene glycol in humans is approximately 1.4 mL/kg.

Methanol, the main component of windshield washer fluid, is slowly metabolized via alcohol dehydrogenase to formaldehyde with subsequent oxidation via formaldehyde dehydrogenase to yield formic acid. Formic acid is primarily responsible for the metabolic acidosis and visual disturbances (e.g., decreased visual acuity and potential blindness) associated with methanol poisoning. A lethal dose of methanol in humans is approximately 1-2 mL/kg.

Fomepizole has been shown *in vitro* and *in vivo* to block alcohol dehydrogenase enzyme activity in dog, monkey, and human liver. The relative affinity of fomepizole for human ADH is 80,000 times greater than that of methanol and ethylene glycol, and 8,000 times greater than that of ethanol (3, 18, 30). The concentration of fomepizole at which alcohol dehydrogenase is inhibited by 50% *in vitro* is approximately 0.1 $\mu\text{mol/L}$. The plasma concentrations achieved in humans with the proposed dosage regimen are well above this, with peak concentrations of fomepizole between 100-300 $\mu\text{mol/L}$ (8.2-24.6 mg/L). These levels are achieved with oral or IV fomepizole doses of 10-20 mg/kg. Fomepizole is most effective when given in close proximity to the ethylene glycol or methanol ingestion before significant target organ damage occurs.

Pharmacokinetics:

The plasma half-life of fomepizole varies with dose, even in patients with normal renal function, and has not been calculated.

Distribution

After intravenous infusion, fomepizole rapidly distributes to total body water. The volume of distribution is between 0.6 L/kg and 1.02 L/kg.

Metabolism

In healthy volunteers, only 1-3.5% of the administered dose of fomepizole (7-20 mg/kg oral and IV) was excreted unchanged in the urine, indicating that metabolism is the major route of elimination. In humans, the primary metabolite of fomepizole is 4-carboxypyrazole (approximately 80-85% of administered dose), which is excreted in the urine. Other metabolites of fomepizole observed in the urine are 4-hydroxymethylpyrazole and the N-glucuronide conjugates of 4-carboxypyrazole and 4-hydroxymethylpyrazole.

Excretion

After a single dose, the elimination of fomepizole is best characterized by Michaelis-Menten kinetics with saturable elimination occurring at plasma concentrations of 100-300 $\mu\text{mol/L}$ (8.2-24.6 mg/L).

With multiple doses, fomepizole rapidly induces its own metabolism via the cytochrome P450 mixed-function oxidase system, which produces a significant increase in the elimination rate after about 30-40 hours. After enzyme induction, elimination follows first-order kinetics.

Special Populations

No special pharmacokinetic studies have been performed with respect to pediatric, geriatric, hepatically-impaired, or renally-impaired patients.

Gender: Possible gender differences were not investigated therefore dose adjustments for patient subgroups cannot be recommended.

INDICATIONS AND CLINICAL USE

Fomepizole for Injection is indicated as an antidote for ethylene glycol (such as antifreeze) or methanol (such as windshield washer fluid) poisoning, or for use in suspected ethylene glycol or methanol ingestion, either alone or in combination with hemodialysis (see **DOSAGE AND ADMINISTRATION**).

CONTRAINDICATIONS

Fomepizole for Injection should not be administered to patients with a documented serious hypersensitivity reaction to fomepizole or other pyrazoles.

PRECAUTIONS

General:

Fomepizole for Injection should not be given undiluted or by bolus injection. Venous irritation and phlebosclerosis were noted in two of six normal volunteers given bolus injections of fomepizole (over 5 minutes) at a concentration of 25 mg/mL.

Patients should be closely monitored for anaphylaxis symptoms (such as dyspnea, wheezing, flushing, etc.) as serious, life-threatening hypersensitivity reaction (anaphylaxis) has been reported following fomepizole administration. If such a reaction occurs, fomepizole therapy

should be discontinued immediately and adequate medical treatment should be initiated (see **CONTRAINDICATIONS** and **ADVERSE REACTIONS**).

Minor allergic reactions (mild rash, eosinophilia) have also been reported in a few patients receiving Fomepizole for Injection (see **ADVERSE REACTIONS**). Therefore, patients should be monitored for signs of allergic reactions.

Drug Interactions:

Oral doses of fomepizole (10-20 mg/kg) significantly reduced the rate of elimination of ethanol (by approximately 40%), via alcohol dehydrogenase inhibition, when given to healthy volunteers in moderate doses. Similarly, ethanol decreased the rate of elimination of fomepizole (by approximately 50%) by the same mechanism.

Reciprocal interactions may occur with concomitant use of Fomepizole for Injection and drugs that increase or inhibit the cytochrome P450 system (e.g., phenytoin, carbamazepine, cimetidine, ketoconazole), though this has not been studied.

Pregnancy:

Animal reproduction studies have not been conducted with fomepizole. It is also not known whether Fomepizole for Injection can cause fetal harm when administered to pregnant women or can affect reproduction capacity. Fomepizole for Injection should be given to pregnant women only if clearly needed.

Nursing Mothers:

It is not known whether fomepizole is excreted in human milk. Because many drugs are excreted in human milk, nursing should be discontinued when Fomepizole is administered to nursing women.

Pediatric Use:

Safety and effectiveness in pediatric patients have not been established.

Geriatric Use:

Fomepizole is metabolized by the liver and excreted by the kidneys. The functions of both organs are generally lower in elderly patients; therefore, care should be taken when selecting the dose of fomepizole for elderly patients.

ADVERSE REACTIONS

The most frequent adverse events reported as drug-related or unknown relationship to study drug in the 78 patients and 63 normal volunteers who received fomepizole were headache (14%), nausea (11%), and dizziness, increased drowsiness, and bad taste/metallic taste (6% each). All other adverse events in this population were reported in approximately 3% or fewer of those receiving fomepizole and were as follows:

Body as a Whole: Abdominal pain, fever, pain during fomepizole injection, inflammation at injection site, lumbalgia/backache, hangover.

Cardiovascular: Phlebosclerosis, phlebitis, hypotension.

Gastrointestinal: Vomiting, diarrhea, dyspepsia, heartburn, decreased appetite, transient increase in liver function tests.

Hemic/Lymphatic: Eosinophilia/hypereosinophilia, lymphangitis, anemia.

Central Nervous System (CNS): Lightheadedness, agitation, feeling drunk, facial flush, vertigo, nystagmus, anxiety, “felt strange”, decreased environmental awareness.

Respiratory: Hiccups, pharyngitis.

Skin/Appendages: Application site reaction, rash.

Special Senses: Abnormal smell, speech/visual disturbances, transient blurred vision, roar in ear.

Post-Market Adverse Reactions:

Serious, life-threatening hypersensitivity reaction (anaphylaxis) has been reported following fomepizole administration (see CONTRAINDICATIONS).

SYMPTOMS AND TREATMENT OF OVERDOSAGE

Nausea, dizziness, and vertigo were noted in healthy volunteers receiving 50 and 100 mg/kg

doses of fomepizole (at plasma concentrations of 290-520 mcmol/L, 23.8- 42.7 mg/L). These doses are 3-6 times the recommended dose. This dose-dependent CNS effect was short-lived in most subjects however in one subject it lasted up to 30 hours.

Fomepizole is dialyzable, and hemodialysis may be useful in treating cases of overdose.

DOSAGE AND ADMINISTRATION

Treatment Guidelines:

Treatment of ethylene glycol and methanol poisonings consist of blocking the formation of toxic metabolites using inhibitors of alcohol dehydrogenase, such as fomepizole, and correction of metabolic abnormalities. In patients with high ethylene glycol (≥ 50 mg/dL or ≥ 8.1 mmol/L) or methanol concentrations (≥ 50 mg/dL or ≥ 15.6 mmol/L), significant metabolic acidosis, or renal failure, hemodialysis should be considered in addition to treatment with fomepizole to remove ethylene glycol or methanol and the respective toxic metabolites of these alcohols.

Treatment with Fomepizole for Injection:

Begin Fomepizole for Injection treatment immediately upon suspicion of ethylene glycol or methanol ingestion based on patient disclosure or history and/or anion gap metabolic acidosis, increased osmolar gap, visual disturbances, or oxalate crystals in the urine, **OR** a documented serum ethylene glycol greater than 3.2 mmol/L (20 mg/dL) or methanol concentration greater than 6.2 mmol/L (20 mg/dL).

In addition to specific antidote treatment with Fomepizole for Injection, patients intoxicated with ethylene glycol or methanol should be managed as appropriate for metabolic acidosis, acute renal failure (ethylene glycol), adult respiratory distress syndrome, visual disturbances (methanol) and hypocalcemia. At frequent intervals throughout the treatment, patients poisoned with ethylene glycol should be monitored for ethylene glycol concentrations in serum and urine, and the presence of urinary oxalate crystals. Similarly, serum methanol concentrations should be monitored in patients poisoned with methanol. Hepatic enzymes and white blood cell counts should be monitored during treatment, as transient increases in serum transaminase concentrations and eosinophilia have been noted with repeated fomepizole dosing.

Dosing of Fomepizole for Injection:

A loading dose of 15 mg/kg should be administered, followed by doses of 10 mg/kg every 12 hours for 4 doses, then 15 mg/kg every 12 hours thereafter until ethylene glycol or methanol concentrations are undetectable or have been reduced below 20 mg/dL (3.2 mmol/L for ethylene glycol and 6.2 mmol/L for methanol), and the patient is asymptomatic with normal pH. All doses should be administered as a slow intravenous infusion over 30 minutes (see **Preparation for Intravenous Administration**, below).

Dosage with Hemodialysis:

Hemodialysis should be considered in addition to Fomepizole for Injection in the case of renal failure, significant or worsening metabolic acidosis, or a measured ethylene glycol or methanol concentration of greater than or equal to 50 mg/dL (8.1 mmol/L for ethylene glycol and 15.6 mmol/L for methanol). Patients should be dialyzed to correct metabolic abnormalities and to lower the ethylene glycol concentrations below 50 mg/dL (8.1 mmol/L for ethylene glycol and 15.6 mmol/L for methanol).

The following guidelines for administering Fomepizole for Injection during hemodialysis should be used:

Before dialysis: Administer next scheduled dose if > 6 hours since the last dose. During dialysis: Administer doses every 4 hours.

Post-dialysis: If time since last dose is < 1 hour, then give the next scheduled dose 12 hours from the last dose administered, and then follow the normal dosing schedule (see **Dosing of Fomepizole for Injection**).

If time since last dose is ≥ 1 hour but < 3 hours, then immediately administer 50% of the next scheduled dose, and then follow the normal dosing schedule.

If time since last dose is ≥ 3 hours, then immediately administer 100% of the next scheduled dose, and then follow the normal dosing schedule.

Discontinuation of Fomepizole for Injection Treatment:

Treatment with Fomepizole for Injection may be discontinued when ethylene glycol or methanol concentrations are undetectable or have been reduced below 20 mg/dL (3.2 mmol/L for ethylene glycol and 6.2 mmol/L for methanol), and the patient is asymptomatic with normal pH.

Preparation for Intravenous Administration:

When preparing Fomepizole for Injection solution avoid ocular, dermal, or inhalation exposures. In case of eye or skin exposure, flush immediately with copious amounts of water. Seek medical attention if irritation persists. Prepare solution in well-ventilated area. If accidental inhalation occurs, move to fresh air.

Fomepizole for Injection solidifies at temperatures less than 25°C (77°F). If the Fomepizole for Injection solution has become solid in the vial, the solution should be liquefied by placing the vial under warm running water or by holding in the hand. Solidification does not affect the efficacy, safety, or stability of Fomepizole for Injection. Using sterile technique, the appropriate dose of Fomepizole for Injection should be drawn from the vial with a syringe and injected into **at least 100 mL of sterile 0.9% sodium chloride injection or dextrose 5% injection**. Mix well. The entire contents of the resulting solution should be infused over 30 minutes. Fomepizole for Injection, like all parenteral products, should be inspected visually for particulate matter prior to administration.


PHARMACEUTICAL INFORMATION

Drug Substance:

Common name: fomepizole

Chemical name: 4-methylpyrazole

Structural formula:



Molecular Formula: C₄H₆N₂

Molecular weight: 82.11 g/mol

Description: Fomepizole is a colourless to light red semi-solid to liquid. It may be present in a solid form at room temperature. Solidification does not affect the efficacy, safety or stability of fomepizole.

Composition:

The drug product is 100% bulk fomepizole (1 g/mL) (wt/vol) with no excipients or preservatives.

Stability and Storage Recommendations:

Store at controlled room temperature (20 to 25°C).

Fomepizole for Injection vials are for single use only. Any unused portion should be discarded.

For stability information, please refer to **Parenteral Products, Stability**, below.

Parenteral Products:

Stability: Fomepizole for Injection diluted in 0.9% sodium chloride injection or dextrose 5% injection remains stable and sterile for at least 24 hours when stored refrigerated or at room temperature. Fomepizole for Injection does not contain preservatives. Therefore, maintain sterile

conditions, and after dilution do not use beyond 24 hours. Solutions showing haziness, particulate matter, precipitate, discolouration or leakage should not be used.

AVAILABILITY OF DOSAGE FORMS

Fomepizole for Injection is supplied as a sterile, preservative-free solution for intravenous use. Fomepizole for Injection is supplied in a dual pack carton containing two 1.5 mL (1 g/mL) vials of Fomepizole for Injection and in a single pack carton containing one 1.5 mL (1 g/mL) vial of Fomepizole for Injection.

PHARMACOLOGY

Animal Studies:

Fomepizole has been shown *in vitro* and *in vivo* to block alcohol dehydrogenase enzyme activity in dog, monkey, and human liver. The relative affinity of fomepizole for human ADH is 80,000 times greater than that of methanol and ethylene glycol, and 8,000 times greater than that of ethanol (3, 18, 30).

Fomepizole has been shown to inhibit ethylene glycol toxicity in rats (8), dogs (11, 17, 13), monkeys (9), and cats (12). Fomepizole has been shown to inhibit methanol toxicity in monkeys (4, 5, 24, 25, 26). Fomepizole is most effective when given in close proximity to the exposure of ethylene glycol or methanol, before significant renal damage or visual disturbance has occurred. When used in sufficient doses, fomepizole inhibits the metabolism of ethylene glycol or methanol to their toxic metabolites and ameliorates metabolic acidosis before significant target organ damage develops in relevant animal models.

Animal and *in vitro* studies indicate that the main metabolite, 4-carboxypyrazole, is not active pharmacologically nor are any of the minor metabolites active at concentrations found following therapeutic doses of fomepizole.

In a study of dogs given a lethal dose of ethylene glycol, three animals each were administered fomepizole, ethanol, or left untreated (control group). The three animals in the untreated group

became progressively obtunded, moribund, and died. At necropsy, all three dogs had severe renal tubular damage. Fomepizole or ethanol, given 3 hours after ethylene glycol ingestion, attenuated the metabolic acidosis and prevented the renal tubular damage associated with ethylene glycol intoxication.

Several studies have demonstrated that fomepizole plasma concentrations of approximately 10 mmol/L (0.82 mg/L) in monkeys are sufficient to inhibit methanol metabolism to formate, which is also mediated by alcohol dehydrogenase. Based on these results, peak plasma concentrations of fomepizole in humans in the range of 100 to 300 mmol/L (8.2-24.6 mg/L) have been targeted to assure adequate plasma concentrations for the effective inhibition of alcohol dehydrogenase.

Clinical Studies:

Clinical studies demonstrate that fomepizole is rapidly and widely distributed to the total body water after IV infusion and undergoes non-linear Michaelis-Menten elimination kinetics after a single dose. Following 36 hours of chronic treatment consisting of repeated supplemental doses at 12 hour intervals, fomepizole can induce its own metabolism and larger supplemental doses are needed to maintain blood levels in the therapeutic range. After metabolism is induced, the elimination approximates a first order kinetic model. Fomepizole is readily dialyzed, and dosing recommendations must provide for increased dosing during dialysis.

The main metabolite, 4-carboxypyrazole, is formed from fomepizole at the same rate regardless of the size of the dose of fomepizole, suggesting that saturation kinetics arise in the pathway leading to the formation of 4-carboxypyrazole.

Fomepizole can inhibit the metabolism of ethanol and ethylene glycol, and these substances in turn inhibit the elimination of fomepizole. All three substances are substrate inhibitors of alcohol dehydrogenase. One may expect drug interactions with other alcohol dehydrogenase inhibitors, and induction of the elimination of fomepizole by other enzyme inducers that affect the cytochrome P450 enzyme system.

The efficacy of fomepizole in the treatment of ethylene glycol and methanol intoxication was studied in two prospective, U.S. clinical trials without concomitant control groups. In these trials, 14/16 patients in the ethylene glycol trial and 7/11 patients in the methanol trial underwent

hemodialysis in combination with fomepizole therapy. (See **DOSAGE AND ADMINISTRATION**). All patients received fomepizole shortly after admission.

The results of these two studies provide evidence that fomepizole blocks ethylene glycol and methanol metabolism mediated by alcohol dehydrogenase in the clinical setting. In both studies, plasma concentrations of toxic metabolites of ethylene glycol and methanol (glycolate and formate, respectively) failed to rise. Even though the relationship to fomepizole therapy was confounded by treatment with hemodialysis and significant blood ethanol concentrations, in post-dialysis periods, when ethanol concentrations were insignificant (< 100 mg/dL) and the concentrations of ethylene glycol or methanol ranged between 26 to 60 mg/dL administration of fomepizole alone prevented any rise in glycolate or formate concentrations.

In a separate French trial, 5 patients presented with ethylene glycol concentrations ranging from 46.5 to 345 mg/dL, insignificant ethanol blood concentrations, and normal renal function. These patients were treated with fomepizole alone without hemodialysis, and none developed signs of renal injury.

Ethanol (intravenous and oral) has been used as an alcohol dehydrogenase inhibitor to treat ethylene glycol poisoning. However, the amount given must be carefully controlled, since ethanol itself is a hepatotoxin and CNS depressant. In contrast, fomepizole exhibits a slow and reproducible rate of elimination and has minimal CNS depressant effects.

TOXICOLOGY

Acute Toxicity:

An acute toxicology study has been conducted in the mouse and rat over the dose range of 0.75 to 2.00 g/kg. The oral LD₅₀ (lethal dose, 50%) of fomepizole in the mouse and rat was 1.3 g/kg and 1.4 g/kg, respectively. The most pronounced toxic symptoms were hypnosis in high doses and sedation in low doses. Death occurred between 2 and 24 hours. One rat in the 1.75 g/kg dose group died between 24 and 48 hours post-dose.

However, in another study the oral LD₅₀'s of fomepizole were reported as 0.54 and 0.64 g/kg in the mouse and rat, respectively. The intravenous LD₅₀ was reported to be 0.32 g/kg in both the

mouse and the rat. These dose levels represent significant margins over those proposed in the human therapeutic dose regimen.

Subacute Toxicity:

Based on a dose ranging study, fomepizole was evaluated for systemic toxicity in dogs following 14-days intravenous infusion at doses of 10, 20 and 30 mg/kg/day for 30 minutes every 12 hours. This infusion regimen is generally consistent with the planned label indication, but the duration of this study (14 days) exceeds the maximum therapeutic use in humans.

The results of this study indicated that such a regimen in dogs produces no signs of systemic toxicity or local vascular irritation at a dose of 10 mg/kg (no effect level) twice daily for 14 days. At 20 mg/kg, which is above the maximum planned clinical dose, effects were minimal and were limited to a decrease in serum triglycerides, an increase in serum sodium (males only), an increase in bicarbonate, and an increase in urine output (increased volume and decreased specific gravity). This dose of 20 mg/kg was considered a no-toxic-effect level in the dog. A dose of 30 mg/kg was a clear effect level, with the liver identified as the primary target organ (increased alkaline phosphatase and alanine transaminase (ALT), decreased triglycerides and increased liver weights). Also noted at this dose were decreased food consumption, decreased potassium, increased sodium and bicarbonate, and increased urine output. Even at this dose level, none of the effects were life threatening, and most (all except alkaline phosphatase and ALT) were reversible following a 28 day recovery period.

Plasma concentrations determined in this study indicate that, at the no effect level of 10 mg/kg, drug levels observed during the infusion were similar to those seen in humans.

The results of this study indicate that, under the planned conditions of clinical use and considering the short term use and life-saving potential of the drug, fomepizole is safe for the intended indication and does not represent a significant risk to the patient.

Chronic Toxicity:

A series of experiments were conducted to evaluate the chronic toxicity in young male Cynomolgus monkeys. The objective of the first series of experiments was to assess the toxicity

of fomepizole after dosing for six weeks in 12 monkeys. Toxicity assessments included clinical signs, hematology, and blood chemistry, and gross and microscopic pathology were evaluated.

Additionally, ophthalmoscopy with assessment of the fundus structures and recordings of the electroretinogram (ERG) were performed. The objective of the second series of experiments was to identify any direct effect of fomepizole on the eye of the monkey as assessed by ERG.

Except for one monkey who died during anesthesia, the administration of fomepizole induced no clinical toxic reactions, regardless of the dose administered and no visual disturbances were observed. The results of these experiments supported that an initial loading dose of fomepizole administered at a dose of 15-20 mg/kg inhibited methanol oxidation in the monkey model. Furthermore, it was concluded that the results of these experiments clearly indicated that fomepizole would be an attractive substitute for ethanol in the treatment of methanol poisoning.

Carcinogenesis:

There have been no long-term studies performed in animals to evaluate carcinogenic potential.

Mutagenesis:

There was a positive Ames test result in the *Escherichia coli* tester strain WP2uvrA and the *Salmonella typhimurium* tester strain TA102 in the absence of metabolic activation using fomepizole concentrations of 100, 333, 1000, 3330, and 5000 mcg/plate.

In addition, the mutagenicity of fomepizole was tested in the *in vivo* mouse micronucleus assay. In the 35 animals tested, dosed at 75, 150, and 300 mg/kg, the bone marrow showed that fomepizole did not induce a significant increase in the frequency of micronucleated polychromatic erythrocytes; thus, this test did not indicate bone marrow cytotoxicity. The mutagenicity of fomepizole is considered to be negative in this assay.

Impairment of Fertility:

In rats, fomepizole (110 mg/kg) administered orally for 40 to 42 days resulted in decreased testicular mass (approximately 8% reduction). This dose is approximately 0.6 times the human maximum daily exposure based on surface area (mg/m^2). This reduction was similar for rats treated with either ethanol or fomepizole alone. When fomepizole was given in combination with

ethanol, the decrease in testicular mass was significantly greater (approximately 30% reduction) compared to those rats treated exclusively with fomepizole or ethanol.

BIBLIOGRAPHY

1. Barceloux DG, et al. American Academy of Clinical Toxicology Practice Guidelines on the Treatment of Ethylene Glycol Poisoning. Ad Hoc Committee. *J Tox Clin Tox* 1999; 37(5):537-60.
2. Baud, FJ, Bismuth C, Garnier R, et al. 4-Methylpyrazole may be an alternative to ethanol therapy for ethylene glycol intoxication in man. *Clin Toxicol* 1986-87; 24(6):463-483.
3. Blair AH, Vallee BL. Some catalytic properties of human liver alcohol dehydrogenase. *Biochemistry* 1966; 5(6):2026-2034.
4. Blomstrand R, Ostling-Wintzell H, Lof A, McMartin K, Tolf BR, Hedstrom KG. Pyrazoles as inhibitors of alcohol oxidation and as important tools in alcohol research: an approach to therapy against methanol poisoning. *Proc Natl Acad Sci* 1979; 76(7):3499-3503.
5. Blomstrand R, Ingemansson SO, Jensen M, Hedstrom CG. Normal electroretinogram and no toxicity signs after chronic and acute administration of the alcohol dehydrogenase inhibitor 4-methylpyrazole to the cynomolgus monkey (*Macaca Fascicularis*)-a possible new treatment of methanol poisoning. *Drug and Alcohol Dependence* 1984; 13:9-20.
6. Brent J, et al. Fomepizole for the treatment of ethylene glycol poisoning. *NEJM* 1999; 340:832-838.
7. Brent J, McMartin K, Phillips S, Aaron C, Kulig K. Fomepizole For the Treatment of Methanol Poisoning. *New England Journal Med* 2001;344(6):424-429.
8. Chou JY, Richardson KE. The effect of pyrazole on ethylene glycol toxicity and metabolism in the rat. *Toxicology and Applied Pharmacology* 1978; 43:33-44.
9. Clay KL, Murphy RC. On the metabolic acidosis of ethylene glycol intoxication. *Toxicology and Applied Pharmacology* 1977; 39:39-49.
10. Dang Vu B, Crouzier C, Hubert I, Galliot M, Baud FJ, Bourdon R. Analytical and pharmacokinetic study of 4-methylpyrazole, a new antidote for the treatment of ethylene glycol intoxication. *Ann Fais Exp Chim* 1992; 85(906):99-110.
11. Dial SM, Thrall MAH, Hamar DW. Efficacy of 4-methylpyrazole for treatment of ethylene glycol intoxication in dogs. *Am J Vet Res* 1994a; 55(12):1762-70.
12. Dial SM, Thrall MAH, Hamar DW. Comparison of ethanol and 4-methylpyrazole as treatments for ethylene glycol intoxication in cats. *Am J Vet Res* 1994b; 55(12):1771-82.
13. Dial SM, Thrall MA, Hamar DW. 4-methylpyrazole as treatment for naturally acquired ethylene glycol intoxication in dogs. *JAVMA* 1989; 195(1):73-6.

14. Donvan JW, et al. A comparison of fomepizole with hemodialysis vs. fomepizole alone in therapy of severe ethylene glycol toxicity. *J Tox Clin Tox* 1998; 36(5):451-452.
15. Feierman DE, Cederbaum AI. Increased content of cytochrome P-450 and 4-methylpyrazole binding spectrum after 4-methylpyrazole treatment. *Biochem Biophys Res Comm* 1985; 126(3):1076-1081.
16. Gavaler JS, Gay V, Egler K, Van Theil DH. Evaluation of the differential *in vivo* toxic effects of ethanol and acetaldehyde on the hypothalamic-pituitary-gonadal axis using 4-methylpyrazole. *Alcoholism Clin Exper Res* 1983; 7(3):332-336.
17. Grauer GF, Thrall MAH, Henre BA, Hjelle JJ. Comparison of the effects of ethanol and 4-methylpyrazole on the pharmacokinetics and toxicity of ethylene glycol in the dog. *Toxicology Letters* 1987; 35:307-14.
18. Hantson P, Wallemacq P, Brau M, Vanbinst R, Haufroid V, Mahieu P. Two cases of acute methanol poisoning partially treated by oral 4-methylpyrazole. *Intensive Care Med* 1999; 25(5):528-531.
19. Harry P, Turcant A, Bouachour G, Houze P, Alquier P, Allain P. Efficacy of 4-methylpyrazole in ethylene glycol poisoning: clinical and toxicokinetic aspects. *Human & Exper Toxicol* 1994; 13:61-64.
20. Jacobsen D. New treatment for ethylene glycol poisoning editorial (comment). *N Engl J Med* 1999; 340(11):879-81.
21. Jacobsen D, Barron SK, Sebastian S, Blomstrand R, McMartin KE. Non-linear kinetics of 4-methylpyrazole in healthy human subjects. *Eur J Clin Pharmacol* 1989; 37:599-604.
22. Jacobsen D, Sebastian CS, Barron SK, Carriere EW, McMartin KE. Effects of 4-methylpyrazole, methanol/ethylene glycol antidote, in healthy humans. *J Emergency Med* 1990; 8:455-461.
23. Jobard E, Harry P, Turcant A, Roy PM, Allain P. 4-Methylpyrazole and hemodialysis in ethylene glycol poisoning. *Clin Toxicol* 1996; 34(4):373-377.
24. McMartin KE, Makar AB, Martin A, Palese M, Tephly TR. Methanol poisoning I. The role of formic acid in the development of metabolic acidosis in the monkey and the reversal by 4-methylpyrazole. *Biochemical Medicine* 1975; 13:319-33.
25. McMartin KE, Martin-Amat G, Makar AB, Tephly TR. Methanol poisoning: Role of formate metabolism in the monkey. In: Thurman RG, Williamson JR, Drott H, Chance B eds. *Alcohol & Aldehyde Metabolizing Systems*. New York, NY: Academic Press; 1977b; 429-440.
26. McMartin KE, Hedstrom KG, Tolf B-R, Ostling-Wintzell H, Blomstrand R. Studies on the metabolic interactions between 4-methylpyrazole and methanol using the monkey as

- an animal model. *Archives of Biochemistry and Biophysics* 1980; 199(2):606-614.
27. McMartin KE, Brent J, and Meta Study Group. Pharmacokinetics of fomepizole (4MP) in patients. *J Tox Clin Tox* 1998; 36(5):450-451.
 28. Magnusson G, Nyberg JA, Bodin NO, Hansson E. Toxicity of pyrazole and 4-methylpyrazole in mice and rats. *Experientia* 1972; 28(10):1198-200.
 29. Moreau CL, et al. Glycolate kinetics and hemodialysis clearance in ethylene glycol poisoning. META Study Group. *J Tox Clin Tox* 1998; 36(7):659-66.
 30. Pietruszko R. Human liver alcohol dehydrogenase – inhibition of methanol activity by pyrazole, 4-methylpyrazole, 4-hydroxymethylpyrazole and 4-carboxypyrazole. *Biochemical Pharmacology* 1975; 24:1603-1607.
 31. Sivilotti M, et al. Pharmacokinetics of ethylene glycol and methanol during fomepizole therapy: Results of the meta trial. *J Tox Clin Tox* 1998; 36(5):451.
 32. Sivilotti M, Burns M, McMartin K, Brent J. Reversible blindness in methanol poisoning treated with fomepizole. *Journal of Toxicology - Clinical Toxicology* 1998; 36(5):514.
 33. Wax P, et al. Effect of fomepizole (4MP) on ethanol elimination in ethylene glycol (EG) and methanol poisoned patients. *J Tox Clin Tox* 1998; 36(5):451-452.
 34. Antizol® Product Monograph. Paladin Labs Inc., dated October 23, 2017. Control No. 207700.

READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE

PATIENT MEDICATION INFORMATION

Pr Fomepizole for Injection

Read this carefully. This document is a summary and will not tell you everything about this drug. Talk to your healthcare professional about your medical condition and treatment and ask if there is any new information about Fomepizole for Injection.

What is Fomepizole for Injection used for?

- Fomepizole for Injection is an antidote which is a medicine used to counter the effects of a poison.
- It is used as an antidote for ethylene glycol and methanol poisoning.
- Ethylene glycol or methanol poisoning can happen from swallowing substances like antifreeze, coolant or windshield washer fluid.

How does Fomepizole for Injection work?

Fomepizole for Injection blocks an enzyme in the body called alcohol dehydrogenase. This prevents it from breaking down ethylene glycol and methanol into their toxic forms that harm the body.

What are the ingredients in Fomepizole for Injection?

Medicinal ingredient: Fomepizole

Non-medicinal ingredients: None

Fomepizole for Injection comes in the following dosage forms:

Each 1.5 mL vial of Fomepizole for Injection contains 1.5 g of the medicinal ingredient fomepizole (1 g / mL).

Do not use Fomepizole for Injection if:

- you are allergic to fomepizole or other medicines called pyrazoles. An example of a pyrazole is celecoxib, a medicine use to treat pain. If you are not sure whether a drug is a pyrazole, ask your healthcare professional.

To help avoid side effects and ensure proper use, talk to your healthcare professional before you receive Fomepizole for Injection. Talk about any health conditions or problems you may have, including if you are:

- pregnant or think you might be pregnant.
- breastfeeding since you should not breastfeed while taking Fomepizole for Injection.

Tell your healthcare professional about all the medicines you take, including any drugs, vitamins, minerals, natural supplements or alternative medicines.

The following may interact with Fomepizole for Injection:

- Alcohol.
- Drugs that use a system called CYP450 in the body. If you are not sure whether a drug uses the CYP450 system, ask a healthcare professional. Some examples of these drugs are:
 - Cimetidine, a medicine used to treat heartburn and stomach ulcers.
 - Ketoconazole, a medicine used to treat fungal infections.
 - Phenytoin and carbamazepine, medicines used to control seizures.

How to take Fomepizole for Injection:

- Fomepizole for Injection will be given to you by a healthcare professional.
- It will be infused directly into your vein.
- It will be infused over 30 minutes.
- Follow all instruction given to you by your healthcare professional.

Usual dose:

Your doctor will determine the dose of Fomepizole for Injection that is right for you. They will also decide how many doses of Fomepizole for Injection you should get. Your doctor may decide to use Fomepizole for Injection alone or in combination with kidney dialysis.

Overdose:

If you think you have received too much Fomepizole for Injection, contact your healthcare professional, hospital emergency department or regional poison control centre immediately, even if there are no symptoms.

What are possible side effects from using Fomepizole for Injection?

These are not all the possible side effects you may feel when receiving Fomepizole for Injection. If you experience any side effects not listed here, contact your healthcare professional.

Side effects may include:

- abnormal smell
- back pain
- bad or metallic taste in your mouth
- blurred vision
- decreased appetite
- diarrhea
- dizziness, light-headedness
- facial flushing
- feeling anxious or nervous
- feeling drunk or hung-over
- fever
- headache
- heartburn

- hiccups
- low blood pressure
- nausea, vomiting
- pain, redness and swelling at injection site
- rapid, involuntary eye movements
- rash
- roar in the ear
- sleepiness
- sore throat
- speech or visual disturbances
- stomach pain
- upset stomach

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
RARE Anaphylaxis (life-threatening allergic reaction): difficulty breathing, itching, skin rash or hives, swelling of your tongue or throat, warm red skin, wheezing.			✓

If you have a troublesome symptom or side effect that is not listed here or becomes bad enough to interfere with your daily activities, talk to your healthcare professional.

Reporting Side Effects

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada/adverse-reaction-reporting.html>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

Storage:

Store at controlled room temperature (20 to 25 °C). Fomepizole for Injection vials are for single use only. Unused portions should be discarded.

If you want more information about Fomepizole for Injection:

- Talk to your healthcare professional.
- Find the full product monograph that is prepared for healthcare professionals and includes this Patient Medication Information by visiting the Health Canada website (<https://www.canada.ca/en/health-canada.html>); the manufacturer's website (<http://www.sterimaxinc.com>) or by calling SteriMax Inc. at 1-800-881-3550.

This document was prepared by SteriMax Inc.

Last Revised: June 12, 2018