

PRODUCT MONOGRAPH

^{Pr} **TEVA-BETAHISTINE**

Betahistine Dihydrochloride Tablets

8 mg, 16 mg and 24 mg

Anti-vertigo Agent

Teva Canada Limited
30 Novopharm Court
Toronto, Ontario
Canada M1B 2K9
www.tevacanada.com

Date of Revision:
September 10, 2018

Control Number: 218390

Table of Contents

PART I: HEALTH PROFESSIONAL INFORMATION.....	3
SUMMARY PRODUCT INFORMATION	3
INDICATIONS AND CLINICAL USE.....	3
CONTRAINDICATIONS	3
WARNINGS AND PRECAUTIONS.....	4
ADVERSE REACTIONS.....	4
DRUG INTERACTIONS	6
DOSAGE AND ADMINISTRATION	6
OVERDOSAGE	7
ACTION AND CLINICAL PHARMACOLOGY	7
STORAGE AND STABILITY.....	10
DOSAGE FORMS, COMPOSITION AND PACKAGING	10
PART II: SCIENTIFIC INFORMATION.....	11
PHARMACEUTICAL INFORMATION.....	11
CLINICAL TRIALS.....	12
DETAILED PHARMACOLOGY	13
TOXICOLOGY	15
REFERENCES	17
PART III: CONSUMER INFORMATION.....	18

Pr **TEVA-BETAHISTINE**

Betahistine Dihydrochloride Tablets
8 mg, 16 mg and 24 mg

PART I: HEALTH PROFESSIONAL INFORMATION

SUMMARY PRODUCT INFORMATION

Route of Administration	Dosage Form / Strength	ALL Relevant Nonmedicinal Ingredients
Oral	Tablets / 8 mg, 16 mg and 24 mg	Mannitol, microcrystalline cellulose, and sodium stearyl fumarate.

INDICATIONS AND CLINICAL USE

TEVA-BETAHISTINE (betahistine dihydrochloride) is indicated for:

- reducing the episodes of recurrent vertigo associated with Ménière's disease.

Geriatrics (>65 years of age):

Limited data from clinical studies in this patient group is available. However, as with all drugs, caution should be exercised in this patient population (see **WARNINGS AND PRECAUTIONS, Special Populations, Geriatrics (> 65 years of age)**).

Pediatrics (< 18 years of age):

TEVA-BETAHISTINE (betahistine dihydrochloride) is not recommended for use in children below 18 years of age due to insufficient data on safety and efficacy (see **WARNINGS AND PRECAUTIONS, Special Populations, Pediatrics (< 18 years of age)**).

CONTRAINDICATIONS

TEVA-BETAHISTINE is contraindicated in:

- patients with known hypersensitivity to betahistine or to any of the tablet constituents. For a complete listing, see the **DOSAGE FORMS, COMPOSITION AND PACKAGING** section.
- patients with peptic ulcer and in patients with a history of this condition (see **WARNINGS AND PRECAUTIONS, Gastrointestinal**).
- patients with pheochromocytoma.

WARNINGS AND PRECAUTIONS

Respiratory

Although clinical intolerance to betahistine dihydrochloride tablets has been shown in a relatively few patients with bronchial asthma, caution should be exercised when giving the product to asthmatic patients.

Gastrointestinal

Several patients with a history of peptic ulcer have experienced an exacerbation of symptoms while using betahistine dihydrochloride tablets. Although experiments in animals and in humans have shown that the gastrointestinal side effects associated with betahistine dihydrochloride are not related to gastric acid production, TEVA-BETAHISTINE is contraindicated in the presence of patients with peptic ulcer and in patients with a history of this condition.

Special Populations

Pregnant Women

The safety of betahistine dihydrochloride in human pregnancy has not been established. Animal studies are insufficient with respect to effects on pregnancy, embryonal / foetal development, parturition and postnatal development. The potential risk for humans is unknown. Betahistine dihydrochloride should not be used during pregnancy unless the potential benefits to the mother outweigh the potential risk to the fetus.

Nursing Women

There are no clinical studies of the excretion of betahistine dihydrochloride into human milk. Betahistine accumulates in the milk of lactating rat dams. The importance of the drug to the mother should be weighed against the benefits of nursing and the potential risks for the child.

Pediatrics (< 18 years of age)

The safety and efficacy of betahistine dihydrochloride in pediatric patients below 18 years of age have not been evaluated, therefore its use in this population is not recommended.

Geriatrics (> 65 years of age)

Limited data from clinical studies suggest that a dosage adjustment is unlikely to be required in this patient group. In general, however, the risk of adverse reactions to any drug may be greater in elderly patients as they are more likely to have decreased renal and/or hepatic function, and are more likely to be taking concomitant medications. Therefore, as with all drugs, caution should be exercised in this patient population.

Effect on Ability to Drive and Use Machines

Although not studied in patients with Ménière's disease, in clinical studies using normal healthy volunteers without Ménière's disease, 72 mg of betahistine dihydrochloride taken three times daily for three days, had no or negligible effects on driving skills in a driving task. However, Ménière's disease itself can negatively affect the ability to drive and use machines.

Monitoring and Laboratory Tests

There is no known interference between betahistine dihydrochloride and laboratory tests.

ADVERSE REACTIONS

Adverse Drug Reaction Overview

The most common adverse reactions experienced with betahistine dihydrochloride tablets are nausea, dyspepsia and headache.

Clinical Trial Adverse Drug Reactions

Because clinical trials are conducted under very specific conditions the adverse reaction rates observed in the clinical trials may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse drug reaction information from clinical trials is useful for identifying drug-related adverse events and for approximating rates.

Table 1. Undesirable Effects Experienced in Betahistine Dihydrochloride–Treated Patients in Placebo-Controlled Clinical Trials*

	<u>Betahistine Dihydrochloride</u> N = 770 (%)	<u>Placebo</u> N = 751 (%)
<u>Gastrointestinal Disorders</u>		
Dyspepsia	1.0	0.5
Nausea	1.8	0.5
<u>Nervous System Disorders</u>		
Headache	5.3	4.4

* Data represent adverse events reported from 23 placebo-controlled studies where the frequency of the adverse event was reported $\geq 1\%$ in the Betahistine group and at a frequency at least $\geq 0.5\%$ higher than placebo.

Post-Market Adverse Drug Reactions

In addition to those events reported during clinical trials, the following undesirable effects have been reported spontaneously during post-marketing use and in scientific literature. A frequency cannot be estimated from the available data and is therefore classified as 'not known'.

Cardiac Disorders:	Ventricular extrasystoles*, hypotension*, including orthostatic and postural hypotension, tachycardia*
Gastrointestinal Disorders:	Mild gastric complaints (e.g. vomiting, gastrointestinal pain, abdominal distension and bloating)
Immune System Disorders:	Hypersensitivity reactions (e.g. anaphylaxis).
Nervous System Disorders:	Somnolence*, convulsions*, paraesthesia*, confusion and hallucinations*
Respiratory System:	Dyspnoea*
Skin and Subcutaneous Tissue Disorders:	Angioneurotic oedema, urticaria, skin rashes of various types, Pruritus and Stevens Johnson syndrome*

* The causal relationship between betahistine dihydrochloride and the emergence of these events has not been established.

DRUG INTERACTIONS

Overview

No in vivo interaction studies have been performed. In vitro data revealed no inhibition of Cytochrome P450 enzymes.

Drug-Drug Interactions

Table 2. Established or Potential Drug-Drug Interactions

Betahistine Dihydrochloride	Ref	Effect	Clinical comment
Antihistamines	T	Efficacy of either drug affected.	Caution is recommended.
Drugs that inhibit monoamino-oxidase (MAO) including MAO subtype B (e.g. selegiline)	In vitro	Inhibition of betahistine dihydrochloride metabolism.	Caution is recommended when using betahistine dihydrochloride and MAO inhibitors (including MAO-B selective) concomitantly.

Legend: C = Case Study; CT = Clinical Trial; T = Theoretical

Drug-Food Interactions

Following administration of betahistine dihydrochloride to healthy male subjects under fed conditions, the rate of absorption was slowed down and maximum concentrations (C_{max}) decreased, whereas the extent of absorption (AUC_{inf}) was not affected by food.

Drug-Herb Interactions

Interactions with herbal products have not been established.

Drug-Laboratory Interactions

Interactions with laboratory tests have not been established.

DOSAGE AND ADMINISTRATION

Dosing Considerations

Mild gastric complaints (e.g. vomiting, gastrointestinal pain, abdominal distension and bloating), have been observed. These can normally be minimized or eliminated by taking the dose during meals or by lowering the dose.

Hepatic Insufficiency

Betahistine dihydrochloride is highly metabolized primarily by the liver. No pharmacokinetic data are available in patients with hepatic impairment. Caution is advised with use of betahistine dihydrochloride in this population.

Recommended Dose and Dosage Adjustment

The usual daily dosage range is 24-48 mg administered orally in divided doses.

Three Times Daily Dosing:

8 mg tablets: 1 to 2 tablets three times daily.

16 mg tablets: ½ to 1 tablet three times daily.

Twice Daily Dosing

24 mg tablets: 1 tablet twice daily.

Missed Dose

If a dose is missed, the missed dose should not be taken. The next dose should be taken at the usual time. A double dose should never be taken to make up for a forgotten dose.

OVERDOSAGE

For management of a suspected drug overdose, contact your regional Poison Control Centre.

A few overdose cases (up to 640 mg), with mild to moderate symptoms of nausea, dry mouth, dyspepsia, abdominal pain and somnolence have been reported. Presumably, more serious complications (e.g. convulsion, pulmonary or cardiac complications) may occur in cases of intentional overdose of TEVA-BETAHISTINE (betahistine dihydrochloride) above 640 mg, especially in combination with other overdosed drugs. Standard overdose protocol / supportive measures should be followed.

ACTION AND CLINICAL PHARMACOLOGY

Mechanism of Action

The mechanism of action of betahistine dihydrochloride is only partly understood. There are several plausible hypotheses that are supported by animal studies and human data:

Betahistine dihydrochloride affects the histaminergic system: Betahistine dihydrochloride acts both as a partial histamine H₁-receptor agonist and histamine H₃-receptor antagonist in neuronal tissue, and has negligible H₂-receptor activity. Betahistine dihydrochloride increases histamine turnover and release by blocking presynaptic H₃-receptors and inducing H₃-receptor downregulation.

Betahistine dihydrochloride may increase blood flow to the cochlear region: Pharmacological testing in animals has shown that the blood circulation in the striae vascularis of the inner ear improves, probably by means of a relaxation of the precapillary sphincters of the microcirculation of the inner ear.

Betahistine dihydrochloride alters neuronal firing in the vestibular nuclei: Betahistine dihydrochloride was also found to have a dose dependent inhibiting effect on spike generation of neurons in lateral and medial vestibular nuclei.

Pharmacodynamics

In ten healthy male volunteers, single oral doses of 8, 16, and 32 mg of betahistine dihydrochloride given in a placebo-controlled, double-blind cross-over study produced dose-related effects on the vestibular system, as measured by electronystagmography (ENG). Maximal effects on the slow nystagmus phase were found 3 to 4 hours after drug intake. Nystagmus duration was reduced by a mean value of 35% (after 8 mg), 48% (16 mg), or 59% (32 mg); all reductions were statistically significant ($p < 0.0005$).

Eleven patients with Meniere's disease were treated in a three month, open-label study of the pharmacological effects of betahistine dihydrochloride on hearing and ENG-recorded, rotation-induced nystagmus. The study participants took one, 8 mg tablet three times a day (total daily dose, 24 mg). The speed of the quick phase of eye shift pre-treatment versus that achieved at the end of the three month treatment period was used as the parameter of effectiveness in this study. Hearing was evaluated pre- and post-treatment using three pure tone hearing levels (250, 500, 1000 Hz).

Hearing loss was less after treatment but the difference did not achieve statistical significance. At some rates of acceleration and at all rates of deceleration, there was an increase in the mean eye shift per second; this increase reached statistical significance in six of the 12 tests.

Pharmacokinetics

The pharmacokinetic profile of betahistine dihydrochloride was studied in six healthy male volunteers. Tablets containing 8 mg of radio-labelled betahistine dihydrochloride were administered to the subjects following an overnight fast, 30 minutes before a standard breakfast. Urine was collected for at least 56 hours after dosing, and five blood samples were drawn from each volunteer at 1, 2, 3, 8, and 25 hours for the first two volunteers, and at 1, 2, 3, 5.5, and 8 hours for the next four subjects.

Total urinary excretion of the radio-label was 90.7%, and the urinary half life was 3.5 hours. More than 85% of the administered dose was excreted in the urine within 24 hours. Only one primary urinary metabolite was identified - 2-pyridylacetic acid. Maximum plasma levels of radioactivity were attained by the 1 hour sampling time; the plasma half life of the radio-label was 3.4 hours.

Absorption:

Orally administered betahistine dihydrochloride is readily and almost completely absorbed from all parts of the gastro-intestinal tract. Under fed conditions C_{max} is lower compared to fasted conditions. However, total absorption of betahistine dihydrochloride is similar under both

conditions, indicating that food intake only slows down the absorption of betahistine dihydrochloride.

Distribution

The percentage of betahistine dihydrochloride that is bound by blood plasma proteins is less than 5%.

Metabolism:

After absorption, betahistine dihydrochloride is rapidly and almost completely metabolized into 2-pyridylacetic acid (2-PAA; which has no pharmacological activity). After oral administration of betahistine dihydrochloride the plasma (and urinary) levels of betahistine dihydrochloride are very low. Pharmacokinetic analyses are therefore based on 2-PAA measurements in plasma and urine. The plasma concentration of 2-PAA reaches its maximum 1 hour after intake and declines with a half-life of about 3.5 hours.

Excretion:

2-PAA is readily excreted in the urine. In the dose range between 8 and 48 mg, about 85% of the original dose is recovered in the urine. Renal or fecal excretion of betahistine dihydrochloride itself is of minor importance.

Recovery rates are constant over the oral dose range of 8 to 48 mg indicating that the pharmacokinetics of betahistine dihydrochloride are linear, and suggesting that the involved metabolic pathway is not saturated.

Special Populations and Conditions

Pediatrics

TEVA-BETAHISTINE is not recommended for use in children below 18 years of age (see **WARNINGS and PRECAUTIONS, Pediatrics (< 18 years of age)**).

Geriatrics

Limited data from clinical studies in this patient group is available, however, as with all drugs, caution should be exercised in this patient population (see **WARNINGS and PRECAUTIONS, Geriatrics (> 65 years of age)**).

Gender

No gender related pharmacokinetic differences have been observed in adult patients studied.

Race

Pharmacokinetic differences due to race have not been identified.

Hepatic Insufficiency

No pharmacokinetic data are available in patients with hepatic impairment.

Renal Insufficiency

No pharmacokinetic data are available in patients with renal impairment.

Genetic Polymorphism

No data is available on genetic polymorphism.

STORAGE AND STABILITY

Store tablets at controlled room temperature (15-30°C) and protect from exposure to heat and moisture.

DOSAGE FORMS, COMPOSITION AND PACKAGING

TEVA-BETAHISTINE (betahistine dihydrochloride) 8 mg, 16 mg and 24 mg tablets contain 8 mg, 16 mg and 24 mg of betahistine dihydrochloride respectively. Non-medicinal ingredients include: mannitol, microcrystalline cellulose, and sodium stearyl fumarate.

TEVA-BETAHISTINE (betahistine dihydrochloride) tablets are available as follows:

8 mg tablets: White to almost white, round, flat-faced, bevelled-edge tablets, engraved "N" on one side and "8" on the other side.

16 mg tablets: White to almost white, round, biconvex, scored tablets, engraved "N" on one side and "16" on the scored side.

24 mg tablets: White to almost white, round, biconvex, scored tablets, engraved "N" on one side and "24" on the scored side.

TEVA-BETAHISTINE (betahistine dihydrochloride) 8 mg, 16 mg and 24 mg tablets are available in bottles of 100 tablets and 16 mg and 24 mg tablets are available in blisters of 100 tablets.

PART II: SCIENTIFIC INFORMATION

PHARMACEUTICAL INFORMATION


Drug Substance:

Common Name: Betahistine dihydrochloride

Chemical Name: N-Methyl-2-(2-pyridyl) ethylamine dihydrochloride

Molecular Formula & Molecular Mass: $C_8H_{12}N_2 \cdot 2HCl$ 209.1 g/mol

Structural Formula:



Physicochemical Properties:

Betahistine dihydrochloride is a white to light yellow crystalline powder which is very hygroscopic; with a partition coefficient $\text{Log } P$ (octanol/water) = 0.7. Betahistine dihydrochloride is very soluble in ethanol; freely soluble in water and practically insoluble in isopropyl alcohol. The melting range is between 148 °C and 154°C. The pKa values are, $\text{pKa}_1=3.5$ and $\text{pKa}_2=9.7$.

CLINICAL TRIALS

Comparative Bioavailability Study

This is a blinded, single-dose, randomized, two-period, two-sequence, two-treatment, crossover study to evaluate the comparative bioavailability between Teva-Betahistine 24 mg Tablets (Teva Canada Limited, Canada) and Serc[®] 24 mg Tablets (BGP Pharma ULC, formally Solvay Pharma Inc., Canada) after a single-dose in 24 healthy subjects under fasting conditions. A summary of the bioavailability data based on measurements of betahistine dihydrochloride's major metabolite 2-Pyridyl-Acetic Acid is presented in the following table.

2-Pyridyl-Acetic Acid (1 x 24 mg) From measured data uncorrected for potency Geometric Mean Arithmetic Mean (CV %)				
Parameter	Test*	Reference†	% Ratio of Geometric Means	90 % Confidence Interval
AUC _T (ng*h/mL)	6766.9 6956.2 (25)	6500.5 6623.3 (19)	104.10	99.63 - 108.77
AUC _I (ng*h/mL)	6878.9 7082.6 (25)	6604.7 6734.7 (20)	104.15	99.66 - 108.85
C _{max} (ng/mL)	1374.9 1415.7 (24)	1300.6 1321.3 (19)	105.71	98.53 - 113.42
T _{max} [§] (h)	0.74 (46)	0.76 (45)		
T _{1/2} [§] (h)	3.95 (17)	3.89 (16)		

* Teva-Betahistine 24 mg Tablets (Teva Canada Limited, Canada)

† Serc[®] 24 mg Tablets (BGP Pharma ULC, formally Solvay Pharma Inc., Canada) were purchased in Canada.

§ Expressed as the arithmetic mean (CV%) only

DETAILED PHARMACOLOGY

Animal:

Pharmacodynamics

In vitro:

In studies on a variety of isolated organs and isolated tissues, betahistine dihydrochloride produced responses similar to those induced by histamine. Betahistine dihydrochloride had little or no affinity for histamine H₂-receptors as confirmed by its activity in tests using rat uterine muscle, rabbit and guinea pig hearts and atrial pairs,³ and isolated guinea pig ileum.

In vivo:

In evaluations of its effects on the circulation of the inner ear, betahistine dihydrochloride was found to produce greater effects than histamine. Betahistine dihydrochloride increased blood flow in the labyrinthine arteries of dogs by 60.9% following a dose of 100 mcg. Circulation in the stria vascularis and spiral ligament of guinea pigs and chinchillas was increased by a mean of 50% when betahistine dihydrochloride was administered at doses as low as 0.1 mg/animal. In guinea pigs, cochlear blood flow was increased for 30 minutes following a dose of 0.2 mg/kg.

In cats, betahistine dihydrochloride was also found to have a dose dependent inhibiting effect on spike generation of neurons in lateral and medial vestibular nuclei. Betahistine dihydrochloride at 60 and 120 times the normal human dose accelerates the vestibular recovery after unilateral neurectomy. Taken together these properties may contribute to its beneficial therapeutic effects in Ménière's disease and vestibular vertigo.

Betahistine dihydrochloride increases histamine turnover and release likely by blocking presynaptic H₃-receptors and inducing H₃-receptor downregulation. This effect on the histaminergic system could provide an explanation for the efficacy of betahistine dihydrochloride in the treatment of vertigo and vestibular diseases.

Studies of the effects on the cardiovascular system, the pulmonary system, the renal system, the gastrointestinal system, and the central nervous system all indicated that betahistine dihydrochloride produced effects similar to, but less potent than, those of histamine.

Following rapid intravenous (iv) administration of betahistine dihydrochloride, a brief fall in blood pressure was evoked in rats, guinea pigs, dogs, and cats. This hypotensive effect could be blocked by the administration of histamine H₁-antagonists, but not by histamine H₂-antagonists. When betahistine dihydrochloride was injected slowly into anaesthetized dogs, general blood pressure decreased while basilar blood flow increased by up to 200%. Pronounced increases in blood flow were found in the coronary (225%), labyrinthine (161%), and communicating hepatic arteries (156%).

Intravenous doses of 0.2 to 0.4 mg/kg given to anaesthetized guinea pigs produced an increase in pulmonary resistance while large doses induced bronchoconstriction. Parenteral administration to

guinea pigs at a dose of 0.2 mg/kg induced vasopermeability. Administration of an intraperitoneal dose of 40 mg/kg caused death by respiratory failure.

In the perfused baboon kidney, the addition of betahistine dihydrochloride to the perfusate produced increases in urine flow, osmotic clearance, urea and creatinine clearance.

Betahistine dihydrochloride doses of 80 to 1600 mcg/kg/min administered as a continuous infusion to dogs with Heidenhain pouches produced a slight increase in the rate of acid secretion corresponding to 8.8% to 17.6% of the maximum response to histamine. In dogs with gastric fistulae, an increase in acid secretion was obtained with a subcutaneous dose of 20 mg/kg betahistine dihydrochloride. This increase corresponded with that produced by 30 mcg/kg of histamine.

The effect of betahistine dihydrochloride on continuous avoidance behaviour in rats was compared with that of histamine. Betahistine dihydrochloride injected intraventricularly at a dose of 0.32 mg/animal and histamine at a dose of 0.08 mg/animal produced a significant increase in the avoidance rate; an increase was also observed after an intraperitoneal dose of 4.0 mg/kg betahistine dihydrochloride, but this did not achieve statistical significance. Betahistine dihydrochloride did not affect the righting reflex when given to newborn chicks at a dose of 100 mg/kg, while a dose of 50 mg/kg histamine produced sleep characterized by loss of the righting reflex.

Pharmacokinetics

The absorption, distribution, metabolism, and excretion of betahistine dihydrochloride were studied in female rats.

Absorption

After oral administration, betahistine dihydrochloride is completely absorbed from the gastrointestinal tract.

Distribution

The distribution of radio-labelled betahistine dihydrochloride was evaluated at 0.5, 1, 3, 6, 24 or 48 hours after the oral or intravenous administration of a 1 mg dose. Following intravenous administration, radioactivity was distributed rapidly throughout the body, with immediate and intensive secretion into the stomach and intestines. There was a transient accumulation of radioactivity in the liver and the portal vein. After oral administration, radioactivity was distributed throughout the body, with high accumulation in the stomach and intestines. Levels of radioactivity in excess of blood levels were observed in the bronchial epithelium, the eye, and the preputial gland. At 24 hours, only the preputial gland and the alimentary system still showed evidence of accumulated radioactivity. By 48 hours, there was no remaining activity.

Metabolism

The metabolite pattern in rat urine showed only one main metabolite – 2 pyridylacetic acid.

Excretion

Following the oral or intravenous administration of 0.5 mg radio-labelled compound, total excretion of the radioactive label was 80 to 90%, about 67% of which was in the urine. Of the total excretion which took place, 98.5% appeared in the urine within 24 hours.

TOXICOLOGY

Acute Toxicity:

The oral LD₅₀ (Lethal Dose; LD₅₀ is the amount of compound, given at a single dose, which causes the death of 50% [one half] of a group of test animals) for betahistine dihydrochloride is 3040 mg/kg in the albino rat. The intravenous LD₅₀ is 5.1 mg/kg in the rabbit. Side effects in the nervous system were seen in dogs and baboons after intravenous doses at and above 120 mg/kg. Signs of toxicity included ataxia, salivation, inactivity, hyperpnoea, tremors, and cyanosis. Severe gastroenteritis was noted during pathology.

Long-Term Toxicity

Rat and Dog

In a six month study, dogs were given doses of up to 25 mg/kg/day. There were no significant abnormalities noted in any of the parameters assessed. In rats given doses of up to 500 mg/kg/day for 18 months, there were no significant abnormalities noted in any of the parameters assessed. Oral dosing up to and above 250 mg/kg, in dogs and in rats respectively, of betahistine dihydrochloride administered during 3 months did not result in adverse effects.

In investigational studies with betahistine dihydrochloride in rats over 6 months with doses starting at 13 mg/kg and above, hyperemia in some tissues was reported, namely liver, spleen and kidneys. Data presented in the literature are limited, therefore, the impact of this finding is not clear.

Dog and Baboon

Emesis was observed at 300 mg/kg and 120 mg/kg following oral and iv dosing respectively in dogs and sporadically in baboons.

Mutagenicity and Carcinogenicity

In the studies conducted with betahistine dihydrochloride, no mutagenic effects have been observed.

Special carcinogenicity studies were not performed with betahistine dihydrochloride. However, in two 18 month chronic toxicity studies in rats there was no indication of any tumors, neoplasms or hyperplasia in the histopathological examination. Therefore, betahistine dihydrochloride up to a dose of 500 mg/kg did not show any evidence for carcinogenicity in these limited 18 month studies.

Reproduction and Teratology:

Limited data are available for betahistine dihydrochloride on reproduction. In a one-generation study in rats, an oral dose of approximately 250 mg/kg/day betahistine dihydrochloride had no adverse effect on male and female fertility, implantation of foetuses, parturition and viability of pups during lactation. No abnormalities were noted in weaned rats. In pregnant rabbits treated orally with 10 or 100 mg/kg betahistine dihydrochloride from mating (gestation day 0) to gestation day 28, slight increases in fetal loss were observed in the two test groups and a slight increase in the incidence of reduced or unossified sternebrae and extra ribs were observed compared to untreated rabbits. Both effects were non-significant and within historical control data. No adverse effects were noted on implantations, vitality or weight of foetuses.

REFERENCES

1. Bertrand R.A. Meniere's disease: Subjective and objective evaluation of medical treatment with betahistine HCl. *Acta Otolaryngol Suppl* 1970; 305:48-69.
2. Burkin A. Betahistine treatment of Meniere's syndrome. *Clin Med* 1967; 74:41-48.
3. Frew I.J.C., Menon G.N. Betahistine hydrochloride in Menière's disease. *Postgrad Med J* 1984; 52:501-503.
4. Martinez D.M. The effect of SERC (betahistine dihydrochloride) on the circulation of the inner ear in experimental animals. *Acta Otolaryngol* 1970; Suppl 305:29-47.
5. Oosterveld W.J. Betahistine dihydrochloride in the treatment of vertigo of peripheral vestibular origin. A double-blind placebo-controlled study. *J Laryngol Otol* 1984; 98:37-41.
6. Wilmot T.J. An objective study of the effect of betahistine dihydrochloride on hearing and vestibular function tests in patients with Meniere's disease. *J Laryngol Otol* 1971; 85:369-73.
7. Wilmot T.J., Menon G.N. Betahistine in Menière's disease. *J Laryngol Otol* 1976; 90:833-840.
8. Product Monograph for Serc[®] (Betahistine Dihydrochloride) Control # 203694 by BGP Pharma ULC. Date of Revision: August 1, 2017
9. Comparative bioavailability study between Teva-Betahistine 24 mg Tablets (Teva Canada Limited, Canada) and Serc[®] 24 mg Tablets (Solvay Pharma Inc., Canada) after a single-dose in healthy subjects under fasting conditions (Study #: 2004-864).

PART III: CONSUMER INFORMATION

Pr TEVA-BETAHISTINE
Betahistine Dihydrochloride Tablets

This leaflet is part III of a three-part "Product Monograph" published when TEVA-BETAHISTINE was approved for sale in Canada and is designed specifically for Consumers. This leaflet is a summary and will not tell you everything about TEVA-BETAHISTINE. Contact your doctor, nurse, or pharmacist if you have any questions about the drug.

ABOUT THIS MEDICATIONWhat the medication is used for:

TEVA-BETAHISTINE is used for reducing the episodes of recurrent vertigo (dizziness) associated with Ménière's disease.

What it does:

TEVA-BETAHISTINE is a type of medicine called a histamine analogue. It is thought to work by helping the blood flow in your inner ear, which lowers the build up of pressure. It is this build up of pressure in the ear which is thought to cause feelings of dizziness (vertigo), feelings of being sick (nausea and vomiting), ringing in the ears (tinnitus), and hearing loss or hearing difficulty suffered by people with Ménière's disease.

When it should not be used:

Do not take TEVA-BETAHISTINE if:

- you are allergic (hypersensitive) to betahistine dihydrochloride or to any of the non-medicinal ingredients in the product
- you have a peptic (stomach) ulcer or have a history of this condition.
- your doctor has told you that you have an adrenal gland tumour (called pheochromocytoma).

What the medicinal ingredient is:

Betahistine Dihydrochloride

What the nonmedicinal ingredients are:

Mannitol, microcrystalline cellulose, and sodium stearyl fumarate.

What dosage forms it comes in:

Tablets: 8 mg, 16 mg and 24 mg tablets

WARNINGS AND PRECAUTIONS

BEFORE you use TEVA-BETAHISTINE talk to your doctor, nurse or pharmacist if you:

- have or have had health problems in the past.
- you are taking medications, including ones you can take without a doctor's prescription e.g. natural or herbal remedies.

- suffer from bronchial asthma, since asthmatic patients should use TEVA-BETAHISTINE with caution.
- have liver problems.
- are pregnant, plan to become pregnant, or are breastfeeding
- are less than 18 years old.

Driving and using machines: TEVA-BETAHISTINE is not likely to affect your ability to drive or use tools or machinery. However, remember that the disease for which you are being treated (recurrent vertigo associated with Ménière's disease) can make you feel dizzy or sick, and can affect your ability to drive and use machines. Do not drive or engage in hazardous activities if you are experiencing these symptoms.

INTERACTIONS WITH THIS MEDICATION

As with most medicines, interactions with other drugs are possible. Tell your doctor, nurse, or pharmacist about all the medicines you take, including drugs prescribed by other doctors, vitamins, minerals, natural supplements, or alternative medicines.

The following may interact with TEVA-BETAHISTINE:

- antihistamines - these may lower the effect of TEVA-BETAHISTINE. In addition, TEVA-BETAHISTINE may lower the effect of antihistamines.
- monoamine-oxidase inhibitors (MAOIs) - used to treat depression or Parkinson's Disease. These may increase the amount of TEVA-BETAHISTINE available in your bloodstream and your doctor may have to change your dose of TEVA-BETAHISTINE.

PROPER USE OF THIS MEDICATION

It is very important to take TEVA-BETAHISTINE exactly as your doctor has instructed. If you are not sure when or how many tablets to take each day, check with your doctor, nurse or pharmacist. It is recommended to take your dose at about the same time every day. TEVA-BETAHISTINE is not recommended for use in children below 18 years of age.

TEVA-BETAHISTINE can be taken with or without food. If TEVA-BETAHISTINE causes upset stomach, take it with meals.

Usual adult dose:

The usual daily dose is 24 – 48 mg daily in divided doses.

TEVA-BETAHISTINE 8 mg tablets - one to two tablets three times a day.

TEVA-BETAHISTINE 16 mg tablets - half to one tablet three times a day.

or

TEVA-BETAHISTINE 24 mg tablets - one tablet twice a day.

IMPORTANT: PLEASE READ

Overdose:

If you think you have taken too much TEVA-BETAHISTINE contact your doctor, nurse, pharmacist, hospital emergency department or regional Poison Control Centre immediately, even if there are no symptoms.

Missed Dose:

If you forget a dose, skip the missed dose. Take your next dose at the usual time. Do not take a double dose to make up for a forgotten dose.

SIDE EFFECTS AND WHAT TO DO ABOUT THEM

Like all medicines, TEVA-BETAHISTINE can cause side effects in some people. Most people have few or no side effects from betahistine dihydrochloride. Side effects that do occur tend to be generally mild and do not last a long time.

Side effects include:

- skin rashes and itching
- stomach pain
- nausea, vomiting
- bloating, indigestion
- headache

After the release of betahistine dihydrochloride on the market, there were few reports of patients experiencing drowsiness, heart palpitations or lightheadedness, faintness or dizziness from a drop in blood pressure during treatment with betahistine dihydrochloride. It is unclear whether they resulted from the use of betahistine dihydrochloride or other unrelated factors.

If any of these affect you severely, tell your doctor, nurse or pharmacist.

SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM

Symptom / effect		Talk with your doctor, nurse or pharmacist		Stop taking drug and seek immediate medical attention
		Only if severe	In all cases	
Rare	Severe allergic reaction: rash, hives, itching, swelling of the face, lips, tongue or throat, difficulty swallowing or breathing			✓

This is not a complete list of side effects. For any unexpected effects while taking TEVA-BETAHISTINE, contact your doctor, nurse or pharmacist.

HOW TO STORE IT

Keep your tablets at room temperature (15 to 30°C), protect from exposure to heat and moisture. Keep out of reach and sight of children.

Reporting Side Effects

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada/adverse-reaction-reporting.html>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

MORE INFORMATION

This document plus the full product monograph, prepared for health professionals can be found by contacting, Teva Canada Limited by:

Phone: 1-800-268-4127 ext. 3;
 Email: druginfo@tevacanada.com; or
 Fax: 1-416-335-4472

This leaflet was prepared by:
 Teva Canada Limited
 30 Novopharm Court
 Toronto, Ontario
 Canada, M1B 2K9
www.tevacanada.com

Last revised: September 10, 2018