

## PRODUCT MONOGRAPH

### ISENTRESS<sup>®</sup>

\* Raltegravir tablets 400 mg

\* Raltegravir chewable tablets, 25 mg, 100 mg

(\* as raltegravir potassium)

### ISENTRESS HD<sup>®</sup>

Raltegravir tablets 600 mg

(as raltegravir potassium)

Human immunodeficiency virus integrase strand transfer inhibitor

Date of Revision:

September 19, 2018


**Merck Canada Inc.**

16750 route Transcanadienne  
Kirkland, QC Canada H9H 4M7  
www.merck.ca

**Submission Control No: 210317**

## Table of Contents

<b>PART I: HEALTH PROFESSIONAL INFORMATION .....</b>	<b>3</b>
SUMMARY PRODUCT INFORMATION.....	3
INDICATIONS AND CLINICAL USE .....	3
CONTRAINDICATIONS.....	4
WARNINGS AND PRECAUTIONS .....	4
ADVERSE REACTIONS .....	7
DRUG INTERACTIONS .....	22
DOSAGE AND ADMINISTRATION .....	29
OVERDOSAGE.....	31
ACTION AND CLINICAL PHARMACOLOGY .....	31
STORAGE AND STABILITY .....	37
DOSAGE FORMS, COMPOSITION AND PACKAGING.....	37
<b>PART II: SCIENTIFIC INFORMATION .....</b>	<b>39</b>
PHARMACEUTICAL INFORMATION.....	39
CLINICAL TRIALS .....	40
TOXICOLOGY.....	56
REFERENCES.....	57
<b>PART III: CONSUMER INFORMATION.....</b>	<b>59</b>

 **ISENTRESS<sup>®</sup>**  
 \* Raltegravir tablets 400 mg  
 \* Raltegravir chewable tablets, 25 mg, 100 mg  
 (\* as raltegravir potassium)

 **ISENTRESS HD<sup>®</sup>**  
 Raltegravir tablets 600 mg  
 (as raltegravir potassium)

## PART I: HEALTH PROFESSIONAL INFORMATION

### SUMMARY PRODUCT INFORMATION

Route of Administration	Dosage Form / Strength	Clinically Relevant Nonmedicinal Ingredients
Oral	Film coated tablets 400 mg and 600 mg raltegravir (as raltegravir potassium)	Lactose monohydrate
	Chewable tablets 25 mg and 100 mg raltegravir (as raltegravir potassium)	Ethanol (trace) Phenylalanine, a component of aspartame Soy Lecithin
		<i>For a complete listing see DOSAGE FORMS, COMPOSITION AND PACKAGING section.</i>

### INDICATIONS AND CLINICAL USE

#### Adults

ISENTRESS<sup>®</sup> and ISENTRESS HD<sup>®</sup> (raltegravir) are indicated in combination with other antiretroviral agents for the treatment of human immunodeficiency virus (HIV-1) infection in adult patients.

#### Pediatrics

ISENTRESS<sup>®</sup> is indicated in combination with other antiretroviral agents for the treatment of HIV-1 infection in children and adolescents 2 years of age and older with body weight at least 7 kg.

ISENTRESS HD<sup>®</sup> is indicated in combination with other antiretroviral agents for the treatment of HIV-1 infection in pediatric patients weighing at least 40 kg.

(see WARNINGS AND PRECAUTIONS, Special Populations, Pediatrics and DOSAGE AND ADMINISTRATION).

## CONTRAINDICATIONS

ISENTRESS<sup>®</sup> is contraindicated in patients who are hypersensitive to any component of this medicine. For a complete listing of components, see the DOSAGE FORMS, COMPOSITION AND PACKAGING section of the product monograph.

## WARNINGS AND PRECAUTIONS

### Skin

#### **Severe Skin and Hypersensitivity Reactions:**

Severe, potentially life-threatening, and fatal skin reactions have been reported. These include cases of Stevens-Johnson syndrome and toxic epidermal necrolysis. Hypersensitivity reactions have also been reported and were characterized by rash, constitutional findings, and sometimes, organ dysfunction, including hepatic failure. Discontinue ISENTRESS<sup>®</sup> and other suspect agents immediately if signs or symptoms of severe skin reactions or hypersensitivity reactions develop (including, but not limited to, severe rash or rash accompanied by fever, general malaise, fatigue, muscle or joint aches, blisters, oral lesions, conjunctivitis, facial edema, hepatitis, eosinophilia, angioedema). Clinical status including liver aminotransferases should be monitored and appropriate therapy initiated. Delay in stopping ISENTRESS<sup>®</sup> treatment or other suspect agents after the onset of severe rash may result in a life-threatening reaction.

### General

Do not substitute ISENTRESS<sup>®</sup> chewable tablets or ISENTRESS<sup>®</sup> 400 mg film-coated tablets for ISENTRESS HD<sup>®</sup> 600 mg film-coated tablets.

### **Drug Interactions**

#### Antacids

It is not recommended to administer antacids (containing aluminum and/or magnesium) with ISENTRESS<sup>®</sup> 400 mg twice daily (see DRUG INTERACTIONS, Drug-Drug Interactions, Effect of Other Agents on the Pharmacokinetics of Raltegravir).

Coadministration of ISENTRESS<sup>®</sup> 1200 mg (2 x 600 mg) once daily with calcium carbonate and aluminum/magnesium containing antacids resulted in reduced raltegravir plasma levels therefore coadministration is not recommended.

#### Atazanavir

Coadministration of ISENTRESS<sup>®</sup> 1200 mg (2 x 600 mg) once daily with atazanavir resulted in increased raltegravir plasma levels therefore coadministration is not recommended (see DRUG INTERACTIONS).

#### Tipranavir/ritonavir

Coadministration of ISENTRESS<sup>®</sup> 1200 mg (2 x 600 mg) once daily with tipranavir/ritonavir could result in decreased raltegravir trough plasma levels therefore coadministration is not recommended (see DRUG INTERACTIONS).

### Strong inducers of drug metabolizing enzymes

Caution should be used when coadministering ISENTRESS<sup>®</sup> 400 mg twice daily with strong inducers of uridine diphosphate glucuronosyltransferase (UGT) 1A1 (e.g., rifampin) due to reduced plasma concentrations of raltegravir (see DRUG INTERACTIONS).

Strong inducers of drug metabolizing enzymes (e.g., rifampin) have not been studied with ISENTRESS<sup>®</sup> 1200 mg (2 x 600 mg) once daily but could result in decreased raltegravir trough plasma level therefore coadministration with ISENTRESS<sup>®</sup> 1200 mg (2 x 600 mg) once daily is not recommended (see DRUG INTERACTIONS).

### Immune Reconstitution Inflammatory Syndrome

During the initial phase of treatment, patients responding to antiretroviral therapy may develop an inflammatory response to indolent or residual opportunistic infections (such as *Mycobacterium avium* complex, cytomegalovirus, *Pneumocystis jirovecii* pneumonia, and tuberculosis, or reactivation of varicella zoster virus), which may necessitate further evaluation and treatment.

Autoimmune disorders (such as Graves' disease, polymyositis and Guillain-Barre syndrome) have also been reported to occur in the setting of immune reconstitution, however the time to onset is more variable, and can occur many months after initiation of treatment.

### Musculoskeletal

Grade 2–4 creatine kinase laboratory abnormalities were observed in patients treated with raltegravir (see ADVERSE REACTIONS). Cases of myopathy and rhabdomyolysis have been reported with raltegravir. A relationship to raltegravir is not clear in a majority of these cases; however there have been isolated post-market reports of myopathy and rhabdomyolysis either with an association to raltegravir or where a relationship to raltegravir could not be ruled out. Use with caution in patients at increased risk of myopathy or rhabdomyolysis, such as patients receiving concomitant medications known to cause these conditions.

### Phenylketonurics

ISENTRESS<sup>®</sup> Chewable Tablets contain phenylalanine, a component of aspartame. Phenylalanine can be harmful to patients with phenylketonuria.

### Special Populations

**Pregnant Women:** Developmental toxicity studies were performed in rabbits (at doses up to 1000 mg/kg/day) and rats (at doses up to 600 mg/kg/day). The highest doses in these studies produced systemic exposures in these species approximately 3-to 4-fold above the exposure at the recommended human dose. No treatment related external, visceral, or skeletal changes were observed in rabbits. Treatment-related increases over controls in the incidence of supernumerary ribs were seen in rats at 600 mg/kg/day (exposures 4.4- fold above the exposure at the recommended human dose). In both rabbits and rats, no treatment related effects on embryonic/fetal survival or fetal weights were observed.

In rats, at a maternal dose of 600 mg/kg/day, mean drug concentrations in fetal plasma were approximately 1.5- to 2.5-fold greater than in maternal plasma at 1 hour and 24 hours postdose,

respectively. In rabbits, at a maternal dose of 1000 mg/kg/day, mean drug concentrations in fetal plasma were approximately 2% of the mean maternal concentration at both 1 and 24 hours postdose. Toxicokinetic studies demonstrated placental transfer of drug in both species.

### **Antiretroviral Pregnancy Registry (APR)**

To monitor maternal-fetal outcomes of pregnant patients exposed to ISENTRESS<sup>®</sup>, an International Antiretroviral Pregnancy Registry has been established.

Physicians are encouraged to report pregnancy cases for inclusion in the registry by calling Merck Canada at 1-800-567-2594.

**Antiretroviral Pregnancy Registry Data:** Based on prospective reports from the APR of over 500 exposures to ISENTRESS<sup>®</sup> 400 mg twice daily during pregnancy resulting in live births (including over 250 exposures in the first trimester), there was no difference between the overall risk of birth defects for raltegravir compared with the background birth defect rate of 2.7% in the U.S. reference population of the Metropolitan Atlanta Congenital Defects Program (MACDP).

There are no data on the use of ISENTRESS<sup>®</sup> 1200 mg (2 x 600 mg) once daily in pregnant women.

**Nursing Women:** It is not known whether raltegravir is secreted in human milk. However, raltegravir is secreted in the milk of lactating rats. In rats, at a maternal dose of 600 mg/kg/day, mean drug concentrations in milk were approximately 3-fold greater than in maternal plasma. Breast-feeding is not recommended while taking ISENTRESS<sup>®</sup>. In addition, it is recommended that HIV-infected mothers not breast-feed their infants to avoid risking postnatal transmission of HIV.

**Pediatrics:** Based on an open-label, multicenter clinical trial in pediatric patients, the safety profile of ISENTRESS<sup>®</sup> 25 mg and 100 mg chewable tablets, and 400 mg tablets in children 2 years of age and older was comparable to that observed in adults (see ADVERSE REACTIONS, Pediatric Adverse Reactions). Safety and effectiveness of ISENTRESS<sup>®</sup> in children under 2 years of age have not been established.

ISENTRESS<sup>®</sup> 1200 mg (2 X 600 mg) once daily has not been studied in pediatric patients. However, population PK modeling and simulation support the use of 1200 mg (2 x 600 mg) once daily in pediatric patients weighing at least 40 kg (see DOSAGE and ADMINISTRATION and CLINICAL PHARMACOLOGY sections).

Use of once daily dosing regimen 1200 mg (2 x 600 mg) is NOT recommended for pediatric patients below 40 kg.

**Geriatrics (≥65 years of age):** Clinical studies of ISENTRESS<sup>®</sup> did not include sufficient numbers of patients aged 65 and over to determine whether they respond differently from younger patients. In general, dose selection for an elderly patient should be cautious, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy.

## ADVERSE REACTIONS

### Clinical Trial Adverse Drug Reactions

*Because clinical trials are conducted under very specific conditions the adverse reaction rates observed in the clinical trials may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse drug reaction information from clinical trials is useful for identifying drug-related adverse events and for approximating rates.*

#### **Adverse Experiences in Treatment-Experienced Adults**

The safety assessment of ISENTRESS<sup>®</sup> in treatment-experienced patients is based on the pooled safety data from the randomized clinical studies P018 and P019, reported using the recommended dose of ISENTRESS<sup>®</sup> 400 mg twice daily in combination with optimized background therapy (OBT) in 462 patients, in comparison to 237 patients taking placebo in combination with OBT. During double-blind treatment, the total follow-up was 1051 patient-years in the group receiving ISENTRESS<sup>®</sup> 400 mg b.i.d. and 322 patient-years in the group receiving placebo.

For patients in the group receiving ISENTRESS<sup>®</sup> 400 mg twice daily + OBT (mean follow-up 118.7 weeks) and the comparator group placebo + OBT (mean follow-up 71.0 weeks) in the pooled analysis for studies P018 and P019, the most commonly reported clinical adverse experiences (>10% in either group), of all intensities and regardless of causality were: diarrhea in 26.6% and 24.9%, nausea in 13.6% and 16.0%, headache in 12.1% and 13.5%, nasopharyngitis in 14.3% and 8.9%, fatigue in 12.1% and 5.9%, upper respiratory tract infection in 15.8% and 10.1%, bronchitis in 12.1% and 6.8%, pyrexia in 9.7% and 13.9%, vomiting in 8.9% and 11.0% of patients, respectively.

Clinical adverse events of all intensities and regardless of causality occurring in  $\geq 2\%$  of treatment-experienced adult patients are presented in Table 1.

**Table 1 – Percentage of Patients with Adverse Experience of All Intensities and Regardless of Causality Occurring in ≥2% of Treatment-Experienced Adult Patients in Either Treatment Group**

System Organ Class, Preferred Term	Randomized Studies P018 and P019			
	ISENTRESS® 400 mg b.i.d. + OBT (N = 462)		Placebo + OBT (N = 237)	
	n	(%)	n	(%)
Patients With One Or More Adverse Experiences	433	(93.7)	213	(89.9)
Patients With No Adverse Experience	29	(6.3)	24	(10.1)
<b>Blood And Lymphatic System Disorders</b>	52	(11.3)	23	(9.7)
Anemia	16	(3.5)	11	(4.6)
Lymphadenopathy	25	(5.4)	7	(3.0)
Neutropenia	6	(1.3)	5	(2.1)
<b>Eye Disorders</b>	44	(9.5)	18	(7.6)
Conjunctivitis	16	(3.5)	1	(0.4)
<b>Gastrointestinal Disorders</b>	262	(56.7)	126	(53.2)
Abdominal Discomfort	6	(1.3)	6	(2.5)
Abdominal Distension	17	(3.7)	8	(3.4)
Abdominal Pain	35	(7.6)	13	(5.5)
Abdominal Pain Upper	21	(4.5)	11	(4.6)
Aphthous Stomatitis	6	(1.3)	5	(2.1)
Constipation	22	(4.8)	2	(0.8)
Diarrhea	123	(26.6)	59	(24.9)
Dyspepsia	15	(3.2)	3	(1.3)
Flatulence	19	(4.1)	8	(3.4)
Gastritis	11	(2.4)	8	(3.4)
Gastroesophageal Reflux Disease	10	(2.2)	3	(1.3)
Hemorrhoids	14	(3.0)	6	(2.5)
Nausea	63	(13.6)	38	(16.0)
Vomiting	41	(8.9)	26	(11.0)
<b>General Disorders And Administration Site Conditions</b>	194	(42.0)	94	(39.7)
Asthenia	20	(4.3)	9	(3.8)
Chest Pain	17	(3.7)	4	(1.7)
Fatigue	56	(12.1)	14	(5.9)
Oedema Peripheral	16	(3.5)	7	(3.0)
Pyrexia	45	(9.7)	33	(13.9)
<b>Infections And Infestations</b>	328	(71.0)	153	(64.6)
Anogenital Warts	18	(3.9)	4	(1.7)
Bronchitis	56	(12.1)	16	(6.8)
Cellulitis	14	(3.0)	5	(2.1)
Folliculitis	16	(3.5)	2	(0.8)
Gastroenteritis	26	(5.6)	8	(3.4)
Genital Herpes	10	(2.2)	6	(2.5)
Herpes Simplex	16	(3.5)	5	(2.1)
Herpes Zoster	34	(7.4)	4	(1.7)
Influenza	33	(7.1)	10	(4.2)
Nasopharyngitis	66	(14.3)	21	(8.9)
Oesophageal Candidiasis	4	(0.9)	6	(2.5)
Onychomycosis	9	(1.9)	5	(2.1)
Oral Candidiasis	10	(2.2)	23	(9.7)
Pharyngitis	18	(3.9)	5	(2.1)
Pneumonia	33	(7.1)	12	(5.1)
Respiratory Tract Infection	19	(4.1)	1	(0.4)
Rhinitis	9	(1.9)	6	(2.5)
Sinusitis	32	(6.9)	10	(4.2)
Tooth Infection	4	(0.9)	5	(2.1)
Upper Respiratory Tract Infection	73	(15.8)	24	(10.1)
Urinary Tract Infection	15	(3.2)	12	(5.1)
<b>Investigations</b>	38	(8.2)	23	(9.7)
Weight Decreased	16	(3.5)	9	(3.8)



System Organ Class, Preferred Term	Randomized Studies P018 and P019			
	ISENTRESS® 400 mg b.i.d. + OBT (N = 462)		Placebo + OBT (N = 237)	
	n	(%)	n	(%)
<b>Metabolism And Nutrition Disorders</b>	75	(16.2)	30	(12.7)
Anorexia	11	(2.4)	7	(3.0)
Decreased Appetite	12	(2.6)	3	(1.3)
Hyperlipidemia	12	(2.6)	1	(0.4)
<b>Musculoskeletal And Connective Tissue Disorders</b>	151	(32.7)	55	(23.2)
Arthralgia	30	(6.5)	10	(4.2)
Back Pain	33	(7.1)	10	(4.2)
Muscle Spasms	17	(3.7)	8	(3.4)
Musculoskeletal Pain	13	(2.8)	2	(0.8)
Myalgia	17	(3.7)	10	(4.2)
Pain In Extremity	31	(6.7)	10	(4.2)
<b>Neoplasms Benign, Malignant And Unspecified (Incl Cysts And Polyps)</b>	70	(15.2)	19	(8.0)
Skin Papilloma	31	(6.7)	9	(3.8)
<b>Nervous System Disorders</b>	156	(33.8)	68	(28.7)
Dizziness	33	(7.1)	6	(2.5)
Headache	56	(12.1)	32	(13.5)
Hypoesthesia	10	(2.2)	4	(1.7)
Neuropathy Peripheral	21	(4.5)	9	(3.8)
Paresthesia	11	(2.4)	5	(2.1)
<b>Psychiatric Disorders</b>	80	(17.3)	42	(17.7)
Anxiety	16	(3.5)	9	(3.8)
Depression	18	(3.9)	12	(5.1)
Insomnia	32	(6.9)	13	(5.5)
<b>Reproductive System And Breast Disorders</b>	52	(11.3)	17	(7.2)
Erectile Dysfunction	11	(2.4)	2	(0.8)
<b>Respiratory, Thoracic And Mediastinal Disorders</b>	99	(21.4)	48	(20.3)
Asthma	8	(1.7)	5	(2.1)
Cough	32	(6.9)	14	(5.9)
Oropharyngeal Pain	15	(3.2)	9	(3.8)
Sinus Congestion	5	(1.1)	5	(2.1)
<b>Skin And Subcutaneous Tissue Disorders</b>	171	(37.0)	70	(29.5)
Eczema	8	(1.7)	8	(3.4)
Erythema	11	(2.4)	4	(1.7)
Lipodystrophy Acquired	16	(3.5)	5	(2.1)
Night Sweats	15	(3.2)	5	(2.1)
Pruritus	18	(3.9)	10	(4.2)
Rash	41	(8.9)	10	(4.2)
Skin Lesion	16	(3.5)	2	(0.8)
<b>Vascular Disorders</b>	54	(11.7)	20	(8.4)
Hypertension	35	(7.6)	9	(3.8)

## Discontinuations

In the pooled analyses for studies P018 and P019, the rates of discontinuation of therapy due to adverse experiences (clinical and laboratory) were 4.5% in patients receiving ISENTRESS® + OBT and 5.5% in patients receiving placebo + OBT.

## Serious Events

The following serious drug related adverse reactions were reported in clinical studies: gastritis, hepatitis, renal failure, genital herpes, accidental overdose.

## Laboratory Abnormalities

### TREATMENT-EXPERIENCED

The percentages of treatment experienced adult patients receiving either ISENTRESS<sup>®</sup> 400 mg twice daily or placebo (both with OBT) in P018 and P019 with selected Grades 2 to 4 laboratory abnormalities that represent a worsening from baseline are presented in Table 2.

**Table 2 – Selected Laboratory Abnormalities Reported in Treatment-Experienced Adult Patients (≥2%)**

Laboratory Parameter Preferred Term	Limit	Grade	Randomized Studies P018 and P019	
			ISENTRESS <sup>®</sup> 400 mg b.i.d. (N = 462) (%)	Placebo (N = 237) (%)
<b>hematology laboratory test</b>				
absolute neutrophil count (10 <sup>9</sup> /L)	0.75–0.999	Grade 2	(4.1)	(5.9)
	0.50–0.749	Grade 3	(3.0)	(3.4)
hemoglobin (mmol/L)	1.16–1.31	Grade 2	(1.3)	(2.5)
platelet count (10 <sup>9</sup> /L)	50–99.999	Grade 2	(3.5)	(5.1)
<b>blood chemistry test</b>				
fasting (non-random) serum LDL-C (mmol/L)	4.13–4.90	Grade 2	(14.5)	(8.2)
	≥4.91	Grade 3	(6.5)	(6.5)
fasting (non-random) serum cholesterol (mmol/L)	6.20–7.77	Grade 2	(20.6)	(16.9)
	>7.77	Grade 3	(11.0)	(6.2)
fasting (non-random) serum triglyceride (mmol/L)	5.65–8.48	Grade 2	(9.6)	(10.3)
	8.49–13.56	Grade 3	(6.3)	(5.8)
	>13.56	Grade 4	(4.5)	(2.2)
fasting (non-random) serum glucose test (mmol/L)	6.95–13.88	Grade 2	(11.3)	(7.5)
	13.89–27.75	Grade 3	(2.9)	(1.3)
total serum bilirubin	1.6–2.5 x ULN	Grade 2	(5.6)	(3.0)
	2.6–5.0 x ULN	Grade 3	(3.0)	(2.5)
serum creatinine	1.4–1.8 x ULN	Grade 2	(3.3)	(3.0)
serum aspartate aminotransferase	2.6–5.0 x ULN	Grade 2	(9.5)	(8.5)
	5.1–10.0 x ULN	Grade 3	(4.3)	(3.0)
serum alanine aminotransferase	2.6–5.0 x ULN	Grade 2	(10.8)	(9.7)
	5.1–10.0 x ULN	Grade 3	(4.8)	(2.5)
serum alkaline phosphatase	2.6–5.0 x ULN	Grade 2	(2.2)	(0.4)
serum pancreatic amylase test	2.1–5.0 x ULN	Grade 3	(4.6)	(3.0)
serum lipase test	1.6–3.0 x ULN	Grade 2	(5.9)	(3.8)
	3.1–5.0 x ULN	Grade 3	(2.0)	(0.8)
serum creatine kinase	6.0–9.9 x ULN	Grade 2	(2.6)	(2.1)
	10.0–19.9 x ULN	Grade 3	(4.1)	(2.5)
	≥20.0 x ULN	Grade 4	(3.0)	(1.3)
ULN = Upper limit of normal range.				

### **Adverse Experiences in Treatment-Naïve Adults**

The safety of ISENTRESS<sup>®</sup> was evaluated in HIV-infected treatment-naïve subjects in 2 Phase III studies: STARTMRK (Protocol 021) evaluated ISENTRESS<sup>®</sup> 400 mg twice daily versus efavirenz, both in combination with emtricitabine (+) tenofovir disoproxil fumarate and ONCEMRK (Protocol 292) evaluated ISENTRESS<sup>®</sup> 1200 mg (2 x 600 mg) once daily versus ISENTRESS<sup>®</sup> 400 mg twice daily, both in combination with emtricitabine (+) tenofovir disoproxil fumarate.

### **STARTMRK (Protocol 021; ISENTRESS<sup>®</sup> 400 mg twice daily)**

The following safety assessment of ISENTRESS<sup>®</sup> in treatment-naïve patients is based on the randomized double-blind active controlled study of treatment-naïve patients, STARTMRK (Protocol 021) with ISENTRESS<sup>®</sup> 400 mg twice daily in combination with a fixed dose of emtricitabine 200 mg (+) tenofovir disoproxil fumarate 245 mg, (N = 281) versus efavirenz (EFV) 600 mg at bedtime in combination with emtricitabine (+) tenofovir disoproxil fumarate (N = 282). During double-blind treatment, the total follow-up for patients receiving ISENTRESS<sup>®</sup> 400 mg twice daily + emtricitabine (+) tenofovir disoproxil fumarate was 1104 patient-years and 1036 patient-years for patients receiving efavirenz 600 mg at bedtime + emtricitabine (+) tenofovir disoproxil fumarate.

Numbers (%) of patients with clinical adverse experiences and with drug-related adverse experiences in the group receiving ISENTRESS<sup>®</sup>, were less frequent than in the group receiving efavirenz based on the nominal p-values (0.325 and <0.001 respectively).

The most commonly reported clinical adverse experiences (>10% in either group), of all intensities and regardless of causality in patients treated with ISENTRESS<sup>®</sup> + emtricitabine (+) tenofovir disoproxil fumarate versus the patients treated with efavirenz + emtricitabine (+) tenofovir disoproxil fumarate, were: diarrhea in 25.6 and 27.0%, nausea in 16.7% and 14.5%, vomiting in 8.2% and 10.6%, headache in 26.0% and 28.4%, fatigue in 9.3% and 13.5%, influenza in 11.7% and 13.5%, nasopharyngitis in 26.7% and 22.3%, upper respiratory tract infection in 21.4% and 20.2%, arthralgia in 8.5% and 11.7%, back pain in 12.1% and 9.9%, dizziness in 16.4% and 38.3%, abnormal dreams in 8.2% and 13.1%, anxiety in 8.9% and 11.0%, depression in 10.3% and 11.7%, insomnia in 15.7% and 14.9%, cough in 16.7% and 12.1%, pyrexia in 15.7% and 13.8% and rash in 7.8% and 13.8% of patients, respectively.

Clinical adverse events of all intensities and regardless of causality occurring in  $\geq 2\%$  of treatment-naïve adult patients are presented in Table 3.

**Table 3 – Percentage of Patients with Adverse Experience of All Intensities and Regardless of Causality Occurring in (≥2%) of Treatment-Naïve Adult Patients in Either Treatment Group**

System Organ Class, Preferred Term	Randomized Study Protocol 021			
	ISENTRESS® 400 mg b.i.d. (N = 281)		Efavirenz 600 mg q.h.s. (N = 282)	
	n	(%)	n	(%)
Patients With One Or More Adverse Experiences	271	(96.4)	276	(97.9)
Patients With No Adverse Experience	10	(3.6)	6	(2.1)
<b>Blood And Lymphatic System Disorders</b>	25	(8.9)	14	(5.0)
Lymphadenopathy	13	(4.6)	5	(1.8)
<b>Ear And Labyrinth Disorders</b>	26	(9.3)	29	(10.3)
Vertigo	7	(2.5)	14	(5.0)
<b>Eye Disorders</b>	18	(6.4)	40	(14.2)
Conjunctivitis	7	(2.5)	14	(5.0)
<b>Gastrointestinal Disorders</b>	169	(60.1)	175	(62.1)
Abdominal Discomfort	6	(2.1)	4	(1.4)
Abdominal Distension	10	(3.6)	7	(2.5)
Abdominal Pain	25	(8.9)	20	(7.1)
Abdominal Pain Upper	8	(2.8)	19	(6.7)
Constipation	5	(1.8)	10	(3.5)
Diarrhea	72	(25.6)	76	(27.0)
Dyspepsia	25	(8.9)	14	(5.0)
Flatulence	14	(5.0)	19	(6.7)
Gastritis	14	(5.0)	12	(4.3)
Gastroesophageal Reflux Disease	9	(3.2)	7	(2.5)
Gingivitis	8	(2.8)	7	(2.5)
Hemorrhoids	11	(3.9)	8	(2.8)
Nausea	47	(16.7)	41	(14.5)
Toothache	10	(3.6)	5	(1.8)
Vomiting	23	(8.2)	30	(10.6)
<b>General Disorders And Administration Site Conditions</b>	101	(35.9)	126	(44.7)
Asthenia	17	(6.0)	16	(5.7)
Chest Pain	8	(2.8)	13	(4.6)
Chills	8	(2.8)	7	(2.5)
Fatigue	26	(9.3)	38	(13.5)
Influenza Like Illness	10	(3.6)	11	(3.9)
Malaise	5	(1.8)	6	(2.1)
Oedema Peripheral	4	(1.4)	9	(3.2)
Pain	11	(3.9)	7	(2.5)
Pyrexia	44	(15.7)	39	(13.8)
<b>Infections And Infestations</b>	222	(79.0)	216	(76.6)
Acarodermatitis	7	(2.5)	6	(2.1)
Acute Sinusitis	6	(2.1)	5	(1.8)
Body Tinea	3	(1.1)	8	(2.8)
Bronchitis	29	(10.3)	30	(10.6)
Ear Infection	6	(2.1)	8	(2.8)
Folliculitis	9	(3.2)	7	(2.5)
Furuncle	2	(0.7)	6	(2.1)
Gastroenteritis	18	(6.4)	19	(6.7)
Gastroenteritis Viral	1	(0.4)	8	(2.8)
Genital Herpes	12	(4.3)	15	(5.3)
Herpes Simplex	10	(3.6)	11	(3.9)
Herpes Zoster	14	(5.0)	16	(5.7)
Influenza	33	(11.7)	38	(13.5)
Nasopharyngitis	75	(26.7)	63	(22.3)
Onychomycosis	13	(4.6)	14	(5.0)
Oral Candidiasis	9	(3.2)	6	(2.1)
Oral Herpes	7	(2.5)	9	(3.2)
Otitis Media	6	(2.1)	7	(2.5)

System Organ Class, Preferred Term	Randomized Study Protocol 021			
	ISENTRESS® 400 mg b.i.d. (N = 281)		Efavirenz 600 mg q.h.s. (N = 282)	
	n	(%)	n	(%)
Pharyngitis	27	(9.6)	26	(9.2)
Pneumonia	8	(2.8)	11	(3.9)
Respiratory Tract Infection	3	(1.1)	6	(2.1)
Rhinitis	9	(3.2)	11	(3.9)
Secondary Syphilis	7	(2.5)	1	(0.4)
Sinusitis	23	(8.2)	23	(8.2)
Syphilis	14	(5.0)	14	(5.0)
Tinea Pedis	8	(2.8)	7	(2.5)
Tonsillitis	10	(3.6)	8	(2.8)
Tooth Abscess	5	(1.8)	6	(2.1)
Upper Respiratory Tract Infection	60	(21.4)	57	(20.2)
Urethritis	7	(2.5)	2	(0.7)
Urinary Tract Infection	8	(2.8)	14	(5.0)
<b>Injury, Poisoning And Procedural Complications</b>	56	(19.9)	56	(19.9)
Contusion	6	(2.1)	9	(3.2)
Laceration	6	(2.1)	5	(1.8)
Ligament Sprain	4	(1.4)	7	(2.5)
Muscle Strain	2	(0.7)	6	(2.1)
<b>Investigations</b>	18	(6.4)	26	(9.2)
Weight Decreased	6	(2.1)	10	(3.5)
<b>Metabolism And Nutrition Disorders</b>	39	(13.9)	54	(19.1)
Decreased Appetite	12	(4.3)	19	(6.7)
Hyperlipidemia	2	(0.7)	10	(3.5)
Hypertriglyceridemia	1	(0.4)	6	(2.1)
<b>Musculoskeletal And Connective Tissue Disorders</b>	95	(33.8)	105	(37.2)
Arthralgia	24	(8.5)	33	(11.7)
Back Pain	34	(12.1)	28	(9.9)
Muscle Spasms	3	(1.1)	7	(2.5)
Musculoskeletal Pain	5	(1.8)	10	(3.5)
Myalgia	11	(3.9)	15	(5.3)
Neck Pain	8	(2.8)	5	(1.8)
Pain in Extremity	18	(6.4)	15	(5.3)
<b>Neoplasms Benign, Malignant And Unspecified (Incl Cysts And Polyps)</b>	33	(11.7)	43	(15.2)
Anogenital Warts	8	(2.8)	13	(4.6)
Kaposi's Sarcoma AIDS Related	2	(0.7)	6	(2.1)
Skin Papilloma	11	(3.9)	11	(3.9)
<b>Nervous System Disorders</b>	134	(47.7)	179	(63.5)
Dizziness	46	(16.4)	108	(38.3)
Headache	73	(26.0)	80	(28.4)
Hypoesthesia	5	(1.8)	7	(2.5)
Memory Impairment	6	(2.1)	0	(0.0)
Migraine	9	(3.2)	3	(1.1)
Paresthesia	8	(2.8)	13	(4.6)
Somnolence	3	(1.1)	22	(7.8)
<b>Psychiatric Disorders</b>	99	(35.2)	127	(45.0)
Abnormal Dreams	23	(8.2)	37	(13.1)
Anxiety	25	(8.9)	31	(11.0)
Depression	29	(10.3)	33	(11.7)
Insomnia	44	(15.7)	42	(14.9)
Nightmare	10	(3.6)	15	(5.3)
Sleep Disorder	2	(0.7)	7	(2.5)
<b>Renal And Urinary Disorders</b>	25	(8.9)	24	(8.5)
Dysuria	9	(3.2)	3	(1.1)

System Organ Class, Preferred Term	Randomized Study Protocol 021			
	ISENTRESS <sup>®</sup> 400 mg b.i.d. (N = 281)		Efavirenz 600 mg q.h.s. (N = 282)	
	n	(%)	n	(%)
<b>Reproductive System And Breast Disorders</b>	35	(12.5)	41	(14.5)
Erectile Dysfunction	13	(4.6)	5	(1.8)
<b>Respiratory, Thoracic And Mediastinal Disorders</b>	105	(37.4)	85	(30.1)
Asthma	3	(1.1)	6	(2.1)
Cough	47	(16.7)	34	(12.1)
Dyspnea	6	(2.1)	9	(3.2)
Nasal Congestion	13	(4.6)	8	(2.8)
Oropharyngeal Pain	22	(7.8)	15	(5.3)
Productive Cough	8	(2.8)	4	(1.4)
Rhinitis Allergic	18	(6.4)	6	(2.1)
Rhinorrhea	10	(3.6)	10	(3.5)
Sinus Congestion	5	(1.8)	10	(3.5)
<b>Skin And Subcutaneous Tissue Disorders</b>	98	(34.9)	140	(49.6)
Acne	10	(3.6)	5	(1.8)
Alopecia	6	(2.1)	5	(1.8)
Dermatitis	9	(3.2)	8	(2.8)
Dermatitis Allergic	2	(0.7)	8	(2.8)
Eczema	9	(3.2)	5	(1.8)
Night Sweats	9	(3.2)	2	(0.7)
Pruritus	12	(4.3)	15	(5.3)
Rash	22	(7.8)	39	(13.8)
Rash Maculo-Papular	2	(0.7)	9	(3.2)
Rash Papular	3	(1.1)	6	(2.1)
Seborrheic Dermatitis	3	(1.1)	9	(3.2)
Skin Lesion	8	(2.8)	7	(2.5)
<b>Vascular Disorders</b>	25	(8.9)	29	(10.3)
Hypertension	18	(6.4)	18	(6.4)

### Discontinuations

In the study P021, the rate of discontinuation of therapy due to adverse reactions (clinical and laboratory) was 5.0% in patients receiving ISENTRESS<sup>®</sup> + emtricitabine (+) tenofovir disoproxil fumarate and 10.0% in patients receiving efavirenz + emtricitabine (+) tenofovir disoproxil fumarate.

### Serious Events

The following serious drug-related adverse reactions were reported in the clinical study, P021: anemia, immune reconstitution inflammatory syndrome, mental disorder, suicide attempt, depression.

### ONCEMRK (Protocol 292; ISENTRESS<sup>®</sup> 1200 mg [2 x 600 mg] once daily)

The safety of ISENTRESS<sup>®</sup> 1200 mg (2 x 600 mg) once daily was based on the analyses of 48-and 96-week data from a randomized double-blind active controlled study in 797 treatment-naïve HIV-1 infected patients, comparing 531 patients receiving ISENTRESS<sup>®</sup> 1200 mg (2 x 600 mg) once daily with 266 patients receiving ISENTRESS<sup>®</sup> 400 mg twice daily, each in combination with emtricitabine (+) tenofovir disoproxil fumarate. The total follow-up for patients on ISENTRESS<sup>®</sup> 1200 mg (2 x 600 mg) once daily was 515.6 patient-years at 48 weeks and 913.3 patient-years at 96 weeks and for ISENTRESS<sup>®</sup>

400 mg twice daily was 257.7 patient years at 48 weeks and 450.1 patient-years at 96 weeks.

The proportion of patients at week 48 with drug-related clinical and laboratory adverse experiences in the group receiving ISENTRESS<sup>®</sup> 1200 mg (2 x 600 mg) once daily, and the group receiving ISENTRESS<sup>®</sup> 400 mg twice daily were generally similar (24.5%, 1.5% versus 25.6%, 1.5%, respectively).

The rates of discontinuation of therapy due to clinical and laboratory adverse experiences at week 48 were 0.8% and 0.4% in patients receiving ISENTRESS<sup>®</sup> 1200 mg (2 x 600 mg) once daily and 2.3% and 0% in patients receiving ISENTRESS<sup>®</sup> 400 mg twice daily.

The most commonly reported clinical adverse experiences at week 48 (>10% in either treatment group), of all intensities and regardless of causality, respectively, were headache (13.4% versus 10.9%), nausea (11.3% versus 9.8%), and diarrhea (10.9% versus 11.3%).

There were no drug-related clinical adverse reactions of moderate to severe intensity occurring in  $\geq 2\%$  of patients reported in either treatment group.

The rates of serious clinical adverse experiences at week 48 were similar between patients receiving ISENTRESS<sup>®</sup> 1200 mg (2 x 600 mg) once daily and in patients receiving ISENTRESS<sup>®</sup> 400 mg twice daily (5.8% versus 9.4%, respectively). The rates of serious drug related clinical adverse experiences were also similar between the treatment groups (0.2% versus 0.8%, respectively).

Overall, the clinical adverse experiences at week 96 were consistent with those observed at week 48.

### **CNS Events**

In treatment-naïve patients (P021) central nervous system (CNS) adverse reactions, as measured by proportion of patients with 1 or more CNS symptoms (described below) were reported significantly less frequently in the group receiving ISENTRESS<sup>®</sup> + emtricitabine (+) tenofovir disoproxil fumarate as compared with the group receiving efavirenz + emtricitabine (+) tenofovir disoproxil fumarate,  $p < 0.001$ ,  $< 0.001$ ,  $< 0.001$  and  $< 0.001$  for cumulative events through Weeks 8, 48, 96 and 156, respectively. In the group receiving ISENTRESS<sup>®</sup>, the percentage of patients with 1 or more CNS symptoms was 20.3% compared to 52.1% in the group receiving efavirenz by Week 8, and 26.3% compared to 58.5% by Week 48, and 28.8% compared to 60.6% by Week 96, and 31.3% compared to 62.4% by Week 156. CNS adverse reactions for this analysis were dizziness, insomnia, concentration impaired, somnolence, depression, nightmare, confusional state, suicidal ideation, nervous system disorder, psychotic disorder, abnormal dreams, suicide attempt, acute psychosis, delirium, depressed level of consciousness, hallucination, auditory hallucination, completed suicide and major depression.

**Table 4 – Number (%) of Patients With Specific Clinical Adverse Experiences (Incidence >0% in One or More Treatment Groups) by System Organ Class – Nervous System Disorders and Psychiatric Disorders – Weeks 96 and 240 respectively**

	Randomized Study Protocol 021							
	ISENTRESS® 400 mg b.i.d.		Efavirenz 600 mg q.h.s.		ISENTRESS® 400 mg b.i.d.			Efavirenz 600 mg q.h.s.
	(N = 281)		(N = 282)		(N = 281)			(N = 282)
	n	(%)	n	(%)	n	(%)	n	(%)
Patients With One Or More Adverse Experiences	81	(28.8)	171	(60.6)	110	(39.1)	181	(64.2)
Patients With No Adverse Experience	200	(71.2)	111	(39.4)	171	(60.9)	101	(35.8)
<b>Nervous System Disorders</b>	24	(8.5)	120	(42.6)	47	(16.7)	124	(44.0)
Dizziness	23	(8.2)	104	(36.9)	46	(16.4)	108	(38.3)
Nervous System Disorder	0	(0.0)	3	(1.1)	0	(0.0)	3	(1.1)
Somnolence	3	(1.1)	22	(7.8)	3	(1.1)	22	(7.8)
<b>Psychiatric Disorders</b>	66	(23.5)	86	(30.5)	84	(29.9)	99	(35.1)
Abnormal Dreams	21	(7.5)	37	(13.1)	23	(8.2)	37	(13.1)
Confusional State	1	(0.4)	1	(0.4)	1	(0.4)	1	(0.4)
Depressed Mood	2	(0.7)	5	(1.8)	2	(0.7)	5	(1.8)
Depression	17	(6.0)	17	(6.0)	29	(10.3)	33	(11.7)
Depressive Symptom	0	(0.0)	1	(0.4)	0	(0.0)	1	(0.4)
Hallucination	0	(0.0)	0	(0.0)	0	(0.0)	1	(0.4)
Hallucination, Auditory	0	(0.0)	1	(0.4)	0	(0.0)	1	(0.4)
Hallucination, Visual	0	(0.0)	1	(0.4)	0	(0.0)	1	(0.4)
Insomnia	34	(12.1)	31	(11.0)	44	(15.7)	42	(14.9)
Major Depression	2	(0.7)	0	(0.0)	2	(0.7)	0	(0.0)
Nightmare	8	(2.8)	14	(5.0)	10	(3.6)	15	(5.3)
Psychotic Disorder	1	(0.4)	0	(0.0)	1	(0.4)	0	(0.0)
Suicidal Behaviour	0	(0.0)	1	(0.4)	0	(0.0)	1	(0.4)
Suicidal Ideation	0	(0.0)	1	(0.4)	1	(0.4)	1	(0.4)
Suicide Attempt	1	(0.4)	0	(0.0)	4	(1.4)	0	(0.0)

## Laboratory Abnormalities

### TREATMENT-NAÏVE

#### STARTMRK (Protocol 021; ISENTRESS® 400 mg twice daily)

The percentages of treatment-naïve adult patients receiving either ISENTRESS® 400 mg twice daily or efavirenz (both with emtricitabine (+) tenofovir disoproxil fumarate) in P021 with selected Grades 2 to 4 laboratory abnormalities that represent a worsening from baseline are presented in Table 5.



**Table 5 – Selected Laboratory Abnormalities Reported in Treatment-Naïve Adult Patients (≥2%)**

Laboratory Parameter Preferred Term	Limit	Grade	Randomized Study P021	
			ISENTRESS® 400 mg b.i.d. (N = 281) (%)	Efavirenz 600 mg q.h.s. (N = 282) (%)
<b>hematology laboratory test</b>				
absolute neutrophil count (10 <sup>9</sup> /L)	0.75–0.999	Grade 2	(2.8)	(5.0)
	0.50–0.749	Grade 3	(2.8)	(1.4)
<b>blood chemistry test</b>				
fasting (non-random) serum LDL-C (mmol/L)	4.14–4.90	Grade 2	(8.5)	(11.5)
	≥4.91	Grade 3	(3.0)	(10.3)
fasting (non-random) serum cholesterol (mmol/L)	6.21–7.76	Grade 2	(12.3)	(19.1)
	>7.76	Grade 3	(0.0)	(6.4)
fasting (non-random) serum triglyceride (mmol/L)	8.47–8.48	Grade 2	(0.7)	(4.9)
fasting (non-random) serum glucose test (mmol/L)	6.99–13.88	Grade 2	(6.6)	(6.0)
total serum bilirubin	1.6–2.5 x ULN	Grade 2	(4.6)	(0.4)
serum aspartate aminotransferase	2.6–5.0 x ULN	Grade 2	(7.5)	(10.4)
	5.1–10.0 x ULN	Grade 3	(4.6)	(2.9)
serum alanine aminotransferase	2.6–5.0 x ULN	Grade 2	(11.0)	(11.8)
	5.1–10.0 x ULN	Grade 3	(1.8)	(2.2)
serum alkaline phosphatase (IU(alk phos)/L)	2.6–5.0 x ULN	Grade 2	(1.1)	(3.2)
ULN = Upper limit of normal range				

Lipids, Change from Baseline

Changes from baseline in fasting lipids are shown in Table 6.

**Table 6 – Lipid Values, Mean Change from Baseline, Protocol 021**

Laboratory Parameter Preferred Term	ISENTRESS <sup>®</sup> 400 mg Twice Daily + Emtricitabine (+) Tenofovir Disoproxil Fumarate N = 207			Efavirenz 600 mg At Bedtime + Emtricitabine (+) Tenofovir Disoproxil Fumarate N = 187		
			Change from Baseline at Week 240			Change from Baseline at Week 240
	Baseline Mean (mmol/L) [mg/dL]	Week 240 Mean (mmol/L) [mg/dL]	Mean Change (mmol/L) [mg/dL]	Baseline Mean (mmol/L) [mg/dL]	Week 240 Mean (mmol/L) [mg/dL]	Mean Change (mmol/L) [mg/dL]
LDL-Cholesterol*	(2,49) [96]	(2,74) [106]	(0,26) [10]	(2,41) [93]	(3,06) [118]	(0,65) [25]
HDL-Cholesterol*	(0,98) [38]	(1,14) [44]	(0,16) [6]	(0,98) [38]	(1,32) [51]	(0,34) [13]
Total Cholesterol*	(4,12) [159]	(4,53) [175]	(0,41) [16]	(4,06) [157]	(5,20) [201]	(1,14) [44]
Triglyceride*	(1,45) [128]	(1,47) [130]	(0,02) [2]	(1,59) [141]	(2,01) [178]	(0,42) [37]
Total: HDL-C ratio	(4,20) [4,18]	(3,97) [3,98]	(2,56) [2,67]	(4,15) [4,13]	(3,95) [3,95]	(3,35) [3,38]
Non-HDL-C	(3,13) [121]	(3,39) [131]	(0,26) [10]	(3,08) [119]	(3,89) [150]	(0,80) [31]

\* Fasting (non-random) laboratory tests at Week 240.

Notes:

N = total number of subjects per treatment group with at least one lipid test result available. The analysis is based on all available data.

If subjects initiated or increased serum lipid-reducing agents, the last available lipid values prior to the change in therapy were used in the analysis. If the missing data was due to other reasons, subjects were censored thereafter for the analysis. At baseline, serum lipid-reducing agents were used in 5% of subjects in the group receiving ISENTRESS<sup>®</sup> and 3% in the efavirenz group. Through Week 240, serum lipid-reducing agents were used in 9% of subjects in the group receiving ISENTRESS<sup>®</sup> and 15% in the efavirenz group.

Through 240 weeks of therapy, ISENTRESS<sup>®</sup> demonstrated minimal effects on serum lipids with small increases in total cholesterol, HDL-C, LDL-C, triglycerides and non-HDL-C. The group treated with efavirenz had a significantly higher mean change from baseline in total cholesterol, HDL-C, LDL-C, triglycerides and non-HDL-C.

### **ONCEMRK (Protocol 292; ISENTRESS<sup>®</sup> 1200 mg [2 x 600 mg] once daily)**

The percentages of patients receiving either ISENTRESS<sup>®</sup> 1200 mg (2 x 600 mg) once daily or ISENTRESS<sup>®</sup> 400 mg twice daily in P292 with selected Grade 2 to 4 laboratory abnormalities that represent a worsening Grade from baseline are presented in Table 7. Overall, the laboratory abnormalities at week 96 were consistent with those observed at week 48.

**Table 7 - Selected Laboratory Abnormalities Reported in Treatment-Naïve Adult Patients (≥2%)**

		Randomized Study P292 week 48	
Laboratory Parameter Preferred Term (Unit)	Limit	Raltegravir 1200 mg Once Daily (N = 531)	Raltegravir 400 mg Twice Daily (N = 266)
Blood Chemistry Test			
Aspartate aminotransferase			
Grade 2	2.6-5.0 x ULN	3.0%	1.9%
Alanine aminotransferase			
Grade 2	2.6-5.0 x ULN	2.5%	0.8%
Lipase			
Grade 2	1.6-3.0 x ULN	5.1%	4.5%
Creatine Kinase			
Grade 2	6.0-9.9 x ULN	3.2%	2.3%
Grade 3	10.0-19.9 x ULN	1.1%	2.6%
ULN = Upper limit of normal range Note: Raltegravir 1200 mg once daily and raltegravir 400 mg twice daily were administered with Truvada*			

**Less Common Clinical Trial Adverse Drug Reactions (<2%) in Adults**

Drug-related clinical adverse reactions occurring in less than 2% of treatment-experienced patients (n = 462) receiving ISENTRESS<sup>®</sup> + OBT and treatment-naïve patients (n = 281) receiving ISENTRESS<sup>®</sup> + emtricitabine (+) tenofovir disoproxil fumarate and of moderate to severe intensity are listed below by system organ class:

**Blood and Lymphatic System Disorders:**

lymph node pain, neutropenia, anemia, lymphadenopathy

**Cardiac Disorders:**

ventricular extrasystoles

**Ear and Labyrinth Disorders:**

vertigo, tinnitus

**Eye Disorders:**

visual impairment

**Gastrointestinal Disorders:**

diarrhea, nausea, abdominal pain, abdominal distension, abdominal pain upper, vomiting, constipation, abdominal discomfort, dyspepsia, flatulence, gastritis, gastroesophageal reflux disease, dry mouth, eructation, erosive duodenitis

**General Disorders and Administration Site Conditions:**

asthenia, fatigue, pyrexia, chills, face edema, peripheral edema, submandibular mass

**Hepatobiliary Disorders:**

hepatitis, hepatitis alcoholic

**Immune System Disorders:**

drug hypersensitivity, immune reconstitution inflammatory syndrome

**Infections and Infestations:**

herpes simplex, genital herpes, gastroenteritis, herpes zoster, folliculitis, lymph node abscess

**Investigations:**

weight increased, weight decreased

**Metabolism and Nutrition Disorders:**

diabetes mellitus, dyslipidemia, increased appetite, decrease appetite, hypercholesterolemia and body fat disorder

**Musculoskeletal and Connective Tissue Disorders:**

arthralgia, myalgia, back pain, musculoskeletal pain, osteoporosis, polyarthrititis, arthritis, neck pain

**Nervous System Disorders:**

dizziness, neuropathy peripheral, paresthesia, somnolence, tension headache, tremor, hypersomnia, memory impairment

**Psychiatric Disorders:**

depression, insomnia, anxiety, abnormal dreams, nightmare, mental disorder, confusional state, major depression, suicide attempt

**Renal and Urinary Disorders:**

nephritis, nephrolithiasis, nocturia, renal failure, tubulointerstitial nephritis

**Reproductive System and Breast Disorders:**

gynecomastia, erectile dysfunction

**Respiratory, Thoracic and Mediastinal Disorders:**

epistaxis

**Skin and Subcutaneous Tissue Disorders:**

lipodystrophy acquired, rash, hyperhidrosis, dermatitis acneiform, erythema, lipohypertrophy, night sweats, rash macular, rash maculo-papular, rash pruritic, xeroderma, prurigo, lipoatrophy, pruritus, acne, alopecia, skin lesion, lipoatrophy

## **Selected Adverse Experiences**

### **Additional neoplasms, benign, malignant and unspecified**

In studies of ISENTRESS<sup>®</sup> 400 mg twice daily, cancers were observed in treatment-experienced patients who initiated ISENTRESS<sup>®</sup> or placebo, both with OBT and in treatment-naïve patients who initiated ISENTRESS<sup>®</sup> or efavirenz, both with emtricitabine (+) tenofovir disoproxil fumarate; several were recurrent. The types and rates of specific cancers were those expected in a highly immunodeficient population (many had CD4+ cell counts below 50 cells/mm<sup>3</sup> and most had prior AIDS diagnoses). The risk of developing cancer in these studies was similar in the group receiving ISENTRESS<sup>®</sup> and the group receiving the comparator.

### **Additional musculoskeletal and connective tissue disorders**

Grade 2–4 creatine kinase laboratory abnormalities were observed in patients treated with ISENTRESS<sup>®</sup> (see Table 2). Myopathy and rhabdomyolysis have been reported. Use with caution in patients at increased risk of myopathy or rhabdomyolysis, such as patients receiving concomitant medications known to cause these conditions.

### **Rash**

Rash occurred more commonly in treatment-experienced patients receiving regimens containing ISENTRESS<sup>®</sup> + darunavir compared to patients receiving ISENTRESS<sup>®</sup> without darunavir or darunavir without ISENTRESS<sup>®</sup>. However, rash that was considered drug related occurred at similar rates for all three groups. These rashes were mild to moderate in severity and did not limit therapy; there were no discontinuations due to rash.

### **Patients with Co-existing Conditions**

#### **Patients Co-infected with Hepatitis B and/or Hepatitis C Virus**

In Phase III studies of ISENTRESS<sup>®</sup>, patients with chronic (but not acute) active hepatitis B and/or hepatitis C co-infection were permitted to enroll provided that baseline liver function tests did not exceed 5 times the upper limit of normal (ULN). In the treatment experienced studies, BENCHMRK 1 and BENCHMRK 2 (Protocol 018 and Protocol 019), 16% of all patients (114/699) were co-infected; in the treatment-naïve studies, STARTMRK (Protocol 021) and ONCEMRK (Protocol 292), 6% (34/563) and 2.9% (23/797), respectively, were co-infected. In general the safety profile of ISENTRESS<sup>®</sup> in patients with hepatitis B and/or hepatitis C virus co-infection was similar to that in patients without hepatitis B and/or hepatitis C virus co-infection, although the rates of AST and ALT abnormalities were somewhat higher in the subgroup with hepatitis B and/or hepatitis C virus co-infection for both treatment groups. At 96-weeks, in treatment-experienced patients, Grade 2 or higher laboratory abnormalities that represent a worsening Grade from baseline of AST, ALT or total bilirubin occurred in 29%, 34% and 13%, respectively, of co-infected patients treated with ISENTRESS<sup>®</sup> as compared to 11%, 13% and 9% of all other patients treated with ISENTRESS<sup>®</sup>. At 240-weeks, in treatment-naïve patients (P021), Grade 2 or higher laboratory abnormalities that represent a worsening Grade from baseline of AST, ALT or total bilirubin occurred in 22%, 44% and 17%, respectively, of co-infected patients treated with ISENTRESS<sup>®</sup> as compared to 13%, 13% and 5% of all other patients treated with ISENTRESS<sup>®</sup>.

### **Pediatric Adverse Reactions**

ISENTRESS<sup>®</sup> has been studied in 126 antiretroviral treatment-experienced HIV-1 infected children and adolescents 2 through 18 years of age, in combination with other antiretroviral

agents in IMPAACT P1066 (see WARNINGS AND PRECAUTIONS, Special Populations, Pediatrics (<16 years of age) and CLINICAL TRIALS, Pediatric Patients). Of the 126 patients, 96 received the recommended dose of ISENTRESS<sup>®</sup>.

In these 96 children and adolescents, the frequency, type and severity of drug related adverse reactions through Week 24 were comparable to those observed in adults.

One patient experienced drug related clinical adverse reactions of Grade 3 psychomotor hyperactivity, abnormal behavior and insomnia; one patient experienced a Grade 2 serious drug related allergic rash.

One patient experienced drug related laboratory abnormalities, Grade 4 AST and Grade 3 ALT, which were considered serious.

### **Post-Market Adverse Drug Reactions**

The following additional adverse experiences have been reported in post-marketed experience without regard to causality:

#### **Blood and Lymphatic System Disorders:**

thrombocytopenia

#### **Hepatobiliary Disorders:**

hepatic failure (with and without associated hypersensitivity) in patients with underlying liver disease and/or concomitant medications

#### **Musculoskeletal and Connective Tissue Disorders:**

rhabdomyolysis

#### **Nervous System Disorders:**

cerebellar ataxia

#### **Psychiatric Disorders:**

depression (particularly in patients with a pre-existing history of psychiatric illness), including suicidal ideation and behaviors, paranoia

#### **Skin and Subcutaneous Tissue Disorders:**

Stevens-Johnson syndrome, drug rash with eosinophilia and systemic symptoms (DRESS)

## **DRUG INTERACTIONS**

### **Overview**

Raltegravir does not inhibit (IC<sub>50</sub>>100 μM) CYP1A2, CYP2B6, CYP2C8, CYP2C9, CYP2C19, CYP2D6 or CYP3A *in vitro*. Moreover, *in vitro*, raltegravir did not induce CYP3A4. A midazolam drug interaction study confirmed the low propensity of raltegravir to alter the

pharmacokinetics of agents metabolized by CYP3A4 *in vivo* by demonstrating a lack of meaningful effect of raltegravir on the pharmacokinetics of midazolam, a sensitive CYP3A4 substrate.

Similarly, raltegravir is not an inhibitor ( $IC_{50} > 50 \mu M$ ) of the UDP-glucuronosyltransferases (UGT) tested (UGT1A1, UGT2B7), and raltegravir does not inhibit P-glycoprotein-mediated transport. Based on these data, ISENTRESS<sup>®</sup> is not expected to affect the pharmacokinetics of drugs that are substrates of these enzymes or P-glycoprotein (e.g., protease inhibitors, NNRTIs, methadone, opioid analgesics, statins, azole antifungals, proton pump inhibitors and anti-erectile dysfunction agents).

### **Drug-Drug Interactions**

#### **Effect of Raltegravir on the Pharmacokinetics of Other Agents**

In drug interaction studies performed using the 400 mg twice daily dose, raltegravir did not have a clinically meaningful effect on the pharmacokinetics of the following: hormonal contraceptives, methadone, maraviroc, tenofovir, midazolam, lamivudine, etravirine, darunavir/ritonavir, telaprevir and boceprevir. In a multiple-dose drug interaction study, ethinyl estradiol and norelgestromin AUC values were 98% and 114%, respectively, when coadministered with raltegravir as compared to when administered without raltegravir. In a multiple-dose drug interaction study, tenofovir AUC and trough concentrations when coadministered with raltegravir were 90% and 87% of values obtained with tenofovir disoproxil fumarate monotherapy. In another drug interaction study, midazolam AUC from coadministration was 92% of the value obtained with midazolam alone. In a Phase II study, lamivudine pharmacokinetics were similar in patients receiving combinations with raltegravir versus with efavirenz.

#### **Effect of Other Agents on the Pharmacokinetics of Raltegravir**

Raltegravir is not a substrate of cytochrome P450 (CYP) enzymes.

Based on *in vivo* and *in vitro* studies, raltegravir is eliminated mainly by metabolism via a UGT1A1-mediated glucuronidation pathway.

#### **Inducers of Drug Metabolizing Enzymes**

Coadministration of ISENTRESS<sup>®</sup> 400 mg twice daily with drugs that are potent inducers of UGT1A1, such as rifampin (an inducer of numerous drug metabolizing enzymes), reduces plasma concentrations of raltegravir. Caution should be used when coadministering ISENTRESS<sup>®</sup> 400 mg twice daily with rifampin or other strong inducers of UGT1A1. (see WARNINGS AND PRECAUTIONS). The impact of other potent inducers of drug metabolizing enzymes, such as phenytoin and phenobarbital, on UGT1A1 is unknown. Other less potent inducers (e.g., efavirenz, nevirapine, rifabutin, glucocorticoids, St. John's Wort, pioglitazone) may be used with the recommended dose of ISENTRESS<sup>®</sup> 400 mg twice daily.

The impact of drugs that are strong inducers of UGT1A1 such as rifampin on ISENTRESS<sup>®</sup> 1200 mg (2 x 600 mg) once daily is unknown, but co-administration is likely to decrease raltegravir trough levels based on the reduction in trough concentrations observed with ISENTRESS<sup>®</sup> 400 mg twice daily; therefore coadministration with ISENTRESS<sup>®</sup>

1200 mg (2 x 600 mg) once daily is not recommended. The impact of other strong inducers of drug metabolizing enzymes, such as phenytoin and phenobarbital, on UGT1A1 is unknown therefore coadministration with ISENTRESS<sup>®</sup> 1200 mg (2 x 600 mg) once daily is not recommended. In drug interaction studies, efavirenz did not have a clinically meaningful effect on the pharmacokinetics of ISENTRESS<sup>®</sup> 1200 mg (2 x 600 mg) once daily, therefore other less potent inducers (e.g., efavirenz, nevirapine, rifabutin, glucocorticoids, St. John's wort, pioglitazone) may be used with ISENTRESS<sup>®</sup> 1200 mg (2 x 600 mg) once daily.

#### Inhibitors of UGT1A1

Coadministration of ISENTRESS<sup>®</sup> 400 mg twice daily with drugs that are known to be potent UGT1A1 inhibitors (e.g., atazanavir; see Table 8) increases plasma levels of raltegravir. However, the degree of increase is modest and combination therapy with these inhibitors was well tolerated in the clinical studies such that no dose adjustment is required for ISENTRESS<sup>®</sup> 400 mg twice daily.

Coadministration of atazanavir with ISENTRESS<sup>®</sup> 1200 mg (2 x 600 mg) once daily significantly increased plasma levels of raltegravir therefore coadministration of ISENTRESS<sup>®</sup> 1200 mg (2 x 600 mg) once daily and atazanavir is not recommended.

#### Antacids

In a drug interaction study in HIV patients, concomitant administration of an antacid (containing divalent metal cations, e.g., aluminium and magnesium) within 6 hours of ISENTRESS<sup>®</sup> 400 mg twice daily reduced raltegravir absorption by chelation, resulting in significantly decreased raltegravir plasma levels. Administration of antacids (containing aluminium and/or magnesium) with ISENTRESS<sup>®</sup> 400 mg twice daily is not recommended. Coadministration of antacids (containing calcium carbonate) with ISENTRESS<sup>®</sup> 400 mg twice daily decreased raltegravir plasma levels; however, this interaction is not considered clinically meaningful. Therefore, antacids (containing calcium carbonate) may be coadministered with ISENTRESS<sup>®</sup> 400 mg twice daily.

Coadministration of ISENTRESS<sup>®</sup> 1200 mg (2 x 600 mg) once daily with aluminum/magnesium and calcium carbonate containing antacids are likely to result in clinically meaningful reductions in the plasma trough levels of raltegravir. Based on these findings, coadministration of aluminum/magnesium and calcium carbonate containing antacids with ISENTRESS<sup>®</sup> 1200 mg 2 x 600 mg) once daily, is not recommended.

#### Agents that Increase Gastric pH

In a drug interaction study in HIV patients, coadministration of agents that increase gastric pH, e.g., famotidine (H2 blocker) or omeprazole (proton pump inhibitor) with ISENTRESS<sup>®</sup> 400 mg twice daily resulted in increased raltegravir plasma levels due to increased solubility of raltegravir. However, since concomitant use of proton pump inhibitors (PPIs) or H2 blockers with ISENTRESS<sup>®</sup> 400 mg twice daily did not result in a unique safety signal in Phase 3 studies, famotidine or omeprazole may be coadministered with ISENTRESS<sup>®</sup>.



### Additional Considerations

In drug interaction studies of ISENTRESS<sup>®</sup> 400 mg twice daily, atazanavir, efavirenz, ritonavir, tenofovir, tipranavir/ritonavir and telaprevir did not have a clinically meaningful effect on the pharmacokinetics of raltegravir. Rifampin, which is a strong inducer of drug metabolizing enzymes, caused a decrease in trough levels of raltegravir (see subsections Inducers of Drug Metabolizing Enzymes and Inhibitors of UGT1A1 above).

No studies have been conducted to evaluate the drug interactions of ritonavir, tipranavir/ritonavir, boceprevir or etravirine with ISENTRESS<sup>®</sup> 1200 mg (2 x 600 mg) once daily. While the magnitudes of change on raltegravir exposure from ISENTRESS<sup>®</sup> 400 mg twice daily by ritonavir, boceprevir or etravirine were small, the impact from tipranavir/ritonavir was greater (GMR C<sub>trough</sub>=0.45, GMR AUC=0.76). Coadministration of ISENTRESS<sup>®</sup> 1200 mg (2 x 600 mg) once daily and tipranavir/ritonavir is not recommended.

Previous studies of ISENTRESS<sup>®</sup> 400 mg twice daily showed that coadministration of tenofovir disoproxil fumarate (a component of Truvada\*) increased raltegravir exposure. Truvada\* was identified to increase raltegravir 1200 mg (2 x 600 mg) once daily bioavailability by 12%, however its impact is not clinically meaningful. Therefore, coadministration of Truvada\* and ISENTRESS<sup>®</sup> 1200 mg (2 x 600 mg) once daily is permitted.

Findings from clinical studies conducted for ISENTRESS<sup>®</sup> 400 mg twice daily to evaluate the effect of raltegravir on coadministered drugs and presented in Table 8 may be used as references to raltegravir 1200 mg once daily, unless otherwise noted.

All interaction studies were performed in adults. Drug interactions are further described below in Table 8.

**Table 8 – Effect of Other Agents on the Pharmacokinetics of Raltegravir in Adults**

Coadministered Drug/Class Name	Coadministered Drug Dose/Schedule	Raltegravir Dose/Schedule	Ratio (90% Confidence Interval) of Raltegravir Pharmacokinetic Parameters with/without Coadministered Drug; No Effect = 1.00				Clinical Comment
			n	C <sub>max</sub>	AUC	C <sub>min</sub>	
<b>Metal Cation Antacids</b>							
aluminum and magnesium hydroxide antacid	20 mL single dose given with raltegravir	400 mg twice daily	25	0.56 (0.42, 0.73)	0.51 (0.40, 0.65)	0.37 (0.29, 0.48)	Concomitant use or staggered administration of antacids (containing aluminium and/or magnesium) with ISENTRESS <sup>®</sup> 400 mg twice daily is not recommended
	20 mL single dose given 2 hours before raltegravir		23	0.49 (0.33, 0.71)	0.49 (0.35, 0.67)	0.44 (0.34, 0.55)	
	20 mL single dose given 2 hours after raltegravir		23	0.78 (0.53, 1.13)	0.70 (0.50, 0.96)	0.43 (0.34, 0.55)	
	20 mL single dose given 6 hours before raltegravir		16	0.90 (0.58, 1.40)	0.87 (0.64, 1.18)	0.50 (0.39, 0.65)	
	20 mL single dose given 6 hours after raltegravir		16	0.90 (0.58, 1.41)	0.89 (0.64, 1.22)	0.51 (0.40, 0.64)	
aluminum and magnesium hydroxide antacid	20 mL single dose given 12 hours after raltegravir	1200 mg single dose	19	0.86 (0.65, 1.15)	0.86 (0.73, 1.03)	0.42 (0.34, 0.52)	Concomitant use of antacids (containing aluminium and/or magnesium) with ISENTRESS <sup>®</sup> 1200 mg once daily is not recommended
calcium carbonate	3000 mg single dose	400 mg twice daily	24	0.48 (0.36, 0.63)	0.45 (0.35, 0.57)	0.68 (0.53, 0.87)	No dose adjustment required for ISENTRESS <sup>®</sup> 400 mg twice daily
calcium carbonate antacid	3000 mg single dose given with raltegravir	1200 mg single dose	19	0.26 (0.21, 0.32)	0.28 (0.24, 0.32)	0.52 (0.45, 0.61)	Concomitant use of antacids (containing calcium

Coadministered Drug/Class Name	Coadministered Drug Dose/Schedule	Raltegravir Dose/Schedule	Ratio (90% Confidence Interval) of Raltegravir Pharmacokinetic Parameters with/without Coadministered Drug; No Effect = 1.00				Clinical Comment
			n	C <sub>max</sub>	AUC	C <sub>min</sub>	
	3000 mg single dose given 12 hours after raltegravir	1200 mg single dose	19	0.98 (0.81, 1.17)	0.90 (0.80, 1.03)	0.43 (0.36, 0.51)	carbonate) with ISENTRESS <sup>®</sup> 1200 mg once daily is not recommended
<b>H2 Receptor Antagonist</b>							
famotidine	20 mg daily <sup>†</sup>	400 mg twice daily	18	1.60 (1.14, 2.24)	1.44 (1.09, 1.90)	1.06 (0.85, 1.34)	No dose adjustment required for ISENTRESS <sup>®</sup> 400 mg twice daily and 1200 mg once daily
<b>Proton Pump Inhibitors</b>							
omeprazole	20 mg daily	400 mg twice daily	18	1.51 (0.98, 2.35)	1.37 (0.99, 1.89)	1.24 (0.95, 1.62)	No dose adjustment required for ISENTRESS <sup>®</sup> 400 mg twice daily and 1200 mg once daily
<b>Antibacterial</b>							
rifampin	600 mg daily	400 mg single dose	9	0.62 (0.37, 1.04)	0.60 (0.39, 0.91)	0.39 (0.30, 0.51)	Use with caution with ISENTRESS <sup>®</sup> 400 mg twice daily (see Warnings and Precautions, <u>General</u> )  Concomitant use with ISENTRESS <sup>®</sup> 1200 mg once daily is not recommended
rifampin	600 mg daily	800 mg twice daily	14	1.62 <sup>‡</sup> (1.12, 2.33)	1.27 <sup>‡</sup> (0.94, 1.71)	0.47 <sup>‡</sup> (0.36, 0.61)	
<b>HIV – Antiviral: Protease Inhibitors (PI)</b>							
atazanavir	400 mg daily	100 mg single dose	10	1.53 (1.11, 2.12)	1.72 (1.47, 2.02)	1.95 (1.30, 2.92)	No dosage adjustment required for ISENTRESS <sup>®</sup> 400 mg twice daily

Coadministered Drug/Class Name	Coadministered Drug Dose/Schedule	Raltegravir Dose/Schedule	Ratio (90% Confidence Interval) of Raltegravir Pharmacokinetic Parameters with/without Coadministered Drug; No Effect = 1.00				Clinical Comment
			n	C <sub>max</sub>	AUC	C <sub>min</sub>	
atazanavir	400 mg daily	1200 mg single dose	14	1.16 (1.01, 1.33)	1.67 (1.34, 2.10)	1.26 (1.08, 1.46)	Not recommended with ISENTRESS <sup>®</sup> 1200 mg once daily
atazanavir/ritonavir	300 mg/ 100 mg daily	400 mg twice daily	10	1.24 (0.87, 1.77)	1.41 (1.12, 1.78)	1.77 (1.39, 2.25)	No dosage adjustment required for ISENTRESS <sup>®</sup> 400 mg twice daily
ritonavir	100 mg twice daily	400 mg single dose	10	0.76 (0.55, 1.04)	0.84 (0.70, 1.01)	0.99 (0.70, 1.40)	No dosage adjustment required for ISENTRESS <sup>®</sup> 400 mg twice daily and 1200 mg once daily
tipranavir/ritonavir	500 mg/ 200 mg twice daily	400 mg twice daily	15 (14 for C <sub>min</sub> )	0.82 (0.46, 1.46)	0.76 (0.49, 1.19)	0.45 (0.31, 0.66)	No dosage adjustment required for ISENTRESS <sup>®</sup> 400 mg twice daily  Co-administration of ISENTRESS <sup>®</sup> (1200 mg once daily) is not recommended.
<b>HIV – Antiviral: Non-Nucleoside Reverse Transcriptase Inhibitor (NNRTI)</b>							
efavirenz	600 mg daily	400 mg single dose	9	0.64 (0.41, 0.98)	0.64 (0.52, 0.80)	0.79 (0.49, 1.28)	No dosage adjustment required for ISENTRESS <sup>®</sup> 400 mg twice daily
efavirenz	600 mg daily	1200 mg single dose	21	0.91 (0.70, 1.17)	0.86 (0.73, 1.01)	0.94 (0.76, 1.17)	No dosage adjustment required for ISENTRESS <sup>®</sup> 1200 mg once daily

Coadministered Drug/Class Name	Coadministered Drug Dose/Schedule	Raltegravir Dose/Schedule	Ratio (90% Confidence Interval) of Raltegravir Pharmacokinetic Parameters with/without Coadministered Drug; No Effect = 1.00				Clinical Comment
			n	C <sub>max</sub>	AUC	C <sub>min</sub>	
etravirine	200 mg twice daily	400 mg twice daily	19	0.89 (0.68, 1.15)	0.90 (0.68, 1.18)	0.66 (0.34, 1.26)	No dosage adjustment required for ISENTRESS <sup>®</sup> 400 mg twice daily and 1200 mg once daily
<b>HIV – Antiviral: Nucleotide Analog Reverse Transcriptase Inhibitor (NRTI)</b>							
tenofovir disoproxil fumarate	300 mg daily	400 mg twice daily	9	1.64 (1.16, 2.32)	1.49 (1.15, 1.94)	1.03 (0.73, 1.45)	No dosage adjustment required for ISENTRESS <sup>®</sup> 400 mg twice daily
† Famotidine was administered 2 hours prior to ISENTRESS <sup>®</sup> administration ‡ Compared to 400 mg twice daily administered alone							

Rash occurred frequently in healthy subjects in a drug-interaction study when darunavir/ritonavir was coadministered with ISENTRESS<sup>®</sup>. Darunavir/ritonavir should be coadministered with ISENTRESS<sup>®</sup> only if the benefits outweigh the risks.

## DOSAGE AND ADMINISTRATION

### Dosing Considerations

ISENTRESS<sup>®</sup> is to be given in a combination regimen with other antiretroviral agents.

ISENTRESS<sup>®</sup> can be administered with or without food.

ISENTRESS HD<sup>®</sup> is available in the following dose strength:

- 600 mg film-coated tablet for once daily use

ISENTRESS<sup>®</sup> is available in the following dose strengths:

- 400 mg film-coated tablet for twice daily use
- chewable tablet in 100 mg (scored) and 25 mg strengths for twice daily use

Because the formulations have different pharmacokinetic profiles, do not substitute the 400 mg tablet for the 600 mg tablet to create a 1200 mg once daily dose and do not substitute the chewable tablet for the 400 mg or 600 mg tablet (see ACTION and CLINICAL PHARMACOLOGY, Pharmacokinetics).

During coadministration of ISENTRESS<sup>®</sup> 400 mg tablets with rifampin, the recommended dosage of ISENTRESS<sup>®</sup> is 800 mg twice daily in adults. There are no data to guide coadministration of ISENTRESS<sup>®</sup> with rifampin in patients below 18 years of age (see DRUG INTERACTIONS).

**Recommended Dose and Dosage Adjustment**

The maximum dose of the chewable tablet is 300 mg twice daily.

For the treatment of patients with HIV-1 infection the dosage of ISENTRESS<sup>®</sup> or ISENTRESS HD<sup>®</sup> is as follows:

**ISENTRESS<sup>®</sup> Twice Daily Dosing:**

**Adults and adolescents at least 12 years of age:** One 400 mg tablet twice daily, orally

**Children 2 to less than 12 years of age:** Chewable tablets: weight based to maximum dose 300 mg, twice daily, as specified in Table 9

**Table 9 – Recommended Dose for ISENTRESS<sup>®\*</sup> Chewable Tablets for Pediatric Patients 2 to less than 12 Years of Age, weighing from 7 kg**

Body Weight (kg)	Dose	Number of Chewable Tablets per dose
7 to <10	50 mg twice daily	0.5 x 100 mg <sup>†</sup> or 2 X 25 mg
10 to <14	75 mg twice daily	3 x 25 mg or 0,5 X 100 mg <sup>†</sup> + 1 X 25 mg
14 to <20	100 mg twice daily	1 x 100 mg
20 to <28	150 mg twice daily	1.5 x 100 mg <sup>†</sup> or 1 X 100 mg + 2 X 25 mg
28 to <40	200 mg twice daily	2 x 100 mg
At least 40	300 mg twice daily	3 x 100 mg

\* The weight based dosing recommendation for the chewable tablet is based on approximately 6 mg/kg/dose twice daily.  
<sup>†</sup> The 100 mg chewable tablet can be divided into equal halves

**ISENTRESS HD<sup>®</sup> Once Daily Dosing:**

**Adults and adolescents with body weight of at least 40 kg, who are either treatment-naïve or virologically suppressed on an initial regimen of ISENTRESS<sup>®</sup> 400 mg twice daily:**

Two 600 mg tablets once daily, orally

Do not substitute the 400 mg tablet for the 600 mg tablet to create a 1200 mg once daily dose.

**Hepatic/biliary/Pancreatic**

There were no clinically important pharmacokinetic differences between patients with moderate hepatic insufficiency and healthy subjects. No dosage adjustment is necessary for patients with mild to moderate hepatic insufficiency. The effect of severe hepatic insufficiency on the pharmacokinetics of raltegravir has not been studied.

## **Renal**

There were no clinically important pharmacokinetic differences between patients with severe renal insufficiency and healthy subjects. No dosage adjustment is necessary.

## **OVERDOSAGE**

For management of a suspected drug overdose, contact your regional Poison Control Centre.

Activated charcoal may be administered to aid in the removal of unabsorbed drug. The extent to which ISENTRESS<sup>®</sup> may be dialyzable is unknown.

No specific information is available on the treatment of overdose with ISENTRESS<sup>®</sup>. Multiple doses as high as 1800 mg (3 x 600 mg) q.d. for 28 days and 800 mg b.i.d. were studied in Phase I without evidence of toxicity. Occasional doses of 2400 mg per day were taken in Phase III studies without evidence of toxicity. Based upon available data, raltegravir appears to be well tolerated at doses up to 800 mg b.i.d. and when administered with drugs that increase exposure by 50–70% (such as tenofovir and atazanavir).

## **ACTION AND CLINICAL PHARMACOLOGY**

### **Mechanism of Action**

ISENTRESS<sup>®</sup> contains raltegravir, a human immunodeficiency virus integrase strand transfer inhibitor. Raltegravir inhibits the catalytic activity of HIV integrase, an HIV-encoded enzyme that is required for viral replication. Inhibition of integrase prevents the covalent insertion, or integration, of the HIV genome into the host cell genome during the early phase of infection. HIV genomes that fail to integrate cannot direct the production of new infectious viral particles, so inhibiting integration prevents propagation of the viral infection. Raltegravir did not significantly inhibit human phosphoryltransferases including DNA polymerases  $\alpha$ ,  $\beta$ , and  $\gamma$ .

### **Pharmacodynamics**

#### **Microbiology**

Raltegravir at concentrations of  $31 \pm 20$  nM resulted in 95% inhibition ( $IC_{95}$ ) of viral spread (relative to an untreated virus-infected culture) in human T-lymphoid cell cultures infected with the cell-line adapted HIV-1 variant H9IIIB. In addition, raltegravir at concentrations of 6 to 50 nM resulted in 95% inhibition of viral spread in cultures of mitogen-activated human peripheral blood mononuclear cells infected with diverse, primary clinical isolates of HIV-1, including isolates from 5 non-B subtypes, and isolates resistant to reverse transcriptase inhibitors and protease inhibitors. In a single-cycle infection assay, raltegravir inhibited infection of 23 HIV isolates representing 5 non-B subtypes and 5 circulating recombinant forms with  $IC_{50}$  values ranging from 5 to 12 nM. Raltegravir also inhibited replication of an HIV-2 isolate when tested in CEMx174 cells ( $IC_{95} = 6$  nM). Additive to synergistic antiretroviral activity was observed when human T-lymphoid cells infected with the H9IIIB variant of HIV-1 were incubated with

raltegravir in combination with nucleoside analog reverse transcriptase inhibitors (zidovudine, zalcitabine, stavudine, abacavir, tenofovir, didanosine, or lamivudine); non-nucleoside reverse transcriptase inhibitors (efavirenz, nevirapine, or delavirdine); protease inhibitors (indinavir, saquinavir, ritonavir, amprenavir, lopinavir, nelfinavir, or atazanavir); or the entry inhibitor enfuvirtide.

### **Drug Resistance**

The mutations observed in HIV-1 integrase that contributed to raltegravir resistance (evolved either *in vitro* or in patients treated with raltegravir) generally included a substitution at either Y143 (changed to C, H or R) or Q148 (changed to H, K, or R) or N155 (changed to H) plus one or more additional mutations (e.g., L74I/M, E92Q, T97A, E138A/K, G140A/S, V151I, G163R, H183P, Y226C/D/F/H, S230R, and D232N).

Recombinant viruses containing a single primary mutation (Q148H, K or R, or N155H) displayed decreased replication capacity and reduced susceptibility to raltegravir *in vitro*. Secondary mutations further decreased susceptibility to raltegravir and sometimes acted as compensatory mutations for viral replication capacity.

Mutations conferring resistance to raltegravir generally also confer resistance to the integrase strand transfer inhibitor elvitegravir. Mutations at amino acid 143 confer greater resistance to raltegravir than to elvitegravir, and the E92Q mutation confers greater resistance to elvitegravir than to raltegravir. Viruses harboring a mutation at amino acid 148, along with one or more other raltegravir resistance mutations, may also have clinically significant resistance to dolutegravir.

#### Treatment-Naïve Subjects:

By Week 96 in the STARTMRK trial, the primary raltegravir resistance-associated substitutions were observed in 4 (2 with Y143H/R and 2 with Q148H/R) of the 10 virologic failure subjects with evaluable genotypic data from paired baseline and raltegravir treatment-failure isolates.

#### Treatment-Experienced Subjects:

By Week 96 in the BENCHMRK trials, at least one of the primary raltegravir resistance-associated substitutions, Y143C/H/R, Q148H/K/R, and N155H, was observed in 76 of the 112 virologic failure subjects with evaluable genotypic data from paired baseline and raltegravir treatment-failure isolates. The emergence of the primary raltegravir resistance-associated substitutions was observed cumulatively in 70 subjects by Week 48 and 78 subjects by Week 96, 15.2% and 17% of the raltegravir recipients, respectively. Some (n = 58) of those HIV-1 isolates harboring one or more of the primary raltegravir resistance-associated substitutions were evaluated for raltegravir susceptibility yielding a median decrease of 26.3-fold (mean  $48.9 \pm 44.8$ -fold decrease, ranging from 0.8- to 159-fold) compared to the wild-type reference.

### **Cardiac Electrophysiology**

In a randomized, placebo-controlled, crossover study, 31 healthy subjects were administered a single oral supratherapeutic dose of raltegravir 1600 mg and placebo. There was no effect on the QTc interval. Peak raltegravir plasma concentrations were approximately 4-fold higher than the peak concentrations following a 400-mg dose.



## **Pharmacokinetics**

**Absorption – Adults:** As demonstrated in healthy volunteers administered single oral doses of raltegravir (400 mg film coated tablet) in the fasted state, raltegravir 400 mg twice daily is rapidly absorbed with a  $T_{max}$  of approximately 3 hours postdose in the fasted state. Raltegravir AUC and  $C_{max}$  increase dose proportionally over the dose range 100 mg to 1600 mg. Raltegravir  $C_{trough}$  increases dose proportionally over the dose range of 100 mg to 800 mg and increases slightly less than dose proportionally over the dose range 100 mg to 1600 mg. In patients on 400 mg twice daily monotherapy, raltegravir drug exposures were characterized by a geometric mean  $AUC_T$  of  $14.3\mu M \cdot hr$  and  $C_{trough}$  of 142 nM. With twice-daily dosing, pharmacokinetic steady state is achieved rapidly, within approximately the first 2 days of dosing. There is little to no accumulation in AUC and  $C_{max}$  and evidence of slight accumulation in  $C_{trough}$ . The absolute bioavailability of raltegravir has not been established.

Following administration of a single 1200 mg dose (2 x 600 mg tablet), raltegravir is also rapidly absorbed with a median  $T_{max}$  of approximately 1.5 to 2 hours in the fasted state. The bioavailability of raltegravir following once daily multiple dose administration of a 2 x 600 mg tablet is not comparable to the bioavailability of raltegravir following once daily multiple dose administration of a 3 x 400 mg tablet such that there were increases of 21% and 46% for  $AUC_T$  and  $C_{max}$  respectively while  $C_{trough}$  values were similar with the 600 mg tablets. Following administration of a 1200 mg once daily dosing of the 600 mg tablets in patients, steady state  $AUC_T$  was  $53.7 h \cdot \mu M$ ,  $C_{trough}$  was 75.6 nM, and median  $T_{max}$  was 1.50 h.

### **Effect of Food on Oral Absorption**

ISENTRESS<sup>®</sup> may be administered with or without food. Raltegravir was administered without regard to food in the pivotal safety and efficacy studies in HIV-infected patients. The effect of consumption of low-, moderate- and high-fat meals on steady-state raltegravir pharmacokinetics was assessed in healthy volunteers (Table 10). Administration of multiple doses of raltegravir 400 mg twice daily following a moderate-fat meal did not affect raltegravir  $AUC_T$  to a clinically meaningful degree with an increase of 13% relative to fasting. Raltegravir  $C_{trough}$  was 66% higher and  $C_{max}$  was 5% higher following a moderate-fat meal compared to fasting. Administration of 400 mg twice daily raltegravir following a high-fat meal increased  $AUC_T$  and  $C_{max}$  by approximately 2-fold and increased  $C_{trough}$  by 4.1-fold. Administration of 400 mg twice daily raltegravir following a low-fat meal decreased  $AUC_T$  and  $C_{max}$  by 46% and 52%, respectively;  $C_{trough}$  was essentially unchanged. Food appears to increase pharmacokinetic variability relative to fasting.

**Table 10 – Comparison of Raltegravir Plasma Pharmacokinetics in Healthy Adult Male and Female Subjects Administered Multiple Oral Doses of 400 mg Twice Daily Raltegravir in the Fasted State and After a Specified Meal**

	AUC <sub>T</sub> (μM•hr) <sup>a</sup>	C <sub>max</sub> (μM) <sup>a</sup>	C <sub>12hr</sub> (nM) <sup>a</sup>	T <sub>max</sub> (hr) <sup>b</sup>
Fasted	10.0	2.71	110	3.0
Low-Fat Meal	5.38	1.31	94	3.0
Moderate-Fat Meal	11.3	2.85	182	4.0
High-Fat Meal	21.2	5.32	453	4.0

<sup>a</sup> Geometric Mean  
<sup>b</sup> Median  
AUC<sub>T</sub> = AUC over a steady state dosing interval

Following administration of a single 1200 mg dose of the 600 mg tablets under low fat, low calorie fed conditions, there were decreases in AUC<sub>t</sub>, C<sub>max</sub> and C<sub>24hr</sub> by approximately 42%, 52% and 16%, respectively when compared to administration under fasting conditions. When a single 1200 mg dose of the 600 mg tablets was administered under high fat, high calorie fed conditions, AUC<sub>t</sub> was similar; however, there were decreases in C<sub>24hr</sub> and C<sub>max</sub> by approximately 12% and 28%, respectively when compared to administration under fasting conditions (Table 11).

**Table 11 – Comparison of Raltegravir Plasma Pharmacokinetics in Healthy Adult Male and Female Subjects Following Administration of a Single 1200 mg Oral Dose of the 600 mg Raltegravir Tablets in the Fasted State and After a Specified Meal**

	AUC <sub>t</sub> (μM.hr) <sup>a</sup>	C <sub>max</sub> (μM) <sup>a</sup>	C <sub>24hr</sub> (nM) <sup>a</sup>	T <sub>max</sub> (hr) <sup>b</sup>
Fasted	56.5	22.6	57.7	1.50
Low-Fat Meal	32.9	10.8	48.2	2.00
High-Fat Meal	57.6	16.3	50.7	3.00

<sup>a</sup> Geometric Mean  
<sup>b</sup> Median  
AUC<sub>t</sub> = AUC to the last timepoint with a detectable concentration value

**Distribution – Adults:** Raltegravir is approximately 83% bound to human plasma protein over the concentration range of 2 to 10 μM.

In two studies of HIV-1 infected subjects who received raltegravir 400 mg twice daily, raltegravir was readily detected in the cerebrospinal fluid. In the first study (n = 18), the median cerebrospinal fluid concentration was 5.8% (range 1 to 53.5%) of the corresponding plasma concentration. In the second study (n = 16), the median cerebrospinal fluid concentration was 3% (range 1 to 61%) of the corresponding plasma concentration. These median proportions are approximately 3- to 6-fold lower than the free fraction of raltegravir in plasma.

**Metabolism and Excretion – Adults:** The apparent terminal half-life of raltegravir is approximately 9 hours, with a shorter α-phase half-life (~1 hour) accounting for much of the AUC. Following administration of an oral dose of radiolabeled raltegravir, approximately 51 and 32% of the dose was excreted in feces and urine, respectively. In feces, only raltegravir was present, most of which is likely derived from hydrolysis of raltegravir-glucuronide secreted in bile as observed in preclinical species. Two components, namely raltegravir and raltegravir-glucuronide, were detected in urine and accounted for approximately 9 and 23% of the dose, respectively. The major circulating entity was raltegravir and represented approximately 70% of

the total radioactivity; the remaining radioactivity in plasma was accounted for by raltegravir-glucuronide. Studies using isoform-selective chemical inhibitors and cDNA-expressed UGT show that UGT1A1 is the main enzyme responsible for the formation of raltegravir-glucuronide. Thus, the data indicate that the major mechanism of clearance of raltegravir in humans is UGT1A1-mediated glucuronidation.

### **Special Populations and Conditions**

**Pediatrics:** Based on a formulation comparison study in healthy adult volunteers, the chewable tablet has higher oral bioavailability compared to the 400 mg tablet. In this study, administration of the chewable tablet with a high fat meal led to an average 6% decrease in AUC, 62% decrease in  $C_{max}$ , and 188% increase in  $C_{12hr}$  compared to administration in the fasted state. Administration of the chewable tablet with a high fat meal does not affect raltegravir pharmacokinetics to a clinically meaningful degree and the chewable tablet can be administered without regard to food.

The doses recommended for HIV-infected children and adolescents 2 through 18 years of age (see DOSAGE AND ADMINISTRATION, Recommended Dose and Dosage Adjustment) resulted in a pharmacokinetic profile of raltegravir similar to that observed in adults receiving 400 mg twice daily. Table 12 displays pharmacokinetic parameters in the 400 mg tablet (6 through 18 years of age) and the chewable tablet (2 to less than 12 years of age).

**Table 12 – Raltegravir Pharmacokinetic Parameters IMPAACT P1066 Following Administration of Doses in Dosage and Administration**

Age	Formulation	Dose	N*	Geometric Mean (%CV) AUC <sub>0-12hr</sub> (μM*hr)	Geometric Mean (%CV) C <sub>12hr</sub> (nM)
12 through 18 years	400 mg tablet	400 mg twice daily, regardless of weight**	11	15.7 (98%)	333 (78%)
6 to less than 12 years	400 mg tablet	400 mg twice daily, for patients ≥25 kg	11	15.8 (120%)	246 (221%)
6 to less than 12 years	Chewable tablet	Weight based dosing, see Table 9	10	22.6 (34%)	130 (88%)
2 to less than 6 years	Chewable tablet	Weight based dosing, see Table 9	12	18.0 (59%)	71 (55%)

\* Number of patients with intensive pharmacokinetic (PK) results at the final recommended dose.  
 \*\* Patients in this age group received approximately 8 mg/kg dose at time of intensive PK which met PK and safety targets. Based on review of the individual profiles and receipt of a mean dose of 390 mg, 400 mg twice daily was selected as the recommended dose for this age group.

The pharmacokinetics of raltegravir in children less than 2 years of age has not been established.

ISENTRESS<sup>®</sup> 1200 mg (2 x 600 mg) once daily was not evaluated in a pediatric clinical study, however, population PK modeling and simulation analyses were conducted. Given that all the pediatric simulated exposures are within the adult exposures observed from Phase III ONCEMRK (Protocol 292), and that there are no safety concerns at the same exposure values, a weight cutoff of 40 kg is deemed adequate to achieve a safe administration of ISENTRESS<sup>®</sup> 1200 mg (2 x 600 mg) once daily while maintaining clinical efficacy. These results support the use of ISENTRESS<sup>®</sup> 1200 mg (2 x 600 mg) once daily in pediatric patients weighing at least 40 kg.

**Geriatrics:** The effect of age (18 years and older) on the pharmacokinetics of raltegravir was evaluated in the composite analysis and the population pharmacokinetic (PK) analysis. There was no clinically meaningful effect of age on raltegravir pharmacokinetics. No dosage adjustment is necessary.

**Gender:** A study of the pharmacokinetics of raltegravir 400 mg twice daily was performed in young adult healthy males and females. Additionally, the effect of gender was evaluated in a composite analysis of pharmacokinetic data from 103 healthy subjects and 28 HIV patients receiving raltegravir monotherapy with fasted administration. The effect of gender was also evaluated in a population PK analysis of concentration data from 80 healthy subjects and HIV patients receiving raltegravir alone or in combination with other drugs and in fasted and fed conditions. There were no clinically important pharmacokinetic differences due to gender. For raltegravir 1200 mg (2 x 600 mg) once daily, based on population pharmacokinetic analysis, there were no clinically important pharmacokinetic differences due to gender. No dosage adjustment is necessary.

**Race and ethnicity:** The effect of race on the pharmacokinetics of raltegravir was evaluated in the composite analysis for ISENTRESS<sup>®</sup> 400 mg twice daily, and no clinically meaningful effect of race on raltegravir pharmacokinetics was concluded. For ISENTRESS<sup>®</sup> 1200 mg (2 x 600 mg) once daily, population PK analysis also demonstrated that the impacts of race and ethnicity are not clinically meaningful. No dosage adjustment is necessary.

**Body Mass Index (BMI) and Body Weight:** The composite analysis assessed the effect of BMI on the pharmacokinetics of raltegravir in adults. There was no clinically meaningful effect of BMI on raltegravir pharmacokinetics. Additionally, no clinically meaningful effect of body weight on raltegravir pharmacokinetics was identified in the population PK analysis for both ISENTRESS 400 mg twice daily and ISENTRESS<sup>®</sup> 1200 mg (2 x 600 mg) once daily. No dosage adjustment is necessary.

**Hepatic Insufficiency:** Raltegravir is eliminated primarily by glucuronidation in the liver. A study of the pharmacokinetics of raltegravir was performed in adult patients with moderate hepatic insufficiency. Additionally, hepatic insufficiency was evaluated in the composite pharmacokinetic analysis. There were no clinically important pharmacokinetic differences between patients with moderate hepatic insufficiency and healthy subjects. No dosage adjustment is necessary for patients with mild to moderate hepatic insufficiency. No hepatic impairment study has been conducted with ISENTRESS<sup>®</sup> 1200 mg (2 x 600 mg) once daily; however, based on results with ISENTRESS<sup>®</sup> 400 mg twice daily tablet, no clinically meaningful effect is expected for mild and moderate hepatic impairment. The effect of severe hepatic insufficiency on the pharmacokinetics of raltegravir has not been studied.

**Renal Insufficiency:** Renal clearance of unchanged drug is a minor pathway of elimination. A study of the pharmacokinetics of raltegravir was performed in adult patients with severe renal insufficiency. Additionally, renal insufficiency was evaluated in the composite pharmacokinetic analysis. There were no clinically important pharmacokinetic differences between patients with severe renal insufficiency and healthy subjects. No dosage adjustment is necessary. No renal impairment study was conducted with ISENTRESS<sup>®</sup> 1200 mg (2 x 600 mg) once daily; however, based on results with ISENTRESS<sup>®</sup> 400 mg twice daily tablet, no clinically meaningful effect is

anticipated. Because the extent to which ISENTRESS<sup>®</sup> may be dialyzable is unknown, dosing before a dialysis session should be avoided.

**Genetic Polymorphism:** There is no evidence that common UGT1A1 polymorphisms alter raltegravir pharmacokinetics to a clinically meaningful extent. In a comparison of 30 adult subjects with \*28/\*28 genotype (associated with reduced activity of UGT1A1) to 27 adult subjects with wild-type genotype, the geometric mean ratio (90% CI) of AUC was 1.41 (0.96, 2.09).

## STORAGE AND STABILITY

### 400 mg Tablets

Store at room temperature (15°C–30°C).

Keep the desiccant in the bottle.

### 600 mg Tablets, Chewable Tablets 25 mg and 100 mg

Store at room temperature (15°C–30°C) in the original package with the bottle tightly closed.

Keep the desiccant in the bottle to protect from moisture.

## DOSAGE FORMS, COMPOSITION AND PACKAGING

### Dosage Forms

ISENTRESS HD<sup>®</sup> film-coated 600-mg tablets are yellow, oval-shaped tablets debossed with Merck logo and “242” on one side. The tablets are packaged in white, HDPE bottles and enclosed with a child resistant closure. Each bottle contains 60 tablets and silica gel desiccant.

ISENTRESS<sup>®</sup> film-coated 400-mg tablets are pink, oval-shaped, with 227 on one side. They are supplied as unit-of-use bottles of 60.

ISENTRESS<sup>®</sup> chewable tablets 100-mg are pale orange, oval shaped scored tablet, debossed with the Merck logo on one side of the score and 477 on the other, and scored on the other side of the tablet. They are supplied in unit-of-use bottles of 60.

ISENTRESS<sup>®</sup> chewable tablets 25-mg are pale yellow, round, flat faced, beveled edge tablet debossed with the Merck logo on one side and 473 on the other side of the tablet. They are supplied in unit-of-use bottles of 60.

### Composition

ISENTRESS HD<sup>®</sup> 600 mg film-coated tablets contains 651.6 mg of raltegravir (as potassium salt), equivalent to 600 mg of raltegravir free phenol.

ISENTRESS<sup>®</sup> 400 mg film-coated tablets contains 434.4 mg of raltegravir (as potassium salt), equivalent to 400 mg of raltegravir free phenol.

ISENTRESS<sup>®</sup> 100 mg chewable tablets contains 108.6 mg of raltegravir (as potassium salt), equivalent to 100 mg of raltegravir free phenol.

ISENTRESS<sup>®</sup> 25 mg chewable tablets contains 27.16 mg of raltegravir (as potassium salt), equivalent to 25 mg of raltegravir free phenol.

Non-medicinal ingredients:

ISENTRESS HD<sup>®</sup> 600 mg film-coated tablets:

croscarmellose sodium, hypromellose 2910, magnesium stearate, microcrystalline cellulose. In addition, the film coating contains the following inactive ingredients: black iron oxide, carnauba wax, hypromellose 2910, iron oxide yellow, lactose monohydrate, titanium dioxide, and triacetin.

ISENTRESS<sup>®</sup> 400 mg film-coated tablets:

calcium phosphate dibasic anhydrous, hypromellose 2208, lactose monohydrate, magnesium stearate, microcrystalline cellulose, poloxamer 407 (contains 0.01% butylated hydroxytoluene as antioxidant), sodium stearyl fumarate. In addition, the film coating contains the following inactive ingredients: black iron oxide and red iron oxide, polyethylene glycol 3350, polyvinyl alcohol, talc and titanium dioxide.

ISENTRESS<sup>®</sup> 25 mg and 100 mg chewable tablets:

ammonium hydroxide, crospovidone, ethylcellulose 20 cP, fructose, hydroxypropyl cellulose, hypromellose 2910/6cP, macrogol/PEG 400, magnesium stearate, mannitol, medium chain triglycerides, monoammonium glycyrrhizinate, natural and artificial flavors (orange, banana, and masking), oleic acid, phenylalanine (as part of the artificial sweetener, aspartame), saccharin sodium, sodium citrate dihydrate, sodium stearyl fumarate, sorbitol, sucralose and yellow iron oxide.

ISENTRESS<sup>®</sup> 100 mg also contain red iron oxide.



## CLINICAL TRIALS

### Description of Clinical Trials

#### Adults

This indication is based on the evidence of durable efficacy of ISENTRESS<sup>®</sup> 400 mg twice daily from the original analysis of 48 week data from 3, randomized, double-blind, controlled trials.

The efficacy of ISENTRESS<sup>®</sup> 400 mg twice daily is supported by data from two randomized, double-blind, controlled trials in antiretroviral treatment-experienced subjects, BENCHMRK 1 (48 and 96 weeks), and BENCHMRK 2 (48 and 96 weeks); and by data from one randomized, double-blind, controlled trial in antiretroviral treatment-naive subjects, STARTMRK (48, 96, 156 and 240 weeks).

The efficacy of ISENTRESS<sup>®</sup> 1200 mg once daily is supported by data from one randomized, double-blind, controlled trial in antiretroviral treatment-naive subjects, ONCEMRK (48 and 96 weeks).

#### Treatment-Experienced Adult Patients

BENCHMRK- 1 and BENCHMRK- 2 are Phase III studies to evaluate the safety and antiretroviral activity of ISENTRESS<sup>®</sup> 400 mg b.i.d. in combination with an optimized background therapy (OBT), versus OBT alone, in HIV-infected patients, 16 years or older, with documented resistance to at least 1 drug in each of 3 classes (NRTIs, NNRTIs, PIs) of antiretroviral therapies. Randomization was stratified by degree of resistance to PI (1PI vs. >1PI) and the use of enfuvirtide in the OBT. Prior to randomization, OBT was selected by the investigator based on genotypic/phenotypic resistance testing and prior ART history.



Table 14 shows the demographic characteristics between patients in the group receiving ISENTRESS<sup>®</sup> 400 mg b.i.d. and patients in the group receiving placebo.

**Table 14 – Baseline Population Characteristics**

<b>BENCHMRK-1 and -2 Pooled</b>	<b>ISENTRESS<sup>®</sup> 400 mg b.i.d. + OBT (N = 462)</b>	<b>Placebo + OBT (N = 237)</b>
<b>Gender n (%)</b>		
Male	405 (87.7)	210 (88.6)
Female	57 (12.3)	27 (11.4)
<b>Race n (%)</b>		
White	301 (65.2)	173 (73.0)
Black	65 (14.1)	26 (11.0)
Asian	16 (3.5)	6 (2.5)
Hispanic	53 (11.5)	19 (8.0)
Others	27 (5.8)	13 (5.5)
<b>Age (years)</b>		
Median (min, max)	45.0 (16 to 74)	45.0 (17 to 70)
<b>CD4 Cell Count</b>		
Median (min, max), cells/mm <sup>3</sup>	119 (1 to 792)	123 (0 to 759)
≤50 cells/mm <sup>3</sup> , n (%)	146 (31.6)	78 (32.9)
>50 and ≤200 cells/mm <sup>3</sup> , n (%)	173 (37.4)	85 (35.9)
<b>Plasma HIV RNA</b>		
Median (min, max), log <sub>10</sub> copies/mL	4.8 (2.3 to 5.9)	4.7 (2.3 to 5.9)
≥100,000 copies/mL, n (%)	165 (35.7)	78 (32.9)
<b>History of AIDS n (%)</b>		
Yes	427 (92.4)	215 (90.7)
<b>Prior Use of ART, Median (1<sup>st</sup> Quartile, 3<sup>rd</sup> Quartile)</b>		
Years of ART Use	10.1 (7.3 to 12.1)	10.2 (7.9 to 12.4)
Number of ART	12.0 (9 to 15)	12.0 (9 to 14)
<b>Hepatitis Co-infection* n (%)</b>		
No Hepatitis B or C	385 (83.3)	200 (84.4)
Hepatitis B only	36 (7.8)	7 (3.0)
Hepatitis C only	37 (8.0)	28 (11.8)
Co-infection of Hepatitis B and C	4 (0.9)	2 (0.8)
<b>Stratum n (%)</b>		
Enfuvirtide in OBT	175 (37.9)	89 (37.6)
Resistant to ≥2 PI	447 (96.8)	226 (95.4)
* Hepatitis B surface antigen positive or hepatitis C antibody positive.		

Table 15 compares the characteristics of optimized background therapy at baseline in the group receiving ISENTRESS<sup>®</sup> 400 mg b.i.d. and patients in the control group.

**Table 15 – Characteristics of Optimized Background Therapy at Baseline**

<b>BENCHMRK-1 and -2 Pooled</b>	<b>ISENTRESS<sup>®</sup> 400 mg b.i.d. + OBT (N = 462)</b>	<b>Placebo + OBT (N = 237)</b>
<b>Number of ARTs in OBT</b>		
Median (min, max)	4 (1 to 7)	4 (2 to 7)
<b>Number of Active PI in OBT by Phenotypic Resistance Test*</b>		
0	165 (35.7)	96 (40.5)
1 or more	278 (60.2)	137 (57.8)
<b>Phenotypic Sensitivity Score (PSS)<sup>†</sup></b>		
0	67 (14.5)	43 (18.1)
1	144 (31.2)	71 (30.0)
2	142 (30.7)	66 (27.8)
3 or more	85 (18.4)	48 (20.3)
<b>Genotypic Sensitivity Score (GSS)<sup>†</sup></b>		
0	116 (25.1)	65 (27.4)
1	177 (38.3)	95 (40.1)
2	111 (24.0)	49 (20.7)
3 or more	51 (11.0)	23 (9.7)
* Darunavir use in OBT in darunavir naïve patients was counted as one active PI.		
<sup>†</sup> The Phenotypic Sensitivity Score (PSS) and the Genotypic Sensitivity Score (GSS) were defined as the total oral ARTs in OBT to which a patient's viral isolate showed phenotypic sensitivity and genotypic sensitivity, respectively, based upon phenotypic and genotypic resistance tests. Enfuvirtide use in OBT in enfuvirtide-naïve patients was counted as one active drug in OBT in the GSS and PSS. Similarly, darunavir use in OBT in darunavir-naïve patients was counted as one active drug in OBT.		

Week 96 outcomes for the 699 patients randomized and treated with the recommended dose of ISENTRESS<sup>®</sup> 400 mg b.i.d. or comparator in the pooled BENCHMRK 1 and 2 studies are shown in Table 16.

**Table 16 – Virologic Outcomes of Randomized Treatment of Protocols 018 and 019 at 96 Weeks (Pooled Analysis)**

	<b>ISENTRESS<sup>®</sup> 400 mg Twice Daily + OBT (N = 462)</b>	<b>Placebo + OBT (N = 237)</b>
<b>Subjects with HIV-1 RNA less than 50 copies/mL</b>	55%	27%
<b>Virologic Failure*</b>	35%	66%
<b>No virologic data at Week 96 Window</b>		
<b>Reasons</b>		
<b>Discontinued study due to AE or death<sup>†</sup></b>	3%	3%
<b>Discontinued study for other reasons<sup>‡</sup></b>	4%	4%
<b>Missing data during window but on study</b>	4%	<1%
<p>* Includes subjects who switched to open-label raltegravir after Week 16 due to the protocol-defined virologic failure, subjects who discontinued prior to Week 96 for lack of efficacy, subjects changed OBT due to lack of efficacy prior to Week 96, or subjects who were <math>\geq 50</math> copies in the 96 week window.</p> <p><sup>†</sup> Includes subjects who discontinued due to AE or Death at any time point from Day 1 through the Week 96 window if this resulted in no virologic data on treatment during the Week 96 window.</p> <p><sup>‡</sup> Other includes: withdrew consent, loss to follow-up, moved etc., if the viral load at the time of discontinuation was &lt;50 copies/mL.</p>		

The mean changes in plasma HIV-1 RNA from baseline were -1.81 log<sub>10</sub> copies/mL in the group receiving ISENTRESS<sup>®</sup> 400 mg b.i.d. and -0.75 log<sub>10</sub> copies/mL for the control group. The mean increase from baseline in CD4+ cell counts was higher in the group receiving ISENTRESS<sup>®</sup> 400 mg b.i.d. (118 cells/mm<sup>3</sup>) than in the control group (47 cells/mm<sup>3</sup>).

**Table 17 – Proportion of Patients With HIV RNA <50 Copies/mL Over Time – Protocols 018 and 019 Combined (Non-Completer = Failure Approach<sup>†</sup>)**

Endpoint	Visit	Response				Difference in Percent Response [Group A Minus Group B] <sup>‡</sup> (95% CI)
		ISENTRESS <sup>®</sup> 400 mg b.i.d. (Group A)		Placebo (Group B)		
		n/N	% (95% CI)	n/N	% (95% CI)	
Proportions with HIV RNA <50 copies/mL	Week 2	102/462	22.1 (18.4, 26.1)	24/237	10.1 (6.6, 14.7)	12.0 (6.3, 17.2)
	Week 4	195/459	42.5 (37.9, 47.2)	43/237	18.1 (13.5, 23.7)	24.3 (17.4, 30.8)
	Week 8	247/458	53.9 (49.2, 58.6)	66/236	28.0 (22.3, 34.2)	26.0 (18.4, 33.1)
	Week 12	275/460	59.8 (55.1, 64.3)	74/237	31.2 (25.4, 37.5)	28.6 (21.0, 35.7)
	Week 16	283/457	61.9 (57.3, 66.4)	82/236	34.7 (28.7, 41.2)	27.2 (19.5, 34.5)
	Week 24	289/461	62.7 (58.1, 67.1)	80/237	33.8 (27.8, 40.2)	28.9 (21.3, 36.2)
	Week 32	282/453	62.3 (57.6, 66.7)	78/237	32.9 (27.0, 39.3)	29.3 (21.7, 36.6)
	Week 40	290/458	63.3 (58.7, 67.7)	78/237	32.9 (27.0, 39.3)	30.4 (22.8, 37.6)
	Week 48	285/459	62.1 (57.5, 66.5)	78/237	32.9 (27.0, 39.3)	29.2 (21.5, 36.4)
	Week 60	281/456	61.6 (57.0, 66.1)	72/236	30.5 (24.7, 36.8)	31.1 (23.5, 38.2)
	Week 72	269/460	58.5 (53.8, 63.0)	70/237	29.5 (23.8, 35.8)	28.9 (21.4, 36.1)
	Week 84	265/460	57.6 (52.9, 62.2)	66/237	27.8 (22.2, 34.0)	29.8 (22.3, 36.8)
	Week 96	262/460	57.0 (52.3, 61.5)	62/237	26.2 (20.7, 32.2)	30.8 (23.4, 37.7)

<sup>†</sup> Approach to handling missing values: Non-Completer = Failure (NC = F) Approach.  
<sup>‡</sup> A positive value means ISENTRESS<sup>®</sup> is better than Placebo.  
 Note: ISENTRESS<sup>®</sup> and Placebo were administered with Optimized Background Therapy (OBT).  
 N = Number of patients in each treatment group.  
 n = Number of patients in each subcategory.

Virologic responses at Week 96 by baseline genotypic and phenotypic sensitivity score are shown in Table 18.

**Table 18 – Virologic Response at 96 Week Window by Baseline Genotypic/Phenotypic Sensitivity Score**

	Percent with HIV-1 RNA <50 copies/mL At Week 96			
	n	ISENTRESS® 400 mg Twice Daily + OBT (N = 462)	n	Placebo + OBT (N = 237)
<b>Phenotypic Sensitivity Score (PSS)*</b>				
0	67	43	43	5
1	144	58	71	23
2	142	61	66	32
3 or more	85	48	48	42
<b>Genotypic Sensitivity Score (GSS)*</b>				
0	116	39	65	5
1	177	62	95	26
2	111	61	49	53
3 or more	51	49	23	35
* The Phenotypic Sensitivity Score (PSS) and the Genotypic Sensitivity Score (GSS) were defined as the total oral ARTs in OBT to which a patient's viral isolate showed phenotypic sensitivity and genotypic sensitivity, respectively, based upon phenotypic and genotypic resistance tests. Enfuvirtide use in OBT in enfuvirtide-naïve patients was counted as one active drug in OBT in the GSS and PSS. Similarly, darunavir use in OBT in darunavir-naïve patients was counted as one active drug in OBT.				

### Switch of Suppressed Patients from Lopinavir (+) Ritonavir to Raltegravir

The SWITCHMRK 1 & 2 studies evaluated HIV-infected patients receiving suppressive (screening HIV RNA <50 copies/ml; stable regimen >3 months) therapy with lopinavir 200 mg (+) ritonavir 50 mg 2 tablets twice daily plus at least 2 nucleoside reverse transcriptase inhibitors and randomized them 1:1 to continue lopinavir (+) ritonavir 2 tablets twice daily (n = 174 and n = 178, respectively) or replace lopinavir (+) ritonavir with raltegravir 400 mg twice daily (n = 174 and n = 176, respectively). Patients with a prior history of virological failure were not excluded and the number of previous antiretroviral therapies was not limited.

These studies were terminated after the primary efficacy analysis at Week 24 because they failed to demonstrate non-inferiority of raltegravir versus lopinavir (+) ritonavir. In both studies at Week 24, suppression of HIV RNA to less than 50 copies/ml was maintained in 84.4% of the raltegravir group versus 90.6% of the lopinavir (+) ritonavir group, (Non-completer = Failure). In patients who had never experienced virological failure before study entry, similar virologic response rates were seen in the raltegravir and the lopinavir (+) ritonavir groups.

## Treatment-Naïve Adult Patients

### STARTMRK (Protocol 021; ISENTRESS® 400 mg twice daily)

STARTMRK is a Phase III study to evaluate the safety and antiretroviral activity of ISENTRESS® 400 mg twice daily + emtricitabine (+) tenofovir disoproxil fumarate versus efavirenz 600 mg at bedtime plus emtricitabine (+) tenofovir disoproxil fumarate in treatment-naïve HIV-infected patients with HIV RNA >5000 copies/mL. Randomization was stratified by screening HIV RNA level ( $\leq 50,000$  copies/mL; and  $> 50,000$  copies/mL) and by hepatitis status.

Table 19 shows the demographic characteristics between patients in the group receiving ISENTRESS® 400 mg twice daily and patients in the group receiving efavirenz.

**Table 19 – Patient Baseline Characteristics**

	<b>ISENTRESS® 400 mg Twice Daily (N = 281)</b>	<b>Efavirenz 600 mg At Bedtime (N = 282)</b>	<b>Total (N = 563)</b>
<b>Gender n (%)</b>			
Male	227 (80.8)	231 (81.9)	458 (81.3)
Female	54 (19.2)	51 (18.1)	105 (18.7)
<b>Race n (%)</b>			
White	116 (41.3)	123 (43.6)	239 (42.5)
Black	33 (11.7)	23 (8.2)	56 (9.9)
Asian	36 (12.8)	32 (11.3)	68 (12.1)
Hispanic	60 (21.4)	67 (23.8)	127 (22.6)
Native American	1 (0.4)	1 (0.4)	2 (0.4)
Multiracial	35 (12.5)	36 (12.8)	71 (12.6)
<b>Region n (%)</b>			
Latin America	99 (35.2)	97 (34.4)	196 (34.8)
Southeast Asia	34 (12.1)	29 (10.3)	63 (11.2)
North America	82 (29.2)	90 (31.9)	172 (30.6)
EU/Australia	66 (23.5)	66 (23.4)	132 (23.4)
<b>Age (years)</b>			
18–64 n (%)	279 (99.3)	278 (98.6)	557 (98.9)
$\geq 65$ n (%)	2 (0.7)	4 (1.4)	6 (1.1)
Mean (SD)	37.6 (9.0)	36.9 (10.0)	37.2 (9.5)
Median (min, max)	37.0 (19 to 67)	36.0 (19 to 71)	37.0 (19 to 71)
<b>CD4 Cell Count (cells/microL)</b>			
N <sup>†</sup>	281	281	562
Mean (SD)	218.9 (124.2)	217.4 (133.6)	218.1 (128.8)
Median (min, max)	212.0 (1 to 620)	204.0 (4 to 807)	207.5 (1 to 807)

<b>Plasma HIV RNA (log<sub>10</sub> copies/mL)</b>			
N <sup>†</sup>	281	282	563
Mean (SD)	5.0 (0.6)	5.0 (0.6)	5.0 (0.6)
Median (min, max)	5.1 (2.6 to 5.9)	5.0 (3.6 to 5.9)	5.0 (2.6 to 5.9)
<b>Plasma HIV RNA (copies/mL)</b>			
N <sup>†</sup>	281	282	563
Geometric Mean	103,205	10,6215	104,702
Median (min, max)	114,000 (400 to 750,000)	104,000 (4,410 to 750,000)	110,000 (400 to 750,000)
<b>History of AIDS n (%)</b>			
Yes	52 (18.5)	59 (20.9)	111 (19.7)
<b>Stratum n (%)</b>			
Screening HIV RNA ≤ 50,000	75 (26.7)	80 (28.4)	155 (27.5)
Hepatitis B or C Positive <sup>‡</sup>	18 (6.4)	16 (5.7)	34 (6.0)
<b>Viral Subtype n (%)</b>			
Clade B	219 (77.9)	230 (81.6)	449 (79.8)
Non-Clade B <sup>§</sup>	59 (21.0)	47 (16.7)	106 (18.8)
Missing	3 (1.1)	5 (1.8)	8 (1.4)
<b>Baseline Plasma HIV RNA<sup>†</sup> n (%)</b>			
≤ 50,000 copies/mL	79 (28.1)	84 (29.8)	163 (29.0)
>50,000 copies/mL	202 (71.9)	198 (70.2)	400 (71.0)
≤ 100,000 copies/mL	127 (45.2)	139 (49.3)	266 (47.2)
>100,000 copies/mL	154 (54.8)	143 (50.7)	297 (52.8)
<b>Baseline CD4 Cell Counts n (%)</b>			
≤ 50 cells/mm <sup>3</sup>	27 (9.6)	31 (11.0)	58 (10.3)
>50 cells/mm <sup>3</sup> and ≤200 cells/mm <sup>3</sup>	104 (37.0)	105 (37.2)	209 (37.1)
>200 cells/mm <sup>3</sup>	150 (53.4)	145 (51.4)	295 (52.4)
missing	0 (0.0)	1 (0.4)	1 (0.2)
<sup>†</sup> Patients with missing results excluded. <sup>‡</sup> Evidence of hepatitis B surface antigen or evidence of HCV RNA by polymerase chain reaction (PCR) quantitative test for hepatitis C Virus. <sup>§</sup> Non-Clade B Subtypes (# of patients): Clade A (4), A/C (1), A/G (2), A1(1), AE (29), AG (12), BF (6), C (37), D (2), F (2), F1 (5), G (2), Complex (3)			
Notes: ISENTRRESS <sup>®</sup> and efavirenz were administered with emtricitabine (+) tenofovir disoproxil fumarate N = Number of patients in each group. n (%) = Number (percent) of patients in each sub-category.			

Patients receiving ISENTRESS<sup>®</sup> achieved viral suppression (HIV RNA < 50 copies/mL) earlier than those receiving efavirenz, both in combination with emtricitabine (+) tenofovir disoproxil fumarate.

In the STARTMRK trial of combination antiretroviral therapy in treatment-naïve patients, ISENTRESS<sup>®</sup> with emtricitabine (+) tenofovir disoproxil fumarate demonstrated through

156 weeks consistent virologic and immunologic efficacy non-inferior to that of efavirenz with emtricitabine (+) tenofovir disoproxil fumarate with respect to the number of subjects with HIV-1 RNA <50 copies/mL [76% vs. 68%; 7.4% (-0.1% , 14.7%)]. In addition, ISENTRESS<sup>®</sup> with emtricitabine (+) tenofovir disoproxil fumarate demonstrated comparable efficacy relative to efavirenz with emtricitabine, tenofovir disoproxil fumarate across demographic and baseline prognostic factors, including: baseline plasma HIV RNA level >100,000 copies/mL, baseline CD4 cells ≤50 cells/mm<sup>3</sup>, demographic groups (including age, gender, region, and race), viral subtypes (comparing non-clade B as a group to clade B), and viral hepatitis co-infection status (hepatitis B and/or C).

**Table 20 – Virologic Outcomes of Randomized Treatment of Protocol 021 at 240 Weeks**

	<b>ISENTRESS<sup>®</sup> 400 mg Twice Daily (N = 281)</b>	<b>Efavirenz 600mg At bedtime (N = 282)</b>	<b>Difference (ISENTRESS<sup>®</sup> – Efavirenz) (CI)</b>
<b>Subjects with HIV-1 RNA less than 50 copies/mL</b>	66%	60%	6.64% (-1.4%, 14.5%)
<b>Virologic Failure*</b>	8%	15%	
<b>No virologic data at Week 240 Window</b>			
<b>Reasons</b>			
<b>Discontinued study due to AE or death<sup>†</sup></b>	5%	10%	
<b>Discontinued study for other reasons<sup>‡</sup></b>	15%	14%	
<b>Missing data during window but on study</b>	6%	2%	
* Includes subjects who discontinued prior to Week 240 for lack of efficacy or subjects who are ≥50 copies/mL in the 240-week window (+/-6 weeks).			
† Includes subjects who discontinued due to AE or Death at any time point from Day 1 through the Week 240 window if this resulted in no virologic data on treatment during Week 240 visit window.			
‡ Other includes: withdrew consent, loss to follow-up, moved etc., if the viral load at the time of discontinuation was <50 copies/mL.			

The mean changes in CD4 count from baseline were 295 cells/mm<sup>3</sup> in the group receiving ISENTRESS<sup>®</sup> 400 mg twice daily and 236 cells/mm<sup>3</sup> in the group receiving efavirenz 600 mg at bedtime.

In the STARTMRK trial of combination antiretroviral therapy in treatment-naïve patients, ISENTRESS<sup>®</sup> with emtricitabine (+) tenofovir disoproxil fumarate demonstrated through 240 weeks consistent virologic and immunologic efficacy relative to efavirenz with emtricitabine (+) tenofovir disoproxil fumarate across demographic and baseline prognostic factors, including: baseline plasma HIV RNA level >100,000 copies/mL, baseline CD4 cells ≤50 cells/mm<sup>3</sup>, demographic groups (including age, gender, region, and race), viral subtypes (comparing non-clade B as a group to clade B), and viral hepatitis co-infection status (hepatitis B and/or C).



## Long-Term Treatment-Naïve Results

**Table 21 – Proportion of Patients With HIV RNA <50 Copies/mL Over Time – Protocol 021 (Non-Completer = Failure Approach<sup>†</sup>)**

Endpoint	Visit	Response				Difference in Percent Response [Group A Minus Group B] <sup>‡</sup>
		ISENTRESS <sup>®</sup> 400 mg b.i.d. (Group A)		Efavirenz 600 mg q.h.s. (Group B)		
		n/N	% (95% CI)	n/N	% (95% CI)	(95% CI)
Proportions with HIV RNA <50 copies/mL	Week 2	62/281	22.1 (17.4, 27.4)	6/282	2.1 (0.8, 4.6)	20.3 (15.1, 26.0)
	Week 4	144/279	51.6 (45.6, 57.6)	33/282	11.7 (8.2, 16.0)	40.6 (34.0, 47.1)
	Week 8	209/281	74.4 (68.9, 79.4)	107/282	37.9 (32.3, 43.9)	37.0 (29.6, 44.1)
	Week 12	227/278	81.7 (76.6, 86.0)	169/282	59.9 (54.0, 65.7)	22.1 (15.1, 29.2)
	Week 16	242/281	86.1 (81.5, 89.9)	219/281	77.9 (72.6, 82.6)	8.4 (2.2, 14.8)
	Week 24	244/279	87.5 (83.0, 91.1)	239/282	84.8 (80.0, 88.7)	2.7 (-3.1, 8.5)
	Week 32	241/278	86.7 (82.1, 90.5)	239/280	85.4 (80.7, 89.3)	1.3 (-4.5, 7.2)
	Week 40	239/280	85.4 (80.7, 89.3)	234/281	83.3 (78.4, 87.4)	2.1 (-4.0, 8.2)
	Week 48	241/280	86.1 (81.5, 89.9)	230/281	81.9 (76.8, 86.2)	4.2 (-1.9, 10.3)
	Week 60	231/281	82.2 (77.2, 86.5)	225/282	79.8 (74.6, 84.3)	2.4 (-4.1, 8.9)
	Week 72	241/281	85.8 (81.1, 89.6)	231/282	81.9 (76.9, 86.2)	3.8 (-2.3, 10.0)
	Week 84	234/280	83.6 (78.7, 87.7)	223/281	79.4 (74.2, 83.9)	4.2 (-2.3, 10.7)
	Week 96	228/281	81.1 (76.1, 85.5)	222/282	78.7 (73.5, 83.4)	2.4 (-4.3, 9.0)
	Week 108	228/281	81.1 (76.1, 85.5)	211/279	75.6 (70.2, 80.5)	5.4 (-1.4, 12.2)
	Week 120	220/277	79.4 (74.2, 84.0)	213/281	75.8 (70.4, 80.7)	3.5 (-3.4, 10.4)
	Week 132	214/279	76.7 (71.3, 81.5)	207/281	73.7 (68.1, 78.7)	2.9 (-4.2, 10.1)
	Week 144	217/280	77.5 (72.2, 82.3)	197/281	70.1 (64.4, 75.4)	7.3 (0.0, 14.5)
	Week 156	212/281	75.4 (70.0, 80.4)	194/282	68.8 (63.0, 74.2)	6.6 (-0.8, 14.0)
	Week 168	208/281	74.0 (68.5, 79.0)	192/282	68.1 (62.3, 73.5)	5.9 (-1.7, 13.3)
	Week 180	206/280	73.6 (68.0, 78.6)	192/280	68.6 (62.8, 74.0)	4.9 (-2.6, 12.4)
Week 192	214/281	76.2 (70.7, 81.0)	189/282	67.0 (61.2, 72.5)	9.0 (1.6, 16.4)	
Week 204	205/280	73.2 (67.6, 78.3)	187/281	66.5 (60.7, 72.0)	6.5 (-1.0, 14.1)	
Week 216	205/277	74.0 (68.4, 79.1)	186/282	66.0 (60.1, 71.5)	7.9 (0.3, 15.4)	
Week 228	197/280	70.4 (64.6, 75.6)	178/281	63.3 (57.4, 69.0)	6.8 (-1.0, 14.5)	
Week 240	198/279	71.0 (65.3, 76.2)	171/279	61.3 (55.3, 67.0)	9.5 (1.7, 17.3)	

<sup>†</sup> Approach to handling missing values: Non-Completer = Failure (NC = F) Approach.

<sup>‡</sup> A positive value means ISENTRESS<sup>®</sup> is better than Efavirenz. The 95% CIs were calculated using Miettinen and Nurminen's method with weights proportional to the size of each stratum (screen HIV RNA >50,000 copies/mL or ≤50,000 copies/mL).

Note: ISENTRESS<sup>®</sup> and Efavirenz were administered with emtricitabine (+) tenofovir disoproxil fumarate.

N = Number of patients in each treatment group.

n = Number of patients in each subcategory.

**ONCEMRK (Protocol 292; ISENTRESS® 1200 mg [2 x 600 mg] once daily)**

ONCEMRK is a Phase III study to evaluate the safety and antiretroviral activity of ISENTRESS® 1200 mg (2 x 600 mg) once daily versus ISENTRESS® 400 mg twice daily, both in combination with emtricitabine (+) tenofovir disoproxil fumarate, in treatment-naïve HIV-infected patients with HIV RNA  $\geq$  1000 copies/mL. Randomization was stratified by screening HIV RNA level ( $\leq$ 100,000 copies/mL; and  $>$ 100,000 copies/mL) and by hepatitis status.

Table 22 shows the demographic characteristics for both treatment groups.

**Table 22 - Subject Baseline Characteristics by Treatment Group**

	<b>Raltegravir 1200 mg Once daily (N = 531) n (%)</b>	<b>Raltegravir 400 mg Twice daily (N = 266) n (%)</b>	<b>Total (N = 797) n (%)</b>
<b>Gender n (%)</b>			
Male	440 (82.9)	234 (88.0)	674 (84.6)
Female	91(17.1)	32 (12.0)	123 (15.4)
<b>Race n (%)</b>			
American Indian or Alaska Native	3 (0.6)	3 (1.1)	6 (0.8)
Asian	83 (15.6)	40 (15.0)	123 (15.4)
Black or African American	98 (18.5)	36 (13.5)	134 (16.8)
Multiple	46 (8.7)	14 (5.3)	60 (7.5)
Native Hawaiian or Other Pacific Islander	0 (0.0)	1 (0.4)	1 (0.1)
White	301 ( 56.7)	172 (64.7)	473 ( 59.3)
<b>Ethnicity n (%)</b>			
Hispanic or Latino	126 (23.7)	52 (19.5)	178 (22.3)
Not Hispanic or Latino	380 (71.6)	205 (77.1)	585 (73.4)
Not Reported	19 (3.6)	8 (3.0)	27 (3.4)
Unknown	6 (1.1)	1 (0.4)	7 (0.9)
<b>Region n (%)</b>			
Africa	43 (8.1)	13 (4.9)	56 (7.0)
Asia/Pacific	86 (16.2)	46 (17.3)	132 (16.6)
Europe	200 (37.7)	112 (42.1)	312 (39.1)
Latin America	77 (14.5)	26 (9.8)	103 (12.9)
North America	125 (23.5)	69 (25.9)	194 (24.3)

<b>Age (years)</b>			
18 to 64	527 (99.2)	263 (98.9)	790 (99.1)
>=65	4 (0.8)	3 (1.1)	7 (0.9)
Mean (SD)	35.4 (10.3)	36.9 (11.0)	35.9 (10.5)
Median (min, max)	34.0 (18,66)	35.0 (19,84)	34.0 (18,84)
<b>Baseline CD4 Cell Count (cells/mm<sup>3</sup>)</b>			
N <sup>†</sup>	531	266	797
Mean (SD)	407.6 (213.7)	428.9 (217.3)	414.7 (215.0)
Median (min, max)	380.0 (19,1836)	415.5 (19,1130)	390.0 (19,1836)
<b>Baseline CD4 Cell Counts n (%)</b>			
<=50 cells/mm <sup>3</sup>	9 (1.7)	6 (2.3)	15 (1.9)
>50 cells/mm <sup>3</sup> and <=200 cells/mm <sup>3</sup>	60 (11.3)	31 (11.7)	91 (11.4)
>200 cells/mm <sup>3</sup>	462 (87.0)	229 (86.1)	691 (86.7)
<b>Baseline Plasma HIV RNA (log<sub>10</sub> copies/mL)</b>			
N <sup>†</sup>	531	266	797
Mean (SD)	4.6 (0.7)	4.6 (0.7)	4.6 (0.7)
Median (min, max)	4.6 (1.6, 6.6)	4.6 (2.7, 6.2)	4.6 (1.6, 6.6)
<b>Baseline Plasma HIV RNA (copies/mL)</b>			
N <sup>†</sup>	531	266	797
Geometric Mean	40518.8	40733.2	40590.2
Median (min, max)	43890.0 (39, 3910386)	40631.0 (454, 1466713)	42424.0 (39,3910386)
<b>Baseline Plasma HIV RNA n (%)</b>			
<=100,000 copies/mL	382 (71.9)	189 (71.1)	571 (71.6)
>100,000 copies/mL	149 (28.1)	77 (28.9)	226 (28.4)
<b>Baseline Plasma HIV RNA n (%)</b>			
<=500,000 copies/mL	506 (95.3)	251 (94.4)	757 (95.0)
>500,000 copies/mL	25 (4.7)	15 (5.6)	40 (5.0)
<b>History of AIDS n (%)</b>			
Yes	79 (14.9)	28 (10.5)	107 (13.4)
No	452 (85.1)	238 (89.5)	690 (86.6)

<b>Stratum n (%)</b>			
Screening HIV RNA ≤ 100,000	382 (71.9)	190 (71.4)	572 (71.8)
Hepatitis B and/or C Positive <sup>††</sup>	15 (2.8)	8 (3.0)	23 (2.9)
<b>Baseline Hepatitis Status</b>			
Hep B Positive Only	11 (2.1)	3 (1.1)	14 (1.8)
Hep C Positive Only	4 (0.8)	4 (1.5)	8 (1.0)
Both Hep B and Hep C Positive	0 (0.0)	1 (0.4)	1 (0.1)
<b>Viral Subtype n (%)</b>			
Clade B	335 (63.1)	186 (69.9)	521 (65.4)
Non-Clade B	194 (36.5)	77 (28.9)	271 (34.0)
Missing	2 (0.4)	3 (1.1)	5 (0.6)
<p>† Subjects with missing results excluded.</p> <p>†† Evidence of hepatitis B surface antigen or evidence of HCV RNA by polymerase chain reaction (PCR) quantitative test for hepatitis C Virus. Nineteen subjects previously classified as hepatitis B or C positive were subsequently identified based on lab tests as being hepatitis B or C negative. Three subjects previously classified as hepatitis B or C negative were subsequently identified based on lab tests as being hepatitis B or C positive.</p> <p>Note: Raltegravir 1200 mg once daily and raltegravir 400 mg twice daily were administered with Truvada*.  N = Number of patients randomized and treated in each treatment group.  n (%) = Number (percent) of patients in each sub-category.</p>			

The ISENTRESS<sup>®</sup> 1200 mg (2 x 600 mg) once daily regimen was non-inferior to the ISENTRESS<sup>®</sup> 400 mg twice daily regimen at both Weeks 48 and 96. At Week 48, 88.9% versus 88.3% of once-daily and twice-daily patients, respectively, had HIV RNA <40 copies/mL. At Week 96, 81.5% versus 80.1% of once-daily and twice-daily patients, respectively, had HIV RNA <40 copies/mL. A summary of antiretroviral response and immunologic effect at Week 48 is shown in Table 23.

**Table 23 - Efficacy Analysis by Treatment Group at Week 48 (Non-Completer = Failure Approach<sup>†</sup>)**

Parameter	Unadjusted Data Summary by Treatment Group		Treatment Difference (once daily – twice daily) <sup>**</sup>
	Raltegravir 1200 mg once daily n/N (%)	Raltegravir 400 mg twice daily n/N (%)	Estimated Difference (95% CI)
<b>Primary</b>			
Proportion of Patients with HIV RNA <40 copies/mL <sup>†</sup>	472/531 (88.9)	235/266 (88.3)	0.5 (-4.2, 5.2) <sup>§</sup>
<b>Supportive</b>			
Proportion of Patients with HIV RNA <50 copies/mL <sup>†</sup>	477/531 (89.8)	240/266 (90.2)	-0.4 (-4.9, 4.0)
Proportion of Patients with HIV RNA <200 copies/mL <sup>†</sup>	484/531 (91.1)	243/266 (91.4)	-0.2 (-4.4, 4.0)
	Mean (95% CI)	Mean (95% CI)	Mean Difference (95% CI)
<b>Secondary</b>			
Change from Baseline in CD4 Cell Count (cells/mm <sup>3</sup> ) <sup>‡</sup>	232 (215, 249)	234 (213, 255)	-2.1 (-31, 27)
<p><sup>**</sup> The 95% CIs for the treatment differences in percent response were calculated using stratum-adjusted Mantel-Haenszel method with the difference weighted by the harmonic mean of sample size per arm for each stratum (screening HIV-1 RNA ≤100,000 copies/mL or HIV-1 RNA &gt;100,000 copies/mL). The 95% CI for mean difference in CD4 change was based on t-distribution.</p> <p><sup>†</sup> Approach to handling missing values: Non-Completer = Failure (NC = F) Approach.</p> <p><sup>‡</sup> OF: Observed Failure approach.</p> <p><sup>§</sup> Raltegravir 1200 mg once daily is concluded non-inferior to raltegravir 400 mg twice daily if the lower bound of the 95% CI for the difference in percent response is above -10 percentage points.</p> <p>Note: Raltegravir 1200 mg once daily and raltegravir 400 mg twice daily were administered with Truvada* N = Number of subjects in each treatment group.</p>			


Virologic outcomes by the Snapshot Approach at Week 48 are shown in Table 24.

**Table 24 - Virologic Outcomes of Randomized Treatment of Protocol 292 at Week 48**

<b>Outcome</b>	<b>Raltegravir 1200 mg once daily (N=531) n (%)</b>	<b>Raltegravir 400 mg twice daily (N=266) n (%)</b>
HIV RNA <40 copies/mL	472 (88.9)	235 (88.3)
HIV RNA ≥ 40 copies/mL**	29 (5.5)	16 (6.0)
No Virologic Data at Week 48 Window	30 (5.6)	15 (5.6)
Reasons		
Discontinued study due to AE or Death†	6 (1.1)	6 (2.3)
Discontinued study for Other Reasons‡	20 (3.8)	7 (2.6)
On study but missing data in window	4 (0.8)	2 (0.8)
<p>** Includes subjects who changed any component of background therapy to a new drug class or changed background components that were not permitted per protocol or changed any background drug in the regimen because of lack of efficacy (perceived or documented) before Week 48, subjects who discontinued study drug or study before Week 48 for lack or loss of efficacy and subjects with HIV RNA equal to or above 40 copies/mL in the Week 48 window (relative day 295 - 378).</p> <p>† Includes subjects who discontinued because of adverse event (AE) or death at any time point from Day 1 through the time window if this resulted in no virologic data on treatment during the specified window.</p> <p>‡ Other Reasons includes: lost to follow-up, non-compliance with study drug, physician decision, pregnancy, withdrawal by subject.</p> <p>Note: Raltegravir 1200 mg once daily and raltegravir 400 mg twice daily were administered with Truvada*. n (%) = Number (Percent) of subjects in each category.</p>		

Figure 1 presents the proportion of patients with HIV RNA <40 copies/mL over time by treatment group.

**Figure 1 - Proportion of Subjects With HIV RNA < 40 copies/mL Over Time (95% CI) Snapshot Approach (Non-Completer=Failure)**



In the ONCEMRK trial, ISENTRESS® 1200 mg (2 x 600 mg) once daily demonstrated consistent virologic and immunologic efficacy relative to ISENTRESS® 400 mg twice daily, both in combination with emtricitabine (+) tenofovir disoproxil fumarate, across demographic and baseline prognostic factors, including: baseline HIV RNA levels >100,000 copies/mL and >500,000 copies/mL, baseline CD4 cells  $\leq 50$  cells/mm<sup>3</sup>, demographic groups (including age, gender, race, ethnicity and region), viral hepatitis co-infection status (hepatitis B and/or C), concomitant proton pump inhibitors/H2 blockers use and viral subtypes (comparing non-clade B as a group to clade B).

Consistent efficacy in patients receiving ISENTRESS® 1200 mg (2 x 600 mg) once daily was observed across HIV subtypes with 94.6% (296/313) and 93.6% (175/187) of patients with B and non-B subtypes, respectively, achieving HIV RNA <40 copies/mL at week 48 (OF approach). Consistent results were observed at week 96.

## **Pediatric Patients**

### **2 to 18 Years of Age**

IMPAACT P1066 is a Phase I/II open label multicenter trial to evaluate the pharmacokinetic profile, safety, tolerability, and efficacy of raltegravir in HIV infected children. This study enrolled 126 treatment experienced children and adolescents 2 through 18 years of age. Patients were stratified by age, enrolling adolescents first and then successively younger children. Patients received either the 400 mg tablet formulation (6 through 18 years of age) or the chewable tablet formulation (2 through 11 years of age). Raltegravir was administered with an optimized background regimen.

The initial dose finding stage included intensive pharmacokinetic evaluation. Dose selection was based upon achieving similar raltegravir plasma exposure and trough concentration as seen in adults, and acceptable short term safety. After dose selection, additional patients were enrolled for evaluation of long term safety, tolerability and efficacy. Of the 126 patients, 96 received the recommended dose of ISENTRESS<sup>®</sup> (see DOSAGE AND ADMINISTRATION).

These 96 patients had a median age of 13 (range 2 through 18) years, were 51% Female, 34% Caucasian, and 59% Black. At baseline, mean plasma HIV-1 RNA was 4.3 log<sub>10</sub> copies/mL, median CD4 cell count was 481 cells/mm<sup>3</sup> (range: 0–2361) and median CD4% was 23.3% (range: 0–44). Overall, 8% had baseline plasma HIV-1 RNA >100,000 copies/mL and 59% had a CDC HIV clinical classification of category B or C. Most patients had previously used at least one NNRTI (78%) or one PI (83%).

Ninety-three (97%) patients 2 through 18 years of age completed 24 weeks of treatment (3 discontinued due to non-compliance). At Week 24, 72% achieved  $\geq 1$  log<sub>10</sub> HIV RNA drop from baseline or <400 copies/mL; 54% achieved HIV RNA <50 copies/mL. The mean CD4 count (percent) increase from baseline to Week 24 was 119 cells/mm<sup>3</sup> (3.8%).

Seventy-two (95%) patients 6 through 18 years of age completed 48 weeks of treatment (4 discontinued due to non-compliance). At Week 48, 77% achieved  $\geq 1$  log<sub>10</sub> HIV RNA drop from baseline or <400 copies/mL; 56% achieved HIV RNA <50 copies/mL. The mean CD4 count (percent) increase from baseline to Week 48 was 155 cells/mm<sup>3</sup> (4.7%).

## **TOXICOLOGY**

### **Acute Toxicity**

In dogs, an intravenous 3-day rising dose escalation study caused mortality at high doses; this is considered to result from cardiac arrhythmia secondary to the excessive potassium salt administered in the drug formulation. Mild physical signs were noted at lower doses. In a 7-day intravenous study in dogs, at 100 mg/kg/day (exposure approximately 23-fold above the exposure at the recommended human dose), treatment-related effects were limited to physical signs which included body weight loss; minimal increases in serum urea nitrogen; increases in alanine aminotransferase activity, alkaline phosphatase activity, and cholesterol; and very slight renal tubular dilatation.



## **Chronic Toxicity**

Chronic repeat dose toxicity studies were conducted in rats (6 month duration) and dogs (1 year duration). In dogs, transient and/or intermittent emesis and weight loss were observed at 360 mg/kg/day (exposure 9 fold above the exposure at the recommended human dose). In rats, mortality, preceded by physical signs of drug intolerance, was seen at 600 mg/kg/day (exposures 4.8 fold above the exposure at the recommended human dose), but not at 120 mg/kg/day. In rats, inflammation of the nasal cavity and degeneration of the stomach mucosa occurred at 120 mg/kg/day (exposures 1.6 fold above the exposure at the recommended human dose) and is suggestive of irritative properties of the drug.

## **Carcinogenesis, Mutagenesis, Impairment of Fertility**

Carcinogenicity studies of raltegravir in mice did not show any carcinogenic potential. At the highest dose levels, 400 mg/kg/day in females and 250 mg/kg/day in males, systemic exposure was approximately 2-fold greater (females) or equal to (males) the AUC (54  $\mu\text{M}\cdot\text{hr}$ ) at the 400-mg twice daily dose. In rats, carcinogenic potential considered to be specific for this species was identified, but is regarded as having minimal relevance for humans. In rats, tumors (squamous cell carcinoma) of the nose/nasopharynx were identified in high- and mid-dose group animals. These neoplasms are considered to result from local deposition and/or aspiration of drug on the mucosa of the nose/nasopharynx during dosing and are an expected consequence of chronic irritation and inflammation. Consistent with this, the increased incidence of these neoplasms correlated with oral dosing of high concentrations of raltegravir (>300 mg/kg) instead of systemic exposure. However, at the NOAEL, systemic exposure was 1.4 to 1.7 fold greater than the AUC (54  $\mu\text{M}\cdot\text{hr}$ ) at the clinical 400-mg twice daily dose.

All genotoxicity studies to evaluate mutagenicity and clastogenicity were negative.

No effect on fertility was seen in male and female rats at doses up to 600 mg/kg/day which resulted in 3 fold exposure above the exposure at the recommended human dose.

## **Development**

Oral administration of up to 600 mg/kg/day to juvenile rats resulted in drug irritation effects in the stomach which were similar to those seen in adult rats. No additional toxicities were noted in juvenile rats indicating that juvenile rats were no more sensitive to drug effects than adult rats.

Treatment-related increases over controls in the incidence of supernumerary ribs were seen in rats at 600 mg/kg/day (exposures 4.4- fold above the exposure at the recommended human dose).

## **REFERENCES**

1. Cahn, P, Kaplan R, Sax P, Squires, K, Molina JM, Avihingsanon A, et al. Raltegravir (RAL) 1200 mg Once Daily (QD) is Non-Inferior to RAL 400 mg Twice Daily (BID), in Combination with Tenofovir/Emtricitabine, in Treatment-Naïve HIV-1 Infected Subjects: Week 48 Results. Abstract Submitted. Conference: 21<sup>st</sup> International AIDS Conference · Reference No: A-792-0138-10312.

2. Croteau D, Letendre S, Best B.M., Ellis RJ, Breidinger S, Clifford D, et al. Total raltegravir concentrations in cerebrospinal fluid exceed the 50-percent inhibitory concentration for wild-type HIV-1. *Antimicrob Agents Chemother* 2010; 54(12):5156-60.
3. Grinsztejn B, Nguyen B-Y, Katlama C, Gatell JM, Lazzarin A, Vittecoq D, Gonzalez CJ, Chen J, Harvey C, Isaacs RD. Safety and efficacy of the HIV-1 integrase inhibitor raltegravir (MK0518) in treatment-experienced patients with multidrug-resistant virus: a phase II randomised controlled trial. *Lancet* 2007; 369:1261-69.
4. Markowitz M, Morales-Ramirez JO, Nguyen B-Y, Kovacs CM, Steigbigel RT, Cooper DA, Liporace R, Schwartz R, Isaacs R, Gilde LR, Wenning L, Zhao J, Teppler H. Antiretroviral activity, pharmacokinetics, and tolerability of MK-0518, a novel inhibitor of HIV-1 integrase, dosed as monotherapy for 10 days in treatment-naïve HIV-1-infected individuals. 2006. *J Acquir Immune Defic Syndr* 2006;43:509-515.
5. Nachman S, Zheng N, Acosta E, Teppler H, Homony B, Graham B, et al. Pharmacokinetics, Safety and 48 Week Efficacy of oral Raltegravir in Human Immunodeficiency Virus type-1 (HIV) infected Children 2 through 18 years of Age. *Clinical Infectious Diseases Advance Access published October 21, 2013.*
6. Rockstroh JK, DeJesus E, Lennox JL, Yazdanpanah Y, Saag MS, Wan H, Rodgers AJ, et al. Durable Efficacy and Safety of Raltegravir Versus Efavirenz When Combined With Tenofovir/Emtricitabine in Treatment-Naïve HIV-1–Infected Patients: Final 5-Year Results From STARTMRK. *J Acquir Immune Defic Syndr* 2013; 63, No 1: 77-85.
7. Yilmaz A, Gisslen M, Spudich S, Lee E, Jayewardene A, Aweeka F, et al. Raltegravir cerebrospinal fluid concentrations in HIV-1 infection. *PLoS ONE* 2009;4(9):e6877.

## PART III: CONSUMER INFORMATION

**ISENTRESS<sup>®</sup>**

\* Raltegravir tablets 400 mg

\* Raltegravir chewable tablets, 25 mg, 100 mg  
(\* as raltegravir potassium)**ISENTRESS HD<sup>®</sup>**Raltegravir tablets 600 mg  
(as raltegravir potassium)

This leaflet is part III of a three-part “Product Monograph” published when ISENTRESS<sup>®</sup> was approved for sale in Canada and is designed specifically for Consumers. This leaflet is a summary and will not tell you everything about ISENTRESS<sup>®</sup>. Contact your physician or pharmacist if you have any questions about the drug.

**ABOUT THIS MEDICATION**What is HIV?

HIV is a disease that destroys the body’s ability to protect itself from infection.

- It is caused by a virus (called HIV) that attacks certain white blood cells and weakens the immune system.
- HIV is spread by contact with blood of or sexual contact with an infected person.

What the medication is used for:

- ISENTRESS<sup>®</sup> is a medicine that helps control HIV infection, in combination with other antiretroviral medications.
- Your physician has prescribed ISENTRESS<sup>®</sup> to help control your HIV infection.
- ISENTRESS<sup>®</sup> may be used to treat children and adolescents that are 2 years of age and older.

What it does:

ISENTRESS<sup>®</sup> is a medicine that helps to control HIV infection. The term HIV stands for Human Immunodeficiency Virus. It is the virus that causes AIDS (Acquired Immune Deficiency Syndrome).

**How does ISENTRESS<sup>®</sup> work?**

- ISENTRESS<sup>®</sup> blocks an enzyme which the virus (HIV) needs in order to make more virus. The enzyme that ISENTRESS<sup>®</sup> blocks is called HIV integrase.
- ISENTRESS<sup>®</sup> is used in combination with other HIV medicines to treat adults, adolescents, and children 2 years of age and older who are infected with HIV. Anti-HIV medicines are also called “antiretroviral medicines” or an-ti-re-tro-vi-ral medicines.

**When used with other anti-HIV medicines, ISENTRESS<sup>®</sup> may do two things:**

1. It may reduce the amount of HIV in your blood. This is called your “viral load”.
  - Reducing the amount of HIV in the blood may keep your immune system healthy.
  - This in turn, can help your immune system to fight infection.
2. It may also increase the number of white blood cells that help fight the virus (HIV).
  - Physicians call them CD4 (T) cells.

ISENTRESS<sup>®</sup> may not have these effects in all patients.

When it should not be used:

- Do not take ISENTRESS<sup>®</sup> if you are hypersensitive to any of its ingredients (see What the non-medicinal ingredients are).

What the medicinal ingredient is:

Raltegravir potassium

What the non-medicinal ingredients are:600 mg film-coated tablets:

croscarmellose sodium, hypromellose 2910, magnesium stearate, microcrystalline cellulose. In addition, the film coating contains the following inactive ingredients: black iron oxide, carnauba wax, hypromellose 2910, iron oxide yellow, lactose monohydrate, titanium dioxide, and triacetin.

400 mg raltegravir film-coated tablets

calcium phosphate dibasic anhydrous, hypromellose 2208, lactose monohydrate, magnesium stearate, microcrystalline cellulose, poloxamer 407 (contains 0.01% butylated hydroxytoluene as antioxidant), sodium stearyl fumarate.

In addition, the film coating contains the following inactive ingredients: black iron oxide, polyethylene glycol 3350, polyvinyl alcohol, red iron oxide, talc and titanium dioxide.

25 mg and 100 mg raltegravir chewable tablets:

ammonium hydroxide, crospovidone, ethylcellulose 20 cP, fructose, hydroxypropyl cellulose, hypromellose 2910/6cP, macrogol/PEG 400, magnesium stearate, mannitol, medium chain triglycerides, monoammonium glycyrrhizinate, natural and artificial flavors (orange, banana, and masking), oleic acid, phenylalanine (as part of the artificial sweetener, aspartame), saccharin sodium, sodium citrate dihydrate, sodium stearyl fumarate, sorbitol, sucralose and yellow iron oxide.

ISENTRESS<sup>®</sup> 100 mg also contain red iron oxide

What dosage forms it comes in:

ISENTRESS HD<sup>®</sup> 600 mg film-coated tablets are yellow, oval shaped tablets debossed with Merck logo and “242” on one side.

ISENTRESS<sup>®</sup> 400 mg film-coated tablets are available as pink, oval shaped, with 227 on one side.

ISENTRESS<sup>®</sup> 100 mg orange-banana flavoured, chewable tablets are available as pale orange, oval shaped scored tablet, debossed with the Merck logo on one side of the score and 477 on the other, and scored on the other side of the tablet.

ISENTRESS<sup>®</sup> 25 mg orange-banana flavoured, chewable tablets are available as pale yellow, round, flat faced, beveled edge tablet debossed with the Merck logo on one side and 473 on the other side of the tablet.

## WARNINGS AND PRECAUTIONS

**Tell your physician about all the medicines you take** (see INTERACTIONS WITH THIS MEDICATION)

BEFORE you use ISENTRESS<sup>®</sup> talk to your physician or pharmacist if:

- You have any allergies.
- You take antacids. Certain antacids (those containing aluminum and/or magnesium) are not recommended with ISENTRESS<sup>®</sup>.
- You have phenylketonuria (PKU). ISENTRESS<sup>®</sup> Chewable Tablets contain phenylalanine as part of the artificial sweetener, aspartame. The artificial sweetener may be harmful to people with PKU.
- You are pregnant or plan to become pregnant.
  - If you take ISENTRESS<sup>®</sup> while you are pregnant, talk to your physician about how you can be included in the Antiretroviral Pregnancy Registry.
- You are breast-feeding or plan to breast-feed.
  - It is recommended that HIV-infected women should not breast-feed their infants. This is because their babies could be infected with HIV through their breast milk.
  - Talk with your physician about the best way to feed your baby.

### Other Warnings

**Does ISENTRESS<sup>®</sup> lower the chance of passing HIV to other people?**

No. ISENTRESS<sup>®</sup> does not reduce the chance of passing HIV to others through sexual contact, sharing needles, or being exposed to your blood.

- Continue to practice safer sex.
- Use latex or polyurethane condoms or other barrier methods to lower the chance of sexual contact with any body fluids. This includes semen from a man, vaginal secretions from a woman, or blood.
- Never re-use, or share needles.
- Ask your physician if you have any questions about safer sex or how to prevent passing HIV to other people.

**What else is there to know?**

- ISENTRESS<sup>®</sup> does not cure HIV infection or AIDS.
- It is very important that you stay under the care of your physician during treatment with ISENTRESS<sup>®</sup>.

## INTERACTIONS WITH THIS MEDICATION

Tell your physician about all the medicines you take including prescription and non-prescription medicines (including antacids), vitamins, and herbal supplements. Some medicines are not recommended with ISENTRESS<sup>®</sup>.

Tell your physician especially if you take:

- antacids containing aluminum, magnesium and calcium carbonate
- atazanavir
- rifampin
- tipranavir/ritonavir

Keep a list of all your medicines to show your physician and pharmacist.

- You can ask your healthcare provider or pharmacist for a list of medicines that interact with ISENTRESS<sup>®</sup>.
- Do not start taking a new medicine without telling your healthcare provider. Your healthcare provider can tell you if it is safe to take ISENTRESS<sup>®</sup> with other medicines.

## PROPER USE OF THIS MEDICATION

Take ISENTRESS<sup>®</sup> exactly as your physician has prescribed.

### Usual dose:

Take ISENTRESS<sup>®</sup> by mouth, with or without food.

ISENTRESS<sup>®</sup> must be used with other anti-HIV medicines.

Do not change your dose, or stop taking ISENTRESS<sup>®</sup> without first talking with your physician. Do not switch from the 400 mg tablet, 600 mg tablets, or the chewable tablets.

### Adult

Take one 400-mg tablet twice daily or two 600 mg tablets once daily as directed by your healthcare provider.

### Children and Adolescents

- Your child's healthcare provider will tell you the right dose and type of ISENTRESS<sup>®</sup> (tablet, chewable tablet) based on your child's weight.
- The 100 mg chewable tablet is scored and can be split into equal halves.
- Do not take more than 300 mg of the chewable tablet twice a day.

### Using different dosage forms/strengths:

The different dosage forms and strengths for ISENTRESS<sup>®</sup> are not interchangeable so:

- a) Use only the 600 mg tablet for the 1200 mg once daily dose.
- b) Use only the 400 mg tablet for the twice daily dose of 400 mg.

c) Use the 25 mg and 100 mg chewable tablets for only the children's dosages up to 300 mg twice daily.

**IMPORTANT: Take ISENTRESS<sup>®</sup> exactly as your physician prescribed and at the right times of day because if you don't:**

- The amount of virus (HIV) in your blood may increase if the medicine is stopped for even a short period of time.
- The virus may develop resistance to ISENTRESS<sup>®</sup> and become harder to treat.
- Your medicines may stop working to fight HIV.
- The activity of ISENTRESS<sup>®</sup> may be reduced (due to resistance).

Overdose:

If you think you have taken too much ISENTRESS<sup>®</sup>, contact your healthcare professional, hospital emergency department or regional poison control centre immediately, even if there are no symptoms.

Missed Dose:

If you miss a dose, take it as soon as you remember. If you do not remember until it is time for your next dose, skip the missed dose and go back to your regular schedule. Only take ISENTRESS<sup>®</sup> and ISENTRESS HD<sup>®</sup> as instructed.

ISENTRESS<sup>®</sup> must be used with other anti-HIV medicines.

**Be sure to keep a supply of your anti-HIV medicines.**

- When your ISENTRESS<sup>®</sup> supply starts to run low, get more from your physician or pharmacy.
- Do not wait until your medicine runs out to get more.

**SIDE EFFECTS AND WHAT TO DO ABOUT THEM**

Like all medicines, ISENTRESS<sup>®</sup> can cause side effects, although not everybody gets them. In studies, side effects usually were mild and did not cause patients to stop taking ISENTRESS<sup>®</sup>. The side effects reported in patients taking ISENTRESS<sup>®</sup> were similar to side effects in patients treated with a tablet containing no medication (a placebo).

The most common side effects of ISENTRESS<sup>®</sup> include:

- nausea
- headache
- diarrhea
- fever
- vomiting
- fatigue
- dizziness
- difficulty sleeping
- cough
- rash
- tiredness
- upper respiratory tract infection
- inflammation of the nasal passages and throat

- bronchitis
- back pain
- depression

Additionally, while the medicine has been on the market, some further reactions have occurred:

- Depression and suicidal thoughts have been reported. If you develop these feelings, discuss this with your physician.
- Other side effects that have been reported include low blood platelet count, clumsiness and lack of coordination, rash with or without an increase in some white blood cells, severe skin reaction, liver failure. If you develop any of these reactions, discuss with your physician.

In some patients with advanced HIV infection (AIDS), signs and symptoms of inflammation from opportunistic infections may occur when combination antiretroviral treatment is started. Tell your physician immediately if you notice any symptoms of infection.

Phenylketonuria (PKU). ISENTRESS<sup>®</sup> Chewable Tablets contain phenylalanine as part of the artificial sweetener, aspartame. The artificial sweetener may be harmful to people with PKU.

Contact your physician promptly if you experience unexplained muscle pain, tenderness, or weakness while taking ISENTRESS<sup>®</sup>.

Contact your physician promptly if you develop a rash. Severe and life-threatening skin reactions and allergic reactions have been reported in some patients taking ISENTRESS<sup>®</sup>.

Changes in your immune system (Immune Reconstitution Inflammatory Syndrome) can happen when you start taking HIV medicines. Your immune system may get stronger and begin to fight infections that have been hidden in your body for a long time, or you could develop an autoimmune disease in which your immune system reacts against your own body (e.g. Grave's disease (which affects the thyroid gland), Guillain-Barre syndrome (which affects the nervous system) or polymyositis (which affects the muscles) and it may develop at any time, sometimes months later after the start of HIV therapy). Sometimes symptoms can be severe, so if you develop high temperature (fever), joint or muscle pain, redness, rash, swelling, or fatigue or any new symptoms contact your physician right away.

Tell your physician if you develop any unusual side effect or if any known side effect does not go away or gets worse. For more information, ask your physician or pharmacist.

Keep ISENTRESS<sup>®</sup> and all medicines out of the reach and sight of children.

**SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM**

Symptoms / effects	Talk with your physician or pharmacist		Stop taking drug and call your physician or pharmacist
	Only if severe	In all cases	
<b>Uncommon</b>			
<u>Severe skin reactions and allergic reactions</u> : occasionally life-threatening, with symptoms such as rash, itching or hives on the skin, swelling of the face, lips, tongue or other parts of the body, shortness of breath, wheezing or trouble breathing			✓
persistent fatigue	✓		
<u>lack of white blood cells</u> : frequent infections such as fever, severe chills, sore throat or mouth ulcers	✓		
<u>lack of red blood cells</u> : tiredness, headaches, being short of breath when exercising, dizziness and looking pale	✓		
severe chest pain			✓
<u>stomach problems</u> : pain, nausea, vomiting, heartburn	✓		
<u>liver disease</u> : liver disease with nausea, vomiting, loss of appetite, feeling generally unwell, fever, itching, yellowing of the skin and eyes, and dark coloured urine		✓	
<u>kidney disease</u> : nausea, loss of appetite and weakness, pass little or no urine, breathlessness			✓
depression, suicidal thoughts and actions		✓	
shaking		✓	
speech disorders		✓	
disturbance in attention		✓	

*This is not a complete list of side effects. For any unexpected effects while taking ISENTRESS<sup>®</sup>, contact your physician or pharmacist.*

**HOW TO STORE IT**

Store ISENTRESS<sup>®</sup> and ISENTRESS HD<sup>®</sup> at room temperature (15°C–30°C).

400 mg tablets:

Keep the desiccant in the bottle.

600 mg tablets, Chewable tablets 25 mg and 100 mg: Store in the original package with the bottle tightly closed. Keep the desiccant in the bottle to protect from moisture.

**Reporting Side Effects**

You can help improve the safe use of health products for Canadians by reporting serious and unexpected side effects to Health Canada. Your report may help to identify new side effects and change the product safety information.

**3 ways to report:**

- Online at MedEffect (<https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada.html>);
  - By calling 1-866-234-2345 (toll-free);
  - By completing a Consumer Side Effect Reporting Form and sending it by:
    - Fax to 1-866-678-6789 (toll-free), or
    - Mail to: Canada Vigilance Program  
Health Canada, Postal Locator 1908C  
Ottawa, ON  
K1A 0K9
- Postage paid labels and the Consumer Side Effect Reporting Form are available at MedEffect ([https://www.canada.ca/en/health-products/medeffect-canada.html](https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada.html)).

*NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.*

**MORE INFORMATION****If you want more information about ISENTRESS<sup>®</sup>:**

- Talk to your healthcare professional
- Find the full product monograph that is prepared for healthcare professionals and includes this Consumer Information by visiting the Health Canada website (<https://www.canada.ca/en/health-canada/services/drugs-health-products/drug-products/drug-product-database.html>) or Merck Canada website [www.merck.ca](http://www.merck.ca) or by calling Merck Canada at 1-800-567-2594.

To report an adverse event related to ISENTRESS<sup>®</sup>, please contact 1-800-567-2594.

This leaflet was prepared by Merck Canada Inc.

Last revised: September 20, 2018

<sup>®</sup> Merck Sharp & Dohme Corp. Used under license.

\* All other trademarks are the property of their respective owners.

© 2011, 2018 Merck Canada Inc. All rights reserved.