

PRODUCT MONOGRAPH

<sup>Pr</sup>**SANDOZ VALPROIC**

Valproic Acid

250 mg Capsules

Mfr. Std.

Antiepileptic

Sandoz Canada Inc.  
110 Rue de Lauzon  
Boucherville, QC, Canada  
J4B 1E6

Date of Revision: December 18, 2018

Submission Control No: 222665

## Table of Contents

<b>PART I: HEALTH PROFESSIONAL INFORMATION .....</b>	<b>3</b>
SUMMARY PRODUCT INFORMATION.....	3
INDICATIONS AND CLINICAL USE .....	3
CONTRAINDICATIONS.....	4
WARNINGS AND PRECAUTIONS .....	4
ADVERSE REACTIONS .....	19
DRUG INTERACTIONS .....	23
DOSAGE AND ADMINISTRATION .....	31
OVERDOSAGE.....	33
ACTION AND CLINICAL PHARMACOLOGY .....	34
STORAGE AND STABILITY .....	36
DOSAGE FORMS, COMPOSITION AND PACKAGING.....	36
<b>PART II: SCIENTIFIC INFORMATION .....</b>	<b>37</b>
PHARMACEUTICAL INFORMATION .....	37
CLINICAL TRIALS .....	38
DETAILED PHARMACOLOGY .....	40
TOXICOLOGY.....	40
REFERENCES.....	43
<b>PART III: CONSUMER INFORMATION .....</b>	<b>45</b>

**Pr SANDOZ VALPROIC**

Valproic Acid

**PART I: HEALTH PROFESSIONAL INFORMATION**

**SUMMARY PRODUCT INFORMATION**

<b>Route of Administration</b>	<b>Dosage Form / Strength</b>	<b>Non-medicinal Ingredients</b>
Oral	Capsule / 250 mg	Corn oil, FD&C Yellow No. 6, gelatin, glycerin, methylparaben*, propylparaben*, titanium dioxide  * methylparaben and propylparaben may cause allergic reactions (possibly delayed).

**INDICATIONS AND CLINICAL USE**

Sandoz Valproic (valproic acid) is indicated for:

- use as sole or adjunctive therapy in the treatment of simple or complex absence seizures, including petit mal, and is useful in primary generalized seizures with tonic-clonic manifestations.
- use adjunctively in patients with multiple seizure types which include either absence or tonic-clonic seizures.

In accordance with the International Classification of Seizures, simple absence is defined as a very brief clouding of the sensorium or loss of consciousness (lasting usually 2 to 15 seconds), accompanied by certain generalized epileptic discharges without other detectable clinical signs. Complex absence is the term used when other signs are also present.

See CONTRAINDICATIONS and WARNINGS AND PRECAUTIONS, Hepatic/Biliary/Pancreatic, Serious or Fatal Hepatotoxicity, for statement regarding serious or fatal hepatic dysfunction.

**Geriatrics (≥ 65 years of age):** The safety and efficacy of valproic acid in elderly patients with epilepsy has not been evaluated in clinical trials. Caution should thus be exercised in dose selection for an elderly patient, recognizing the more frequent hepatic and renal dysfunctions, and limited experience with valproic acid in this population. For a brief discussion, see WARNINGS AND PRECAUTIONS, Special Populations, Geriatrics (≥ 65 years of age); DOSAGE AND ADMINISTRATION and ACTION AND CLINICAL PHARMACOLOGY, Special Populations and Conditions, Geriatrics.

**Pediatrics (< 18 years of age):** When Sandoz Valproic is used in children under the age of 2 years, it should be used with extreme caution and as a sole agent. Above the age of 2 years, experience in epilepsy has indicated that the incidence of fatal hepatotoxicity decreases considerably in progressively older patient groups. For a brief discussion, see WARNINGS AND PRECAUTIONS, Special Populations, Pediatrics (< 18 years of age).

## CONTRAINDICATIONS

Sandoz Valproic (valproic acid) is contraindicated in:

- patients with hepatic disease or significant hepatic dysfunction (see WARNINGS AND PRECAUTIONS, Serious Warnings and Precautions, Hepatotoxicity and WARNINGS AND PRECAUTIONS, Hepatic/Biliary/Pancreatic, Serious or Fatal Hepatotoxicity).
- patients known to have mitochondrial disorders caused by mutations in mitochondrial DNA polymerase  $\gamma$  (POLG; e.g., Alpers-Huttenlocher Syndrome) and children under two years of age who are suspected of having a POLG-related disorder (see WARNINGS AND PRECAUTIONS, Hepatic/Biliary/Pancreatic).
- patients with known hypersensitivity to the drug, any ingredient in the formulation or component of the container. For a complete listing of ingredients (see the DOSAGE FORMS, COMPOSITION AND PACKAGING).
- patients with known urea cycle disorders (see WARNINGS AND PRECAUTIONS, Endocrine and Metabolism, Urea Cycle Disorders).
- patients with known porphyria.

## WARNINGS AND PRECAUTIONS

### Serious Warnings and Precautions

- **Hepatotoxicity:** Hepatic failure resulting in fatalities has occurred in patients receiving valproic acid. These incidences usually occurred during the first 6 months of treatment with valproic acid. Caution should be observed when administering Sandoz Valproic (valproic acid) to patients with a prior history of hepatic disease. Patients on multiple anticonvulsants, children, those with congenital metabolic disorders, those with severe seizure disorders accompanied by mental retardation, and those with organic brain disease may be at particular risk. Experience has indicated that children under the age of 2 years are at a considerably increased risk of developing fatal hepatotoxicity, especially those on multiple anticonvulsants (see WARNINGS AND PRECAUTIONS, Hepatic/Biliary/Pancreatic, Serious or Fatal Hepatotoxicity).
- **Female children/Female adolescents/Women of childbearing potential/Pregnancy (Teratogenicity):** Sandoz Valproic (valproic acid) should not be used in female children, in female adolescents, in women of childbearing potential and pregnant women unless alternative treatments are ineffective or not tolerated because of its high teratogenic potential

and risk of developmental disorders in infants exposed in utero to valproate. Sandoz Valproic can produce teratogenic effects such as neural tube defects (e.g., spina bifida) in a dose-dependent manner. In addition, valproate can cause decreased IQ scores following in utero exposure. The benefit and risk should be carefully reconsidered at regular treatment reviews, at puberty and urgently when a woman of childbearing potential treated with Sandoz Valproic plans a pregnancy or if she becomes pregnant.

- Women of childbearing potential must use effective contraception during treatment and be informed of the risks associated with the use of Sandoz Valproic during pregnancy. In women planning to become pregnant every effort should be made to switch to appropriate alternative treatment prior to conception (see WARNINGS AND PRECAUTIONS, Special Populations, Women of Childbearing Potential and WARNINGS AND PRECAUTIONS, Special Populations, Pregnant Women.)
- **Patients with Mitochondrial Disease:** There is an increased risk of valproate-induced acute liver failure and resultant deaths in patients with hereditary neurometabolic syndromes caused by DNA mutations of the mitochondrial DNA Polymerase  $\gamma$  (POLG) gene (e.g. Alpers Huttenlocher Syndrome). Valproic acid is contraindicated in patients known to have mitochondrial disorders caused by POLG mutations and children under two years of age who are clinically suspected of having a mitochondrial disorder (see CONTRAINDICATIONS). In patients over two years of age who are clinically suspected of having a hereditary mitochondrial disease, Sandoz Valproic should only be used after other anticonvulsants have failed. This older group of patients should be closely monitored during treatment with valproic acid for the development of acute liver injury with regular clinical assessments and serum liver testing. POLG mutation screening should be performed in accordance with current clinical practice (see WARNINGS AND PRECAUTIONS, Hepatic/Biliary/Pancreatic, Patients with Mitochondrial Disease).
- **Pancreatitis:** Cases of life-threatening pancreatitis have been reported in both children and adults receiving valproic acid. Some of the cases have been described as hemorrhagic with a rapid progression from initial symptoms to death. Patients and guardians should be warned that abdominal pain, nausea, vomiting and/or anorexia can be symptoms of pancreatitis that require prompt medical attention. If pancreatitis is diagnosed, Sandoz Valproic should ordinarily be discontinued. Alternative treatment for the underlying medical condition should be initiated as clinically indicated. Some cases have occurred shortly after initial use as well as after several years of use (see WARNINGS AND PRECAUTIONS, Hepatic/Biliary/Pancreatic, Pancreatitis).

### **General**

Antiepileptic drugs (AEDs), including Sandoz Valproic (valproic acid), should be withdrawn gradually to minimize the potential for seizures or increased seizure frequency (see DOSAGE AND ADMINISTRATION, Recommended Dose and Dosage Adjustment).

**Interaction with Carbapenem Antibiotics:** Carbapenem antibiotics (ertapenem, imipenem, meropenem, doripenem) can reduce serum valproic acid concentrations to sub-therapeutic levels.

This can result in loss of seizure control in epileptic patients or loss of efficacy in non-epileptics. In some cases of co-administration in epileptic patients, breakthrough seizures have occurred. Increasing valproic acid dose may not be sufficient to overcome this interaction. If co-administration is essential, serum valproic acid concentrations should be monitored daily after initiating carbapenem therapy. Alternative antibacterial or anticonvulsant therapy should be considered if serum valproic acid concentrations drop significantly or seizure control deteriorates (see DRUG INTERACTIONS, Drug-Drug Interactions, Table 1).

**Effects of Valproate on HIV and CMV Viruses Replication:** There are *in vitro* studies that suggest valproate stimulates the replication of the HIV (Human Immunodeficiency Virus) and CMV (Cytomegalovirus) viruses under certain experimental conditions. The clinical relevance of these *in vitro* data is unknown. Additionally, the relevance of these *in vitro* findings is uncertain for patients receiving maximally suppressive antiretroviral therapy. Nevertheless, these data should be borne in mind when interpreting the results from regular monitoring of the viral load in HIV infected patients receiving valproate or when following CMV infected patients clinically.

### **Carcinogenesis and Mutagenesis**

Long-term animal toxicity studies indicate that valproic acid is a weak carcinogen or promoter in rats and mice. The significance of these findings for humans is unknown at present (see TOXICOLOGY, Mutagenicity and Carcinogenicity).

### **Endocrine and Metabolism**

**Urea Cycle Disorders:** Sandoz Valproic (valproic acid) is contraindicated in patients with known urea cycle disorders. Hyperammonemic encephalopathy, sometimes fatal, has been reported following initiation of valproic acid in patients with urea cycle disorders, a group of uncommon genetic abnormalities, particularly ornithine transcarbamylase deficiency. Prior to initiation of Sandoz Valproic (valproic acid), evaluation for urea cycle disorders (UCD) should be considered in the following patients:

- 1) those with a history of unexplained encephalopathy or coma, encephalopathy associated with protein load, pregnancy-related or postpartum encephalopathy, unexplained mental retardation, or history of elevated plasma ammonia or glutamine;
- 2) those with signs and symptoms of UCD, for example, cyclical vomiting and lethargy, episodic extreme irritability, ataxia, low blood urea nitrogen (BUN), protein avoidance;
- 3) those with a family history of UCD or a family history of unexplained infant deaths (particularly males);
- 4) those with other signs or symptoms of UCD. Patients receiving Sandoz Valproic (valproic acid) who develop symptoms of unexplained hyperammonemic encephalopathy should receive prompt treatment (including discontinuation of Sandoz Valproic (valproic acid)) and be evaluated for underlying urea cycle disorders (see CONTRAINDICATIONS and WARNINGS AND PRECAUTIONS, Endocrine and Metabolism, Hyperammonemia and Hyperammonemia and Encephalopathy Associated with Concomitant Topiramate Use).

**Hyperammonemia:** Hyperammonemia has been reported in association with valproic acid and

may be present despite normal liver function tests. In patients who develop unexplained lethargy and vomiting or changes in mental status, hyperammonemic encephalopathy should be considered as a possible cause and serum ammonia level should be measured. Hyperammonemia should also be considered in patients with hypothermia (see WARNINGS AND PRECAUTIONS, Endocrine and Metabolism, Hypothermia). If serum ammonia is increased, Sandoz Valproic (valproic acid) should be discontinued. Appropriate interventions for treatment of hyperammonemia should be initiated, and such patients should undergo investigation for underlying urea cycle disorders (see CONTRAINDICATIONS and WARNINGS AND PRECAUTIONS, Endocrine and Metabolism, Urea Cycle Disorders and Hyperammonemia and Encephalopathy Associated with Concomitant Topiramate Use).

Asymptomatic elevations of serum ammonia are more common and, when present, require close monitoring of serum ammonia levels. If the elevation persists, discontinuation of Sandoz Valproic (valproic acid) should be considered.

**Hyperammonemia and Encephalopathy Associated with Concomitant use of Topiramate, Acetazolamide, Phenobarbital or Phenytoin:** Concomitant administration of topiramate, acetazolamide, phenobarbital or phenytoin and valproic acid has been associated with hyperammonemia with or without encephalopathy in patients who have tolerated either drug alone. Clinical symptoms of hyperammonemic encephalopathy often include acute alterations in level of consciousness and/or cognitive function with lethargy or vomiting. Hypothermia can also be a manifestation of hyperammonemia (see WARNINGS AND PRECAUTIONS, Endocrine and Metabolism, Hypothermia). In most cases, symptoms and signs abated with discontinuation of either drug. This adverse event is not due to a pharmacokinetic interaction.

It is not known if topiramate, acetazolamide, phenobarbital or phenytoin monotherapy is associated with hyperammonemia.

**Patients with inborn errors of metabolism or reduced hepatic mitochondrial activity may be at an increased risk for hyperammonemia with or without encephalopathy. Although not studied, an interaction of topiramate, acetazolamide, phenobarbital or phenytoin and Sandoz Valproic (valproic acid) may exacerbate existing defects or unmask deficiencies in susceptible persons (see CONTRAINDICATIONS and WARNINGS AND PRECAUTIONS, Endocrine and Metabolism, Urea Cycle Disorders and Hyperammonemia).**

**Hypothermia:** Hypothermia, defined as an unintentional drop in core body temperature to < 35°C (95°F), has been reported in association with valproic acid both in conjunction with and in the absence of hyperammonemia. This adverse reaction can also occur in patients using concomitant topiramate with valproic acid after starting topiramate treatment or after increasing the daily dose of topiramate (see DRUG INTERACTIONS, Drug-Drug Interactions, Table 1). Hypothermia may be manifested by a variety of clinical abnormalities including, lethargy, confusion, coma, and significant alterations in other major organ systems such as the cardiovascular and respiratory systems. Clinical management and assessment should include examination of blood ammonia levels. Consideration should be given to stopping Sandoz Valproic (valproic acid) in patients who develop hypothermia (see WARNINGS AND PRECAUTIONS, Endocrine and Metabolism, Hyperammonemia).

## **Hematologic**

**Thrombocytopenia:** Because of reports of thrombocytopenia and inhibition of the second phase of platelet aggregation, and abnormal coagulation parameters (e.g., low fibrinogen), platelet counts and coagulation tests are recommended before initiating therapy and at periodic intervals. It is recommended that patients receiving Sandoz Valproic (valproic acid) be monitored for platelet count and coagulation parameters prior to planned surgery. Clinical evidence of hemorrhage, bruising or a disorder of hemostasis/coagulation is an indication for reduction of Sandoz Valproic (valproic acid) dosage or withdrawal of therapy (see also WARNINGS AND PRECAUTIONS, Hematologic, Dosing-related Adverse Reactions: Thrombocytopenia).

In addition, the findings from a crossover clinical trial conducted with divalproex sodium extended-release tablets, in 44 epilepsy patients, indicate that the frequency of treatment-emergent mild thrombocytopenia (platelet count between 100 to 150 x 10<sup>9</sup>/L) was significantly higher after 12 weeks of treatment with divalproex sodium extended-release tablets than after 12 weeks of treatment with divalproex sodium (7 versus 3 low counts, respectively).

**Dosing-related Adverse Reactions: Thrombocytopenia:** The frequency of adverse effects thrombocytopenia (particularly elevated liver enzymes and thrombocytopenia) may be dose-related. In a clinical trial of divalproex sodium as monotherapy in patients with epilepsy, 34/126 patients (27%) receiving approximately 50 mg/kg/day on average, had at least one value of platelets  $\leq 75 \times 10^9/L$ . Approximately half of these patients had treatment discontinued with return of platelet counts to normal. In the remaining patients, platelet counts normalized with continued treatment. In this study, the probability of thrombocytopenia appeared to increase significantly at total valproate concentrations of  $\geq 110$  mcg/mL (females) or  $\geq 135$  mcg/mL (males). The therapeutic benefit which may accompany the higher doses should therefore be weighed against the possibility of a greater incidence of adverse events.

## **Hepatic/Biliary/Pancreatic**

**Serious or Fatal Hepatotoxicity:** Hepatic failure resulting in fatalities has occurred in patients receiving valproic acid and its derivatives. These incidences usually have occurred during the first 6 months of treatment with valproic acid. Caution should be observed when administering Sandoz Valproic (valproic acid) to patients with a prior history of hepatic disease. Patients on multiple anticonvulsants, children, those with congenital metabolic disorders, those with severe seizure disorders accompanied by mental retardation, and those with organic brain disease may be at particular risk.

Experience has indicated that children under the age of 2 years are at a considerably increased risk of developing fatal hepatotoxicity, especially those on multiple anticonvulsants, those with congenital metabolic disorders, those with severe seizure disorders accompanied by mental retardation, and those with organic brain disease. The risk in this age group decreased considerably in patients receiving valproic acid as monotherapy. Similarly, patients aged 3 to 10 years were at somewhat greater risk if they received multiple anticonvulsants than those who received only valproic acid. Above the age of 2 years, experience in epilepsy has indicated that the incidence of fatal hepatotoxicity decreases considerably in progressively older patients. No deaths have been reported in patients over 10 years of age who received valproic acid alone.

If Sandoz Valproic (valproic acid) is to be used in children 2 years old or younger, it should be



used with extreme caution and as a sole agent. The benefits of therapy should be weighed against the risk (see WARNINGS AND PRECAUTIONS, Special Populations, Pediatrics [ $< 18$  years of age]).

Serious or fatal hepatotoxicity may be preceded by non-specific symptoms such as loss of seizure control, malaise, weakness, lethargy, facial edema, anorexia and vomiting. Patients should be monitored closely for appearance of these symptoms. Patients and parents should be instructed to report such symptoms. Because of the non-specific nature of some of the early signs, hepatotoxicity should be suspected in patients who become unwell, other than through obvious cause, while taking Sandoz Valproic (valproic acid).

Liver function tests should be performed prior to therapy and at frequent intervals thereafter especially during the first 6 months. However, physicians should not rely totally on serum biochemistry since these tests may not be abnormal in all instances, but should also consider the results of careful interim medical history and physical examination.

In high-risk patients, it might also be useful to monitor serum fibrinogen and albumin for decreases in concentration and serum ammonia for increases in concentration. If changes occur, Sandoz Valproic (valproic acid) should be discontinued. Dosage should be titrated to and maintained at the lowest dose consistent with optimal seizure control.

The drug should be discontinued immediately in the presence of significant hepatic dysfunction, suspected or apparent. In some cases, hepatic dysfunction has progressed in spite of discontinuation of drug. The frequency of adverse effects (particularly elevated liver enzymes and thrombocytopenia) may increase with increasing dose. The therapeutic benefit which may accompany the higher doses should therefore be weighed against the possibility of a greater incidence of adverse effects (see CONTRAINDICATIONS).

**Patients with Mitochondrial Disease:** Valproate induced acute liver failure and liver-related deaths have been reported in patients with hereditary neurometabolic syndromes caused by mutations in the gene for mitochondrial DNA polymerase  $\gamma$  (POLG) (e.g., Alpers-Huttenlocher Syndrome) at a higher rate than those without these syndromes (see CONTRAINDICATIONS).

POLG-related disorders should be suspected in patients with a family history or suggestive symptoms of a POLG-related disorder, including but not limited to unexplained encephalopathy, refractory epilepsy (focal, myoclonic), status epilepticus at presentation, developmental delays, psychomotor regression, axonal sensorimotor neuropathy, myopathy cerebellar ataxia, ophthalmoplegia, or complicated migraine with occipital aura. POLG mutation testing should be performed in accordance with current clinical practice for the diagnostic evaluation of such disorders. The A467T and W748S mutations are present in approximately 2/3 of patients with autosomal recessive POLG-related disorders.

In patients over two years of age who are clinically suspected of having a hereditary mitochondrial disease, valproic acid should only be used after other anticonvulsants have failed. This older group of patients should be closely monitored during treatment with Sandoz Valproic

(valproic acid) for the development of acute liver injury with regular clinical assessments and liver function test monitoring.

In the presence of significant hepatic dysfunction, suspected or apparent, Sandoz Valproic (valproic acid) should be discontinued and alternative therapy initiated. In some cases, hepatic dysfunction has progressed in spite of discontinuation of the drug (see CONTRAINDICATIONS and WARNINGS AND PRECAUTIONS).

**Pancreatitis: Cases of life-threatening pancreatitis have been reported in both children and adults receiving valproic acid. Some of the cases have been described as hemorrhagic with a rapid progression from initial symptoms to death. Some cases have occurred shortly after initial use as well as after several years of use. The rate based upon the reported cases exceeds that expected in the general population and there have been cases in which pancreatitis recurred after rechallenge with valproic acid. In clinical trials, there were 2 cases of pancreatitis without alternative etiology in 2416 patients, representing 1044 patient-years experience. Patients and guardians should be warned that abdominal pain, nausea, vomiting, and/or anorexia can be symptoms of pancreatitis that require prompt medical evaluation. If pancreatitis is diagnosed, Sandoz Valproic (valproic acid) should ordinarily be discontinued. Alternative treatment for the underlying medical condition should be initiated as clinically indicated.**

#### **Muscle Effects /Rhabdomyolysis**

Rare cases of rhabdomyolysis, independent of neuroleptic malignant syndrome, have been reported to occur in patients treated with valproic acid. Cases have included renal failure and fatalities.

Patients should be carefully monitored for muscle pain, tenderness or weakness, particularly if accompanied by malaise or fever or tea-coloured urine. Blood creatine phosphokinase (CPK) levels should be assessed in patients experiencing these symptoms and valproic acid therapy should be discontinued if markedly elevated CPK levels are measured or if the patient develops signs and symptoms indicative of rhabdomyolysis.

Caution should be exercised in prescribing Sandoz Valproic to patients with predisposing/risk factors, including: prior history of muscular disorders such as CPT II deficiency (carnitine palmitoyltransferase type II); uncontrolled hypothyroidism; hepatic or renal impairment; concomitant medications that are known to be associated with rhabdomyolysis (eg statins, antipsychotics, diuretics, some antidepressants).

#### **Neurologic**

**Brain Atrophy:** There have been postmarketing reports of reversible and irreversible cerebral and cerebellar atrophy with neurological symptoms, in children, adults, and the elderly, receiving valproate therapy. A temporal relationship between valproate therapy and the development of cerebral atrophy and associated signs and symptoms was also demonstrated. In some cases, symptoms disappeared after valproate discontinuation but patients recovered with permanent sequelae (see ADVERSE REACTIONS). The motor and cognitive functions of patients on valproate should be routinely monitored and drug should be discontinued in the presence of suspected or apparent signs of brain atrophy.

**Neurological Problems in Children after In-utero Exposure to Valproate:** Reports of cerebral atrophy with various forms of neurological problems including developmental delays, psychomotor impairment and decreased IQ scores have been reported in children who were exposed in-utero to valproate products (see WARNINGS AND PRECAUTIONS, Special Populations, Pregnant Women).

**Driving and Hazardous Occupations:** Sandoz Valproic (valproic acid) may produce central nervous system (CNS) depression especially when combined with another CNS depressant, such as alcohol. Therefore, patients should be advised not to engage in hazardous occupations, such as driving a car or operating dangerous machinery, until it is known that they do not become drowsy from the drug.

**Aggravated convulsions:** As with other antiepileptic drugs, some patients may experience a worsening of convulsion frequency and severity, or the onset of new types of convulsions with valproate. Postmarketing reports of serious aggravated seizures have been reported for valproic acid including status epilepticus and death. In case of aggravated convulsions, patients should be advised to consult their physician immediately.

### **Psychiatric**

**Suicidal Behaviour and Ideation:** Suicidal ideation and behaviour have been reported in patients treated with antiepileptic agents in several indications.

All patients treated with antiepileptic drugs (AEDs), irrespective of indication, should be monitored for signs of suicidal ideation and behaviour and appropriate treatment should be considered. Patients (and caregivers of patients) should be advised to seek medical advice should signs of suicidal ideation or behaviour emerge.

An FDA meta-analysis of randomized placebo controlled trials, in which AEDs were used for various indications, has shown a small increased risk of suicidal ideation and behaviour in patients treated with these drugs. The mechanism of this risk is not known.

There were 43,892 patients treated in the placebo controlled clinical trials that were included in the meta-analysis. Approximately 75% of patients in these clinical trials were treated for indications other than epilepsy and, for the majority of non-epilepsy indications the treatment (AED or placebo) was administered as monotherapy. Patients with epilepsy represented approximately 25% of the total number of patients treated in the placebo controlled clinical trials and, for the majority of epilepsy patients, treatment (AED or placebo) was administered as adjunct to other antiepileptic agents (i.e., patients in both treatment arms were being treated with one or more AED). Therefore, the small increased risk of suicidal ideation and behaviour reported from the meta-analysis (0.43% for patients on AEDs compared to 0.24% for patients on placebo) is based largely on patients that received monotherapy treatment (AED or placebo) for non-epilepsy indications. The study design does not allow an estimation of the risk of suicidal ideation and behaviour for patients with epilepsy that are taking AEDs, due both to this population being the minority in the study, and the drug-placebo comparison in this population being confounded by the presence of adjunct AED treatment in both arms.

**Behavioural Disorders:** There have been postmarketing reports of behavioural disorders, including aggression, agitation, abnormal behaviour, disturbance in attention, and learning disorders. Although patients of all ages were affected, including the elderly and those exposed to valproate products *in-utero*, a large number of cases were reported in children. There was no clear trend with respect to valproate dose. In some cases, patients improved or recovered following valproate discontinuation.

### **Renal**

**Renal Impairment:** Renal impairment is associated with an increase in the unbound fraction of valproate. In several studies, the unbound fraction of valproate in plasma from renally impaired patients was approximately double that for subjects with normal renal function. Accordingly, monitoring of total concentrations in patients with renal impairment may be misleading since free concentrations may be substantially elevated whereas total concentrations may appear to be normal. Hemodialysis in renally impaired patients may remove up to 20% of the circulating valproate.

### **Sensitivity/Resistance**

**Multi-organ Hypersensitivity Reaction:** Multi-organ hypersensitivity reactions have been rarely reported in close temporal association to the initiation of valproic acid in adult and pediatric patients (median time to detection 21 days; range 1 to 40). Although there have been a limited number of reports, many of these cases resulted in hospitalization and at least one death has been reported. Signs and symptoms of this disorder were diverse; however, patients typically, although not exclusively, presented with fever and rash associated with other organ system involvement. Other associated manifestations may include lymphadenopathy, hepatitis, liver function test abnormalities, hematological abnormalities (e.g., eosinophilia, thrombocytopenia, neutropenia), pruritus, nephritis, oliguria, hepato-renal syndrome, arthralgia, and asthenia. Because the disorder is variable in its expression, other organ system symptoms and signs, not noted here may occur. If this reaction is suspected, Sandoz Valproic (valproic acid) should be discontinued and an alternative treatment started. Although the existence of cross sensitivity with other drugs that produce this syndrome is unclear, the experience amongst drugs associated with multi-organ hypersensitivity would indicate this to be a possibility.

### **Sexual Function/Reproduction**

**Fertility:** The effect of valproic acid on the development of the testis and on sperm production and fertility in humans is unknown (see TOXICOLOGY, Reproduction and Teratology, Fertility, for results in animal studies).

Amenorrhea, polycystic ovaries and increased testosterone levels have been reported in women using valproate. Valproate administration may also impair fertility in men. Case reports indicate that effects on fertility dysfunctions are reversible after treatment discontinuation.

### **Skin**

**Serious Skin Reactions:** The dose of lamotrigine should be reduced when co-administered with Sandoz Valproic (valproic acid). Serious skin reactions (such as Stevens-Johnson syndrome and Toxic Epidermal Necrolysis) have been reported with concomitant lamotrigine and valproic acid administration (see Lamotrigine Product Monograph for details on lamotrigine dosing with concomitant valproic acid administration).

## **Special Populations**

### **Female children/Female adolescents/Women of childbearing potential/Pregnancy**

**Sandoz Valproic (valproic acid) can cause fetal harm when administered to pregnant women. In comparison with some other antiepileptics drugs (AEDs), Sandoz Valproic (valproic acid) use during pregnancy is associated with an increased risk of severe birth defects such as neural tube defects (e.g., spina bifida), craniofacial defects, cleft palate, cardiovascular malformations, hypospadias, etc. (see WARNINGS AND PRECAUTIONS, Special Populations, Pregnant Women, Birth Defects).**

Valproic acid should not be used in female children, in female adolescents, in women of childbearing potential and pregnant women unless alternative treatments are ineffective or not tolerated because of its high teratogenic potential and risk of developmental disorders in infants exposed in utero to valproate. The benefit and risk should be carefully re-assessed at regular treatment reviews, at puberty and urgently when a woman of childbearing potential treated with valproic acid plans a pregnancy or becomes pregnant. This is especially important when Sandoz Valproic (valproic acid) use is considered for an indication/a condition that is not usually associated with permanent injury or death.

Since some of the congenital malformations occur in the first trimester of pregnancy before many women know that they are pregnant, all women of childbearing potential should be informed of the potential hazard to the fetus from exposure to Sandoz Valproic (valproic acid).

Women of childbearing potential must use an effective method of contraception while using Sandoz Valproic (valproic acid).

Estrogen-containing products, including estrogen-containing hormonal contraceptives, may increase the clearance of valproate, which may result in decreased serum concentration of valproate and potentially increased seizure frequency. Prescribers should monitor valproate plasma levels and clinical response (Seizure control) when initiating or discontinuing estrogen-containing products (see DRUG INTERACTIONS, Drug-Drug Interactions, Table 1)

In women planning to become pregnant, every effort should be made to switch to appropriate alternative treatment prior to conception.

### **If a Woman wants to plan a Pregnancy**

- During pregnancy, maternal tonic-clonic seizures and status epilepticus with hypoxia may carry a particular risk of death for the mother and the unborn child.
- In women planning to become pregnant or who are pregnant, valproate therapy should be reassessed.

Valproate therapy should not be discontinued without a reassessment of the benefits and risks of treatment by a physician who is experienced in the management of epilepsy. Antiepileptic drugs should not be abruptly discontinued in patients to whom the drug is administered to prevent

major seizures because of the strong possibility of precipitating status epilepticus with attendant hypoxia and risks to both the mother and the unborn child.

If based on a careful evaluation of the risks and benefits, valproate treatment is continued during pregnancy, the following are recommended:

- Use the lowest effective dose and divide the daily valproate dose into several small doses to be taken throughout the day.
- Folate supplementation before pregnancy may decrease the risk of neural tube defects common to all pregnancies. However, available evidence does not suggest that folate prevents birth defects or malformations due to valproate exposure.
- Institute specialized prenatal monitoring in order to detect the possible occurrence of neural tube defects or other malformations.

With regard to drugs given for minor seizures, the risks of discontinuing medication prior to or during pregnancy should be weighed against the risk of congenital defects in each particular case and with the particular family history. In individual cases where the severity and frequency of the seizure disorder are such that the removal of medication does not pose a serious threat to the patient, discontinuation of the drug may be considered prior to and/or during pregnancy, and alternative therapy considered, although it cannot be said with any confidence that even minor seizures do not pose some hazard to the developing embryo or fetus.

Women of childbearing age should be encouraged to seek the counsel of their physician and should report a positive pregnancy test promptly. Where the necessity for continued use of antiepileptic medication is in doubt, appropriate consultation is indicated. Current best practice guidelines should be considered in order to provide the optimal counsel to patients regarding the teratogenic risks associated with Sandoz Valproic (valproic acid).

### **Pregnancy Exposure Risk Related to Valproate**

Both valproate monotherapy and polytherapy are associated with abnormal pregnancy outcomes. Available data suggest that antiepileptic polytherapy including valproate is associated with a greater risk of congenital malformations than valproate monotherapy.

Tests to detect neural tube and other defects using current accepted procedures should be considered as part of routine prenatal care in pregnant women receiving Sandoz Valproic (valproic acid).

### Pregnancy Registry

Pregnant patients taking Sandoz Valproic (valproic acid) should be encouraged to enroll in the North American Antiepileptic Drug (NAAED) Pregnancy Registry. This can be done by calling the toll free number 1-888-233-2334, and must be done by patients themselves. Information on the registry can also be found at the following website: <http://www.aedpregnancyregistry.org/>.

### Risk in the neonate

Cases of hemorrhagic syndrome have been reported very rarely in neonates whose mothers have taken valproate during pregnancy. This hemorrhagic syndrome is related to thrombocytopenia,

hypofibrinogenemia and/or to a decrease in other coagulation factors.

Afibrinogenemia has also been reported and can be fatal. However, this syndrome must be distinguished from the decrease of vitamin-K factors induced by phenobarbital and other enzymes. Therefore, in neonates, platelet count, plasma levels of fibrinogen, coagulation tests, and coagulation factors should be investigated.

Cases of hypoglycemia have been reported in neonates whose mothers have taken valproate during the third trimester of pregnancy.

Cases of hypothyroidism have been reported in neonates whose mothers have taken valproate during pregnancy.

Withdrawal syndrome (symptoms include: agitation, irritability, hyperexcitability, jitteriness, hyperkinesia, tonic disorders, tremor, convulsions and feeding disorders) may occur, in the days following birth, in neonates whose mothers have taken valproate during the last trimester of pregnancy.

### Birth Defects

There are multiple reports in the clinical literature that indicate the use of antiepileptic drugs during pregnancy results in an increased incidence of birth defects in the offspring. The incidence of congenital malformations in the general population is regarded to be approximately 2%; in children of treated epileptic women, this incidence may be increased 2- to 3-fold. The increase is largely due to specific defects such as congenital malformations of the heart, cleft lip and/or palate, craniofacial abnormalities and neural tube defects. Nevertheless, the great majority of mothers receiving antiepileptic medications deliver normal infants.

The data described below were gained almost exclusively from women who received valproate to treat epilepsy. Data from Pregnancy Registries indicate an increased risk of congenital anomalies in infants exposed to valproic acid monotherapy during the first trimester of pregnancy as compared to other antiepileptic drugs. Based on Pregnancy Registry data and the United States Centers for Disease Control (CDC), the estimated risk of valproate-exposed women having children with spina bifida, oral clefts, neural tube defects, and hypospadias is approximately 1 to 2% as compared to the risk of spina bifida in the general population which is about 0.06 to 0.07%.

In a study using NAAED Pregnancy Registry data, 16 cases of major malformations following prenatal valproate exposure were reported among offspring of 149 enrolled women who used valproate during pregnancy. Three of the 16 cases were neural tube defects; the remaining cases included craniofacial defects, cardiovascular malformations and malformations of varying severity involving other body systems. The NAAED Pregnancy Registry has reported a major malformation rate of 10.7% in the offspring of women exposed to valproate monotherapy during pregnancy (average daily dose: 1000 mg; range: 500 – 2000 mg/day) as compared to major malformation rate of 2.9% among 1048 epileptic women who received any other antiepileptic drug monotherapy during pregnancy. These data show a four-fold increased risk for any major malformation following valproate exposure *in utero* compared to the risk following exposure *in utero* to any other antiepileptic drug monotherapy.

Data derived from a meta-analysis (including registries and cohort studies) has shown that 10.73% of children of epileptic women exposed to valproate monotherapy during pregnancy suffer from congenital malformations (95% CI: 8.16-13.29). This is a greater risk of major malformations than for the general population, for whom the risk is about 2-3%. The risk is dose-dependent but a threshold dose below which no risk exists cannot be established.

Available data show an increased incidence of minor and major malformations. The most common types of malformations include neural tube defects, facial dysmorphism, cleft lip and palate, craniostenosis, cardiac, renal and urogenital defects, limb defects (including bilateral aplasia of the radius), and multiple anomalies involving various body systems.

#### Risk of Neurological Problems from In-Utero Exposure

**Cerebral Atrophy:** Exposure *in-utero* to valproate products has been associated with cerebral atrophy with varying degrees/manifestations of neurological compromise, including developmental delays and psychomotor impairment (see ADVERSE REACTIONS and WARNINGS AND PRECAUTIONS, Neurologic, Brain Atrophy).

**Developmental Delay, Decreased IQ, Autism and/or Autism Spectrum Disorders:** Available data to date show that exposure to valproate in utero can have adverse effects on mental and physical development of the exposed children. The risk seems to be dose-dependent but a threshold dose below which no risk exists, cannot be established. The exact gestational period for risk of these effects is uncertain, and it is possible that the risk exists throughout the entire pregnancy.

Studies in preschool children exposed in utero to valproate show that up to 30-40% experience delays in their early development such as talking and walking later, lower intellectual abilities, poor language skills (speaking and understanding) and memory problems.

Intelligence quotient (IQ) measured in school aged children (age 6) with a history of valproate exposure in utero was on average 7-10 points lower than those children exposed to other antiepileptics. There is evidence in children exposed to valproate that the risk of intellectual impairment may be independent of maternal IQ.

There are limited data on long term outcomes. Available data show that children exposed to valproate in utero are at increased risk of autistic spectrum disorder (approximately three-fold) and childhood autism (approximately five-fold) compared to the general study population.

Limited data suggests that children exposed to valproate in utero may be more likely to develop symptoms of Attention Deficit/Hyperactivity Disorder (ADHD).

#### Coagulation Abnormalities

There have been reported postmarketing cases of coagulation abnormalities in patients of all ages receiving valproate therapy. These include thrombocytopenia, hypofibrinogenemia, and/or decrease in other coagulation factors, which can lead to bleeding and other complications, especially in the cases of decrease in factors VII, VIII and XIII. These abnormalities may not necessarily be dose-dependent. Some of the hemorrhage manifestations may include mucosal bleeding (e.g., menorrhagia, epistaxis, hematuria, melena), easy-bruising, soft-tissue hematoma,



hemarthrosis, and intracranial hemorrhage. Caution should be taken in patients taking valproate and anticoagulants and in cases of injury or surgery to avoid life-threatening or fatal bleeding (see WARNINGS AND PRECAUTIONS, Hematologic, Thrombocytopenia).

Pregnant women taking Sandoz Valproic (valproic acid) may also develop coagulation abnormalities, which may result in hemorrhagic complications in the neonate including death (see WARNINGS AND PRECAUTIONS, Hematologic, Thrombocytopenia). If Sandoz Valproic (valproic acid) is used in pregnancy, the coagulation parameters should be monitored carefully.

#### Hepatic Failure

Hepatic failure, resulting in the death of a newborn and of an infant has been reported following the use of valproate during pregnancy.

#### Hypoglycemia

Serious postmarketing reports of hypoglycemia have been received for neonates whose mothers received valproic acid treatment during pregnancy. In most cases, valproic acid was the only reported antiepileptic drug (AED). Most of these neonates also displayed other congenital anomalies including hypospadias, complex facial dysmorphism, limb anomalies, severe cardiac anomalies, etc. Therefore, if a decision has been made to use valproic acid during pregnancy, or if the patient becomes pregnant while taking this drug, the patient should be made aware of the potential hazard to the fetus.

#### Thyroid Gland Abnormalities

Cases of hypothyroidism have been reported in neonates whose mothers have taken valproate during pregnancy. There have also been reported cases of increased serum thyroid stimulating hormone or decreased serum thyroxine levels in children receiving valproate therapy. In addition, there have been reported cases of hypothyroidism and hyperthyroidism in adults and children receiving valproate monotherapy.

#### Teratogenicity in Animals

Animal studies have demonstrated valproic acid induced teratogenicity (see TOXICOLOGY, Reproduction and Teratology), and studies in human females have demonstrated placental transfer of the drug. Increased frequencies of malformations, as well as intrauterine growth retardation and death, have been observed in mice, rats, rabbits, and monkeys following prenatal exposure to valproate. Malformations of the skeletal system are the most common structural abnormalities produced in experimental animals, but neural tube closure defects have been seen in mice exposed to maternal plasma valproate concentrations exceeding 230 mcg/mL (2.3 times the upper limit of the human therapeutic range for epilepsy) during susceptible periods of embryonic development.

Administration of an oral dose of 200 mg/kg/day or greater (50% of the maximum human daily dose or greater on a mg/m<sup>2</sup> basis) to pregnant rats during organogenesis produced malformations (skeletal, cardiac and urogenital) and growth retardation in the offspring. These doses resulted in peak maternal plasma valproate levels of approximately 340 mcg/mL or greater (3.4 times the upper limit of the human therapeutic range for epilepsy or greater). Behavioural deficits have been reported in the offspring of rats given a dose of 200 mg/kg/day throughout most of pregnancy.

An oral dose of 350 mg/kg/day (approximately 2 times the maximum human daily dose on a mg/m<sup>2</sup> basis) produced skeletal and visceral malformations in rabbits exposed during organogenesis. Skeletal malformations, growth retardation, and death were observed in rhesus monkeys following administration of an oral dose of 200 mg/kg/day (equal to the maximum human daily dose on a mg/m<sup>2</sup> basis) during organogenesis. This dose resulted in peak maternal plasma valproate levels of approximately 280 mcg/mL (2.8 times the upper limit of the human therapeutic range for epilepsy).

**Nursing Women:** Valproic acid is secreted in breast milk. Concentrations in breast milk have been reported to be 1 to 10% of maternal serum concentrations. As a general rule, nursing should not be undertaken while a patient is receiving valproic acid. Based on literature and clinical experience, hematological disorders have been shown in breastfed newborns/infants of treated women.

**Pediatrics (< 18 years of age):** Experience has indicated that children under the age of 2 years are at a considerably increased risk of developing fatal hepatotoxicity, especially those with the aforementioned conditions (see WARNINGS AND PRECAUTIONS, Hepatic/Biliary/Pancreatic, Serious or Fatal Hepatotoxicity). When Sandoz Valproic (valproic acid) is used in this patient group, it should be used with extreme caution and as a sole agent. The benefits of therapy should be weighed against the risks (see CONTRAINDICATIONS).

Above the age of 2 years, experience in epilepsy has indicated that the incidence of fatal hepatotoxicity decreases considerably in progressively older patient groups.

In patients over two years of age who are clinically suspected of having a hereditary mitochondrial disease, Sandoz Valproic (valproic acid) should only be used after other anticonvulsants have failed (see WARNINGS AND PRECAUTIONS, Hepatic/Biliary/Pancreatic, Patients with Mitochondrial Disease).

Younger children, especially those receiving enzyme-inducing drugs, will require larger maintenance doses to attain targeted total and unbound valproate concentrations. The variability in free fraction limits the clinical usefulness of monitoring total serum valproate concentrations. Interpretation of valproate concentrations in children should include consideration of factors that affect hepatic metabolism and protein binding.

**Geriatrics (≥ 65 years of age):** Alterations in the kinetics of unbound valproate in the elderly indicate that the initial dosage should be reduced in this population (see DOSAGE AND ADMINISTRATION and ACTION AND CLINICAL PHARMACOLOGY, Special Populations and Conditions, Geriatrics).

The safety and efficacy of valproic acid in elderly patients with epilepsy has not been evaluated in clinical trials. Caution should thus be exercised in dose selection for an elderly patient, recognizing the more frequent hepatic and renal dysfunctions, and limited experience with valproic acid in this population.

A study of elderly patients revealed valproate-related somnolence and discontinuation of valproic acid for this adverse event (see WARNINGS AND PRECAUTIONS, Special Populations,

Geriatrics, Somnolence in the Elderly). The starting dose should be reduced in elderly patients, and dosage reductions or discontinuation should be considered in patients with excessive somnolence (see DOSAGE AND ADMINISTRATION).

#### Somnolence in the elderly

In a group of elderly patients (mean age 83 years old, n = 172), valproic acid doses were increased by 125 mg/day to a target dose of 20 mg/kg/day. Compared to placebo a significantly higher number of valproate-treated patients had somnolence, and although not statistically significant, a higher number of valproate-treated patients experienced dehydration. Discontinuations for somnolence were also significantly higher in valproate-treated patients compared to placebo. In approximately one-half of the patients with somnolence, there was also associated reduced nutritional intake and weight loss. In elderly patients, dosage should be increased more slowly and with regular monitoring for fluid intake, dehydration, somnolence, urinary tract infection and other adverse events. Dose reductions or discontinuation of Sandoz Valproic (valproic acid) should be considered in patients with decreased food or fluid intake and in patients with excessive somnolence (see DOSAGE AND ADMINISTRATION).

#### Monitoring and Laboratory Tests

Since Sandoz Valproic (valproic acid) may interact with concurrently administered drugs which are capable of enzyme induction, periodic plasma concentration determinations of valproate and concomitant drugs are recommended during the early course of therapy and whenever enzyme-inducing drugs are introduced or withdrawn (see DRUG INTERACTIONS).

#### Monitoring Valproate Concentrations

Protein binding of valproate is reduced in the elderly, in patients with renal impairment, and in the presence of other drugs (e.g., acetylsalicylic acid). Accordingly, measurements of plasma levels of valproate may be misleading in these patients, as actual drug exposure may be higher than measured values (see WARNINGS AND PRECAUTIONS, Hepatic/Biliary/Pancreatic, WARNINGS AND PRECAUTIONS, Endocrine and Metabolism, Hyperammonemia, WARNINGS AND PRECAUTIONS, Hematologic, Thrombocytopenia and DRUG INTERACTIONS, Drug-Drug Interactions, Table 1).

### **ADVERSE REACTIONS**

#### Adverse Drug Reaction Overview

The most commonly reported adverse reactions are nausea, vomiting and indigestion. Since valproic acid has usually been used with other antiepileptics, it is not possible in most cases to determine whether the adverse reactions mentioned in this section are due to valproic acid alone or to the combination of drugs.

Adverse events that have been reported with valproic acid from epilepsy trials, spontaneous reports, and other sources are listed below by system organ class.

Blood and Lymphatic  
System Disorders:

Thrombocytopenia and inhibition of the secondary phase of platelet aggregation may be reflected in altered bleeding time,

petechiae, bruising, hematoma formation, epistaxis, and hemorrhage (see WARNINGS AND PRECAUTIONS, Hematologic, Thrombocytopenia). Relative lymphocytosis, macrocytosis and hypofibrinogenemia have been noted. Leukopenia and eosinophilia have also been reported. Anemia, including macrocytic with or without folate deficiency, aplastic anemia, pancytopenia, bone marrow suppression, agranulocytosis and acute intermittent porphyria have been reported.

- Cardiac Disorders: Bradycardia
- Ear and Labyrinth Disorders: Hearing loss, either reversible or irreversible, has been reported; however, a cause and effect relationship has not been established. Ear pain has also been reported.
- Gastrointestinal Disorders: Nausea, vomiting and indigestion are the most commonly reported side effects at the initiation of therapy. These effects are usually transient and rarely require discontinuation of therapy. Diarrhea, abdominal cramps, constipation and gingival disorder (mainly gingival hyperplasia) have also been reported. There have been reports of acute pancreatitis, including rare fatal cases, occurring in association with valproic acid therapy (see WARNINGS AND PRECAUTIONS, Hepatic/Biliary/Pancreatic, Pancreatitis).
- Parotid gland swelling has also been reported in patients receiving valproic acid.
- General Disorders and Administration Site Conditions: Edema of the extremities, fever and hypothermia
- Hepatobiliary Disorders: Minor elevations of transaminases (e.g., serum glutamic-oxaloacetic transaminase (SGOT) and serum glutamic-pyruvic transaminase (SGPT)) and lactate dehydrogenase (LDH) are frequent and appear to be dose-related. Occasionally, laboratory tests also show increases in serum bilirubin and abnormal changes in other liver function tests. These results may reflect potentially serious hepatotoxicity (see WARNINGS AND PRECAUTIONS, Hepatic/Biliary/Pancreatic, Serious or Fatal Hepatotoxicity).
- Immune System Disorder: Allergic reaction, anaphylaxis
- Infections and Infestations: Pneumonia and otitis media

Investigations: Abnormal thyroid function tests (including both hyperthyroidism and hypothyroidism) (see WARNINGS AND PRECAUTIONS, Special Populations, Pregnant Women, Thyroid Gland Abnormalities and DRUG INTERACTIONS, Drug-Laboratory Interactions).

#### Metabolism and Nutrition

Disorders: Hyperammonemia (see WARNINGS AND PRECAUTIONS, Endocrine and Metabolism, Hyperammonemia), hyponatremia, biotin deficiency/biotinidase deficiency and inappropriate antidiuretic hormone (ADH) secretion. There have been rare reports of Fanconi syndrome (proximal renal tubular dysfunction) occurring primarily in children. Decreased carnitine concentrations have been reported although the clinical relevance is undetermined. Hyperglycinemia has been reported and associated with a fatal outcome in patient with pre-existing nonketotic hyperglycinemia.

Anorexia with some weight loss and increased appetite with some weight gain have also been reported.

Obesity has been reported in post-marketing experience.

#### Musculoskeletal and Connective Tissue Disorders:

Weakness, rhabdomyolysis and bone pain have been reported (see WARNINGS AND PRECAUTIONS, Muscle Effects /Rhabdomyolysis).

Reports have been received of decreased bone mass, potentially leading to osteoporosis and osteopenia, during long-term therapy with some anticonvulsant medications, including valproic acid. Some studies have indicated that supplemental calcium and vitamin D may be of benefit to patients who are on chronic valproic acid therapy.

A lupus erythematosus-like syndrome has been reported rarely.

#### Neoplasms Benign, Malignant and Unspecified (including cysts and polyps):

Myelodysplastic syndrome in both adults and children (all children were on valproate monotherapy). In some cases in adults and/or children, myelodysplastic syndrome was reversible upon valproate discontinuation.

#### Nervous System Disorders:

Sedative effects have been noted in patients receiving valproic acid alone but occur most often in patients on combination

therapy. Sedation usually disappears upon reduction of other anti-epileptic medication.

Hallucination, ataxia, headache, nystagmus, diplopia, asterixis, "spots before the eyes", tremor (may be dose-related), confusion, dysarthria, dizziness, hypesthesia, vertigo, incoordination, memory impairment, cognitive disorder, and extrapyramidal disorders including parkinsonism have been reported with the use of valproate. Rare cases of coma have been reported in patients receiving valproic acid alone or in conjunction with phenobarbital.

Encephalopathy, with or without fever or hyperammonemia, has been reported without evidence of hepatic dysfunction or inappropriate valproate plasma levels. Most patients recovered, with noted improvement of symptoms, upon discontinuation of the drug.

There have been postmarketing reports of reversible and irreversible cerebral and cerebellar atrophy temporally associated with the use of valproate products. In some cases the patients recovered with permanent sequelae (see WARNINGS AND PRECAUTIONS, Neurologic, Brain Atrophy). Cerebral atrophy seen in children exposed to valproate *in utero* led to various forms of neurological events, including developmental delays and psychomotor impairment (see WARNINGS AND PRECAUTIONS, Special Populations, Pregnant Women).

Congenital malformations and developmental disorders have also been reported.

Aggravated convulsions (increase in number of seizures or appearance of new seizure type or worsening of seizures) have been reported in patients with epilepsy treated with valproate monotherapy.

Psychiatric Disorders:

Emotional upset, depression, psychosis, aggression, psychomotor hyperactivity, hostility, agitation, disturbance in attention, abnormal behaviour, learning disorder and behavioural deterioration.

(see WARNINGS AND PRECAUTIONS, Psychiatric).

Renal and Urinary Disorders:

Enuresis, urinary incontinence, acute renal failure, tubulointerstitial nephritis and urinary tract infection.

Reproductive System and Breast Disorders:

There have been reports of irregular menses, secondary amenorrhea, breast enlargement and galactorrhea in patients receiving valproic acid. Hyperandrogenism (hirsutism, virilism, acne, male pattern alopecia, and/or androgen increased). There have been rare spontaneous reports of polycystic ovary disease. A cause and effect relationship has not been established.

Respiratory, Thoracic and Mediastinal Disorders:

Increased cough, pleural effusion.

Skin and Subcutaneous Tissue Disorders:

Transient and/or dose related alopecia (hair loss), hair disorders (such as hair texture abnormal, hair colour changes, hair growth abnormal), have been observed. Skin rash, photosensitivity, generalized pruritus, erythema multiforme, Stevens-Johnson syndrome, and petechiae have rarely been noted.

Rare cases of Toxic Epidermal Necrolysis (TEN) have been reported including a fatal case of a 6 month old infant taking valproic acid and several other concomitant medications. An additional case of Toxic Epidermal Necrolysis resulting in death was reported in a 35 year old patient with AIDS taking several concomitant medications and with a history of multiple cutaneous drug reactions.

Serious skin reactions have been reported with concomitant administration of lamotrigine and valproic acid (see DRUG INTERACTIONS, Drug-Drug Interactions, Table 1).

Cutaneous vasculitis has also been reported.

Nail and nail bed disorders have also been reported in post-marketing experience.

**Adverse Events in Elderly Patients**

In elderly patients (above 65 years of age), there were more frequent reports of accidental injury, infection, pain, and to a lesser degree, somnolence and tremor, when compared to patients 18 to 65 years of age. Somnolence and tremor tended to be associated with the discontinuation of valproic acid.

**DRUG INTERACTIONS**

**Serious Drug Interactions**

- Rare cases of coma have been reported in patients receiving valproic acid alone or in conjunction with phenobarbital (see Drug-Drug Interactions, Table 1).
- Serious skin reactions (such as Stevens-Johnson syndrome and Toxic Epidermal Necrolysis) have been reported with concomitant lamotrigine and valproic acid administration (see Drug-Drug Interactions, Table 1).

### **Overview**

Valproic acid has been found to be a weak inhibitor of some P<sub>450</sub> isozymes, epoxide hydase, and glucuronyl transferases.

Drugs that affect the level of expression of hepatic enzymes, particularly those that elevate levels of glucuronyl transferases (such as ritonavir; see Table 1 below), may increase the clearance of valproate. For example, phenytoin, carbamazepine, and phenobarbital (or primidone) can double the clearance of valproate. Thus, patients on valproic acid monotherapy will generally have longer half-lives and higher concentrations than patients receiving polytherapy with antiepilepsy drugs.

In contrast, drugs that are inhibitors of cytochrome P<sub>450</sub> isozymes, such as antidepressants, may be expected to have little effect on valproate clearance because cytochrome P<sub>450</sub> microsomal mediated oxidation is a relatively minor secondary metabolic pathway compared to glucuronidation and beta-oxidation.

The concomitant administration of valproic acid with drugs that exhibit extensive protein binding (e.g., acetylsalicylic acid, carbamazepine, dicumarol, warfarin, tolbutamide, and phenytoin) may result in alteration of serum drug levels.

Since valproic acid may interact with concurrently administered drugs which are capable of enzyme induction, periodic plasma concentration determinations of valproate and concomitant drugs are recommended during the early course of therapy and whenever enzyme-inducing drugs are introduced or withdrawn.

### **Drug-Drug Interactions**

Table 1 provides information about the potential influence of several commonly prescribed medications on valproic acid pharmacokinetics as well as the potential influence of valproic acid on the pharmacokinetics and pharmacodynamics of several commonly prescribed medications. The list is not exhaustive nor could it be, since new interactions are continuously being reported. Please note that drugs may be listed under specific name, family or pharmacologic class. Reading the entire section is recommended.



<b>Table 1: Summary of Drug-Drug Interaction Studies Including Important Interactions, Non-clinically Important Interactions and No Observed Interactions</b>			
<b>Concomitant Drug</b>	<b>Ref</b>	<b>Effect</b>	<b>Clinical Comment</b>
Acetaminophen	CT	↔ acetaminophen	Valproic acid had no effect on any of the pharmacokinetic parameters of acetaminophen when it was concurrently administered to three epileptic patients.
Acetazolamide			Concomitant administration of valproate and acetazolamide has been associated with encephalopathy and/or hyperammonemia. Patients treated with those two drugs should be carefully monitored for signs and symptoms of hyperammonemic encephalopathy (see WARNINGS AND PRECAUTIONS, Endocrine and Metabolism).
Acetylsalicylic Acid	CT	↑ valproate	A study involving the co-administration of acetylsalicylic acid at antipyretic doses (11 to 16 mg/kg) with valproic acid to pediatric patients (n = 6) revealed a decrease in protein binding and an inhibition of metabolism of valproate. Valproate free fraction was increased 4-fold in the presence of acetylsalicylic acid compared to valproic acid alone. The beta-oxidation pathway consisting of 2-E-valproic acid, 3-OH-valproic acid, and 3-keto valproic acid was decreased from 25% of total metabolites excreted on valproic acid alone to 8.3% in the presence of acetylsalicylic acid. Caution should be observed when valproic acid is administered with drugs affecting coagulation, [e.g., acetylsalicylic acid and warfarin] (see ADVERSE REACTIONS).
Alcohol	T	No pharmacokinetic (PK) interaction	Valproic acid may potentiate the CNS depressant action of alcohol.
Amitriptyline / Nortriptyline	CT	In general: ↓ amitriptyline ↓ nortriptyline  Rarely: ↑ amitriptyline ↑ nortriptyline	Administration of a single oral 50 mg dose of amitriptyline to 15 normal volunteers (10 males and 5 females) who received valproic acid (500 mg twice daily) resulted in a 21% decrease in plasma clearance of amitriptyline and a 34% decrease in the net clearance of nortriptyline.  Rare post-marketing reports of concurrent use of valproic acid and amitriptyline resulting in an increased amitriptyline and nortriptyline levels have been received. Concurrent use of valproic acid and amitriptyline has rarely been associated with toxicity. Monitoring of amitriptyline levels should be considered for patients taking valproic acid concomitantly with amitriptyline. Consideration should be given to lowering the dose of amitriptyline/nortriptyline in the presence of valproic acid.
Antacids	CT	↔ valproate	A study involving the co-administration of valproic acid 500 mg with commonly administered antacids (Maalox <sup>®</sup> , Trisogel, and Titalac <sup>™</sup> - 160 milliequivalent doses) did not reveal any effect on the extent of absorption of valproic acid.
Other - Antipsychotics, Monoamine Oxidase Inhibitors (MAOIs) and Tricyclic Antidepressants			In addition to enhancing CNS depression when used concurrently with valproic acid, antipsychotics, tricyclic antidepressants and MAOIs may lower the seizure threshold. Dosage adjustments may be necessary to control seizures.

<b>Table 1: Summary of Drug-Drug Interaction Studies Including Important Interactions, Non-clinically Important Interactions and No Observed Interactions</b>			
<b>Concomitant Drug</b>	<b>Ref</b>	<b>Effect</b>	<b>Clinical Comment</b>
Antiretroviral agents - Ritonavir - Lopinavir - Zidovudine - Lamivudine	C CT	↓ valproate ↑ zidovudine	Protease inhibitors such as lopinavir and ritonavir decrease valproate plasma level when co-administered with valproate.  Reduction of therapeutic effect of valproate was observed in a patient with bipolar disorder with the initiation of HIV treatment with lopinavir/ritonavir, zidovudine, and lamivudine.  In 6 patients who were seropositive for HIV, the clearance of zidovudine (100 mg every 8 hours) was decreased by 38% after administration of valproate (250 or 500 mg every 8 hours); the half-life of zidovudine was unaffected.
Benzodiazepines			Valproic acid may decrease oxidative liver metabolism of some benzodiazepines, resulting in increased serum concentrations (see Table 1, Diazepam and Lorazepam).
Carbamazepine / carbamazepine-10,11-epoxide	CT	↓ CBZ ↑ CBZ-E  ↓ valproate	Concomitant use of carbamazepine (CBZ) with valproic acid may result in decreased serum concentrations and half-life of valproate due to increased metabolism induced by hepatic microsomal enzyme activity. Monitoring of serum concentrations is recommended when either medication is added to or withdrawn from an existing regimen. Changes in the serum concentration of the 10,11-epoxide (CBZ-E) metabolite of carbamazepine, however, will not be detected by routine serum carbamazepine assay.  Serum levels of carbamazepine decreased 17% while that of carbamazepine-10,11-epoxide increased by 45% upon co-administration of valproic acid and CBZ to epileptic patients.
Carbapenem Antibiotics		↓ valproate	Carbapenem antibiotics (ertapenem, imipenem, meropenem, doripenem) can reduce serum valproic acid concentrations to sub-therapeutic levels. This can result in loss of seizure control in epileptic patients or loss of efficacy in nonepileptics. In some cases of co-administration in epileptic patients, breakthrough seizures have occurred. Increasing valproic acid dose may not be sufficient to overcome this interaction. If co-administration is essential, serum valproic acid concentrations should be monitored daily. Alternative antibacterial or anticonvulsant therapy should be considered if serum valproic acid concentrations drop significantly or seizure control deteriorates (see WARNINGS AND PRECAUTIONS, General, Interaction with Carbapenem Antibiotics).
Chlorpromazine	CT	↑ valproate	A study involving the administration of 100 to 300 mg/day of chlorpromazine to schizophrenic patients already receiving valproic acid (200 mg twice daily) revealed a 15% increase in trough plasma levels of valproate. This increase is not considered clinically important.
Cholestyramine		↓ valproate	Cholestyramine may lead to a decrease in plasma levels of valproate when co-administered.

<b>Table 1: Summary of Drug-Drug Interaction Studies Including Important Interactions, Non-clinically Important Interactions and No Observed Interactions</b>			
<b>Concomitant Drug</b>	<b>Ref</b>	<b>Effect</b>	<b>Clinical Comment</b>
Cimetidine	T	↑ valproate	Cimetidine may decrease the clearance and increase the half-life of valproic acid by altering its metabolism. In patients receiving valproic acid, serum valproic acid levels should be monitored when treatment with cimetidine is instituted, increased, decreased, or discontinued. The valproic acid dose should be adjusted accordingly.
Clonazepam	T	No PK interaction	The concomitant use of valproic acid and clonazepam may induce absence status in patients with a history of absence type seizures.
Clozapine	CT	No interaction	In psychotic patients (n = 11), no interaction was observed when valproic acid was co-administered with clozapine.
Diazepam	CT	↑ diazepam	Valproate displaces diazepam from its plasma albumin binding sites and inhibits its metabolism. Co-administration of valproic acid (1500 mg daily) increased the free fraction of diazepam (10 mg) by 90% in healthy volunteers (n = 6). Plasma clearance and volume of distribution for free diazepam were reduced by 25% and 20%, respectively, in the presence of valproate. The elimination half-life of diazepam remained unchanged upon addition of valproate.
Estrogen-Containing Products	C CT T	↓ valproate	Estrogen-containing products, including estrogen-containing hormonal contraceptives, may increase the clearance of valproate, which may result in decreased serum concentration of valproate and potentially increased seizure frequency. Prescribers should monitor valproate plasma levels and clinical response (seizure control) when adding or discontinuing estrogen-containing products
Ethosuximide	CT	↑ ethosuximide	Valproate inhibits the metabolism of ethosuximide. Administration of a single ethosuximide dose of 500 mg with valproic acid (800 to 1600 mg/day) to healthy volunteers (n = 6) was accompanied by a 25% increase in elimination half-life of ethosuximide and a 15% decrease in its total clearance as compared to ethosuximide alone. Patients receiving valproic acid and ethosuximide, especially along with other anticonvulsants, should be monitored for alterations in serum concentrations of both drugs.
Felbamate	CT	↑ valproate	A study involving the co-administration of 1200 mg/day of felbamate with valproic acid to patients with epilepsy (n = 10) revealed an increase in mean valproate peak concentration by 35% (from 86 to 115 mcg/mL) compared to valproic acid alone. Increasing the felbamate dose to 2400 mg/day increased the mean valproate peak concentration to 133 mcg/mL (another 16% increase). A decrease in valproic acid dosage may be necessary when felbamate therapy is initiated. Lower doses of valproic acid may be necessary when used concomitantly with felbamate.
Haloperidol	CT	↔ valproate	A study involving the administration of 6 to 10 mg/day of haloperidol to schizophrenic patients already receiving valproic acid (200 mg twice daily) revealed no significant changes in valproate trough plasma levels.
Lamotrigine	CT	↑ lamotrigine ↓ valproate	The effects of valproic acid on lamotrigine were investigated in 6 healthy male subjects. Each subject received a single oral dose of lamotrigine alone and with valproic acid 200 mg every

**Table 1: Summary of Drug-Drug Interaction Studies Including Important Interactions, Non-clinically Important Interactions and No Observed Interactions**

Concomitant Drug	Ref	Effect	Clinical Comment
			<p>8 hours for 6 doses starting 1 hour before the lamotrigine dose was given. Valproic acid administration reduced the total clearance of lamotrigine by 21% and increased the plasma elimination half-life from 37.4 hours to 48.3 hours (<math>p &lt; 0.005</math>). Renal clearance of lamotrigine was unchanged. In a steady-state study involving 10 healthy volunteers, the elimination half-life of lamotrigine increased from 26 to 70 hours with valproic acid co-administration (a 165% increase).</p> <p>In a study involving 16 epileptic patients, valproic acid doubled the elimination half-life of lamotrigine. In an open-labelled study, patients receiving enzyme inducing AEDs (e.g., carbamazepine, phenytoin, phenobarbital, or primidone) demonstrated a mean lamotrigine plasma elimination half-life of 14 hours while the elimination half-life was 30 hours in patients taking valproic acid plus an enzyme inducing antiepileptic agent. The latter value is similar to the lamotrigine half-life during monotherapy indicating that valproic acid may counteract the effect of the enzyme inducer. If valproic acid is discontinued in a patient receiving lamotrigine and an enzyme inducing antiepileptic serum lamotrigine concentrations may decrease. Patients receiving combined antiepileptic therapy require careful monitoring when another agent is started, stopped or when the dose is altered.</p> <p>Serious skin reactions (such as Stevens-Johnson syndrome and Toxic Epidermal Necrolysis) have been reported with concomitant lamotrigine and valproic acid administration.</p>
Lithium	CT	↔ lithium	<p>In a double-blind placebo-controlled multiple dose crossover study in 16 healthy male volunteers, pharmacokinetic parameters of lithium were not altered by the presence or absence of valproate. The presence of lithium, however, resulted in an 11 to 12% increase in the AUC and <math>C_{max}</math> of valproate. <math>T_{max}</math> was also reduced. Although these changes were statistically significant, they are not likely to have clinical importance.</p> <p>Co-administration of valproic acid (500 mg twice daily) and lithium carbonate (300 mg three times daily) to normal male volunteers (<math>n = 16</math>) had no effect on the steady-state kinetics of lithium.</p>
Lorazepam	CT	↑ lorazepam	<p>Concomitant administration of valproic acid (500 mg twice daily) and lorazepam (1 mg twice daily) in normal male volunteers (<math>n = 9</math>) was accompanied by a 17% decrease in the plasma clearance of lorazepam. This decrease is not considered clinically important.</p>
Nimodipine	CT	↑ nimodipine	<p>Concomitant treatment of nimodipine with valproic acid may increase nimodipine plasma concentration by 50%.</p>

<b>Table 1: Summary of Drug-Drug Interaction Studies Including Important Interactions, Non-clinically Important Interactions and No Observed Interactions</b>			
<b>Concomitant Drug</b>	<b>Ref</b>	<b>Effect</b>	<b>Clinical Comment</b>
Olanzapine	CT	↓ olanzapine	Valproic acid may decrease the olanzapine plasma concentration. Administration of a single 5 mg dose of olanzapine to 10 healthy, non-epileptic volunteers with divalproex sodium extended-release tablets 1000 mg QD did not affect olanzapine C <sub>max</sub> and elimination half-life. However, olanzapine AUC was 35% lower in the presence of divalproex sodium extended-release tablets. The clinical significance of these observations is unknown.
Oral Contraceptive Steroids	CT	No PK interaction	Evidence suggests that there is an association between the use of certain AEDs capable of enzyme induction and failure of oral contraceptives. One explanation for this interaction is that enzyme-inducing drugs effectively lower plasma concentrations of the relevant steroid hormones, resulting in unimpaired ovulation. However, other mechanisms, not related to enzyme induction, may contribute to the failure of oral contraceptives. Valproic acid is not a significant enzyme inducer and would not be expected to decrease concentrations of steroid hormones. However, clinical data about the interaction of valproic acid with oral contraceptives are minimal.  Administration of a single-dose of ethinylloestradiol (50 mcg)/levonorgestrel (250 mcg) to 6 women on valproic acid (200 mg twice daily) therapy for 2 months did not reveal any pharmacokinetic interaction.
Phenobarbital	CT C	↑ phenobarbital ↓ valproate	Phenobarbital increases the metabolism of valproic acid and hence, increases valproic acid metabolite levels. Therefore patients treated with this drug should be carefully monitored for signs and symptoms of hyperammonemia.  Valproate was found to inhibit the metabolism of phenobarbital. Co-administration of valproic acid (250 mg twice daily for 14 days) with phenobarbital to normal subjects (n = 6) resulted in a 50% increase in half-life and a 30% decrease in plasma clearance of phenobarbital (60 mg single-dose). The fraction of phenobarbital dose excreted unchanged increased by 50% in the presence of valproate.  There is evidence for severe CNS depression, with or without significant elevations of barbiturate or valproate serum concentrations. All patients receiving concomitant barbiturate therapy should be closely monitored for neurological toxicity. Serum barbiturate concentrations should be obtained, if possible, and the barbiturate dosage decreased, if appropriate.
Phenytoin	CT C	↑ phenytoin ↓ valproate	Phenytoin increases the metabolism of valproic acid and hence, increases valproic acid metabolite levels. Therefore patients treated with this drug should be carefully monitored for signs and symptoms of hyperammonemia.  Valproate displaces phenytoin from its plasma albumin binding sites and inhibits its hepatic metabolism. Co-administration of valproic acid (400 mg three times daily) with phenytoin (250 mg) in normal volunteers (n = 7) was

**Table 1: Summary of Drug-Drug Interaction Studies Including Important Interactions, Non-clinically Important Interactions and No Observed Interactions**

Concomitant Drug	Ref	Effect	Clinical Comment
			<p>associated with a 60% increase in the free fraction of phenytoin. Total plasma clearance and apparent volume of distribution of phenytoin increased 30% in the presence of valproate. Both the clearance and apparent volume of distribution of free phenytoin were reduced by 25%.</p> <p>In patients with epilepsy, there have been reports of breakthrough seizures occurring with the combination of valproic acid and phenytoin. The dosage of phenytoin should be adjusted as required by the clinical situation.</p>
Primidone	T	↑ phenobarbital	Primidone is metabolized into a barbiturate (phenobarbital), and therefore, may also be involved in a similar or identical interaction with valproic acid as phenobarbital.
Propofol		↑ propofol	Valproic acid may inhibit the metabolism of propofol, thus increasing propofol exposure. Reductions in propofol dose of 26 - 35% have been observed when co-administered with valproic acid. The normal dose of propofol may be excessive for patients receiving oral valproic acid treatment and may induce complications or delay recovery from anesthesia in electroconvulsive therapy (ECT).
Quetiapine			Co-administration of valproate and quetiapine may increase the risk of neutropenia/leucopenia.
Rifampin	CT	↓ valproate	A study involving the administration of a single dose of valproic acid (7 mg/kg) 36 hours after 5 nights of daily dosing with rifampin (600 mg) revealed a 40% increase in the oral clearance of valproate. Valproic acid dosage adjustment may be necessary when it is co-administered with rifampin.
Rufinamide		↑ rufinamide	Valproic acid may lead to an increase in plasma levels of rufinamide in a dose-dependent manner. This increase is dependent on concentration of valproic acid. Caution should be exercised particularly in children, as this effect is larger in the pediatric population.
Selective Serotonin Re-Uptake Inhibitors (SSRIs)	C	↑ valproate	Some evidence suggests that SSRIs inhibit the metabolism of valproic acid, resulting in higher than expected levels of valproate.
Tolbutamide	T	↑ tolbutamide	From <i>in vitro</i> experiments, the unbound fraction of tolbutamide was increased from 20 to 50% when added to plasma samples taken from patients treated with valproic acid. The clinical relevance of this displacement is unknown.
Topiramate	CT	Effect unknown	<p><b>Hyperammonemia:</b> Concomitant administration of valproate and topiramate has been associated with encephalopathy and/or hyperammonemia. Patients treated with those two drugs should be carefully monitored for signs and symptoms of hyperammonemic encephalopathy (see also CONTRAINDICATIONS, patients with known urea cycle disorders and WARNINGS AND PRECAUTIONS, Endocrine and Metabolism).</p> <p><b>Hypothermia:</b> Concomitant administration of topiramate with valproic acid has also been associated with hypothermia in patients who have tolerated either drug alone. Blood ammonia levels should be measured in patients with reported onset of</p>

<b>Table 1: Summary of Drug-Drug Interaction Studies Including Important Interactions, Non-clinically Important Interactions and No Observed Interactions</b>			
<b>Concomitant Drug</b>	<b>Ref</b>	<b>Effect</b>	<b>Clinical Comment</b>
			hypothermia (see WARNINGS AND PRECAUTIONS, Endocrine and Metabolism, Hypothermia).
Warfarin	T	Effect unknown	In an <i>in vitro</i> study, valproate increased the unbound fraction of warfarin by up to 32.6%. The therapeutic relevance of this is unknown, however, coagulation tests should be monitored if valproic acid is instituted in patients taking anticoagulants.  Caution is recommended when valproic acid is administered with drugs affecting coagulation (see ADVERSE REACTIONS).

Legend: C = Case Study; CT = Clinical Trial; T = Theoretical

### **Drug-Food Interactions**

Co-administration of valproic acid with food should cause no clinical problems in the management of patients with epilepsy.

### **Drug-Herb Interactions**

Interactions with herbal products have not been established.

### **Drug-Laboratory Interactions**

Valproic acid is partially eliminated in the urine as a ketone-containing metabolite which may lead to a false interpretation of the urine ketone test.

There have been reports of altered thyroid function tests associated with valproic acid; the clinical significance of these is unknown. See WARNINGS AND PRECAUTIONS, Special Populations, Pregnant Women, Thyroid Gland Abnormalities.

### **Drug-Lifestyle Interactions**

Refer to WARNINGS AND PRECAUTIONS, Neurologic, Driving and Hazardous Occupations for details.

## **DOSAGE AND ADMINISTRATION**

### **Dosing Considerations**

Patients receiving combined antiepileptic therapy require careful monitoring when another agent is started, stopped or when the dose is altered (see DRUG INTERACTIONS).

As the dosage of Sandoz Valproic (valproic acid) is titrated upward, blood concentrations of phenobarbital and/or phenytoin may be affected (see DRUG INTERACTIONS).

AEDs should not be abruptly discontinued in patients in whom the drug is administered to prevent major seizures because of the strong possibility of precipitating status epilepticus with attendant hypoxia and threat to life.

Any changes in dosage and administration, or the addition or discontinuance of concomitant drugs, should ordinarily be accompanied by close monitoring of clinical status and valproate plasma concentrations.

When changing therapy involving drugs known to induce hepatic microsomal enzymes (e.g., carbamazepine) or other drugs with valproate interactions (see DRUG INTERACTIONS), it is advisable to monitor serum valproate concentrations.

**Dosing in Elderly Patients:** Due to a decrease in unbound clearance of valproate and possibly a greater sensitivity to somnolence in the elderly, the starting dose should be reduced. Dosage should be increased more slowly and with regular monitoring for fluid and nutritional intake, dehydration, somnolence, urinary tract infection, and other adverse events. Dose reductions or discontinuation of Sandoz Valproic should be considered in patients with decreased food or fluid intake and in patients with excessive somnolence. The ultimate therapeutic dose should be achieved on the basis of clinical response (see WARNINGS AND PRECAUTIONS, Special Populations, Geriatrics [ $\geq 65$  years of age]).

**Dose-Related Adverse Events:** The frequency of adverse events (particularly elevated liver enzymes and thrombocytopenia) may be dose related. The probability of thrombocytopenia appears to increase significantly at total valproate concentration of  $\geq 110$  mcg/mL [females] or  $\geq 135$  mcg/mL [males] (see WARNINGS AND PRECAUTIONS, Hematologic, Dosing-related Adverse Reactions, Thrombocytopenia). Therefore, the benefit of improved therapeutic effect with higher doses should be weighed against the possibility of a greater incidence of adverse effects.

**Recommended Dose and Dosage Adjustment**

Sandoz Valproic (valproic acid) is administered orally. The recommended initial dosage is 15 mg/kg/day, increasing at one-week intervals by 5 to 10 mg/kg/day until seizures are controlled or side effects preclude further increases.

The maximum recommended dosage is 60 mg/kg/day. When the total daily dose exceeds 250 mg, it should be given in a divided regimen (Table 2).

Table 2 Initial Doses by Weight (based on 15 mg/kg/day)					
Weight		Total Daily Dose (mg)	Number of 250 mg Capsules		
kg	lb		Dose 1	Dose 2	Dose 3
10 - 24.9	22 - 54.9	250	0	0	1
25 - 39.9	55 - 87.9	500	1	0	1
40 - 59.9	88 - 131.9	750	1	1	1
60 - 74.9	132 - 164.9	1000	1	1	2
75 - 89.9	165 - 197.9	1250	2	1	2

**Therapeutic Blood Levels:** A good correlation has not been established between daily dose, total serum valproate concentration and therapeutic effect. However, therapeutic valproate serum concentrations for most patients with epilepsy will range from 50 to 100 mcg/mL (350 to 700 micromole/L). Some patients may be controlled with lower or higher serum concentrations



(see WARNINGS AND PRECAUTIONS).

**Conversion from valproic acid to divalproex sodium:** Divalproex sodium enteric-coated tablets dissociate to the valproate ion in the gastrointestinal tract. Divalproex sodium tablets are uniformly and reliably absorbed, however, because of the enteric coating, absorption is delayed by an hour when compared to valproic acid.

The bioavailability of both types of divalproex sodium tablets (regular and extended release) is equivalent to that of valproic acid capsules.

In patients previously receiving valproic acid therapy, divalproex sodium should be initiated at the same daily dosing schedule. After the patient is stabilized on divalproex sodium, a dosing schedule of two or three times a day may be elected in selected patients. Changes in dosage administration of divalproex sodium or concomitant medications should be accompanied by increased monitoring of plasma concentrations of valproate and other medications, as well as the patient's clinical status.

**Female children, female adolescents, women of childbearing potential and pregnant women**

Valproic acid treatment should be initiated and supervised by a specialist experienced in the management of epilepsy. Treatment should only be initiated if other treatments are ineffective or not tolerated and the benefit and risk should be carefully re-assessed at regular treatment reviews. Preferably valproic acid should be prescribed at the lowest effective dose. The daily dose should be divided into at least two single doses. Available data suggest that antiepileptic polytherapy including valproate is associated with a greater risk of congenital malformations than valproate monotherapy. Therefore monotherapy should be favoured when possible.

**Missed Dose**

The patient should not abruptly stop taking their medication because of the risk of increasing their seizures.

If the patient misses a dose, they should not try to make up for it by doubling up on their next dose. They should take their next regularly scheduled dose and try not to miss any more doses.

**Administration**

Sandoz Valproic may be taken with or without food.

Patients who experience gastrointestinal irritation may benefit from administration of the drug with food or by a progressive increase of the dose from an initial low level. The capsules should be swallowed without chewing to avoid local irritation of the mouth and throat. Co-administration of Sandoz Valproic (valproic acid) with food should cause no clinical problems in the management of patients with epilepsy.

**OVERDOSAGE**

For management of a suspected drug overdose, contact your regional Poison Control Centre.
---

Overdosage with valproic acid may result in somnolence, muscular hypotonia, hyporeflexia, miosis, impaired respiratory function, hypotension, metabolic acidosis, heart block, deep coma and circulatory collapse/shock. Cases of intracranial hypertension related to cerebral oedema have been reported. Fatalities have been reported; however, patients have recovered from valproate levels as high as 2120 mcg/mL.

In a reported case of overdosage with valproic acid after ingesting 36 g in combination with phenobarbital and phenytoin, the patient presented in deep coma. An electroencephalogram (EEG) recorded diffuse slowing, compatible with the state of consciousness. The patient made an uneventful recovery.

In overdose situations, the fraction of drug not bound to protein is high and hemodialysis or tandem hemodialysis plus hemoperfusion may result in significant removal of drug. The benefit of gastric lavage or emesis will vary with the time since ingestion. As valproic acid is absorbed very rapidly, gastric lavage may be of limited value. General supportive measures should be applied with particular attention to the prevention of hypovolemia and the maintenance of adequate urinary output.

Naloxone has been reported to reverse the CNS depressant effects of valproic acid overdosage. Because naloxone could theoretically also reverse the antiepileptic effects of valproic acid, it should be used with caution.

## **ACTION AND CLINICAL PHARMACOLOGY**

Pharmacotherapeutic group: Anticonvulsant and mood-stabilizing drug; ATC-Code: N03AG01.

### **Mechanism of Action**

Valproic acid has anticonvulsant properties. Although its mechanism of action has not yet been established, it has been suggested that its activity is related to increased brain levels of gamma-aminobutyric acid (GABA). The effect on the neuronal membrane is unknown.

### **Pharmacodynamics**

A good correlation has not been established between daily dose, serum level and therapeutic effect. In epilepsy, the therapeutic plasma concentrations range is believed to be from 50 to 100 mcg/mL (350 to 700 micromole/L) of total valproate. Occasional patients may be controlled with serum levels lower or higher than this range (see DOSAGE AND ADMINISTRATION).

### **Pharmacokinetics**

**Absorption:** Valproic acid is rapidly absorbed after oral administration. Peak serum levels occur approximately 1 to 4 hours after a single oral dose. A slight delay in absorption occurs when the drug is administered with meals but this does not affect the total absorption.

**Distribution:** Valproic acid is rapidly distributed throughout the body and the drug is strongly bound (90%) to human plasma proteins. Increases in dose may result in decreases in the extent of protein binding and variable changes in valproic acid clearance and elimination.

### Protein Binding

The plasma protein binding of valproate is concentration dependent and the free fraction increases from approximately 10% at 40 mcg/mL to 18.5% at 130 mcg/mL. Protein binding of valproate is reduced in the elderly, in patients with chronic hepatic diseases, in patients with renal impairment, in hyperlipidemic patients, and in the presence of other drugs (e.g., acetylsalicylic acid). Conversely, valproate may displace certain protein-bound drugs (e.g., phenytoin, carbamazepine, warfarin, and tolbutamide) (see DRUG INTERACTIONS for more detailed information on the pharmacokinetic interactions of valproate with other drugs).

### CNS Distribution

Valproate concentrations in cerebrospinal fluid (CSF) approximate unbound concentrations in plasma (ranging from 7 to 25% of total concentration).

**Metabolism:** Valproate is metabolized almost entirely by the liver. In adult patients on monotherapy, 30 to 50% of an administered dose appears in urine as a glucuronide conjugate. Mitochondrial (beta)-oxidation is the other major metabolic pathway, typically accounting for over 40% of the dose. Usually, less than 15 to 20% of the dose is eliminated by other oxidative mechanisms. Less than 3% of an administered dose is excreted unchanged in urine.

Due to the saturable plasma protein binding, the relationship between dose and total valproate concentration is nonlinear; concentration does not increase proportionally with the dose, but rather increases to a lesser extent. The kinetics of unbound drug are linear.

**Excretion:** Mean plasma clearance and volume of distribution for total valproate are 0.56 L/hr/1.73 m<sup>2</sup> and 11 L/1.73 m<sup>2</sup>, respectively. Mean plasma clearance and volume of distribution for free valproate are 4.6 L/hr/1.73 m<sup>2</sup> and 92 L/1.73 m<sup>2</sup>, respectively. These estimates cited apply primarily to patients who are not taking drugs that affect hepatic metabolizing enzyme systems. For example, patients taking enzyme-inducing AEDs (carbamazepine, phenytoin, and phenobarbital) will clear valproate more rapidly. Because of these changes in valproic acid clearance, monitoring of valproate and concomitant drug concentrations should be intensified whenever enzyme-inducing drugs are introduced or withdrawn.

Elimination of valproic acid and its metabolites occurs principally in the urine, with minor amounts in the feces and expired air. Very little unmetabolized parent drug is excreted in the urine.

The serum half-life ( $t_{1/2}$ ) of valproic acid is typically in the range of 6 to 16 hours. Half-lives in the lower part of the above range are usually found in patients taking other AEDs capable of enzyme induction.

### Special Populations and Conditions

**Neonates/ Infants:** Within the first 2 months of life, infants have a markedly decreased ability to eliminate valproate compared to children and adults. This is a result of reduced clearance (perhaps due to delay in development of glucuronosyltransferase and other enzyme systems involved in valproate elimination) as well as increased volume of distribution (in part due to decreased plasma protein binding). For example, in one study, the half-life in neonates under

10 days ranged from 10 to 67 hours, compared to a range of 7 to 13 hours in children greater than 2 months.

**Pediatrics:** Patients between 3 months and 10 years have 50% higher clearances expressed on weight (i.e., L/min/kg) than do adults. Over the age of 10 years, children have pharmacokinetic parameters that approximate those of adults.

**Geriatrics:** The capacity of elderly patients (age range: 68 to 89 years) to eliminate valproate has been shown to be reduced compared to younger adults (age range: 22 to 26 years). Intrinsic clearance is reduced by 39%; the free fraction is increased by 44% (see DOSAGE AND ADMINISTRATION).

**Gender:** There are no differences in unbound clearance (adjusted for body surface area) between males and females ( $4.8 \pm 0.17$  and  $4.7 \pm 0.07$  L/hr per  $1.73 \text{ m}^2$ , respectively).

**Race:** The effects of race on the kinetics of valproate have not been studied.

**Hepatic Insufficiency:** See CONTRAINDICATIONS and WARNINGS AND PRECAUTIONS, Hepatic/Biliary/Pancreatic, Serious or Fatal Hepatotoxicity, for statements regarding hepatic dysfunction and associated fatalities.

**Renal Insufficiency:** See WARNINGS AND PRECAUTIONS, Renal, Renal Impairment.

**Genetic Polymorphism:** No data available on genetic polymorphism.

## STORAGE AND STABILITY

Sandoz Valproic capsules 250 mg should be stored between 15 and 30°C, away from heat and humidity. Keep bottle tightly closed when not in use.

## DOSAGE FORMS, COMPOSITION AND PACKAGING

### Composition

Sandoz Valproic (valproic acid) 250 mg capsules contain the following nonmedicinal ingredients: corn oil, gelatin, glycerin, methylparaben\*, propylparaben\*, titanium dioxide, FD&C yellow #6.

\* methylparaben and propylparaben may cause allergic reactions (possibly delayed).

### Availability of Dosage Forms

Each orange-coloured soft gelatin capsule contains valproic acid 250 mg and is imprinted with the mark "SZ 250". Bottles of 100 and 500 capsules.

## PART II: SCIENTIFIC INFORMATION


### PHARMACEUTICAL INFORMATION

Proper name: valproic acid

Chemical name: 2-propylpentanoic acid or dipropylacetic acid

Molecular formula and molecular mass:  $C_8H_{16}O_2$ ; 144.21 g/mol

Structural formula:



Physicochemical properties: Valproic acid is a colourless, oily liquid which is very soluble in organic solvents and slightly soluble in water.

## CLINICAL TRIALS

The following tables present the summaries of the comparative bioavailability data for Sandoz Valproic acid capsules and the Canadian reference product, under fasted and fed conditions, respectively. The study consisted of a randomized, single oral dose, 4 x 4 Latin Square Cross-over design with a one week wash out period. 24 male, volunteers, non smokers, were used, all of which completed the study.

### Fasting Conditions

<p><b>Valproic Acid</b>  <b>(1 x 250 mg)</b>  <b>From measured data</b></p> <p><b>Geometric Mean</b>  <b>Arithmetic Mean (CV %)</b></p>
---

Parameter	Test Sandoz Valproic	Reference Depakene <sup>®†</sup>	% Ratio of Geometric Means	95 % Confidence Interval
AUC <sub>T</sub> mcg/ml·h	417.11 426.0 (20.4)	411.84 420.1 (19.9)	102.2	98.0-106.5
AUC <sub>∞</sub> mcg/ml·h	460.24 472.6 (23.2)	455.63 468.7 (24.7)	101.7	97.1-106.5
C <sub>max</sub> mcg/ml·h	28.52 29.18 (23.2)	28.77 29.45 (23.2)	98.5	91.3-106.4
T <sub>max</sub> <sup>§</sup> (h)	1.83 (59.5)	2.63 (57.3)		
T <sub>1/2</sub> <sup>§</sup> (h)	13.95 (18.7)	13.83 (27.1)		

<sup>†</sup> Depakene<sup>®</sup> Capsules 250 mg was manufactured by Abbott Canada.

<sup>§</sup> Expressed as the arithmetic mean (CV%) only.

**Fed Conditions**

<b>Valproic Acid (1 x 250 mg) From measured data</b>
<b>Geometric Mean Arithmetic Mean (CV %)</b>

<b>Parameter</b>	<b>Test Sandoz Valproic</b>	<b>Reference Depakene<sup>®†</sup></b>	<b>% Ratio of Geometric Means</b>	<b>95 % Confidence Interval</b>
AUC <sub>T</sub> mcg/ml·h	391.10 397.4 (17.9)	404.21 410.5 (17.7)	96.5	92.6-100.6
AUC <sub>∞</sub> mcg/ml·h	425.14 432.9 (18.9)	450.22 462.6 (25.8)	97.1	92.7-101.8
C <sub>max</sub> mcg/ml·h	23.81 24.38(24.8)	24.90 25.19 (15.6)	94.8	87.8-102.4
Tmax <sup>§</sup> (h)	4.71 (26.4)	4.46 (26.9)		
T <sub>1/2</sub> <sup>§</sup> (h)	13.53 (17.6)	13.76 (30.7)		

<sup>†</sup> Depakene<sup>®</sup> Capsules 250 mg was manufactured by Abbott Canada.

<sup>§</sup> Expressed as the arithmetic mean (CV%) only.

## **DETAILED PHARMACOLOGY**

### **Animal**

Valproic acid has been shown to be effective against several types of chemically and electrically induced convulsions in a variety of animal species. These included maximal electroshock, low frequency electroshock, CO<sub>2</sub> withdrawal, pentylene tetrazole, cobalt, bemegride, bicuculline and l-glutamate. Many forms of photic and auditory induced seizures are also effectively blocked by valproic acid.

In animal studies, valproic acid at doses of 175 mg/kg or less had no effect on locomotor activity and conditioned responses to positive reinforcement.

Doses greater than 175 mg/kg inhibited spontaneous and conditioned behaviour in mice and rats and interfered with coordination of hind limbs in rats. Suppression of spontaneous and evoked brain potentials was also demonstrated at these higher dose levels.

Valproic acid at doses of 175 mg/kg or less had little or no effect on the autonomic nervous system, cardiovascular system, respiration, body temperature, inflammatory responses, smooth muscle contraction or renal activity. Intravenous doses of 22, 43 and 86 mg/kg in animals caused very transient decreases followed by compensatory increases in blood pressure.

Sodium valproate injectable caused decreased activity, ataxia, dyspnea, prostration and death in rats and mice acutely exposed to dosages exceeding 200 mg/kg.

## **TOXICOLOGY**

The initial animal testing was done with sodium valproate, whereas most of the recent research has been with valproic acid. The conversion factor is such that 100 mg of the sodium salt is equivalent to 87 mg of the acid. References to dosage are in terms of valproic acid activity.

### **Acute Toxicity**

Acute toxicity has been determined in several animal species using oral, intravenous, intraperitoneal and subcutaneous routes. The oral median lethal dose in adult rats and dogs was about 1 to 2 g/kg. Toxicity was similar for both sexes; however, it tended to be greater in newborn and 14-day old rats and in young adult rats. The signs of toxicity were those of central nervous system depression. Specific organ damage was limited to cellular debris in reticuloendothelial tissue and slight fatty degeneration of the liver.

Large oral doses (more than 500 mg/kg) produced irritation of the gastrointestinal tract of rats.

In adult male mice, the oral medial lethal dose of divalproex sodium was 1.66 g/kg (equal to approximately 1.54 g/kg valproic acid).

Pulverized divalproex sodium enteric-coated tablets (equivalent to 250 mg valproic acid), suspended in 0.2% methylcellulose, were administered orally to mice and rats of both sexes (10/sex/species/group) in dosages ranging from 1.74 to 4.07 g/kg. The oral median lethal dose



(LD<sub>50</sub>) ranged from 2.06 to 2.71 g/kg. No consistent sex-related or species-related differences were observed.

Signs of central nervous system depression, such as decreased activity, ataxia, and sleep, were observed. At necropsy, discolouration and/or thickening of the glandular mucosa were observed in only 2 female rats treated with 2.71 g/kg that died acutely.

When mature rats and dogs were administered up to 240 mg/kg/day or 120 mg/kg/day, respectively, for at least four consecutive weeks, no significant toxicologic effects were reported. However, significant reductions in testicular weights and total white cell counts in rats given 240 mg/kg/day were considered as evidence of subtle toxicity from sodium valproate injectable. Therefore, 90 mg/kg/day in rats and 120 mg/kg/day in dogs were considered the highest nontoxic doses.

The acute intravenous toxicity of sodium valproate injectable formulation containing the equivalent of 100 mg valproic acid/mL was evaluated in both sexes of mice and rats. Groups of mice and rats (five/sex/species/group) were treated at dosages ranging from 0.5 to 9.0 mL/kg (50 to 900 mg valproate/kg). No overt signs of toxicity were present in rats and mice given 0.5 mL/kg (50 mg valproate/kg). LD<sub>50</sub> values for the test solution in mice and rats (data combined for both sexes) were 7.3 and 7.0 mL/kg (730 and 700 mg valproate/kg), respectively.

### **Subacute and Chronic Toxicity**

Subacute and chronic toxicity studies consisted of 1, 3, 6 and 18 months studies in rats and 3, 6 and 12 months studies in dogs. Pathologic changes included suppression of the hematopoietic system, depletion of lymphocytes from lymphoid tissues and the loss of germinal epithelial cells from seminiferous tubules. Reduced spermatogenesis and testicular atrophy occurred in dogs at doses greater than 90 mg/kg/day and in rats at doses greater than 350 mg/kg/day. In rats, the first indication of toxicity at 350 mg/kg/day was decreased food consumption and growth.

### **Mutagenicity and Carcinogenicity**

**Mutagenicity:** Valproate was not mutagenic in an *in vitro* bacterial assay (Ames test), did not produce dominant lethal effects in mice, and did not increase chromosome aberration frequency in an *in vivo* cytogenetic study in rats. Increased frequencies of sister chromatid exchange (SCE) have been reported in a study of epileptic children taking valproate, but this association was not observed in another study conducted in adults. There is some evidence that increased SCE frequencies may be associated with epilepsy. The biological significance of increase in SCE frequency is not known.

**Carcinogenicity:** Two hundred rats were given valproic acid in the diet for 107 weeks. Mean doses consumed in the treatment period were: 81 mg/kg/day (males) and 85 mg/kg/day (females), in the low dose group; 161 mg/kg/day (males) and 172 mg/kg/day (females) in the high dose group (approximately 10 to 50% of the maximum human daily dose on a mg/m<sup>2</sup> basis). Control animals received corn oil in the diet. The chief finding in the study was an increased incidence of skin fibrosarcomas in treated males of the high-dose group. There were 2 such neoplasms in the low dose group, 5 in the high dose group and none in control males. Fibrosarcomas in rats are relatively infrequent, usually occurring in less than 3% of animals.

Valproic acid was also administered in the diet to female mice for nearly 19 months at doses of 81 and 163 mg/kg/day and to male mice for nearly 23 months at doses of 80 and 159 mg/kg/day. A significant dose related trend occurred in male mice in the incidence of bronchoalveolar adenomas, and when the data were adjusted for the times of death, the incidence in the high dose group was significantly increased.

Depending on the method of statistical analysis, the incidence of hepatocellular carcinomas and/or adenomas also showed significant or almost significant increases for the corresponding observations. The results of these two studies indicate that valproic acid is a weak carcinogen or promoter in rats and mice. The significance of these findings for humans is unknown at present.

### **Reproduction and Teratology**

Studies in rats have shown placental transfer of the drug. Doses greater than 65 mg/kg/day given to rats, mice and rabbits produced an increased incidence of skeletal abnormalities of the ribs, vertebrae and palate.

Doses greater than 150 mg/kg/day given to pregnant rabbits produced fetal resorptions and (primarily) soft-tissue abnormalities in the offspring.

In rats, there was a dose related delay in onset of parturition. Post-natal growth and survival of the progeny were adversely affected, particularly when drug administration spanned the entire gestation and early lactation period. Embryo lethality or major developmental abnormalities occurred in rats and rabbits at doses of 350 mg/kg/day.

Survival among pups born to the high dose females was very poor but was improved when pups were transferred to control dams shortly after birth.

**Fertility:** Chronic toxicity studies in juvenile and adult rats and dogs demonstrated reduced spermatogenesis and testicular atrophy at oral doses of valproic acid of 400 mg/kg/day or greater in rats (approximately equivalent to or greater than the maximum human daily dose on a mg/m<sup>2</sup> basis) and 150 mg/kg/day or greater in dogs (approximately 1.4 times the maximum human daily dose or greater on a mg/m<sup>2</sup> basis). Segment I fertility studies in rats have shown that oral doses up to 350 mg/kg/day (approximately equal to the maximum human daily dose on a mg/m<sup>2</sup> basis) for 60 days have no effect on fertility.

The effect of valproate on testicular development and on sperm production and fertility in humans is unknown.

## REFERENCES

- 1 Godin Y, Heiner L, Mark J, Mandel P. Effects of di-n-propyl-acetate, An Anticonvulsive Compound on GABA Metabolism. *J. Neurochem.* 1969;16:869.
- 2 Simler S, Ciesielski L, Maitre M. Effects of Sodium n-dipropyl-acetate on Audiogenic Seizures and Brain Aminobutyric Acid Level. *Biochem. Pharmacol.* 1973;22:1701.
- 3 Eadie MJ. Plasma Level Monitoring of Anticonvulsants. *Clin Pharmacokinet.* 1976;1:52.
- 4 Gram L, Wulff K, Kasmussen KE, Flachs H, Wuertz-Jorgensen A, Somerbeck KW, Lohren V. Valproate Sodium: A Controlled Clinical Trial Including Monitoring of Drug Levels. *Epilepsia.* 1977;18:141.
- 5 Benoit JM, Besson JM, Bivot JB, Aleonard P. Effects of Na Dipropylacetate (DPA) on Dorsal Root Potentials Induced by Cortical Seizures. *J. Pharmacol.* 1971;2:23.
- 6 Frey HH, Loscher W. Di-n-propyl-acetic Acid. Profile of Anticonvulsant Activity in Mice. *Arzneim-Forsch.* 1976;26:299.
- 7 Kupferberg H, Lust WD, Penry JK. Anticonvulsant Activity of Dipropylacetic acid (DPA) in Relation to GABA and CGMP Brain Levels in Mice. *Fed. Proc.* 1975;34:283.
- 8 Misslin R, Ropartz PH, Mandel P. The Effects of n-dipropylacetate on the Acquisition of Conditioned Behaviour with Negative Reinforcement in Mice. *Psychopharmacologie.* 1975;44:263.
- 9 Shearer DE, Fleming DE, Bigler ED, Wilson CE. Suppression of Photically Evoked After-discharge Bursting Following Administration of Anticonvulsants in Waking Rats, *Pharmacol. Biochem. Behav.* 1974;2:839.
- 10 Shuto K, Nishigaki T. The Pharmacological Studies on Sodium Dipropylacetate Anticonvulsant Activities and General Pharmacological Actions. *Oyo Yakuri (Pharmacometrics).* 1970;4:937.
- 11 Simon D, Perry JK. Sodium di-n-propylacetate (DPA) in the Treatment of Epilepsy. A Review. *Epilepsia.* 1975;16:549.
- 12 Robert E, Guibaud P. Maternal Valproic Acid and Congenital Neural Tube Defects. *The Lancet* 2(8304): October 23, 1982;937.
- 13 Shakir RA. Comparison of Sodium Valproate and Phenytoin as Single Drug Treatment in Epilepsy. *Epilepsia.* 1981;22:27-33.
- 14 Shakir RA. Comparative Trial of Valproate Sodium and Clonazepam in Chronic Epilepsy. *Arch Neurol.* 1979;36:301-304.

- 15 Turnbull DM, Rawlins MD, Weightman D, Chadwick DW. A Comparison of Phenytoin and Valproate in Previously Untreated Adult Epileptic Patients. *J. Neurol Neurosur. Psychiat.* 1982;45:55-59.
- 16 Gugler R, vonUnruh GE. Clinical Pharmacokinetics of Valproic Acid. *Clinical Pharmacokinetics.* 1980;5:67-83.
- 17 Cramer JA, Mattson RH. Valproic Acid: In Vitro Plasma Protein Binding and Interaction with Phenytoin. *Ther. Drug. Monit.* 1979;1(1):105-116.
- 18 Bowdle TA, Patel JH, Levy RH. Valproic Acid Dosage and Plasma Protein Binding and Clearance. *Clin. Pharmacol. Ther.* 1980;28(4):487-492.
- 19 Gram L, Flachs H. Sodium Valproate, Serum Level and Clinical Effect in Epilepsy: A Controlled Study. *Epilepsia.* 1979;20:303-312.
- 20 Kapetamovic IM, Kupferberg HJ. Inhibition of Microsomal Phenobarbital Metabolism of Valproic Acid. *Biochem. Pharmacol.* 1981;30(11):1361-1363.
- 21 Patel IH, Levy RH. Phenobarbital - Valproic Acid Interaction. *Clin. Pharmacol. Ther.* 1979;27(4):515-521.
- 22 Bruni J, Wilder BJ. Valproic Acid and Plasma Levels of Phenobarbital. *Neurol.* 1980;30:94-97.
- 23 Pinder PM, Brogden RN. Sodium Valproate: A Review of Its Pharmacological Properties and Therapeutic Efficacy in Epilepsy. *Drugs.* 1977;B:81-123.
- 24 Millet Y, Sinty JM. Problems Created by Combining Phenobarbital and Sodium Dipropylacetate Therapeutically: Case Report. *Europ. Toxicol.* 1976;9(6):381-383.
- 25 Rai PV. Acute Intoxication During a Combined Treatment of Sodium Valproate and Phenobarbitone. In: Meinardi H and Rowan AJ, editors. *Advances in epileptology*, Swets Zertlinger, Amsterdam, pp. 366-369, 1978.
- 26 BGP Pharma ULC, Product Monograph: DEPAKENE. Control #: 217988, Date of Revision: October 9, 2018.

**PART III: CONSUMER INFORMATION**

**Pr SANDOZ VALPROIC  
valproic acid capsules**

**This leaflet is part III of a three-part "Product Monograph" published when Sandoz Valproic was approved for sale in Canada and is designed specifically for consumers. This leaflet is a summary and will not tell you everything about Sandoz Valproic. Contact your doctor or pharmacist if you have any questions about the drug.**

**ABOUT THIS MEDICATION**

**What the medication is used for:**

Sandoz Valproic has been prescribed to you to control your epilepsy. Please follow your doctor's recommendations carefully.

**What it does:**

Sandoz Valproic has anticonvulsant properties. The mechanism of action has not yet been established. It has been suggested that its activity in epilepsy is related to increased brain concentrations of gamma-aminobutyric acid (GABA).

**When it should not be used:**

Sandoz Valproic should not be taken by:

- patients with liver disease or significant liver dysfunction
- patients with mitochondrial diseases (e.g. Alpers or Alpers-Huttenlocher disease)
- patients who are allergic to valproic acid or any of the other ingredients in Sandoz Valproic
- patients with known urea cycle disorders (a genetic disorder)
- patients with known porphyria (a genetic disorder)

**What the medicinal ingredient is:**

Valproic acid

**What the non-medicinal ingredients are:**

Sandoz Valproic 250 mg capsules contain the following nonmedicinal ingredients: corn oil, gelatin, glycerin, methylparaben, propylparaben, titanium dioxide, FD&C yellow #6.

**What dosage forms it comes in:**

Sandoz Valproic is available as a capsule containing 250 mg of valproic acid.

**WARNINGS AND PRECAUTIONS**

**Serious Warnings and Precautions**

- **Hepatotoxicity:** liver failure resulting in death has occurred in patients receiving valproic acid. These incidents usually

occurred during the first 6 months of treatment with valproic acid. Patients taking several anticonvulsant drugs, children, those with a history of liver disease, metabolic disorders, severe seizure disorders accompanied by mental retardation, and those with brain disease may be at particular risk. Experience has indicated that children under the age of 2 years are at a considerably increased risk of developing fatal hepatotoxicity, especially those on multiple anticonvulsants.

- **Birth Defects:** Sandoz Valproic can cause birth defects and problems with early development of the child if it is taken during pregnancy. Sandoz Valproic should not be used in female children, in female adolescents, in women of childbearing potential and pregnant women unless other treatments do not work or are not tolerated. If you are a female of childbearing age you should use an effective method of birth control while you are taking Sandoz Valproic. Tell your doctor right away if you become pregnant or think you might be pregnant.
- **Pancreatitis:** cases of life threatening pancreas disorder have been reported in both children and adults receiving valproic acid. Some cases have occurred shortly after first use as well as after several years of use. Abdominal pain, nausea, vomiting and/or anorexia can be symptoms of pancreatitis that require immediate medical evaluation.

**BEFORE you use Sandoz Valproic talk to your doctor or pharmacist if you:**

- have a history of, or suffer from a liver disease, such as jaundice (yellowing of the skin and eyes).
- have ever had an unusual or allergic reaction to Sandoz Valproic (including fever or rash).
- are a female child or female adolescent, a woman of childbearing potential, or pregnant.
- are breast-feeding (nursing); Sandoz Valproic passes into breast milk. You must discuss with your doctor whether you should breastfeed or take Sandoz Valproic. You cannot do both.
- are male and thinking about fathering a child. Sandoz Valproic can make you less fertile.
- are taking any other prescription or over the counter medicine.
- have kidney disease.
- have other medical conditions including a history of unexplained coma, intellectual disability or any type of brain dysfunction.
- have a psychiatric disorder or have thoughts of suicide.
- consume alcohol on a regular basis.

**Precautions while taking Sandoz Valproic:**

- Your doctor will monitor your response to Sandoz Valproic on a regular basis. However, if your seizures get worse, you should tell your doctor immediately.
- Since Sandoz Valproic may cause poor coordination and/or drowsiness, you should not engage in hazardous activities, such as driving and operating machinery, until you know that

you don't become drowsy from the drug.

- You should not stop taking your medication unless directed by your doctor. You should always check that you have an adequate supply of Sandoz Valproic. You should remember that this medicine was prescribed only for you; it should never be given to anyone else.
- As with other drugs used to treat epilepsy, some patients may experience an increase in the number of seizures and the severity (including status epilepticus), or the onset of new types of seizures with Sandoz Valproic instead of an improvement. If you start having more seizures, new types of seizures or your seizures get worse contact your doctor immediately.

### Female Children, Female Adolescents and Women of Childbearing Potential

- All female children, female adolescents and women of childbearing age who are being treated with Sandoz Valproic should talk to their healthcare providers about using other possible treatments instead of Sandoz Valproic. If you are a female capable of becoming pregnant you should only take Sandoz Valproic if nothing else works for you. If the decision is made to use Sandoz Valproic, you must use an effective method of birth control (contraception). You should talk to your doctor about the best kind of birth control to use while you are taking Sandoz Valproic.
- No longer getting your period, fluid filled sacs (cysts) on the ovaries and increased testosterone levels have been reported in women taking Sandoz Valproic.
- Before prescribing this medicine to you, your doctor will have explained what might happen to your baby if you become pregnant while taking Sandoz Valproic. If you decide later you want to have a child you should not stop taking your medicine until you have discussed this with your doctor and agreed on a plan for switching to another medication if this is possible.
- Ask your doctor about taking folic acid when trying to get pregnant. Folic acid can lower the general risk of birth defects in the spine (spina bifida) and early miscarriage that exists with all pregnancies. However, it is unlikely that folic acid will reduce the risk of birth defects associated with Sandoz Valproic use.

### Pregnant Women

- Sandoz Valproic carries a risk if taken during pregnancy. The higher the dose, the higher the risks, though all doses carry a risk.
- If you take Sandoz Valproic during pregnancy, your child has a serious risk of birth defects and problems with development which can be seriously debilitating such as lower IQ and problems with brain development. Birth defects which have been reported include spina bifida (where the bones of the spine are not properly developed); problems with the development of the bones of the face and skull; and problems with the development of the heart, kidney, urinary tract and

sexual organs, arms and legs. These can begin early in the pregnancy, even before you know that you are pregnant.

- It is estimated that up to 30-40% of preschool children whose mothers took Sandoz Valproic during pregnancy may have problems with early childhood development. This means these children can be slow to walk and talk, have lower intelligence (lower IQ scores) than other children, and have difficulty with language and memory.
- Children born to mothers who took Sandoz Valproic during pregnancy are also more likely to have Autism spectrum disorders and more likely to develop symptoms of Attention Deficit Hyperactivity Disorder (ADHD).
- There may be other medications to treat your condition that have a lower chance of birth defects.
- If you are planning to become pregnant, or if you become pregnant while taking Sandoz Valproic, you should promptly inform your doctor. Do not suddenly stop taking the drug. Appropriate treatment options will need to be discussed with your physician to ensure the benefits outweigh the risks.
- Pregnancy Registry: If you become pregnant while taking Sandoz Valproic, talk to your doctor about registering with the North American Antiepileptic Drug Pregnancy Registry. You can enroll in this registry by calling 1-888-233-2334. The purpose of this registry is to collect information about the safety of antiepileptic medicines during pregnancy. Information on the registry can also be found at the following website:  
<http://www.aedpregnancyregistry.org/>.

### INTERACTIONS WITH THIS MEDICATION

#### Serious Drug Interactions

- Rare cases of coma have been reported in patients receiving valproic acid alone or when taken with phenobarbital.
- Serious skin reactions (such as conditions called Stevens-Johnson syndrome and Toxic Epidermal Necrolysis) have been reported when valproic acid and lamotrigine were taken together.

#### Drugs that may interact with Sandoz Valproic include:

- Anticonvulsants such as carbamazepine, lamotrigine, primidone, topiramate, felbamate, phenytoin, ethosuximide, phenobarbital, olanzapine, rufinamide;
- Anticoagulants such as acetylsalicylic acid, warfarin, dicumarol;
- Benzodiazepines such as diazepam, lorazepam, clonazepam;
- Some medicines used to treat infections such as rifampin;
- Some medicines used to treat diabetes such as tolbutamide;
- Some HIV-antiviral medication such as zidovudine, ritonavir, lopinavir, lamivudine;
- Any of the group of antibiotics in the carbapenem class such as doripenem, ertapenem, imipenem, meropenem;
- Some medicines used to treat heartburn and peptic ulcers such as cimetidine;

- Medicines used to treat depression such as Selective Serotonin Re-Uptake Inhibitors (SSRIs), Monoamine Oxidase Inhibitors (MAOIs), Tricyclic antidepressants such as amitriptyline, nortriptyline;
- Acetazolamide a medicine used to treat glaucoma and epilepsy;
- Cholestyramine, a medicine used to lower cholesterol;
- Propofol, a drug used to relax you before and after surgery;
- Nimodipine;
- Antipsychotics;
- Estrogen-containing products (including contraceptives that contain estrogen).

**PROPER USE OF THIS MEDICATION**

Sandoz Valproic treatment must only be started and supervised by a doctor specialised in the treatment of epilepsy. Please consult your doctor before taking any other medication, including over-the-counter medicines. Some drugs can produce various side effects when they are used in combination with Sandoz Valproic.

It is important to keep your appointments for medical checkups.

The doctor may need to take blood tests to measure the amount of Sandoz Valproic in your blood when adjusting your medications.

Do not stop taking Sandoz Valproic suddenly as this can cause a serious increase in the number of seizures and their severity, including status epilepticus.

**Usual dose:**

It is very important to take Sandoz Valproic exactly as instructed by your doctor.

The recommended starting dose of Sandoz Valproic will be decided by your doctor based on your weight, your seizures and your concomitant medications. Be sure to tell your doctor all the prescription and over the counter medications that you are currently taking. Your doctor will gradually increase the dosage until your condition is well controlled without experiencing side effects. You should carefully follow the instructions that were given to you and not change your dose without consulting with your doctor.

Sandoz Valproic may be taken with or without food.

**Overdose:**

If you think you have taken too much Sandoz Valproic, contact your healthcare professional, hospital emergency department or regional poison control centre immediately, even if there are no symptoms.

**Missed Dose:**

If you miss a dose, you should not try to make up for it by

doubling up on your next dose. You should take your next regularly scheduled dose and try not to miss any more doses.

**SIDE EFFECTS AND WHAT TO DO ABOUT THEM**

You should check with your doctor or pharmacist right away if you notice any bothersome or unusual effects while taking Sandoz Valproic.

The most commonly reported adverse reactions include nausea, vomiting, indigestion sleepiness, headache, diarrhea, weakness, tremor and dizziness. Changes in hair are also reported, such as hair loss or an increase in hair on face, chest and back. If any of these affect you severely, contact your doctor or pharmacist.

You should know that this does not mean that you will experience such effects, because people can react in different ways to the same medicine.

**SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM**

Symptom / effect		Talk with your doctor or pharmacist right away		Seek Emergency medical attention
		Only if severe	In all cases	
Common	<b>Hallucinations:</b> seeing or hearing something that is not there.	✓		
	<b>Urinary incontinence</b>		✓	
Uncommon	<b>Brain dysfunction from high ammonia levels in the blood:</b> tiredness, vomiting, abnormal walking, extreme irritability <sup>†</sup> , combative/bizarre behaviour <sup>††</sup> , refusal to eat meat or high protein products <sup>††</sup>		✓	

**SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM**

Symptom / effect	Talk with your doctor or pharmacist right away		Seek Emergency medical attention
	Only if severe	In all cases	
<b>Decreased number of platelets in the blood:</b> may result in easy bruising and bleeding from the skin or other areas		✓	
<b>Liver disorder:</b> weakness, tiredness, abdominal pain, diarrhea, facial swelling, loss of appetite, yellowing of the skin or eyes, dark urine, nausea and vomiting		✓	
<b>Pancreas disorder:</b> abdominal pain, nausea, vomiting, and/or loss of appetite		✓	
<b>Thoughts of suicide or hurting yourself:</b> symptoms of depression or unusual changes in mood or behaviour		✓	
Rare <b>Muscle disorder:</b> unexplained muscle pain or tenderness, with a fever or “tea-coloured” urine, or reduced urination		✓	
† In young children †† In older children or adults			

*This is not a complete list of side effects. For any unexpected effects while taking Sandoz Valproic, contact your doctor or pharmacist.*

**HOW TO STORE IT**

Sandoz Valproic capsules 250 mg should be stored between 15 and 30°C, away from heat and humidity. Keep bottle tightly closed when not in use.

**Keep out of reach and sight of children.**

**Reporting Side Effects**

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on [Adverse Reaction Reporting](https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada/adverse-reaction-reporting.html) (<https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada/adverse-reaction-reporting.html>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

*NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.*

**MORE INFORMATION**

This document, plus the full product monograph prepared for health professionals, can be obtained by contacting the sponsor, Sandoz Canada Inc., at: 1-800-361-3062

or  
 by written request at:  
 110 Rue de Lauzon  
 Boucherville, (QC), Canada  
 J4B 1E6

or by e-mail at :  
 medinfo@sandoz.com  
 This leaflet was prepared by Sandoz Canada Inc.  
 Last revised: December 18, 2018