

PRODUCT MONOGRAPH

^{Pr}**TEVA-QUINAPRIL/HCTZ**

Quinapril Hydrochloride and Hydrochlorothiazide

10 mg/12.5 mg, 20 mg/12.5 mg and 20 mg/25 mg Tablets

Angiotensin Converting Enzyme Inhibitor/Diuretic

Teva Canada Limited
30 Novopharm Court
Toronto, Ontario
M1B 2K9

Date of Preparation:
December 17, 2018

Submission Control No: 103220, 161914, 222643

Table of Contents

PART I: HEALTH PROFESSIONAL INFORMATION..... 3
SUMMARY PRODUCT INFORMATION 3
INDICATIONS AND CLINICAL USE 3
CONTRAINDICATIONS 3
WARNINGS AND PRECAUTIONS 4
ADVERSE REACTIONS 12
DRUG INTERACTIONS 15
DOSAGE AND ADMINISTRATION 20
OVERDOSAGE 22
STORAGE AND STABILITY 25
DOSAGE FORMS, COMPOSITION AND PACKAGING 25

PART II: SCIENTIFIC INFORMATION 26
PHARMACEUTICAL INFORMATION 26
CLINICAL TRIALS 28
DETAILED PHARMACOLOGY 31
TOXICOLOGY 32
REFERENCES 40

PART III: CONSUMER INFORMATION..... 42

PrTEVA-QUINAPRIL/HCTZ

Quinapril Hydrochloride and Hydrochlorothiazide

PART I: HEALTH PROFESSIONAL INFORMATION

SUMMARY PRODUCT INFORMATION

Route of Administration	Dosage Form / Strength	Nonmedicinal Ingredients
Oral	Tablet / 10/12.5 mg, 20/12.5 mg and 20/25 mg	Calcium phosphate (dibasic) anhydrous, Crospovidone, Hydroxypropyl cellulose, Hydroxypropyl methylcellulose, Iron oxide red, Iron oxide yellow, Magnesium Carbonate, Magnesium Stearate, Polyethylene glycol, Povidone and Titanium Dioxide.

INDICATIONS AND CLINICAL USE

TEVA-QUINAPRIL/HCTZ (quinapril hydrochloride and hydrochlorothiazide) is indicated for the treatment of essential hypertension in patients for whom combination therapy is appropriate.

TEVA-QUINAPRIL/HCTZ is not indicated for initial therapy. Patients in whom quinapril and hydrochlorothiazide are initiated simultaneously can develop symptomatic hypotension (see **WARNINGS AND PRECAUTIONS, Cardiovascular – Hypotension and DRUG INTERACTIONS**).

Patients should be titrated on the individual drugs. If the fixed combination represents the dosage determined by this titration, the use of TEVA-QUINAPRIL/HCTZ may be more convenient in the management of patients. If during maintenance therapy dosage adjustment is necessary, it is advisable to use individual drugs.

Pediatrics (<18 years of age)

The safety and effectiveness of TEVA-QUINAPRIL/HCTZ in children have not been established, therefore use in this age group is not recommended.

CONTRAINDICATIONS

TEVA-QUINAPRIL/HCTZ is contraindicated in:

- Patients who are hypersensitive to the drug or to any ingredient in the formulation. For a complete listing, see the Dosage Forms, Composition and Packaging section of the product monograph.

- Patients with a history of angioedema related to previous treatment with an angiotensin converting enzyme (ACE) inhibitor (see **WARNINGS AND PRECAUTIONS, General, Angioedema**).
- Combination with sacubitril/valsartan due to increased risk of angioedema
- Patients hypersensitive to other sulfonamide-derived drugs because of the hydrochlorothiazide component.
- Patients with anuria
- Women who are pregnant, intend to become pregnant, or of childbearing potential who are not using adequate contraception (see **WARNINGS AND PRECAUTIONS, Special Populations, Pregnant Women** and **ADVERSE REACTIONS**).
- Nursing women (see **WARNINGS AND PRECAUTIONS, Special Populations, Nursing Women**).
- Combination with aliskiren-containing medicines in patients with
 - diabetes mellitus (type 1 or type 2),
 - moderate to severe kidney insufficiency (GFR < 60 mL/min/1.73 m²),
 - hyperkalemia (> 5 mMol/L) or
 - congestive heart failure who are hypotensive (see **WARNINGS AND PRECAUTIONS, Dual blockade of the Renin-Angiotensin System (RAS)** and **Renal Impairment**, and **DRUG INTERACTIONS, Aliskiren-containing medicines and Angiotensin receptor blockers (ARBs) or other ACE inhibitors**).
- Combination with angiotensin receptor blockers (ARBs) or other ACE inhibitors in patients with
 - diabetes with end organ damage,
 - moderate to severe kidney insufficiency (GFR < 60 mL/min/1.73m²),
 - hyperkalemia (> 5 mMol/L) or
 - congestive heart failure who are hypotensive (see **DRUG INTERACTIONS, Angiotensin receptor blockers (ARBs) or other ACE inhibitors**).

WARNINGS AND PRECAUTIONS

Serious Warnings and Precautions

When used in pregnancy, angiotensin converting enzyme (ACE) inhibitors can cause injury or even death of the developing fetus. When pregnancy is detected, TEVA-QUINAPRIL/HCTZ should be discontinued as soon as possible.

General

Angioedema

Angioedema has been reported in patients treated with ACE inhibitors, including quinapril. Angioedema associated with laryngeal involvement may be fatal. If laryngeal stridor or angioedema of the face, tongue, or glottis occurs, treatment with TEVA-QUINAPRIL/HCTZ should be discontinued immediately, the patient treated appropriately in accordance with accepted medical care, and carefully observed until the swelling disappears. In instances where swelling is confined to the face and lips, the condition generally resolves without treatment, although antihistamines may be useful in relieving symptoms. Where there is involvement of the tongue,

glottis or larynx, likely to cause airway obstruction, appropriate therapy (including but not limited to 0.3 to 0.5 mL of subcutaneous epinephrine solution 1:1000) should be administered promptly (see **ADVERSE REACTIONS**).

The incidence of angioedema during ACE inhibitor therapy has been reported to be higher in black than in non-black patients.

Patients taking a concomitant mTOR inhibitor (e.g. temsirolimus), DPP-4 inhibitor (e.g. sitagliptin) or neutral endopeptidase (NEP) inhibitor may be at increased risk for angioedema.

Caution should be used when either initiating ACE inhibitor therapy in patients already taking a mTOR inhibitor, DPP-4 inhibitor or NEP inhibitor or vice versa (see **DRUG INTERACTIONS**).

Patients with a history of angioedema related or unrelated to ACE inhibitor therapy may be at increased risk of angioedema while receiving an ACE inhibitor (see **CONTRAINDICATIONS**).

Cardiovascular

Dual Blockade of the Renin-Angiotensin System (RAS)

Dual blockade of the Renin-Angiotensin System (RAS): There is evidence that co-administration of angiotensin converting enzyme (ACE) inhibitors, such as TEVA-QUINAPRIL/HCTZ, or of angiotensin receptor antagonists (ARBs) with aliskiren increases the risk of hypotension, syncope, stroke, hyperkalemia and deterioration of renal function, including acute renal failure, in patients with diabetes mellitus (type 1 or type 2) and/or moderate to severe kidney insufficiency (GFR < 60 mL/min/1.73 m²). Therefore, the use of TEVA-QUINAPRIL/HCTZ in combination with aliskiren-containing drugs is contraindicated in these patients (see **CONTRAINDICATIONS**).

Further, co-administration of ACE inhibitors, including TEVA-QUINAPRIL/HCTZ, with other agents blocking the RAS, such as ARBs or aliskiren-containing drugs, is generally not recommended in other patients, since such treatment has been associated with an increased incidence of severe hypotension, acute renal failure, and hyperkalemia. Administration should be limited to individually defined cases with close monitoring of renal function and blood potassium levels (see **CONTRAINDICATIONS**).

Hypotension

Symptomatic hypotension has occurred after administration of quinapril, usually after the first or second dose or when the dose was increased. It is more likely to occur in patients who are volume depleted by diuretic therapy, dietary salt restriction, dialysis, diarrhea, or vomiting. In patients with ischemic heart or cerebrovascular disease, an excessive fall in blood pressure could result in a myocardial infarction or cerebrovascular accident (see **ADVERSE REACTIONS**). Because of the potential fall in blood pressure in these patients, therapy with TEVA-QUINAPRIL/HCTZ should be started under close medical supervision. Such patients should be followed closely for the first weeks of treatment and whenever the dose of TEVA-QUINAPRIL/HCTZ is increased. In patients with severe congestive heart failure, with or without associated renal insufficiency, excessive hypotension has been observed and may be associated with oliguria and/or progressive azotemia, and rarely with acute renal failure and/or death.

If hypotension occurs, the patient should be placed in supine position and, if necessary, receive an intravenous infusion of 0.9% sodium chloride. A transient hypotensive response is not a contraindication to further doses which usually can be given without difficulty once the blood pressure has increased after volume expansion. If symptoms persist, the dosage should be reduced or the drug discontinued.

Valvular Stenosis

There is concern on theoretical grounds that patients with aortic stenosis might be at particular risk of decreased coronary perfusion when treated with vasodilators because they do not develop as much afterload reduction.

Endocrine and Metabolism

Initial and periodic determination of serum electrolytes should be performed at appropriate intervals to detect possible electrolyte imbalance.

Hyperkalemia/Hypokalemia

Quinapril: Elevated serum potassium (> 5.7 mMol/L) was observed in approximately 2% of patients receiving quinapril. In most cases these were isolated values which resolved despite continued therapy. Hyperkalemia was a cause of discontinuation of therapy in < 0.1% of hypertensive patients. Risk factors for the development of hyperkalemia include renal insufficiency, diabetes mellitus, and the concomitant use of agents to treat hypokalemia, potassium-sparing diuretics, potassium supplements, potassium-containing salt substitutes, or other drugs known to raise serum potassium levels (see **WARNINGS AND PRECAUTIONS, Monitoring and Laboratory Tests, Serum Electrolytes, ADVERSE REACTIONS and DRUG INTERACTIONS, Agents Increasing Serum Potassium, Trimethoprim-containing products**). The addition of a potassium-sparing diuretic to TEVA-QUINAPRIL/HCTZ, which contains a diuretic, is not recommended.

Hydrochlorothiazide: Treatment with thiazide diuretics has been associated with hypokalemia. Hypokalemia can also sensitize or exaggerate the response of the heart to the toxic effects of digitalis. The risk of hypokalemia is greatest in patients with cirrhosis of the liver, in patients experiencing a brisk diuresis, in patients who are receiving inadequate oral intake of electrolytes, and in patients receiving concomitant therapy with corticosteroids or ACTH or with other drugs known to increase the risk of hypokalemia induced by thiazide diuretics (e.g. aminoglycoside antibiotics, cisplatin, foscarnet, amphotericin B and loop diuretics (furosemide)).

Quinapril/Hydrochlorothiazide: The opposite effects of hydrochlorothiazide and quinapril on serum potassium may approximately balance each other in many patients so that no net effect will be seen. In other patients, one or the other effect may be dominant.

Other electrolytes imbalances

Hydrochlorothiazide: In addition to hypokalemia, treatment with thiazide diuretics has also been associated with hyponatremia and hypochloremic alkalosis. These disturbances have sometimes been manifest as one or more of the following: dryness of mouth, thirst, weakness, lethargy,

drowsiness, restlessness, muscle pain or cramps, muscular fatigue, hypotension, oliguria, tachycardia, nausea, confusion, seizures and vomiting.

Chloride deficits secondary to thiazide therapy are generally mild and require specific treatment only under extraordinary circumstances (e.g. in liver disease or renal disease). Dilutional hyponatremia may occur in edematous patients, especially in hot weather; appropriate therapy is water restriction rather than administration of salt, except when the hyponatremia is life threatening. In actual salt depletion, replacement of salt is the therapy of choice.

Thiazides may decrease calcium excretion. Thiazides may cause intermittent and slight elevation of serum calcium in the absence of known disorders of calcium metabolism. Marked hypercalcemia may be evidence of hidden hypoparathyroidism. In a few patients on prolonged thiazide therapy, pathological changes in the parathyroid gland have been observed, with hypercalcemia and hypophosphatemia. More serious complications of hyperparathyroidism (renal lithiasis, bone resorption, and peptic ulceration) have not been seen. Thiazides should be discontinued before performing tests for parathyroid function.

Thiazides increase the urinary excretion of magnesium, and hypomagnesaemia may result.

Hypoglycemia/Hyperglycemia and Diabetes

Quinapril: ACE inhibitors may reduce insulin resistance and may lead to hypoglycemia in diabetic patients on insulin or oral hypoglycemic agents; closer monitoring of diabetic patients may be required.

Hydrochlorothiazide: Thiazide-induced hyperglycemia may compromise blood sugar control. Depletion of serum potassium augments glucose intolerance. Monitor glycemic control, supplement potassium, if necessary, to maintain appropriate serum potassium levels, and adjust diabetes medications as required (see **DRUG INTERACTIONS, Antidiabetic agents (e.g. insulin, oral hypoglycemic agents, sitagliptin)**).

Overt diabetes may be precipitated in susceptible individuals.

Other metabolic parameters

Hyperuricemia may occur, or acute hyperuricemia may be precipitated, in certain patients receiving thiazide therapy.

Increase in cholesterol, triglyceride and glucose levels may be associated with thiazide diuretic therapy.

Thiazides may decrease serum PBI levels without signs of thyroid disturbance.

Hematologic

Neutropenia/Agranulocytosis

Agranulocytosis and bone marrow depression have been caused by ACE inhibitors.

Agranulocytosis did occur during quinapril treatment in one patient with a history of neutropenia

during previous captopril therapy. Periodic monitoring of white blood cell counts should be considered, especially in patients with collagen vascular disease and/or renal disease.

Hepatic

Impairment of Liver Function

Hepatitis (hepatocellular and/or cholestatic), elevations of liver enzymes and/or serum bilirubin have occurred during therapy with other ACE inhibitors in patients with or without pre-existing liver abnormalities. In most cases the changes were reversed on discontinuation of the drug.

Elevations of liver enzymes and/or serum bilirubin have been reported for quinapril hydrochloride and hydrochlorothiazide tablets (see **ADVERSE REACTIONS**). Should the patient receiving TEVA-QUINAPRIL/HCTZ experience any unexplained symptoms particularly during the first weeks or months of treatment, it is recommended that a full set of liver function tests and any other necessary investigation be carried out. Discontinuation of TEVA-QUINAPRIL/HCTZ should be considered when appropriate.

There are no adequate studies in patients with cirrhosis and/or liver dysfunction. TEVA-QUINAPRIL/HCTZ should be used with particular caution in patients with pre-existing liver abnormalities. In such patients baseline liver function tests should be obtained before administration of the drug and close monitoring of response and metabolic effects should apply.

Patients with Impaired Liver Function

TEVA-QUINAPRIL/HCTZ should be used with caution in patients with impaired hepatic function or progressive liver disease, since minor alterations of fluid and electrolyte balance may precipitate hepatic coma. Also, since the metabolism of quinapril to quinaprilat is normally dependent upon hepatic esterase, patients with impaired liver function could develop markedly elevated plasma levels of quinapril.

Immune

Anaphylactoid Reactions during Membrane Exposure

Anaphylactoid reactions have been reported in patients dialysed with high-flux membranes (e.g.: polyacrylonitrile [PAN]) and treated concomitantly with an ACE inhibitor. Dialysis should be stopped immediately if symptoms such as nausea, abdominal cramps, burning, angioedema, shortness of breath and severe hypotension occur. Symptoms are not relieved by antihistamines. In these patients consideration should be given to using a different type of dialysis membrane or a different class of antihypertensive agent.

Anaphylactoid Reactions during LDL Apheresis

Rarely, patients receiving ACE inhibitors during low density lipoprotein apheresis with dextran sulfate have experienced life-threatening anaphylactoid reactions. These reactions were avoided by temporarily withholding the ACE inhibitor therapy prior to each apheresis.

Anaphylactoid Reactions during Desensitization

There have been isolated reports of patients experiencing sustained life threatening anaphylactoid reactions while receiving ACE inhibitors during desensitizing treatment with hymenoptera (bees, wasps) venom. In the same patients, these reactions have been avoided when ACE inhibitors were temporarily withheld for at least 24 hours, but they have reappeared upon inadvertent rechallenge to an ACE inhibitor.

Hypersensitivity to Hydrochlorothiazide

Sensitivity reactions to hydrochlorothiazide may occur in patients with or without a history of allergy or bronchial asthma.

Nitritoid Reactions-Gold

Nitritoid reactions (symptoms include facial flushing, nausea, vomiting and symptomatic hypotension) have been reported rarely in patients on therapy with injectable gold (sodium aurothiomalate) and concomitant ACE inhibitor therapy including TEVA-QUINAPRIL/HCTZ (see **DRUG INTERACTIONS**).

Systemic Lupus Erythematosus

Thiazide diuretics have been reported to cause exacerbation or activation of systemic lupus erythematosus.

Ophthalmologic

Acute Myopia and Secondary Angle-Closure Glaucoma related to Hydrochlorothiazide

Hydrochlorothiazide, a sulfonamide, can cause an idiosyncratic reaction, resulting in acute transient myopia and acute angle-closure glaucoma. Symptoms include acute onset of decreased visual acuity or ocular pain and typically occur within hours to weeks of drug initiation. Untreated acute angle-closure glaucoma can lead to permanent vision loss.

The primary treatment is to discontinue hydrochlorothiazide as rapidly as possible. Prompt medical or surgical treatments may need to be considered if the intraocular pressure remains uncontrolled. Risk factors for developing acute angle-closure glaucoma may include a history of sulfonamide or penicillin allergy.

Peri-Operative Considerations

Surgery/Anaesthesia

In patients undergoing major surgery or during anaesthesia with agents that produce hypotension, ACE inhibitors will block angiotensin II formation secondary to compensatory renin release. If hypotension occurs and is considered to be due to this mechanism, it can be corrected by volume expansion.

Renal

Azotemia

Azotemia may be precipitated or increased by hydrochlorothiazide. Cumulative effects of the drug may develop in patients with impaired renal function. If increasing azotemia and oliguria occur,

TEVA-QUINAPRIL/HCTZ should be discontinued.

Renal Impairment

The use of ACE inhibitors, including TEVA-QUINAPRIL/HCTZ, with ARBs or aliskiren-containing drugs is contraindicated in patients with moderate to severe kidney insufficiency (GFR < 60 mL/min/1.73m²) (see **CONTRAINDICATIONS** and **DRUG INTERACTIONS, Aliskiren-containing medicines and Angiotensin receptor blockers (ARBs) or other ACE inhibitors**).

As a consequence of inhibiting the renin-angiotensin-aldosterone system (RAAS), changes in renal function have been seen in susceptible individuals. In patients whose renal function may depend on the activity of the RAAS, such as patients with bilateral renal artery stenosis, unilateral renal artery stenosis to a solitary kidney, or severe congestive heart failure, treatment with agents that inhibit this system has been associated with oliguria, progressive azotemia, and rarely, acute renal failure and/or death. In susceptible patients, concomitant diuretic use may further increase risk (see **ADVERSE REACTIONS** and **DOSAGE ADMINISTRATION**).

Use of TEVA-QUINAPRIL/HCTZ should be followed by the appropriate assessment of renal function (see **ADVERSE REACTIONS** and **DOSAGE AND ADMINISTRATION**).

Thiazides may not be appropriate diuretics for use in patients with renal impairment and are ineffective at creatinine clearance values of ≤ 30 mL/min (i.e. moderate or severe renal insufficiency) (see **ADVERSE REACTIONS** and **DOSAGE AND ADMINISTRATION**).

Respiratory

Cough

A dry, persistent cough, which usually disappears only after withdrawal or lowering of the dose of quinapril has been reported. Such possibility should be considered as part of the differential diagnosis of the cough.

Special Populations

Pregnant Women

Quinapril is contraindicated in pregnancy (see **CONTRAINDICATIONS** and **ADVERSE REACTIONS**). ACE inhibitors can cause fetal and neonatal morbidity and mortality when administered to pregnant women. Several dozen cases have been reported in the world literature. When pregnancy is detected, TEVA-QUINAPRIL/ HCTZ should be discontinued as soon as possible.

The use of ACE inhibitors during the second and third trimesters of pregnancy has been associated with fetal and neonatal injury including hypotension, neonatal skull hypoplasia, anuria, reversible or irreversible renal failure, and death. Oligohydramnios has also been reported, presumably resulting from decreased fetal renal function; oligohydramnios in this setting has been associated with fetal limb contractures, craniofacial deformation, and hypoplastic lung development.

Prematurity, and patent ductus arteriosus and other structural cardiac malformations, as well as neurological malformations, have also been reported following exposure in the first trimester of pregnancy.

Infants with a history of *in utero* exposure to ACE inhibitors should be closely observed for hypotension, oliguria, and hyperkalemia. If oliguria occurs, attention should be directed toward support of blood pressure and renal perfusion. Exchange transfusion or dialysis may be required as a means of reversing hypotension and/or substituting for impaired renal function; however, limited experience with those procedures has not been associated with significant clinical benefit.

If oligohydramnios is observed, a non-stress test (NST), and/or a biophysical profiling (BPP) may be appropriate, depending upon the week of pregnancy. If concerns regarding fetal well-being still persist, a contraction stress test (CST) should be considered. Patients and physicians should be aware, however, that oligohydramnios may not appear until after the fetus has sustained irreversible injury.

Thiazides cross the placental barrier and appear in cord blood. Although studies in humans have not been done, effects to the fetus may include fetal or neonatal jaundice, thrombocytopenia, and possibly other adverse reactions which have occurred in the adult.

Animal Data: No fetotoxic or teratogenic effects were observed in rats at quinapril doses as high as 300 mg/kg/day (180x maximum daily human dose), despite maternal toxicity at 150 mg/kg/day. Offspring body weights were reduced in rats treated late in gestation and during lactation with doses of ≥ 25 mg/kg/day. Quinapril hydrochloride was not teratogenic in rabbits; however, maternal and embryo toxicity were seen in some rabbits at doses as low as 0.5 mg/kg/day and 1 mg/kg/day, respectively.

No adverse effects on fertility or reproduction were observed in rats at quinapril dose levels ≤ 100 mg/kg/day (60x maximum daily human dose).

Nursing Women

The presence of concentrations of ACE inhibitor has been reported in human milk. Thiazides also appear in human milk. The use of TEVA-QUINAPRIL/HCTZ is contraindicated during breast-feeding (see **CONTRAINDICATIONS**).

Pediatrics (< 18 years of age)

The safety and effectiveness of TEVA-QUINAPRIL/HCTZ in children have not been established, therefore use in this age group is not recommended.

Monitoring and Laboratory Tests

Serum Electrolytes: Variations of serum electrolytes levels have been observed with ACCURETIC. Initial and periodic determination of serum electrolytes should be performed at appropriate intervals to detect possible electrolyte imbalance (see **CONTRAINDICATIONS** and **WARNINGS and PRECAUTIONS**).

Creatinine and Blood Nitrogen: Increases (>1.25x ULN) in serum creatinine and blood urea nitrogen (BUN) were observed in 3% and 4% respectively, of patients treated with quinapril and hydrochlorothiazide tablets (see **WARNINGS AND PRECAUTIONS** and **DOSAGE AND ADMINISTRATION**).

Hepatic: Elevations of liver enzymes and/or serum bilirubin have occurred in patients receiving quinapril and hydrochlorothiazide tablets. If a patient receiving TEVA-QUINAPRIL/HCTZ experience any unexplained symptoms, particularly during the first weeks or months of treatment, a full set of liver function tests and any other investigation should be carried out. Discontinuation of TEVA-QUINAPRIL/HCTZ should be considered when appropriate. In patients with pre-existing liver abnormalities, baseline liver function tests should be obtained before administration of the drug. The response and metabolic effects should be closely monitored (see **WARNINGS AND PRECAUTIONS**).

Glucose: Elevations in glucose values have occurred. Monitor glycemic control, supplement potassium, if necessary, to maintain appropriate serum potassium levels, and adjust diabetes medications as required (see **WARNINGS AND PRECAUTIONS**, **Endocrine and Metabolism**, **Hypoglycemia/Hyperglycemia and Diabetes** and **DRUG INTERACTIONS, Antidiabetic agents (e.g. insulin, oral hypoglycemic agents, sitagliptin)**).

Triglyceride: Elevations in triglyceride values have occurred (see **WARNINGS AND PRECAUTIONS**, **Endocrine and Metabolism**, **Other metabolic parameters**).

Serum Uric Acid: Elevations in serum uric acid values have occurred (see **WARNINGS AND PRECAUTIONS**, **Endocrine and Metabolism**, **Other metabolic parameters**).

Hematology: Possibly clinically important increases and decreases in hematology parameters have occurred. Periodic monitoring of white blood cell counts should be considered; especially in patients with collagen vascular disease and/or renal disease (see **WARNINGS AND PRECAUTIONS**, **Hematologic**, **Neutropenia/Agranulocytosis**).

Other laboratory test values with clinically important deviations during controlled and uncontrolled trials included: Magnesium, Cholesterol, PBI, Parathyroid Function Tests and Calcium (see **WARNINGS AND PRECAUTIONS**, **Endocrine and Metabolism**, **Other electrolytes imbalances and Other metabolic parameters**).

ADVERSE REACTIONS

Clinical Trial Adverse Drug Reactions

Because clinical trials are conducted under very specific conditions the adverse reaction rates observed in the clinical trials may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse drug reaction information from clinical trials is useful for identifying drug-related adverse events and for approximating rates.

Quinapril hydrochloride and hydrochlorothiazide was evaluated for safety in 1571 patients with essential hypertension, including 943 patients in controlled studies (see Table 1), 345 patients in placebo-controlled trials, and 517 patients who were treated with quinapril hydrochloride/hydrochlorothiazide for ≥ 1 year. Adverse reactions were limited to those reported previously with quinapril or hydrochlorothiazide (HCTZ) when used separately for the treatment of hypertension.

Serious or clinically significant adverse reactions observed in $< 0.2\%$ of patients treated with quinapril and HCTZ were: hematemesis, gout, syncope and angioedema. Therapy was discontinued in 2.1% of patients due to an adverse event (AE). Headache (0.5%) and dizziness (0.3%) were the most frequent reasons for withdrawal.

The most frequent adverse experiences in controlled trials were headache (6.7%), dizziness (4.8%), cough (3.2%), and fatigue (2.9%). The cough is characteristically non-productive, persistent and resolves after discontinuation of therapy (see **WARNINGS AND PRECAUTIONS, Respiratory, Cough**).

Table 1: Adverse Events in $\geq 1\%$ of Quinapril/Hydrochlorothiazide Patients in Controlled Clinical Studies

Adverse Events	<u>Quinapril / HCTZ</u> N = 943 (% Patients)	<u>Quinapril</u> N = 799 (% Patients)
Body as a Whole		
Asthenia	1.1	1.2
Fatigue	2.9	2.0
Headache	6.7	4.8
Back pain	1.5	0.7
Chest pain	1.0	1.2
Viral infection	1.9	2.0
Cardiovascular System		
Vasodilatation	1.0	0.4
Digestive System		
Dyspepsia	1.2	1.9
Nausea and/or vomiting	1.8	2.0
Diarrhea	1.4	1.7
Abdominal pain	1.7	1.6
Musculoskeletal System		
Myalgia	2.4	0.9
Nervous System		
Dizziness	4.8	2.7
Insomnia	1.2	1.5
Somnolence	1.2	0.9
Vertigo	1.0	0.3

Adverse Events	<u>Quinapril / HCTZ</u> N = 943 (% Patients)	<u>Quinapril</u> N = 799 (% Patients)
Respiratory System		
Pharyngitis	1.1	1.4
Rhinitis	2.0	3.0
Bronchitis	1.2	1.3
Coughing	3.2	2.7
Upper respiratory infection	1.3	1.1

Clinical AEs regardless of relationship to therapy, occurring in $\geq 0.5\%$ to $< 1.0\%$ of patients treated with quinapril plus HCTZ in controlled and uncontrolled trials and less frequent clinically significant events seen in clinical trials or in post marketing experience included:

- Cardiovascular:** Hypotension, palpitations, tachycardia
- Gastrointestinal:** Dry mouth or throat, flatulence, pancreatitis
- Respiratory:** Dyspnea, sinusitis
- Integumentary:** Alopecia, erythema multiforme, exfoliative dermatitis, pemphigus, pruritus, rash
- Nervous/Psychiatric:** Nervousness, paresthesia
- Urogenital:** Impotence, urinary tract infection
- Other:** Arthralgia, hemolytic anemia, peripheral edema

Rare AEs, not listed above, which have been reported with either HCTZ, quinapril, or the combination include:

- Cardiovascular:** Atrial flutter, cerebrovascular accident, heart arrest, heart failure, myocardial ischemia, necrotizing angiitis, transient ischemic attack, vasodilation. Orthostatic hypotension may occur, especially in elderly patients with reduced plasma volume, and may be potentiated by alcohol, barbiturates, or narcotics
- Gastrointestinal:** Anorexia, bloody stools, constipation, cramping, gastric irritation, GI hemorrhage, jaundice (intrahepatic cholestatic), pancreatitis, sialadenitis
- Eye Disorders:** Acute myopia and acute angle closure glaucoma (see **WARNINGS AND PRECAUTIONS, Ophthalmologic, Acute Myopia and Acute angle closure glaucoma**)
- Respiratory:** Respiratory distress including pneumonitis, asthma, hoarseness

<i>Integumentary:</i>	Eczema, photosensitivity, rash, Stevens Johnson syndrome, urticaria
<i>Nervous System:</i>	Amnesia, anxiety, confusion, facial paralysis, paresthesias, polyneuritis, xanthopsia
<i>Hematological:</i>	Agranulocytosis, aplastic anemia, hemolytic anemia, leukopenia, purpura, thrombocytopenia
<i>Urogenital:</i>	Dysuria, glycosuria, hematuria, impaired renal function, polyuria
<i>Special Senses:</i>	Taste disturbance, tinnitus, transient blurred vision
<i>Congenital and familial/genetic disorders:</i>	Fetal/neonatal injury including: anuria, hypotension, oligohydramnios, skull hypoplasia, reversible or irreversible renal failure and death (see CONTRAINDICATIONS, WARNINGS AND PRECAUTIONS, Pregnant Women)
<i>Other:</i>	Allergy, anaphylactic reactions, arthritis, chill, dehydration, face edema, fever, fracture, muscle spasm, restlessness, weakness, weight increase
<i>Laboratory Deviations:</i>	Azotemia, hyperglycemia, hyperuricemia, transient hyperlipidemia, WBC decreased

DRUG INTERACTIONS

Drug-Drug Interactions

Proper Name	Reference	Effect	Clinical comment
Agents Affecting Sympathetic Activity	--		Agents affecting sympathetic activity (e.g. ganglionic blocking agents or adrenergic neuron blocking agents) may be used with caution. Beta-adrenergic blocking drugs add some further antihypertensive effect to quinapril.
Agents Increasing Serum Potassium	--	Since quinapril decreases aldosterone production, elevation of serum potassium may occur.	Potassium sparing diuretics such as spironolactone, triamterene or amiloride, potassium supplements or other drugs known to raise serum potassium levels should be given with caution and frequent monitoring of serum potassium, since they may lead to a significant increase in serum potassium. Salt substitutes which contain potassium should also be used with caution.

Proper Name	Reference	Effect	Clinical comment
			Since quinapril/HCTZ contains a diuretic, the addition of a potassium-sparing diuretic is not recommended.
Alcohol, barbiturates or narcotics	C	Potential of orthostatic hypotension may occur in the presence of hydrochlorothiazide.	Avoid alcohol, barbiturates or narcotics, especially with initiation of therapy.
Aliskiren-containing medicines	CT	Dual blockade of the renin-angiotensin-aldosterone system by combining an ACE inhibitor with aliskiren-containing medicines is not recommended since there is an increased risk of hypotension, syncope, stroke, hyperkalemia and changes in renal function, including renal failure.	The use of TEVA-QUINAPRIL/HCTZ in combination with aliskiren-containing medication is contraindicated in patients with: <ul style="list-style-type: none"> • diabetes mellitus (type 1 or type 2), • moderate to severe kidney insufficiency (GFR < 60 mL/min/1.73 m²), • hyperkalemia (> 5mMol/L) or • congestive heart failure who are hypotensive. It is not recommended in other patients (see CONTRAINDICATIONS; WARNINGS AND PRECAUTIONS, Dual Blockade of the Renin-Angiotensin System (RAS))
Amphotericin B	T	Amphotericin B increases the risk of hypokalemia induced by thiazide diuretics.	Monitor serum potassium level.
Angiotensin receptor blockers (ARBs) or other ACE inhibitors	CT	Dual blockade of the renin-angiotensin-aldosterone system by combining an ACE inhibitor with ARBs or other ACE inhibitors is not recommended since there is an increased risk of hypotension, syncope, stroke, hyperkalemia and changes in renal function, including renal failure.	The use of TEVA-QUINAPRIL/HCTZ in combination with ARBs or other ACE inhibitors is contraindicated in patients with <ul style="list-style-type: none"> • diabetes with end organ damage, • moderate to severe kidney insufficiency (GFR < 60 mL/min/1.73 m²), • hyperkalemia (> 5mMol/L) or • congestive heart failure who are hypotensive. It is not recommended in other patients (see CONTRAINDICATIONS, WARNINGS AND PRECAUTIONS, Cardiovascular, Dual blockade of the Renin-Angiotensin System (RAS))
Anion Exchange Resins - Bile acid sequestrants (e.g. cholestyramine)	CT	Bile acid sequestrants bind thiazide diuretics in the gut and impair gastrointestinal absorption by 43-85%. Administration of thiazide 4 hours after a bile acid sequestrant reduced absorption of hydrochlorothiazide by 30-35%.	Give thiazide 2-4 hours before or 6 hours after the bile acid sequestrant. Maintain a consistent sequence of administration. Monitor blood pressure, and increase dose for thiazide, if necessary.
Antidiabetic agents (e.g. insulin, oral hypoglycemic agents, sitagliptin)	CT	ACE inhibitors may reduce insulin resistance and may lead to hypoglycemia in diabetic patients on insulin or oral hypoglycemic agents.	Monitor closely diabetic patients (see WARNINGS AND PRECAUTIONS, Endocrine and Metabolism, Hypoglycemia and Diabetes).
		Thiazide-induced hyperglycemia	Monitor glycemic control, supplement

Proper Name	Reference	Effect	Clinical comment
		may compromise blood sugar control. Depletion of serum potassium augments glucose intolerance.	potassium if necessary, to maintain appropriate serum potassium levels, and adjust diabetes medications as required (see WARNINGS AND PRECAUTIONS, Endocrine and Metabolism, Hypoglycemia and Diabetes).
		Patients taking concomitant DPP-4 inhibitor therapy may be at increased risk for angioedema.	Caution should be used when either initiating ACE inhibitor therapy in patients already taking a DPP-4 inhibitor or <i>vice versa</i> (see WARNINGS AND PRECAUTIONS, General, Angioedema).
Antineoplastic drugs, including cyclophosphamide, methotrexate and mTOR inhibitors (e.g. temsirolimus, everolimus)	C, CT	Concomitant use of thiazide diuretics may reduce renal excretion of cytotoxic agents and enhance their myelosuppressive effects.	Hematological status should be closely monitored in patients receiving this combination. Dose adjustment of cytotoxic agents may be required.
		Patients taking concomitant mTOR inhibitor therapy may be at increased risk for angioedema.	Caution should be used when either initiating ACE inhibitor therapy in patients already taking an mTOR inhibitor or <i>vice versa</i> (see WARNINGS AND PRECAUTIONS, General, Angioedema).
Calcium and vitamin D supplements	C	Thiazides decrease renal excretion of calcium and increase calcium release from bone.	Monitor serum calcium, especially with concomitant use of high doses of calcium supplements. Dose reduction or withdrawal of calcium and/or vitamin D supplements may be necessary.
Carbamazepine	C	Carbamazepine may cause clinically significant hyponatremia. Concomitant use with thiazide diuretics may potentiate hyponatremia.	Monitor serum sodium levels. Use with caution.
Concomitant Diuretic Therapy	--	Patients concomitantly taking ACE inhibitors and diuretics, and especially those in whom diuretic therapy was recently instituted, may occasionally experience an excessive reduction of blood pressure after initiation of therapy.	The possibility of hypotensive effects after the first dose of quinapril can be minimized by either discontinuing the diuretic or increasing the salt intake (except in patients with heart failure) prior to initiation of treatment with quinapril. If it is not possible to discontinue the diuretic, the starting dose of quinapril should be reduced and the patient should be closely observed for several hours following initial dose and until blood pressure has stabilized. (See WARNINGS AND PRECAUTIONS, and DOSAGE AND ADMINISTRATION).
Corticosteroids, and adrenocorticotrophic hormone (ACTH)	T	Intensified electrolyte depletion, particularly hypokalemia, may occur when administered with hydrochlorothiazide.	Monitor serum potassium, and adjust medications, as required.
Digoxin	CT	Thiazide-induced electrolyte	Concomitant administration of

Proper Name	Reference	Effect	Clinical comment
		disturbances, i.e. hypokalemia, hypomagnesemia, increase the risk of digoxin toxicity, which may lead to fatal arrhythmic events.	hydrochlorothiazide and digoxin requires caution. Monitor electrolytes and digoxin levels closely. Supplement potassium or adjust doses of digoxin or thiazides, as required.
Drugs that alter GI motility , i.e., anticholinergic agents, such as atropine and prokinetic agents, such as metoclopramide, domperidone	CT, T	Bioavailability of thiazide diuretics may be increased by anticholinergic agents due to a decrease in gastrointestinal motility and gastric emptying. Conversely, prokinetic drugs may decrease the bioavailability of thiazide diuretics.	Dose of adjustment of thiazide may be required.
Gold	C	Nitritoid reactions (symptoms include facial flushing, nausea, vomiting and hypotension) have been reported rarely in patients on therapy with injectable gold (sodium aurothiomalate) and concomitant ACE inhibitor therapy including TEVA-QUINAPRIL/HCTZ.	
Gout medications (allopurinol, uricosurics, xanthine oxidase inhibitors)	T, RC	Thiazide-induced hyperuricemia may compromise control of gout by allopurinol and probenecid. The co-administration of hydrochlorothiazide and allopurinol may increase the incidence of hypersensitivity reactions to allopurinol.	Dose adjustment of gout medications may be required.
Lithium	CT	Thiazide diuretic agents and ACE inhibitors reduce the renal clearance of lithium and increase the risk of lithium toxicity.	Concomitant use of thiazide diuretics or ACE inhibitors with lithium is generally not recommended. If such use is deemed necessary, reduce lithium dose by 50% and monitor lithium levels closely.
Neutral endopeptidase inhibitor	---	Patients taking concomitant neutral endopeptidase inhibitor may be at increased risk for angioedema.	Caution should be used when either initiating ACE inhibitor therapy in patients already taking a neutral endopeptidase inhibitor or <i>vice versa</i> (see WARNINGS AND PRECAUTIONS, General, Angioedema).
Non-Steroidal Anti-Inflammatory Drugs (NSAID) including selective cyclooxygenase-2 inhibitors (COX-2 inhibitors)	CT	There are two types of interaction between TEVA-QUINAPRIL/HCTZ and NSAIDs: Interaction with ACE-Inhibitor component of TEVA-QUINAPRIL/HCTZ: In patients who are elderly, volume-depleted (including those on diuretic therapy), or with compromised renal function, co-administration of NSAIDs, including selective COX-2 inhibitors, with ACE inhibitors,	If combination use is necessary, monitor renal function, serum potassium, and blood pressure closely. Dose adjustment may be required.

Proper Name	Reference	Effect	Clinical comment
		<p>including quinapril, may result in deterioration of renal function, including possible acute renal failure. These effects are usually reversible.</p> <p>The antihypertensive effect of ACE inhibitors, including quinapril may be attenuated by NSAIDs.</p> <p>Interaction with Diuretic component of TEVA-QUINAPRIL/HCTZ: In some patients, the administration of a NSAID agent can reduce the diuretic, natriuretic, and antihypertensive effects of loop, potassium-sparing, and thiazide diuretics.</p> <p>NSAID-induced inhibition of renal prostaglandins leading to decreases of renal blood flow, along with thiazide-induced decreases in GFR may lead to acute renal failure. Patients with heart failure may be at particular risk.</p>	
Other Antihypertensive Agents	CT	Hydrochlorothiazide may potentiate the action of other antihypertensive drugs (e.g. guanethidine, methyl dopa, beta-blockers, vasodilators, calcium channel blockers, ACEI, ARB, ARB, and direct rennin inhibitors)	
Pressor Amines (e.g. noradrenaline)	--	Possible decreased response to pressor amines may occur in the presence of a thiazide diuretic, but is not sufficient to preclude their use.	
Selective serotonin reuptake inhibitors (SSRIs, e.g. citalopram, escitalopram, sertraline)	T, C	Concomitant use with thiazide diuretics may potentiate hyponatremia.	Monitor serum sodium levels. Use with caution.
Sirolimus (immune-suppressant mTOR inhibitor)	CT	Organ transplant patients taking concomitant sirolimus therapy may be at increased risk for angioedema.	Caution should be used when either initiating ACE inhibitor therapy in patients already taking sirolimus or <i>vice versa</i> (see WARNINGS AND PRECAUTIONS, General, Angioedema).
Skeletal muscle relaxants of the curare family , e.g.,	C	Thiazide drugs may increase responsiveness of some skeletal muscle relaxants, such as curare	

Proper Name	Reference	Effect	Clinical comment
tubocurare		derivatives.	
Tetracycline	--	Concomitant administration of tetracycline with quinapril reduced the absorption of tetracycline in healthy volunteers (by 28-37%) due to the presence of magnesium carbonate as an excipient in the formulation.	This interaction should be considered with concomitant use of TEVA-QUINAPRIL/HCTZ and tetracycline or other drugs which interact with magnesium.
Topiramate	CT	Additive hypokalemia. Possible thiazide-induced increase in topiramate serum concentrations.	Monitor serum potassium and topiramate levels. Use potassium supplements, or adjust topiramate dose as necessary.
Trimethoprim-containing products (sulfamethoxazole/ trimethoprim)	C	In patients who are elderly or have compromised renal function, co-administration of an ACE inhibitor with sulfamethoxazole/trimethoprim has been associated with severe hyperkalemia, likely due to the hyperkalemic effects of trimethoprim.	Quinapril/HCTZ and trimethoprim-containing products should only be co-administered with caution and with appropriate monitoring of serum potassium.

Legend: C=Case Study; RCS=Retrospective Cohort Study; CT=Clinical Trial; T=Theoretical

Drug-Food Interactions

The rate and extent of quinapril absorption are diminished moderately (approximately 25-30%) when administered during a high-fat meal. However, no effect on quinapril absorption occurs when taken during a regular meal.

Drug-Herb Interactions

Interactions with herbal products have not been established.

Drug-Laboratory Test Interactions

Interactions with laboratory products/methods have not been established.

Drug-Lifestyle Interactions

Lifestyle interactions have not been established.

DOSAGE AND ADMINISTRATION

Dosing Considerations

Dosage must be individualized. The fixed combination is not for initial therapy. The dose of TEVA-QUINAPRIL/HCTZ (quinapril hydrochloride and hydrochlorothiazide) should be determined by titration of the individual components.

Once the patient has been successfully titrated with the individual components as described below, TEVA-QUINAPRIL/HCTZ may be substituted if the titrated doses and dosing schedule can be achieved by the fixed combination (see **INDICATIONS AND CLINICAL USE** and **WARNINGS AND PRECAUTIONS**). In some patients a twice daily administration may be required.

Patients do not generally require HCTZ in excess of 50 mg daily, particularly when combined with other antihypertensive agents.

Recommended Dose and Dosage Adjustment

Monotherapy: The recommended initial dose of quinapril in patients not on diuretics is 10 mg once daily. An initial dose of 20 mg once daily can be considered for patients without advanced age, renal impairment, or concomitant heart failure and who are not volume depleted (see **WARNINGS AND PRECAUTIONS, Cardiovascular, Hypotension**). Dosage should be adjusted according to blood pressure (BP) response, generally at intervals of 2-4 weeks. A dose of 40 mg daily should not be exceeded.

In some patients treated once daily, the antihypertensive effect may diminish towards the end of the dosing interval. This can be evaluated by measuring BP just prior to dosing to determine whether satisfactory control is being maintained for 24 hours. If it is not, either 2x daily administration with the same total daily dose, or an increase in dose should be considered. If BP is not controlled with quinapril alone, a diuretic may be added. After the addition of a diuretic, it may be possible to reduce the dose of quinapril.

Concomitant Diuretic Therapy: Symptomatic hypotension occasionally may occur following the initial dose of quinapril and is more likely in patients who are currently being treated with a diuretic. The diuretic should, if possible, be discontinued for 2-3 days before beginning therapy with quinapril to reduce the likelihood of hypotension (see **WARNINGS AND PRECAUTIONS, Cardiovascular, Hypotension**). If the diuretic cannot be discontinued, an initial dose of 5 mg of quinapril should be used with careful medical supervision for several hours and until BP has stabilized. The dosage of quinapril should subsequently be titrated (as described above) to the optimal response.

Dosage Adjustment in Renal Impairment: For use in hemodialysis patients, see **WARNINGS AND PRECAUTIONS** and **ADVERSE REACTIONS - Anaphylactoid Reactions during Membrane Exposure**. Quinapril should be administered on days when dialysis is not performed.

Starting doses should be reduced according to the following guidelines:

Creatinine Clearance (mL/min)	Maximum Recommended Initial Dose (mg)
> 60	10
30 – 60	5
10 - 30	2.5
< 10	Insufficient data for dosage recommendation

Patients should subsequently have dosage titrated (as described above) to the optimal response as described under Monotherapy.

When concomitant diuretic therapy is required in patients with severe renal impairment a loop diuretic rather than a thiazide is preferred for use with quinapril. Therefore, for patients with severe renal dysfunction TEVA-QUINAPRIL/HCTZ is not recommended.

Dosage in the Elderly: The recommended initial dosage of quinapril is 10 mg once daily (depending on renal function), followed by titration to the optimal response as described above under Monotherapy.

OVERDOSAGE

No data are available regarding overdosage with quinapril hydrochloride and hydrochlorothiazide or quinapril. The most likely clinical manifestation would be symptoms attributable to severe hypotension, which should be normally treated by intravenous volume expansion with 0.9% sodium chloride. Hemodialysis and peritoneal dialysis have little effect on the elimination of quinapril and quinaprilat.

The most common signs and symptoms observed for HCTZ monotherapy overdosage are those caused by electrolyte depletion (hypokalemia, hyponatremia, hyponatremia) and dehydration resulting from excessive diuresis. If digitalis has also been administered, hypokalemia may accentuate cardiac arrhythmias.

For management of a suspected drug overdose, contact your regional Poison Control Centre.

ACTION AND CLINICAL PHARMACOLOGY

Mechanism of Action

TEVA-QUINAPRIL/HCTZ (quinapril hydrochloride and hydrochlorothiazide) is a fixed-combination tablet which combines the antihypertensive actions of an angiotension-converting enzyme (ACE) inhibitor, quinapril hydrochloride and a diuretic, hydrochlorothiazide (HCTZ). In clinical studies, administration of this combination produced greater reductions in blood pressure (BP) than the single agents given alone.

Pharmacodynamics

Quinapril: Quinapril is a nonpeptide, nonsulphydryl ACE inhibitor. ACE is a peptidyl dipeptidase that catalyzes the conversion of angiotensin I to the vasoconstrictor angiotensin II. After absorption, quinapril is rapidly de-esterified to quinaprilat (quinapril diacid), its principal active metabolite. Its primary mode of action is to inhibit circulating and tissue ACE, thereby decreasing vasopressor activity and aldosterone secretion. Although the decrease in aldosterone is small, it results in a small increase in serum K⁺ (see **WARNINGS AND PRECAUTIONS**). Removal of angiotensin II negative feedback on renin secretion leads to increased plasma renin activity.

ACE is identical to kininase II. Thus, quinapril may interfere with the degradation of bradykinin, a potent peptide vasodilator. However, it is not known whether this system contributes to the therapeutic effects of quinapril.

In animal studies, the antihypertensive effect of quinapril outlasts its inhibitory effect on circulating ACE. Tissue ACE inhibition more closely correlates with the duration of antihypertensive effects and this may be related to enzyme binding characteristics.

Administration of 10-40 mg of quinapril to patients with essential hypertension results in a reduction of both sitting and standing BP with minimal effect on heart rate. Antihypertensive activity commences within 1 hour with peak effects usually achieved by 2- 4 hours after dosing. Achievement of maximum BP lowering effects may require 2 - 4 weeks of therapy in some patients. At the recommended doses, antihypertensive effects are maintained throughout the 24-hour dosing interval in most patients. While the dose response relationship is relatively flat, a dose of 40 mg was somewhat more effective at trough than 10-20 mg, and 2x daily dosing tended to give a somewhat lower BP than 1x daily dosing with the same total daily dose. The antihypertensive effect of quinapril was maintained during long-term therapy with no evidence of loss of effectiveness.

Hemodynamic assessments in patients with essential hypertension indicate that BP reduction produced by quinapril is accompanied by a reduction in total peripheral resistance and renal vascular resistance with little or no change in heart rate and cardiac index. There was an increase in renal blood flow which was not significant. Little or no change in glomerular filtration rate (GFR) or filtration fraction was observed.

Hydrochlorothiazide: HCTZ acts directly on the kidney to increase excretion of sodium and chloride, and an accompanying volume of water. HCTZ also increases the excretion of potassium and bicarbonate and decreases calcium excretion.

As a result of its diuretic effect, HCTZ increases plasma renin activity, increases aldosterone secretion, decreases serum potassium, and increases urinary potassium loss. Administration of quinapril inhibits the renin-angiotensin-aldosterone axis and tends to attenuate the potassium decrease associated with HCTZ.

The mechanism underlying the antihypertensive activity of diuretics is unknown. During chronic administration peripheral vascular resistance is reduced; however, this may be secondary to changes in sodium balance.

Quinapril/Hydrochlorothiazide: When quinapril and HCTZ are given together, the antihypertensive effects are approximately additive.

Pharmacokinetics

Quinapril: Following oral administration of quinapril, peak plasma concentrations of quinapril occur within 1 hour. Based on the recovery of quinapril and its metabolites in urine, the extent of absorption is $\geq 60\%$. Following absorption, quinapril is de-esterified to its major active

metabolite, quinaprilat (quinapril diacid), a potent ACE inhibitor, and to minor inactive metabolites. Quinapril has an apparent half-life in plasma of approximately 1 hour. Peak plasma quinaprilat concentrations occur approximately 2 hours after an oral dose of quinapril. Quinaprilat is eliminated primarily by renal excretion and has an effective accumulation half-life of approximately 3 hours. Quinaprilat has an elimination half-life in plasma of approximately 2 hours with a prolonged terminal phase of 25 hours. Approximately 97% of either quinapril or quinaprilat circulating in plasma is bound to proteins.

The rate and extent of quinapril absorption are diminished moderately (approximately 25-30%) when administered during a high-fat meal. However, no effect on quinapril absorption occurs when taken during a regular meal.

Studies in rats indicate that quinapril and its metabolites do not cross the blood-brain barrier.

Hydrochlorothiazide: After oral administration of HCTZ, diuresis begins within 2 hours, peaks in about 4 hours, and lasts about 6-12 hours; the extent of absorption is approximately 50-80%. HCTZ is excreted unchanged by the kidney. When plasma levels have been followed for ≥ 24 hours, the plasma half-life has been observed to vary between 4-15 hours. At least 61% of the oral dose is eliminated unchanged within 24 hours. HCTZ crosses the placental but not the blood-brain barrier.

Quinapril/Hydrochlorothiazide: Concomitant administration of quinapril and HCTZ has little or no effect on the bioavailability or the pharmacokinetics of either drug.

Special Populations and Conditions

Geriatrics:

Quinapril: Therapeutic effects appear to be the same for elderly (> 65 years of age) and younger adult patients given the same daily dosages, with no increase in AEs in elderly patients.

Race:

Quinapril: The antihypertensive effect of ACE inhibitors is generally lower in black than in non-black patients.

Hepatic Insufficiency:

Quinapril: Pharmacokinetic studies in patients with end-stage renal disease or chronic hemodialysis or continuous ambulatory peritoneal dialysis indicate that dialysis has little effect on the elimination of quinapril and quinaprilat.

The disposition of quinapril and quinaprilat in patients with renal insufficiency is similar to that in patients with normal renal function until creatinine clearance is ≤ 60 mL/min. With creatinine clearance < 60 mL/min, peak and trough quinaprilat concentrations increase, apparent half-life increases, and time to steady state may be delayed. The elimination of quinaprilat may be reduced in elderly patients (> 65 years) and in those with heart failure; this reduction is attributable to decrease in renal function (see **DOSAGE AND ADMINISTRATION**). Quinaprilat concentrations are reduced in patients with alcoholic cirrhosis due to impaired de-esterification of quinapril.

STORAGE AND STABILITY

Store at controlled room temperature between 15° - 30°C.

DOSAGE FORMS, COMPOSITION AND PACKAGING

TEVA-QUINAPRIL/HCTZ is available as fixed combination tablets in 3 strengths of quinapril hydrochloride with hydrochlorothiazide:

- **TEVA-QUINAPRIL/HCTZ 10 mg/12.5 mg:** Contains 10 mg of quinapril as quinapril hydrochloride and 12.5 mg of hydrochlorothiazide. Pink, oval, biconvex, film-coated tablet, scored on both sides, engraved with N scoreline N on one side and 10 scoreline 12.5 on the other side.
- **TEVA-QUINAPRIL/HCTZ 20 mg/12.5 mg:** Contains 20 mg of quinapril as quinapril hydrochloride and 12.5 mg of hydrochlorothiazide. Pink, triangular, biconvex, film-coated tablet, engraved with 20 and 12.5 on one side and N on the other side.
- **TEVA-QUINAPRIL/HCTZ 20 mg/25 mg:** Contains 20 mg of quinapril as quinapril hydrochloride and 25 mg of hydrochlorothiazide. Pink, round, biconvex, film-coated tablet, unscored, engraved with N on one side and 20 over 25 on the other side.

Available in blisters of 30 tablets.

Composition

TEVA-QUINAPRIL/HCTZ contain the following inactive ingredients: Calcium phosphate (dibasic) anhydrous, Crospovidone, Hydroxypropyl cellulose, Hydroxypropyl methylcellulose, Iron oxide red, Iron oxide yellow, Magnesium Carbonate, Magnesium Stearate, Polyethylene glycol, Povidone and Titanium Dioxide.

PART II: SCIENTIFIC INFORMATION

PHARMACEUTICAL INFORMATION


Drug Substance: Quinapril Hydrochloride

- Common name: Quinapril Hydrochloride
- Chemical name:
1. [3S-[2[R*(R*)],3R*]]-2-[2-[[1-(Ethoxycarbonyl)-3-phenylpropyl]amino]-1-oxopropyl]-1,2,3,4-tetrahydro-3-isoquinolinecarboxylic acid hydrochloride
 2. (S)-2-[(S)-1-Carboxy-3-phenylpropyl]alanyl]-1,2,3,4-tetrahydro-3-isoquinoline-carboxylic acid 1-ethyl ester hydrochloride

Molecular formula: $C_{25}H_{30}N_2O_5 \cdot HCl$

Molecular mass: 474.99

Structural formula:



- Physiochemical properties: White to off-white solid with a pink cast at times.
- Solubility: Freely soluble in aqueous solvents.
- Melting point: Melting starts at 120°C and the product decomposes.
- Dissociation Constants: $pK_{a1} = 5.3$; $pK_{a2} = 3.4$; $pK_{a3} = 3.1$

Quantitative aqueous pH solubility profile data:

Buffer pH	Solubility in mg/ml	Dose/Solubility* (mL) (for 40 mg tablet)
1	> 2	< 20
2	> 2	< 20
3	> 2	< 20
4	> 2	< 20
5	> 2	< 20
6	> 2	< 20
7	> 2	< 20
8	> 2	< 20

Drug Substance: Hydrochlorothiazide


Common name: Hydrochlorothiazide

Chemical name: 1. 2H-1,2,4-Benzothiadiazine-7-sulfonamide,6-chloro-3,4-dihydro-,1,1-dioxide
2. 6-Chloro-3,4 dihydro-2H-1,2,4-benzothiadiazine-7-sulfonamide 1,1-dioxide

Molecular formula: $C_7H_8ClN_3O_4S_2$

Molecular mass: 297.74

Structural formula:



Physicochemical properties: Hydrochlorothiazide is a white to off-white, crystalline powder

Solubility: Practically insoluble in water, but freely soluble in sodium hydroxide solution.

Dissociation Constants: $pK_{a1} = 7.9$; $pK_{a2} = 9.2$

Quantitative aqueous pH solubility profile data:

Solvent	Buffer pH	Solubility in mg/mL	Dose/Solubility* (mL) (for 25 mg hydrochlorothiazide tablet)
0.1N HCl	1.0	0.608	< 50
0.1N Acetate Buffer	4.5	0.623	< 50
Simulated intestinal fluid	7.5	1.09	< 25
0.05 M borate buffer	8.9	1.03	< 25

CLINICAL TRIALS

A blind, randomized, 3-way crossover, bioequivalence study of Quinapril-Hydrochlorothiazide 20-12.5 mg tablets (Teva Canada Limited, Canada) and Accuretic 20-12.5 mg (Pfizer Canada Inc.) in 48 male healthy subjects under fasting conditions.

SUMMARY TABLE OF THE COMPARATIVE BIOAVAILABILITY DATA

<p style="text-align: center;">Quinapril (1 x 20-12.5 mg) From measured data uncorrected for potency Geometric Mean Arithmetic Mean (CV %)</p>				
Parameter	Test*	Reference†	% Ratio of Geometric Means	90% Confidence Interval (%)
AUC _{0-t} (ng.h/mL)	140.28	140.23	100.04	95.59-104.70
	146.83 (34.93)	144.19 (26.68)		
AUC _{0-inf} (ng.h/mL)	146.08	146.10	99.98	95.67-104.48
	152.54 (33.70)	149.95 (25.62)		
C _{max} (ng/mL)	128.89	130.53	98.74	88.89-109.69
	137.27 (35.21)	138.89 (36.37)		
T _{max} [§] (h)	0.703 (34.30)	0.700 (36.60)		
T _{1/2} [§] (h)	0.87 (22.30)	0.92 (20.00)		

* TEVA-QUINAPRIL/HCTZ 20/12.5 mg Tablets (Manufactured for Teva Canada Limited)

† Accuretic® 20/12.5 mg Tablets (Pfizer Canada Inc., Canada), purchased in Canada

§ Expressed as either the arithmetic mean (CV%) only or the median (range) only

SUMMARY TABLE OF THE COMPARATIVE BIOAVAILABILITY DATA

<p style="text-align: center;">Hydrochlorothiazide (1 x 20-12.5 mg) From measured data uncorrected for potency Geometric Mean Arithmetic Mean (CV %)</p>				
Parameter	Test*	Reference†	% Ratio of Geometric Means	90% Confidence Interval (%)
AUC _{0-t} (ng.h/mL)	490.17	482.37	101.62	98.71-104.60
	502.90 (19.10)	495.44 (18.87)		
AUC _{0-inf} (ng.h/mL)	515.29	506.80	101.67	98.95-104.47
	528.88 (19.60)	520.31 (18.84)		
C _{max} (ng/mL)	76.06	75.77	100.39	94.88-106.21
	78.31 (22.85)	77.93 (22.13)		
T _{max} [§] (h)	2.26 (39.47)	2.19 (41.82)		
T _½ [§] (h)	9.22 (10.28)	9.15 (7.34)		

* TEVA-QUINAPRIL/HCTZ 20/12.5 mg Tablets (Manufactured for Teva Canada Limited)

† Accuretic® 20/12.5 mg Tablets (Pfizer Canada Inc., Canada), purchased in Canada

§ Expressed as either the arithmetic mean (CV%) only or the median (range) only

A blind, randomized, 3-way crossover, bioequivalence study of Quinapril-Hydrochlorothiazide 20-25 mg tablets (Teva Canada Limited, Canada) and Accuretic 20-25 mg (Pfizer Canada Inc.) in 49 male healthy subjects under fasting conditions.

SUMMARY TABLE OF THE COMPARATIVE BIOAVAILABILITY DATA

<p style="text-align: center;">Quinapril (1 x 20-25 mg) From measured data uncorrected for potency Geometric Mean Arithmetic Mean (CV %)</p>				
Parameter	Test*	Reference†	% Ratio of Geometric Least Square Means	90% Confidence Interval (%)
AUC _{0-t} (ng.h/mL)	136.72	136.11	100.45	95.80-105.32
	144.53 (36.27)	143.74 (33.40)		
AUC _{0-inf} (ng.h/mL)	140.35	140.43	99.95	95.38-104.74
	148.10 (35.55)	148.27 (33.04)		
C _{max} (ng/mL)	136.95	136.63	100.23	89.88-111.78
	150.08 (43.16)	147.42 (42.42)		
T _{max} [§] (h)	0.672 (47.40)	0.638 (35.14)		
T _{1/2} [§] (h)	0.85 (15.47)	0.89 (19.11)		

* TEVA-QUINAPRIL/HCTZ 20/25 mg Tablets (Manufactured for Teva Canada Limited)

† Accuretic® 20/25 mg Tablets (Pfizer Canada Inc., Canada), purchased in Canada

§ Expressed as either the arithmetic mean (CV%) only or the median (range) only

<p style="text-align: center;">Hydrochlorothiazide (1 x 20-25 mg) From measured data uncorrected for potency Geometric Mean Arithmetic Mean (CV %)</p>				
Parameter	Test*	Reference†	% Ratio of Geometric Least Square Means	90% Confidence Interval (%)
AUC _{0-t} ‡ (ng.h/mL)	888.47	905.14	98.16	95.08-101.34
	902.66 (18.97)	913.91 (15.73)		
AUC _{0-inf} (ng.h/mL)	945.37	958.56	98.62	95.65-101.69
	961.39 (19.57)	968.14 (15.92)		
C _{max} (ng/mL)	143.58	147.74	97.18	90.68-104.16
	148.26 (25.42)	152.25 (26.29)		
T _{max} § (h)	2.07 (37.33)	2.14 (40.68)		
T _{1/2} § (h)	9.90 (12.80)	9.75 (10.64)		

* TEVA-QUINAPRIL/HCTZ 20/25 mg Tablets (Manufactured for Teva Canada Limited)

† Accuretic® 20/25 mg Tablets (Pfizer Canada Inc., Canada), purchased in Canada

§ Expressed as either the arithmetic mean (CV%) only or the median (range) only

DETAILED PHARMACOLOGY

Quinapril: Mechanism of Action

In Vitro Studies: Quinapril was shown to be an inhibitor of angiotensin converting enzyme (ACE) in both plasma and tissue. In assays utilizing human plasma as sources of ACE, the diacid form of quinapril (quinaprilat) exhibited greater inhibition of ACE activity than quinapril (6.4×10^{-10} M and 8.4×10^{-8} M respectively). In rabbit and rat aortic strips, quinapril (10^{-7} M, 10^{-5} M) specifically suppressed the contractile responses elicited by angiotensin I (50% contraction at approximately 10^{-7} M and 10^{-6} M angiotensin I respectively), but had no effect on contractions induced by angiotensin II and potassium chloride.

In Vivo Studies: Following oral dosing of quinapril, captopril or enalapril (0.1-3 mg/kg) to conscious normotensive rats, plasma ACE inhibition was assessed in vivo by the decrease in pressor response to intravenous (IV) angiotensin I, angiotensin II, norepinephrine and bradykinin. Quinapril produced a dose-dependent reduction (44% at 0.1 mg/kg, 81% at 0.3 mg/kg) of angiotensin I (0.32 µg/kg IV) pressor response and potentiated the response to bradykinin (154% after 0.3 mg/kg quinapril), but had no effect on angiotensin II and norepinephrine responses. Quinapril was equipotent to captopril and enalapril, but had a longer duration of action than

captopril. In the conscious dog, oral administration of quinapril (0.1-3 mg/kg) resulted in plasma ACE inhibition comparable to that of enalapril and captopril.

In human subjects, quinapril at single oral doses of 10-20 mg/day produced 95-100% inhibition of plasma ACE activity at 0.5 hour post-dose, with > 80% inhibition persisting at 24 hours post-dose. Multiple oral doses of quinapril to humans for 12-weeks (20-80 mg/day) confirmed the inhibitory effect on plasma ACE and showed that it produces corresponding decreases in angiotensin II with significant increases in plasma renin activity. Once or 2x daily dosing did not alter the results.

Hydrochlorothiazide

For a discussion of the pharmacology of HCTZ, consult the corresponding product monograph.

Quinapril Hydrochloride and Hydrochlorothiazide: Antihypertensive Activity

Animal Studies: In spontaneously hypertensive rats the combination of quinapril/HCTZ administered for 3 days was more efficacious than quinapril or HCTZ monotherapy. Quinapril alone (0.3 mg/kg PO) and HCTZ (30 mg/kg PO) alone reduced blood pressure (BP) by ≤ 17 and 16%, respectively ($p > 0.05$). Plasma renin activity was elevated 3x by HCTZ. The combination of quinapril/ HCTZ (0.3 and 30 mg/kg PO) produced significant reductions in BP on days 1 and 2 and a sustained reduction (28- 32%) in BP on day 3.

In chronic two kidney, perinephritic hypertensive dogs neither quinapril nor enalapril had antihypertensive activity at oral doses ≤ 10 mg/kg. Plasma renin activity was only transiently elevated in this model. Co-administration of HCTZ (10 mg/kg PO) and quinapril (10 mg/kg PO) in HCTZ pretreated dogs (10 mg/kg for 1 day) reduced mean arterial BP by 35 mmHg. HCTZ monotherapy had no significant effect on BP over a 24-hour period.

Clinical Studies: Two controlled studies evaluated the efficacy and safety of quinapril and HCTZ combination therapy compared with each drug given as monotherapy in patients with essential hypertension. The combination therapy caused a statistically significant greater fall in diastolic blood pressure (DBP) than each drug given as monotherapy. In a placebo controlled study, when quinapril hydrochloride (10 mg, 40 mg) and HCTZ (12.5 mg, 25 mg) were administered alone or in combination, mean reductions in DBP (at trough) produced by quinapril monotherapy ranged from 7.3-10.3 mmHg, by HCTZ monotherapy from 7.2-11.4 mmHg, and by combination therapy from 8.2 -14.9 mmHg. Placebo produced a mean reduction in DBP of 2.2 mmHg.

TOXICOLOGY

Quinapril Hydrochloride

Acute Toxicity: The acute oral (PO) and intravenous (IV) toxicity of quinapril are summarized in Table 2.

Table 2: Acute Toxicity of Quinapril

Species	Sex	Route of Administration	Median Lethal Dose (mg/kg)
Mouse	Male	PO	1492 – 2150
	Female	PO	1440 – 2005
	Male	IV	504
	Female	IV	523
Rat	Male	PO	4280
	Female	PO	3541
	Male	IV	158 – 300
	Female	IV	108 – 273
Dog	Male & Female	PO	>400

Quinapril showed a low order of acute toxicity. Clinical signs of toxicity in both mice and rats were depression or hypoactivity, prostration and ataxia. Peak mortality occurred within 24 hours in oral studies and within 15 minutes in IV studies. Asymptomatic oral dose levels were 500 mg/kg in mice and 1000 mg/kg in rats.

In the dog study, escalating oral doses of 50- 400 mg/kg were given over 13-consecutive days. Vomiting occurred after doses of > 150 mg/kg. BPs decreased with increasing dose. At 400 mg/kg, the female had elevated creatinine and blood urea nitrogen (BUN) levels, decreased sodium and chloride levels, and granular casts in the urine. Gastric erosions and ulcers were seen in both animals and renal tubular dilatation was noted in the female.

The results of quinapril toxicity from subacute, chronic, reproductive, genetic, and carcinogenicity studies are given in Tables 3- 7, respectively.

Table 8 summarizes the results of toxicity studies with quinaprilat, the major active metabolite of quinapril.

TABLE 3: Subacute Toxicity Studies of Quinapril

Species	Duration (Week)	No. of Animals/ Sex/Group	Route	Doses (mg/kg/day)	Results
Mouse	2	10	PO	VC ¹ , 125, 250, 500 750	One drug-related death at 750 mg/kg; reduced food consumption and body weight gain. MTD ² about 500 mg/kg.
Mouse	13	10	PO	VC, 50, 125, 250 500	Body weight gain suppression, decreases in heart weight, hyperplasia of juxtaglomerular apparatus (JGA). MTD between 50 and 125 mg/kg.
Rat	2	5	PO	VC, 200, 400, 800, 1200	Deaths at 400, 800, and 1200 mg/kg; salivation, reduced food consumption, body weight gain suppression, pulmonary, renal, and gastric lesions.
Rat	2	10	PO	UC ¹ , VC, 100, 400 800	Deaths at 400 and 800 mg/kg; respiratory signs, salivation, increased BUN, decreased RBC, Hgb, and Hct; increased liver weights, decreased heart weights; pulmonary edema and focal gastric erosions. MTD was between 400 and 800 mg/kg
Rat	13	12	PO	UC, VC, 50, 250 500	Deaths at 250 and 500 mg/kg; salivation; slightly increased BUN, CPK, and LDH; decreased RBC, Hct, and Hgb; decreased heart weight, pulmonary and gastric lesions at \geq 250 mg/kg; increased renin granules in JG cells. MTD between 50 and 250 mg/kg.
Dog	2	2	PO	VC, 25, 125, 250 (125 b.i.d.)	No deaths; emesis, mild focal erosions and inflammation of the stomach at 125 mg/kg. MTD estimated as 250 mg/kg.
Dog	13	3	PO	VC, 25, 125, 250 (125 b.i.d.)	Sporadic emesis and anorexia; mild to moderate reversible increase in BUN and mild depression of RBC, Hct, Hgb at 250 mg/kg; focal gastric erosions at \geq 125 mg/kg, increases in renin granules in JG cells; hypertrophy and hyperplasia of JGA. MTD was between 25 and 125 mg/kg

1. VC = Vehicle Control; UC = Untreated Control.
 2. MTD = Maximum Tolerated Dose; RBC = red blood cell count H b =hemoglobin; Hct = hematocrit; CPK = creatine phosphokinase; LDH = lactate dehydrogenase; JG = juxtaglomerular apparatus; BUN = blood urea nitrogen.

TABLE 4: Chronic Toxicity Studies of Quinapril					
Species	Duration (Week)	No. of Animals/ Sex/Group	Route	Doses (mg/kg/day)	Results
Rat	57 ¹	30	PO	UC ² , VC ² , 10, 50, 100	No drug-related deaths; transient post-dose salivation, body weight gain suppression, increased BUN, decreased glucose, increased plasma renin level, decreased heart weight, JGA hypertrophy and hyperplasia with increased granules; degenerative changes in kidneys.
Dog	52	4	PO	VC 10, 50, 100	No deaths; elevation of plasma renin and liver enzyme levels, focal areas of chronic active inflammation in the liver at 100 mg/kg; gastric erosion at 50 mg/kg, and hypertrophy/hyperplasia of renal JGA.
¹ 52 weeks treatment plus 4 weeks without treatment for some animals ² UC = Untreated Control; VC = Vehicle Control; BUN = blood urea nitrogen; JGA = juxtaglomerular apparatus					

Table 5: Reproductive Toxicology Studies of Quinapril					
Species	No. of Animals/ Sex/Group	Route	Doses (mg/kg/day)	Duration of Dosing	Results
<u>Fertility</u> Rat	12 Male 24 Female	PO	VC ¹ , 10, 50, 100	<u>Males</u> 60 days prior to mating <u>Females</u> 14 days prior to mating until weaning of offspring	No effects on fertility, no adverse effects on F ₁ offspring parameters and no teratogenic effects.
<u>Teratology</u> Rat	5 Female	PO	100, 200, 400, 600, 800	Days 6 to 15 of gestation	No teratogenicity. Maternal deaths at 600 and 800 mg/kg; decreased fetal body weights at 200 mg/kg and higher.
Rat	20 Female	PO	UC ¹ , VC, 50, 150, 300	Days 6 to 15 of gestation	No fetotoxic or teratogenic effects. Reversible maternal toxicity.
Rabbit	5 – 7 Female	PO	10, 15, 25, 50, 100, 200, 400	Days 6 to 18 of gestation	Severe materno- and fetotoxicity.
Rabbit	5 Female	PO	VC, 1, 2, 4, 6, 8	Days 6 to 18 of gestation	Abortions and maternal deaths at 4, 6, and 8 mg/kg; materno- and fetotoxicity at doses > 1 mg/kg.
Rabbit	14 Female	PO	VC 0.5, 1.0, 1.5	Days 6 to 18 of gestation	Not teratogenic. Maternal weight loss; increased incidence of postimplantation loss (embryotoxicity) at 1.0 and 1.5 mg/kg.
<u>Perinatal/ Postnatal</u> Rat	20 Female	PO	VC, 25, 75, 150	Day 15 of gestation to Day 20 of lactation	Reduction in offspring body weights from birth to Day 21 postnatally at 25, 75 and 150 mg/kg.

¹ UC = Untreated Control; VC = Vehicle Control

	Test	Dosage Range	Results
Mutagenicity			
1) In Vitro	a) Initial cytotoxicity in <i>Salmonella</i> strain b) Mutagenesis assay in <i>Salmonella</i>	Up to 10,000 µg/plate 625 to 10,000 µg/plate	Non-cytotoxic. Negative-with or without metabolic activation.
2) In Vitro	a) Initial cytotoxicity assay b) Point mutation assay in Chinese hamster lung cells	Up to 44,300 µg/mL 175 to 1400 µg/mL	Cytotoxic at ≥ 1400 µg/mL. Negative-did not manifest direct acting or promutagen activity.
Cytogenetics			
1) In Vitro	a) Initial cytotoxicity assay b) Sister chromatid exchange (SCE) assay in Chinese hamster ovary cells	Up to 44,300 µg/mL 10.94 to 1400 µg/mL	Cytotoxic at concentrations > 700 µg/mL No increase in SCE at toxicity-limited doses ≤ 700 µg/mL in the presence of metabolic activation or ≤ 1400 µg/mL in the absence of metabolic activation.
2) In Vitro	a) Initial cytotoxicity assay b) Structural chromosomal aberration (SCA) assay in Chinese hamster lung cells	Up to 2700 µg/mL 800 to 1800 µg/mL	Cytotoxic at ≥ 1200 µg/mL. Slight, statistically significant increase in SCA with metabolic activation; not considered biologically significant.
3) In Vivo	a) Mouse micronucleus assay	1 to 1430 µg/kg	Not clastogenic; no increased frequency of micronuclei.

Species	Duration (week)	No. of Animals/ Sex/Group	Route	Doses (mg/kg/day)	Results
Mouse	104	50	PO	UC ¹ , VC ¹ , 5, 35, 75	No evidence of tumorigenic potential. Reduced heart weight, nephritis, and JGA hypertrophy/hyperplasia.
Rat	104	65	PO	UC, VC, 10, 50 100	No evidence of tumorigenic potential. Reduced RBC, JGA hypertrophy/hyperplasia and renal degenerative changes.

¹ UC = Untreated Control; VC = Vehicle Control; JGA = juxtaglomerular apparatus; RBC = red blood cell count

Table 8: Toxicity Studies of Quinaprilat					
Species	Duration (week)	No. of Animals/ Sex/Group	Route	Doses (mg/kg/day)	Results
A. Acute Studies					
Mouse	Single-dose	10	IV	VC ¹ , 250, 500, 1000	No deaths; MLD > 1000 mg/kg. No clinical or gross pathological changes.
Rat	Single-dose	10	IV	VC, 50, 100, 200, 300, 400	No deaths; MLD > 400 mg/kg. No clinical or gross pathological changes
Dog	Escalating dose	1	IV	Escalating; 1-240	No deaths; MLD > 240 mg/kg. Reduced food consumption, weight loss, and slight increase in myeloid to erythroid ratio.
B. Subacute Studies					
Rat	2	5	IV	VC, 25, 50, 100, 200	No deaths, clinical signs or adverse pathological findings.
Rat	4	10	IV	VC, 20, 100, 200	No drug-related deaths or clinical signs; reduced heart weights.
Dog	2	1	IV	VC, 10, 50, 100	Sporadic increases in heart rate.
Dog	4	3	IV	VC, 10, 50, 100	No clinical or gross pathologic findings; JGA hypertrophy/hyperplasia.
¹ VC = Vehicle Control; MLD = median lethal dose; JGA = juxtaglomerular apparatus					

C. Genotoxicity Studies:

	Test	Dosage Range	Results
<u>Mutagenicity</u> 1) In Vitro	a) Initial cytotoxicity in <i>Salmonella</i> b) Mutagenesis assay in <i>Salmonella</i>	≤ 1200 µg/plate 75-1200 µg/plate	Non-cytotoxic Negative-with or without metabolic activation

Quinapril Hydrochloride and Hydrochlorothiazide

The 14 day Median Lethal Dose (MLD) in mice was 1068/667 mg/kg quinapril/ HCTZ; in rats it was 4640/2896 mg/kg. For quinapril alone, the oral MLD ranged from 1440-2150 mg/kg for mice, and from 3531-4280 mg/kg in rats. In dogs, no drug-related clinical signs of toxicity were observed at doses of 125-250x the maximum human dose of quinapril given in combination with HCTZ, and 60-120x the maximum human dose of HCTZ in combination (100/60 mg/kg in males, 200/120 mg/kg in females).

REFERENCES

1. Ferry J, Horvath A, Sedman A, et al. Influence of food on the pharmacokinetics of quinapril and its active diacid metabolite, CI-928. *J Clin Pharmacol* 1987; 27:397-399.
2. Ferry J, Cetnarowski A, Sedman A, et al. Multiple-Dose Cimetidine Administration Does Not Influence the Single-Dose Pharmacokinetics of Quinapril and Its Active Metabolite CI-928. *J Clin Pharmacol* 1988; 28:48-51.
3. Fabris B, Chen B Pusic V, et al. Inhibition of angiotensin-converting enzyme (ACE) in plasma and tissue. *J Cardiovasc Pharmacol* 1990; 15(Suppl 2):S6-S13.
4. Frank G. Overview of the clinical development of quinapril. *Clin Cardiol* 1990; 13(Suppl 7):13-18.
5. Frishman W. The safety and efficacy of quinapril in the treatment of mild to moderate essential hypertension. *Clin Cardiol* 1990; 13(Suppl 7):19-25.
6. Goldstein R. The treatment of moderate to severe hypertension with ACE inhibitors. *J Cardiovasc Pharmacol* 1990; 15(Suppl2):S29-S35.
7. Gupta R, Kjeldsen S, Krause L, et al. Hemodynamic effects of quinapril a novel angiotensin-converting enzyme inhibitor. *Clin Pharmacol Ther* 1990; 48:41-49.
8. Kaplan H, Taylor D, Olson S. Quinapril: Overview of preclinical data. *Clin Cardiol* 1990; 13(Suppl7):4-12.
9. Knapp L, Frank G, McLain R, et al. The safety and tolerability of quinapril. *J Cardiovasc Pharmacol* 1990; 15(Suppl2):S47-S55.
10. Schnaper H. The management of hypertension in older patients. *J Cardiovasc Pharmacol* 1990; 15(Suppl2):S56-S61.
11. Swartz R, Stermann B Horvath A, et al. Pharmacokinetics of Quinapril and its active metabolite quinaprilat during continuous ambulatory peritoneal dialysis. *J Clin Pharmacol* 1990; 30:1136-41.
12. Taylor S. The treatment of mild to moderate hypertension with ACE inhibitors. *J Cardiovasc Pharmacol* 1990; 15(Suppl2):S24-S28.
13. Verresen L, Waer M Vanrenterhem Y, et al. Angiotensin converting enzyme inhibitors and anaphylactoid reactions to high-flux membrane dialysis. *Lancet* 1990; 336:1360-1362.
14. Wadworth A, Brogden R. Quinapril: A review of its pharmacological properties and therapeutic efficacy in cardiovascular disorders. *Drugs* 1991; 41:378-399.

15. Halstenson C, Opsahl J, Rachael K, et al. The pharmacokinetics of quinapril and its active metabolite, quinaprilat, in patients with various degrees of renal function. *J Clin Pharmacol* 1992; 32:344-350.
16. Canter D, Frank G, et al. Quinapril and hydrochlorothiazide combination for control of hypertension: assessment by factorial design. *J Human Hypertension* 1994; 8:155-162.
17. Lenz T, Schulte K, et al. Quinapril, hydrochlorothiazide, and combination in patients with moderate to severe hypertension. *Eur Heart J* 1994; 15:940-946.
18. A blind, randomized, 3-way crossover, bioequivalence study of Quinapril-Hydrochlorothiazide 20-12.5 mg tablets (Teva Canada Limited, Canada) and Accuretic 20-12.5 mg (Pfizer Canada Inc.) in healthy subjects under fasting conditions. Data on file at Teva Canada Limited.
19. A blind, randomized, 3-way crossover, bioequivalence study of Quinapril-Hydrochlorothiazide 20-25 mg tablets (Teva Canada Limited, Canada) and Accuretic 20-25 mg (Pfizer Canada Inc., Canada) in healthy subjects under fasting conditions. Data on file at Teva Canada Limited.
20. ACCURETIC Product Monograph by Pfizer Canada Inc., Control No. 201238. Date of Revision: March 21, 2017.

PART III: CONSUMER INFORMATION

Pr^rTEVA-QUINAPRIL/HCTZ

(Quinapril hydrochloride/Hydrochlorothiazide Tablets)

Read this carefully before you start taking TEVA-QUINAPRIL/HCTZ and each time you get a refill. This leaflet is a summary and will not tell you everything about TEVA-QUINAPRIL/HCTZ. Talk to your doctor, nurse or pharmacist about your medical condition and treatment and ask if there is any new information about TEVA-QUINAPRIL/HCTZ.

ABOUT THIS MEDICATION

What the medication is used for:

TEVA-QUINAPRIL/HCTZ (Quinapril hydrochloride/Hydrochlorothiazide) lowers high blood pressure.

What it does:

TEVA-QUINAPRIL/HCTZ contains a combination of 2 drugs, quinapril hydrochloride and hydrochlorothiazide:

- Quinapril hydrochloride is an angiotensin converting enzyme (ACE) inhibitor. You can recognize ACE inhibitors because their medicinal ingredient ends in “PRIL”. It lowers blood pressure.
- Hydrochlorothiazide is a diuretic or “water pill” that increases urination. This lowers blood pressure.

This medicine does not cure high blood pressure. It helps to control it. Therefore, it is important to continue taking TEVA-QUINAPRIL/HCTZ regularly even if you feel fine.

When it should not be used:

Do not take TEVA-QUINAPRIL/HCTZ if you:

- Are allergic to quinapril hydrochloride or hydrochlorothiazide or to any non-medicinal ingredients in the formulation
- Are allergic to any sulfonamide-derived drugs (sulfa drugs); most of them have a medicinal ingredient that ends in “-MIDE”
- Have experienced an allergic reaction (angioedema) with swelling of the hands, feet, or ankles, face, lips, tongue, throat, or sudden difficulty breathing or swallowing, to any ACE inhibitor or without a known cause. Be sure to tell your doctor, nurse, or pharmacist that this has happened to you
- Have been diagnosed with hereditary angioedema: an increased risk of getting an allergic reaction that is passed down through families. This can be triggered by different factors, such as surgery, flu, or dental procedures
- Are taking Entresto (sacubitril/valsartan), due to the increased risk of serious allergic reaction which causes swelling of the face or throat (angioedema)

- when taken with TEVA-QUINAPRIL/HCTZ.
- Have difficulty urinating or produce no urine
- Are pregnant or intend to become pregnant. Taking TEVA-QUINAPRIL/HCTZ during pregnancy can cause injury and even death to your baby
- Are breastfeeding, TEVA-QUINAPRIL/HCTZ passes into breast milk
- Are taking aliskiren-containing medicines, such as Rasilez, **and** have one of the following conditions:
 - Diabetes
 - Kidney disease
 - High levels of potassium
 - Congestive heart failure combined with hypotension
- Are taking an angiotensin receptor blocker (ARB), another medicine to treat your high blood pressure, or another ACE inhibitor and have one of the following conditions:
 - Diabetes with end organ damage
 - Kidney disease
 - High levels of potassium
 - Congestive heart failure combined with hypotension

You can recognize ARBs because their medicinal ingredient ends in “-SARTAN”.

What the medicinal ingredients are:

Quinapril hydrochloride and Hydrochlorothiazide

What the non-medicinal ingredients are:

Calcium phosphate (dibasic) anhydrous, Crospovidone, Hydroxypropyl cellulose, Hydroxypropyl methylcellulose, Iron oxide red, Iron oxide yellow, Magnesium Carbonate, Magnesium Stearate, Polyethylene glycol, Povidone and Titanium Dioxide.

What dosage forms it comes in:

10 mg/12.5 mg, 20 mg/12.5 mg and 20 mg/25 mg Tablets

WARNINGS AND PRECAUTIONS

Serious Warnings and Precautions - Pregnancy

TEVA-QUINAPRIL/HCTZ should not be used during pregnancy. If you discover that you are pregnant while taking TEVA-QUINAPRIL/HCTZ, stop the medication and contact your doctor, nurse, or pharmacist as soon as possible.

Before you use TEVA-QUINAPRIL/HCTZ, talk to your doctor, nurse or pharmacist if you:

- Are allergic to any drug used to lower blood pressure or penicillin
- Have recently received or are planning to get allergy shots for bee or wasp stings
- Have narrowing of an artery or a heart valve
- Have had a heart attack or stroke
- Have heart failure

- Have diabetes, liver or kidney disease
- Have lupus or gout
- Are on dialysis or receiving LDL apheresis (treatment to remove “bad cholesterol” from the blood)
- Are dehydrated or suffer from excessive vomiting, diarrhea, or sweating
- Are taking a salt substitute that contains potassium, potassium supplements, or a potassium-sparing diuretic (a specific kind of “water pill”)
- Are taking an antibiotic containing trimethoprim
- Are on a low-salt diet
- Are receiving gold (sodium aurothiomalate) injections
- Are less than 18 years old
- Are taking a neutral endopeptidase inhibitor. The combination with TEVA-QUINAPRIL/HCTZ is not recommended.
- Are taking an aliskiren-containing medicine, such as Rasilez, used to lower high blood pressure. The combination with TEVA-QUINAPRIL/HCTZ is not recommended. Are taking an angiotensin receptor blocker (ARB). You can recognize an ARB because its medicinal ingredient ends in “-SARTAN”
- Are taking a medicine that contains aliskiren, such as Rasilez, an angiotensin receptor blocker (ARB), or another ACE inhibitor (in addition to TEVA-QUINAPRIL/HCTZ). The combination with TEVA-QUINAPRIL/HCTZ is not recommended.
- Are currently taking anti-cancer (temsirolimus, everolimus), anti-rejection (sirolimus) or anti-diabetic (gliptins) drugs. Use of ACE inhibitors, such as TEVA-QUINAPRIL/HCTZ, with these drugs may increase the chance of having an allergic reaction.

Hydrochlorothiazide in TEVA-QUINAPRIL/HCTZ can cause sudden eye disorders:

- **Myopia:** sudden nearsightedness or blurred vision
 - **Glaucoma:** an increased pressure in your eye, eye pain. Untreated, it may lead to permanent vision loss
- These eye disorders are related and can develop within hours to weeks of starting TEVA-QUINAPRIL/HCTZ.

You may become sensitive to the sun while taking TEVA-QUINAPRIL/HCTZ. Exposure to sunlight should be minimized until you know how you respond.

If you are going to have surgery and will be given an anesthetic. Be sure to tell your doctor or dentist that you are taking TEVA-QUINAPRIL/HCTZ.

Driving and using machines: before you perform tasks which may require special attention, wait until you know how you respond to TEVA-QUINAPRIL/HCTZ. Dizziness, lightheadedness, or fainting can especially occur after the first dose and when the dose is increased.

INTERACTIONS WITH THIS MEDICATION

As with most medicines, interactions with other drugs are possible. Tell your doctor, nurse, or pharmacist about the medicines you take, including drugs prescribed by other doctors, vitamins, minerals, natural supplements, or alternative medicines.

The following may interact with TEVA-QUINAPRIL/HCTZ:

- Adrenocorticotrophic hormone (ACTH) used to treat West Syndrome
- Alcohol, narcotics (strong pain medications) or barbiturates (sleeping pills). They may cause low blood pressure and dizziness when you go from lying down or sitting to standing up
- Amphotericin B, an antifungal drug
- Anticancer drugs, including cyclophosphamide, methotrexate, temsirolimus and everolimus
- Anti-rejection drugs, such as sirolimus (Rapamune)
- Antidepressants, in particular selective serotonin reuptake inhibitors (SSRIs), including citalopram, escitalopram, and sertraline
- Anti-diabetic drugs including insulin and oral medicines (e.g. metformin, gliptins, sulfonylureas)
- Bile acid resins used to lower cholesterol
- Calcium or vitamin D supplements
- Corticosteroids used to treat joint pain and swelling
- Digoxin, a heart medication
- Drugs known to increase the potassium level in the blood such as a salt substitute that contains potassium, potassium supplements, potassium-sparing diuretic (a specific kind of “water pill”) (e.g. spironolactone, triamterene, amiloride, sulfamethoxazole/trimethoprim).
- Drugs that slow down or speed up bowel function, including atropine, metoclopramide, and domperidone
- Drugs used to treat epilepsy, including carbamazepine and topiramate
- Gold for the treatment of rheumatoid arthritis
- Gout medications, including allopurinol and probenecid
- Lithium used to treat bipolar disease
- Non-steroidal anti-inflammatory drugs (NSAIDs) used to reduce pain and swelling. Examples include ibuprofen, naproxen, and celecoxib
- Blood pressure lowering drugs, including diuretics (“water pills”), aliskiren-containing products (e.g. Rasilez), angiotensin receptor blockers (ARBs) or other ACE inhibitors (in addition to TEVA-QUINAPRIL/HCTZ).
- Pressor amines (drugs which increase blood pressure, such as adrenaline)
- Skeletal muscle relaxants used to relieve muscle spasms, including tubocurarine
- Tetracycline (a type of antibiotic)

PROPER USE OF THIS MEDICATION

TEVA-QUINAPRIL/HCTZ is not for initial therapy. You must first be stabilized on the individual medicinal ingredients (quinapril hydrochloride and hydrochlorothiazide) of TEVA-QUINAPRIL/HCTZ. If your dosage matches the dosages in TEVA-QUINAPRIL/HCTZ, your doctor may prescribe TEVA-QUINAPRIL/HCTZ taken once a day (instead of each medicinal ingredient as a separate pill).

Take TEVA-QUINAPRIL/HCTZ exactly as prescribed. It is recommended to take your dose at about the same time every day.

TEVA-QUINAPRIL/HCTZ can be taken with or without food. If TEVA-QUINAPRIL/HCTZ causes upset stomach, take it with food or milk.

Usual Adult dose:

The recommended starting dose is one 10 mg/12.5 mg tablet daily.

Overdose:

If you think you have taken too much TEVA-QUINAPRIL/HCTZ contact your doctor, nurse, pharmacist, hospital emergency department or regional Poison Control Centre immediately, even if there are no symptoms.

Missed Dose:

If you have forgotten to take your dose during the day, carry on with the next one at the usual time. Do NOT double dose.

SIDE EFFECTS AND WHAT TO DO ABOUT THEM

Side effects may include:

- Dizziness, headache, trouble sleeping
- Drowsiness, fatigue, weakness
- Cough, stuffy and runny nose
- Rash, itching
- Abdominal pain, upset stomach, decreased appetite, constipation
- Muscle pain, spasms, back pain, restlessness
- Pins and needles in your fingers
- Nausea, vomiting, diarrhea
- Sore throat
- Reduced libido
- Rash, red patches on the skin

If any of these affects you severely, tell your doctor, nurse or pharmacist.

TEVA-QUINAPRIL/HCTZ can cause abnormal blood test results. Your doctor will decide when to perform blood tests and will interpret the results.

SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM

Symptom/ Effect		Talk with your doctor or pharmacist		Stop taking drug and seek immediate medical help
		Only if severe	In all cases	
Common	Low blood pressure: dizziness, fainting, light-headedness may occur when you go from lying or sitting to standing up	√		
	Decreased or increased levels of potassium in the blood: Irregular heartbeats, muscle weakness and generally feeling unwell		√	
Uncommon	Allergic reactions including angioedema: Rash, hives. Swelling of the face, lips, tongue or throat, difficulty swallowing or breathing			√
	Kidney disorder: Decreased urination, nausea, vomiting, swelling of extremities, fatigue		√	
	Liver disorder: Yellowing of the skin or eyes, dark urine, abdominal pain, nausea,		√	

SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM				
	vomiting, loss of appetite			
Uncommon	Increased blood sugar: Frequent urination, thirst and hunger	√		
	Electrolyte imbalance: Weakness, drowsiness, muscle pain or cramps, irregular heartbeat		√	
Rare	Decreased platelets: Bruising, bleeding, fatigue and weakness		√	
	Decreased white blood cells: Infections, fatigue, fever, aches, pains and flu-like symptoms		√	
	Edema: Swelling of the hands, ankles or feet		√	
	Vomiting blood			√
	High nitrogen compound found in blood (Azotemia): rapid heart rate, high blood pressure, fatigue, confusion, lightheadedness, dizziness, decreased urine production			√
	Chest pain Heart attack			√
Very rare	Toxic epidermal necrolysis: severe skin peeling, especially in the mouth and eyes			√
Unknown	Eye disorders:			√

SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM				
	Myopia: Sudden near sightedness or blurred vision			
	Glaucoma: Increased pressure in your eyes, eye pain			
	Anemia: Fatigue, loss of energy, weakness, shortness of breath		√	
	Inflammation of the pancreas: Abdominal pain that lasts and gets worse when you lie down, nausea, vomiting		√	
	Tachycardia: Fast heart beat		√	

This is not a complete list of side effects. For any unexpected effects while taking TEVA-QUINAPRIL/HCTZ, contact your doctor, nurse or pharmacist.

HOW TO STORE IT

Store TEVA-QUINAPRIL/HCTZ at room temperature, between 15°C and 30°C. Protect from moisture. Keep in well closed container.

Keep TEVA-QUINAPRIL/HCTZ out of reach and sight of children.

Reporting Side Effects

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada/adverse-reaction-reporting.html>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

MORE INFORMATION

This document plus the full product monograph, prepared for health professionals can be found by contacting Teva Canada Limited by:

Phone: 1-800-268-4127 ext. 3;

Email: druginfo@tevacanada.com; or

Fax: 1-416-335-4472

This leaflet was prepared by:

Teva Canada Limited

30 Novopharm Court

Toronto, Ontario

Canada, M1B 2K9

Prepared on: December 17, 2018