

PRODUCT MONOGRAPH

Pr ACCEL-SILDENAFIL

sildenafil as sildenafil citrate

Tablets 25 mg, 50 mg and 100 mg

House Standard

cGMP-Specific Phosphodiesterase Type 5 Inhibitor

Treatment of Erectile Dysfunction

Accel Pharma Inc.
99 Place Frontenac
Pointe-Claire, Quebec
H9R 4Z7

Date of Revision:
December 27, 2018

Submission Control No: 222236

Table of Contents

PART I: HEALTH PROFESSIONAL INFORMATION	3
SUMMARY PRODUCT INFORMATION	3
INDICATIONS AND CLINICAL USE.....	3
CONTRAINDICATIONS	3
WARNINGS AND PRECAUTIONS.....	4
ADVERSE REACTIONS.....	7
DRUG INTERACTIONS	12
DOSAGE AND ADMINISTRATION	15
OVERDOSAGE	16
ACTION AND CLINICAL PHARMACOLOGY	17
STORAGE AND STABILITY.....	20
SPECIAL HANDLING INSTRUCTIONS	20
DOSAGE FORMS, COMPOSITION AND PACKAGING	21
PART II: SCIENTIFIC INFORMATION	22
PHARMACEUTICAL INFORMATION.....	22
CLINICAL TRIALS.....	22
DETAILED PHARMACOLOGY	26
TOXICOLOGY	29
REFERENCES	41
PART III: CONSUMER INFORMATION.....	46

Pr **ACCEL-SILDENAFIL**
Sildenafil Citrate Tablets
25 mg, 50 mg and 100 mg

PART I: HEALTH PROFESSIONAL INFORMATION

SUMMARY PRODUCT INFORMATION

Route of Administration	Dosage Form / Strength	Clinically Relevant Nonmedicinal Ingredients
Oral	Tablets 25 mg, 50 mg and 100 mg	The tablets also contain the following non-medicinal ingredients: calcium hydrogen phosphate anhydrous, FD&C Blue #2/indigo carmine aluminum lake, macrogol, magnesium stearate, microcrystalline cellulose, polyvinyl alcohol, sodium croscarmellose, talc and titanium dioxide.

INDICATIONS AND CLINICAL USE

ACCEL-SILDENAFIL (sildenafil citrate) is indicated for:

- the treatment of erectile dysfunction, which is the inability to achieve or maintain a penile erection sufficient for satisfactory sexual performance.

CONTRAINDICATIONS

Sildenafil citrate has been shown to potentiate the hypotensive effects of nitrates in healthy volunteers and in patients, and is therefore contraindicated in patients who are taking any type of nitrate drug therapy, or who utilize short-acting nitrate-containing medications, due to the risk of developing potentially life-threatening hypotension. The use of organic nitrates, either regularly and/or intermittently, in any form (e.g. oral, sublingual, transdermal, by inhalation) is absolutely contraindicated (see **ACTION AND CLINICAL PHARMACOLOGY, DOSAGE AND ADMINISTRATION**).

After patients have taken sildenafil citrate, it is unknown when nitrates, if necessary, can be safely administered. Plasma levels of sildenafil at 24 hours post-dose are much lower (2 ng/mL) than at peak concentration (440 ng/mL). In the following patients: age >65, hepatic impairment (e.g. cirrhosis), severe renal impairment (e.g. CLcr <30 mL/min), and concomitant use of potent cytochrome P-450 3A4 inhibitors (erythromycin), plasma levels of sildenafil at 24 hours post-dose have been found to be 3 to 8 times higher than those seen in healthy volunteers. Although plasma levels of sildenafil at 24 hours post-dose are much lower than at peak concentration, it is unknown whether nitrates can be safely coadministered at this time point (see **DETAILED**

PHARMACOLOGY, Pharmacodynamic Studies).

Treatments for erectile dysfunction should not be generally used in men for whom sexual activity is inadvisable (see also **WARNINGS AND PRECAUTIONS**).

Sildenafil citrate is contraindicated in patients with a known hypersensitivity to any component of the tablet (see **PHARMACEUTICAL INFORMATION**).

ACCEL-SILDENAFIL is contraindicated in patients with erectile dysfunction with previous episode of non-arteritic anterior ischaemic optic neuropathy (NAION) (see **WARNINGS AND PRECAUTIONS**).

The co-administration of PDE5 inhibitors, including **ACCEL-SILDENAFIL**, with guanylate cyclase stimulators, such as riociguat, is contraindicated as it may lead to potentially life-threatening episodes of symptomatic hypotension or syncope.

WARNINGS AND PRECAUTIONS

General

The evaluation of erectile dysfunction should include a determination of potential underlying causes and the identification of appropriate treatment following a complete medical assessment.

Cardiovascular

As with all treatments for erectile dysfunction, there is a potential cardiac risk of sexual activity in patients with pre-existing cardiovascular disease, including hypertension (BP >140/90). Therefore, treatments for erectile dysfunction, including sildenafil citrate, should not be generally administered in men for whom sexual activity is inadvisable because of their underlying cardiovascular status.

There are no controlled clinical data on the safety or efficacy of sildenafil citrate in the following groups, if prescribed, this should be done with caution.

- Patients who have suffered a myocardial infarction, stroke, or life-threatening arrhythmia within the last 6 months
- Patients with resting hypotension (BP <90/50 at rest) or hypertension (BP >170/110 at rest)
- Patients with cardiac failure or coronary artery disease causing unstable angina

(see **ACTION AND CLINICAL PHARMACOLOGY**)

Caution is advised when sildenafil is administered to patients taking an alpha-blocker, as the coadministration may lead to symptomatic hypotension in a few susceptible individuals (see **DRUG INTERACTIONS**). In order to minimize the potential for developing postural hypotension, patients should be hemodynamically stable on alpha-blocker therapy prior to initiating sildenafil treatment. Initiation of sildenafil at lower doses should be considered. In addition, physicians should advise patients what to do in the event of postural hypotensive symptoms.

Hematologic

In clinical trials, sildenafil has been shown to have systemic vasodilatory properties that result in

transient decreases in blood pressure (see **DETAILED PHARMACOLOGY**). This is of little or no consequence in most patients. However, prior to prescribing sildenafil, physicians should carefully consider whether their patients with certain underlying conditions could be adversely affected by such vasodilatory effects, especially in combination with sexual activity. Patients with increased susceptibility to vasodilators include those with left ventricular outflow obstruction (e.g., aortic stenosis, hypertrophic obstructive cardiomyopathy), or those with the rare syndrome of multiple system atrophy manifesting as severely impaired autonomic control of blood pressure.

In humans, sildenafil citrate has no effect on bleeding time when taken alone or with acetylsalicylic acid. *In vitro* studies with human platelets indicate that sildenafil potentiates the antiaggregatory effect of sodium nitroprusside (a nitric oxide donor). The combination of heparin and sildenafil citrate had an additive effect on bleeding time in the anesthetized rabbit, but this interaction has not been studied in humans (see **ACTION AND CLINICAL PHARMACOLOGY**).

There is no safety information on the administration of sildenafil citrate to patients with bleeding disorders or active peptic ulceration. Therefore, **ACCEL-SILDENAFIL** should be administered with caution to these patients.

Hepatic/Biliary/Pancreatic

In volunteers with hepatic cirrhosis (Child-Pugh A and B), sildenafil clearance was reduced, resulting in increases in AUC (84%) and C_{max} (47%) compared to age-matched volunteers with no hepatic impairment.

A starting dose of 25 mg should be considered in patients with hepatic impairment (see **ACTION AND CLINICAL PHARMACOLOGY, DOSAGE AND ADMINISTRATION**).

Ophthalmologic

Patients should stop taking PDE5 inhibitors, including sildenafil citrate, and consult their physician immediately if they experience a decrease in, or sudden loss of, vision in one or both eyes. Postmarketing reports of sudden loss of vision have occurred rarely, in temporal association with the use of PDE5 inhibitors. An observational study evaluated whether recent use of PDE5 inhibitors, as a class, was associated with acute onset of NAION. The results suggest an approximate 2-fold increase in the risk of NAION within 5 half-lives of PDE5 inhibitor use.

Individuals who have already experienced NAION are at increased risk of NAION recurrence. PDE5 inhibitors, including sildenafil citrate, are not recommended in patients with male erectile dysfunction with a previous episode of NAION (see **CONTRAINDICATIONS**). There are no controlled clinical data on the safety or efficacy of sildenafil citrate in patients with retinitis pigmentosa (a minority of these patients have genetic disorders of retinal phosphodiesterases). If prescribed, this should be done with caution. (see **ACTION AND CLINICAL PHARMACOLOGY**).

A small percentage of patients experience visual effects (e.g. impairment of colour discrimination, increased perception to light, blurred vision, eye pain, ocular redness) after taking sildenafil citrate. If this happens, then the patient should not operate a motor vehicle or any heavy machinery until the adverse effects disappear (see **ACTION AND CLINICAL PHARMACOLOGY**).

Rare cases of central serous chorioretinopathy have been reported during the post-marketing period in temporal association with the use of sildenafil citrate. It is not known if medical and other factors were reported that may have also played a role in the development of the condition. It is not possible to determine whether the development of the condition was related directly to the use of sildenafil, to the patient's possible underlying risk factors, a combination of these factors, or to other factors. These cases of central serous chorioretinopathy in patients receiving sildenafil did not provide evidence of serious or permanent alteration in visual function. (See **Post-Market Adverse Drug Reactions**).

Otologic

Sudden decrease or loss of hearing has been reported in a few number of postmarketing and clinical trials cases with the use of PDE5 inhibitors, including sildenafil citrate. These events, which may be accompanied by tinnitus and dizziness, have been reported in temporal association to the intake of PDE5 inhibitors, including sildenafil citrate. In some of the cases, medical conditions and other factors were reported that may have also played a role in the otologic adverse events. In many cases, medical follow-up information was limited. It is not possible to determine whether these events are related directly to the use of PDE5 inhibitors or to other factors (see **ADVERSE REACTIONS, Post-Market Adverse Drug Reactions** and **PART III: CONSUMER INFORMATION**). Physicians should advise patients to stop taking **ACCEL-SILDENAFIL** and seek prompt medical attention in case of sudden decrease or loss of hearing.

Renal

In volunteers with mild ($CL_{Cr} = 50-80$ mL/min) and moderate ($CL_{Cr} = 30-49$ mL/min) renal impairment, the pharmacokinetics of a single oral dose of sildenafil citrate (50 mg) was not altered. In volunteers with severe ($CL_{Cr} < 30$ mL/min) renal impairment, sildenafil clearance was reduced, resulting in increases in AUC (100%) and C_{max} (88%) compared to age-matched volunteers with no renal impairment.

A starting dose of 25 mg should be considered in patients with severe renal impairment (see **ACTION AND CLINICAL PHARMACOLOGY, DOSAGE AND ADMINISTRATION**).

Sexual Function/Reproduction

Although **priapism** had not been reported during clinical trials, prolonged erection greater than 4 hours and priapism (painful erections greater than 6 hours in duration) have been reported infrequently during the post-marketing surveillance of sildenafil citrate. In the event of an erection that persists longer than 4 hours, the patient should seek immediate medical assistance. If priapism is not treated immediately, penile tissue damage and permanent loss of potency could result (see **ADVERSE REACTIONS**).

Agents for the treatment of erectile dysfunction should be used with caution in patients with anatomical deformation of the penis (such as angulation, cavernosal fibrosis or Peyronie's disease) or in patients who have conditions which may predispose them to priapism (such as sickle cell anemia, multiple myeloma or leukemia).

The safety and efficacy of combinations of sildenafil citrate with other PDE5 inhibitors, or other

pulmonary arterial hypertension (PAH) treatments containing sildenafil, or other agents for the treatment of erectile dysfunction have not been studied. Therefore, the use of such combinations is not recommended.

Skin/Appendages

Rare cases of Stevens-Johnson's Syndrome (SJS) and Toxic Epidermal Necrolysis (TEN) have been reported during the post-marketing period.

Special Populations

Women, Nursing Mothers, Pregnancy: Sildenafil citrate is not indicated for use in women. There are no adequate and well-controlled studies in pregnant or lactating women.

Pediatrics: Sildenafil citrate is not indicated for use in children.

Geriatrics (> 65 years of age): Healthy elderly volunteers had a reduced clearance of sildenafil, with free plasma concentrations approximately 40% greater than those seen in younger volunteers (18 to 45 years). Since higher plasma levels may increase both the pharmacological action and incidence of some adverse events, a starting dose of 25 mg should be considered (see **[ACTION AND CLINICAL PHARMACOLOGY, DOSAGE AND ADMINISTRATION](#)**).

Driving and Operating Machinery: As dizziness and altered vision were reported in clinical trials with sildenafil, patients should be aware of how they react to **ACCEL-SILDENAFIL**, before driving or operating machinery. The effect of sildenafil on the ability to drive and use machinery has not been studied.

ADVERSE REACTIONS

Adverse Drug Reaction Overview

Pre-Marketing Experience:

Sildenafil citrate was administered to over 3700 patients (aged 19-87 years) during clinical trials worldwide. Over 550 patients were treated for longer than one year.

In placebo-controlled clinical studies, the discontinuation rate due to adverse events for sildenafil citrate (2.5%) was not significantly different from placebo (2.3%). The adverse events were generally transient and mild to moderate in nature.

Clinical Trial Adverse Drug Reactions

Because clinical trials are conducted under very specific conditions the adverse reaction rates observed in the clinical trials may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse drug reaction information from clinical trials is useful for identifying drug-related adverse events and for approximating rates.

In trials of all designs, adverse events reported by patients receiving sildenafil citrate were generally similar. In fixed-dose studies, the incidence of some adverse events increased with dose. The nature of the adverse events in flexible-dose studies, which more closely reflect the recommended dosage regimen, was similar to that for fixed-dose studies.

When sildenafil citrate was taken as recommended (on an as-needed basis) in flexible-dose, placebo-controlled clinical trials, the following adverse events were reported:

Table 1. Adverse Events Reported by $\geq 2\%$ of Patients Treated with sildenafil citrate or Placebo in PRN Flexible-Dose Phase II/III Studies

Adverse Event	Percentage of Patients Reporting	
	SILDENAFIL CITRATE (n=734)	PLACEBO (n=725)
Headache	15.8%	3.9%
Flushing	10.5%	0.7%
Dyspepsia	6.5%	1.7%
Nasal Congestion	4.2%	1.5%
Respiratory Tract Infection	4.2%	5.4%
Flu Syndrome	3.3%	2.9%
Urinary Tract Infection	3.1%	1.5%
Abnormal Vision*	2.7%	0.4%
Diarrhea	2.6%	1.0%
Dizziness	2.2%	1.2%
Rash	2.2%	1.4%
Back Pain	2.2%	1.7%
Arthralgia	2.0%	1.5%

*Abnormal Vision: Mild and transient changes, predominantly impairment of colour discrimination (blue/green), but also increased perception to light or blurred vision.

At doses above the recommended dose range, adverse events were similar to those detailed above but generally were reported more frequently.

Less Common Clinical Trial Adverse Drug Reactions (<2%)

The following events occurred in <2% of patients in in phase II/III controlled clinical trials where a causal relationship is uncertain:

Autonomic: sweating, dry mouth;

<i>Cardiovascular:</i>	abnormal electrocardiogram, angina pectoris, arrhythmia, AV block, cardiac arrest, cardiomyopathy, heart failure, hypertension, hypotension, palpitation, postural hypotension, myocardial ischemia, syncope, tachycardia, varicose vein, vascular anomaly;
<i>Central & Peripheral Nervous System:</i>	tremor, abnormal dreams, anxiety, agitation, ataxia, depression, insomnia, nervousness, somnolence, paresthesia, vertigo, speech disorder, reflexes decreased, hyperesthesia, neuropathy, migraine, myasthenia, oculogyric crisis, neuralgia, hypertonia;
<i>Gastrointestinal:</i>	vomiting, gastritis, gastrointestinal disorder, flatulence, increased appetite, gastroenteritis, stomatitis, eructation, dysphagia, colitis, glossitis, constipation, rectal hemorrhage, mouth ulceration, esophagitis, rectal disorder, gingivitis, tooth disorder;
<i>Hematopoietic:</i>	anemia and leukopenia;
<i>Liver/Biliary:</i>	liver function tests abnormal, ALT increased;
<i>Metabolic/Nutritional:</i>	edema, thirst, gout, hyperuricemia, hypoglycemic reaction, unstable diabetes, hyperglycemia, hyperlipidemia, hypernatremia;
<i>Musculoskeletal:</i>	myalgia, bone disorder, arthrosis, arthritis, tendon rupture, tenosynovitis, bone pain, joint disorder, synovitis;
<i>Respiratory:</i>	asthma, dyspnea, laryngitis, pharyngitis, sinusitis, bronchitis, respiratory disorder, carcinoma of lung, sputum increased, cough increased;
<i>Skin/Appendages:</i>	skin carcinoma, skin disorder, skin hypertrophy, skin ulcer, contact dermatitis, exfoliative dermatitis, pruritus, urticaria, photosensitivity reaction, nail disorder, acne, herpes simplex, furunculosis;
<i>Special Senses:</i>	Sudden decrease or loss of hearing, mydriasis, conjunctivitis, photophobia, eye pain, tinnitus, ear pain, lacrimation disorder, eye disorder, eye hemorrhage, ear disorder, cataract, dry eyes;
<i>Urogenital:</i>	penile erection, other sexual dysfunction, cystitis, nocturia, balanitis, urinary frequency, breast enlargement, prostatic disorder, testis disorder, urinary incontinence, urinary tract disorder, urine abnormality, abnormal ejaculation, genital edema and anorgasmia;
<i>Vascular Disorders:</i>	cerebrovascular disorder, cerebral thrombosis;

General: face edema, peripheral edema, chills, allergic reaction, asthenia, pain, infection, shock, hernia, accidental fall, abdominal pain, chest pain, accidental injury, intentional overdose.

Myocardial Infarction and Cardiovascular Mortality

In an analysis of double blind placebo controlled clinical trials encompassing over 700 person-years of observation on placebo and over 1300 person-years on sildenafil, there were no differences in the incidence rate of myocardial infarction (MI) or in the rate of cardiovascular mortality for patients receiving sildenafil compared to those receiving placebo. The rates of MI were 1.1 per 100 person-years for men receiving sildenafil and for those receiving placebo. The rates of cardiovascular mortality were 0.3 per 100 person-years for men receiving sildenafil and those receiving placebo.

Clinical Trial Adverse Drug Reactions Reported in 74 Double-Blind Placebo-Controlled Phase II/III/IV Studies

When sildenafil citrate was taken as recommended in 74 randomized double-blind, placebo-controlled (DBPC) Phase II/III/IV studies, adverse reactions reported by $\geq 2\%$ of patients treated with sildenafil citrate (n=9,570) and more frequently than placebo (n=7,237) were headache, flushing, dyspepsia, nasal congestion and dizziness. The nature and frequency of adverse reactions reported by $\geq 2\%$ of patients in this pooled dataset of 74 DBPC studies was consistent with the adverse reactions reported in the 6 flexible-dose studies detailed above in **Table 1**.

The following adverse reactions occurred in $< 2\%$ of patients in the 74 DBPC clinical trials.

Cardiac disorders: palpitations, tachycardia;

Eye disorders: vision blurred, chromatopsia, cyanopsia, photophobia, visual disturbance, photopsia, ocular hyperaemia, eye pain, visual brightness, abnormal sensation in eye, asthenopia, conjunctival hyperaemia, dry eye, erythroptosis, eye disorder, eye irritation, eye edema, eyelid edema, eye swelling, halo vision, xanthopsia;

Gastrointestinal disorders: nausea, dry mouth, abdominal pain upper, vomiting, gastroesophageal reflux disease, oral hypoesthesia;

General conditions and administration site conditions: feeling hot, irritability;

Immune system disorders: hypersensitivity;

Infections and infestations: rhinitis;

Investigations:	heart rate increased;
Musculoskeletal and connective tissue disorders:	pain in extremity, myalgia;
Nervous system disorders:	syncope, somnolence;
Reproductive system and breast disorders:	erection increased;
Respiratory, thoracic and mediastinal disorders:	epistaxis, sinus congestion, nasal oedema, nasal dryness, throat tightness;
Skin and subcutaneous tissue disorders:	rash;
Vascular disorders:	hot flush, hypotension.

Post-Market Adverse Drug Reactions

Reports of adverse events temporally associated with sildenafil citrate during post-marketing surveillance that are not listed above and for which the causal relationship is unknown, include the following:

Cardiovascular:

Epistaxis; serious cardiovascular events - including myocardial infarction, sudden cardiac death, ventricular arrhythmia, cerebrovascular hemorrhage, and transient ischemic attack - have been reported. Most of these patients had pre-existing cardiovascular risk factors. Many of these events were reported to occur during or shortly after sexual activity, and a few were reported to occur shortly after the use of sildenafil citrate without sexual activity. Others were reported to have occurred hours to days after the use of sildenafil citrate with sexual activity. It is not possible to determine whether these events are related directly to sildenafil citrate, to sexual activity, to the patient's underlying cardiovascular disease, to combination of these factors, or to other factors (see **WARNINGS AND PRECAUTIONS**).

Central & Peripheral Nervous System: seizure, seizure recurrence, transient global amnesia;

Gastrointestinal: vomiting;

Urogenital: prolonged erection, priapism (see **WARNINGS AND PRECAUTIONS**) and hematuria;

Skin / Appendages: Stevens-Johnson Syndrome (SJS) and Toxic Epidermal Necrolysis

(TEN)

Special Senses:

diplopia, temporary vision loss/decreased vision, blurred vision, Non-Arteritic Anterior Ischemic Optic Neuropathy (NAION), retinal vein occlusion, visual field defect, eye pain, ocular redness or bloodshot appearance, ocular burning, ocular swelling/pressure, increased intraocular pressure, retinal vascular disease of bleeding, vitreous detachment/traction and paramacular edema.

Cases of sudden decrease or loss of hearing have been reported post-marketing in temporal association with the use of PDE5 inhibitors, including sildenafil citrate. In some of the cases, medical conditions and other factors were reported that may have also played a role in the otologic adverse events. In many cases, medical follow-up information was limited. It is not possible to determine whether these events are related directly to the use of sildenafil citrate, to the patient's underlying risk factors for hearing loss, a combination of these factors, or to other factors (see **WARNINGS AND PRECAUTIONS, ADVERSE REACTIONS** and **PART III: CONSUMER INFORMATION**).

Rare cases of central serous chorioretinopathy have been reported during the post-marketing period in temporal association with the use of sildenafil citrate. It is not known if medical and other factors were reported that may have also played a role in the development of the condition. It is not possible to determine whether the development of the condition was related directly to the use of sildenafil, to the patient's possible underlying risk factors, a combination of these factors, or to other factors. These cases of central serous chorioretinopathy in patients receiving sildenafil did not provide evidence of serious or permanent alteration in visual function. (see **WARNINGS AND PRECAUTIONS**).

DRUG INTERACTIONS

Serious Drug Interactions

- Use of organic nitrates in any form is absolutely contraindicated (see **CONTRAINDICATIONS** section)

Overview

***In vitro* studies:**

Sildenafil metabolism is principally mediated by the cytochrome P-450 (CYP) isoforms 3A4 (major route) and 2C9 (minor route) (see **ACTION AND CLINICAL PHARMACOLOGY**). Therefore, inhibitors of these isoenzymes may reduce sildenafil clearance and inducers of these isoenzymes may increase sildenafil clearance.

Sildenafil is a weak inhibitor of the cytochrome P-450 isoforms 1A2, 2C9, 2C19, 2D6, 2E1 and 3A4 (IC₅₀>150 μM). Given sildenafil peak plasma concentrations of approximately 1 μM after recommended doses, it is unlikely that sildenafil citrate will alter the clearance of the substrates of

these isoenzymes.

***In vivo* studies:**

Sildenafil (50 mg) did not potentiate the hypotensive effect of alcohol in healthy volunteers with mean maximum blood alcohol levels of 0.08%.

Drug-Drug Interactions

Effects of Other Drugs on sildenafil citrate

Pharmacokinetic data from patients in clinical trials showed no effect on sildenafil pharmacokinetics of CYP2C9 inhibitors (such as tolbutamide, warfarin), CYP2D6 inhibitors (such as selective serotonin reuptake inhibitors, tricyclic antidepressants), thiazide and related diuretics, angiotensin converting enzyme (ACE) inhibitors, and calcium channel blockers. The AUC of the active metabolite, N-desmethyl sildenafil, was increased 62% by loop and potassium-sparing diuretics and 102% by nonspecific beta-blockers. These effects on the metabolite are not expected to be of clinical consequence.

In healthy male volunteers, there was no evidence of an effect of azithromycin (500 mg daily for 3 days) on the AUC, C_{max} , T_{max} , elimination rate constant, or subsequent half-life of sildenafil or its principle circulating metabolite.

CYP3A4 Inhibitors

The concomitant use of potent cytochrome P-450 3A4 inhibitors (e.g. erythromycin, saquinavir, ritonavir, ketoconazole, itraconazole) as well as the non-specific CYP inhibitor, cimetidine, is associated with increased plasma levels of sildenafil (see **DOSAGE AND ADMINISTRATION, DETAILED PHARMACOLOGY**).

When a single 100 mg dose of sildenafil citrate was administered with erythromycin, a moderate CYP3A4 inhibitor, at steady state (500 mg b.i.d. for 5 days), there was a 182% increase in sildenafil systemic exposure (AUC).

When the dose of sildenafil for subjects receiving potent CYP3A4 inhibitors was administered as recommended, the maximum free plasma sildenafil concentration did not exceed 200 nM for any individual and was consistently well tolerated.

In a study of healthy male volunteers, co-administration of the endothelin antagonist bosentan, (an inducer of CYP3A4 [moderate], CYP2C9 and possibly of CYP2C19) at steady state (125 mg twice a day) with sildenafil at steady state (80 mg three times a day) resulted in 62.6% and 55.4% decrease in sildenafil AUC and C_{max} , respectively. Sildenafil increased bosentan AUC and C_{max} by 49.8% and 42%, respectively. Concomitant administration of strong CYP3A4 inducers, such as rifampin, is expected to cause greater decreases in plasma concentrations of sildenafil.

Cimetidine (800 mg), a cytochrome P450 inhibitor and a non-specific CYP3A4 inhibitor, caused a 56% increase in plasma sildenafil concentrations when co-administered with sildenafil citrate (50 mg) to healthy volunteers.

Population pharmacokinetic analysis of clinical trial data indicated a reduction in sildenafil clearance when co-administered with CYP3A4 inhibitors (such as ketoconazole, erythromycin, cimetidine). However, there was no increased incidence of adverse events in these patients.

HIV Protease Inhibitor

In addition, coadministration of the HIV protease inhibitor saquinavir, also CYP3A4 inhibitor, at steady state (1200 mg t.i.d) with sildenafil (100 mg single dose) resulted in a 140 % increase in sildenafil C_{max} and a 210% increase in sildenafil AUC. Sildenafil had no effect on saquinavir pharmacokinetics. Stronger CYP3A4 inhibitors such as ketoconazole, itraconazole would be expected to have still greater effects (see **DOSAGE AND ADMINISTRATION**).

Coadministration with the HIV protease inhibitor ritonavir, which is a highly potent P-450 inhibitor, at steady state (500 mg b.i.d) with sildenafil (100 mg single dose) resulted in a 300% (4-fold) increase in sildenafil C_{max} and a 1000% (11-fold) increase in sildenafil plasma AUC. At 24 hours the plasma levels of sildenafil were still approximately 200 ng/mL, compared to approximately 5 ng/mL when sildenafil was dosed alone. This is consistent with the marked effects of ritonavir on a broad range of P-450 substrates. Sildenafil had no effect on ritonavir pharmacokinetics (see **DOSAGE AND ADMINISTRATION**).

CYP3A4 Inducers

It can be expected that concomitant administration of CYP3A4 inducers, such as rifampin, will decrease plasma levels of sildenafil.

CYP2C9 Substrate

No significant interactions were shown with tolbutamide (single 250 mg dose) or warfarin (single 40 mg dose), both of which are metabolized by CYP2C9, when co-administered with 50 mg sildenafil.

Antacids

In normal healthy male volunteers, co-administration of single doses of antacid (magnesium hydroxide/aluminium hydroxide) with sildenafil did not affect the AUC, C_{max} , elimination rate constant, or subsequent half-life of sildenafil. The T_{max} was reduced by 0.42 hours.

Effect of sildenafil citrate on Other Drugs

Alpha-blockers

In three specific drug-drug interaction studies, the alpha-blocker doxazosin (4 mg and 8 mg) and sildenafil (25 mg, 50 mg, or 100 mg) were administered simultaneously to patients with benign prostatic hyperplasia (BPH) stabilized on doxazosin therapy. In these study populations, mean additional reductions of supine blood pressure of 7/7 mmHg, 9/5 mmHg, and 8/4 mmHg, and mean additional reductions of standing blood pressure of 6/6 mmHg, 11/4 mmHg, and 4/5 mmHg, for 25 mg, 50 mg, or 100 mg respectively, were observed. When sildenafil and doxazosin were administered simultaneously to patients stabilized on doxazosin therapy, there were infrequent reports of patients who experienced symptomatic postural hypotension. These reports included dizziness and lightheadedness, but not syncope. Concomitant administration of sildenafil to patients taking alpha-blocker therapy may lead to symptomatic hypotension in a few susceptible individuals (see **WARNINGS AND PRECAUTIONS**).

Some alpha-blockers and antidepressants have reported priapism or prolonged/painful erections in their labels.

Bleeding Time

Sildenafil citrate (50 mg) did not potentiate the increase in bleeding time, measured using a standard simplate method, caused by acetylsalicylic acid (150 mg).

Use with Other Concomitant Therapies:

Antihypertensives

When sildenafil citrate (100 mg) was co-administered with amlodipine, 5 mg or 10 mg, in hypertensive patients, the mean additional reduction of supine blood pressure was 8 mm Hg systolic and 7 mm Hg diastolic (see [ACTION AND CLINICAL PHARMACOLOGY](#)).

Patients on multiple antihypertensive medications were included in the pivotal clinical trials for sildenafil citrate. Analysis of the safety database was carried out after pooling of the following classes of antihypertensive medication: diuretics, beta-blockers, ACE inhibitors, angiotensin II antagonists, antihypertensive medicinal products (vasodilator and centrally-acting), adrenergic neurone blockers, calcium channel blockers and alpha-adrenoceptor blockers. The analysis showed no differences in the adverse effect profile of patients taking sildenafil citrate with and without antihypertensive medication.

A large controlled study was performed in men with erectile dysfunction and arterial hypertension who were taking combinations of diuretics, beta blockers, ACE inhibitors and calcium channel blockers. The incidence rate of all adverse events, including those possibly related to hypotensive episodes, was consistent with observations in other patient populations. Also, there was no evidence of an increased incidence rate of any adverse event in the subgroups taking 2 antihypertensive agents and 3 or more antihypertensive agents. There was no indication of additional safety risk of sildenafil use in this subject population (see [DETAILED PHARMACOLOGY](#)).

Bosentan

Sildenafil at steady state (80 mg three times a day) resulted in a 49.8% increase in bosentan AUC and a 42% increase in bosentan C_{max} (125 mg twice a day) (see [DRUG INTERACTIONS](#)).

Drug-Food Interactions

Grapefruit juice being a weak inhibitor of CYP3A4 gut wall metabolism may give rise to modest increases in plasma levels of sildenafil.

ACCEL-SILDENAFIL (sildenafil citrate) can be taken with or without food. However, when sildenafil citrate is taken with a high-fat meal, the rate of absorption is reduced with a mean delay in T_{max} of 60 minutes and a mean reduction in C_{max} of 29%. AUC is decreased by 11%. The patient may find that it takes longer to work if taken with a high-fat meal (see [ACTION AND CLINICAL PHARMACOLOGY](#)).

DOSAGE AND ADMINISTRATION

Dosing Considerations

The following factors are associated with increased plasma levels (AUC) of sildenafil:

- age 65 years or over (40%)
- hepatic impairment (e.g. cirrhosis: 84%)
- severe renal impairment (e.g. creatinine clearance <30 mL/min: 100%)
- concomitant use of potent cytochrome P-450 3A4 inhibitors (e.g. erythromycin: 182%; saquinavir: 210%; ritonavir: 1000%). It can also be expected that more potent cytochrome P-450 3A4 inhibitors such as ketoconazole and itraconazole would result in increased levels of sildenafil.

(see **Recommended Dose and Dosage Adjustment, ACTION AND CLINICAL PHARMACOLOGY, WARNINGS AND PRECAUTIONS**).

Sildenafil citrate has been shown to potentiate the hypotensive effects of nitrates in healthy volunteers and in patients, and is therefore contraindicated in patients who are taking any type of nitrate drug therapy, or who utilize short-acting nitrate-containing medications, due to the risk of developing potentially life-threatening hypotension. The use of organic nitrates, either regularly and/or intermittently, in any form (e.g. oral, sublingual, transdermal, by inhalation) is absolutely contraindicated (see **ACTION AND CLINICAL PHARMACOLOGY, CONTRAINDICATIONS**).

Recommended Dose and Dosage Adjustment

For most patients, the recommended dose of **ACCEL-SILDENAFIL** is 50 mg taken as needed. The maximum recommended dose is 100 mg. Dosage may be decreased to 25 mg if necessary.

Since higher plasma levels may increase both efficacy and the incidence of adverse events, a starting dose of 25 mg should be considered in patients, age 65 years or over, on concomitant CYP3A4 inhibitors, with severe renal impairment, with hepatic impairment and on ritonavir (see **Dosing Considerations** above, **ACTION AND CLINICAL PHARMACOLOGY, WARNINGS AND PRECAUTIONS**).

The concomitant use of the potent cytochrome P-450 3A4 inhibitor, ritonavir is associated with a 1000% (11-fold) increase in plasma levels (AUC) of sildenafil. Given the extent of the interaction with patients receiving concomitant therapy with ritonavir, it is recommended not to exceed a maximum single dose of 25 mg of sildenafil in a 48-hour period (see **WARNINGS AND PRECAUTIONS**).

Administration

To be taken as needed approximately 30 – 60 minutes before sexual activity. However, **ACCEL-SILDENAFIL** may be taken anywhere from 0.5 hour to 4 hours before sexual activity. The maximum recommended dosing frequency is once per day.

OVERDOSAGE

For management of a suspected drug overdose, contact your regional Poison Control Centre immediately.

In studies with healthy volunteers of single doses of up to 800 mg, adverse events were similar to those seen at lower doses but incidence rates and severities were increased. In cases of overdose, standard supportive measures should be adopted as required. Renal dialysis is not expected to accelerate clearance as sildenafil is highly bound to plasma proteins and not eliminated in the urine.

Treatment of Priapism

Patients should be instructed to report any erections persisting for more than 4 hours to a physician. The treatment of priapism/prolonged erection should be according to established medical practice. Physicians may refer to two suggested protocols for detumescence presented below.

Detumescence Protocols

- 1) Aspirate 40 to 60 mL blood from either left or right *corpora* using vacutainer and holder for drawing blood. Patient will often detumescence while aspirating. Apply ice for 20 minutes post aspiration if erection remains.

If procedure 1) is unsuccessful, then try procedure 2).

- 2) Put patient in supine position. Dilute 10 mg phenylephrine into 20 mL distilled water for injection (0.05%). With an insulin syringe, inject 0.1 to 0.2 mL (50-100 µg) into the corpora every 2 to 5 minutes, until the detumescence occurs. The occasional patient may experience transient bradycardia and hypertension when given phenylephrine injections, therefore monitor patient's blood pressure and pulse every 10 minutes. Patients at risk include those with cardiac arrhythmias and diabetes. Refer to the prescribing information for phenylephrine before use. **Do not give phenylephrine to patients on MAO inhibitors.** When phenylephrine is used within the first 12 hours of erection, the majority of patients will respond.

If procedure 2) is unsuccessful, then try procedure 3).

- 3) If the above measures fail to detumescence the patient, a urologist should be consulted as soon as possible, especially if the erection has been present for many hours. If priapism is not treated immediately, penile tissue damage and/or permanent loss of potency may result.

ACTION AND CLINICAL PHARMACOLOGY

Mechanism of Action

Sildenafil citrate is a cGMP-specific phosphodiesterase type 5 (PDE5) inhibitor, used for the treatment of male erectile dysfunction.

The physiological mechanism responsible for erection of the penis involves the release of nitric

oxide (NO) in the *corpus cavernosum* in response to sexual stimulation. Nitric oxide then activates the enzyme guanylate cyclase, which results in increased levels of cyclic guanosine monophosphate (cGMP), producing smooth muscle relaxation in the *corpus cavernosum* and allowing inflow of blood.

Sildenafil has no direct relaxant effect on isolated human *corpus cavernosum*, but enhances the effect of NO by inhibiting PDE5, which is responsible for the biodegradation of cGMP in the *corpus cavernosum*. When sexual stimulation causes local release of NO, inhibition of PDE5 by sildenafil produces increased levels of cGMP in the *corpus cavernosum*, resulting in smooth muscle relaxation and increased inflow of blood to the *corpus cavernosum*. Sildenafil, at recommended doses, has no effect in the absence of sexual stimulation.

Studies *in vitro* have shown that sildenafil has between 10 and 10,000-fold greater selectivity for PDE5 than for other phosphodiesterase isoforms namely PDEs 1, 2, 3, 4, and 6 and greater than 700-fold effect on PDE7-PDE11. In particular, sildenafil has greater than 4,000-fold selectivity for PDE5 over PDE3, the cAMP-specific phosphodiesterase isoform involved in the control of cardiac contractility. Sildenafil is about 10-fold as potent for PDE5 compared to PDE6, an isoenzyme found in the retina; this lower selectivity is thought to be the basis for colour vision abnormalities observed with higher doses or plasma levels of sildenafil (see **WARNINGS AND PRECAUTIONS, DETAILED PHARMACOLOGY**).

PDE5 is also found in lower concentrations in platelets, vascular and visceral smooth muscles, and skeletal muscle. The sildenafil-induced inhibition of PDE5 in these tissues appears to be the basis for the enhanced platelet antiaggregatory activity of nitric oxide observed *in vitro*, and inhibition of platelet thrombus formation *in vivo*, and peripheral arterial-venous dilation *in vivo* (see **WARNINGS AND PRECAUTIONS**).

Pharmacodynamics

Effects of sildenafil citrate on Blood Pressure (BP):

Single oral doses of sildenafil (100 mg) administered to healthy volunteers produced decreases in supine blood pressure (mean maximum decrease of 8.3/5.3 mm Hg). The decrease in blood pressure was most notable approximately 1-2 hours after dosing. The effects are not related to dose or plasma levels. Larger effects were recorded among patients receiving concomitant nitrates (see **CONTRAINDICATIONS**).

Effects of sildenafil citrate on Cardiac Parameters:

Single oral doses of sildenafil citrate up to 100 mg in healthy volunteers produced no clinically relevant effects on ECG.

Effects of sildenafil citrate on Erectile Response:

Sildenafil citrate was studied in clinical trials of various designs. In fixed-dose clinical trials, 62%, 74%, and 82% of patients on 25 mg, 50 mg and 100 mg of sildenafil citrate, respectively, reported an improvement in their erections, compared to 25% on placebo (p <0.0001, see **CLINICAL TRIALS**).

In eight double-blind, placebo-controlled, cross-over studies using RigiScan[®] (a device used to objectively measure penile rigidity and duration of erections), erections during sexual stimulation improved significantly on sildenafil citrate compared to placebo. These studies included patients with organic etiologies (such as spinal cord injury and diabetes mellitus), and patients without an established organic cause. Most studies assessed the efficacy of sildenafil citrate approximately 60 minutes post-dose.

All eight studies investigating the effects of sildenafil on penile plethysmography (RigiScan[®]) after visual sexual stimulation (VSS) under laboratory conditions, consistently showed that doses of up to 100 mg resulted in statistically significant improvements in duration of erections of 60% rigidity (considered hard enough for penetrative sexual intercourse), compared with placebo. In patients who respond, the median time to onset of erections (60% rigidity) in response to VSS, was 25 minutes after an oral dose of 50 mg sildenafil. The mean total duration of erections 60% rigidity at the base of the penis was 3, 24 and 32 minutes for subjects receiving placebo, 25 mg and 50 mg doses, respectively, when exposed to VSS for 2 hours.

Sildenafil citrate increases couples' ability to have sexual intercourse (see [CLINICAL TRIALS](#)).

Pharmacokinetics

Absorption: Sildenafil is rapidly absorbed. Maximum observed plasma concentrations are reached within 30 to 120 minutes (median 60 minutes) of oral dosing in the fasted state. The mean absolute bioavailability is 41% (range 25%-63%). The oral pharmacokinetics of sildenafil citrate is proportional over the recommended dose range studied (25 mg to 100 mg).

Sildenafil inhibits the human PDE5 enzyme *in vitro* by 50% at a concentration of 3.5 nM. In man, the mean maximum free plasma concentration of sildenafil following a single oral dose of 100 mg is approximately 18 ng/mL, or 38 nM.

When sildenafil citrate was administered with a high-fat meal, the rate of absorption was significantly decreased, with a 29% reduction in C_{max} and a 60-minute delay in T_{max} . The patient may find that it takes longer to work if taken with a high-fat meal. However, although it was statistically significant (AUC decreased by 11%), the decrease in the extent of absorption was not clinically relevant. The relative bioavailability fed/fasted was 89% (90% CI; 84-94%) (see [DRUG INTERACTIONS](#)).

Distribution: The mean steady state volume of distribution (V_{ss}) for sildenafil is 105 litres, indicating distribution into the tissues. Sildenafil and its major circulating N-desmethyl metabolite are both approximately 96% bound to plasma proteins. Protein binding is independent of total drug concentrations.

Based upon measurements of sildenafil in the semen of healthy volunteers, less than 0.001% of the ingested dose may appear in the semen of patients 90 minutes after drug intake.

Metabolism: Sildenafil is cleared predominantly by the CYP3A4 (major route) and CYP2C9 (minor route) hepatic microsomal isoenzymes. The major circulating metabolite results from

N-demethylation of sildenafil at the N-methyl piperazine moiety. This metabolite has a PDE selectivity profile similar to sildenafil and an *in vitro* potency against PDE5 approximately 50% that of the parent drug. Plasma concentrations of this metabolite are approximately 40% of those seen for sildenafil. The N-desmethyl metabolite is further metabolised, with a terminal half-life of approximately 4 hours.

Excretion: The total body clearance of sildenafil is 41 L/h with a resultant terminal phase half-life of 3-5 hours. After either oral or intravenous administration, sildenafil is excreted as metabolites predominantly in the feces (approximately 80% of administered dose) and to a lesser extent in the urine (approximately 13% of the administered dose).

Special Populations and Conditions

Geriatrics: Healthy elderly subjects (65 years or older) had a reduced clearance of sildenafil, resulting in approximately 90 % higher plasma concentrations of sildenafil and the active N-desmethyl metabolite compared to those seen in healthy younger volunteers (18-45 years). Due to age-differences in plasma protein binding, the corresponding increase in free sildenafil plasma concentration was approximately 40 %.

Hepatic Insufficiency: In volunteers with hepatic cirrhosis (Child-Pugh A and B), sildenafil clearance was reduced, resulting in increases in AUC (85%) and C_{max} (47%) compared to age-matched volunteers with no hepatic impairment. The pharmacokinetics of sildenafil in patients with severe hepatic impairment (Child-Pugh class C) have not been studied.

Since sildenafil clearance is reduced in geriatric patients (65 years or older), patients with renal impairment or patients with hepatic impairment, a starting dose of 25 mg should be considered. Based on efficacy and toleration, the dose may be increased to 50 mg or 100 mg (see **WARNINGS AND PRECAUTIONS, DOSAGE AND ADMINISTRATION**).

Renal Insufficiency: In volunteers with mild ($CL_{cr} = 50-80$ mL/min) and moderate ($CL_{cr} = 30-49$ mL/min) renal impairment, the pharmacokinetics of a single oral dose of sildenafil citrate (50 mg) were not altered. In volunteers with severe ($CL_{cr} < 30$ mL/min) renal impairment, sildenafil clearance was reduced, resulting in increases in AUC (100%) and C_{max} (88%) compared to age-matched volunteers with no renal impairment.

In addition, N-desmethyl metabolite AUC and C_{max} values were significantly increased by 200 % and 79 % respectively in subjects with severe renal impairment compared to subjects with normal renal function.

STORAGE AND STABILITY

Store at controlled room temperature between 15 and 30°C.

SPECIAL HANDLING INSTRUCTIONS

Not Applicable.

DOSAGE FORMS, COMPOSITION AND PACKAGING

ACCEL-SILDENAFIL - 25 mg tablets (sildenafil citrate equivalent to 25 mg of sildenafil per tablet) are elliptic, blue, film-coated tablets, embossed with a “-“ on one side, and supplied as follows:

- Cartons of 4 tablets

ACCEL-SILDENAFIL - 50 mg tablets (sildenafil citrate equivalent to 50 mg of sildenafil per tablet) are elliptic, blue, film-coated tablets, and supplied as follows:

- Cartons of 4 tablets or 8 tablets

ACCEL-SILDENAFIL - 100 mg tablets (sildenafil citrate equivalent to 100 mg of sildenafil per tablet) are elliptic, blue, film-coated tablets, embossed with a “+“ on one side, and supplied as follows:

- Cartons of 4 tablets or 8 tablets

Each tablet contains the following non-medicinal ingredients: calcium hydrogen phosphate anhydrous, FD&C Blue #2/indigo carmine aluminum lake, macrogol, magnesium stearate, microcrystalline cellulose, polyvinyl alcohol, sodium croscarmellose, talc and titanium dioxide.

PART II: SCIENTIFIC INFORMATION

PHARMACEUTICAL INFORMATION

Drug Substance


Proper name: sildenafil citrate

Chemical name:

1-[[3-(6,7-dihydro-1-methyl-7-oxo-3-propyl-1H-pyrazolo[4,3-d]pyrimidin-5-yl)-4-ethoxyphenyl]sulfonyl]-4-methylpiperazine citrate

Molecular formula and molecular mass: $C_{22}H_{30}N_6O_4S \cdot C_6H_8O_7$; 666.7

Structural formula:



Physicochemical properties:	Sildenafil citrate is a white or almost white, slightly hygroscopic, crystalline powder.	
p _{Ka} :	protonation of tertiary amine	6.53
	deprotonation of pyrimidinone moiety	9.17
Partition coefficient:	octanol/water	2.7
Solubility:	Water	3.5 mg/mL
	1M HCl	5.8 mg/mL
	1M NaOH	42.3 mg/mL

CLINICAL TRIALS

Comparative Bioavailability

A randomized, double blind, two-treatment, two-sequence, two-period, single-dose, crossover study comparing ACCEL-SILDENAFIL (sildenafil citrate) 100 mg tablets (Accel Pharma Inc.) with Viagra (sildenafil citrate) 100 mg tablets (Pfizer Canada Inc.) was conducted in 24 healthy,

adult, male subjects under fasting conditions. A summary of the bioavailability data is presented in the following table.

Sildenafil (1 x 100 mg sildenafil citrate) From measured data Geometric Mean Arithmetic Mean (CV%)				
Parameter	Test*	Reference†	% Ratio of Geometric Means	90% Confidence Interval
AUC _T (ng·h/mL)	1355.94 1507.34 (47.13)	1324.75 1449.16 (44.77)	102.35	93.76 – 111.74
AUC _I (ng·h/mL)	1371.74 1524.43 (47.07)	1341.76 1467.32 (44.62)	102.23	93.71 – 111.53
C _{max} (ng/mL)	384.26 428.95 (51.52)	366.78 404.05 (47.17)	104.76	92.08 – 119.20
T _{max} § (h)	0.92 (0.50 – 3.50)	1.13 (0.50 – 4.00)		
T _{1/2} € (h)	4.23 (29.37)	4.29 (25.11)		

* Accel-Sildenafil 100 mg Tablets

† Viagra (sildenafil citrate) 100 mg Tablets (Pfizer Canada Inc.) were purchased in Canada.

§ Expressed as the median (range) only

€ Expressed as the arithmetic mean (CV%) only

Study demographics and trial design

Sildenafil citrate was evaluated at doses including 25 mg, 50 mg and 100 mg in 21 randomized, double-blind, placebo-controlled trials of up to 6 months duration. In these studies, sildenafil citrate was administered to more than 3000 patients aged 19 to 87 years, with ED of various etiologies (organic, psychogenic, mixed) with a mean duration of 5 years.

This patient population included men with the following concomitant conditions: angina, benign prostatic hyperplasia (BPH), depression, type I and type II diabetes mellitus, hypertension, previous myocardial infarction, radical prostatectomy, spinal cord injury, transurethral resection of the prostate (TURP), and vasculogenic disease.

Efficacy was demonstrated in all 21 studies and results were consistent regardless of baseline severity, etiology and age. Efficacy was maintained over the long-term (1 year). Sildenafil citrate was effective in a broad range of ED patients, including those with a history of coronary artery disease (myocardial infarction, angina), hypertension, other cardiac disease (arrhythmias, congestive heart failure), peripheral vascular disease, diabetes mellitus, depression, coronary artery bypass graft (CABG), radical prostatectomy and TURP, and in patients taking antidepressants, antihypertensives, antipsychotics, and diuretics.

Study results

Sildenafil citrate was studied in clinical trials of various designs. In fixed-dose clinical trials, 62%, 74%, and 82% of patients on 25 mg, 50 mg and 100 mg of sildenafil citrate, respectively, reported an improvement in their erections, compared to 25% on placebo (see [Figure 1](#)).




Figure 1 - Percentage of Patients reporting an Improvement in Erections

The primary efficacy endpoints were the ability to both achieve and maintain an erection sufficient for sexual intercourse, as measured by patient responses to the International Index of Erectile Function (IIEF), a sexual function questionnaire. The results from the partner questionnaire corroborated the data from the study subjects, with analyses showing clear treatment related improvements in the ability to achieve and maintain erections.

Responses to the IIEF are scored on a five-point scale ranging from ‘almost never/never’ (1) to ‘almost always/always’ (5), with a score of (0) assigned for no attempts at sexual intercourse. During IIEF validation, scores for the primary efficacy endpoints for men without erectile dysfunction were 4.38 and 4.34, respectively. Compared to baseline treatment over 12 weeks, sildenafil citrate patients reported the following statistically significant changes (see [Figure 2](#)).




Figure 2 - Effect of sildenafil citrate on Ability to Achieve and Maintain an Erection Sufficient for Sexual Intercourse

Men with untreated ED have lower scores (Figure 3, Bar 1) for all sexual function domains of the IIEF (erection, orgasm, desire, overall satisfaction, intercourse satisfaction). In these men, sildenafil citrate restores the values of the domains (Figure 3, Bar 2) towards the values of age matched controls without ED (Figure 3, Bar 3).




Figure 3 - Effect of sildenafil citrate on Male Sexual Function Domains of the IIEF

Sildenafil citrate increases couples' ability to have sexual intercourse. With sildenafil citrate, 64%, 67% and 72% of attempts resulted in successful sexual intercourse on doses of 25 mg, 50 mg, and 100 mg, respectively, compared to 23% on placebo. Of sildenafil citrate patients with one or more successful attempt at intercourse, 81% of attempts were successful.

The efficacy of sildenafil citrate was maintained over time. In a long-term, open-label trial of 12-month duration, 88% (256/292) of patients reported that sildenafil citrate treatment improved their erections.

Eighty-seven percent (87%) of patients completed the one-year study. When these patients were

followed for an additional year (total exposure of 24-months), oral sildenafil was an effective, well tolerated treatment for erectile dysfunction of organic, psychogenic or mixed aetiology.

In a controlled clinical study which reflects the recommended dosage regimen, 74% of patients were taking sildenafil citrate 100 mg after 12 weeks of treatment, compared to 23% and 3% taking sildenafil citrate 50 mg and 25 mg, respectively.

Other Patient Populations:

Across all trials, sildenafil citrate improved the erections of 59% of diabetic patients, and 43% of radical prostatectomy patients (versus 16% and 15% on placebo, respectively). This was assessed using the GAQ.

In a study of patients with spinal cord injury, sildenafil citrate improved the ability to have sexual intercourse in 80% of patients versus 10% on placebo.

Subgroup analyses of responses to a global improvement question in patients with psychogenic etiology in two fixed-dose studies and two titrations studies showed 84% of sildenafil citrate patients reported improvement in erections compared with 26% of placebo patients.

These studies confirm that sildenafil enhances the erectile response to sexual stimulation in subjects with erectile dysfunction (ED) of psychogenic and broad-spectrum etiology, including patients with diabetes mellitus and with spinal cord injury.

Use with Other Concomitant Therapies:

Antihypertensives

A large (n=568) randomized, double-blind, placebo-controlled, parallel group, multicenter, flexible-dose study (sildenafil citrate up to 100 mg) in males with erectile dysfunction and arterial hypertension taking 2 or more antihypertensive agents was conducted (the majority of these were diuretics, beta blockers, ACE inhibitors and calcium channel blockers). Fifty-eight percent of the patients were taking 2 antihypertensive agents and 42% were taking 3 or more antihypertensive agents composed of similar groups of drugs for blood pressure control. Sildenafil citrate improved the erections in 71% of men compared to 18% in the placebo group, and 62% of attempts at sexual intercourse were successful with sildenafil citrate compared to 26% on placebo. The incidence rate of all adverse events, including those possibly related to hypotensive episodes, was consistent with observations in other patient populations. Also, there was no evidence of an increased incidence rate of any adverse event in the subgroups taking 2 antihypertensive agents and 3 or more antihypertensive agents. There was no indication of additional safety risk of sildenafil use in this subject population (see **WARNINGS AND PRECAUTIONS**).

DETAILED PHARMACOLOGY

Human

Pharmacodynamic Studies

Oral doses of sildenafil of 50 mg, 100 mg and 200 mg produced statistically significant decreases in

supine systolic and diastolic blood pressure (a mean maximum decrease of approximately 8 mm Hg and 5 mm Hg, respectively) compared with placebo, with no effect on pulse rate. The mean maximum fall in systolic and diastolic blood pressure occurred at peak plasma levels (approximately 1 hour post-dose), and there was a tendency for blood pressure to return to baseline values by 4 hours post-dose.

In healthy volunteers, there were no clinically significant changes in cardiac index (derived from bio-impedance measures of cardiac output) up to 12 hours post-dose for sildenafil administered orally (100 mg, 150 mg and 200 mg), nor intravenously (20-80 mg), compared with placebo. Sildenafil has both arteriodilator and venodilator effects on the peripheral vasculature.

In a study of the hemodynamic effects of a single oral 100 mg dose of sildenafil in 14 patients with severe coronary artery disease (CAD) (>70% stenosis of at least one coronary artery), the mean resting systolic and diastolic blood pressures decreased by 7% and 6%, respectively compared to baseline. Mean pulmonary systolic blood pressure decreased by 9%. Sildenafil had no effect on cardiac output, and did not impair blood flow through the stenosed coronary arteries, and resulted in improvement (approximately 13%) in adenosine-induced coronary flow reserve (in both stenosed and reference arteries).

In a double-blind study, 144 patients with erectile dysfunction and chronic stable angina limited by exercise, not receiving chronic oral nitrates, were randomized to a single dose of placebo or sildenafil citrate 100 mg 1 hour prior to exercise testing. The primary endpoint was time to limiting angina in the evaluable cohort. The mean times (adjusted for baseline) to onset of limiting angina were 423.6 and 403.7 seconds for sildenafil (N=70) and placebo, respectively. These results demonstrated that the effect of sildenafil citrate on the primary endpoint was statistically non-inferior to placebo. It should be noted that the results presented were from a controlled clinical research trial in which selected patients were carefully screened and monitored.

After patients have taken sildenafil citrate, it is unknown when nitrates, if necessary, can be safely administered. Based on the pharmacokinetic profile of a single 100 mg oral dose given to healthy normal volunteers, the plasma levels of sildenafil at 24 hours post-dose are approximately 2 ng/mL (compared to peak plasma levels of approximately 440 ng/mL). In the following patients: age >65, hepatic impairment (e.g. cirrhosis), severe renal impairment (e.g. creatine clearance <30 mL/min), and concomitant use of potent cytochrome P-450 3A4 inhibitors (erythromycin), plasma levels of sildenafil at 24 hours post-dose have been found to be 3 to 8 times higher than those seen in healthy volunteers. Although plasma levels of sildenafil at 24 hours post-dose are much lower than at peak concentration, it is unknown whether nitrates can be safely coadministered at this time point (see **CONTRAINDICATIONS**).

Single oral doses of sildenafil above 15 mg were generally associated with a potentiation of the antiaggregatory effects of sodium nitroprusside (SNP) on ADP aggregation of *ex vivo* platelets. Sildenafil had no effect on other *ex vivo* tests (ADP-induced platelet aggregation of whole blood and ADP-induced aggregation of platelet-rich plasma in the absence of SNP). Sildenafil therefore has no direct effect on platelet function *ex vivo*, but potentiates the action of a nitric oxide (NO) donor, SNP. This confirms the need for an NO drive before sildenafil will produce its pharmacological effects. These modest effects on platelet activity, *ex vivo*, did not result in a

clinically significant effect on bleeding time in healthy volunteers.

Effect of sildenafil citrate on Sperm Motility:

Sildenafil had no effect on sperm motility, morphology, count, density, vitality, ejaculate volume or viscosity. The concentrations of sildenafil in the ejaculate, 1.5 hours and 4 hours post-dose, were 18% and 17%, respectively, of the concentrations in plasma at the same time points. The concentrations of the metabolite, at the same time points were 5% and 15%, respectively.

There was no effect on sperm motility or morphology after single 100 mg oral doses of sildenafil citrate in healthy volunteers.

Effects of sildenafil citrate on Vision:

At single oral doses of 100 mg and 200 mg, transient dose-related impairment of colour discrimination (blue/green) was detected using the Farnsworth-Munsell 100-hue test, with peak effects near the time of peak plasma levels. This finding is consistent with the inhibition of PDE6, which is involved in phototransduction in the retina. An evaluation of visual function at doses up to twice the maximum recommended dose revealed no effects of sildenafil citrate on visual acuity, contrast sensitivity, ERGs, intraocular pressure, or pupillometry. In flexible titration studies of 4 to 26 weeks, 3% of patients on sildenafil reported visual disturbances: mild and transient impairment of colour discrimination (predominantly blue/green), and also increased perception to light or blurred vision (see **WARNINGS AND PRECAUTIONS**).

In healthy volunteers aged 40-65 years, single doses of sildenafil up to 200 mg had no clinically relevant effect on visual acuity, contrast sensitivity, pupil diameter and constriction velocity, visual fields, recovery time following dazzle, electroretinogram or intraocular pressure. Modest, transient changes in colour discrimination were observed (Farnsworth-Munsell 100 Hue test) after 100 mg and 200 mg doses, but not at 50 mg. At 100 mg, this effect was apparent only at one hour after dose and at 200 mg, up to two hours after dose.

In a placebo-controlled, crossover study of patients with documented early age-related macular degeneration (n=9), sildenafil (single dose, 100mg) was well-tolerated and demonstrated no clinically significant changes in the visual tests conducted (visual acuity, Amsler grid, color discrimination, simulated traffic light, Humphrey perimeter and photostress).

Pharmacokinetic Studies

When administered orally to healthy male volunteers in the fasted state, sildenafil was rapidly absorbed, with maximum observed plasma concentrations (C_{max}) occurring 0.5-2 hours after dosing in most subjects. C_{max} and areas under the plasma concentration time curve to infinite time (AUC) increased in a proportional manner with dose over the clinical dose range 25-100 mg.

Sildenafil has an apparent volume of distribution at steady state (V_{ss}) of 105 litres and a mean plasma clearance (CL) of 41 L/h. Both V_{ss} and CL were shown to be significantly correlated to body weight. The absolute oral bioavailability was 41%. Sildenafil has a terminal half-life of approximately 4 hours (range 2-8 hours). Approximately 96% of sildenafil is bound to plasma proteins.

Metabolism and Elimination

The major circulating metabolite of sildenafil, results from N-demethylation of sildenafil at the N-methyl piperazine moiety. It has a similar selectivity for PDE isozymes as sildenafil, but exhibits around 50% of the potency of sildenafil. The metabolism of sildenafil occurs in human hepatic microsomes and is mediated by two cytochrome P-450 isoforms [CYP2C9 (minor route) and CYP3A4 (major route)].

The concomitant use of potent cytochrome P-450 3A4 inhibitors (e.g. erythromycin, saquinavir, ritonavir, ketoconazole, itraconazole) as well as the non-specific CYP inhibitor, cimetidine, is associated with increased plasma levels of sildenafil (see **WARNINGS AND PRECAUTIONS, DOSAGE AND ADMINISTRATION**).

TOXICOLOGY

Acute Toxicity Mice and Rats:

Species	Route	Dose mg/kg/day	#Animals / dose level	Duration	Findings
Single dose oral toxicity in mice and rats (90155/56)					
Sprague-Dawley rat CD1 mice	Oral (gavage)	rat: 300 500 1000 mice: 500 1000	5/sex	1 day	At 1000 mg/kg one male mouse died within 24 hours after drug administration. In rats, mortality occurred in three females at 1000 mg/kg and in one female at 500 mg/kg. The dose of 1000 mg/kg induced clinical signs in both species, generally within 24 hours following the administration, which persisted less than 24-48 hours. Some of these signs were similar in mice and rats and consisted of partially-closed eyes, hunched posture, tremours, depression, coldness to the touch (with pallor of ears and paws in rats) and prostration. Female rats were more affected than male rats. Dyspnea was limited to one mouse, and chromodacryorrhea to four female rats. Clinical signs at 500 mg/kg included partially-closed eyes in one mouse and subdued behaviour in the female rat which died. No clinical signs were observed in rats at 300 mg/kg. In both species, the doses administered induced no changes in body weight gain and there were no treatment- related macroscopical changes at gross necropsy. These results indicate that the no observed adverse effect level (NOAEL) was 500 mg/kg in mice and 300 mg/kg in rats.
Single dose intravenous toxicity in mice and rats (91045/046)					
Sprague-Dawley rat CD1 mice	I.V.	rat: 10 mice: 20	5/sex	1 day	All animals survived the treatment and gained weight over the 14-day study period. There were no clinical signs during the study and no abnormalities at necropsy. Under the conditions of this study, the no observed effect level (NOAEL) after intravenous administration was 20 mg/kg in mice and 10 mg/kg in rats.

Long-Term Toxicity - Mice:

Species	Route	Dose mg/kg/day	#Animals / dose level	Duration	Findings
3-Month oral (gavage) prechronic toxicity study in mice (94049)					
CD1	Oral (gavage)	10 50 100 200	10/sex	3 months	<p>The exposure to sildenafil and its metabolite UK-103,320 was similar in males and females and approximately dose-related. Treatment-related mortality occurred in 3/20 animals in each group given 50, 100 or 200 mg/kg. A marked gastrointestinal dilation was the cause of the death and was associated with a number of clinical signs, in particular dyspnea and/or swollen abdomen. This dilation resulted in gastrointestinal inflammation, fatty changes and focal/multifocal necrosis in the liver, atrophy of adipose tissues and hemoconcentration. There was also a mild gastrointestinal dilation in a few survivors of these groups. In males treated with 50, 100 or 200 mg/kg, there was an apparent decrease in body weight gain. However, in the absence of dose relationship and consistent statistical significance, the association with treatment is questionable. Plasma cholesterol was slightly increased in females treated with 50, 100 or 200 mg/kg and plasma triglycerides were slightly decreased in males treated with 100 or 200 mg/kg. However we consider these changes to be of minor toxicological importance.</p> <p>The NOAEL in this study was 10 mg/kg, given the mortality and gastrointestinal dilation at higher doses.</p>
3-Month oral (gavage) exploratory toxicity study in mice (94101)					
CD1	Oral (gavage)	20 40 100	10/sex	3 months	<p>The exposure to sildenafil and its metabolite UK-103,320 was similar in males and females and increased superproportionally with dose level. Treatment-related mortality occurred in 1/20 animals in each group given 40 or 100 mg/kg. A marked gastrointestinal dilation was the cause of the death and was associated with a number of clinical signs, in particular dyspnea and/or swollen abdomen. There was also a transient abdominal swelling in a few survivors of these groups.</p> <p>The NOAEL in this study was 20 mg/kg, given the mortality and gastrointestinal dilation at higher doses.</p>

Long-Term Toxicity - Rats:

Species	Route	Dose mg/kg/day	#Animals / dose level	Duration	Findings
10-Day oral range-finding toxicity in rats (90080)					
Sprague Dawley	Oral (gavage)	50 150 500	5/sex	10 days	<p>Measurement of plasma concentrations of sildenafil and UK-103,320 showed that females were exposed predominantly to the drug while males were exposed mainly to the metabolite, UK-103,320, and a lower level of unchanged compound. Concentrations of UK-95,340 were generally below the limit of determination (30 ng/mL). Exposure increased with dose but not in linear manner. At 500 mg/kg, 1/5 females died after the second dose with no apparent cause of death. Of the animals used for plasma drug determination, 1/10 rats at 150 mg/kg and 2/10 rats at 500 mg/kg died after the first or second dose. As these animals died after taking blood samples, they were not considered in the analysis of mortality. Food consumption was decreased between day 1 and 4 in mid- and high-dose males and in all treated female groups. A dose-related decrease of plasma triglycerides occurred in males, and an increase of plasma cholesterol was seen in high-dose females. Blood urea increased in mid- and high-dose males and in the 3 treated female groups. Relative heart weight was slightly increased in high-dose males. Kidney and liver weights were increased in mid- and high-dose females, and in high-dose males. The increase of liver weight was associated with centrilobular hypertrophy. Changes in red blood cell parameters were seen in females. They indicate a decrease of circulating red blood cells at the 3 dose levels, with some evidence of regenerative response at the high dose. An increase of white blood cell counts was recorded at the mid dose in females and at the high dose in both sexes. Changes at the dose of 50 mg/kg were considered minor.</p> <p>The NOAEL in this study was 150 mg/kg, based on the mortality at 500 mg/kg.</p>

Species	Route	Dose mg/kg/day	#Animals / dose level	Duration	Findings
1-Month oral toxicity in rats (90143)					
Sprague Dawley	Oral (gavage)	10 45 200	10/sex	1 month	<p>Plasma concentrations of sildenafil were higher in females than in males, while concentrations of the metabolite, UK-103,320, were higher in males than in females. As a result, females were exposed predominantly to the unchanged drug and males to an almost equal balance of drug and metabolite. These data indicate that N-demethylation of sildenafil to UK-103,320 is an important route of sildenafil biotransformation in male rats. Concentrations of UK-95,340 were generally below the limit of determination (30 ng/mL).</p> <p>One of the high-dose females used for plasma drug level determination died after the first dose, before blood samples had been taken. Clinical signs were limited to a few high-dose animals and consisted of chromodacryorrhea and palpebral closure. Slight increases in water and food intake were seen generally in mid- and high-dose animals. A mild dose-related decrease in circulating red blood cells with evidence of a regenerative response was found in mid- and high-dose females and, to a smaller extent, in high-dose males. A moderate neutrophilia was seen in high-dose males, while a moderate lymphocytosis occurred in mid- and high-dose females. Plasma chemistry changes at the high dose consisted of increases in urea, decreases in triglycerides (males) and increases in cholesterol (females), but remained within our normal range of values. Doses of 45 and/or 200 mg/kg were associated with an increase in liver weight and centrilobular hypertrophy in both sexes. Hypertrophy of the zona glomerulosa of the adrenal glands was seen in the high-dose males and in the mid- and high-dose females. Thyroid follicular hypertrophy occurred at the high dose in both sexes. In addition, mesenteric arteritis was found in two mid-dose and one high-dose males, but was not considered to be related to the treatment. The NOAEL was 45 mg/kg in this study.</p>
28-Day oral exploratory toxicity study in rats (94085)					
Sprague-Dawley	Oral (gavage)	0 60 120	10 males/group	28 days	<p>A 2-year rat carcinogenicity study with sildenafil citrate at a contract laboratory (Study No. 911/002), at doses of 1.5, 5 and 60 mg/kg, was terminated after unexpectedly high mortality and severe toxic effects in high-dose males during weeks 3 and 4. An exploratory study was performed to confirm that the batch of sildenafil used at the contract laboratory did not induce severe toxicity.</p> <p>The only treatment-related effects were a mild dose-related increase in liver and kidney weights and possibly a slight decrease in body weight gain. Importantly, the absence of death in this study confirms the results of previous studies up to 200 mg/kg, and contrasts with the results of the study at the contract laboratory. Subsequently, it was shown that the mortality in the carcinogenicity study (Study No. 911/002) was due to dosing with a cytotoxic compound from another company and not sildenafil. Consequently, the contracted carcinogenicity study was invalid.</p>

Species	Route	Dose mg/kg/day	#Animals / dose level	Duration	Findings
Investigation of the relationship between liver enzyme induction and thyroxine clearance in rats (96010)					
Sprague-Dawley	Oral (gavage)	200	10 females	1 month	<p>Following the appearance of thyroid follicular hypertrophy in rats, an investigative study was conducted to examine the relationship between liver enzyme induction and thyroid hypertrophy in rats. Two groups of 10 female rats were treated orally with sildenafil citrate at 200 mg/kg for 29 days, and two control groups received the vehicle alone. One treated group and one control group were used for assessment of exogenous thyroxine clearance. The other treated group and the other control group were used for measurement of plasma TSH and thyroid hormones, for histopathological examination of the liver and thyroid, and for determination of UDP-glucuronyl transferase (UDPGT) activity in the liver.</p> <p>The treatment caused the deaths of 2/20 rats on days 2 or 3. In the treated group, there was an increase in the weight of liver and thyroid, associated with minimal centrilobular hypertrophy of the liver and thyroid follicular cell hypertrophy. There was also an increase in hepatic UDPGT activity, an increase in TSH, and a decrease in T3 and T4 hormones. In addition, the clearance of exogenous thyroxine was increased in treated animals.</p> <p>These results are consistent with the view that the thyroid hypertrophy associated with treatment of rats with sildenafil was due to induction of hepatic UDPGT which increased the clearance of thyroid hormone and consequently caused a compensatory increase in plasma TSH which stimulated the thyroid gland.</p>
6-Month oral toxicity study in rats (91098)					
Sprague-Dawley	Oral (gavage)	3 12 60	20/sex	6 months	<p>Drug and metabolite plasma level determinations showed that females were exposed predominantly to sildenafil while males were exposed almost exclusively to the metabolite. No treatment-related deaths were recorded. Chromodacryorrhea was seen in the 3 treated groups. Body weight gain and food consumption were increased at the low dose and, to a lesser extent, at the mid dose. A trend towards a reduced body weight gain was seen at the high dose; however, the relationship to compound administration cannot be ascertained. Decreases of plasma bilirubin and triglycerides, and increases in plasma urea, total proteins and cholesterol were seen at the high dose. These changes suggest compound-induced metabolic changes in the liver. Increase in liver weight associated with mild centrilobular hypertrophy indicate an adaptive response. Thyroid hypertrophy occurred at the high dose in both sexes and at a lower incidence in mid-dose males. This change was considered to be a secondary phenomenon related to increased hepatic clearance of thyroid hormone. Although thyroid hormones and hepatic clearance were not measured in this study, changes in these parameters were demonstrated in an exploratory study (Study No. 96010). Hypertrophy of the zona glomerulosa of the adrenal gland occurred with a dose-related incidence at the mid and high doses and was associated with an increase in the weight of the organ at 60 mg/kg.</p> <p>The NOAEL in this study was 60 mg/kg.</p>
13-Day intravenous range-finding in rats (90139)					
Sprague-Dawley	I.V.	2.5 5 10	5/sex	13 days	<p>No deaths occurred during the treatment period. The only clinical sign noted was a transient redness of the ears in a few treated animals, notably in the high-dose male group. The NOAEL in this study was 10 mg/kg.</p>
1-Month intravenous toxicity study in rats (91044)					
Sprague-Dawley	I.V.	0.5 2 4	10/sex	1 month	<p>No compound-related changes were seen at the doses of 0.5 and 2 mg/kg. At the dose of 4 mg/kg, the incidence and severity of mild myocardial inflammation was slightly increased compared to the control group; the relationship to treatment cannot be ascertained. The NOAEL in this study was 2 mg/kg.</p>

Long-Term Toxicity - Dogs:

Species	Route	Dose mg/kg/day	#Animals / dose level	Duration	Findings
10-Day oral range-finding toxicity in dogs (90081)					
Beagle	Oral (gavage)	10 30 100	1 male 2 females	10 days	Plasma concentrations of sildenafil and UK-103,320 were similar in males and females and increased with dose, although subproportionally at the high dose. The proportion of UK-103,320 relative to sildenafil varied minimally (18-24%) over the dose range examined and indicates no detectable saturation of this metabolic pathway. Concentrations of UK-95,340 were generally below the limit of determination (30 ng/mL). Emesis and salivation occurred at the dose of 100 mg/kg, and lacrimation, conjunctival redness and a transient decrease in amplitude of the pupillary reflex were seen at all dose levels. There was no evidence of a convincing change in blood pressure, given the spontaneous variation in this parameter. Heart rate was increased at 30 and 100 mg/kg, and probably represents a reflex response to the vasodilating properties of the compound. Decreases in PQ and QT intervals of the ECG at these doses were secondary to the heart rate changes. Two high-dose animals showed a moderate increase of plasma cholesterol which was not considered to be toxicologically important. An arteritis of an extramural branch of a coronary artery was found in one high-dose female. This is considered to be a spontaneous finding considering the morphological features and the background incidence in Beagle dogs in our laboratories. The NOAEL in this study was therefore 100 mg/kg.
1-Month oral toxicity study in dogs (90125)					
Beagle	Oral (gavage)	5 20 80	3/sex	1 month	The dogs were exposed to concentrations of sildenafil and UK-103,320, which increased with dose, although subproportionally at the high dose. The proportion of UK-103,320 relative to sildenafil varied minimally (15-19%) over the dose range examined and indicates no detectable saturation of this metabolic pathway. Concentrations of UK-95,340 were generally below the limit of determination (30 ng/mL). At the mid and high doses, the compound induced a low incidence of emesis and transient salivation. A moderate incidence of soft and liquid feces was noted at all doses. There was no evidence of consistent changes in blood pressure, although there were increases in heart rate at 20 and 80 mg/kg. Changes in the ECG (increased P-wave amplitude and decreases in PQ and QT intervals) were expected from the increases in heart rate. There was a moderate increase in plasma cholesterol at the high dose. A mild coronary arteritis was seen in one high-dose animal, but considering the morphological features of this finding, and the high background incidence in Beagle dogs in our laboratories, this was not thought to be treatment-related. The NOAEL was 80 mg/kg in this study.
6-Month oral toxicity in dogs (91099)					
Beagle	Oral (gavage)	3 15 50	4/sex	6 months	Analyses of plasma sildenafil and UK-103,320 showed dose-related concentrations in the dog. The proportion of UK-103,320 relative to sildenafil varied minimally (15-23%) as the dose increased, indicating no saturation of this process. Salivation, emesis and resistance to compound administration were seen when the animals were treated with an initial high dose of 80 mg/kg, and reflected gastric intolerance to the compound at this dose level. These signs were rare after reducing the high dose to 50 mg/kg. A moderate increase in heart rate, associated with decreases in PQ and QT intervals, occurred at the high dose and is considered to be a reflex response to the vasodilatory properties of the drug. Increases in plasma cholesterol and in liver weight were seen in animals treated with 15 and 50 mg/kg. A high-dose male showed a number of clinical signs and changes in hematological parameters and plasma chemistry associated with a disseminated arteritis. These changes correspond to Idiopathic Juvenile Arteritis Syndrome (Beagle Pain Syndrome) which occurs sporadically in Beagle dogs. Another high-dose male showed arteritis in the thymus which indicated a less severe expression of the same disease. It is probable that the high dose precipitated the expression of this latent spontaneous disorder. The NOAEL in this study was 15 mg/kg, given the appearance of Idiopathic Juvenile Arteritis Syndrome at higher doses.

Species	Route	Dose mg/kg/day	#Animals / dose level	Duration	Findings
12-Month oral toxicity study in dogs (95039)					
Beagle	Oral (gavage)	3 10 50	4/sex	12 months	<p>The dogs were exposed to approximately dose-related concentrations of sildenafil and its N-demethylated metabolite, UK-103,320. The proportion of UK-103,320 relative to sildenafil varied minimally as the dose increased. Features typical of a syndrome of Idiopathic Juvenile Arteritis occurred in all high-dose males. In 3/4 high-dose males, there was arteritis which affected several organs. In one of these dogs, arteritis was associated with a number of clinical signs, body weight loss and hematological changes. In the other two animals, there were no clinical or hematological correlates to arteritis. In addition, the fourth high-dose male presented clinical signs and clinical pathology changes typical of the syndrome though no vascular lesion was found at histopathology. Focal coronary arteritis occurred in one low-dose and one high-dose female; neither finding was considered treatment-related. The treatment produced an increase in the amount of lipogenic pigments in renal tubular epithelium in 1/8 animals at the mid dose and 7/8 animals at the high dose, a dose-related decrease in plasma creatine kinase, mainly in males, and a decrease in plasma myosin in high-dose animals. However, these changes were considered of no toxicological importance. A dose-related increase in heart rate occurred at the high and mid doses, and was considered to be due to compensatory mechanisms occurring in response to the vasodilatory properties of the compound.</p> <p>The NOAEL in this study was 10 mg/kg, given the appearance of Idiopathic Juvenile Arteritis Syndrome at higher doses.</p>
14-Day intravenous range-finding toxicity in dogs (90142)					
Beagle	I.V.	2.5 5 10	2 males 1 female	14 days	<p>The doses of 5 and 10 mg/kg were associated with liquid feces and an inhibition of the pupillary reflex. An increase in heart rate was observed at the high dose and, to a lesser extent, at the mid dose. This change was probably related to the vasodilator effect of the compound. Evidence of vasodilatation was provided by the peripheral redness seen in two high-dose animals. An increase in plasma cholesterol occurred in 2/3 high-dose animals but was not considered to be toxicologically important. At the dose of 2.5 mg/kg, there were no treatment-related changes. The NOAEL was 10 mg/kg in this study.</p>
1-Month intravenous toxicity in dogs (91041)					
Beagle	I.V.	0 0.5 2 4	3/sex	1 month	<p>The treatment induced no adverse effects. The NOAEL is therefore 4 mg/kg in this study.</p>

Bioequivalence - Dogs:

Species	Route	Dose mg/kg/day	#Animals / dose level	Duration	Findings
Bioequivalence between base and citrate in dogs (91058)					
Beagle	Oral	300	1 male 1 female	N/A	<p>The aim of the current study was to assess, in the dog, the oral bioequivalence of a suspension of the base, and of capsules of the citrate. The base was suspended in a 5% aqueous solution of methylcellulose 4000 cps containing 0.1% Tween 80 and acidified with hydrochloric acid 0.1M (final concentration). The citrate salt was administered in gelatin capsules.</p> <p>On day 1, a first group of one male and one female beagle dogs was treated with the base and the second group of one male and one female was treated with the citrate. On day 8, the first group received the citrate, and the second group the base. The animals were regularly examined for clinical signs and weighed before each administration. Blood was sampled 0.25,0.5,1,1.5,2,3,4,6,8,11 and 24 hours after each administration. Plasma levels of UK-92,480 and two metabolites, UK-95-340 and UK-103,320, were measured.</p> <p>One male dog vomited after each administration and its drug and metabolite plasma concentrations were therefore considered not to be relevant. In other dogs, maximal plasma concentrations and AUCs of UK-92,480 and of UK-103,320, observed after administration of the citrate in capsules were similar to or higher than those seen after administration of the base in a suspension. All the plasma concentration of UK-95,340 were below the limit of detection of the assay. These data indicate that bioavailability of the citrate in the dog is identical to or better than that of the base.</p>

Carcinogenesis and Mutagenesis

Sildenafil was not carcinogenic when administered to rats for 24 months at a dose resulting in systemic drug exposure (AUC) of 110- and 146-times, respectively, for male (unbound sildenafil and its major metabolite) and female (unbound sildenafil) rats. The exposures observed in humans given the Recommended Human Dose (RHD) of 20 mg t.i.d. sildenafil was not carcinogenic when administered to mice for 18-21 months at dosages up to the Maximum Tolerated Dose (MTD) of 10 mg/kg/day, approximately 1.1 times the RHD on a mg/m² basis.

Sildenafil has been studied in a comprehensive battery of tests designed to detect genotoxic activity. Sildenafil did not display mutagenic activity in bacterial or mammalian cells *in vitro*, or clastogenic activity *in vitro* or *in vivo*.

As the clinical dose is administered three times daily, the clinical free AUC used to calculate exposure multiples was 19 ng-h/mx³, and compared with the AUC 0-24 hours in the preclinical species.

Species	Route	Dose mg/kg/day	#Animals/dose level	Duration	Findings
Pharmacokinetic study in rats (94067)					
Sprague Dawley	Oral (gavage)	60	5/sex	14 days	This study was conducted to provide an estimate of the pharmacokinetic exposure of rats over 24 hours. Plasma concentrations of sildenafil were higher in females than in males, while concentrations of the metabolite, UK-103,320, were higher in males than in females.
Oral toxicity and carcinogenicity study in mice (95007)					
CD1	Oral (gavage)	3 10 30	55/sex	<u>3 &</u> 10 mg: males 649 days females 558 days <u>30 mg:</u> males 453 days females 404 days	<p>The exposure to the parent compound and the demethylated metabolite, UK-103,320 was dose-related. The compound produced an increase in mortality rate with consequent decreases in survival times and percent of survival.</p> <p>The effect was marked at the mid dose in females and at the high dose in both sexes. In addition, the percent of survival was also slightly decreased in mid-dose males, at the end of the study. Because of the lower survival in mid- and high-dose animals interim sacrifices were decided. When the survival in the high-dose group reached about 20%, the survivors were sacrificed, on day 405 (females) or 454 (males). Control, low- and mid-dose groups were sacrificed on day 559 (females) or 650 (males), when the survival at the mid dose was about 20%. In a number of animals, especially high-dose males (40%), unscheduled death was preceded by abdominal swelling and/or dyspnea. Gastrointestinal dilation and gavage accident were identified as causes of unscheduled death related to treatment. Additionally, the number of deaths without explanatory macroscopic or histopathological changes was higher in mid- and high-dose groups than in the control groups. In high-dose males and females, there was also a trend to body weight decrease compared to controls (10 and 18%, respectively). In addition, there was an abrupt body weight loss in most animals dying prematurely which was more marked in mid- and high-dose females. The treatment produced no increase in the incidence of neoplastic lesions. Furthermore, in the animals sacrificed at the various interim and final sacrifices, there were no differences in the incidence of non-neoplastic lesions between control and treated groups.</p> <p>In conclusion, the doses of 10 and 30 mg/kg produced signs of toxicity consisting mainly of a dose-related increase in mortality. At the dose of 3 mg/kg, although there was no compound effect on group mortality, 2 animals died from gastrointestinal dilation. There were no carcinogenic effects at any dose.</p>

Species	Route	Dose mg/kg/day	#Animals/dose level	Duration	Findings
24-Month oral toxicity and carcinogenicity study in rats study (94092)					
Sprague-Dawley	Oral (gavage)	1.5 5 60	60/sex	24 month	<p>The rats were exposed to plasma concentrations of sildenafil and UK-103,320 that increased with dose levels. Male rats were exposed predominantly to UK-103,320, whereas unchanged drug was the major circulating form in females. Overall, the total exposure to drug and metabolite was higher in females than in males.</p> <p>The treatment produced no mortality. Survival at the end of the study ranged between 18 and 42% in males and between 15 and 25% in females.</p> <p>The body weight was decreased in high-dose animals, compared to controls. A transient decrease in body weight occurred also in mid-dose females. There was a dose-related decrease in plasma bilirubin which, in our view, is related to the enzyme-inducing properties of the compound. In high-dose males there was an increased incidence of proliferative changes in the thyroid which was mainly related to an increase in follicular cell hyperplasia. We consider that these changes are the consequence of an increased turnover of thyroid hormones due to hepatic enzyme induction and bear no relevance to man.</p> <p>To conclude, the dose of 60 mg/kg was associated with a toxicologically significant decrease in body weight and with an increase in follicular proliferative changes in the thyroid in males. At 5 mg/kg there was only an inconsistent decrease in the body weight of females. There were no compound effects at 1.5 mg/kg. There were no indications of a carcinogenic potential of sildenafil.</p>

Mutagenicity studies (90817-01/02)			
Study Type	Strain	Dose	Results
<i>in vitro</i> bacterial mutagenicity	<i>S. typhimurium</i> TA 1535, 1537, 98, 100	0.002 - 1 mg/plate	negative
<i>in vitro</i> mammalian cell mutagenicity	Chinese Hamster Ovary / HGPR1	65-240 µg/mL	negative
<i>in vitro</i> clastogenicity	Human lymphocytes	10, 20, 25 µg/mL -S9 100, 125, 250 µg/mL + S9	negative
<i>in vivo</i> clastogenicity	Mouse bone marrow	0, 500, 1000, 2000 mg/kg	negative

Reproduction and Teratology

No evidence of teratogenicity, embryotoxicity or fetotoxicity was observed in rats and rabbits which received up to 200 mg/kg/day during organogenesis. These doses represent, respectively, about 32 and 68 times the RHD on a mg/m² basis in a 50 kg subject. In the rat pre- and postnatal development study, the no observed adverse effect dose was 30 mg/kg/day given for 36 days. In the non-pregnant rat the AUC at this dose was about 24 times unbound human AUC.

Species	Route	Dose mg/kg/day	#Animals/dose level	Duration	Findings
Maternal toxicity study in rats by the oral route (92020)					
Sprague-Dawley	Oral (gavage)	10 50 200	7 females	Gestation days 6-17	Hematological, biochemical (plasma) and pathological changes were recorded only at 200 mg/kg. Hematological changes consisted of a moderate decrease in hemoglobin, red blood cell count and packed cell volume accompanied by an increase in the mean red blood cell distribution width. The only variation observed in plasma chemistry was a decrease in mean plasma triglycerides. Finally, a mild hepatic weight increase with hepatic centrilobular hypertrophy was noted after pathological examination. With regard to the fetuses, there was a decrease in the mean male body weight at 200 mg/kg. In male fetuses at 10 and 50 mg/kg and in female fetuses at all dose levels, the mean body weights were similar to those of the control group. The NOAEL was 50 mg/kg in dams and fetuses given the changes in plasma chemistry and fetal weight of males at 200 mg/kg.
Study of fertility and early embryonic development to implantation in rats by the oral route (94081)					
Sprague-Dawley	Oral (gavage)	3 12 60	20/sex	<u>Males:</u> from 9 weeks before mating to gestation day 20 <u>Females:</u> from 2 weeks before mating to gestation day 6	The treatment produced no adverse effects on the fertility of either sex. In addition, there was no evidence of maternal, embryo- or fetotoxicity. The only finding was a moderate reduction in plasma triglycerides in females treated with 60 mg/kg. Therefore the NOAEL in this study was 60 mg/kg.

Species	Route	Dose mg/kg/day	#Animals/dose level	Duration	Findings
Study for effects on pre- and post-natal development, including maternal function, in rats by the oral route (95068/95095)					
Sprague-Dawley	Oral (gavage)	10 30 60	20 females	from gestation day 6 until 20 days after birth	<p>The only noteworthy finding was a toxicologically significant decrease in the ratio of viable pups at birth, with consequently a decreased litter size of viable pups, at 60 mg/kg. At this high-dose level, there was a toxicologically significant decrease in the 4-day survival index, in the F₁ pups body weight on day 1 p.p. and some delay in a developmental landmark, the appearance of upper incisors. There were no findings in the reproductive function of the F₁ generation, and in the F₂ generation.</p> <p>The NOAEL was 30 mg/kg for F₀ females and F₁ pups, given the minimal maternal toxicity and the effect on pup development during the first 2 weeks of life. The NOAEL for the F₂ generation is 60 mg/kg.</p>

Species	Route	Dose mg/kg/day	#Animals/dose level	Duration	Findings
Study for effects on embryo-foetal development in rats by the oral route (95058/95059)					
Sprague-Dawley	Oral (gavage)	10 50 200	20 females	Gestation days 6-17	<p>There were detectable levels of sildenafil and UK-103,320 in maternal plasma, amniotic fluid and fetal homogenates at all dose levels. Treatment at 200 mg/kg produced salivation and a reduction in mean body weight gain between days 6 and 9 p.c., accompanied by a decrease in food intake on day 9 p.c. On day 18 p.c., the mean food consumption increased. Hematological changes consisted of a slight decrease in hemoglobin, red blood cell count and hematocrit accompanied by an increase in the mean red blood cell distribution width at 200 mg/kg. A dose-related increase in the reticulocyte count was present, reaching statistical significance at the high-dose only. The only variation in plasma chemistry was a dose-related decrease in mean plasma triglycerides, at most moderate and statistically significant at the high-dose only. The body weight of male fetuses was reduced at 200 mg/kg. There were no treatment-related external, skeletal or visceral anomalies.</p> <p>Treatment with 200 mg/kg produced a slight maternal toxicity without embryotoxicity but a slight toxicity in male fetuses only. There was no maternal, fetal or embryotoxicity after treatment with 10 or 50 mg/kg. There were no teratological effects at any dose.</p> <p>The NOAEL in this study was 50 mg/kg in dams and fetuses, given the slight toxicity at 200 mg/kg.</p>

Rabbits:

Species	Route	Dose mg/kg/day	#Animals/dose level	Duration	Findings
Maternal toxicity study in rabbits by the oral route (95003/95004)					
New Zealand White	Oral (gavage)	50 100 200	7 females	Gestation days 6-18	<p>Pregnant females and fetuses were exposed to the drug. The only noteworthy findings in dams were an increase in plasma glucose and a decrease in plasma cholesterol at the high dose. This is indicative of a minimal toxicity in dams. There were no adverse effects on embryo or fetal development.</p> <p>The NOAEL was 100 mg/kg in dams given the changes in plasma chemistry values at 200 mg/kg. The NOEL was 200 mg/kg in the developing embryos and fetuses.</p>

Species	Route	Dose mg/kg/day	#Animals/dose level	Duration	Findings
Study for effects on embryo-foetal development in rabbits by the oral route (95043/44)					
New Zealand White	Oral (gavage)	10 50 200	20 females	Gestation days 6-18	<p>Sildenafil and UK-103,320 were found in the plasma of pregnant females. The presence of sildenafil was also detected in amniotic fluid. At the high-dose, there were reductions in body weight and body weight gain late in gestation, compared to the control group, which are indicative of minimal maternal toxicity. A reduction in food intake in high-dose females during the same period may have contributed to the body weight changes. The plasma chemistry changes, encountered in the preliminary study, were not found in this study. The treatment had no adverse effects on the developing conceptus.</p> <p>The NOAEL in this study was 50 mg/kg for dams, given the effect on body weight at 100 mg/kg. The NOEL was 100 mg/kg in the developing embryos and fetuses.</p>

Special Studies:

Species	Route	Dose mg/kg/day	#Animals/dose level	Duration	Findings
Antigenicity study in guinea pigs (95-29-81)					
Hartley Guinea Pigs	Oral	4 mg/mL 20 mg/mL	5/group	N/A	<p>In the active systemic anaphylaxis test, male guinea pigs that received daily doses of 4 or 20 mg/kg sildenafil orally 5 days a week for 3 weeks showed no signs of systemic anaphylaxis reactions after intravenous injection of sildenafil 19 days later as challenge antigen. Similarly, when male guinea pigs sensitized subcutaneously with 2 or 10 mg sildenafil/guinea pig (given on 4 occasions at 1 week intervals) were challenged 16 days later with intravenous injection of sildenafil, they showed no signs of systemic anaphylaxis.</p> <p>In the passive cutaneous anaphylaxis test, guinea pigs were challenged with sildenafil (30 mg/guinea pig). No positive PCA reactions were observed against anti-sera obtained from guinea pigs immunized orally or subcutaneously with sildenafil.</p>
	sub-cutaneous (with Freund's complete adjuvant)	2 mg/mL 10 mg/mL			
Intra-arterial irritation in rabbits (91073)					
New Zealand White	Intra-arterial	1 mg/animal	4 females	1 day	Sildenafil (1 mg/animal) was administered into the central ear artery of rabbits in a volume of 0.5 mL to examine the potential irritant reactions. The single injection produced no arterial irritation over a 21-day observation period.

REFERENCES

1. Agelink MW, Schmitz T, Rembrink K, et al. Cardiovascular effects of sildenafil citrate (Viagra®): A naturalistic cross-over study. *Eur J Med Res* 2001;6(11):459-64.
2. Arruda-Olson AM, Mahoney DW, Nehra A, et al. Cardiovascular effects of sildenafil during exercise in men with known or probable coronary artery disease. *JAMA* 2002;287(6):719-25.
3. Ballard SA, Burslem FMF, Gingell CJC, et al. *In vitro* profile of UK-92,480, an inhibitor of cyclic GMP-specific phosphodiesterase 5 for the treatment of male erectile dysfunction. *J Urol* 1996;155(No 5, Suppl):1462.
4. Ballard SA, Gingell, CJ, Tang K, et al. Effects of sildenafil on the relaxation of human *corpus cavernosum* tissue *in vitro* and on the activities of cyclic nucleotide phosphodiesterase isozymes. *J Urol* 1998;159(6):2164-71.
5. Bernard F, Carrier S, Lee JC, Talwar V, Defoy I. Men with Mild Erectile Dysfunction Benefit from Sildenafil Treatment. *J Sex Med* 2010; 7: 3725-3735.
6. Boolell M, Allen MJ, Ballard SA, et al. Sildenafil: an orally active type 5 cyclic GMP-specific phosphodiesterase inhibitor for the treatment of penile erectile dysfunction. *Int J Impot Res* 1996;8(2):47-52.
7. Boshier A, Wilton LV, Shakir SA. Evaluation of the safety of sildenafil for male erectile dysfunction: experience gained in general practice use in England in 1999. *BJU Int.* 2004; 93(6): 796-801.
8. Bush, HS. Safe use of sildenafil in patients with coronary artery disease. *Cleve Clin J Med*, 2001; 68(4):349-52.
9. Buvat J, Gingell CJ, Jardin A, et al. Sildenafil (Viagra™), an oral treatment for erectile dysfunction: a 1-year, open-label, extension study. *J Urol* 1997;157(4 Suppl):204.
10. Carson C, Burnett A, Levine L, et al. The efficacy of sildenafil citrate (Viagra®) clinical populations: an update. *Urology* 2002; 60 (2B Suppl): 12-27.
11. Carter AJ, Ballard SA and Naylor AM. Effect of selective phosphodiesterase type 5 inhibitor sildenafil on erectile function in the anesthetized dog. *J Urol* 1998;160(1):242-6.
12. Chang G, Hermann HC. Issues and recommendations for treating erectile dysfunction in men with ischemic heart disease. *Cardiovasc Rev Rep* 2001;22(5):285-91.
13. Chuang AT, Strauss JD, Murphy RA, et al. Sildenafil, a type-5 GMP phosphodiesterase inhibitor, specifically amplifies endogenous cGMP-dependent relaxation in rabbit *corpus cavernosum* smooth muscle *in vitro*. *J Urol* 1998;160(1):257-61.

14. DeBusk RF, Pepine CJ, Glasser DB, et al. Efficacy and safety of sildenafil citrate in men with erectile dysfunction and stable coronary artery disease. *Am J Cardiol* 2004;93:147-53.
15. Derry F, Gardner BP, Glass C, et al. Sildenafil (ViagraTM): a double-blind, placebo-controlled, single-dose, two-way crossover study in men with erectile dysfunction caused by traumatic spinal cord injury. *J Urol* 1997;157(4 Suppl):181.
16. Derry F, Glass C, Dinsmore WW, et al. Sildenafil (ViagraTM): An oral treatment for men with erectile dysfunction caused by traumatic spinal cord injury - A 28-day, double-blind, placebo-controlled, parallel-group, dose-response study. 49th Annual Meeting of the American Academy of Neurology. April 12-19, 1997; Boston, Massachusetts, USA. *Neurology* 1997;48(3 Suppl.2):A215.
17. Dogterom P, Zbinden G. Cardiotoxicity of Vasodilators and Positive Inotropic/Vasodilating Drugs in Dogs: An Overview. *Critical Reviews in Toxicology*, 22:203-241(1992).
18. Eardley I, Morgan RJ, Dinsmore WW, et al. UK-92,480, a new oral therapy for erectile dysfunction, a double-blind, placebo controlled trial with treatment taken as required. *J Urol* 1996; 155(No 5, Suppl):495A.
19. Eardley I, Ellis P, Boolel M, et al. Onset and duration of action of sildenafil citrate for the treatment of erectile dysfunction. *Br J Clin Pharmacol* 2002;53:61S-65S.
20. Fox KM, Thadani U, Ma PTS, et al. Sildenafil citrate does not reduce exercise tolerance in men with erectile dysfunction and chronic stable angina. *Eur Heart J* 2003;24:2206-12.
21. Gingell C, Jardin A, Olsson A, et al. UK-92,480, a new oral treatment for erectile dysfunction: a double-blind, placebo-controlled, once daily dose response study. *J Urol* 1996;155(No 5,Suppl):495A.
22. Gingell C, Sultana S, Wulff M, et al. Duration of action sildenafil citrate in men with erectile dysfunction. *J Sex Med* 2004; 1: 179-84.
23. Glass C, Derry F, Dinsmore WW, et al. Sildenafil (VIAGRATM): an oral treatment for men with erectile dysfunction caused by traumatic spinal cord injury - a double-blind, placebo-controlled, single-dose, two-way crossover study using RigiScan. *J Spinal Cord Med* 1997;20:145. American Spinal Injury Association Annual Meeting; March 24-26, 1997; Houston, TX.
24. Goldstein I, Lue TF, Padma-Nathan H, et al. Oral sildenafil in the treatment of erectile dysfunction. *N Engl J Med* 1998;338(20):1397-404.
25. Guay AT, Perez JB, Jacobson J, et al. Efficacy and safety of sildenafil citrate for treatment of erectile dysfunction in a population with associated organic risk factors. *J Androl* 2001;22(5):793-7.

26. Hatzichristou DG. Sildenafil citrate: lessons learned from 3 years of clinical experience. *Int J Impot Res* 2002;14(Suppl 1):S43-S52.
27. Hayes TJ, Roberts GKS, Halliwell WH. An Idiopathic Febrile Necrotizing Arteritis Syndrome in the Dog: Beagle Pain Syndrome. *Toxicologic Pathology* 1989;17:129-37.
28. Herrmann HC, Chang G, Klugherz BD, et al. Hemodynamic effects of sildenafil in men with severe coronary artery disease. *N Engl J Med* 2000;342:1622-6.
29. Jackson, G. Sildenafil (ViagraTM): new data, new confidence in treating erectile dysfunction in the cardiovascular patient. *IJCP* 2002;56(2);75.
30. Jetter A, Kinzing-Schippers M, Walchner-Bonjean M, et al. Sildenafil bioavailability is moderately increased by grapefruit juice. *Eur J Clin Pharmacol* 2001;57(8):A35 (abstract).
31. Jetter A, Kinzing-Schippers M, Walchner-Bonjean M, et al. Effects of grapefruit juice on the pharmacokinetics of sildenafil. *Clin Pharmacol Ther* 2002;71(1):21-9.
32. Kloner, RA, Zusman RM. Cardiovascular effects of sildenafil citrate and recommendations for its use. *Am J Cardiol* 1999;84 (5B):11N-17N.
33. Kloner RA. Cardiovascular risk and sildenafil. *Am J Cardiol* 2000;86(2A): 57F-61F.
34. Kloner RA, Morris B, Prisant LM, et al. Effect of sildenafil in patients with erectile dysfunction taking antihypertensive therapy. *Am J Hyperten* 2001;14:70-3.
35. Krenzelok, EP. Sildenafil: clinical toxicology profile. *J Clin Toxicol* 2000;38(6):645-51.
36. Langtry, HD, Markham A. Sildenafil: a review of its use in erectile dysfunction. *Drugs* 1999;57(6):967-89.
37. Laties AM, Fraunfelder FT. Ocular safety of Viagra. *Trans Am Ophtalmol Soc* 1999;97:115-25.
38. Lee JC, Bernard F, Carrier S, Talwar V, Defoy I. Do Men with Mild Erectile Dysfunction have the Same Risk Factors as the General Erectile Dysfunction Clinical Trial Population? *BJU International* 2010 Oct. 15 [Epub ahead of print].
39. Lue TF, Sildenafil Study Group. A study of Sildenafil (ViagraTM), a new oral agent for the treatment of male erectile dysfunction. 92nd Annual Meeting of the American Urological Association. April 12-17, 1997; New Orleans, Louisiana, USA. *J Urol* 1997; 157(4 Suppl):181.
40. McCullough A, Barada J, Fawzy A, et al. Achieving treatment optimisation with sildenafil citrate (Viagra[®]) in patients with erectile dysfunction. *Urology* 2002; 60 (2B Suppl): 28-38.

41. Morales A, Gingell, C, Collins M, et al. Clinical safety of oral sildenafil citrate (VIAGRA™) in the treatment of erectile dysfunction. *Int J Impot Res* 1998;10:69-74.
42. Morales A, Gajewski JB, Pommerville PJ, et al. 3-year efficacy and safety of Viagra® in men with erectile dysfunction: results of Canadian long-term extension study. *Can J Urol* 2003; 10(3):1826 [abstract].
43. Moreland RB, Goldstein I, Traish A. Sildenafil, a novel inhibitor of phosphodiesterase type 5 in human *corpus cavernosum* smooth muscle cells. *Life Sci* 1998;62(20):309-18.
44. Muirhead GJ, Allen MJ, James GC, et al. Pharmacokinetics of sildenafil (Viagra™), a selective cGMP PDE5 inhibitor, after single oral doses in fasted and fed healthy volunteers. *Br J Clin Pharmacol* 1996;42(No 2):268P.
45. Muirhead GJ, Faulkner S, Harness JA, et al. The effects of steady-state erythromycin and azithromycin on the pharmacokinetics of sildenafil citrate in healthy volunteers. *Br J Clin Pharmacol* 2002;53:37S-43S.
46. Muller, JE. Triggering of Cardiac events by sexual activity: findings from crossover analysis. *Am J Cardiol* 2000;86(2A):14F-18F.
47. Olson, AM, Persson CA. Efficacy and safety of sildenafil citrate for the treatment of erectile dysfunction in men with cardiovascular disease. *Int J Clin Pract* 2001;55(3):171-6.
48. Osterloh, I, Eardley, I, Carson, C, et al. Sildenafil: A selective phosphodiesterase (PDE)5 inhibitor in the treatment of erectile dysfunction (ED). In: Carson, C., Kirby, R., Goldstein, I., eds. *Textbook of Erectile Dysfunction*. Oxford, UK: Isis Media LTD; 1999: 285-307.
49. Padma-Nathan H, Steers WD, Wicker PA. Efficacy and safety of oral sildenafil in the treatment of erectile dysfunction: a double-blind, placebo-controlled study of 329 patients. *Int J Clin Prac* 1998;52(6):1-5.
50. Padma-Nathan H, Eardley I, Kloner R, et al. A 4-year update on the safety of sildenafil citrate (Viagra®). *Urology* 2002; 60 (2B Suppl): 67-90.
51. Partizi R, Leonardo F, Pelliccia F, et al. Effect of sildenafil citrate upon myocardial ischemia in patients with chronic stable angina in therapy with beta-blockers. *Ital Heart J* 2001;2(11):841-4.
52. Pfizer Dept. of Medicine. Sildenafil (Viagra®): A new drug in trial for the treatment of erectile dysfunction. *Urologe [B]* 1997;37:51.
53. Piccirillo G, Nocco M, Lionetti M, et al. Effects of sildenafil citrate (Viagra) on cardiac repolarization and on automatic control in subjects with chronic heart failure. *Am Heart J* 2002;143(4):703-10.

54. Pickering T, Shepard A, Puddey I, et al. Sildenafil citrate for erectile dysfunction in men receiving multiple antihypertensive agents - a randomized controlled trial. *Am J Hypertens* 2004; 17:1135-42.
55. Price D, Wareham K, Gingell CJ, et al. Sildenafil (Pfizer UK-92,480), a novel oral treatment for erectile dysfunction (ED) in patients with diabetes. *Diabetes* 1996; 45(Suppl 2):6A.
56. Rosen RC, Riley A, Wagner G, et al. The International Index of Erectile Function (IIEF): A multidimensional scale for assessment of erectile dysfunction. *Urology* 1997;49(6):822-30.
57. Shakir SAW, Wilton LV, Boshier A, et al. Cardiovascular events in users of sildenafil: results from first phase of prescription event monitoring in England. *B Med J* 2001; 322:651-2.
58. Steers W, Guay AT, Leriche A, et al. Assessment of the efficacy and safety of Viagra[®] (sildenafil citrate) in men with erectile dysfunction during long-term treatment. *Int J Impot Res* 2001;13:261-7.
59. Steif CG, Ückert S, Becker AJ, et al. The effect of the specific phosphodiesterase (PDE) inhibitors on human and rabbit cavernous tissue *in vitro* and *in vivo*. *J Urol* 1998;159(4):1390-3.
60. Terrett NK, Bell AS, Brown D, et al. UK-92,480, a potent and selective inhibitor of type V_A cGMP phosphodiesterase with utility for the treatment of male erectile dysfunction. *Abstr Papers Am Chem Soc* 1995; 210 Meet(Pt 2):MED 229.
61. Terrett NK, Bell AS, Brown D, et al. Sildenafil (ViagraTM), a potent and selective inhibitor of type 5 GMP phosphodiesterase with utility for the treatment of male erectile dysfunction. *Bioorganic & Medicinal Chemistry Letters* 1996; 6(15): 1819-24.
62. Vitezic, D. A risk-benefit assessment of sildenafil in the treatment of erectile dysfunction. *Drug Safety* 2001; 24(4): 255-65.
63. Warrington, JS, Shader, RI, von-Moltke, LL, et al. *In vitro* biotransformation of sildenafil (Viagra): identification of human cytochromes and potential drug interactions. *Drug Metab Dispos* 2000; 28(4):392-7.
64. Product monograph of PrViagra[®] (sildenafil citrate tablets), Pfizer Canada Inc. Submission Control No: 218856, Date of Revision: 18 September 2018.

PART III: CONSUMER INFORMATION

Pr ACCEL-SILDENAFIL
sildenafil citrate tablets
25 mg, 50 mg and 100 mg

This leaflet is part III of a three-part "Product Monograph" published when ACCEL-SILDENAFIL was approved for sale in Canada and is designed specifically for Consumers. This leaflet is a summary and will not tell you everything about ACCEL-SILDENAFIL. Contact your doctor or pharmacist if you have any questions about the drug.

ABOUT THIS MEDICATION

What the medication is used for:

ACCEL-SILDENAFIL (sildenafil citrate) is a treatment for erectile dysfunction in male adults. This is when a man cannot achieve or maintain a hard, erected penis for satisfactory sexual activity.

It is important to note that ACCEL-SILDENAFIL works only with sexual stimulation. ACCEL-SILDENAFIL alone does not increase your sex drive.

What it does:

ACCEL-SILDENAFIL belongs to a class of medicine called phosphodiesterase type 5 (PDE5) inhibitors. Following sexual stimulation, ACCEL-SILDENAFIL works by helping to relax the blood vessels in your penis by allowing blood to flow into your penis. This results in improved erectile function.

When it should not be used:

- If you are taking any medicines containing nitrates in any form (oral, sublingual [under the tongue], skin patch, or by inhalation [spray]). Although ACCEL-SILDENAFIL is used occasionally, nitrates must never be used. Nitrates are found in many prescription medicines that are used in the treatment of angina pectoris (chest pain due to heart disease), such as nitroglycerin, isosorbide mononitrate, or isosorbide dinitrate. If you do not understand what nitrates are, or are unsure about whether a medication you are taking is a "nitrate", ask your doctor or pharmacist.

If you take ACCEL-SILDENAFIL with nitrate-containing medicines or any other nitrates (e.g., amyl nitrite "poppers"), your blood pressure could suddenly drop to a life-threatening level. You could get dizzy, faint, or even have a heart attack or stroke.

- If you have loss of vision in one or both eyes from an eye disease called non-arteritic anterior ischaemic optic neuropathy (NAION)
- If you have ever had an allergic reaction to sildenafil

citrate or the nonmedicinal ingredients in ACCEL-SILDENAFIL.

- ACCEL-SILDENAFIL is not to be used in men for whom sexual activity is inadvisable.
- Do not take ACCEL-SILDENAFIL with guanylate cyclase stimulators, such as riociguat.

What the medicinal ingredient is:

Sildenafil citrate.

What the nonmedicinal ingredients are:

Calcium hydrogen phosphate anhydrous, indigo carmine aluminium lacquer, macrogol 3350, magnesium stearate, microcrystalline cellulose, polyvinilic alcohol, sodium croscarmellose, talc and titanium dioxide.

What dosage forms it comes in:

ACCEL-SILDENAFIL 25 mg, tablets, are elliptic, blue, film-coated tablets, embossed with a "-" on one side.

ACCEL-SILDENAFIL 50 mg, tablets, are elliptic, blue, film-coated tablets.

ACCEL-SILDENAFIL 100 mg, tablets, are elliptic, blue, film-coated tablets, embossed with a "+" on one side.

WARNINGS AND PRECAUTIONS

This medicine has been prescribed for you personally and you should not pass it on to others. It may harm them, even if their symptoms are the same as yours.

BEFORE you use ACCEL-SILDENAFIL talk to your doctor or pharmacist if you:

- have heart problems (irregular heartbeats, angina, chest pain, or had a heart attack). If you have heart problems, ask your doctor if your heart is healthy enough to handle the extra strain of having sex. If you have chest pain, dizziness or nausea during sex, stop exerting yourself. You should **not** use nitrates but you should seek immediate medical assistance.
- are 65 years of age or over
- have had a stroke
- have low blood pressure or uncontrolled high blood pressure
- have liver or kidney problems
- have sickle cell anemia (abnormality of the red blood cells), multiple myeloma (cancer of the bone marrow) or leukaemia (cancer of the white blood cells)
- are allergic to sildenafil or any of the other ingredients of ACCEL-SILDENAFIL tablets
- have a deformed penis or Peyronie's disease
- have ever had an erection that lasted more than 4 hours
- have stomach ulcers or other bleeding problems.
- have a rare inherited eye disease called retinitis

pigmentosa

- have had temporary, decrease, or permanent loss of vision in one or both eyes, including a condition called Non-Arteritic Anterior Ischemic Optic Neuropathy (NAION). The specific type of vision decrease or loss, called non-arteritic anterior ischemic optic neuropathy (NAION), seems to occur rarely when blood flow to the optic nerve is reduced or blocked. Vision decrease or loss may be partial or complete, in one or very occasionally both eyes. While in some cases the condition may improve over time, it can also be irreversible.
- Rare but serious skin reactions of Stevens-Johnson's Syndrome (SJS) and Toxic Epidermal Necrolysis (TEN) have been reported during the post-marketing period.

ACCEL-SILDENAFIL is not recommended for use in children under 18 years of age and in women.

ACCEL-SILDENAFIL does not protect against sexually transmitted diseases (STD), including Human Immunodeficiency Virus (HIV).

Alcohol consumption may decrease the ability to get an erection.

If you are taking **ACCEL-SILDENAFIL** and experience temporary, decrease, or permanent loss of vision, stop taking **ACCEL-SILDENAFIL** and call your doctor.

In case of chest pain occurring during or after sexual activity you should **not** use nitrates but you should seek immediate medical assistance.

Sudden decrease or loss of hearing has been reported in a few number of postmarketing and clinical trial cases with the use of PDE5 inhibitors, including sildenafil citrate. It has not been established whether these are related directly to the use of these medications or to other factors. If you experience these symptoms, stop taking **ACCEL-SILDENAFIL** and call your doctor.

Driving and using machines: Before you perform tasks which may require special attention, wait until you know how you respond to **ACCEL-SILDENAFIL**. Dizziness or altered vision can occur while using **ACCEL-SILDENAFIL**.

INTERACTIONS WITH THIS MEDICATION

You should tell your doctor or pharmacist about any other medications that you are taking. **ACCEL-SILDENAFIL** may interfere with some drugs. Only take **ACCEL-SILDENAFIL** with other drugs if recommended by your doctor.

Drugs that may interact with ACCEL-SILDENAFIL include:

- any drugs that contain nitrates in any form (oral, sublingual [under the tongue], skin patch or by inhalation [spray]). Nitrates are found in many prescriptions that are used to treat angina pectoris (chest pain due to heart disease). You should not take **ACCEL-SILDENAFIL** if you are taking these drugs.
- alpha-blockers (drugs used to treat prostate problems or high blood pressure)
- ketoconazole or itraconazole (drugs used to treat fungal infections)
- erythromycin (a drug used to treat bacterial infections)
- ritonavir, saquinavir or other drugs for the treatment of HIV
- cimetidine (a drug generally used to treat duodenal or gastric problems)
- bosentan (a drug used in the treatment of high blood pressure in the blood vessels between the heart and the lungs)
- drugs to treat erectile dysfunction. These drugs are Cialis, Levitra, Adcirca and Staxyn.
- other drugs that contain sildenafil. They can be called Revatio. They are used in the treatment of high blood pressure in the blood vessels between the heart and the lungs.

Grapefruit juice may increase the levels of **ACCEL-SILDENAFIL** in your blood.

PROPER USE OF THIS MEDICATION

Usual dose:

Your doctor can determine the dose that works best for you. Always take **ACCEL-SILDENAFIL** as prescribed by your doctor. You should speak with your doctor or pharmacist if you are unsure. A dose above 100 mg per day is not recommended.

Swallow the tablet whole, with some water.

You should **not** take more than one dose of **ACCEL-SILDENAFIL** per day.

If you have serious liver or kidney problems or you are 65 years of age or over, your doctor may start you at the lowest dose (25 mg) of **ACCEL-SILDENAFIL**.

How to optimize your use of ACCEL-SILDENAFIL:

You should take **ACCEL-SILDENAFIL** approximately 30 to 60 minutes before sexual activity. You can engage in sexual activity within 30 minutes of taking the tablet and for up to 4 hours. The amount of time it takes to have an effect varies slightly from person to person. Remember, sexual stimulation is required for **ACCEL-SILDENAFIL**.

to work.

You should avoid excessive drinking of alcohol, since alcohol can temporarily impair the ability to get an erection.

ACCEL-SILDENAFIL can be taken with or without food. However, you may find that it takes longer for **ACCEL-SILDENAFIL** to work if you take it with a high-fat meal.

ACCEL-SILDENAFIL may not work the first time or every time. If **ACCEL-SILDENAFIL** did not work for you on one occasion, try again on another day. You will learn how well **ACCEL-SILDENAFIL** works for you through your personal experience. The first few times may be charged with emotion or anxiety. If after a few separate attempts, you do not get the results you expect, talk to your doctor.

Overdose:

You should **not** take more than one dose of **ACCEL-SILDENAFIL** per day.

If you have taken more **ACCEL-SILDENAFIL** than you should, contact your doctor or a poison control centre immediately.

SIDE EFFECTS AND WHAT TO DO ABOUT THEM

As with most drugs, **ACCEL-SILDENAFIL** can cause some side effects. These effects are usually mild to moderate in nature and do not last for a long time.

Side effects may include:

- headache, facial flushing
- nausea, vomiting, indigestion, abdominal pain,
- dizziness
- dry, stuffy, or swollen nose
- throat tightness, dry mouth, decreased sensitivity of the mouth
- pain in arms and legs, myalgia (muscle pain)
- somnolence
- erection increased

If you notice any side effects not mentioned above, or any of the above-mentioned side effects persist or become bothersome, please contact your doctor or pharmacist.

SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM				
	Symptom / effect	Talk with your doctor or pharmacist		Stop taking drug and call your doctor or pharmacist
		Only if severe	In all cases	
Less common	Effect on hearing: sudden decrease or loss of hearing		✓	
	Effects on vision: colour tinge, increased brightness of light, blurred vision			✓
	Impaired or sudden loss of vision: decreased eyesight or unable to see with one or both eyes			✓
	Detached retina: a decrease in, or sudden loss of vision in one or both eyes			✓
Rare	Serious skin reactions: rash, blisters, peeling skin and pains.			✓
Very rare	Priapism: erection lasting more than 4 hours			✓
Unknown	Cough		✓	
	Allergic reactions: rash, hives, itch, swelling of the face, lips, tongue or throat, difficulty swallowing or breathing			✓
	Nosebleed		✓	
	<i>Noted in patients taking sildenafil citrate for pulmonary hypertension:</i>		✓	
	Fever			
	Shortness of breath		✓	
	Seizure, seizure recurrence			✓
	Transient global amnesia: temporary memory loss		✓	
	Heart attack (myocardial infarction): chest pain, shortness of breath			✓
	Chest pain (unstable angina)			✓
	Arrhythmia/tachycardia, palpitations: fast or irregular heartbeat, heart rate increased			✓
	Hypotension (low blood pressure): dizziness, fainting, lightheadedness	✓		
	Stroke (cerebrovascular hemorrhage): bleeding in the brain, vision changes, difficulty speaking, weakness on one side of the body, dizziness, lack of			✓

SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM

Symptom / effect	Talk with your doctor or pharmacist		Stop taking drug and call your doctor or pharmacist
	Only if severe	In all cases	
coordination or poor balance			
Transient ischaemic attack: temporary loss of vision, difficulty speaking, weakness on one side of the body, numbness or tingling usually on one side of the body, dizziness, lack of coordination or poor balance.			✓
Pulmonary Hemorrhage (acute bleeding from the lung): oozing of bloody fluid from the nose and respiratory tract, accompanied by rapid worsening of patient respiration, turning blue and in severe cases, shock)			✓

This is not a complete list of side effects. For any unexpected effects while taking ACCEL-SILDENAFIL, contact your doctor or pharmacist

HOW TO STORE IT

- Store between 15°C and 30°C, in the original package. Do not freeze.
- Do not take ACCEL-SILDENAFIL after the expiry date shown on the package.
- Always keep ACCEL-SILDENAFIL out of reach and sight of children.

REPORTING SIDE EFFECTS

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada/adverse-reaction-reporting.html>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

MORE INFORMATION

This document plus the full product monograph, prepared for health professionals can be found at <http://www.accelpharma.com> or by calling 1-877-822-2235.

This leaflet was prepared by Accel Pharma Inc.

Last revised: December 27, 2018